A Berry-Esseen bound of order $1/\sqrt{n}$ for martingales

Songqi Wu^a Xiaohui Ma^a Hailin Sang^b Xiequan Fan^a

^aCenter for Applied Mathematics, Tianjin University, Tianjin, China ^bDepartment of Mathematics, The University of Mississippi, University, MS 38677, USA

Received *****; accepted after revision +++++

Presented by

Abstract

Renz [13] has established a rate of convergence $1/\sqrt{n}$ in the central limit theorem for martingales with some restrictive conditions. In the present paper a modification of the methods, developed by Bolthausen [2] and Grama and Haeusler [6], is applied for obtaining the same convergence rate for a class of more general martingales. An application to linear processes is discussed. To cite this article: A. Name1, A. Name2, C. R. Acad. Sci. Paris, Ser. I 340 (2005).

Résumé

Une limite de Berry-Esseen de l'ordre $1/\sqrt{n}$ pour les martingales. Renz [13] a établi un taux de convergence $1/\sqrt{n}$ dans le théorème de la limite centrale pour les martingales avec certaines conditions restrictives. Dans le présent article, une modification des méthodes, développées par Bolthausen [2] et Grama et Haeusler [6], a appliqué pour obtenir le même taux de convergence pour une classe de martingales plus générales. Une application aux processus linéaires est discutée. Pour citer cet article : A. Name1, A. Name2, C. R. Acad. Sci. Paris, Ser. I 340 (2005).

1. Introduction and main result

For $n \in \mathbf{N}$, let $(\xi_i, \mathcal{F}_i)_{i=0,...,n}$ be a finite sequence of martingale differences defined on some probability space $(\Omega, \mathcal{F}, \mathbf{P})$, where $\xi_0 = 0$ and $\{\emptyset, \Omega\} = \mathcal{F}_0 \subseteq ... \subseteq \mathcal{F}_n \subseteq \mathcal{F}$ are increasing σ -fields. Denote

$$X_0 = 0, \quad X_k = \sum_{i=1}^k \xi_i, \ k = 1, ..., n.$$

17 janvier 2022

Email address: fanxiequan@hotmail.com (Xiequan Fan).

Preprint submitted to the Académie des sciences

Then $X = (X_k, \mathcal{F}_k)_{k=0,\dots,n}$ is a martingale. Denote by $\langle X \rangle$ the conditional variance of X :

$$\langle X \rangle_0 = 0, \quad \langle X \rangle_k = \sum_{i=1}^k \mathbf{E} [\xi_i^2 | \mathcal{F}_{i-1}], \ k = 1, ..., n$$

Define

$$D(X_n) = \sup_{x \in \mathbf{R}} |\mathbf{P}(X_n \le x) - \Phi(x)|,$$

where $\Phi(x)$ is the distribution function of the standard normal random variable. Denote by $\xrightarrow{\mathbf{P}}$ the convergence in probability as $n \to \infty$. According to the martingale central limit theorem, the "conditional Lindeberg condition"

$$\sum_{i=1}^{n} \mathbf{E} \left[\xi_{i}^{2} \mathbf{1}_{\{|\xi_{i}| \geq \epsilon\}} \middle| \mathcal{F}_{i-1} \right] \xrightarrow{\mathbf{P}} 0, \text{ for each } \epsilon > 0,$$

and the "conditional normalizing condition" $\langle X \rangle_n \xrightarrow{\mathbf{P}} 1$ together implies asymptotic normality of X_n , that is, $D(X_n) \to 0$ as $n \to \infty$.

The convergence rate of $D(X_n)$ has attracted a lot of attentions. For instance, Bolthausen [2] proved that if $|\xi_i| \leq \epsilon_n$ for a number ϵ_n and $\langle X \rangle_n = 1$ a.s., then $D(X_n) \leq c\epsilon_n^3 n \log n$, where, here and after, c is an absolute constant not depending on ϵ_n and n. El Machkouri and Ouchti [3] improved the factor $\epsilon_n^3 n \log n$ in Bolthausen's bound to $\epsilon_n \log n$ under the following more general condition

$$\mathbf{E}\left[\left|\xi_{i}\right|^{3}\left|\mathcal{F}_{i-1}\right]\right] \leq \epsilon_{n} \mathbf{E}\left[\xi_{i}^{2}\left|\mathcal{F}_{i-1}\right] \quad a.s. \text{ for all } i=1,2,...,n.$$

For more related results, we refer to Ouchti [12] and Mourrat [11]. Recently, Fan [4] proved that if there exist a positive constant ρ and a number ϵ_n , such that

$$\mathbf{E}\left[|\xi_i|^{2+\rho} \big| \mathcal{F}_{i-1}\right] \le \epsilon_n^{\rho} \mathbf{E}\left[\xi_i^2 \big| \mathcal{F}_{i-1}\right] \quad a.s. \text{ for all } i = 1, 2, ..., n,$$

and $\langle X \rangle_n = 1$ a.s., then $D(X_n) \leq c_\rho \hat{\epsilon}_n$, where

$$\hat{\epsilon}_n = \begin{cases} \epsilon_n^{\rho}, & \text{if } \rho \in (0,1) \\ \epsilon_n |\log \epsilon_n|, & \text{if } \rho \ge 1 \end{cases}$$

and c_{ρ} is a constant depending only on ρ . Fan also showed that this Berry-Esseen bound is optimal. In particular, if $\epsilon_n \approx 1/\sqrt{n}$, then we have $\epsilon_n |\log \epsilon_n| \approx (\log n)/\sqrt{n}$. Thus, we cannot obtain the classical convergence rate $1/\sqrt{n}$ for general martingales.

However, the convergence rate $1/\sqrt{n}$ for martingales is possible to be attained with some additional restrictive conditions. For instance, Renz [13] proved that if there exists a constant $\rho > 0$ such that

$$\mathbf{E}[\xi_{i}^{2}|\mathcal{F}_{i-1}] = 1/n, \quad \mathbf{E}[\xi_{i}^{3}|\mathcal{F}_{i-1}] = 0 \quad \text{and} \quad \mathbf{E}[|\xi_{i}|^{3+\rho}|\mathcal{F}_{i-1}] \le cn^{-(3+\rho)/2}, \quad \text{a.s.},$$
(1)

then it holds

$$D(X_n) = O\left(\frac{1}{\sqrt{n}}\right). \tag{2}$$

He also showed that this result is not true for $\rho = 0$. More martingale Berry-Esseen bounds of convergence rate $1/\sqrt{n}$ can be found in Bolthausen [2] and Kir'yanova and Rotar [10].

In this paper we are interested in extending (2) to a class of more general martingales. The following theorem is our main result.

Theorem 1.1 Assume that there exist some numbers $\rho \in (0, +\infty)$, $\epsilon_n \in (0, \frac{1}{2}]$ and $\delta_n \in [0, \frac{1}{2}]$ such that for all $1 \le i \le n$,

$$\left|\langle X\rangle_n - 1\right| \le \delta_n^2,\tag{3}$$

$$\mathbf{E}\left[\xi_{i}^{3}\big|\mathcal{F}_{i-1}\right] = 0\tag{4}$$

and

$$\mathbf{E}\left[\left|\xi_{i}\right|^{3+\rho}\left|\mathcal{F}_{i-1}\right] \leq \epsilon_{n}^{1+\rho} \mathbf{E}\left[\xi_{i}^{2}\left|\mathcal{F}_{i-1}\right] \quad a.s.$$

$$\tag{5}$$

Then

$$D(X_n) \le c_{\rho}(\epsilon_n + \delta_n),$$

where c_{ρ} depends only on ρ . In addition, it holds $c_{\rho} = O(\rho^{-1}), \rho \to 0$.

Notice that under the conditions of Renz [13], the conditions of Theorem 1.1 are satisfied with $\delta_n = 0$ and $\epsilon_n \simeq 1/\sqrt{n}$. Thus Theorem 1.1 extends Renz's result to a class of more general martingales.

Thanks to the additional condition (4), the Berry-Esseen bound (6) improves the bound of Fan [4] by replacing $\epsilon_n |\log \epsilon_n|$ with ϵ_n .

Relaxing the condition (3), we have the following analogue estimation of Fan (cf. (26) of [4]). **Theorem 1.2** Assume that there exist some numbers $\rho \in (0, +\infty)$ and $\epsilon_n \in (0, \frac{1}{2}]$ such that for all $1 \leq i \leq n$,

$$\mathbf{E}\left[\xi_i^3 \middle| \mathcal{F}_{i-1}\right] = 0$$

and

$$\mathbf{E}[|\xi_i|^{3+\rho} | \mathcal{F}_{i-1}] \le \epsilon_n^{1+\rho} \mathbf{E}[\xi_i^2 | \mathcal{F}_{i-1}] \quad a.s$$

Then, for all $p \geq 1$,

$$D(X_n) \le c_{\rho} \epsilon_n + c_p \left(\mathbf{E} \Big[\left| \langle X \rangle_n - 1 \right|^p \Big] + \mathbf{E} \Big[\max_{1 \le i \le n} |\xi_i|^{2p} \Big] \right)^{1/(2p+1)}, \tag{6}$$

where c_{ρ} and c_{p} depend only on ρ and p, respectively.

It is easy to see that when $p \to \infty$,

$$\left(\mathbf{E}\left[\left|\langle X\rangle_{n}-1\right|^{p}\right]\right)^{1/(2p+1)} \to ||\langle X\rangle_{n}-1||_{\infty}^{1/2},$$

which coincides with δ_n of Theorem 1.1.

2. Application

We first extend Theorem 1.1 to triangular arrays with infinity many terms in each line. For $n \in \mathbf{N}$, let $(\xi_{n,i}, \mathcal{F}_{n,i})_{i=-\infty}^n$ be a sequence of martingale differences defined on some probability space $(\Omega, \mathcal{F}, \mathbf{P})$, where the adapted filtration is $\{\emptyset, \Omega\} = \mathcal{F}_{-\infty} \subset ... \subset \mathcal{F}_{n,n-1} \subset \mathcal{F}_{n,n} \subset \mathcal{F}$. Denote $X_{n,k} = \sum_{i=-\infty}^{k} \xi_{n,i}, k \leq n$.

Then $(X_{n,k}, \mathcal{F}_{n,k})_{k=-\infty}^n$ is a martingale. Let $\langle X \rangle_{n,k} = \sum_{i=-\infty}^k \mathbf{E}[\xi_{n,i}^2 | \mathcal{F}_{n,i-1}], k \leq n$. In particular, denote $X_n := X_{n,n}$ and $\langle X \rangle_n := \langle X \rangle_{n,n}$.

With some slight modification on the proof, Theorem 1.1 still holds in this new setting. Now we apply Theorem 1.1 with this new setting to the partial sum of linear processes. Let $(\varepsilon_i)_{i \in \mathbf{Z}}$ be a sequence of identically distributed martingale differences adapted to the filtration $(\mathcal{F}_i)_{i \in \mathbf{Z}}$. We consider the causal linear process in the form

$$Y_k = \sum_{j=-\infty}^k a_{k-j}\varepsilon_j,\tag{7}$$

where the martingale differences have finite variance and the sequence of real coefficients satisfies $\sum_{i=0}^{\infty} a_i^2 < 1$ ∞ . Without loss of generality, let the variance of the martingale difference to be 1. We say the linear process has long memory if $\sum_{i=0}^{\infty} |a_i| = \infty$. In this case, we assume that $a_0 = 1$ and

$$a_i = l(i)i^{-\alpha}, \ i > 0, \ \text{with} \ 1/2 < \alpha < 1.$$
 (8)

Here $l(\cdot)$ is a slowly varying function. On the other hand, we say the linear process has short memory if $\sum_{i=0}^{\infty} |a_i| < \infty \text{ and } \sum_{i=0}^{\infty} a_i \neq 0. \text{ The third case is } \sum_{i=0}^{\infty} |a_i| < \infty \text{ and } \sum_{i=0}^{\infty} a_i = 0.$

The long memory linear processes covers the well-known fractional ARIMA processes (cf. Granger and Joyeux [7]; Hosking [9]), which play an important role in financial time series modeling and application. As a special case, let 0 < d < 1/2 and B be the backward shift operator with $B\varepsilon_k = \varepsilon_{k-1}$ and consider

$$Y_k = (1-B)^{-d} \varepsilon_k = \sum_{i=0}^{\infty} a_i \varepsilon_{k-i}, \text{ where } a_i = \frac{\Gamma(i+d)}{\Gamma(d)\Gamma(i+1)}$$

For this example we have $\lim_{n\to\infty} a_n/n^{d-1} = 1/\Gamma(d)$. Note that these processes have long memory because $\sum_{j=0}^{\infty} |a_j| = \infty.$

The partial sum $S_n = \sum_{k=1}^n Y_k$ of causal linear process (7) can be written as $S_n = \sum_{-\infty}^n b_{n,i}\varepsilon_i$, where $b_{n,i} = \sum_{j=0}^{n-i} a_j$ for $0 < i \le n$, and $b_{n,i} = \sum_{j=1-i}^{n-i} a_j$ for $i \le 0$. The variance of S_n is $B_n^2 =$ $var(S_n) = \sum_{i=-\infty}^{n} b_{n,i}^2. \text{ Now let } X_{n,k} = \sum_{i=-\infty}^{k} b_{n,i}\varepsilon_i/B_n. \text{ Then } X_n = X_{n,n} = S_n/B_n \text{ and } \langle X \rangle_n = \sum_{i=-\infty}^{n} b_{n,i}^2 \mathbf{E}[\varepsilon_i^2 | \mathcal{F}_{i-1}]/B_n^2. \text{ If we assume } |\langle X \rangle_n - 1| \leq \delta_n^2 \text{ for some } \delta_n \in [0, \frac{1}{2}], \mathbf{E}[\varepsilon_i^3 | \mathcal{F}_{i-1}] = 0 \text{ and}$ $\mathbf{E}[|\varepsilon_i|^{3+\rho}|\mathcal{F}_{i-1}] \leq d_{\rho}^{1+\rho} \mathbf{E}[\varepsilon_i^2|\mathcal{F}_{i-1}]$ a.s. for all $i \in \mathbf{Z}$ and some constant d_{ρ} , then, by Theorem 1.1,

$$\sup_{x \in \mathbf{R}} |\mathbf{P}(S_n/B_n \le x) - \Phi(x)| \le c_{\rho}(\epsilon_n + \delta_n),$$

where $\epsilon_n = d_\rho \sup_{i \le n} |b_{n,i}|/B_n$. In the case that $\sum_{i=0}^{\infty} |a_i| < \infty$, $\sup_{i \le n} |b_{n,i}| \le \sum_{i=0}^{\infty} |a_i| < \infty$ and it is well known that B_n^2 has order n. Hence ϵ_n has order $1/\sqrt{n}$ in this case. In the long memory case $\sum_{i=0}^{\infty} |a_i| = \infty$, if we assume (8), B_n^2 has order $n^{3-2\alpha}l^2(n)$ (e.g., Wu and Min [14]) and $\sup_{i\le n} |b_{n,i}|$ has order $n^{1-\alpha}l(n)$ (see Beknazaryan et al. [1] for upper bound and Fortune et al. [5] for lower bound in the case d = 1). Hence in this case ϵ_n also has order $1/\sqrt{n}$. In either case the Berry-Esseen bound has order $1/\sqrt{n}$ if $\delta_n = O(n^{-1/2})$. In particular if we in addition assume that the innovations $(\varepsilon_i)_{i \in \mathbb{Z}}$ are independent, then $\delta_n = 0$ and the Berry-Esseen bound $\sup_{x \in \mathbf{R}} |\mathbf{P}(S_n/B_n \leq x) - \Phi(x)|$ has order $1/\sqrt{n}$. Here the condition $\mathbf{E}[\varepsilon_i^3|\mathcal{F}_{i-1}] = 0$ is needed to have the Berry-Esseen bound of order $1/\sqrt{n}$. We cannot have this order from the result of Fan [4].

3. Proofs of theorems

3.1. Preliminary lemmas

In the proofs of theorems, we need the following technical lemmas. The first two lemmas can be found in Fan [4] (cf. Lemmas 3.1 and 3.2 therein). Lemma 3.1 If there exists an s > 3 such that

$$\mathbf{E}[|\xi_i|^s | \mathcal{F}_{i-1}] \le \epsilon_n^{s-2} \mathbf{E}[\xi_i^2 | \mathcal{F}_{i-1}],$$

then, for any $t \in [3, s)$,

 $\mathbf{E}[|\xi_i|^t | \mathcal{F}_{i-1}] \le \epsilon_n^{t-2} \mathbf{E}[\xi_i^2 | \mathcal{F}_{i-1}].$

Lemma 3.2 If there exists an s > 3 such that

$$\mathbf{E}[|\xi_i|^s | \mathcal{F}_{i-1}] \le \epsilon_n^{s-2} \mathbf{E}[\xi_i^2 | \mathcal{F}_{i-1}],$$

then

$$\mathbf{E}[\xi_i^2 | \mathcal{F}_{i-1}] \le \epsilon_n^2.$$

The next two technical lemmas are due to Bolthausen (cf. Lemmas 1 and 2 of [2]). Lemma 3.3 Let X and Y be random variables. Then

$$\sup_{u} \left| \mathbf{P} \left(X \le u \right) - \Phi(u) \right| \le c_1 \sup_{u} \left| \mathbf{P} \left(X + Y \le u \right) - \Phi(u) \right| + c_2 \left| \left| \mathbf{E} [Y^2 | X] \right| \right|_{\infty}^{1/2},$$

where c_1 and c_2 are two positive constants.

Lemma 3.4 Let G(x) be an integrable function on \mathbf{R} of bounded variation $||G||_V$, X be a random variable and $a, b \neq 0$ are real numbers. Then

$$\mathbf{E}\left[G\left(\frac{X+a}{b}\right)\right] \le ||G||_V \sup_{u} \left|\mathbf{P}\left(X \le u\right) - \Phi(u)\right| + ||G||_1|b|,$$

where $||G||_1$ is the $L_1(\mathbf{R})$ norm of G(x).

In the proof of Theorem 1.2, we also need the following lemma of El Machkouri and Ouchti [3]. Lemma 3.5 Let X and Y be two random variables. Then, for $p \ge 1$,

$$D(X+Y) \le 2D(X) + 3 \left\| \mathbf{E} [Y^{2p} | X] \right\|_{1}^{1/(2p+1)}.$$
(9)

3.2. Proof of Theorem 1.1

By Lemma 3.1, we only need to consider the case of $\rho \in (0, 1]$. We follow the method of Grama and Haeusler [6]. Let $T = 1 + \delta_n^2$. We introduce a modification of the conditional variance $\langle X \rangle_n$ as follows :

$$V_k = \langle X \rangle_k \mathbf{1}_{\{k < n\}} + T \mathbf{1}_{\{k = n\}}.$$
(10)

It is easy to see that $V_0 = 0, V_n = T$, and that $(V_k, \mathcal{F}_k)_{k=0,\dots,n}$ is a predictable process. Set

$$\gamma = \epsilon_n + \delta_n.$$

Let c_* be some positive and sufficient large constant. Define the following non-increasing discrete time predictable process

$$A_k = c_*^2 \gamma^2 + T - V_k, \quad k = 1, ..., n.$$
(11)

Obviously, we have $A_0 = c_*^2 \gamma^2 + T$ and $A_n = c_*^2 \gamma^2$. In addition, for $u, x \in \mathbf{R}$, and y > 0, denote

$$\Phi_u(x,y) = \Phi\left(\frac{u-x}{\sqrt{y}}\right). \tag{12}$$

Let $\mathcal{N} = \mathcal{N}(0, 1)$ be a standard normal random variable, which is independent of X_n . Using a smoothing procedure, by Lemma 3.3, we deduce that

$$\sup_{u} \left| \mathbf{P} (X_{n} \leq u) - \Phi(u) \right| \leq c_{1} \sup_{u} \left| \mathbf{P} (X_{n} + c_{*} \gamma \mathcal{N} \leq u) - \Phi(u) \right| + c_{2} \gamma \\
= c_{1} \sup_{u} \left| \mathbf{E} [\Phi_{u} (X_{n}, A_{n})] - \Phi(u) \right| + c_{2} \gamma \\
\leq c_{1} \sup_{u} \left| \mathbf{E} [\Phi_{u} (X_{n}, A_{n})] - \mathbf{E} [\Phi_{u} (X_{0}, A_{0})] \right| \\
+ c_{1} \sup_{u} \left| \mathbf{E} [\Phi_{u} (X_{0}, A_{0})] - \Phi(u) \right| + c_{2} \gamma \\
= c_{1} \sup_{u} \left| \mathbf{E} [\Phi_{u} (X_{n}, A_{n})] - \mathbf{E} [\Phi_{u} (X_{0}, A_{0})] \right| \\
+ c_{1} \sup_{u} \left| \mathbf{E} [\Phi_{u} (X_{n}, A_{n})] - \mathbf{E} [\Phi_{u} (X_{0}, A_{0})] \right| \\
+ c_{1} \sup_{u} \left| \Phi \left(\frac{u}{\sqrt{c_{*}^{2} \gamma^{2} + T}} \right) - \Phi(u) \right| + c_{2} \gamma.$$
(13)

It is obvious that

$$\left|\Phi\left(\frac{u}{\sqrt{c_*^2\gamma^2 + T}}\right) - \Phi(u)\right| \le c_3 \left|\frac{1}{\sqrt{c_*^2\gamma^2 + T}} - 1\right| \le c_4\gamma.$$
(14)

Returning to (13), we get

$$\sup_{u} \left| \mathbf{P} \left(X_n \le u \right) - \Phi(u) \right| \le c_1 \sup_{u} \left| \mathbf{E} \left[\Phi_u \left(X_n, A_n \right) \right] - \mathbf{E} \left[\Phi_u \left(X_0, A_0 \right) \right] \right| + c_5 \gamma.$$
(15)

By a simple telescoping, we know that

$$\mathbf{E}\left[\Phi_u(X_n, A_n)\right] - \mathbf{E}\left[\Phi_u(X_0, A_0)\right] = \mathbf{E}\left[\sum_{k=1}^n \left(\Phi_u(X_k, A_k) - \Phi_u(X_{k-1}, A_{k-1})\right)\right].$$
(16)

Taking into account the fact that

$$\frac{\partial^2}{\partial x^2} \Phi_u(x,y) = 2\frac{\partial}{\partial y} \Phi_u(x,y),$$

we get

$$\mathbf{E}\left[\Phi_u(X_n, A_n)\right] - \mathbf{E}\left[\Phi_u(X_0, A_0)\right] = J_1 + J_2 - J_3,\tag{17}$$

where

$$J_{1} = \mathbf{E} \bigg[\sum_{k=1}^{n} \bigg(\Phi_{u}(X_{k}, A_{k}) - \Phi_{u}(X_{k-1}, A_{k}) - \frac{\partial}{\partial x} \Phi_{u}(X_{k-1}, A_{k}) \xi_{k} - \frac{1}{2} \frac{\partial^{2}}{\partial x^{2}} \Phi_{u}(X_{k-1}, A_{k}) \xi_{k}^{2} - \frac{1}{6} \frac{\partial^{3}}{\partial x^{3}} \Phi_{u}(X_{k-1}, A_{k}) \xi_{k}^{3} \bigg) \bigg],$$
(18)

$$J_2 = \frac{1}{2} \mathbf{E} \bigg[\sum_{k=1}^n \frac{\partial^2}{\partial x^2} \Phi_u(X_{k-1}, A_k) \Big(\bigtriangleup \langle X \rangle_k - \bigtriangleup V_k \Big) \bigg], \tag{19}$$

$$J_3 = \mathbf{E} \bigg[\sum_{k=1}^n \bigg(\Phi_u(X_{k-1}, A_{k-1}) - \Phi_u(X_{k-1}, A_k) - \frac{\partial}{\partial y} \Phi_u(X_{k-1}, A_k) \bigtriangleup V_k \bigg) \bigg],$$
(20)

where $\triangle \langle X \rangle_k = \langle X \rangle_k - \langle X \rangle_{k-1}$. Now, we need to give some estimates of J_1, J_2 and J_3 . To this end, we introduce some notations. Denote by ϑ_i some random variables satisfying $0 \le \vartheta_i \le 1$, which may represent different values at different places. For the rest of the paper, φ stands for the density function of the standard normal random variable.

Control of J_1 : For convenience's sake, let $T_{k-1} = (u - X_{k-1})/\sqrt{A_k}, k = 1, 2, ..., n$. It is easy to see that

$$B_{k} =: \Phi_{u}(X_{k}, A_{k}) - \Phi_{u}(X_{k-1}, A_{k}) - \frac{\partial}{\partial x} \Phi_{u}(X_{k-1}, A_{k})\xi_{k}$$
$$-\frac{1}{2}\frac{\partial^{2}}{\partial x^{2}}\Phi_{u}(X_{k-1}, A_{k})\xi_{k}^{2} - \frac{1}{6}\frac{\partial^{3}}{\partial x^{3}}\Phi_{u}(X_{k-1}, A_{k})\xi_{k}^{3}$$
$$= \Phi\left(T_{k-1} - \frac{\xi_{k}}{\sqrt{A_{k}}}\right) - \Phi(T_{k-1}) + \Phi'(T_{k-1})\frac{\xi_{k}}{\sqrt{A_{k}}}$$
$$-\frac{1}{2}\Phi''(T_{k-1})\left(\frac{\xi_{k}}{\sqrt{A_{k}}}\right)^{2} + \frac{1}{6}\Phi'''(T_{k-1})\left(\frac{\xi_{k}}{\sqrt{A_{k}}}\right)^{3}.$$

To estimate the right hand side of the last equality, we distinguish two cases.

Case 1: $|\xi_k/\sqrt{A_k}| \le 2 + |T_{k-1}|/2$. By a four-term Taylor expansion, it is obvious that if $|\xi_k/\sqrt{A_k}| \le 1$, then

$$B_{k} = \left| \frac{1}{24} \Phi^{(4)} \left(T_{k-1} - \vartheta \frac{\xi_{k}}{\sqrt{A_{k}}} \right) \left| \frac{\xi_{k}}{\sqrt{A_{k}}} \right|^{4} \right|$$
$$\leq \left| \Phi^{(4)} \left(T_{k-1} - \vartheta \frac{\xi_{k}}{\sqrt{A_{k}}} \right) \right| \left| \frac{\xi_{k}}{\sqrt{A_{k}}} \right|^{3+\rho}.$$

If $|\xi_k/\sqrt{A_k}| > 1$, by a three-term Taylor expansion, then

$$\begin{aligned} \left| B_k \right| &\leq \frac{1}{2} \left(\left| \Phi^{\prime\prime\prime} \left(T_{k-1} - \vartheta \frac{\xi_k}{\sqrt{A_k}} \right) \right| + \left| \Phi^{\prime\prime\prime} (T_{k-1}) \right| \right) \left| \frac{\xi_k}{\sqrt{A_k}} \right|^3 \\ &\leq \left| \Phi^{\prime\prime\prime} \left(T_{k-1} - \vartheta^\prime \frac{\xi_k}{\sqrt{A_k}} \right) \right| \left| \frac{\xi_k}{\sqrt{A_k}} \right|^3 \\ &\leq \left| \Phi^{\prime\prime\prime} \left(T_{k-1} - \vartheta^\prime \frac{\xi_k}{\sqrt{A_k}} \right) \right| \left| \frac{\xi_k}{\sqrt{A_k}} \right|^{3+\rho}, \end{aligned}$$

where

$$\vartheta' = \begin{cases} \vartheta, \text{ if } \left| \Phi''' \left(T_{k-1} - \vartheta \frac{\xi_k}{\sqrt{A_k}} \right) \right| \ge \left| \Phi'''(T_{k-1}) \right|, \\ 0, \text{ if } \left| \Phi''' \left(T_{k-1} - \vartheta \frac{\xi_k}{\sqrt{A_k}} \right) \right| < \left| \Phi'''(T_{k-1}) \right|. \end{cases}$$

Using the inequality $\max\{|\Phi^{\prime\prime\prime}(t)|,|\Phi^{\prime\prime\prime\prime}(t)|\}\leq \varphi(t)(2+t^4),$ we find that

$$\left| B_{k} \mathbf{1}_{\{|\xi_{k}/\sqrt{A_{k}}| \leq 2+|T_{k-1}|/2\}} \right| \leq \varphi \left(T_{k-1} - \vartheta_{1} \frac{\xi_{k}}{\sqrt{A_{k}}} \right) \left(2 + \left(T_{k-1} - \vartheta_{1} \frac{\xi_{k}}{\sqrt{A_{k}}} \right)^{4} \right) \left| \frac{\xi_{k}}{\sqrt{A_{k}}} \right|^{3+\rho} \leq g_{1}(T_{k-1}) \left| \frac{\xi_{k}}{\sqrt{A_{k}}} \right|^{3+\rho},$$

$$(21)$$

where

$$g_1(z) = \sup_{|t-z| \le 2+|z|/2} \varphi(t)(2+t^4).$$

Case 2 : $|\xi_k/\sqrt{A_k}| > 2 + |T_{k-1}|/2$. It is obvious that, for $|\bigtriangleup x| > 1 + |x|/2$,

$$\begin{split} \left| \Phi(x - \bigtriangleup x) - \Phi(x) + \Phi'(x) \bigtriangleup x - \frac{1}{2} \Phi''(x) (\bigtriangleup x)^2 + \frac{1}{6} \Phi'''(x) (\bigtriangleup x)^3 \right| \\ &\leq \left(\left| \frac{\Phi(x - \bigtriangleup x) - \Phi(x)}{|\bigtriangleup x|^3} \right| + |\Phi'(x)| + |\Phi''(x)| + |\Phi'''(x)| \right) |\bigtriangleup x|^3 \\ &\leq \left(8 \left| \frac{\Phi(x - \bigtriangleup x) - \Phi(x)}{(2 + |x|)^3} \right| + |\Phi'(x)| + |\Phi''(x)| + |\Phi'''(x)| \right) |\bigtriangleup x|^3 \\ &\leq \left(\frac{\tilde{c}}{(2 + |x|)^3} + |\Phi'(x)| + |\Phi''(x)| + |\Phi'''(x)| \right) |\bigtriangleup x|^3 \\ &\leq \frac{\hat{c}}{(2 + |x|)^3} |\bigtriangleup x|^3 \\ &\leq \frac{\hat{c}}{(2 + |x|)^3} |\bigtriangleup x|^{3+\rho}. \end{split}$$

Hence, we have

$$\left| B_k \mathbf{1}_{\{|\xi_k/\sqrt{A_k}|>2+|T_{k-1}|/2\}} \right| \le g_2(T_{k-1}) \left| \frac{\xi_k}{\sqrt{A_k}} \right|^{3+\rho},\tag{22}$$

where

$$g_2(z) = \frac{\hat{c}}{(2+|z|)^3}.$$

Denote

$$G(z) = g_1(z) + g_2(z).$$

Combining (21) and (22) together, we get

$$|B_k| \le G(T_{k-1}) \left| \frac{\xi_k}{\sqrt{A_k}} \right|^{3+\rho}.$$
(23)

Therefore,

$$\left|J_{1}\right| = \left|\mathbf{E}\left[\sum_{k=1}^{n} B_{k}\right]\right| \le \mathbf{E}\left[\sum_{k=1}^{n} G(T_{k-1}) \left|\frac{\xi_{k}}{\sqrt{A_{k}}}\right|^{3+\rho}\right].$$
(24)

Next, we consider conditional expectation of $|\xi_k|^{3+\rho}$. By condition (5), we get

$$\mathbf{E}\left[\left|\xi_{k}\right|^{3+\rho}\left|\mathcal{F}_{k-1}\right] \le \epsilon_{n}^{1+\rho} \bigtriangleup \langle X \rangle_{k},\tag{25}$$

where $riangle \langle X \rangle_k = \langle X \rangle_k - \langle X \rangle_{k-1}$. And we know that

$$\Delta \langle X \rangle_k = \Delta V_k = V_k - V_{k-1}, \ 1 \le k < n, \ \Delta \langle X \rangle_n \le \Delta V_n, \tag{26}$$

then

$$\mathbf{E}\left[\left|\xi_{k}\right|^{3+\rho}\left|\mathcal{F}_{k-1}\right] \le \epsilon_{n}^{1+\rho} \bigtriangleup V_{k}.$$
(27)

By (24) and (27), we obtain

$$|J_1| \le R_1 := \epsilon_n^{1+\rho} \bigg[\sum_{k=1}^n \frac{G(T_{k-1})}{A_k^{(3+\rho)/2}} \bigtriangleup V_k \bigg].$$
(28)

To estimate R_1 , we introduce the time change τ_t as follow : for any real $t \in [0, T]$,

 $\tau_t = \min\{k \le n : V_k \ge t\}, \text{ where } \min \emptyset = n.$ (29)

Obviously, for any $t \in [0, T]$, the stopping time τ_t is predictable. In addition, $(\sigma_k)_{k=1,...,n+1}$ (with $\sigma_1 = 0$) stands for the increasing sequence of moments when the increasing and stepwise function $\tau_t, t \in [0, T]$, has jumps. It is easy to see that $\Delta V_k = \int_{[\sigma_k, \sigma_{k+1})} dt$, and that $k = \tau_t$ for $t \in [\sigma_k, \sigma_{k+1})$. Since $\tau_T = n$, we have

$$\sum_{k=1}^{n} \frac{G(T_{k-1})}{A_{k}^{(3+\rho)/2}} \bigtriangleup V_{k} = \sum_{k=1}^{n} \int_{[\sigma_{k},\sigma_{k+1})} \frac{G(T_{\tau_{t}-1})}{A_{\tau_{t}}^{(3+\rho)/2}} dt = \int_{0}^{T} \frac{G(T_{\tau_{t}-1})}{A_{\tau_{t}}^{(3+\rho)/2}} dt.$$
(30)

Let $a_t = c_*^2 \gamma^2 + T - t$. Because of $\Delta V_{\tau_t} \leq 2\epsilon_n^2 + 2\delta_n^2$ (cf. Lemma 3.2), we know that

$$t \le V_{\tau_t} = V_{\tau_t - 1} + \triangle V_{\tau_t} \le t + 2\epsilon_n^2 + 2\delta_n^2, \quad t \in [0, T].$$
(31)

Assume $c_* \geq 2$, then we have

$$\frac{1}{2}a_t \le A_{\tau_t} = c_*^2 \gamma^2 + T - V_{\tau_t} \le a_t, \quad t \in [0, T].$$
(32)

Note that G(z) is symmetric and is non-increasing in $z \ge 0$. The last bound implies that

$$R_1 \le 2^{(3+\rho)/2} \epsilon_n^{1+\rho} \int_0^T \frac{1}{a_t^{(3+\rho)/2}} \mathbf{E} \Big[G\Big(\frac{u - X_{\tau_t - 1}}{a_t^{1/2}}\Big) \Big] dt.$$
(33)

Note also that G(z) is a symmetric integrable function of bounded variation. By Lemma 3.4, it is obvious that

$$\mathbf{E}\left[G\left(\frac{u-X_{\tau_t-1}}{a_t^{1/2}}\right)\right] \le c_6 \sup_{z} \left|\mathbf{P}\left(X_{\tau_t-1} \le z\right) - \Phi(z)\right| + c_7 \sqrt{a_t}.$$
(34)

Because of $c_* \geq 2, V_{\tau_t-1} = V_{\tau_t} - \bigtriangleup V_{\tau_t}, V_{\tau_t} \geq t$ and $\bigtriangleup V_{\tau_t} \leq 2\epsilon_n^2 + 2\delta_n^2$, we obtain

$$V_n - V_{\tau_t - 1} = V_n - V_{\tau_t} + \Delta V_{\tau_t} \le 2\epsilon_n^2 + 2\delta_n^2 + T - t \le a_t.$$
(35)

Therefore

$$\mathbf{E}\Big[\left(X_n - X_{\tau_t-1}\right)^2 \Big| \mathcal{F}_{\tau_t-1}\Big] = \mathbf{E}\Big[\sum_{k=\tau_t}^n \mathbf{E}\big[\xi_k^2 \big| \mathcal{F}_{k-1}\big] \Big| \mathcal{F}_{\tau_t-1}\Big]$$
$$= \mathbf{E}\big[\langle X \rangle_n - \langle X \rangle_{\tau_t-1} \big| \mathcal{F}_{\tau_t-1}\big]$$
$$\leq \mathbf{E}[V_n - V_{\tau_t-1} \big| \mathcal{F}_{\tau_t-1}\big]$$
$$\leq a_t.$$

Then, by Lemma 3.3, we deduce that for any $t \in [0, T]$,

$$\sup_{z} \left| \mathbf{P} \left(X_{\tau_t - 1} \le z \right) - \Phi(z) \right| \le c_8 \sup_{z} \left| \mathbf{P} \left(X_n \le z \right) - \Phi(z) \right| + c_9 \sqrt{a_t}.$$
(36)

Combining (28), (33), (34) and (36) together, we get

$$|J_1| \le c_{10}\epsilon_n^{1+\rho} \int_0^T \frac{1}{a_t^{(3+\rho)/2}} dt \sup_z \left| \mathbf{P} \left(X_n \le z \right) - \Phi(z) \right| + c_{11}\epsilon_n^{1+\rho} \int_0^T \frac{1}{a_t^{1+\rho/2}} dt.$$
(37)

Taking some elementary computations, it follows that

$$\int_{0}^{T} \frac{1}{a_t^{(3+\rho)/2}} dt = \int_{0}^{T} \frac{1}{(c_*^2 \gamma^2 + T - t)^{(3+\rho)/2}} dt \le \frac{2}{c_*^{1+\rho} (1+\rho) \gamma^{1+\rho}}$$
(38)

and

$$\int_{0}^{T} \frac{1}{a_t^{1+\rho/2}} dt = \int_{0}^{T} \frac{1}{(c_*^2 \gamma^2 + T - t)^{1+\rho/2}} dt \le \frac{2}{c_*^{\rho} \rho \gamma^{\rho}}.$$
(39)

This yields

$$\left|J_{1}\right| \leq \frac{c_{12}}{c_{*}^{1+\rho}} \sup_{z} \left|\mathbf{P}\left(X_{n} \leq z\right) - \Phi(z)\right| + \frac{c_{\rho,1}\epsilon_{n}}{\rho}.$$
(40)

Control of J_2 : Since $0 \leq \triangle V_k - \triangle \langle X \rangle_k \leq 2\delta^2 \mathbf{1}_{\{k=n\}}$, we have

$$|J_2| \leq \mathbf{E} \Big[\frac{1}{2A_n} \big| \varphi'(T_{n-1}) (\triangle V_n - \triangle \langle X \rangle_n) \big| \Big].$$

Denote $\widetilde{G}(z) = \sup_{|z-t| \leq 1} |\varphi'(t)|$, and then $|\varphi'(z)| \leq \widetilde{G}(z)$ for any real z. Since $A_n = c_*^2 \gamma^2$, then we get the following estimation :

$$|J_2| \le \frac{1}{c_*^2} \mathbf{E} \big[\widetilde{G}(T_{n-1}) \big].$$

Note that \widetilde{G} is non-increasing in $z \ge 0$, and thus it has bounded variation on **R**. By Lemma 3.4, we get

$$|J_2| \le \frac{c_{13}}{c_*^2} \sup_{z} \left| \mathbf{P} \left(X_{n-1} \le z \right) - \Phi(z) \right| + c_{*,2} (\epsilon_n + \delta_n).$$
(41)

Then, by Lemma 3.3, we deduce that

$$\sup_{z} \left| \mathbf{P} \left(X_{n-1} \le z \right) - \Phi(z) \right| \le c_{14} \sup_{z} \left| \mathbf{P} \left(X_n \le z \right) - \Phi(z) \right| + c_{15} \epsilon_n.$$

$$\tag{42}$$

This yields

$$|J_2| \le \frac{c_{16}}{c_*^2} \sup_{z} |\mathbf{P}(X_n \le z) - \Phi(z)| + c_{\rho,2}(\epsilon_n + \delta_n).$$
(43)

Control of J_3 . By a two-term Taylor expansion, it follows that

$$|J_3| = \frac{1}{8} \mathbf{E} \bigg[\sum_{k=1}^n \frac{1}{(A_k - \vartheta_k \bigtriangleup A_k)^2} \varphi^{\prime\prime\prime} \bigg(\frac{u - X_{k-1}}{\sqrt{A_k - \vartheta_k \bigtriangleup A_k}} \bigg) (\bigtriangleup A_k)^2 \bigg].$$

Note that $c_* \geq 2, \triangle A_k \leq 0$ and, by Lemma (3.2), $|\triangle A_k| = \triangle V_k \leq 2\epsilon_n^2 + 2\delta_n^2$. We obtain

$$A_k \le A_k - \vartheta_k \bigtriangleup A_k \le c_*^2 \gamma^2 + T - V_k + 2\epsilon_n^2 + 2\delta_n^2 \le 2A_k.$$

$$\tag{44}$$

Denote $\widehat{G}(z) = \sup_{|t-z| \leq 2} |\varphi'''(t)|$. Then $\widehat{G}(z)$ is symmetric, and is non-increasing in $z \geq 0$. Using (44), we get

$$|J_3| \le (2\epsilon_n^2 + 2\delta_n^2) \mathbf{E} \left[\sum_{k=1}^n \frac{1}{A_k^2} \widehat{G} \left(\frac{T_{k-1}}{\sqrt{2}} \right) \bigtriangleup V_k \right].$$

$$\tag{45}$$

By an argument similar to that of (40), we get

$$|J_{3}| \leq \frac{c_{17}(2\epsilon_{n}^{2} + 2\delta_{n}^{2})}{c_{*}^{2}\gamma^{2}} \sup_{z} |\mathbf{P}(X_{n} \leq z) - \Phi(z)| + \frac{2c_{18}(2\epsilon_{n}^{2} + 2\delta_{n}^{2})}{c_{*}\gamma}$$
$$\leq \frac{c_{19}}{c_{*}^{2}} \sup_{z} |\mathbf{P}(X_{n} \leq z) - \Phi(z)| + \frac{4c_{18}(\epsilon_{n} + \delta_{n})^{2}}{c_{*}\gamma}$$
$$\leq \frac{c_{19}}{c_{*}^{2}} \sup_{z} |\mathbf{P}(X_{n} \leq z) - \Phi(z)| + c_{\rho,3}(\epsilon_{n} + \delta_{n}).$$
(46)

Combining (17), (40), (43) and (46) together, we get

$$\left|\mathbf{E}\left[\Phi_u(X_n, A_n)\right] - \mathbf{E}\left[\Phi_u(X_0, A_0)\right]\right| \le \frac{c_{20}}{c_*^{1+\rho}} \sup_{z} \left|\mathbf{P}(X_n \le z) - \Phi(z)\right| + \frac{\hat{c}_{\rho}}{\rho}(\epsilon_n + \delta_n)$$

By (15), we know that

$$\sup_{z} \left| \mathbf{P} \left(X_n \le z \right) - \Phi(z) \right| \le \frac{c_{21}}{c_*^{1+\rho}} \sup_{z} \left| \mathbf{P} \left(X_n \le z \right) - \Phi(z) \right| + \frac{\tilde{c}_{\rho}}{\rho} (\epsilon_n + \delta_n),$$

from which, choosing $c_*^{1+\rho} = \max{\{2c_{21}, 2^{1+\rho}\}}$, we get

$$\sup_{z} \left| \mathbf{P} \left(X_n \le z \right) - \Phi(z) \right| \le \frac{2\tilde{c}_{\rho}(\epsilon_n + \delta_n)}{\rho}.$$
(47)

3.3. Proof of Theorem 1.2

Following the method of Bolthausen [2], we enlarge the sequence $(\xi_i, \mathcal{F}_i)_{1 \leq i \leq n}$ to $(\hat{\xi}_i, \hat{\mathcal{F}}_i)_{1 \leq i \leq N}$ such that $\langle \hat{X} \rangle_N := \sum_{i=1}^N \mathbf{E} [\hat{\xi}_i^2 | \hat{\mathcal{F}}_{i-1}] = 1$ a.s., and then apply Theorem 1.1 to the enlarged sequence. Consider the stopping time

$$\tau = \sup\{k \le n : \langle X \rangle_k \le 1\}.$$
(48)

Assume that $0 \leq \varepsilon \leq \epsilon_n$. Let $r = \left\lfloor \frac{1 - \langle X \rangle_\tau}{\varepsilon^2} \right\rfloor$, where $\lfloor x \rfloor$ denotes the "integer part" of x. It is easy to see that $r \leq \lfloor \frac{1}{\varepsilon^2} \rfloor$. Set N = n + r + 1. Let $(\zeta_i)_{i \geq 1}$ be a sequence of independent Rademacher random variables, which is independent of the martingale differences $(\xi_i)_{1 \leq i \leq n}$. Consider the random variables $(\hat{\xi}_i, \hat{\mathcal{F}}_i)_{1 \leq i \leq N}$ defined as follows :

$$\hat{\xi}_i = \begin{cases} \xi_i \quad \text{a.s.}, & \text{if } i \leq \tau, \\ \varepsilon \zeta_i \quad \text{a.s.}, & \text{if } \tau + 1 \leq i \leq \tau + r, \\ \left(1 - \langle X \rangle_\tau - r \varepsilon^2\right)^{1/2} \zeta_i \quad \text{a.s., if } i = \tau + r + 1, \\ 0 \quad \text{a.s.,} & \text{if } \tau + r + 1 \leq i \leq N, \end{cases}$$

and $\hat{\mathcal{F}}_i = \sigma(\hat{\xi}_1, \hat{\xi}_2, ..., \hat{\xi}_i).$

Clearly, $(\hat{\xi}_i, \hat{\mathcal{F}}_i)_{1 \le i \le N}$ still forms a martingale difference sequence with respect to the enlarged filtration. Then $\hat{X}_k = \sum_{i=1}^k \hat{\xi}_i$, k = 0, ..., N, with $\hat{X}_0 = 0$, is also a martingale. Moreover, it holds that $\langle \hat{X} \rangle_N = 1$, $\mathbf{E}[\hat{\xi}_i^3 | \hat{\mathcal{F}}_{i-1}] = 0$ and

$$\mathbf{E}[|\hat{\xi}_i|^{3+\rho}|\hat{\mathcal{F}}_{i-1}] \le \epsilon_n^{1+\rho} \mathbf{E}[\hat{\xi}_i^2|\hat{\mathcal{F}}_{i-1}], \quad a.s.$$

By Theorem 1.1, we have

$$D(\hat{X}_N) \le \frac{c_\rho \epsilon_n}{\rho}.$$
(49)

Using Lemma 3.5, we obtain that

$$D(X_n) \le 2D(\hat{X}_N) + 3 \left\| \mathbf{E} \left[\left| X_n - \hat{X}_N \right|^{2p} \right| \hat{X}_N \right] \right\|_1^{1/(2p+1)} \le \frac{2c_{\rho}\epsilon_n}{\rho} + 3 \left(\mathbf{E} \left[\left| \hat{X}_N - X_n \right|^{2p} \right] \right)^{1/(2p+1)}.$$
(50)

Since τ is a stopping time and

$$\hat{X}_N - X_n = \sum_{i=\tau+1}^N \left(\hat{\xi}_i - \xi_i\right), \quad \text{where put } \xi_i = 0 \text{ for } i > n,$$
(51)

 $(\hat{\xi}_i - \xi_i, \hat{\mathcal{F}}_i)_{i \ge \tau+1}$ still forms a martingale difference sequence. Applying Burkhold's inequality (cf. Theorem 2.11 of Hall and Heyde [8]), we get

$$\mathbf{E}\left[\left|\hat{X}_{N}-X_{n}\right|^{2p}\right] \leq \mathbf{E}\left[\max_{\tau+1\leq i\leq N}\left|\hat{X}_{i}-X_{i}\right|^{2p}\right] \\
\leq c_{p}\left(\mathbf{E}\left[\left|\sum_{i=\tau+1}^{N}\mathbf{E}\left[\left(\hat{\xi}_{i}-\xi_{i}\right)^{2}\left|\hat{\mathcal{F}}_{i-1}\right]\right|^{p}\right]+\mathbf{E}\left[\max_{\tau+1\leq i\leq N}\left|\hat{\xi}_{i}-\xi_{i}\right|^{2p}\right]\right).$$
(52)

As ξ_i and $\hat{\xi}_i$ be orthogonal random variables, we have

$$\sum_{i=\tau+1}^{N} \mathbf{E}\left[\left(\hat{\xi}_{i}-\xi_{i}\right)^{2}\middle|\hat{\mathcal{F}}_{i-1}\right] = \sum_{i=\tau+1}^{N} \mathbf{E}\left[\hat{\xi}_{i}^{2}\middle|\hat{\mathcal{F}}_{i-1}\right] + \sum_{i=\tau+1}^{n} \mathbf{E}\left[\xi_{i}^{2}\middle|\hat{\mathcal{F}}_{i-1}\right] = 1 - 2\langle X\rangle_{\tau} + \langle X\rangle_{n}.$$

Noting that $1 - \mathbf{E}[\xi_{\tau+1}^2 | \mathcal{F}_{\tau}] \leq \langle X \rangle_{\tau}$. Consequently, using the inequality $|a+b|^p \leq 2^{p-1} \left(|a|^p + |b|^p \right), p \geq 1$, and Jensen's inequality, we derive that

$$\left|\sum_{i=\tau+1}^{N} \mathbf{E}\left[\left(\hat{\xi}_{i}-\xi_{i}\right)^{2} \middle| \hat{\mathcal{F}}_{i-1}\right]\right|^{p} \leq \left|\langle X \rangle_{n}-1+2\mathbf{E}\left[\xi_{\tau+1}^{2} \middle| \mathcal{F}_{\tau}\right]\right|^{p} \\ \leq 2^{2p-1} \left(\left|\langle X \rangle_{n}-1\right|^{p}+\left|\mathbf{E}\left[\xi_{\tau+1}^{2} \middle| \mathcal{F}_{\tau}\right]\right|^{p}\right) \\ \leq 2^{2p-1} \left(\left|\langle X \rangle_{n}-1\right|^{p}+\mathbf{E}\left[\left|\xi_{\tau+1}\right|^{2p} \middle| \mathcal{F}_{\tau}\right]\right).$$
(53)

Taking expectations on both sides of the last inequality, we deduce that

$$\mathbf{E}\left[\left|\sum_{i=\tau+1}^{N} \mathbf{E}\left[\left(\hat{\xi}_{i}-\xi_{i}\right)^{2}\middle|\hat{\mathcal{F}}_{i-1}\right]\right|^{p}\right] \leq 2^{2p-1}\left(\mathbf{E}\left[\left|\langle X\rangle_{n}-1\right|^{p}\right]+\mathbf{E}\left[|\xi_{\tau+1}|^{2p}\right]\right)\right) \\ \leq 2^{2p-1}\left(\mathbf{E}\left[\left|\langle X\rangle_{n}-1\right|^{p}\right]+\mathbf{E}\left[\max_{1\leq i\leq n}|\xi_{i}|^{2p}\right]\right). \tag{54}$$

Similarly, using the inequality $|a+b|^p \leq 2^{p-1} \left(|a|^p + |b|^p \right), p \geq 1$,

$$\mathbf{E}\left[\max_{\tau+1\leq i\leq N} \left|\hat{\xi}_{i}-\xi_{i}\right|^{2p}\right] \leq 2^{2p-1} \mathbf{E}\left[\max_{\tau+1\leq i\leq N} \left(\left|\xi_{i}\right|^{2p}+\left|\hat{\xi}_{i}\right|^{2p}\right)\right] \\ \leq 2^{2p-1} \left(\mathbf{E}\left[\max_{1\leq i\leq n}\left|\xi_{i}\right|^{2p}\right]+\varepsilon^{2p}\right).$$
(55)

Combining (52), (54) and (55) together, we obtain

$$\mathbf{E}\left[\left|\hat{X}_{N}-X_{n}\right|^{2p}\right] \leq \hat{c}_{p}\left(\mathbf{E}\left[\left|\langle X\rangle_{n}-1\right|^{p}\right]+\mathbf{E}\left[\max_{1\leq i\leq n}|\xi_{i}|^{2p}\right]+\varepsilon^{2p}\right).$$
(56)

.

Finally, applying the last inequality to (50) and let $\varepsilon \to 0$, then we have

$$D(X_n) \le \tilde{c}_{\rho} \frac{\epsilon_n}{\rho} + \tilde{c}_p \left(\mathbf{E} \Big[|\langle X \rangle_n - 1|^p \Big] + \mathbf{E} \Big[\max_{1 \le i \le n} |\xi_i|^{2p} \Big] \right)^{1/(2p+1)}.$$

This completes the proof of Theorem 1.2.

Acknowledgements

The work has been partially supported by the National Natural Science Foundation of China (Grant no. 11601375). The research of Hailin Sang is partial supported by the Simons Foundation (Grant no. 586789).

References

- [1] A. Beknazaryan, H. Sang, Y. Xiao, Cramer type moderate deviations for random fields. J. Appl. Probab. 56 (2019) 223-245.
- [2] E. Bolthausen, Exact convergence rates in some martingale central limit theorems, Ann. Probab. 10 (1982) 672-688.

- [3] M. El Machkouri, L. Ouchti, Exact convergence rates in the central limit theorem for a class of martingales, Bernoulli 13(4) (2007) 981-999.
- X. Fan, Exact rates of convergence in some martingale central limit theorems, J. Math. Anal. Appl. 469 (2019) 1028-1044.
- [5] T. Fortune, M. Peligrad, H. Sang, A local limit theorem for linear random fields, (2020). submitted.
- [6] I. Grama, E. Haeusler, Large deviations for martingales via Cramér's method, Stochastic Process. Appl. 85 (2000) 279-293.
- [7] C.W. Granger, R. Joyeux, An introduction to long-memory time series models and fractional differencing, J. Time Ser. Anal. 1 (1980) 15-29.
- [8] P. Hall, C.C. Heyde, Martingale Limit Theory and Its Applications, Academic, New York, 1980.
- [9] J.R.M. Hosking, Fractional differencing, Biometrika 68 (1981) 165-176.
- [10] L.V. Kir'yanova, V.I. Rotar, Estimates for the rates of convergence in the central limit theorem for martingales, Theory Probab. Appl. 36 (1991) 289-302.
- [11] J.C. Mourrat, On the rate of convergence in the martingale central limit theorem, Bernoulli 19(2) (2013) 633-645.
- [12] L. Ouchti, On the rate of convergence in the central limit theorem for martingale difference sequences, Ann. Inst. Henri Poincaré Probab. Stat. 41(1) (2005) 35-43.
- [13] J. Renz, A note on exact convergence rates in some martingale central limit theorems, Ann. Probab. 24(3) (1996) 1616-1637.
- [14] W.B. Wu, W. Min, On linear processes with dependent innovations. Stochastic Process. Appl. 115 (2005) 939-958.