

Matching Book Embedding of the Cartesian Product of a Complete Graph and a Cycle

Zeling Shao, Yanqing Liu, Zhiguo Li*

School of Science, Hebei University of Technology, Tianjin 300401, China ^{*†}

Abstract

The *book embedding* of a graph G is to place the vertices of G on the spine and draw the edges to the pages so that the edges in the same page do not cross with each other. The book embedding is *matching* if the pages have maximum degree 1. The *matching book thickness* is the minimum number of pages in which graphs can be matching book embedded. In this paper, we show that the matching book thickness of the Cartesian product $K_p \square C_q$ of a complete graph K_p and a cycle C_q is equal to $\Delta(K_p \square C_q) + 1$.

Keywords: Book embedding; Matching book thickness; Cartesian product; Complete graph; Cycle

2000 MR Subject Classification. 05C10

1 Introduction

The concept of a book-embedding of a graph was introduced by Ollmann and Kainen. A *book* of the book embedding consists of a *spine* which is just a line and the *pages* each of which is a half-plane with spine as boundary. The *book thickness* $bt(G)$ is a measure of the quality of a book embedding which is the minimum number of pages in which G can be embedded. Book embedding has been found wide applications in the computer science, VLSI theory, multilayer printed circuit boards, sorting with parallel stacks and turning-machine graphs (see [1, 2]). The Cartesian product of two arbitrary graphs G and B is the graph denoted by $G \square B$ whose vertex set is $V(G) \times V(B)$, the vertex (u_1, v_1) and the vertex (u_2, v_2) are adjacent in $G \square B$ if and only if $u_1 = u_2$ and v_1 is adjacent to v_2 in B , or $v_1 = v_2$ and u_1 is adjacent to u_2 in G (see [3]). The book embedding of graphs has been discussed for a variety of graph families, for examples on complete graphs[4], complete bipartite graphs[5], generalized Petersen graph[6], some Cartesian product graphs[1, 7, 8], some semistrong product graphs[9] and Schrijver graphs[10], etc.

A book embedding of graph G is *matching* if every vertex must have degree at most one on every page. The *matching book thickness* of graph G , denoted by $mbt(G)$, is defined analogously

*Corresponding author. E-mail: zhiguolee@hebut.edu.cn

[†]This work is supported by the Natural Science Foundation of Hebei Province (No. A2019402043), China.

to the book thickness as the minimum number of pages required by any matching book embedding of graph G (see [11 – 13]). A graph G is *dispersable* if $mbt(G) = \Delta(G)$, where $\Delta(G)$ is the maximum degree index (see [4, 11, 13]). Complete bipartite graphs $K_{n,n}$ ($n \geq 1$), even cycles C_{2m} ($m \geq 2$), cubes Q_d ($d \geq 0$) and trees are dispersable [4]. Given an arbitrary graph $G = (V, E)$, let $\chi'(G)$ is the edge chromatic number, by definition, it is easy to know that $mbt(G) \geq \chi'(G) \geq \Delta(G)$. Kainen [14] showed that $mbt(C_p \square C_q)$ is 4, when p, q are both even and $mbt(C_p \square C_q)$ is 5, when p is even and q is odd. But for the case p, q are both odd, $mbt(C_p \square C_q)$ is still undetermined.

S. Overbay [11] showed that any k -regular dispersable graph G is bipartite. Also, she considered the matching book embedding of complete graphs K_n and got $mbt(K_n) = n$. Back in 1979, Bernhart and Kainen conjectured that any k -regular bipartite graph G is dispersable, i.e., $mbt(G) = k$. J.M. Alam, M.A. Bekos, M. Gronemann, M. Kaufmann, and S. Pupyrev [13] disprove this conjecture for the cases $k = 3$, and $k = 4$. In particular, they showed that the Gray graph, which is 3-regular and bipartite, has dispersable book thickness four, while the Folkman graph, which is 4-regular and bipartite, has dispersable book thickness five.

In this paper, we compute the matching book thickness of the Cartesian product of a complete graph K_p and a cycle C_q as follows.

Main Theorem: For $p, q \geq 3$, $mbt(K_p \square C_q) = \Delta(K_p \square C_q) + 1$

2 Proof of the main theorem

The proof will be completed by a sequence of lemmas.

Lemma 2.1. ^[11] *If a graph G is regular and dispersable, then G is bipartite.*

Lemma 2.2. *Let K_p be a complete graph and C_q be a cycle, $p, q \geq 3$, then $mbt(K_p \square C_q) \geq \Delta(K_p \square C_q) + 1$.*

Proof. A graph is not bipartite if and only if it contains an odd cycle. Since the Cartesian product of K_p and C_q contains at least one odd cycle, then $K_p \square C_q$ is not bipartite. Hence the graph $K_p \square C_q$ is not dispersable. Therefore $mbt(K_p \square C_q) \geq \Delta(K_p \square C_q) + 1$. \square

We will compute the matching book thickness of $K_p \square C_q$ from four cases.

Case 1: $K_{2n+1} \square C_{2m+1}$

Lemma 2.3. *If $n \geq 2, m \geq 1$, then $mbt(K_{2n+1} \square C_{2m+1}) = \Delta(K_{2n+1} \square C_{2m+1}) + 1 = 2n + 3$.*

Proof. Since $\Delta(K_{2n+1} \square C_{2m+1}) = 2n + 2$, by Lemma 2.2, we have $mbt(K_{2n+1} \square C_{2m+1}) \geq 2n + 3$.

There are $(2n+1)(2m+1)$ vertices for graph $K_{2n+1} \square C_{2m+1}$, and these vertices can be separated into $2n + 1$ rows and $2m + 1$ columns. Each columns holding a consecutive number from top to

bottom. Let the ordering on spine be as $u_1v_{2n+1}, u_1v_{2n}, u_1v_{2n-1}, \dots, u_1v_1, u_2v_1, u_2v_2, \dots, u_2v_{2n+1}, \dots, u_{2m+1}v_{2n+1}, \dots, u_{2m+1}v_1$.

Page 0: the edge $\{(u_1v_1), (u_{2m+1}v_1)|m \geq 1\}$, and edges $\{(u_iv_a)(u_iv_b)|i = 1 \text{ or } 2m+1; a+b = 2n+3; 1 < a, b \leq 2n+1\}$.

Page 1: the edge $\{(u_1v_2), (u_{2m+1}v_2)|m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b)|i = 1 \text{ or } 2m+1; a+b = 2n+4; 2 < a, b \leq 2n+1\}$.

Page 2: the edge $\{(u_1v_3), (u_{2m+1}v_3)|m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b); (u_iv_1), (u_iv_2)|i = 1 \text{ or } 2m+1; a+b = 2n+5; 3 < a, b \leq 2n+1\}$.

.....

Page $n-1$: the edge $\{(u_1v_n), (u_{2m+1}v_n)|n \geq 2, m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b); (u_iv_c), (u_iv_d)|i = 1 \text{ or } 2m+1; a+b = 3n+2; c+d = n; n < a, b \leq 2n+1; 1 \leq c, d < n\}$.

Page n : the edge $\{(u_1v_{n+1}), (u_{2m+1}v_{n+1})|n \geq 2, m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b); (u_iv_c), (u_iv_d)|i = 1 \text{ or } 2m+1; a+b = 3n+3; c+d = n+1; n+1 < a, b \leq 2n+1; 1 \leq c, d < n+1\}$.

.....

Page $2n-2$: the edge $\{(u_1v_{2n-1}), (u_{2m+1}v_{2n-1})|n \geq 2, m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b); (u_iv_{2n}), (u_iv_{2n+1})|i = 1 \text{ or } 2m+1; a+b = 2n-1; 1 \leq a, b < 2n-1\}$.

Page $2n-1$: the edge $\{(u_1v_{2n}), (u_{2m+1}v_{2n})|n \geq 2, m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b)|i = 1 \text{ or } 2m+1; a+b = 2n; 1 \leq a, b < 2n\}$.

Page $2n$: the edge $\{(u_1v_{2n+1}), (u_{2m+1}v_{2n+1})|n \geq 2, m \geq 1\}$, and edges $\{(u_iv_a), (u_iv_b)|i = 1 \text{ or } 2m+1; a+b = 2n+1; 1 \leq a, b < 2n+1\}$.

Page $2n+1$: edges $\{(u_iv_j), (u_{i+1}v_j)|1 \leq i < 2m+1; j = 1, 2, 3, \dots, 2n+1, i \text{ is odd}\}$, and edges $\{(u_{2m+1}v_a), (u_{2m+1}v_b)|a+b = 2n+2; 1 \leq a, b \leq 2n+1\}$.

Page $2n+2$: edges $\{(u_iv_j), (u_{i+1}v_j)|2 \leq i < 2m+1; j = 1, 2, 3, \dots, 2n+1, i \text{ is even}\}$, and edges $\{(u_1v_a), (u_1v_b)|a+b = 2n+2; 1 \leq a, b \leq 2n+1\}$.

For $k = 0, 1, 2, 3, \dots, 2n$, other edges $\{(u_iv_a), (u_iv_b)|i = 2, 3, \dots, 2m; 1 \leq a, b \leq 2n+1\}$ of $K_{2n+1} \square C_{2m+1}$ are placed on page k , where $k \equiv a+b \pmod{2n+1}$.

So $K_{2n+1} \square C_{2m+1}$ can be matching book embedded in $2n+3$ pages. Therefore the matching book thickness of $K_{2n+1} \square C_{2m+1}$ is $2n+3$ (see Fig.1 for the case $n=2, m=1$). \square




Fig.1 The matching book embedding of $K_5 \square C_3$.

Case 2: $K_{2n} \square C_{2m+1}$

Lemma 2.4. *If $n \geq 2, m \geq 1$, then $mbt(K_{2n} \square C_{2m+1}) = \Delta(K_{2n} \square C_{2m+1}) + 1 = 2n + 2$.*

Proof. Since $\Delta(K_{2n} \square C_{2m+1}) = 2n + 1$, by Lemma 2.2, $mbt(K_{2n} \square C_{2m+1}) \geq 2n + 2$. Vertices of graph $K_{2n} \square C_{2m+1}$ can be separated into $2n$ rows and $2m + 1$ columns. The edges of graph $K_{2n} \square C_{2m+1}$ can be put on $2n + 2$ pages analogously to that of $K_{2n+1} \square C_{2m+1}$ in Lemma 2.3. Hence we have $mbt(K_{2n} \square C_{2m+1}) \leq 2n + 2$. Therefore the result is established (see Fig.2 for the case $n = 3, m = 1$).

□




Fig.2 The matching book embedding of $K_6 \square C_3$.

The following result will be applied to case 3 and case 4, which is a generalization of Theorem 4.3 of [4]:

Lemma 2.5. *Let G be arbitrary and B be a dispersable bipartite graph, then $mbt(G \square B) \leq mbt(G) + \Delta(B)$.*

Proof. Since B is dispersable, then there is a $\Delta(B)$ -edge coloring and a corresponding matching book embedding in a $\Delta(B)$ -page book so that all edges of one color lie in the same page. Since B is bipartite, there is a 2-vertex coloring using colors a and b .

Now we consider the matching book embedding of $G \square B$. Take a matching book embedding of G in $mbt(G)$ pages. Using a matching book embedding of B , replace all vertices which colored a with a copy of this matching book embedding of G and replace all vertices which colored b with the same matching book embedding of G , but in reverse order.

Now each of vertex of B is represent by a copy of G . Since each of these copies are placed on the spine, then the edges of $G \square B$ corresponding to the copies of G can all be matching embedded in $mbt(G)$ pages.

The remaining edges of $G \square B$ connect corresponding vertices in adjacent copies of G . Since B is bipartite, then edges of B connect vertices of different colors. Since the order of vertices in adjacent copies of G is reversed, then edges connect two adjacent copies corresponding to a single

edge of B can be embedded in the appropriate page as concentric semicircles, which makes sure that the book embedding is matching. Since $mbt(B) = \Delta(B)$, the copies of the edges of B can also be embedded in $\Delta(B)$ pages.

Hence all edges of $G \square B$ can be matching book embedded in $mbt(G) + \Delta(B)$ pages. \square

For example, let $G = K_5 - e$, $B = P_3$, the matching book embedding of G and B is shown in Fig.3. The resulting matching book embedding of graph $G \square B$ in 7 pages is shown in Fig.4.




Fig.3 The matching book embedding of G (left) and B (right)




Fig.4 The matching book embedding of $G \square B$

Case 3 and Case 4: $K_{2n} \square C_{2m}, K_{2n+1} \square C_{2m}$

Lemma 2.6. *If $n, m \geq 2$, then*

- (i) $mbt(K_{2n} \square C_{2m}) = \Delta(K_{2n} \square C_{2m}) + 1 = 2n + 2$;
- (ii) $mbt(K_{2n+1} \square C_{2m}) = \Delta(K_{2n+1} \square C_{2m}) + 1 = 2n + 3$.

Proof. Since $K_{2n} \square C_{2m}$ is $2n + 1$ -regular and $K_{2n+1} \square C_{2m}$ is $2n + 2$ -regular, by Lemma 2.2, it can be obtained that $mbt(K_{2n} \square C_{2m}) \geq 2n + 2$, $mbt(K_{2n+1} \square C_{2m}) \geq 2n + 3$.

While the even cycle C_{2m} is dispersable, by Lemma 2.5, for complete graph K_{2n} and K_{2n+1} , we have

$$mbt(K_{2n} \square C_{2m}) \leq mbt(K_{2n}) + 2 = 2n + 2;$$

$$mbt(K_{2n+1} \square C_{2m}) \leq mbt(K_{2n+1}) + 2 = 2n + 3.$$

This gives the upper bound of $K_{2n} \square C_{2m}$ and $K_{2n+1} \square C_{2m}$. Hence the equalities of (i) and (ii) both hold. \square

References

- [1] X. L. Li. Book Embedding of Graphs. Ph. D. Dissertation, ZhengZhou: ZhengZhou University, 2002.
- [2] F. R. K. Chung, F. T. Leighton, A. L. Rosenberg. Embedding Graphs in Books: A Layout Problem with Applications to VLSI Design. Society for Industrial and Applied Mathematics, 1987, 8(1): 33-58.
- [3] J. A. Bondy, U. S. R. Murty. Graph Theory. London: Springer, 2008.
- [4] F. Bernhart, P. C. Kainen. The Book Thickness of a Graph. Journal of Combinatorial Theory, Series B, 1979, 27: 320-331.
- [5] H. Enomoto, T. Nakamigawa, and K. Ota. On the Pagenumber of Complete Bipartite Graphs. Journal of Combinatorial Theory, Series B, 1997, 71: 111-120.
- [6] B. Zhao, W. Xiong, Y. Tian. Embedding Generalized Petersen Graph in Books, Chin. Ann. Math. Ser. B, 2016, 37: 385-394.
- [7] J. Yang, Z. L. Shao, Z. G. Li. Embedding Cartesian Product of Some Graphs in Books. Communications in Mathematical Research, 2018, 34(03): 253-260.
- [8] Z. L. Shao, C. J. Ren, Z. G. Li. Book Embedding and Lovasz Number, submitted for publication.
- [9] B. Zhao, Y. Tian, J. Meng. Embedding Semistrong Product of Paths and Cycles in Books. Journal of Natural Science of Hunan Normal University, 2015, 38(6): 73-77.
- [10] Z. L. Shao, X. L. Hao, Z. G. Li. The Book Embedding of Schrijver Graph. Utilitas Mathematica, 2018, 107: 157-165.
- [11] S. Overbay. Generalized Book Embeddings. Ph. D. Dissertation, Fort Collins: Colorado State University, 1998.
- [12] P. C. Kainen. On Book Embeddings with Degree-1 Pages, submitted for publication.
- [13] J. M. Alam, M. A. Bekos, M. Gronemann, M. Kaufmann, and S. Pupyrev. On Dispersable Book Embeddings. International Workshop on Graph-Theoretic Concepts in Computer Science, 2018, 11159: 1-14.
- [14] P. C. Kainen. Complexity of Products of Even Cycles. Bulletin of the Institute of Combinatorics and Its Applications, 2011, 62: 95-102.