

ON THE MODULI SPACE OF λ -CONNECTIONS

ANOOP SINGH

ABSTRACT. Let X be a compact Riemann surface of genus $g \geq 3$. Let \mathcal{M}_{Hod} denote the moduli space of stable λ -connections over X and $\mathcal{M}'_{Hod} \subset \mathcal{M}_{Hod}$ denote the subvariety whose underlying vector bundle is stable. Fix a line bundle L of degree zero. Let $\mathcal{M}_{Hod}(L)$ denote the moduli space of stable λ -connections with fixed determinant L and $\mathcal{M}'_{Hod}(L) \subset \mathcal{M}_{Hod}(L)$ be the subvariety whose underlying vector bundle is stable. We show that there is a natural compactification of \mathcal{M}'_{Hod} and $\mathcal{M}'_{Hod}(L)$, and study their Picard groups. Let $\mathbb{M}_{Hod}(L)$ denote the moduli space of polystable λ -connections. We investigate the nature of algebraic functions on $\mathcal{M}_{Hod}(L)$ and $\mathbb{M}_{Hod}(L)$. We also study the automorphism group of $\mathcal{M}'_{Hod}(L)$.

1. INTRODUCTION AND STATEMENTS OF THE RESULTS

1.1. Notations and conventions

In what follows, \mathbb{A}^1 denote the affine line over the field of complex numbers \mathbb{C} , X denote the compact connected Riemann surface of genus $g \geq 3$. Unless otherwise stated genus g of X is considered to be ≥ 3 . $J(X)$ denote the Jacobian of X .

1.2. Motivation and results

The Picard group of a moduli space is a very important invariant while studying the classification problems for algebro-geometric objects. The Picard group of moduli space of vector bundles have been studied extensively by several algebraic geometers, for instance [7], [11], [2] to name a few. Also, the Picard group and algebraic functions for the moduli space of logarithmic connections have been studied in [5] and [12]. So, the purpose of this article is to study the Picard group, algebraic function and automorphism group of the moduli spaces of λ -connections. The notion of λ -connection in a vector bundle was introduced by P. Deligne [6] realising the Higgs bundles as a degeneration of vector bundles with connections, where λ varies over the field of complex numbers \mathbb{C} . In the series of papers [16], [15] and [17], Simpson studied the moduli space of λ -connections extensively, where he named the moduli space of λ -connections as Hodge moduli space.

In section (2), we recall the notion of λ -connection in a holomorphic vector bundle E over X of rank n . We also recall the notion of stable and polystable λ -connections. Let $\mathcal{M}_{Hod} = \mathcal{M}_{Hod}(X, n)$ denote the moduli space of stable λ -connections on X of rank n . Let $\mathcal{M}'_{Hod} \subset \mathcal{M}_{Hod}$ be the moduli space whose underlying vector bundle is

2010 *Mathematics Subject Classification.* 14D20, 14C22, 14E05, 14J50.

Key words and phrases. λ -connection, Moduli space, Picard group, algebraic function, automorphism.

stable. We show that there is a natural compactification of the moduli space \mathcal{M}'_{Hod} [see Proposition (3.1)].

Let $\mathcal{U} = \mathcal{U}(X, n)$ denote the moduli space of stable vector bundle of rank n and degree 0, and let

$$p : \mathcal{M}'_{Hod} \rightarrow \mathcal{U} \quad (1.1)$$

be the natural projection sending $(E, \lambda, D) \mapsto E$. The morphism $p : \mathcal{M}'_{Hod} \rightarrow \mathcal{U}$ defined in (1.1) induces a homomorphism

$$p^* : \text{Pic}(\mathcal{U}) \rightarrow \text{Pic}(\mathcal{M}'_{Hod}) \quad (1.2)$$

of Picard groups that sends any line bundle ξ over \mathcal{U} to a line bundle $p^*\xi$ over \mathcal{M}'_{Hod} . Next, let $\iota : \mathcal{M}'_{Hod} \hookrightarrow \mathcal{M}_{Hod}$ be the inclusion morphism. Then ι induces a homomorphism

$$\iota^* : \text{Pic}(\mathcal{M}_{Hod}) \rightarrow \text{Pic}(\mathcal{M}'_{Hod}) \quad (1.3)$$

of Picard groups which restricts any line bundle ξ over \mathcal{M}_{Hod} to \mathcal{M}'_{Hod} . We show that p^* and ι^* are isomorphisms of groups, more concretely [see Theorem (3.3)]

Theorem 1.1. *Let X be a compact Riemann surface of genus $g \geq 3$. Then, the two homomorphism*

$$\text{Pic}(\mathcal{U}) \xrightarrow{p^*} \text{Pic}(\mathcal{M}'_{Hod}) \xleftarrow{\iota^*} \text{Pic}(\mathcal{M}_{Hod})$$

defined in (1.2) and (1.3) are isomorphisms.

Let $\mathbb{M}_{Hod} = \mathbb{M}_{Hod}(X, n)$ (see [9]) denote the coarse moduli space of isomorphism classes of polystable λ -connections on X of rank n . Then we show that $\text{Pic}(\mathbb{M}_{Hod}) \cong \text{Pic}(\mathcal{M}_{Hod})$ [see Proposition (3.4)].

Fix a line bundle L over X of degree zero. For each $\lambda \in \mathbb{C}$, fix a λ -connection D_L^λ on L such that for any $\lambda_1, \lambda_2 \in \mathbb{C}$, we have

$$D_L^{\lambda_1} + D_L^{\lambda_2} = D_L^{\lambda_1 + \lambda_2}.$$

Let $\mathcal{M}_{Hod}(L) = \mathcal{M}_{Hod}(X, n, L)$ denote the moduli space parametrising all stable λ -connection (E, λ, D) such that $\bigwedge^n E \cong L$ and for every $\lambda \in \mathbb{C}$, the λ -connection on $\bigwedge^n E$ induced by D coincides with the given λ -connection D_L^λ on L . Let $\mathcal{M}'_{Hod}(L) \subset \mathcal{M}_{Hod}(L)$ be the moduli spaces whose underlying vector bundle is stable. We show that $\mathcal{M}'_{Hod}(L)$ admits a natural compactification [see Proposition (4.1)] and prove that $\text{Pic}(\mathcal{M}'_{Hod}(L)) \cong \mathbb{Z}$ [see Corollary (4.3)]. Now, we have a natural surjective algebraic morphism

$$\Phi_L : \mathcal{M}_{Hod}(L) \rightarrow \mathbb{A}^1 \quad (1.4)$$

defined by $(E, \lambda, D) \mapsto \lambda$. Then, we have [see Theorem (4.9)]

Theorem 1.2. *Any algebraic function on $\mathcal{M}_{Hod}(L)$ will factor through the map*

$$\Phi_L : \mathcal{M}_{Hod}(L) \rightarrow \mathbb{A}^1$$

defined in (1.4).

Finally, in the last section(5), we prove that the group $\text{Aut}(\mathcal{M}'_{Hod}(L))_0$ of algebraic automorphisms of the moduli space $\mathcal{M}'_{Hod}(L)$ fits into the following short exact sequence

$$0 \rightarrow \mathbb{V} \xrightarrow{\iota} \text{Aut}(\mathcal{M}'_{Hod}(L))_0 \xrightarrow{\rho} \mathbb{W} \times \mathbb{C}^* \rightarrow 0$$

of groups [see Theorem (5.1)].

2. PRELIMINARIES

Definition 2.1. [15] *A λ -connection on X is a triple of the form (E, λ, D) , where $\lambda \in \mathbb{C}$, E is a holomorphic vector bundle on X of rank n such that $\det(E) = \Lambda^n E \in J(X)$ and*

$$D : E \rightarrow \Omega_X^1 \otimes E$$

is a \mathbb{C} -linear map which satisfies (Leibniz property)

$$D(fs) = fD(s) + \lambda df \otimes s \quad (2.1)$$

for every local holomorphic section s of E and every local section f of \mathcal{O}_X .

If (E, λ, D) is a λ -connection on X with $\lambda \neq 0$, then D/λ is a holomorphic connection on E . Therefore by Atiyah [1] and Weil [19], $\text{degree}(E) = 0$. On the other hand, if $\lambda = 0$, then D is an \mathcal{O}_X -linear map, and hence $D \in H^0(X, \Omega_X^1 \otimes \text{End}(E))$. Thus, the triple $(E, 0, D)$ is a Higgs bundle on X with $\text{degree}(E) = 0$ (by Definition (2.1)).

Given two λ -connections (E, λ, D) and (E, λ, D') on X , where E and λ are fixed, we have $D - D' \in H^0(X, \Omega_X^1 \otimes \text{End}(E))$. Conversely, given $\omega \in H^0(X, \Omega_X^1 \otimes \text{End}(E))$, the triple $(E, \lambda, D + \omega)$ is again a λ -connection on X .

Now, for fixed E and fixed $\lambda \in \mathbb{C}$, let $V_\lambda(E)$ denote the set consists of λ -connections (E, λ, D) . Then $V_\lambda(E)$ is an affine space modelled on $H^0(X, \Omega_X^1 \otimes \text{End}(E))$ if $\lambda \neq 0$, and $V_0(E) = H^0(X, \Omega_X^1 \otimes \text{End}(E))$.

Next, fix E . Let (E, λ, D) and (E, λ', D') be two λ -connections on X . Then $(E, \lambda + \lambda', D + D')$ is a λ -connection on X . Moreover, for any $\beta \in \mathbb{C}$, the triple $(E, \beta\lambda, \beta D)$ is a λ -connection on X . The set of all λ -connections in E , denoted by $V(E)$, is a vector space over \mathbb{C} . We have

$$V(E) = \coprod_{\lambda \in \mathbb{C}} V_\lambda(E). \quad (2.2)$$

If E is a stable vector bundle over X , then using Riemann-Roch theorem and Serre duality, the dimension of $V_0(E) = H^0(X, \Omega_X^1 \otimes \text{End}(E))$ is $n^2(g-1) + 1$.

Definition 2.2. *A λ -connection (E, λ, D) over X is said to be semistable (respectively, stable), if for every proper subbundle $F \neq 0$ with $\text{rk}(F) > 0$, invariant under D , that is, $D(F) \subset F \otimes \Omega_X^1$, we have $\text{degree}(F) \leq 0$ (respectively, $\text{degree}(F) < 0$).*

A λ -connection is polystable if it decomposes as a direct sum of stable λ -connections with the same slope.

Let $\mathcal{M}_{\text{Hod}} = \mathcal{M}_{\text{Hod}}(X, n)$ denote the moduli space of stable λ -connections of rank n on X . Then \mathcal{M}_{Hod} is an irreducible smooth quasi-projective variety of dimension $2n^2(g-1) + 2$ (see [16], [14]). We have natural surjective algebraic morphism

$$\Phi : \mathcal{M}_{\text{Hod}} \rightarrow \mathbb{A}^1 \quad (2.3)$$

defined by $(E, \lambda, D) \mapsto \lambda$. Let Φ^λ denote the inverse image $\Phi^{-1}(\lambda)$. Then $\Phi^1 = \mathcal{M}_{\text{DR}}(X, n)$ and $\Phi^0 = \mathcal{M}_{\text{Dol}}(X, n)$ are de Rham moduli space and Dolbeault moduli space, respectively. Let $\mathcal{M}'_{\text{Hod}} \subset \mathcal{M}_{\text{Hod}}$ be the moduli space whose underlying vector

bundle is stable. Then \mathcal{M}'_{Hod} is a Zariski open subvariety of \mathcal{M}_{Hod} follows from [[10], p.635, Theorem 2.8(B)].

3. THE PICARD GROUP OF THE HODGE MODULI SPACE

In this section, we compute the Picard group of the Hodge moduli spaces. Let $\mathcal{U} = \mathcal{U}(X, n)$ denote the moduli space of stable vector bundle of rank n and degree 0, and let

$$p : \mathcal{M}'_{Hod} \rightarrow \mathcal{U} \quad (3.1)$$

be the natural projection sending $(E, \lambda, D) \rightarrow E$. Then p is a surjective algebraic morphism. For $E \in \mathcal{U}$, $p^{-1}(E) = V(E)$, where $V(E)$ is the vector space of all λ -connections in E that can be expressed as disjoint union of $V_\lambda(E)$ for $\lambda \in \mathbb{C}$ described in (2.2). Now, since E is a stable vector bundle, we have

$$\dim_{\mathbb{C}}(V(E)) = n^2(g-1) + 2. \quad (3.2)$$

There is a natural algebraic action of \mathbb{C}^* on \mathcal{M}'_{Hod} . In [18], using this natural \mathbb{C}^* action on \mathcal{M}'_{Hod} , Simpson has proved a relative compactification of \mathcal{M}'_{Hod} over \mathbb{A}^1 . Nevertheless, we give a different proof for the compactification of \mathcal{M}'_{Hod} , which is constructive in nature. The statement and the steps involved in the following Proposition (3.1) is exactly similar to the statement and proof of the Theorem 4.3 in [12]. The main difference is that the fibre of p defined in (3.1) is a finite dimensional vector space, while in [12], the fibre of p [see [12] equation(4.1)] is an affine space modelled over $H^0(X, \Omega_X^1 \otimes \text{End}(E))$. Nonetheless, we shall sketch the construction of algebraic vector bundle \mathcal{Q} over \mathcal{U} , because the morphisms (3.4) and (3.5) in proof of the following Proposition 3.1 are very crucial in proving Theorem (3.3).

Proposition 3.1. *There exists an algebraic vector bundle $\pi : \mathcal{Q} \rightarrow \mathcal{U}$ such that \mathcal{M}'_{Hod} is embedded in $\mathbf{P}(\mathcal{Q})$ with $\mathbf{P}(\mathcal{Q}) \setminus \mathcal{M}'_{Hod}$ as the hyperplane at infinity.*

Proof. By dualizing the fibres of p defined in (3.1), we construct a new vector bundle over \mathcal{U} satisfying certain conditions. For any $E \in \mathcal{U}$, the fiber $p^{-1}(E) = V(E)$ is a finite dimensional vector space over \mathbb{C} . Let $\pi : \mathcal{Q} \rightarrow \mathcal{U}$ be an algebraic vector bundle defined as follows; for every Zariski open subset U of \mathcal{U} ,

$$\mathcal{Q}(U) = \{f : p^{-1}(U) \rightarrow \mathbb{A}^1 \text{ is a regular function} : f|_{V(E)} \in V(E)^\vee\} \quad (3.3)$$

Let $(E, \lambda, D) \in \mathcal{M}'_{Hod}$. Then define a map $\eta_{(E, \lambda, D)} : V(E)^\vee \rightarrow \mathbb{C}$, by $\eta_{(E, \lambda, D)}(\varphi) = \varphi[(E, \lambda, D)]$, which is nothing but the evaluation map. Now, the kernel $\text{Ker}(\eta_{(E, \lambda, D)})$ defines a hyperplane in $V(E)^\vee$ is denoted by $H_{(E, \lambda, D)}$.

Let $\mathbf{P}(\mathcal{Q})$ be the projective bundle defined by hyperplanes in the fiber of π .

Define a map

$$\iota : \mathcal{M}'_{Hod} \rightarrow \mathbf{P}(\mathcal{Q}) \quad (3.4)$$

by sending (E, λ, D) to the hyperplane $H_{(E, \lambda, D)}$, which is clearly an open embedding.

Set $Y = \mathbf{P}(\mathcal{Q}) \setminus \mathcal{M}'_{Hod}$. Let

$$\tilde{\pi} : \mathbf{P}(\mathcal{Q}) \rightarrow \mathcal{U} \quad (3.5)$$

be the natural projection induced from π . Then $\tilde{\pi}^{-1}(E) \cap Y$ is a linear hyperplane in the projective space $\tilde{\pi}^{-1}(E)$ for every $E \in \mathcal{U}$. \square

Let Z be a quasi-projective variety over \mathbb{C} . We recall an elementary fact about Picard groups.

Lemma 3.2. *Let $\iota : U \hookrightarrow Z$ be a Zariski open subset of Z , whose complement has codimension at least 2 in Z . Then the induced homomorphism $\iota^* : \text{Pic}(Z) \rightarrow \text{Pic}(U)$ of Picard groups is an isomorphism.*

Proof. See [8], Chapter II, Proposition 6.5 (b), p.n. 133. \square

The morphism $p : \mathcal{M}'_{Hod} \rightarrow \mathcal{U}$ defined in (3.1) induces a homomorphism

$$p^* : \text{Pic}(\mathcal{U}) \rightarrow \text{Pic}(\mathcal{M}'_{Hod}) \quad (3.6)$$

of Picard groups that sends any line bundle ξ over \mathcal{U} to a line bundle $p^*\xi$ over \mathcal{M}'_{Hod} . Let

$$\iota : \mathcal{M}'_{Hod} \hookrightarrow \mathcal{M}_{Hod} \quad (3.7)$$

be the inclusion map. Then ι induces a homomorphism

$$\iota^* : \text{Pic}(\mathcal{M}_{Hod}) \rightarrow \text{Pic}(\mathcal{M}'_{Hod}) \quad (3.8)$$

of Picard groups which restricts any line bundle ξ over \mathcal{M}_{Hod} to \mathcal{M}'_{Hod} .

Now, we state the following theorem whose first part (that is, $p^* : \text{Pic}(\mathcal{U}) \rightarrow \text{Pic}(\mathcal{M}'_{Hod})$ is an isomorphism) is similar to Theorem 4.4 in [12], but in different context, that is for moduli space of logarithmic connections.

Theorem 3.3. *Let X be a compact Riemann surface of genus $g \geq 3$. Then, the two homomorphism*

$$\text{Pic}(\mathcal{U}) \xrightarrow{p^*} \text{Pic}(\mathcal{M}'_{Hod}) \xleftarrow{\iota^*} \text{Pic}(\mathcal{M}_{Hod}) \quad (3.9)$$

defined in (3.6) and (3.8) are isomorphisms.

Proof. First we show that ι^* is an isomorphism. In view of Lemma (3.2), it is sufficient to show that the complement

$$\mathcal{M}_{Hod} \setminus \mathcal{M}'_{Hod} \subset \mathcal{M}_{Hod}$$

is of codimension at least 2 in \mathcal{M}_{Hod} . Note that the dimension of the moduli space \mathcal{M}_{Hod} is $2n^2(g-1) + 2$. From the first statement in [2][Proposition 1.2, 1], we conclude that the complement $\mathcal{M}_{Hod} \setminus \mathcal{M}'_{Hod}$ has codimension at least 2 in \mathcal{M}_{Hod} .

Next to show that p^* is an isomorphism. we first show that p^* is injective. Let $\xi \rightarrow \mathcal{U}$ be a line bundle such that $p^*\xi$ is a trivial line bundle over \mathcal{M}'_{Hod} . Giving a trivialization of $p^*\xi$ is equivalent to giving a nowhere vanishing section of $p^*\xi$ over \mathcal{M}'_{Hod} . Fix $s \in H^0(\mathcal{M}'_{Hod}, p^*\xi)$ a nowhere vanishing section. Take any point $E \in \mathcal{U}$. Then, from the following commutative diagram

$$\begin{array}{ccc} p^*\xi & \xrightarrow{\tilde{p}} & \xi \\ \downarrow & & \downarrow \\ \mathcal{M}'_{Hod} & \xrightarrow{p} & \mathcal{U} \end{array} \quad (3.10)$$

we get

$$s|_{p^{-1}(E)} : p^{-1}(E) \rightarrow \xi(E)$$

a nowhere vanishing map. Notice that $p^{-1}(E) \cong \mathbb{C}^N$ and $\xi(E) \cong \mathbb{C}$, where $N = n^2(g-1) + 2$. Now, any nowhere vanishing algebraic function on an affine space \mathbb{C}^N is a constant function, that is, $s|_{p^{-1}(E)}$ is a constant function and hence corresponds to a non-zero vector $\alpha_E \in \xi(E)$. Since s is constant on each fiber of p , the trivialization s of $p^*\xi$ descends to a trivialization of the line bundle ξ over \mathcal{U} , and hence giving a nowhere vanishing section of ξ over \mathcal{U} . Thus, ξ is a trivial line bundle over \mathcal{U} .

It remains to show that p^* is surjective. Let $\vartheta \rightarrow \mathcal{M}'_{Hod}$ be an algebraic line bundle. Since $\mathcal{M}'_{Hod} \hookrightarrow \mathbf{P}(\mathcal{Q})$ [follows from (3.4), in the proof of above Proposition (3.1)], we can extend ϑ to a line bundle ϑ' over $\mathbf{P}(\mathcal{Q})$. Again from the morphism $\tilde{\pi} : \mathbf{P}(\mathcal{Q}) \rightarrow \mathcal{U}$ in (3.5) in the above Theorem(3.1) and from [8], Chapter III, Exercise 12.5, p.n. 291., we have

$$\text{Pic}(\mathbf{P}(\mathcal{Q})) \cong \tilde{\pi}^*\text{Pic}(\mathcal{U}) \oplus \mathbb{Z}\mathcal{O}_{\mathbf{P}(\mathcal{Q})}(1). \quad (3.11)$$

Therefore,

$$\vartheta' = \tilde{\pi}^*L \otimes \mathcal{O}_{\mathbf{P}(\mathcal{Q})}(l) \quad (3.12)$$

where L is a line bundle over \mathcal{U} and $l \in \mathbb{Z}$. Since $Y = \mathbf{P}(\mathcal{Q}) \setminus \mathcal{M}'_{Hod}$ is the hyperplane at infinity, using (3.11) the line bundle $\mathcal{O}_{\mathbf{P}(\mathcal{Q})}(Y)$ associated to the divisor Y can be expressed as

$$\mathcal{O}_{\mathbf{P}(\mathcal{Q})}(Y) = \tilde{\pi}^*L_1 \otimes \mathcal{O}_{\mathbf{P}(\mathcal{Q})}(1) \quad (3.13)$$

for some line bundle L_1 over \mathcal{U} . Now, from (3.12) and (3.13), we get

$$\vartheta' = \tilde{\pi}^*(L \otimes (L_1^\vee)^{\otimes l}) \otimes \mathcal{O}_{\mathbf{P}(\mathcal{Q})}(lY). \quad (3.14)$$

Since, the restriction of the line bundle $\mathcal{O}_{\mathbf{P}(\mathcal{Q})}(Y)$ to the compliment $\mathbf{P}(\mathcal{Q}) \setminus Y = \mathcal{M}'_{Hod}$ is the trivial line bundle and restriction of $\tilde{\pi}$ to \mathcal{M}'_{Hod} is the map p defined in (3.1), therefore, we have

$$\vartheta = p^*(L \otimes (L_1^\vee)^{\otimes l}). \quad (3.15)$$

This completes the proof. \square

Let $\mathbb{M}_{Hod} = \mathbb{M}_{Hod}(X, n)$ (see, [9]) denote the coarse moduli space of isomorphism classes of polystable λ -connections of rank n on X . Then \mathbb{M}_{Hod} is a quasi-projective variety that contains \mathcal{M}_{Hod} as a Zariski open dense subset.

Although, we have assumed that genus $g \geq 3$, but the following proposition is true for $g \geq 2$.

Proposition 3.4. *Let X be a compact Riemann surface of genus $g \geq 2$ and $n \geq 2$. Then we have*

$$\text{Pic}(\mathbb{M}_{Hod}) \cong \text{Pic}(\mathcal{M}_{Hod}) \quad (3.16)$$

Proof. In view of Lemma(3.2), it is sufficient to show that the codimension of the closed set $W = \mathbb{M}_{Hod} \setminus \mathcal{M}_{Hod}$ in \mathbb{M}_{Hod} is atleast 2. And that follows from [9], Proposition 3.1. \square

4. ALGEBRAIC FUNCTIONS ON THE HODGE MODULI SPACE

In this section we will study the algebraic functions on the Hodge moduli space with fixed determinant. Fix a line bundle L over the compact Riemann surface X of degree zero. For each $\lambda \in \mathbb{C}$, fix a λ -connection D_L^λ on L such that for any $\lambda_1, \lambda_2 \in \mathbb{C}$, we have

$$D_L^{\lambda_1} + D_L^{\lambda_2} = D_L^{\lambda_1 + \lambda_2}. \quad (4.1)$$

Let $\mathcal{M}_{Hod}(L) = \mathcal{M}_{Hod}(X, n, L)$ denote the moduli space parametrising all stable λ -connections (E, λ, D) such that

- (1) E is a holomorphic vector bundle of rank n over X with $\bigwedge^n E \cong L$.
- (2) for every $\lambda \in \mathbb{C}$, the λ -connection on $\bigwedge^n E$ induced by D coincides with the given λ -connection D_L^λ on L .

Then $\mathcal{M}_{Hod}(L)$ is a smooth quasi-projective closed subvariety of \mathcal{M}_{Hod} of dimension $2(n^2 - 1)(g - 1) + 1$. Let $\mathcal{M}'_{Hod}(L) = \mathcal{M}_{Hod}(L) \cap \mathcal{M}'_{Hod}$, that is, $\mathcal{M}'_{Hod}(L)$ consists of those points of $\mathcal{M}_{Hod}(L)$ whose underlying vector bundle is stable. Then $\mathcal{M}'_{Hod}(L)$ is a Zariski open dense subset of $\mathcal{M}_{Hod}(L)$ (openness follows from [[10], p.635, Theorem 2.8(B)]).

Again, we have a natural surjective algebraic morphism

$$\Phi_L : \mathcal{M}_{Hod}(L) \rightarrow \mathbb{A}^1 \quad (4.2)$$

defined by $(E, \lambda, D) \mapsto \lambda$. Let Φ_L^λ denote the inverse image $\Phi_L^{-1}(\lambda)$. Then, $\Phi_L^1 = \mathcal{M}_{DR}(X, n, L)$ and $\Phi_L^0 = \mathcal{M}_{Dol}(X, n, L)$ are de Rham moduli space and Dolbeault moduli space with fixed determinant L , respectively. Let \mathcal{U}_L be the moduli space parametrising all the stable holomorphic vector bundle E on X of rank n and degree zero such that $\bigwedge^n E \cong L$.

Let

$$p_0 : \mathcal{M}'_{Hod}(L) \rightarrow \mathcal{U}_L \quad (4.3)$$

be the projection defined by sending (E, λ, D) to E . Then from the assumption (4.1) on the fixed family $(L, \lambda, D_L^\lambda)_{\lambda \in \mathbb{C}}$, for every $E \in \mathcal{U}_L$, $p_0^{-1}(E)$ is a vector space of dimension $(n^2 - 1)(g - 1) + 1$. Note that if we remove the assumption (4.1) from the fixed family $(L, \lambda, D_L^\lambda)_{\lambda \in \mathbb{C}}$, $p_0^{-1}(E)$ need not be a vector space. Now, we have following result similar to Proposition (3.1).

Proposition 4.1. *There exists an algebraic vector bundle $\pi' : \Xi' \rightarrow \mathcal{U}_L$ such that $\mathcal{M}'_{Hod}(L)$ is embedded in $\mathbf{P}(\Xi')$ with $\mathbf{P}(\Xi') \setminus \mathcal{M}'_{Hod}(L)$ as the hyperplane at infinity.*

Proof. cf. Proposition (3.1) □

Proposition 4.2. *The homomorphism $p_0^* : \text{Pic}(\mathcal{U}_L) \rightarrow \text{Pic}(\mathcal{M}'_{Hod}(L))$ defined by $\xi \mapsto p_0^* \xi$ is an isomorphism of groups.*

Proof. The proof is similar to the proof of the Theorem (3.3). □

Now, from [11], Proposition 3.4, (ii), we have $\text{Pic}(\mathcal{U}_L) \cong \mathbb{Z}$. Thus, in view of Theorem (4.2), we have

Corollary 4.3. *$\text{Pic}(\mathcal{M}'_{Hod}(L)) \cong \mathbb{Z}$.*

Now, consider $\Phi_L^1 = \mathcal{M}_{DR}(X, n, L)$ the moduli space of 1-connection on X with fixed determinant $(L, 1, D_L^1)$ that is nothing but moduli space of holomorphic connections $(E, 1, D)$ on X with fixed determinant (L, D_L^1) . Let $\mathcal{M}'_{DR}(L) \subset \Phi_L^1$ denote the Zariski open subvariety such that the underlying vector bundle is stable (openness follows from [[10], p.635, Theorem 2.8(B)]).

Let

$$q : \mathcal{M}'_{DR}(L) \rightarrow \mathcal{U}_L \quad (4.4)$$

be the canonical projection map defined by sending (E, D) to E . From Atiyah [1] and Weil [19], being an indecomposable vector bundle of degree zero, any $E \in \mathcal{U}_L$ admits a holomorphic connection. Thus q is a surjective morphism.

First note that for any $E \in \mathcal{U}_L$, the holomorphic cotangent space $\Omega_{\mathcal{U}_L, E}^1$ at E is isomorphic to $H^0(X, \Omega_X^1 \otimes \text{ad}(E))$, where $\text{ad}(E) \subset \text{End}(E)$ is the subbundle consists of endomorphism of E whose trace is zero. Also, $q^{-1}(E)$ is an affine space modelled over $H^0(X, \Omega_X^1 \otimes \text{ad}(E))$. Thus, there is a natural action of $\Omega_{\mathcal{U}_L, E}^1$ on $q^{-1}(E)$, that is,

$$\Omega_{\mathcal{U}_L, E}^1 \times q^{-1}(E) \rightarrow q^{-1}(E)$$

sending (ω, D) to $\omega + D$, which is faithful and transitive. Thus, we have (see [12] for the definition of torsor and similar results for instance [12] Proposition 4.2)

Proposition 4.4. *Let $q : \mathcal{M}'_{DR}(L) \rightarrow \mathcal{U}_L$ be the map as defined in (4.4). Then $\mathcal{M}'_{DR}(L)$ is a $\Omega_{\mathcal{U}_L}^1$ -torsor on \mathcal{U}_L .*

Proposition 4.5. *The homomorphism $q^* : \text{Pic}(\mathcal{U}_L) \rightarrow \text{Pic}(\mathcal{M}'_{DR}(L))$ defined by $\xi \mapsto q^*\xi$ is an isomorphism of groups.*

Proof. The arguments involved in the proof is parallel to the Theorem (3.3). \square

Now, from [11], Proposition 3.4, (ii), we have $\text{Pic}(\mathcal{U}_L) \cong \mathbb{Z}$. Thus, in view of Proposition (4.5), we have

$$\text{Pic}(\mathcal{M}'_{DR}(L)) \cong \mathbb{Z}. \quad (4.5)$$

Let Θ be the ample generator of the group $\text{Pic}(\mathcal{U}_L)$. We have **Atiyah exact sequence**

$$0 \rightarrow \mathcal{O}_{\mathcal{U}_L} \xrightarrow{i} \text{At}(\Theta) \xrightarrow{\sigma} T\mathcal{U}_L \rightarrow 0, \quad (4.6)$$

where $\text{At}(\Theta)$ is called **Atiyah algebra** of holomorphic line bundle Θ , and in this case it is equal to $\mathcal{D}\text{iff}^1(\Theta, \Theta)$, the sheaf of first order differential operators on Θ for more details see [1].

Dualising the above Atiyah exact sequence (4.6), we get following exact sequence,

$$0 \rightarrow \Omega_{\mathcal{U}_L}^1 \xrightarrow{\sigma^*} \text{At}(\Theta)^* \xrightarrow{i^*} \mathcal{O}_{\mathcal{U}_L} \rightarrow 0 \quad (4.7)$$

Consider $\mathcal{O}_{\mathcal{U}_L}$ as trivial line bundle $\mathcal{U}_L \times \mathbb{C}$. Let $t : \mathcal{U}_L \rightarrow \mathcal{U}_L \times \mathbb{C}$ be a holomorphic map defined by $E \mapsto (E, 1)$. Then t is a holomorphic section of the trivial line bundle $\mathcal{U}_L \times \mathbb{C}$.

Let $T = \text{Im}(t) \subset \mathcal{U}_L \times \mathbb{C}$ be the image of t . Then $T \rightarrow \mathcal{U}_L$ is a fibre bundle. Consider the inverse image $i^{*-1}T \subset \text{At}(\Theta)^*$, and denote it by $\mathcal{C}(\Theta)$. Using the same technique as in [12], (see Proposition 5.4), we can show that

$$\mathcal{C}(\Theta) \cong \mathcal{M}'_{DR}(L) \quad (4.8)$$

Lemma 4.6. *Assume that $\text{genus}(X) \geq 3$. Then*

$$H^0(\mathcal{M}'_{DR}(L), \mathcal{O}_{\mathcal{M}'_{DR}(L)}) = \mathbb{C}. \quad (4.9)$$

Proof. In view of the above isomorphism (4.8) of varieties, it is enough to show that $\mathcal{C}(\Theta)$ does not have any non-constant algebraic function. Now, from [5], Theorem 4.3, p.797 (also see [12], Theorem 5.7), we have

$$H^0(\mathcal{C}(\Theta), \mathcal{O}_{\mathcal{C}(\Theta)}) = \mathbb{C}.$$

□

Proposition 4.7. $H^0(\Phi_L^1, \mathcal{O}_{\Phi_L^1}) = \mathbb{C}$.

Proof. Since $\mathcal{M}'_{DR}(L) \subset \Phi_L^1$ is an open dense subvariety. Therefore, from above Lemma (4.6), conclusion follows. □

Next, for any $\lambda \neq 0$, Φ_L^λ is isomorphic to Φ_L^1 , by a morphism sending (E, λ, D) to $(E, 1, \lambda^{-1}D)$. Thus we have

Corollary 4.8. *For any $\lambda \neq 0$, $H^0(\Phi_L^\lambda, \mathcal{O}_{\Phi_L^\lambda}) = \mathbb{C}$.*

Now, we are in position to describe the algebraic functions on the moduli space $\mathcal{M}_{Hod}(L)$, which generalizes a result (see [4] Theorem 4.1, p.1542) to higher rank case on the compact Riemann surface X .

Theorem 4.9. *Any algebraic function on $\mathcal{M}_{Hod}(L)$ will factor through the map*

$$\Phi_L : \mathcal{M}_{Hod}(L) \rightarrow \mathbb{A}^1$$

defined in (4.2).

Proof. Let $g : \mathcal{M}_{Hod}(L) \rightarrow \mathbb{C}$ be an algebraic function. Then, for each $\lambda \neq 0$, the restriction of g to each fibre Φ_L^λ of Φ_L is a constant function, follows from Corollary (4.8). Since

$$\coprod_{0 \neq \lambda \in \mathbb{C}} \Phi_L^\lambda \hookrightarrow \mathcal{M}_{Hod}(L) \quad (4.10)$$

is a dense open subset of $\mathcal{M}_{Hod}(L)$, and hence defining a function from $\mathbb{A}^1 \rightarrow \mathbb{A}^1$. □

Let $\mathbb{M}_{Hod}(L) = \mathbb{M}_{Hod}(X, n, L)$ denote the moduli space of isomorphism classes of polystable λ -connections of rank n on X such that

- (1) E is a holomorphic vector bundle of rank n over X with $\bigwedge^n E \cong L$.
- (2) for every $\lambda \in \mathbb{C}$, the λ -connection on $\bigwedge^n E$ induced by D coincides with the given λ -connection D_L^λ on L .

Then $\mathbb{M}_{Hod}(L)$ is a quasi-projective variety that contains $\mathcal{M}_{Hod}(L)$ as a Zariski open dense subset, and we have the canonical map

$$\tau : \mathbb{M}_{Hod}(L) \rightarrow \mathbb{A}^1 \quad (4.11)$$

defined by sending $(E, \lambda, D) \mapsto \lambda$.

Corollary 4.10. *Any algebraic function on $\mathbb{M}_{Hod}(L)$ will factor through the map*

$$\tau : \mathbb{M}_{Hod}(L) \rightarrow \mathbb{A}^1$$

defined in (4.11).

5. AUTOMORPHISM OF HODGE MODULI SPACE

In [3], the automorphism group of moduli space of rank one λ -connections have been studied. In this section we will obtain a short exact sequence which contains the group of algebraic automorphism of Hodge moduli space. The group of algebraic automorphisms of the moduli spaces $\mathcal{M}'_{Hod}(L)$ and \mathcal{U}_L is denoted by $\text{Aut}(\mathcal{M}'_{Hod}(L))$ and $\text{Aut}(\mathcal{U}_L)$, respectively. Let

$$\text{Aut}(\mathcal{M}'_{Hod}(L))_0 \subset \text{Aut}(\mathcal{M}'_{Hod}(L)) \text{ and } \text{Aut}(\mathcal{U}_L)_0 \subset \text{Aut}(\mathcal{U}_L)$$

be the connected component of the respective groups of algebraic automorphisms containing the identity automorphism.

Let $\Psi : \mathcal{M}'_{Hod}(L) \rightarrow \mathbb{A}^1$ be the canonical projection defined by $(E, \lambda, D) \mapsto \lambda$. Then, for each $\lambda \in \mathbb{C}$, we denote the inverse image $\Psi^{-1}(\lambda)$ by Ψ^λ . For each $\lambda \in \mathbb{C}$, we have canonical projection

$$f^\lambda : \Psi^\lambda \rightarrow \mathcal{U}, \quad (5.1)$$

defined by $(E, \lambda, D) \rightarrow E$. Let \mathbb{V} consists of those $T \in \text{Aut}(\mathcal{M}'_{Hod}(L))_0$ such that for every $\lambda \in \mathbb{C}$, we have

- (1) $T|_{\Psi^\lambda} : \Psi^\lambda \rightarrow \Psi^\lambda$.
- (2) $f^\lambda \circ T|_{\Psi^\lambda} = f^\lambda$.

Then \mathbb{V} is a subgroup of $\text{Aut}(\mathcal{M}'_{Hod}(L))_0$. Let \mathbb{W} denote the group of all algebraic maps from \mathbb{C} to $\text{Aut}(\mathcal{U}_L)_0$. Then we have following

Theorem 5.1. *The group $\text{Aut}(\mathcal{M}'_{Hod}(L))_0$ fits in a short exact sequence*

$$0 \rightarrow \mathbb{V} \xrightarrow{\iota} \text{Aut}(\mathcal{M}'_{Hod}(L))_0 \xrightarrow{\rho} \mathbb{W} \times \mathbb{C}^* \rightarrow 0 \quad (5.2)$$

of groups.

Proof. Let $T \in \text{Aut}(\mathcal{M}'_{Hod}(L))_0$. From Proposition (4.8), Ψ^λ does not admit any nonconstant algebraic function, for any $\lambda \in \mathbb{C}^*$. Hence the composition

$$\Psi^\lambda \xrightarrow{T|_{\Psi^\lambda}} \mathcal{M}'_{Hod}(L) \xrightarrow{\Psi} \mathbb{A}^1, \quad (5.3)$$

is a constant function for all $\lambda \in \mathbb{C}^*$. Note that for every $\lambda \in \mathbb{C}^*$, Ψ^0 is not isomorphic to Ψ^λ , therefore

$$\Psi(T(\Psi^\lambda)) \in \mathbb{C}^*. \quad (5.4)$$

Now, T being an automorphism, (5.4) holds for T^{-1} , and hence the composition in (5.3) is a constant function for every $\lambda \in \mathbb{C}$. This implies that there is an algebraic automorphism $\gamma_0 : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ such that the following diagram

$$\begin{array}{ccc} \mathcal{M}'_{Hod}(L) & \xrightarrow{T} & \mathcal{M}'_{Hod}(L) \\ \downarrow \Psi & & \downarrow \Psi \\ \mathbb{A}^1 & \xrightarrow{\gamma_0} & \mathbb{A}^1 \end{array} \quad (5.5)$$

commutes. Since Ψ^0 is not isomorphic to Ψ^λ for every $\lambda \in \mathbb{C}^*$, therefore $\gamma_0(0) = 0$. Thus, $\gamma_0(z) = \gamma \cdot z$ for some $\gamma \in \mathbb{C}^*$ and $z \in \mathbb{A}^1$. Clearly, $\gamma \in \mathbb{C}^*$ depends on T .

For any $\lambda \in \mathbb{C}$, let

$$\psi^\lambda : \Psi^\lambda \rightarrow \mathcal{U}_L \quad (5.6)$$

be the canonical projection defined by sending $(E, \lambda, D) \rightarrow E$.

For $\lambda \neq 0$ the above map is surjective, and the fibres of ψ^λ is an affine space modelled over $H^0(X, \Omega_X^1 \otimes \text{End}(E))$ and for $\lambda = 0$, the inverse image $(\psi^0)^{-1}(E) = H^0(X, \Omega_X^1 \otimes \text{End}(E))$.

Thus, for given $T \in \text{Aut}(\mathcal{M}'_{\text{Hod}}(L))_0$, there exists a unique automorphism

$$\Upsilon(\lambda) : \mathcal{U}_L \rightarrow \mathcal{U}_L \quad (5.7)$$

such that the following diagram

$$\begin{array}{ccc} \Psi^\lambda & \xrightarrow{T|_{\Psi^\lambda}} & \Psi^{\gamma\lambda} \\ \downarrow \psi^\lambda & & \downarrow \psi^{\gamma\lambda} \\ \mathcal{U}_L & \xrightarrow{\Upsilon(\lambda)} & \mathcal{U}_L \end{array} \quad (5.8)$$

commutes, for every $\lambda \in \mathbb{C}$. Also, note that $\Upsilon(\gamma) \in \text{Aut}(\mathcal{U}_L)_0$. From above phenomenon, we define

$$\rho : \text{Aut}(\mathcal{M}'_{\text{Hod}}(L))_0 \rightarrow \mathbb{W} \times \mathbb{C}^* \quad (5.9)$$

by $T \mapsto (\Upsilon, \gamma)$. It is straightforward to check that ρ is a surjective group homomorphism. Now, the kernel of ρ is \mathbb{V} , which gives the short exact sequence (5.2) of groups.

□

REFERENCES

- [1] Atiyah, M. F., Complex analytic connections in fibre bundles, *Trans. Amer. Math.Soc.* 85(1957), 181-207.
- [2] Bhosle, U. N., Picard group of the moduli space of vector bundles, *Math. Ann.* 314 (1999) 245-263.
- [3] Biswas, I., Heller, S., On the automorphisms of a rank one DeligneHitchin moduli space, *SIGMA* 13 (2017), 072.
- [4] Biswas, I., Hurtubise, J., Raina, A. K., Rank one connections on abelian varieties. *Internat. J. Math.* 22 (2011), no. 11.
- [5] Biswas, I., Raghavendra, N., Line bundles over a moduli space of logarithmic connections on a Riemann surface. *Geom. Funct. Anal.* 15 (2005), 780-808.
- [6] Deligne, P., Various letters to C. Simpson.
- [7] Drézet, J. M., and Narasimhan, M.S. Groupe de Picard des variétés de modules de fibrés semi-stable sur les courbes algébriques. *Invent. Math.*, 97(1):5394, 1989.
- [8] Hartshorne, R., Algebraic geometry, Graduate Texts in Mathematics, Vol. 52. *Springer-Verlag, New York-Heidelberg*, 1977. xvi+496 pp. ISBN: 0-387-90244-9
- [9] Hu, Z., Huang, P., Flat λ -connections, mochiuzuki correspondence and twistor spaces. <https://arxiv.org/pdf/1905.10765.pdf>
- [10] Maruyama, M., Openness of a family of torsion free sheaves, *Jour. Math. Kyoto Univ.* 16(1976), 627-637.
- [11] Ramanan, S., The moduli space of vector bundles over an algebraic curve, *Math. Ann.* 200(1973), 69-84.
- [12] Singh, A., Moduli space of logarithmic connections singular over a finite subset of a compact Riemann surface, arXiv:1912.01288.
- [13] Simpson, C.T.: Moduli of representations of the fundamental group of a smooth projective variety. I. *Inst. Hautes tudes Sci. Publ. Math.* 79, 47129 (1994)

- [14] Simpson, C.T.: Moduli of representations of the fundamental group of a smooth projective variety. II. *Inst. Hautes Etudes Sci. Publ. Math.* 79, 471-29 (1994)
- [15] Simpson, C.T., The Hodge filtration on nonabelian cohomology, in Algebraic geometry-Santa Cruz 1995, 217-281
- [16] Simpson, C.T. A weight two phenomenon for the moduli space of rank one local systems on open varieties. <https://arxiv.org/abs/0710.2800v1>[math.AG], 2007.
- [17] Simpson, C.T., Iterated destability modifications for vector bundles with connection, *Contemp. Math.* 522 (2010), 183-206.
- [18] Simpson, C.T., Nonabelian Hodge theory. Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990), 747-756, Math. Soc. Japan, Tokyo, 1991.
- [19] Weil, A., Generalization de fonctions abeliennes, *J. Math. Pures Appl.* vol. **17**(1938), 47-87.

HARISH-CHANDRA RESEARCH INSTITUTE, HBNI, CHHATNAG ROAD, JHUSI, PRAYAGRAJ 211 019,
INDIA

Email: anoopsingh@hri.res.in