

Fractional matching number and spectral radius of nonnegative matrix of graphs

Ruifang Liu ^{*}; Hong-Jian Lai [†]; Litao Guo [‡]; Jie Xue [§]

Abstract

A fractional matching of a graph G is a function $f : E(G) \rightarrow [0, 1]$ such that for any $v \in V(G)$, $\sum_{e \in E_G(v)} f(e) \leq 1$ where $E_G(v) = \{e \in E(G) : e \text{ is incident with } v \text{ in } G\}$. The fractional matching number of G is $\mu_f(G) = \max\{\sum_{e \in E(G)} f(e) : f \text{ is fractional matching of } G\}$. For any real numbers $a \geq 0$ and $k \in (0, n)$, it is observed that if $n = |V(G)|$ and $\delta(G) > \frac{n-k}{2}$, then $\mu_f(G) > \frac{n-k}{2}$. We determine a function $\varphi(a, n, \delta, k)$ and show that for a connected graph G with $n = |V(G)|$, $\delta(G) \leq \frac{n-k}{2}$, spectral radius $\lambda_1(G)$ and complement \overline{G} , each of the following holds.

- (i) If $\lambda_1(aD(G) + A(G)) < \varphi(a, n, \delta, k)$, then $\mu_f(G) > \frac{n-k}{2}$.
- (ii) If $\lambda_1(aD(\overline{G}) + A(\overline{G})) < (a+1)(\delta+k-1)$, then $\mu_f(G) > \frac{n-k}{2}$.

As corollaries, sufficient spectral condition for fractional perfect matchings and analogous results involving Q -index and A_α -spectral radius are obtained, and former spectral results in [European J. Combin. 55 (2016) 144-148] are extended.

AMS Classification: 05C50

Key words: Signless Laplacian spectral radius; Fractional matching; Fractional perfect matching

1 Introduction

Graphs considered are simple and undirected. We generally follow [4] for undefined terms and notation. For a graph G , and a vertex $v \in V(G)$, define $N_G(v) = \{u \in V(G) : uv \in E(G)\}$ and $E_G(v) = \{e \in E(G) : e \text{ is incident with } v \text{ in } G\}$. If $S \subseteq V(G)$, then define $N_G(S) = \cup_{v \in S} N_G(v)$. As in [4], we use K_n , $\delta(G)$, $\Delta(G)$, $\kappa(G)$ and \overline{G} to denote the complete graph of order n , the minimum degree, the maximum degree, the connectivity and the complement of G , respectively.

^{*}School of Mathematics and Statistics, Zhengzhou University, Zhengzhou, Henan 450001, China. Email: rfliu@zzu.edu.cn

[†]Corresponding author. Department of Mathematics, West Virginia University, Morgantown, WV 26506, USA. E-mail: hjlai@math.wvu.edu

[‡]School of Applied Mathematics, Xiamen University of Technology, Xiamen, Fujian 361024, China. Email: ltguo2012@126.com

[§]School of Mathematics and Statistics, Zhengzhou University, Zhengzhou, Henan 450001, China. Email: jie_xue@126.com

For a subset $S \subseteq V(G)$ or $S \subseteq E(G)$, $G[S]$ denotes the subgraph of G induced by S . A cycle (or path, respectively) on n vertices is often denoted by C_n (or P_n , respectively). A cycle containing all vertices of a graph G is called a Hamilton cycle of G . Let G be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$. The **adjacency matrix** of G is the n by n matrix $A(G) := (a_{ij})$, where a_{ij} is the number of edges joining v_i and v_j in G , and the **diagonal degree matrix** of G is the n by n matrix $D(G) := (d_{ij})$, where $d_{ij} = 0$ if $i \neq j$ and $d_{ii} = d_G(v_i)$, for all i with $1 \leq i \leq n$. For a subset $X \subseteq E(G)$, the **characteristic function** of X is a function $f_X : E \rightarrow \{0, 1\}$ such that $f(e) = 1$ if and only if $e \in X$.

Our approach is motivated by the methods deployed in [11, 13, 14, 15, 16], among others. For any graph G with the adjacency matrix $A(G)$ and the diagonal degree matrix $D(G)$ and for real numbers $a \geq 0$ and $b > 0$, define $J(G : a, b) = aD(G) + bA(G)$. As $J(G : a, b)$ is also real and symmetric, all its eigenvalues are real. Let $\lambda_i(J(G : a, b))$ to be the i th largest eigenvalue of $J(G : a, b)$. In particular, $\lambda_1(J(G : 0, 1)) = \lambda_1(G)$ is known as the **spectral radius** of G ; $\lambda_1(J(G : 1, 1)) = q_1(G)$ is the **Q -index** of G ; and $\lambda_1(J(G : \alpha, 1 - \alpha)) = \lambda_1(A_\alpha(G))$ is called the **A_α -spectral radius** of G with $\alpha \in [0, 1]$, introduced by Nikiforov in [17].

As in [4], an edge subset $M \subseteq E(G)$ is a **matching** of a graph G if $\Delta(G[M]) \leq 1$. The **matching number** of G is $\alpha'(G) = \max\{|M| : M \text{ is a matching of } G\}$. A **fractional matching** of a graph G is a function $f : E(G) \rightarrow [0, 1]$ such that for any $v \in V(G)$, $\sum_{e \in E_G(v)} f(e) \leq 1$. Thus if $f(e) \in \{0, 1\}$ for every edge $e \in E(G)$, then f is the characteristic function of a matching in G . The **fractional matching number** of G is $\mu_f(G) = \max\{\sum_{e \in E(G)} f(e) : f \text{ is fractional matching of } G\}$. The following is known.

Theorem 1.1 *Let G be a graph with $n = |V(G)|$. Each of the following holds.*

- (i) (Lemma 2.1.2 of [20]) *Any fractional matching f satisfies $\mu_f(G) \leq n/2$.*
- (ii) (Theorem 2.1.3 of [20]) *If G is bipartite, then $\mu_f(G) = \alpha'(G)$.*
- (iii) (Theorem 2.1.5 of [20]) *$2\mu_f(G)$ is an integer.*

A fractional matching f of G is a **fractional perfect matching** if $\sum_{e \in E(G)} f(e) = n/2$. If a fractional perfect matching takes values only in $\{0, 1\}$, then it is the characteristic function of a perfect matching.

Let $i(G)$ be the number of isolated vertices of a graph G . In [20], the following are displayed.

Theorem 1.2 (Scheinerman and Ullman [20]) *Let G be a graph. Each of the following holds.*

- (i) $\mu_f(G) \geq \alpha'(G)$.
- (ii) (The fractional Berge-Tutte Formula)

$$\mu_f(G) = \frac{1}{2} (n - \max \{i(G - S) - |S| : \forall S \subseteq V(G)\}). \quad (1)$$

- (iii) (The fractional Tutte's 1-Factor Theorem) *A graph G has a fractional perfect matching if and only if*

$$\forall S \subseteq V(G), i(G - S) \leq |S|. \quad (2)$$

The investigation on the relationship between the eigenvalues and the matching number of a graph was initiated by Brouwer and Haemers [2], in which sufficient conditions on $\lambda_3(G)$ to

assure the existence of a perfect matching were discussed. Cioabă et al. in [5, 6, 7] improved and generalized these results to obtain a best possible upper bound on $\lambda_3(G)$ for the existence of a perfect matching. Furthermore, O and Cioabă [19] determined the relationship between the eigenvalues of a t -edge-connected k -regular graph and its matching number when $t \leq k - 2$.

On the fractional matching number, O [18] studied the connections between the fractional matching number and the spectral radius of a connected graph with given minimum degree. Xue et al. [21] considered the relationship between the fractional matching number and the Laplacian spectral radius of a graph. The main goal of this study is to investigate the relationship between the fractional matching number and the largest eigenvalue of the nonnegative matrix $J(G : a, b)$ of a graph G . This provides a mechanism to have a unified approach to both adjacency eigenvalues and signless Laplacian eigenvalues, and is quite different from those using only the Laplacian matrix. Consequently, analogous results involving the Q -index and A_α -spectral radius of a graph G are also obtained, and former results on spectral radius in [18] are extended.

We first observe that (see Theorem 2.2 in the next section) for any real number $k \in (0, n)$ and a connected graph G with $n = |V(G)|$ and $\delta = \delta(G)$, if $\delta > \frac{n-k}{2}$, then $\mu_f(G) > \frac{n-k}{2}$. Thus we focus on the investigation of $\mu_f(G)$ when $\delta \leq \frac{n-k}{2}$.

For real numbers a, k and an integer δ , with $a \geq 0$, $k \in (0, n)$ and $1 \leq \delta \leq \frac{n-k}{2}$, define

$$\varphi(a, n, \delta, k) = \begin{cases} \delta \sqrt{1 + \frac{2k}{n-k}} & \text{if } a = 0, \\ \frac{2a\delta n}{n-k} & \text{if } a \in (0, 1], \\ \frac{a\delta(n+k)}{n-k} & \text{if } a \in (1, +\infty). \end{cases} \quad (3)$$

Our main results are the following.

Theorem 1.3 *Let a be a real number with $a \geq 0$, $k \in (0, n)$ be a real number, and G be a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$. If*

$$\lambda_1(aD(G) + A(G)) < \varphi(a, n, \delta, k),$$

then $\mu_f(G) > \frac{n-k}{2}$.

The upper bound in Theorem 1.3 is best possible when $a \in \{0, 1\}$, as will be shown in Section 3. A sufficient condition to ensure $\mu_f(G) > \frac{n-k}{2}$ in terms of $\lambda_1(J(\overline{G} : a, 1))$ is obtained in Theorem 1.4 below. The upper bound in Theorem 1.4 is also best possible in some sense.

Theorem 1.4 *Let a be a real number with $a \geq 0$, $k \in (0, n)$ be a real number, and G be a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$. If*

$$\lambda_1(aD(\overline{G}) + A(\overline{G})) < (a+1)(\delta+k-1),$$

then $\mu_f(G) > \frac{n-k}{2}$.

The main results can be applied to obtain sufficient conditions in terms of Q -index $q_1(G)$ and A_α -spectral radius $\lambda_1(A_\alpha(G))$ for a simple graph G to have a large value of $\mu_f(G)$. Detailed justifications of the following corollaries will be presented in Section 4.

Corollary 1.5 Let G be a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$. Each of the following holds.

- (i) If $q_1(G) < \frac{2\delta n}{n-k}$, then $\mu_f(G) > \frac{n-k}{2}$.
- (ii) If $q_1(\overline{G}) < 2(\delta + k - 1)$, then $\mu_f(G) > \frac{n-k}{2}$.

Corollary 1.6 Let G be a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$, and let α be a real number with $0 \leq \alpha \leq 1$. Each of the following holds.

- (i) If $\alpha = 0$ and $\lambda_1(A_\alpha(G)) < \delta \sqrt{1 + \frac{2k}{n-k}}$, then $\mu_f(G) > \frac{n-k}{2}$.
- (ii) If $0 < \alpha \leq \frac{1}{2}$ and $\lambda_1(A_\alpha(G)) < \frac{2\alpha\delta n}{n-k}$, then $\mu_f(G) > \frac{n-k}{2}$.
- (iii) If $\frac{1}{2} < \alpha \leq 1$ and $\lambda_1(A_\alpha(G)) < \frac{\alpha\delta(n+k)}{n-k}$, then $\mu_f(G) > \frac{n-k}{2}$.
- (iv) If $\lambda_1(A_\alpha(\overline{G})) < \delta + k - 1$, then $\mu_f(G) > \frac{n-k}{2}$.

Preliminaries will be presented in the next section. Proofs of the main results and discussions of the consequences will be given in the last two sections.

2 Preliminaries

Throughout this section, we always assume that $k \in (0, n)$ is a real number, and G is a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G)$. A path with end vertices u and v is called a (u, v) -path. We start with some of the well known results of Dirac.

Theorem 2.1 (see also Bondy and Murty [4]) Let G be a simple graph with $n = |V(G)|$ and $\delta = \delta(G) > 0$. Each of the following holds.

- (i) (Dirac [8]) If $\delta \geq n/2$ and $n \geq 3$, then G is hamiltonian.
- (ii) (Dirac [8]) If $\kappa(G) \geq 2$ and $\delta \leq n/2$, then G contains a cycle of length at least 2δ .
- (iii) G contains a path of length δ .

Theorem 2.1 can be applied to prove Theorem 2.2 below, which indicates that the focus of the research should be restricted to the cases when $\delta \leq \frac{n-k}{2}$.

Theorem 2.2 Let $s \geq 1$ be an integer. Each of the following holds.

- (i) If G has a subgraph L with $\Delta(L) \leq 2$ and $|E(L)| \geq 2s$, then $\mu_f(G) \geq s$. In particular, if G has a cycle or path with at least 2δ edges, then $\mu_f(G) \geq \delta$.
- (ii) If $\delta > \frac{n-k}{2}$, then $\mu_f(G) > \frac{n-k}{2}$.

Proof. If $\Delta(L) \leq 2$, then $f = \frac{1}{2}f_{E(L)}$ is a fractional matching, and so $\mu_f(G) \geq \frac{|E(L)|}{2} \geq s$. This proves (i). To show (ii), we observe that $\delta > 0$. As $n \geq n - k$, by Theorem 2.1(i) and Theorem 2.2(i), we may assume that $\frac{n-k}{2} < \delta < \frac{n}{2}$. If $\kappa(G) \geq 2$, then by Theorem 2.1(ii), G contains a cycle C_s with $s = |E(C_s)| \geq 2\delta$. By Theorem 2.2(i), $\mu_f(G) \geq \delta > \frac{n-k}{2}$. Hence we assume that G contains a cut vertex u . Let G_1 and G_2 are two of connected components of $G - u$. Let $v_1, v_2 \in N_G(u)$ with $v_1 \in V(G_1)$ and $v_2 \in V(G_2)$. As G is simple, $\delta(G - u) \geq \delta(G) - 1$.

In particular, $\delta(G_i) \geq \delta(G) - 1$, for $i \in \{1, 2\}$. By Theorem 2.1(iii), for $i \in \{1, 2\}$, each G_i contains a (v_i, v'_i) -path P_i , for some $v'_i \in V(G_i - v_i)$, with $|E(P_i)| \geq \delta - 1$, then G contains a path $P = G[E(P_1) \cup v_1uv_2 \cup E(P_2)]$ of length 2δ , and by Theorem 2.2(i), $\mu_f(G) \geq \delta > \frac{n-k}{2}$. ■

The main tool in our paper is the following eigenvalue interlacing technique. Given two non-increasing real sequences $\theta_1 \geq \theta_2 \geq \dots \geq \theta_n$ and $\eta_1 \geq \eta_2 \geq \dots \geq \eta_m$ with $n > m$, the second sequence is said to **interlace** the first one if $\theta_i \geq \eta_i \geq \theta_{n-m+i}$ for $i = 1, 2, \dots, m$. The interlacing is **tight** if exists an integer $k \in [0, m]$ such that $\theta_i = \eta_i$ for $1 \leq i \leq k$ and $\theta_{n-m+i} = \eta_i$ for $k+1 \leq i \leq m$.

Let M be the following $n \times n$ matrix

$$M = \begin{pmatrix} M_{1,1} & M_{1,2} & \cdots & M_{1,m} \\ M_{2,1} & M_{2,2} & \cdots & M_{2,m} \\ \vdots & \vdots & \ddots & \vdots \\ M_{m,1} & M_{m,2} & \cdots & M_{m,m} \end{pmatrix},$$

whose rows and columns are partitioned into subsets X_1, X_2, \dots, X_m of $\{1, 2, \dots, n\}$. Let $M_{i,j}$ denote the submatrix (called a block) of M by deleting the rows in $\{1, 2, \dots, n\} - X_i$ and deleting the columns in $\{1, 2, \dots, n\} - X_j$. The **quotient matrix** R of the matrix M (with respect to the given partition) is the $m \times m$ matrix whose entries are the average row sums of the blocks $M_{i,j}$ of M . The partition is **equitable** if each block $M_{i,j}$ of M has constant row (and column) sum.

Theorem 2.3 (Brouwer and Haemers [3, 10]) *Let M be a real symmetric matrix. Then the eigenvalues of every quotient matrix of M interlace the ones of M . Furthermore, if the interlacing is tight, then the partition is equitable.*

Theorem 2.4 (Haynsworth [9], You et al. [22]) *Let M be a partitioned matrix, and R be its equitable quotient matrix. Then the eigenvalues of the quotient matrix R are eigenvalues of M . Furthermore, if M is a nonnegative matrix, then the spectral radius of the quotient matrix R equals to the spectral radius of M .*

Let $A = (a_{ij})$ and $B = (b_{ij})$ be two $n \times n$ matrices. Define $A \leq B$ if $a_{ij} \leq b_{ij}$ for all i and j ; and $A < B$ if $A \leq B$ and $A \neq B$.

Theorem 2.5 (Berman and Plemmons [1], Horn and Johnson [12]) *Let $A = (a_{ij})$ and $B = (b_{ij})$ be two $n \times n$ matrices with the spectral radii $\lambda_1(A)$ and $\lambda_1(B)$. If $0 \leq A \leq B$, then $\lambda_1(A) \leq \lambda_1(B)$. Furthermore, if B is irreducible and $0 \leq A < B$, then $\lambda_1(A) < \lambda_1(B)$.*

3 Proof of Theorems 1.3 and 1.4

Following the notation in [4], for $S, T \subseteq V(G)$, define $E_G([S, T]) = \{uv \in E(G) : u \in S \text{ and } v \in T\}$. We will use (3) as the definition of the function $\varphi(a, n, \delta, k)$. Throughout this section, we always assume that n is a positive integer, a and k are real numbers satisfying $a \geq 0$ and $0 < k < n$, and that G is a connected simple graph with $n = |V(G)|$, $\delta = \delta(G) \leq \frac{n-k}{2}$.

3.1 Proof of Theorem 1.3

By contradiction, we assume that $\mu_f(G) \leq \frac{n-k}{2}$. By (1), there exists a vertex subset $S \subseteq V(G)$ satisfying $i(G-S) - |S| \geq k$. Let T be the set of isolated vertices in $G-S$, $|S| = s$ and $|T| = t$. Then $s+t \leq n$ and $t = i(G-S) \geq s+k$, and so $s \leq \frac{n-k}{2}$. As T is the set of isolated vertices in $G-S$, we observe that $N_G(T) \subseteq S$, and so $s \geq \delta$.

Let $X = E_G[S, T]$, $H = G[X]$ be the bipartite subgraph G induced by X , and $r = |E(H)|$. Then $r \geq t\delta$. Accordingly, the quotient matrix $R(aD(H) + A(H))$ of $aD(H) + A(H)$ with respect to the partition (S, T) becomes:

$$R(aD(H) + A(H)) = \begin{pmatrix} \frac{ar}{s} & \frac{r}{s} \\ \frac{r}{t} & \frac{ar}{t} \end{pmatrix}.$$

As the characteristic polynomial of $R(aD(H) + A(H))$ is $\lambda^2 - a(\frac{r}{s} + \frac{r}{t})\lambda + (a^2 - 1)\frac{r^2}{st} = 0$, direct computation yields

$$\begin{aligned} \lambda_1(R(aD(H) + A(H))) &= \frac{1}{2} \left\{ a \left(\frac{r}{s} + \frac{r}{t} \right) + \sqrt{a^2 \left(\frac{r}{s} + \frac{r}{t} \right)^2 - 4(a^2 - 1) \frac{r^2}{st}} \right\} \\ &= \frac{1}{2} \left\{ a \left(\frac{r}{s} + \frac{r}{t} \right) + \sqrt{(a^2 - 1) \left(\frac{r}{s} - \frac{r}{t} \right)^2 + \left(\frac{r}{s} + \frac{r}{t} \right)^2} \right\}. \end{aligned} \quad (4)$$

By Theorems 2.5 and 2.3, to reach a contradiction to the assumption of Theorem 1.3, it suffices to show that in each of the following cases, we always have

$$\lambda_1(R(aD(H) + A(H))) \geq \varphi(a, n, \delta, k), \quad (5)$$

Case 3.1 $a = 0$.

By (4), we have $\lambda_1(R(aD(H) + A(H))) = r\sqrt{\frac{1}{st}}$. Thus

$$\begin{aligned} \lambda_1(aD(G) + A(G)) &\geq \lambda_1(aD(H \cup (n-s-t)K_1) + A(H \cup (n-s-t)K_1)) \\ &= \lambda_1(aD(H) + A(H)) \geq \lambda_1(R(aD(H) + A(H))) \\ &= r\sqrt{\frac{1}{st}} \geq \delta\sqrt{\frac{t}{s}} \geq \delta\sqrt{1 + \frac{k}{s}} \\ &\geq \delta\sqrt{1 + \frac{2k}{n-k}}. \end{aligned}$$

Case 3.2 $0 < a \leq 1$.

This implies that $a^2 - 1 \leq 0$, and so by (4),

$$\lambda_1(R(aD(H) + A(H))) \geq \frac{1}{2} \left\{ a \left(\frac{r}{s} + \frac{r}{t} \right) + \sqrt{(a^2 - 1) \left(\frac{r}{s} + \frac{r}{t} \right)^2 + \left(\frac{r}{s} + \frac{r}{t} \right)^2} \right\} = a \left(\frac{r}{s} + \frac{r}{t} \right).$$

It follows that

$$\begin{aligned} \lambda_1(aD(G) + A(G)) &\geq \lambda_1(aD(H) + A(H)) \geq \lambda_1(R(aD(H) + A(H))) \\ &\geq a \left(\frac{r}{s} + \frac{r}{t} \right) \geq a\delta \left(1 + \frac{t}{s} \right) \geq a\delta \left(2 + \frac{k}{s} \right) \geq \frac{2a\delta n}{n-k}. \end{aligned} \quad (6)$$

Case 3.3 $1 < a < +\infty$.

This implies that $a^2 - 1 > 0$, and so by (4),

$$\lambda_1(R(aD(H) + A(H))) > \frac{1}{2}\left\{a\left(\frac{r}{s} + \frac{r}{t}\right) + \sqrt{(a^2 - 1)\left(\frac{r}{s} - \frac{r}{t}\right)^2 + \left(\frac{r}{s} - \frac{r}{t}\right)^2}\right\} = \frac{ar}{s}.$$

It follows that

$$\begin{aligned} \lambda_1(aD(G) + A(G)) &\geq \lambda_1(aD(H) + A(H)) \geq \lambda_1(R(aD(H) + A(H))) \\ &> \frac{ar}{s} \geq a\delta\frac{t}{s} \geq a\delta\left(1 + \frac{k}{s}\right) \geq \frac{a\delta(n+k)}{n-k}. \end{aligned}$$

As in all the possible values of a , (5) is always obtained, Theorem 1.3 is now justified. \blacksquare

3.2 Sharpness Discussion of Theorem 1.3

Following [4], for disjoint vertex sets X and Y , $G[X, Y]$ denotes a bipartite graph with bipartition (X, Y) . We examine the following example to see that Theorem 1.3 is best possible when $a \in \{0, 1\}$ in some sense.

Example 3.4 Let X and Y be disjoint vertex sets, and k, n and δ be positive integers satisfying $n > k > 0$ and $\delta \leq \min\{|X|, |Y|\}$. Define $\mathcal{B}(\delta, k)$ to be the family of bipartite graphs of the form $G[X, Y]$ with $n = |X| + |Y|$, satisfying each of the following:

- (H1) for any $v \in X$, $d(v)$ is a constant,
- (H2) for any $v \in Y$, $d(v) = \delta$,
- (H3) $|Y| = |X| + k$.

By definition, the complete bipartite graph $K_{\delta, \delta+k} \in \mathcal{B}(\delta, k)$.

Lemma 3.5 For any $H^* \in \mathcal{B}(\delta, k)$, each of the following holds.

- (i) $\mu_f(H^*) = \alpha'(H^*) = |X| = \frac{n-k}{2}$.
- (ii) $a = 0$ or 1 , $\lambda_1(aD(H^*) + A(H^*)) = \varphi(a, n, \delta, k)$.
- (iii) $a \in (0, 1)$ or $(1, +\infty)$, $\lambda_1(aD(H^*) + A(H^*)) > \varphi(a, n, \delta, k)$.

Proof. Let d be the degree of each vertex in X . By (H1) and (H2), $d|X| = |E(H^*)| = \delta|Y| = \delta(|X| + k)$, leading to $d > \delta \geq 1$. For any subset $S \subseteq X$, $|N_{H^*}(S)| = d|S| > |S|$. It follows by Hall's Theorem (Theorem 16.4 of [4]) that H^* has a matching M of size $|X|$. Since H^* is bipartite, by Theorem 1.1(ii), we conclude that $\mu_f(H^*) = \alpha'(H^*) = |X| = \frac{n-k}{2}$, and so (i) follows.

We prove both (ii) and (iii) simultaneously. The quotient matrix $R(aD(H^*) + A(H^*))$ of $aD(H^*) + A(H^*)$ with respect to the partitions X and Y of $V(H^*)$ can be written as

$$R(aD(H^*) + A(H^*)) = \begin{pmatrix} \frac{a\delta|Y|}{|X|} & \frac{\delta|Y|}{|X|} \\ \delta & a\delta \end{pmatrix}.$$

As the characteristic polynomial of $R(aD(H^*) + A(H^*))$ is

$$\lambda^2 - a\delta\left(1 + \frac{|Y|}{|X|}\right)\lambda + (a^2 - 1)\delta^2\frac{|Y|}{|X|} = 0,$$

direct computation indicates that

$$\begin{aligned}\lambda_1(R(aD(H^*) + A(H^*))) &= \frac{1}{2}\left\{a\delta\left(\frac{|Y|}{|X|} + 1\right) + \sqrt{a^2\delta^2\left(\frac{|Y|}{|X|} + 1\right)^2 - 4(a^2 - 1)\delta^2\frac{|Y|}{|X|}}\right\} \\ &= \frac{1}{2}\left\{a\delta\left(\frac{|Y|}{|X|} + 1\right) + \sqrt{(a^2 - 1)\delta^2\left(\frac{|Y|}{|X|} - 1\right)^2 + \delta^2\left(\frac{|Y|}{|X|} + 1\right)^2}\right\}.\end{aligned}\quad (7)$$

As the partition is equitable and $aD(H^*) + A(H^*)$ is nonnegative, it follows by Theorem 2.4 that $\lambda_1(aD(H^*) + A(H^*)) = \lambda_1(R(aD(H^*) + A(H^*)))$. This, together with (7), leads to the following observations.

(A) If $a = 0$, then $\lambda_1(A(H^*)) = \delta\sqrt{\frac{|Y|}{|X|}} = \delta\sqrt{1 + \frac{k}{|X|}} = \delta\sqrt{1 + \frac{2k}{n-k}} = \varphi(0, n, \delta, k)$.

(B) If $a = 1$, then $\lambda_1(D(H^*) + A(H^*)) = \delta\left(\frac{|Y|}{|X|} + 1\right) = \frac{2n\delta}{n-k} = \varphi(1, n, \delta, k)$.

(C) If $0 < a < 1$, then $a^2 - 1 < 0$, and so

$$\begin{aligned}\lambda_1(aD(H^*) + A(H^*)) &> \frac{1}{2}\left\{a\delta\left(\frac{|Y|}{|X|} + 1\right) + \sqrt{(a^2 - 1)\delta^2\left(\frac{|Y|}{|X|} + 1\right)^2 + \delta^2\left(\frac{|Y|}{|X|} + 1\right)^2}\right\} \\ &= a\delta\left(\frac{|Y|}{|X|} + 1\right) = \frac{2a\delta n}{n-k} = \varphi(a, n, \delta, k).\end{aligned}$$

(D) If $1 < a < +\infty$, then $a^2 - 1 > 0$, and so

$$\begin{aligned}\lambda_1(aD(H^*) + A(H^*)) &> \frac{1}{2}\left\{a\delta\left(\frac{|Y|}{|X|} + 1\right) + \sqrt{(a^2 - 1)\delta^2\left(\frac{|Y|}{|X|} - 1\right)^2 + \delta^2\left(\frac{|Y|}{|X|} - 1\right)^2}\right\} \\ &= a\delta\left(\frac{|Y|}{|X|} + 1\right) = \frac{a\delta(n+k)}{n-k} = \varphi(a, n, \delta, k).\end{aligned}$$

These observations (A)-(D) complete the proof of the lemma. \blacksquare

Remark 3.6 By Lemma 3.5, there exist graphs $H^* \in \mathcal{B}(\delta, k)$ with minimum degree δ such that $\lambda_1(A(H^*)) = \varphi(0, n, \delta, k)$, $\mu_f(H^*) = \frac{n-k}{2}$ and $\lambda_1(D(H^*) + A(H^*)) = \varphi(1, n, \delta, k)$, $\mu_f(H^*) = \frac{n-k}{2}$. Hence the upper bound in Theorem 1.3 is best possible when $a \in \{0, 1\}$.

3.3 The Proof and Discussion of Theorem 1.4

By contradiction, assume that $\mu_f(G) \leq \frac{n-k}{2}$. By (1), there exists a vertex subset $S \subseteq V(G)$ such that $i(G-S) - |S| \geq k$. Let T be the set of isolated vertices in $G-S$. Let $|S| = s$ and $|T| = t$. Then $s+t \leq n$ and $t = i(G-S) \geq s+k$. As $\cup_{v \in T} N_G(v) \subseteq S$, we observe that $s \geq \delta$, and so $t \geq s+k \geq \delta+k$. As $K_t \cup (n-t)K_1$ is a spanning subgraph of \overline{G} , it follows by Theorem 2.5 that

$$\begin{aligned}\lambda_1(aD(\overline{G}) + A(\overline{G})) &\geq \lambda_1(aD(K_t \cup (n-t)K_1) + A(K_t \cup (n-t)K_1)) \\ &= (a+1)(t-1) \geq (a+1)(\delta+k-1),\end{aligned}$$

leading to a contradiction to the hypothesis. This completes the proof of Theorem 1.4. \blacksquare

Example 3.7 By definition, the complete bipartite graph $K_{\delta, \delta+k} \in \mathcal{B}(\delta, k)$. Direct computation yields that $\lambda_1(aD(\overline{K_{\delta, \delta+k}}) + A(\overline{K_{\delta, \delta+k}})) = (a+1)(\delta+k-1)$, and $\mu_f(K_{\delta, \delta+k}) = \mu(K_{\delta, \delta+k}) = \delta = \frac{n-k}{2}$. In this sense that the upper bound in Theorem 1.4 is best possible.

4 Corollaries of Theorems 1.3 and 1.4

Throughout this section, we assume that a , b and k are real numbers with $a \geq 0$, $b > 0$ and $k \in (0, n)$. Again, the function $\varphi(a, n, \delta, k)$ is defined in (3). We start by considering the matrix $J(G : a, b) = aD(G) + bA(G)$. Theorems 1.3 and 1.4 have the following seemingly more general forms.

Corollary 4.1 Let G be a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$. If $\lambda_1(aD(G) + bA(G)) < b\varphi(\frac{a}{b}, n, \delta, k)$, then $\mu_f(G) > \frac{n-k}{2}$.

Proof. If $\mu_f(G) \leq \frac{n-k}{2}$, then by Theorem 1.3, $\lambda_1(aD(G) + bA(G)) \geq \varphi(a, n, \delta, k)$. As $aD(G) + bA(G) = b(\frac{a}{b}D(G) + A(G))$, it follows that $\lambda_1(aD(G) + bA(G)) = b\lambda_1(\frac{a}{b}D(G) + A(G)) \geq b\varphi(\frac{a}{b}, n, \delta, k)$, a contradiction. ■

Corollary 4.2 Let G be a connected simple graph of order n with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$. If $\lambda_1(aD(\overline{G}) + bA(\overline{G})) < (a+b)(\delta+k-1)$, then $\mu_f(G) > \frac{n-k}{2}$.

Proof. If $\mu_f(G) \leq \frac{n-k}{2}$, then by Theorem 1.4, $\lambda_1(aD(\overline{G}) + bA(\overline{G})) \geq (a+b)(\delta+k-1)$. As $aD(\overline{G}) + bA(\overline{G}) = b(\frac{a}{b}D(\overline{G}) + A(\overline{G}))$, it follows that $\lambda_1(aD(\overline{G}) + bA(\overline{G})) = b\lambda_1(\frac{a}{b}D(\overline{G}) + A(\overline{G})) \geq b(\frac{a}{b} + 1)(\delta+k-1) = (a+b)(\delta+k-1)$, a contradiction. ■

Finally we observe that Corollary 1.5 follows by letting $a = b = 1$ in Corollaries 4.1 and 4.2, and Corollary 1.6 can be obtained from Corollaries 4.1 and 4.2 by setting $a = \alpha$ and $b = 1 - \alpha$.

4.1 Fractional matching number and eigenvalues

We present an application of Theorem 1.3 in this subsection by showing a relationship between $\mu_f(G)$ and $\lambda_1(aD(G) + A(G))$ for a graph G . Theorem 4.3(i) was proved by O in [18], and is included here for the sake of completeness.

Theorem 4.3 Let G be a connected simple graph with $n = |V(G)|$ and $\delta = \delta(G) \leq \frac{n-k}{2}$. Each of the following holds.

- (i) (O [18]) If $a = 0$, then $\mu_f(G) \geq \frac{n\delta^2}{\lambda_1(G)^2 + \delta^2}$, where the equality holds if and only if $G \in \mathcal{B}(\delta, k)$ and $k = \frac{n(\lambda_1(G)^2 - \delta^2)}{\lambda_1(G)^2 + \delta^2}$ is an integer.
- (ii) If $a = 1$, then $\mu_f(G) \geq \frac{\delta n}{\lambda_1(D(G) + A(G))}$, where the equality holds if and only if $G \in \mathcal{B}(\delta, k)$ and $k = n - \frac{2\delta n}{\lambda_1(D(G) + A(G))}$ is an integer.
- (iii) If $0 < a < 1$, then $\mu_f(G) \geq \frac{a\delta n}{\lambda_1(aD(G) + A(G))}$. Furthermore, the equality holds if $G \in \mathcal{B}(\delta, k)$ and $k = n - \frac{2a\delta n}{\lambda_1(aD(G) + A(G))}$ is an integer.

(iv) If $1 < a < +\infty$, then $\mu_f(G) \geq \frac{a\delta n}{\lambda_1(aD(G)+A(G))+a\delta}$, where the equality holds if $G \in \mathcal{B}(\delta, k)$ and $k = \frac{n(\lambda_1(aD(G)+A(G))-a\delta)}{\lambda_1(aD(G)+A(G))+a\delta}$ is an integer.

Proof. We only prove Theorem 4.3 (ii)-(iv). Throughout the proof, we denote $\lambda_1(aD(G)+A(G))$ and $\mu_f(G)$ by λ_1 and μ_f , respectively. Theorem 1.3 says that if $\lambda_1 < \varphi(a, n, \delta, k)$, then $\mu_f > \frac{n-k}{2}$. Theorem 4.3 follows by applying Theorem 1.3 to each of the following possible values of the number a .

Assume that $0 < a \leq 1$. Define $k_1 = n - \frac{2a\delta n}{\lambda_1}$. As $\frac{1}{n-x}$ is an increasing function of x on $(0, n)$, $\frac{2a\delta n}{n-k}$ decreases towards λ_1 as k decreases towards k_1 , and $\lim_{k \rightarrow k_1} \frac{2a\delta n}{n-k} = \lambda_1$. By Theorem 1.3, $\mu_f > \frac{n-k}{2}$ for each value of $k \in (k_1, n)$. It follows that $\mu_f \geq \lim_{k \rightarrow k_1} \frac{n-k}{2} = \frac{n-k_1}{2} = \frac{a\delta n}{\lambda_1}$. Finally, if $G \in \mathcal{B}(\delta, k)$ and $k = n - \frac{2a\delta n}{\lambda_1(aD(G)+A(G))}$ is an integer, then direct computation yields $\mu_f(G) = \alpha'(G) = \frac{n-k}{2} = \frac{a\delta n}{\lambda_1(aD(G)+A(G))}$. This proves Theorem 4.3(iii) and lower bound part of Theorem 4.3(ii).

Now suppose that $a = 1$ and that $\mu_f(G) = \frac{\delta n}{\lambda_1(D(G)+A(G))}$. To simplify the notation and terms, we shall adopt the definition of the vertex sets S and T , and the definition of the graph H in Subsection 3.1, with $s = |S|$ and $t = |T|$. As $\mu_f(G) = \frac{\delta n}{\lambda_1(D(G)+A(G))}$, every inequality in the arguments of the previous paragraph must be an equality, and so $k = k_1$ and $\lambda_1 = \frac{2\delta n}{n-k}$. Furthermore, every inequality in (6) in Subsection 3.1 must also be an equality. Hence in (6) we must have $a\delta(1 + \frac{t}{s}) = a\delta(2 + \frac{k}{s})$, leading to $k = t - s$, and so k is an integer. Likewise, in (6) we must also have $2 + \frac{k}{s} = \frac{2n}{n-k}$. This, together with $k = t - s$, implies $n = s + t$. It follows that $a = t\delta, t = k + s$. By Theorem 2.3, the partition is equitable, and thus $G \cong H \in \mathcal{B}(\delta, k)$. This completes the proof of Theorem 4.3(ii).

Finally we assume that $a \in (1, +\infty)$. Define $k_2 = \frac{n(\lambda_1 - a\delta)}{\lambda_1 + a\delta}$. As $\frac{n+x}{n-x}$ is an increasing function of x on $(0, n)$, $\frac{a\delta(n+k)}{n-k}$ decreases towards λ_1 as k decreases towards k_2 and $\lim_{k \rightarrow k_2} \frac{a\delta(n+k)}{n-k} = \lambda_1$. By Theorem 1.3, $\mu_f > \frac{n-k}{2}$ for each value of $k \in (k_2, n)$, and so $\mu_f \geq \lim_{k \rightarrow k_2} \frac{n-k}{2} = \frac{n-k_2}{2} = \frac{a\delta n}{\lambda_1 + a\delta}$. If $G \in \mathcal{B}(\delta, k)$ with $k = \frac{n(\lambda_1(aD(G)+A(G))-a\delta)}{\lambda_1(aD(G)+A(G))+a\delta}$ being an integer, then direct computation yields that $\mu_f(G) = \alpha'(G) = \frac{n-k}{2} = \frac{a\delta n}{\lambda_1(aD(G)+A(G))+a\delta}$. This justifies Theorem 4.3(iv), and completes the proof of the theorem. ■

4.2 Fractional perfect matchings and eigenvalues

The following Corollaries 4.4 and 4.5 are immediate consequences of Theorems 1.3 and 1.4, respectively. Letting $k = 1$ in Theorems 1.3 and 1.4, respectively, we conclude that the hypotheses of Corollaries 4.4 and 4.5 respectively imply that $\mu_f(G) > \frac{n-1}{2}$. By Theorem 1.1, $2\mu_f(G)$ is an integer and $\mu_f(G) \leq \frac{n}{2}$. This forces that $\mu_f(G) = \frac{n}{2}$, and so G has a fractional perfect matching.

Corollary 4.4 *Let a be a real number with $a \geq 0$, and G be a simple connected graph of order*

n with minimum degree $\delta \leq \frac{n-1}{2}$. If

$$\lambda_1(aD(G) + A(G)) < \varphi(a, n, \delta, 1),$$

then G has a fractional perfect matching.

Corollary 4.5 *Let a be a real number with $a \geq 0$, G be a connected graph of order n with minimum degree $\delta \leq \frac{n-1}{2}$, and \overline{G} be the complement of G . If*

$$\lambda_1(aD(\overline{G}) + A(\overline{G})) < (a+1)\delta,$$

then G has a fractional perfect matching.

As it is shown in Example 3.7, the upper bound in Corollary 4.5 is best possible. This upper bound in Corollary 4.5 can be slightly improved by excluding some graphs. Following [4], the **join** of two graphs G_1 and G_2 , denoted $G_1 \vee G_2$, is the graph formed from the disjoint union of G_1 and G_2 by adding new edges joining every vertex of G_1 to every vertex of G_2 .

Theorem 4.6 *Let a be a real number with $a \geq 0$, G be a connected graph with $n = |V(G)|$, and $\delta = \delta(G) \leq \frac{n-1}{2}$. If*

$$\lambda_1(aD(\overline{G}) + A(\overline{G})) < (a+1)(\delta+1), \quad (8)$$

then G has a fractional perfect matching unless $G \cong (\delta+1)K_1 \vee H_\delta$, where H_δ is a simple graph of order δ .

Proof. By contradiction, assume that G has no fractional perfect matchings. By (2), there exists a vertex set $S \subseteq V(G)$ such that $i(G-S) - |S| > 0$. Let T be the set of isolated vertices in $G-S$. Then $\cup_{v \in T} N_G(v) \subseteq S$, and so $|S| \geq \delta$. This implies that $|T| \geq |S| + 1 \geq \delta + 1$.

We claim that $V(G) = S \cup T$. By contradiction, assume that there exists a vertex $v \in V(G) - S \cup T$. Then we have $K_{\delta+2} \subseteq \overline{G}[T \cup \{v\}] \subseteq \overline{G}$. By Theorem 2.5,

$$\begin{aligned} \lambda_1(aD(\overline{G}) + A(\overline{G})) &\geq \lambda_1(aD(K_{\delta+2} \cup (n-\delta-2)K_1) + A(K_{\delta+2} \cup (n-\delta-2)K_1)) \\ &= \lambda_1(aD(K_{\delta+2}) + A(K_{\delta+2})) = (a+1)(\delta+1), \end{aligned}$$

contrary to (8).

Furthermore, we claim that $|T| = \delta + 1$. Assume to the contrary that $|T| \geq \delta + 2$. Then \overline{G} contains a clique of order $\delta + 2$, and so $\lambda_1(aD(\overline{G}) + A(\overline{G})) \geq (a+1)(\delta+1)$, contrary to (8) again. Hence $|T| = \delta + 1$, and so $|S| = \delta$. It follows that $G \cong (\delta+1)K_1 \vee H_\delta$. As there exists a vertex subset $V(H_\delta) \subseteq V(G)$ such that $i(G - V(H_\delta)) > |V(H_\delta)|$, by (2), $(\delta+1)K_1 \vee H_\delta$ has no fractional perfect matchings. This completes the proof. \blacksquare

Acknowledgement.

The research of Ruifang Liu is supported by NSFC (No. 11971445), Natural Science Foundation of Henan Province (No. 20A110005). The research of Hong-Jian Lai is supported by NSFC (Nos. 11771039 and 11771443).

References

- [1] A. Berman, R.J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, Academic Press, New York, 1979.
- [2] A.E. Brouwer, W.H. Haemers, Eigenvalues and perfect matchings, *Linear Algebra Appl.* 395 (2005) 155-162.
- [3] A.E. Brouwer, W.H. Haemers, *Spectra of Graphs*, Springer, New York, 2012.
- [4] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, New York, 2008.
- [5] S.M. Cioabă, Perfect matchings, eigenvalues and expansion, *C. R. Math. Acad. Sci. Soc. R. Can.* 27 (2005) 101-104.
- [6] S.M. Cioabă, D.A. Gregory, Large matchings from eigenvalues, *Linear Algebra Appl.* 422 (2007) 308-317.
- [7] S.M. Cioabă, D.A. Gregory, W.H. Haemers, Matchings in regular graphs from eigenvalues, *J. Combin. Theory Ser. B* 99 (2009) 287-297.
- [8] G.A. Dirac, Some theorems on abstract graphs, *Proc. London Math. Soc* 3 (1952) 69-81.
- [9] E.V. Haynsworth, Applications of a theorem on partitioned matrices, *J. Res. Natl. Bur. Stand.* 62 (1959) 73-78.
- [10] W.H. Haemers, Interlacing eigenvalues and graphs, *Linear Algebra Appl.* 226-228 (1995) 593-616.
- [11] Y.M. Hong, X.F. Gu, Q.H. Liu and H.-J. Lai, Fractional spanning tree packing, forest covering and eigenvalues, *Discrete Applied Math.* 213 (2016) 219-223.
- [12] R.A. Horn, C.R. Johnson, *Matrix Analysis*, Cambridge University Press, 1986.
- [13] Q.H. Liu, Y.M. Hong and H.-J. Lai, Edge-disjoint spanning trees and eigenvalues, *Linear Algebra Appl.* 444 (2014) 146-151.
- [14] Q. Liu, Y. Hong, X. Gu and H.-J. Lai, Note on edge-disjoint spanning trees and eigenvalues, *Linear Algebra Appl.* 458 (2014) 128-133.
- [15] R.F. Liu, H.-J. Lai, Y.Z. Tian, Spanning tree packing number and eigenvalues of graphs with given girth, *Linear Algebra Appl.* 578 (2019) 411-424.
- [16] R.F. Liu, H.-J. Lai, Y.Z. Tian, Y. Wu, Vertex-connectivity and eigenvalues of graphs with fixed girth, *Appl. Math. Comput.* 344-345 (2019) 141-149.
- [17] V. Nikiforov, Merging the A - and Q -spectral theories, *Appl. Anal. Discrete Math.* 11 (2017) 81-107.
- [18] S. O, Spectral radius and fractional matchings in graphs, *European J. Combin.* 55 (2016) 144-148.
- [19] S. O, S.M. Cioabă, Edge-connectivity, eigenvalues, and matchings in regular graphs, *SIAM J. Discrete Math.* 24 (2010) 1470-1481.
- [20] E.R. Scheinerman, D.H. Ullman, *Fractional Graph Theory: A Rational Approach to the Theory of Graphs*, Wiley & Sons, 1997.
- [21] J. Xue, M.Q. Zhai, J.L. Shu, Fractional matching number and eigenvalues of a graph, *Linear Multilinear Algebra*, <https://doi.org/10.1080/03081087.2018.1498059>, 2018.
- [22] L.H. You, M. Yang, W. So, W.G. Xi, On the spectrum of an equitable quotient matrix and its application, *Linear Algebra Appl.* 577 (2019) 21-40.