

EXCLUDING A LADDER

TONY HUYNH, GWENAËL JORET, PIOTR MICEK, MICHAŁ T. SEWERYN,
AND PAUL WOLLAN

ABSTRACT. A ladder is a $2 \times k$ grid graph. When does a graph class \mathcal{C} exclude some ladder as a minor? We show that this is the case if and only if all graphs G in \mathcal{C} admit a proper vertex coloring with a bounded number of colors such that for every 2-connected subgraph H of G , there is a color that appears exactly once in H . This type of vertex coloring is a relaxation of the notion of centered coloring, where for every connected subgraph H of G , there must be a color that appears exactly once in H . The minimum number of colors in a centered coloring of G is the treedepth of G , and it is known that classes of graphs with bounded treedepth are exactly those that exclude a fixed path as a subgraph, or equivalently, as a minor. In this sense, the structure of graphs excluding a fixed ladder as a minor resembles the structure of graphs without long paths. Another similarity is as follows: It is an easy observation that every connected graph with two vertex-disjoint paths of length k has a path of length $k + 1$. We show that every 3-connected graph which contains as a minor a union of sufficiently many vertex-disjoint copies of a $2 \times k$ grid has a $2 \times (k + 1)$ grid minor.

Our structural results have applications to poset dimension. We show that posets whose cover graphs exclude a fixed ladder as a minor have bounded dimension. This is a new step towards the goal of understanding which graphs are unavoidable as minors in cover graphs of posets with large dimension.

(T. Huynh) SCHOOL OF MATHEMATICS, MONASH UNIVERSITY, MELBOURNE, AUSTRALIA

(G. Joret) COMPUTER SCIENCE DEPARTMENT, UNIVERSITÉ LIBRE DE BRUXELLES, BRUSSELS, BELGIUM

(P. Micek, M.T. Seweryn) THEORETICAL COMPUTER SCIENCE DEPARTMENT, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, JAGIELLONIAN UNIVERSITY, KRAKÓW, POLAND

(P. Wollan) DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF ROME “LA SAPIENZA”, ROME, ITALY

E-mail addresses: tony.bourbaki@gmail.com, gjoret@ulb.ac.be, piotr.micek@tcs.uj.edu.pl, michal.seweryn@tcs.uj.edu.pl, wollan@di.uniroma1.it.

Date: January 20, 2022.

2010 Mathematics Subject Classification. 05C83, 06A07.

Key words and phrases. graph minor, ladder, treedepth, poset dimension.

T. Huynh is supported by the Australian Research Council. G. Joret is supported by an ARC grant from the Wallonia-Brussels Federation of Belgium. P. Micek was partially supported by the National Science Center of Poland under grant no. 2015/18/E/ST6/00299. M.T. Seweryn was partially supported by Kartezjusz program WND-POWR.03.02.00-00-1001/16-01 funded by The National Center for Research and Development of Poland.

1. INTRODUCTION

Graphs with no long paths are relatively well understood. In particular, if a graph G does not contain a path on $k + 1$ vertices as a subgraph, then G has a centered coloring with at most k colors. Conversely, if G has a centered coloring with at most k colors, then G does not contain a path on 2^k vertices. Here, a *centered coloring* of G is a vertex coloring of G such that for every connected subgraph H of G , some color is assigned to exactly one vertex of H . The minimum number of colors used in a centered coloring of G is known as the *treedepth* of G , denoted $\text{td}(G)$.

In this paper, we show an analogous result for graphs excluding a fixed ladder as a minor. We show that such graphs can be characterized as graphs that admit a 2-connected centered coloring with a bounded number of colors. Here, a *2-connected centered coloring* of a graph G is a vertex coloring of G such that for every connected subgraph H of G having no cutvertex, some color is assigned to exactly one vertex of H .¹ The minimum number of colors in a 2-connected centered coloring of G is denoted $\text{td}_2(G)$.

Before stating our theorem formally, we introduce a related type of coloring. A *cycle centered coloring* of G is a vertex coloring of G such that for every subgraph H of G which is an edge or a cycle, some color is assigned to exactly one vertex of H . The minimum number of colors in a cycle centered coloring of G is denoted $\chi_{\text{cc}}(G)$. While every 2-connected centered coloring of a graph is cycle centered, the converse is not necessarily true.

Let L_k denote the ladder with k rungs (that is, the $2 \times k$ grid graph). Our theorem for graphs excluding a ladder is as follows.

Theorem 1. *For every class \mathcal{C} of graphs, the following properties are equivalent.*

- (i) *There exists an integer $k \geq 1$ such that no graph in \mathcal{C} has an L_k minor.*
- (ii) *There exists an integer $m \geq 1$ such that $\text{td}_2(G) \leq m$ for every graph G in \mathcal{C} .*
- (iii) *There exists an integer $c \geq 1$ such that $\chi_{\text{cc}}(G) \leq c$ for every graph G in \mathcal{C} .*

A second contribution of this paper is as follows. As is well known, every pair of longest paths in a connected graph intersect, or equivalently, if a connected graph contains two vertex disjoint paths of order k , then it contains a path of order $k + 1$. We show a generalization of this statement where paths are replaced with ladders, and ‘two’ with ‘many’.

Theorem 2 (Bumping a ladder). *For every integer $k \geq 1$ there exists an integer $N \geq 1$ with the following property. Every 3-connected graph G containing a union of N vertex-disjoint copies of L_k as a minor contains L_{k+1} as a minor.*

Let us point out the following corollary of Theorem 2. Robertson and Seymour [17] proved that for every fixed planar graph H and every $N \geq 1$, there exists $N' \geq 1$ such that every graph G not containing a union of N

¹We remark that this definition is slightly different than the one given in the abstract but is equivalent. Indeed, if H is connected with no cutvertex then H is either a vertex, an edge, or is 2-connected; edges will make sure that the coloring is proper.

vertex-disjoint copies of H as a minor has a vertex subset X with $|X| \leq N'$ such that $G - X$ has no H minor.

Corollary 3. *For every integer $k \geq 1$ there exists an integer $N' \geq 1$ with the following property. Every 3-connected graph G with no L_{k+1} minor has a vertex subset X with $|X| \leq N'$ such that $G - X$ has no L_k minor.*

We remark that 3-connectivity in Theorem 2 is necessary. See Figure 1. On the other hand, we expect that the dependence on k is not. We conjecture that there exists a constant N_0 such that for every k , Theorem 2 holds true with $N = N_0$. For all we know, this might even be true with $N_0 = 4$.




FIGURE 1. A 2-connected graph with 8 disjoint copies of L_4 and no L_5 -minor.

We conclude this introduction with an application of our results to poset dimension. Let P be a poset. For two elements x and y of P , we say that y *covers* x if $x < y$ in P and there is no element z in P such that $x < z < y$ in P . The *cover graph* of a poset P is the graph on the ground set of P in which two vertices are adjacent if one of them covers the other in P . Informally, the cover graph of P is its Hasse diagram seen as an undirected graph. A *realizer* of P is a set $\{\leq_1, \dots, \leq_d\}$ of linear orders on the ground set of P such that for any two elements x and y of P , we have $x \leq y$ in P if and only if $x \leq_i y$ for every $i \in \{1, \dots, d\}$. The *dimension* of P , denoted $\dim(P)$, is the least size of a realizer of P .

We prove the following result, which shows the relevance of 2-connected centered colorings for studying poset dimension.

Theorem 4. *Let P be a poset with cover graph G and let $m = \text{td}_2(G)$. Then, P has dimension at most $2^{m+1} - 2$.*

When combined with Theorem 1, the above theorem implies the following result. Proving this result was one of our motivations for studying the structure of graphs with no long ladder minor.

Corollary 5. *For every integer $k \geq 1$ there exists an integer $d \geq 1$ such that every poset whose cover graph excludes L_k as a minor has dimension at most d .*

Let us say that a graph H is *unavoidable* if the cover graph of every poset with large enough dimension contains H as a minor. Corollary 5 shows that




FIGURE 2. (Left) Hasse diagram of the poset of Kelly's construction of order k for $k = 4$. Its definition for an arbitrary order k can be inferred from the figure. (Right) A free redrawing of its cover graph.




FIGURE 3.

every ladder is unavoidable. Note that the class of unavoidable graphs is closed under taking minors (thus, fans are also unavoidable, etc.). It is an open problem to obtain a full characterization of unavoidable graphs. Besides ladders, the only other positive result known is that K_4 is unavoidable [10, 18]. As for negative results, a classic construction of Kelly [13], see Figure 2, shows that there are posets with unbounded dimension whose cover graphs are planar and have pathwidth 3. Note that every unavoidable graph must necessarily be a minor of some graph from Kelly's construction. We conjecture that this is precisely the characterization of unavoidable graphs.

Conjecture 6. *A graph H is unavoidable if and only if H is a minor of some graph from Kelly's construction.*

Let us point out the following equivalent reformulation of Conjecture 6: A graph H is unavoidable if and only if H is a minor of some graph obtained by gluing copies of K_4 s along edges in a path-like way, and subdividing all horizontal edges of the K_4 s once; see Figure 3. Indeed, every graph from Kelly's construction is a minor of a graph in the latter family (see Figure 2 (right)), and vice versa.

We end with a short description of related results about poset dimension. Unavoidable graphs are very restricted in nature, since they must be minors of some graph from Kelly's construction. It is thus natural to consider posets excluding some fixed but arbitrary graph H as a minor in their cover graphs, and ask: What kind of unavoidable structure can be found in these posets when dimension is large? Several results have been obtained in the last decade when considering long chains as unavoidable structure. Recall that a *chain* in a poset is a set of pairwise comparable elements. The maximum length of a chain is the *height* of the poset. In general, dimension is not bounded by any function of the height since there are height-2 posets with arbitrarily large

dimension. However, in 2014 Streib and Trotter [19] proved the surprising result that, if the cover graph of a poset is planar, then the poset's dimension is bounded from above by a function of its height (see [4, 11, 14] for improved bounds). This property was then shown to be true more generally if the cover graph excludes a fixed apex graph as a minor [9]. Then Walczak [21] showed that this remains true if any fixed graph H is excluded from the cover graph as a minor, or even as a topological minor (see [15] for a short proof). This was in turn generalized further: Posets with cover graphs belonging to a fixed graph class \mathcal{C} with bounded expansion have dimension bounded from above by a function of their height [12]. A simple proof using a family of graph invariants called weak coloring numbers was given in [8]. Finally, it was shown in [8] that nowhere dense graph classes² can be characterized in terms of dimension of bounded-height posets with cover graphs in the class.

Thus, long chains are unavoidable in posets with large dimension and well-behaved sparse cover graphs. In the case of posets with planar cover graphs, Howard, Streib, Trotter, Walczak, and Wang [7] strengthened this further by showing that two long incomparable chains can be found when dimension is large. They conjectured that this remains true more generally when any fixed graph H is forbidden as a minor from the cover graph.

The paper is organized as follows. We introduce some basic definitions and notations in Section 2. Then we prove Theorem 1 in Section 3 and apply it in Section 4 to prove Theorem 4 about poset dimension. Finally, we prove Theorem 2 in Section 5.

2. PRELIMINARIES

In this section we recall some standard definitions. For a graph G and a subset X of vertices, we denote by $G - X$ the subgraph of G induced on the set of vertices $V(G) \setminus X$. For a vertex $x \in V(G)$, we will use $G - x$ as shorthand notation for $G - \{x\}$. A *cutvertex* of a graph G is a vertex v of G such that $G - v$ has more connected components than G . A *separation* of a graph G is a pair (A_1, A_2) of vertex subsets in G such that $A_1 \cup A_2 = V(G)$ and every edge of G has both endpoints in A_1 or in A_2 . The *order* of the separation is the number $|A_1 \cap A_2|$, and a *k-separation* is a separation of order at most k . A separation (A_1, A_2) of G is *trivial* if $A_1 = V(G)$ or $A_2 = V(G)$, and *nontrivial* otherwise. For $k \geq 1$, we say that a graph is *k-connected* if it has at least $k + 1$ vertices and does not admit a nontrivial $(k - 1)$ -separation.

A *subdivision* of a graph H is a graph obtained by replacing some edges of H with new paths between their endpoints such that none of the paths has an inner vertex in $V(H)$ or on another new path. An *H-model* in a graph G is a function ϕ which assigns to each vertex $u \in V(H)$ a connected subgraph $\phi(u)$ of G , such that

- (i) the graphs $\phi(u)$ and $\phi(v)$ are vertex-disjoint for distinct vertices u and v of H , and
- (ii) G has an edge between $\phi(u)$ and $\phi(v)$ for every edge $uv \in E(H)$.

²See Nešetřil and Ossona de Mendez [16] for background on classes with bounded expansion and on nowhere dense classes.

A graph H is a *minor* of a graph G if and only if there is an H -model in G .

For two vertex subsets A and B in a graph G , an A - B *path* is a path with one endpoint in A and the second in B , and with no internal vertex in $A \cup B$. For two vertices a and b of G , by an a - b *path* we mean an $\{a\}$ - $\{b\}$ path. If u and v are vertices on a path P , then the only u - v path in P is denoted by uPv .

For a positive integer k , the *ladder* L_k is the graph with vertex set $\{1, 2\} \times \{1, \dots, k\}$ in which two vertices (i, j) and (i', j') are adjacent if $|i - i'| + |j - j'| = 1$. Since the maximum degree of L_k is at most 3, the ladder L_k is a minor of a graph G if and only if G has a subgraph isomorphic to a subdivision of L_k .

If (z_1, z_2) is a pair of vertices in a graph H and ϕ is an L_k -model in H such that $z_1 \in V(\phi((1, k)))$ and $z_2 \in V(\phi((2, k)))$, then we say that ϕ is *rooted* at the pair (z_1, z_2) .

3. 2-CONNECTED CENTERED COLORINGS

The goal of this section is to prove Theorem 1. We start by establishing some basic properties of 2-connected centered colorings.

Lemma 7. *For every graph G and subset $X \subseteq V(G)$ we have*

$$\text{td}_2(G - X) \geq \text{td}_2(G) - |X|.$$

Proof. Let φ be a 2-connected centered coloring of $G - X$ using at most $\text{td}_2(G - X)$ colors, and extend it to a coloring φ' of G using at most $\text{td}_2(G - X) + |X|$ colors by assigning new distinct colors to the vertices of X . Consider a connected subgraph H of G which does not have a cutvertex. If H contains a vertex from X , then the color of that vertex is unique in G and thus in H . If $H \subseteq G - X$, then some color is assigned to exactly one vertex of H because φ is a 2-connected centered coloring of $G - X$. Hence φ' is a 2-connected centered coloring of G using $\text{td}_2(G - X) + |X|$ colors, and therefore $\text{td}_2(G) \leq \text{td}_2(G - X) + |X|$, so indeed $\text{td}_2(G - X) \geq \text{td}_2(G) - |X|$. \square

Lemma 8. *If (A_1, A_2) is a separation of order ≤ 1 of a graph G , then*

$$\text{td}_2(G) = \max\{\text{td}_2(G[A_1]), \text{td}_2(G[A_2])\}.$$

Proof. If φ is a 2-connected centered coloring of G , then the restrictions of φ to A_1 and A_2 are 2-connected centered colorings of $G[A_1]$ and $G[A_2]$, respectively, and these colorings use no more colors than φ . Therefore $\text{td}_2(G[A_i]) \leq \text{td}_2(G)$ for $i \in \{1, 2\}$.

Let $m = \max\{\text{td}_2(G[A_1]), \text{td}_2(G[A_2])\}$, and let $\varphi_1: A_1 \rightarrow \{1, \dots, m\}$ and $\varphi_2: A_2 \rightarrow \{1, \dots, m\}$ be 2-connected centered colorings of $G[A_1]$ and $G[A_2]$, respectively. Since $|A_1 \cap A_2| \leq 1$, after permuting the colors in one of the colorings φ_1 or φ_2 we may assume that they agree on $A_1 \cap A_2$. Let $\varphi: V(G) \rightarrow \{1, \dots, m\}$ be a vertex coloring such that its restriction to A_i is φ_i for $i \in \{1, 2\}$. For every connected subgraph H of G which does not have a cutvertex, the separation $(A_1 \cap V(H), A_2 \cap V(H))$ of H must be trivial, so $V(H) \subseteq A_i$ for some $i \in \{1, 2\}$. Since the restriction of φ to A_i is a 2-connected centered coloring of $G[A_i]$, some vertex of H receives a color not used for any other vertex of H . This proves that φ is a 2-connected centered coloring of G and $\text{td}_2(G) \leq m$ as required. \square

Lemma 9. *Every graph G contains a connected subgraph B without a cutvertex such that $\text{td}_2(B) = \text{td}_2(G)$.*

Proof. We prove the lemma by induction on the size of the graph. If G is connected and does not have a cutvertex, then the lemma holds with $B = G$. Otherwise, let (A_1, A_2) be a nontrivial separation of order ≤ 1 of G . Without loss of generality we assume that $\text{td}_2(G[A_1]) \leq \text{td}_2(G[A_2])$. By Lemma 8, we have $\text{td}_2(G[A_2]) = \text{td}_2(G)$. By induction hypothesis applied to $G[A_2]$ (which is a proper subgraph of G), there is a connected subgraph B of $G[A_2]$ without a cutvertex such that $\text{td}_2(B) = \text{td}_2(G[A_2]) = \text{td}_2(G)$, which completes the proof. \square

We will also need the following classical result by Erdős and Szekeres [3].

Theorem 10 (Erdős-Szekeres Theorem). *Let $k \geq 1$ be an integer, let $n = (k-1)^2 + 1$, and let a_1, \dots, a_n be a sequence of distinct integers. Then there exist integers i_1, \dots, i_k with $1 \leq i_1 < \dots < i_k \leq n$ such that $a_{i_1} < \dots < a_{i_k}$ or $a_{i_1} > \dots > a_{i_k}$.*

Recall that L_k denotes the ladder with k rungs, with vertex set $\{1, 2\} \times \{1, \dots, k\}$, and an L_k -model ϕ is rooted at a pair (z_1, z_2) if $z_i \in V(\phi((i, k)))$ for $i \in \{1, 2\}$.

Lemma 11. *Let $k \geq 1$ and $t \geq 1$ be integers, let $s = (k-1)^2 + 2$, let G be a 2-connected graph, and let x_1 and x_2 be distinct vertices of G . If $\text{td}_2(G) > t \cdot s$, then at least one of the following holds:*

- (i) G has an L_k minor, or
- (ii) G has an L_t -model rooted at the pair (x_1, x_2) .

Proof. We prove the lemma by induction on t . Suppose first $t = 1$. Since G is connected, there is an x_1 - x_2 path in G , and so (ii) holds.

Now suppose that $t \geq 2$. By 2-connectivity of G , there exist two internally disjoint x_1 - x_2 paths P and P' . By Menger's Theorem, either there exist $s+1$ disjoint $V(P)$ - $V(P')$ paths in G , or there exists a set of at most s vertices separating $V(P)$ from $V(P')$.

Case 1: there exist $s+1$ disjoint $V(P)$ - $V(P')$ paths Q_1, \dots, Q_{s+1} . We assume that the paths are listed in the order in which they intersect the path P when traversing it from x_1 to x_2 . Let π be a permutation of $\{1, \dots, s+1\}$ such that $Q_{\pi(1)}, \dots, Q_{\pi(s+1)}$ is the order in which the paths intersect P' when traversing it from x_1 to x_2 . The paths P and P' are internally disjoint, so for $2 \leq i \leq s$, the path Q_i has two distinct endpoints. Consider the sequence $\pi(2), \dots, \pi(s)$ of length $s-1 = (k-1)^2 + 1$. By Theorem 10 applied to that sequence, there exist indices $2 \leq i_1 < \dots < i_k \leq s$ such that either $\pi(i_1) < \dots < \pi(i_k)$, or $\pi(i_1) > \dots > \pi(i_k)$. In both cases G has a subgraph isomorphic to a subdivision of L_k obtained as a union of a subpath of P , a subpath of P' and the paths Q_{i_1}, \dots, Q_{i_k} . Hence (i) is satisfied. See Figure 4.

Case 2: there is a set X of at most s vertices separating $V(P)$ from $V(P')$. See Figure 5. We have $\text{td}_2(G) > t \cdot s$, so by Lemma 7, we also have $\text{td}_2(G - X) \geq \text{td}_2(G) - |X| > ts - |X| \geq ts - s$. Hence by Lemma 9 there exists a connected subgraph B of $G - X$ without a cutvertex




FIGURE 4. Case 1: two ways how the paths P , P' and Q_{i_1}, \dots, Q_{i_k} induce a subdivision of L_k .

such that $\text{td}_2(B) > ts - s$. Since $\text{td}_2(B) > ts - s \geq 2s - s \geq 2$, we have $|V(B)| > 2$. By 2-connectivity of G , there exist two disjoint $\{x_1, x_2\} - V(B)$ paths Q_1 and Q_2 in G with $x_i \in V(Q_i)$ for $i \in \{1, 2\}$. Let y_i denote the endpoint of Q_i in B for $i \in \{1, 2\}$. Since $\text{td}_2(B) > ts - s = (t-1)s$, we can apply induction hypothesis to $k, t-1, B, y_1$ and y_2 . If B has an L_k minor, then so does G , so (i) holds. Otherwise, there is an L_{t-1} -model ϕ' in B such that y_i is in $\phi'((i, t-1))$ for $i \in \{1, 2\}$. Since $B \subseteq G - X$ and the set X separates $V(P)$ from $V(P')$, the model ϕ' intersects at most one of the paths P and P' . Without loss of generality, let us assume that ϕ' does not intersect P . For each $i \in \{1, 2\}$, we have $x_i \in V(P) \cap V(Q_i)$, so $V(P) \cap V(Q_i) \neq \emptyset$. Let R be a $V(Q_1) - V(Q_2)$ subpath of P . Since R is disjoint from the model ϕ' , we can see that (ii) is witnessed by an L_t -model ϕ such that $\phi((i, j)) = \phi'((i, j))$ for $(i, j) \in \{1, 2\} \times \{1, \dots, t-1\}$, $\phi((1, t)) = Q_1 - y_1$ and $\phi((2, t)) = (Q_2 - y_2) \cup (R - V(Q_1))$. \square




FIGURE 5. Case 2: An L_3 -model in B extended to an L_4 -model rooted at (x_1, x_2) in G .

Let $k \geq 1$ and let H be a graph isomorphic to a subdivision of L_k . Then the graph H is a union of $k+2$ paths $P_1, P_2, Q_1, \dots, Q_k$, where

- (i) the paths P_1 and P_2 are disjoint,
- (ii) Q_1, \dots, Q_k are disjoint $V(P_1) - V(P_2)$ paths which intersect the paths P_1 and P_2 in the order in which they are listed,
- (iii) the paths Q_1 and Q_k have their endpoints in the endpoints of the paths P_1 and P_2 .

For such paths $P_1, P_2, Q_1, \dots, Q_k$, we call the tuple $(P_1, P_2; Q_1, \dots, Q_k)$ a *subdivision model* of L_k in H .

Lemma 12. *Let $c \geq 1$ be an integer, let H be a graph isomorphic to a subdivision of L_{2^c} with a subdivision model $(P_1, P_2; Q_1, \dots, Q_{2^c})$, and let φ be a cycle centered coloring of H . If the sets of colors used by φ on the paths Q_1, \dots, Q_{2^c} are all the same, then φ uses more than c colors.*

Proof. We prove the lemma by induction on c . It clearly holds true for $c = 1$, so suppose that $c \geq 2$ and the coloring φ on each path Q_i uses exactly the same set of colors, say A . Since φ is a cycle centered coloring, there exists a vertex of unique color on the cycle $P_1 \cup P_2 \cup Q_1 \cup Q_{2^c}$. Let x be such a vertex. We have $\varphi(x) \notin A$, because every color in A appears on both Q_1 and Q_{2^c} . Hence x lies either on P_1 or on P_2 , and its color is unique in the whole graph H . After possibly exchanging P_1 and P_2 , we may assume that $x \in V(P_1)$, and possibly reversing the order of the paths Q_1, \dots, Q_{2^c} , we may assume that x does not lie on the $V(Q_1)$ – $V(Q_{2^{c-1}})$ subpath of P_1 . The graph H contains a subgraph H_1 isomorphic to a subdivision of $L_{2^{c-1}}$ with subdivision model $(P'_1, P'_2; Q_1, \dots, Q_{2^{c-1}})$, where P'_1 is the $V(Q_1)$ – $V(Q_{2^{c-1}})$ subpath of P_1 and P'_2 is the $V(Q_1)$ – $V(Q_{2^{c-1}})$ subpath of P_2 . By induction hypothesis, φ uses more than $c - 1$ colors on H_1 , and all these colors are distinct from $\varphi(x)$. Hence φ uses more than c colors on H . \square

We may now prove Theorem 1, which we restate for convenience.

Theorem 1. *For every class \mathcal{C} of graphs, the following properties are equivalent.*

- (i) *There exists an integer $k \geq 1$ such that no graph in \mathcal{C} has an L_k minor.*
- (ii) *There exists an integer $m \geq 1$ such that $\text{td}_2(G) \leq m$ for every graph G in \mathcal{C} .*
- (iii) *There exists an integer $c \geq 1$ such that $\chi_{\text{cc}}(G) \leq c$ for every graph G in \mathcal{C} .*

Proof. Let us first show the implication (i) \Rightarrow (ii). We prove the contrapositive. Let \mathcal{C} be a class of graphs such that for every integer $m \geq 1$ there exists a graph G in \mathcal{C} with $\text{td}_2(G) > m$. We need to show that for every $k \geq 1$ there exists a graph in \mathcal{C} with an L_k minor. Fix $k \geq 1$ and let $m = k((k-1)^2 + 2) + 1$. Let G be a graph in \mathcal{C} such that $\text{td}_2(G) \geq m$. By Lemma 9, there exists a connected subgraph B of G which does not have a cutvertex such that $\text{td}_2(B) \geq m$. Since $m \geq 2$, the subgraph B has at least two vertices. Let x_1 and x_2 be distinct vertices of B . By Lemma 11 with $t = k$, B has an L_k minor, and thus so does G . The implication (i) \Rightarrow (ii) follows.

The implication (ii) \Rightarrow (iii) is straightforward, since every 2-connected centered coloring is a cycle centered coloring.

It remains to show the implication (iii) \Rightarrow (i). We prove the contrapositive. More precisely, we show that for every integer $c \geq 1$, every graph G containing an L_{4^c} minor satisfies $\chi_{\text{cc}}(G) > c$. Thus, fix an integer $c \geq 1$ and let G be a graph with an L_{4^c} minor. The graph G contains a subgraph isomorphic to a subdivision of L_{4^c} with a subdivision model $(P_1, P_2; Q_1, \dots, Q_{4^c})$.

Towards a contradiction, suppose that there is a cycle centered coloring of G which uses at most c colors. For every $i \in \{1, \dots, 4^c\}$, let A_i denote the set of colors used on the path Q_i . By the pigeonhole principle, for $k = 2^c$, there are indices i_1, \dots, i_k with $1 \leq i_1 < \dots < i_k \leq 4^c$ such that $A_{i_1} = \dots = A_{i_k}$.

Let P'_1 be the $V(Q_{i_1})$ – $V(Q_{i_k})$ subpath of P_1 and let P'_2 be the $V(Q_{i_1})$ – $V(Q_{i_k})$ subpath of P_2 . Then $P'_1 \cup P'_2 \cup Q_{i_1} \cup \dots \cup Q_{i_k}$ is a subgraph isomorphic to a subdivision of L_{2^c} with a subdivision model $(P'_1, P'_2; Q_{i_1}, \dots, Q_{i_k})$ such that the paths Q_{i_j} all use the same set of colors. By Lemma 12, φ uses more than c colors on this subgraph, contradiction. \square

4. AN APPLICATION TO POSET DIMENSION

In this section we show Theorem 4. In a poset $P = (X, \leq_P)$, we consider the relation \leq_P as a subset of $X^2 = X \times X$. A linear order \leq on X is a *linear extension* of P if $\leq_P \subseteq \leq$. For a set $S \subseteq X$, we denote by $P[S]$ the subposet of P induced by S , that is $P[S] = (S, \leq_P \cap S^2)$.

The next lemma is folklore, a proof is included for completeness.

Lemma 13. *Let $d \geq 1$ be an integer and let \mathcal{C} be a class of graphs closed under taking subgraphs such that every poset whose cover graph is in \mathcal{C} has dimension at most d . Let $P = (X, \leq_P)$ be a poset with cover graph G such that $G - z \in \mathcal{C}$ for some vertex z of G . Then $\dim(P) \leq 2d$.*

Proof. Let $U = \{x \in X : x \geq_P z\}$ and let $D = \{x \in X : x \leq_P z\}$. It is easy to observe that the cover graphs of $P[X \setminus U]$ and $P[X \setminus D]$ are (induced) subgraphs of $G - z$, and thus are in \mathcal{C} . Hence $\dim(P[X \setminus U]) \leq d$ and $\dim(P[X \setminus D]) \leq d$. Let \leq_1, \dots, \leq_d be a realizer of $P[X \setminus U]$, and let $\leq_{d+1}, \dots, \leq_{2d}$ be a realizer of $P[X \setminus D]$. Finally, let \leq_U be a linear extension of $P[U]$ and let \leq_D be a linear extension of $P[D]$.

We construct a realizer $\leq'_1, \dots, \leq'_{2d}$ of P as follows:

$$\leq'_i = \begin{cases} \leq_i \cup (X \setminus U) \times U \cup \leq_U & \text{for } i \in \{1, \dots, d\}, \\ \leq_D \cup D \times (X \setminus D) \cup \leq_i & \text{for } i \in \{d+1, \dots, 2d\}. \end{cases}$$

Now it remains to show that $\leq'_1, \dots, \leq'_{2d}$ is a realizer of P . It is straightforward to verify that each \leq'_i is a linear extension of \leq_P , so if $x \leq_P y$, then $x \leq'_i y$ for $i \in \{1, \dots, 2d\}$. It remains to show that if x and y are incomparable in P , then there exists $i \in \{1, \dots, 2d\}$ such that $y <'_i x$. If $\{x, y\} \subseteq X \setminus U$, then there exists $i \in \{1, \dots, d\}$ such that $y <_i x$ and thus $y <'_i x$. Similarly, if $\{x, y\} \subseteq X \setminus D$, then there exists $i \in \{d+1, \dots, 2d\}$ such that $y <_i x$, so $y <'_i x$. Hence we are left with the case when $\{x, y\} \cap U \neq \emptyset$ and $\{x, y\} \cap D \neq \emptyset$. Since $D \times U \subseteq \leq_P$ and the elements x and y are incomparable in P , this implies one of the elements x or y is equal to z . If $x = z$, then $y \notin U$, so $(y, x) \in (X \setminus U) \times U \subseteq \leq'_1$. Similarly, if $y = z$, then $x \notin D$, so $(y, x) \in D \times (X \setminus D) \subseteq \leq'_{d+1}$. Therefore $\leq'_1, \dots, \leq'_{2d}$ is indeed a realizer of P . \square

We will also need the following theorem. Recall that a *block* of a graph G is a maximal connected subgraph of G without a cutvertex. The blocks can be of three types: maximal 2-connected subgraphs, cut edges together with their endpoints, and isolated vertices. Two blocks have at most one vertex in common, and such a vertex is always a cutvertex.

Theorem 14 (Trotter, Walczak and Wang [20]). *Let $d \geq 1$ be an integer and let \mathcal{C} be a class of graphs such that every poset whose cover graph is in*

\mathcal{C} has dimension at most d . If P is a poset such that all blocks of its cover graph are in \mathcal{C} , then $\dim(P) \leq d + 2$.

We are now ready to prove Theorem 4, which we restate first.

Theorem 4. *Let P be a poset with cover graph G and let $m = \text{td}_2(G)$. Then, P has dimension at most $2^{m+1} - 2$.*

Proof. Let us show the following slightly stronger statement, which will help the induction go through: If the cover graph G of a poset P satisfies $\text{td}_2(G) \leq m$, then $\dim(P) \leq 2^{m+1} - 2$; furthermore, $\dim(P) \leq 2^{m+1} - 4$ if G is 2-connected. We prove the statement by induction on m .

For the base case ($m = 1$), G has no edges, and thus P is an *antichain*, that is, a poset in which no pair of distinct elements is comparable. Hence $\dim(P) \leq 2$; indeed, if the elements of P are x_1, \dots, x_n , then P has a realizer $\{\leq, \leq'\}$ where $x_1 < \dots < x_n$ and $x_n <' \dots <' x_1$. As $2^{m+1} - 2 = 2$, the statement holds. (Note that in this case G cannot be 2-connected, so the second part of the statement holds vacuously.)

For the inductive case ($m \geq 2$), we first establish the case where G is 2-connected. Consider a 2-connected centered coloring of G with m colors. There is a vertex z of G whose color is unique in this coloring. Thus $\text{td}_2(G - z) \leq m - 1$. By induction and Lemma 13, we deduce that $\dim(P) \leq 2 \cdot (2^{m-1+1} - 2) = 2^{m+1} - 4$, as desired.

Now we turn to the case that G is not 2-connected. Then each block of G is either 2-connected, or isomorphic to K_1 or K_2 . Using that our claim holds in the 2-connected case, and the obvious fact that a poset whose cover graph is isomorphic to K_1 or K_2 has dimension $1 \leq 2^{m+1} - 4$, we deduce from Theorem 14 that $\dim(P) \leq (2^{m+1} - 4) + 2 = 2^{m+1} - 2$. \square

5. BUMPING A LADDER

In this section, we prove Theorem 2.

In a graph G , if A is the set of cutvertices, and \mathcal{B} is the set of blocks of G , then the *block graph* of G is the bipartite graph on $A \cup \mathcal{B}$, where $a \in A$ is adjacent to $B \in \mathcal{B}$ if $a \in V(B)$. The block graph of a graph is always a forest.

Lemma 15. *Let m and p be positive integers. Let G be a graph with at least p^m vertices and with $\text{td}_2(G) \leq m$. Then there exists a set $Z \subseteq V(G)$ with $|Z| \leq m - 1$ such that $G - Z$ has at least p blocks.*

Proof. We prove the lemma by induction on m . If $m = 1$, then the lemma works with $Z = \emptyset$: since $\text{td}_2(G) \leq 1$, every vertex in G is an isolated vertex forming a block.

Now suppose that $m \geq 2$. Fix a 2-connected centered coloring of G using at most m colors. If G has at least p blocks, then the lemma holds with $Z = \emptyset$. Let us suppose that G has less than p blocks. Then some block of G has at least p^{m-1} vertices. Let B_0 be such a block, and let x be a vertex of unique color in B_0 . Thus, $\text{td}_2(B_0 - x) \leq m - 1$. By induction hypothesis, there exists a vertex subset Y in $B_0 - x$ with $|Y| \leq m - 2$ such that $(B_0 - x) - Y$ has at least p blocks. Let $Z = Y \cup \{x\}$. It remains to show that $G - Z$ has at least p blocks. The graph $(B_0 - x) - Y = B_0 - Z$

is a subgraph of $G - Z$, so every block of $B_0 - Z$ is contained in a block of $G - Z$. Since $B_0 - Z$ has at least p blocks, it suffices to show that no block of $G - Z$ contains two blocks of $B_0 - Z$.

Towards a contradiction, suppose that a block B of $G - Z$ contains two distinct blocks B_1 and B_2 of $B_0 - Z$. In particular, we have $|V(B) \cap V(B_0)| \geq |V(B_1) \cup V(B_2)| \geq 2$. Since two blocks of G have at most one vertex in common, this implies that B_0 is the (unique) block of G containing the block B of $G - Z$. But $B \subseteq B_0 \cap (G - Z) = B_0 - Z$, so B is a block of $B_0 - Z$, so $B_1 \subseteq B$ and $B_2 \subseteq B$ imply $B_1 = B = B_2$, contradiction. This concludes the proof. \square

Observation 16. *If G is a 3-connected graph on at least 5 vertices and e is an edge in G such that G/e is not 3-connected, then G admits a nontrivial separation (A_1, A_2) of order 3 such that both endpoints of e lie in $A_1 \cap A_2$.*

The following lemma is due to Halin [5]. Since we are not aware of a published proof in English³, we include one for the reader's convenience.

Lemma 17 (Halin [5]). *Let G be a 3-connected graph. Let $e \in E(G)$. If neither G/e nor $G - e$ is 3-connected, then some endpoint of e has degree 3.*

Proof. Suppose that G/e and $G - e$ are not 3-connected. As a 3-connected graph, G has at least 4 vertices. If $|V(G)| = 4$, then G is complete and the lemma holds as all vertices are of degree 3. Hence we assume that $|V(G)| \geq 5$.

Since $G - e$ is not 3-connected and has at least 5 vertices, we can fix a nontrivial 2-separation (A_1, A_2) of $G - e$. As a 3-connected graph, G does not admit a nontrivial 2-separation, so neither A_1 nor A_2 contains both endpoints of e . Hence e has an endpoint v_1 in $A_1 \setminus A_2$ and an endpoint v_2 in $A_2 \setminus A_1$.

By Observation 16, we can fix a nontrivial order-3 separation (B_1, B_2) of G with $\{v_1, v_2\} \subseteq B_1 \cap B_2$. Let w denote the vertex of $B_1 \cap B_2$ other than v_1 and v_2 . As (A_1, A_2) is a separation of $G - v_1v_2$ and (B_1, B_2) is a separation of G with $\{v_1, v_2\} \subseteq B_1 \cap B_2$, the pair $(A_i \cap B_j, A_{3-i} \cup B_{3-j})$ is a separation of G for $i, j \in \{1, 2\}$.

We claim that $(A_1 \cap A_2) \setminus B_1 \neq \emptyset$ and $(A_1 \cap A_2) \setminus B_2 \neq \emptyset$. Towards a contradiction, suppose that it is not the case. After possibly swapping B_1 and B_2 we may assume that

$$A_1 \cap A_2 \subseteq B_2. \quad (1)$$

Consider the separations $(A_1 \cap B_1, A_2 \cup B_2)$ and $(A_2 \cap B_1, A_1 \cup B_2)$ of G . The order of those separations is at most 2 as by (1), for $i \in \{1, 2\}$ we have

$$\begin{aligned} (A_i \cap B_1) \cap (A_{3-i} \cup B_2) &= (A_i \cap B_1 \cap A_{3-i}) \cup (A_i \cap B_1 \cap B_2) \\ &\subseteq (A_i \cap B_1 \cap B_2) = \{v_i, w\} \end{aligned}$$

Since G is 3-connected, both separations $(A_1 \cap B_1, A_2 \cup B_2)$ and $(A_2 \cap B_1, A_1 \cup B_2)$ must be trivial. However, by nontriviality of (B_1, B_2) , B_1 is a proper subset of $V(G)$, and thus $A_1 \cap B_1$ and $A_2 \cap B_1$ are proper subsets of

³We note however that the lemma appears as an exercise in Diestel's textbook [2, Chapter 3, exercise 10].

$V(G)$ as well. Therefore it must be the case that $A_2 \cup B_2 = A_1 \cup B_2 = V(G)$, and thus by (1)

$$V(G) = (A_2 \cup B_2) \cap (A_1 \cup B_2) = (A_1 \cap A_2) \cup B_2 = B_2,$$

contradicting the nontriviality of (B_1, B_2) . This contradiction proves our claim that $(A_1 \cap A_2) \setminus B_1 \neq \emptyset$ and $(A_1 \cap A_2) \setminus B_2 \neq \emptyset$.

The order of the separation (A_1, A_2) is at most 2, so the fact that $(A_1 \cap A_2) \setminus B_1 \neq \emptyset$ and $(A_1 \cap A_2) \setminus B_2 \neq \emptyset$ implies that $A_1 \cap A_2 = \{y_1, y_2\}$ for some two vertices $y_1 \in B_1 \setminus B_2$ and $y_2 \in B_2 \setminus B_1$.

Since $w \notin A_1 \cap A_2$, either $w \in A_1 \setminus A_2$ or $w \in A_2 \setminus A_1$. The two cases are symmetric, so let us assume that $w \in A_2 \setminus A_1$. Summarizing, we have $v_i \in B_1 \cap B_2 \cap A_i \setminus A_{3-i}$ and $y_i \in A_1 \cap A_2 \cap B_i \setminus B_{3-i}$ for $i \in \{1, 2\}$, $w \in A_2 \setminus A_1$, and $A_1 \cap A_2 \cap B_1 \cap B_2$ is empty. Hence $(A_1 \cap B_1, A_2 \cup B_2)$ and $(A_1 \cap B_2, A_2 \cup B_1)$ are separations of G with

$$(A_1 \cap B_1) \cap (A_2 \cup B_2) = \{v_1, y_1\} \quad \text{and} \quad (A_1 \cap B_2) \cap (A_2 \cup B_1) = \{v_1, y_2\}.$$

The 3-connectedness of G implies that the two separations are trivial and thus $A_1 = \{v_1, y_1, y_2\}$. As (A_1, A_2) is a separation of $G - e$, this implies that v_1 can be adjacent only to the vertices y_1, y_2 and v_2 , so the degree of v_1 is at most 3. Since G is 3-connected, the degree of v_1 is at least 3, so the vertex $z = v_1$ satisfies the lemma. \square

The next result demonstrates the usefulness of vertices of degree three when attempting to preserve 3-connectivity while taking minors. The result is also due to Halin [6]; see [1] for an alternate proof in English.

Lemma 18 (Halin [6]). *Let G be a 3-connected graph on at least 5 vertices and let $v \in V(G)$ be a vertex of degree 3. Then there exists an edge e incident to v such that G/e is 3-connected.*

Lemma 19. *Let G be a 3-connected graph, let P be an induced path in G , and let $Z \subseteq V(G) \setminus V(P)$ be such that every internal vertex of P has all its neighbors in $V(P) \cup Z$. If the length of P is at least $2|Z| + 3$, then there exists an edge $e \in E(P)$ such that G/e is 3-connected.*

Proof. By the 3-connectivity of G , every internal vertex of P has a neighbor in Z . Hence, when we fix $k = |Z|$ and ℓ to be the length of P , we have $k \geq 1$ and $\ell \geq 2k + 3 \geq 5$.

For each $v \in V(G)$, let $N(v)$ denote the set of neighbors of v in G . Let v_0, v_1, \dots, v_ℓ be the vertices of P in the order in which they appear on the path. For each $i \in \{1, \dots, \ell - 1\}$, let $X_i = \bigcup_{j < i} N(v_j) \cap Z$ and let $Y_i = \bigcup_{j > i} N(v_j) \cap Z$. Since $X_1 \subseteq \dots \subseteq X_{\ell-1}$ and $|X_{\ell-1}| \leq |Z| = k$, there exist at most k indices i with $1 \leq i \leq \ell - 2$ such that $X_i \neq X_{i+1}$. Symmetrically, there are at most k indices i with $1 \leq i \leq \ell - 2$ such that $Y_i \neq Y_{i+1}$. Since $\ell - 2 > 2k$, there exists i such that $1 \leq i \leq \ell - 2$ and $(X_i, Y_i) = (X_{i+1}, Y_{i+1})$. So $N(v_i) \cap Z \subseteq N(\{v_0, \dots, v_{i-1}\})$ and $N(v_{i+1}) \cap Z \subseteq N(\{v_{i+2}, \dots, v_\ell\})$.

Let e be the edge $v_i v_{i+1}$ of P . We claim G/e is 3-connected. Arguing by contradiction, suppose that G/e is not 3-connected. By Observation 16 there is a nontrivial separation (A_1, A_2) of G of order 3 with $v_i, v_{i+1} \in A_1 \cap A_2$. Let $u \notin \{v_i, v_{i+1}\}$ denote the third vertex of $A_1 \cap A_2$. Clearly, we can assume that u does not lie on $v_0 P v_{i-1}$ or that u does not lie on $v_{i+2} P v_\ell$. We will

continue the proof assuming the former; the other case has a symmetric proof.

Lest $(A_1 \setminus \{v_i\}, A_2)$ form a 2-separation in G , there exists a vertex $a \in A_1 \setminus A_2$ such that a is a neighbor of v_i in G . Since v_i is an internal vertex of P we have $a \in Z$ or $a = v_{i-1}$. If $a \in Z$, then since $a \in X_{i+1} = X_i$, there exists an edge of G joining a to a vertex v_{j_1} with $j_1 < i$, and since $a \in A_1 \setminus A_2$, we have $v_{j_1} \in A_1$. If $a = v_{i-1}$, then for $j_1 = i - 1$ the vertex v_{j_1} lies in A_1 . Thus, in either case, there exists $j_1 < i$ such that $v_{j_1} \in A_1$.

Swapping A_1 and A_2 in the above reasoning, there must be a vertex v_{j_2} with $j_2 < i$, such that $v_{j_2} \in A_2$. Therefore $v_{j_1}Pv_{j_2}$ intersects $A_1 \cap A_2$, which contradicts that v_0Pv_{i-1} has no vertices in $A_1 \cap A_2$. We conclude that G/e is 3-connected, as claimed. \square

The following observation will be used in the proof of Lemma 21.

Observation 20. *Let k be an integer with $k \geq 2$, let H be a graph isomorphic to a subdivision of L_k , and let ϕ be an L_k -model in H . For any two distinct vertices x and y of L_k ,*

- (i) *if x and y are adjacent in L_k , then there is exactly one edge between $\phi(x)$ and $\phi(y)$ in H , and*
- (ii) *if x and y are nonadjacent in L_k , then there are no edges between $\phi(x)$ and $\phi(y)$ in H .*

Proof. If this is not true, then there exists an edge e in H such that ϕ is an L_k -model in $H - e$. Hence there exists a proper subgraph of H which is isomorphic to a subdivision of L_k , which is impossible since L_k has no vertices of degree 1. \square

Lemma 21. *Let k be an integer with $k \geq 2$, let H be a graph isomorphic to a subdivision of L_k , and let ϕ be an L_k -model in H . If z_1 and z_2 are distinct vertices of H with $z_1 \in V(\phi((i_1, j_1)))$, $z_2 \in V(\phi((i_2, j_2)))$ and $1 \leq j_1 \leq j_2 \leq k$, then there exists an L_{j_1} -model rooted at (z_1, z_2) and an L_{k-j_2+1} -model rooted at (z_1, z_2) .*

Proof. We only show the existence of an L_{j_1} -model rooted at (z_1, z_2) since an L_{k-j_2+1} -model rooted at (z_1, z_2) can be obtained in a similar way using a symmetry of the ladder.

If $j_1 = 1$, the existence of the desired model follows from connectedness of H . Let us hence assume that $j_1 \geq 2$. Let $A = \bigcup_{1 \leq j < j_1} V(\phi((1, j))) \cup V(\phi((2, j)))$. As H is 2-connected, we can fix two disjoint $A - \{z_1, z_2\}$ paths R_1 and R_2 . By Observation 20, in H there are exactly two edges between A and $V(H) \setminus A$, and thus, since z_1 and z_2 do not lie in A , one of the paths R_1 and R_2 has an endpoint in $V(\phi((1, j_1 - 1)))$, and the other in $V(\phi((2, j_1 - 1)))$. We assume that R_1 has an endpoint v_1 in $V(\phi((1, j_1 - 1)))$ and R_2 has an endpoint v_2 in $V(\phi((2, j_1 - 1)))$. Since $H - A$ is connected, we can fix a $V(R_1) - V(R_2)$ path S in $H - A$. Consider the L_{j_1} -model ϕ' defined as follows.

$$\phi'((i, j)) = \begin{cases} \phi((i, j)) & \text{if } j < j_1, \\ R_1 - v_1 & \text{if } (i, j) = (1, j_1), \\ (R_2 - v_2) \cup (S - V(R_1)) & \text{if } (i, j) = (2, j_1). \end{cases}$$

The model ϕ' is rooted at (z_1, z_2) or (z_2, z_1) , so after possibly swapping all $\phi'((1, j))$ with $\phi'((2, j))$ we obtain the desired model. \square

Lemma 22. *Let k be an integer with $k \geq 2$ and let H be a graph isomorphic to a subdivision of L_k . For every 3-element vertex subset Z in H there exists a pair (z_1, z_2) of distinct vertices in Z and an $L_{\lceil (k+1)/2 \rceil}$ -model in H rooted at (z_1, z_2) .*

Proof. Fix an L_k -model ϕ in H . Let $Z = \{z_1, z_2, z_3\}$, and let (i_1, j_1) , (i_2, j_2) and (i_3, j_3) be three vertices of L_k such that $z_a \in V(\phi((i_a, j_a)))$ for $a \in \{1, 2, 3\}$. Without loss of generality we assume that $j_1 \leq j_2 \leq j_3$. By Lemma 21, there exist an L_{k-j_2+1} -model rooted at (z_1, z_2) and an L_{j_2} -model rooted at (z_2, z_3) . One of these models contains a desired rooted model of $L_{\lceil (k+1)/2 \rceil}$. \square

Lemma 23. *Let k and m be integers with $k \geq 2$ and $m \geq 1$, let G be a graph, and let H^1, \dots, H^{m^2+1} be disjoint subgraphs of G each isomorphic to a subdivision of L_k . If G contains as a subgraph a forest F with at most m components, such that for every $i \in \{1, \dots, m^2 + 1\}$ the graph H^i intersects each component of F in at most one vertex and H^i contains vertices from at least three components of F , then G has an L_{k+1} minor.*

Proof. By Lemma 22, for every $i \in \{1, \dots, m^2 + 1\}$ there exists a pair (z_1^i, z_2^i) of vertices from $V(H^i) \cap V(F)$ such that there exists an $L_{\lceil (k+1)/2 \rceil}$ -model ϕ^i in H^i which is rooted at (z_1^i, z_2^i) . The vertices z_1^i and z_2^i lie in distinct components of F , and by the pigeonhole principle, there exist distinct indices i and j in $\{1, \dots, m^2 + 1\}$ such that the vertices z_1^i and z_1^j lie in the same component of F , and the vertices z_2^i and z_2^j lie in the same component of F . Let P_1 be the $z_1^i - z_1^j$ path in F and let P_2 be the $z_2^i - z_2^j$ path in F . Gluing the $L_{\lceil (k+1)/2 \rceil}$ -models ϕ^i and ϕ^j using paths P_1 and P_2 , we obtain a model of L_{k+1} in G . \square

We may now turn to the proof of Theorem 2, which we restate first.

Theorem 2 (Bumping a ladder). *For every integer $k \geq 1$ there exists an integer $N \geq 1$ with the following property. Every 3-connected graph G containing a union of N vertex-disjoint copies of L_k as a minor contains L_{k+1} as a minor.*

Proof. Let us first observe that if $k = 1$, then the theorem holds trivially because every 3-connected graph contains a cycle of length at least 4, which is isomorphic to a subdivision of L_2 . Thus, we may assume that $k \geq 2$. Fix m to be the minimum positive integer such that every graph G with no L_{k+1} minor satisfies $\text{td}_2(G) \leq m$. Note that m is well defined by Theorem 1. Let λ_0 , λ , and p be positive integers defined as follows.

$$\lambda_0 := ((2m + 2)(4k + 1) + 4)(m^2 + 2k^3(m - 1) + 1);$$

$$\lambda := 6m(\lambda_0 + 3) + 2;$$

$$p := ((m - 1)^2 + 4(m - 1))(m^2 + 4(m - 1))(\lambda + 1).$$

We show that the theorem holds for $N = p^m$. Suppose not and fix G to be a minor minimal 3-connected graph which contains a union of N disjoint

copies of L_k as a minor, but which does not contain L_{k+1} as a minor. Hence we can choose disjoint subgraphs H_1, \dots, H_N of G such that each H_i is isomorphic to a subdivision of L_k . Choose such H_1, \dots, H_N which minimize the value of $|V(H_1 \cup \dots \cup H_N)|$.

Let $H = H_1 \cup \dots \cup H_N$. For each H_i fix a 4-element vertex subset X_i such that the graph obtained from H_i by suppressing every vertex of degree two which is not in X results in a graph isomorphic to L_k . Let $X = X_1 \cup \dots \cup X_N$.

Claim 1. *Every vertex of G adjacent to a vertex not in $V(H)$ has degree 3 and belongs to X .*

Proof. Let $v \in V(G)$ be a vertex adjacent to a vertex $u \in V(G) \setminus V(H)$ in G . Both G/uv and $G - uv$ contain H as a minor, so by minimality, neither G/uv nor $G - uv$ is 3-connected. By Lemmas 17 and 18, there exists $x \in \{u, v\}$ of degree 3 in G and an edge e incident to x such that G/e is 3-connected. If some endpoint of e is not in $V(H)$ then again G/e contains H as a minor, and we have a contradiction to minimality. Thus, both endpoints of e are in $V(H)$, and hence $x = v$. As v has degree 3 in G and $u \in V(G) \setminus V(H)$ is its neighbor, we conclude that v has degree exactly two in H and that $e \in E(H)$. Finally, if $v \notin X$, then H/e contains as a minor N disjoint copies of L_k and we see that G/e contradicts the minimality of our counterexample G . Thus, $v \in X$. \diamond

Since G has no L_{k+1} minor, we have $\text{td}_2(G) \leq m$. We have $|V(G)| \geq N = p^m$, so by Lemma 15, there exists a vertex subset Z of G with $|Z| \leq m - 1$ such that $G - Z$ has at least p blocks. Fix such a set Z .

Let us *mark* each vertex v of $G - Z$ which satisfies

- a) $v \in V(H)$ and v is adjacent in H to a vertex from $Z \cap V(H)$, or
- b) $v \in V(G) \setminus V(H)$ and v is adjacent in G to a vertex from $X \cap Z$.

Since $|Z| \leq m - 1$ and the maximum degree of H is at most 3, the number of vertices which satisfy a) is at most $3(m - 1)$. As every vertex in X with a neighbor in $V(G) \setminus V(H)$ has degree 3 by Claim 1, there are at most $m - 1$ vertices which satisfy b). We deduce the following.

Claim 2. *There are at most $4m - 4$ marked vertices.*

Claim 3. *$G - Z$ has at most $(m - 1)^2 + 4(m - 1)$ components.*

Proof. Suppose to the contrary that $G - Z$ has $(m - 1)^2 + 4m - 4 + 1$ distinct components $C^1, \dots, C^{(m-1)^2+4m-3}$. By Claim 2, we can assume without loss of generality that none of the components $C^1, \dots, C^{(m-1)^2+1}$ contains a marked vertex. For each $i \in \{1, \dots, (m - 1)^2 + 1\}$, choose an arbitrary vertex v of C^i . If v is in H , let H^i denote the (unique) subgraph among H_1, \dots, H_N which contains v . If v is not in H , then, by Claim 1, v has a neighbor $u \in X$. As v is not marked, $u \notin Z$ and therefore u lies in C^i as well. Fix such a neighbor u and let H^i be the component of H containing the vertex u . Since C^i does not contain a marked vertex, H^i must be a subgraph of C^i .

For every $i \in \{1, \dots, (m - 1)^2 + 1\}$ we have $|V(C^i)| \geq 4$, since C^i contains a subdivision of L_k . The 3-connectivity of G implies that the set Z has at least

three vertices. By Menger's theorem, for every $i \in \{1, \dots, (m-1)^2 + 1\}$, we can fix three disjoint $V(H^i)$ - Z paths Q_1^i , Q_2^i and Q_3^i with all internal vertices contained in $V(C^i)$. Let $F = \bigcup_{i=1}^{(m-1)^2+1} (Q_1^i \cup Q_2^i \cup Q_3^i)$. Each component of the graph F is the union of a number of paths which have a common endpoint in Z but are otherwise disjoint. Thus F is a forest, and since $|Z| \leq m-1$, it has at most $m-1$ components. For every $i \in \{1, \dots, (m-1)^2 + 1\}$ the subgraph H^i intersects three components of F , each in one vertex. By Lemma 23, G has an L_{k+1} minor, contradiction. \diamond

Claim 4. *The block graph of every component of $G - Z$ is a tree with at most $m^2 + 4(m-1)$ leaves.*

Proof. Suppose to the contrary that there is a component C of $G - Z$ whose block graph \mathcal{T} is a tree with more than $m^2 + 4(m-1)$ leaves. Let $B^1, \dots, B^{m^2+4(m-1)+1}$ be distinct leaves of \mathcal{T} , and for each $i \in \{1, \dots, m^2 + 4(m-1) + 1\}$, let a^i be the cutvertex of $G - Z$ adjacent to B^i in \mathcal{T} . By Claim 2, we may assume without loss of generality that for $i \in \{1, \dots, m^2 + 1\}$, there is no marked vertex in $B^i - a^i$. For each $i \in \{1, \dots, m^2 + 1\}$ we show there exists a component H^i of H contained in B^i . Fix an index $i \in \{1, \dots, m^2 + 1\}$ and let v be a vertex of $B^i - a^i$. If $v \in V(H)$, fix H^i to be the component of H which contains v . If $v \in V(G) \setminus V(H)$, then v has at least three neighbors all of which must be contained in X . Fix a neighbor $u \in X \setminus \{a_i\}$ of v . As v is not marked, u is a vertex of B^i . In this case, fix H^i to be the component of H which contains u . Since $B^i - a^i$ has no marked vertex, no edge of H^i connects $B^i - a^i$ to Z . Since a^i is the only cutvertex of $G - Z$ in B^i and since H_i is 2-connected, we conclude that H^i is contained in B^i .

Note that $B^i - a^i$ is a component of $G - (Z \cup \{a^i\})$. As G is 3-connected, this implies that $|Z \cup \{a^i\}| \geq 3$. Hence by Menger's theorem, there exist three disjoint paths from $V(H^i)$ to the set $Z \cup \{a^i\}$ with all internal vertices contained in $B^i - a^i$. Fix such three disjoint $V(H^i)$ - $(Z \cup \{a^i\})$ paths Q_1^i , Q_2^i and Q_3^i .

Let T be a tree in $G - Z$ which contains all vertices a^1, \dots, a^{m^2+1} but none of the vertices of $B^i - a^i$ for $i \in \{1, \dots, m^2 + 1\}$, and let $F = T \cup \bigcup_{i=1}^{m^2+1} (Q_1^i \cup Q_2^i \cup Q_3^i)$. Each component of F is either a tree obtained as the union of paths which have a common endpoint in Z but are otherwise disjoint, or a tree obtained as the union of T and paths with one endpoint at $a^i \in V(T)$ for some $i \in \{1, \dots, m^2 + 1\}$. Since $|Z| \leq m-1$, this implies that F is a forest with at most m components. Moreover, for every $i \in \{1, \dots, m^2 + 1\}$, the graph H^i intersects each component of F in at most one vertex, and H^i has a non-empty intersection with at least three components of F . Hence by Lemma 23, G has an L_{k+1} minor, contradiction. \diamond

Recall that the graph $G - Z$ has at least $p \geq ((m-1)^2 + 4(m-1))(m^2 + 4(m-1))(\lambda + 1)$ blocks. Hence by Claim 3, there exists a component C of $G - Z$ with at least $(m^2 + 4(m-1))(\lambda + 1)$ blocks. Let \mathcal{T} be the block graph of C . By Claim 4, \mathcal{T} has at most $(m^2 + 4(m-1))$ leaves, and thus \mathcal{T} contains a path on $\lambda + 1$ blocks and λ cutvertices. Let $B_0 a_1 B_1 a_2 \dots a_\lambda B_\lambda$ be such a path. Let \mathcal{T}' denote the subgraph of \mathcal{T} obtained by removing all edges of the form $a_i B_i$ with $i \in \{1, \dots, \lambda\}$. For every $i \in \{0, \dots, \lambda\}$ let

W_i be the set of all vertices contained in those blocks of C which lie in the same component of \mathcal{T}' as B_i . The sets W_0, \dots, W_λ obtained this way induce connected subgraphs of C whose union is C , and for every $i \in \{1, \dots, \lambda\}$, we have

$$(W_0 \cup \dots \cup W_{i-1}) \cap (W_i \cup \dots \cup W_\lambda) = W_{i-1} \cap W_i = \{a_i\}.$$

See Figure 6.




FIGURE 6. Blocks of the component C of $G - Z$. The block graph \mathcal{T} of C contains a long path $B_0 a_1 B_1 a_2 \dots a_\lambda B_\lambda$. The remaining blocks are grouped into bundles. The bundle W_2 is highlighted in gray.

Claim 5. *There exists an index i_0 with $1 \leq i_0$ and $i_0 + \lambda_0 \leq \lambda - 1$ such that*

- (i) $W_{i_0} \cup \dots \cup W_{i_0 + \lambda_0}$ does not contain a marked vertex, and
- (ii) *there exist two disjoint $(W_0 \cup \dots \cup W_{i_0 - 1}) - (W_{i_0 + \lambda_0 + 1} \cup \dots \cup W_\lambda)$ paths in G which are internally disjoint from C . See Figure 7.*




FIGURE 7. $W_{i_0} \cup \dots \cup W_{i_0 + \lambda_0}$ does not contain a marked vertex, and there are two disjoint paths connecting outside regions and internally disjoint with C .

Proof. For each $z \in Z$, let J_z denote the set of the two largest indices $i \in \{2, \dots, \lambda - 1\}$ such that z is adjacent to a vertex from $W_i \setminus \{a_i\}$ (if there are less than two such indices, let J_z consist of all these indices, possibly $J_z = \emptyset$). Let $J = \bigcup_{z \in Z} J_z$. As $|Z| < m$, we have $|J| < 2m$.

By Claim 2, there are less than $6m$ indices $i \in \{2, \dots, \lambda - 1\}$ such that $i \in J$ or $W_i \setminus \{a_{i+1}\}$ contains a marked vertex. Consider the set $\{2, \dots, \lambda - 1\}$ of $\lambda - 2$ consecutive indices. As $\lambda - 2 = 6m(\lambda_0 + 3)$, by the pigeonhole principle we can find in this set at least $\lambda_0 + 3$ consecutive indices i such that $i \notin J$ and $W_i \setminus \{a_{i+1}\}$ does not contain a marked vertex. We fix a sequence of $\lambda_0 + 2$ such indices: Let i_0 be an index with $2 \leq i_0$ and $i_0 + \lambda_0 + 1 \leq \lambda - 1$ such that $i \notin J$ and $W_i \setminus \{a_{i+1}\}$ does not contain a marked vertex, for every $i \in \{i_0, \dots, i_0 + \lambda_0 + 1\}$. We have

$$\bigcup_{i=i_0}^{i_0+\lambda_0} W_i \subseteq \bigcup_{i=i_0}^{i_0+\lambda_0+1} W_i \setminus \{a_{i+1}\}.$$

By the choice of i_0 , the right-hand side does not contain a marked vertex, therefore the index i_0 satisfies (i).

Partition the set $V(C) \setminus \{a_{i_0}\}$ into the sets U_1 , U_2 and U_3 , where

$$\begin{aligned} U_1 &= (W_0 \cup \dots \cup W_{i_0-1}) \setminus \{a_{i_0}\}, \\ U_2 &= (W_{i_0} \cup \dots \cup W_{i_0+\lambda_0}) \setminus \{a_{i_0}\}, \text{ and} \\ U_3 &= (W_{i_0+\lambda_0+1} \cup \dots \cup W_\lambda) \setminus \{a_{i_0+\lambda_0+1}\}. \end{aligned}$$

Since G is 3-connected, the graph $G - a_{i_0}$ is 2-connected. We have $|U_1| \geq 2$ as $i_0 \geq 2$, and $|U_2 \cup U_3| \geq 2$ as $\lambda - i_0 \geq \lambda_0 \geq 2$, so there exist two disjoint U_1 – $(U_2 \cup U_3)$ paths P_1 and P_2 in $G - a_{i_0}$. For $i \in \{1, 2\}$, let v_i denote the endpoint of P_i contained in $U_2 \cup U_3$, and let z_i denote the vertex adjacent to v_i in P_i . Since a_{i_0} is a cutvertex in C , we deduce that $\{z_1, z_2\} \subseteq Z$.

Each of the paths P_1 and P_2 is internally disjoint from $V(C)$ and has one endpoint in U_1 , but the other endpoint possibly lies in U_2 and not in U_3 as required by (ii). We will show however that if for some $i \in \{1, 2\}$ we have $v_i \in U_2$, then z_i is adjacent to at least two vertices in U_3 . This will imply that by replacing the endpoints of P_1 and P_2 contained in U_2 we can obtain two disjoint U_1 – U_3 paths in G which are internally disjoint from $V(C)$, thus proving that i_0 satisfies (ii).

Let us hence fix an index $i \in \{1, 2\}$ such that $v_i \in U_2$, and let $j \in \{i_0, \dots, i_0 + \lambda_0\}$ be such that $v_i \in W_j \setminus \{a_j\}$. By our choice of i_0 , we have $j \notin J_{z_i}$. Since z_i is adjacent to the vertex $v_i \in W_j \setminus \{a_j\}$, the definition of J_{z_i} implies that $|J_{z_i}| = 2$ and both elements of J_{z_i} are greater than $i_0 + \lambda_0$. Therefore z_i is indeed adjacent to at least two vertices in U_3 , completing the proof of (ii) \diamond

Let us fix an index i_0 as in Claim 5.

Claim 6. *For every index $i \in \{i_0, \dots, i_0 + \lambda_0 - 2\}$, either*

- (i) *some component of H has all its vertices contained in one of the sets W_i , W_{i+1} or W_{i+2} , or*
- (ii) *there is a path in H between a_i and a_{i+1} with all vertices in W_i .*

Proof. For the contrary, suppose that for a fixed index $i \in \{i_0, \dots, i_0 + \lambda_0 - 2\}$ both items do not hold. By Claim 1, $V(G) \setminus V(H)$ is an independent set in G . As W_{i+1} contains two endpoints of an edge in the block B_{i+1} , this implies that $V(H) \cap W_{i+1} \neq \emptyset$. Fix a component H^0 of H such that $V(H^0) \cap W_{i+1} \neq \emptyset$.

Since the set $W_i \cup W_{i+1} \cup W_{i+2}$ does not contain a marked vertex, no edge of H^0 connects a vertex in $W_i \cup W_{i+1} \cup W_{i+2}$ to a vertex in Z . As a_{i+1} and a_{i+2} are cutvertices of $G[W_i \cup W_{i+1} \cup W_{i+2}]$ and H^0 is 2-connected, this implies that if $V(H^0) \subseteq W_i \cup W_{i+1} \cup W_{i+2}$, then (i) holds. Let us hence assume that

$$V(H^0) \setminus (W_i \cup W_{i+1} \cup W_{i+2}) \neq \emptyset.$$

Let $U = (V(G) \setminus (W_i \cup W_{i+1} \cup W_{i+2})) \cup \{a_i, a_{i+3}\}$. Note that $a_i \in V(H^0)$ as otherwise $((W_i \cup W_{i+1} \cup W_{i+2}) \cap V(H^0), U \cap V(H^0))$ is a nontrivial 1-separation of the 2-connected H^0 . Moreover, if $a_{i+1} \notin V(H^0)$, then $((W_{i+1} \cup W_{i+2}) \cap V(H^0), (U \cup W_i) \cap V(H^0))$ is a nontrivial 1-separation of H^0 , so $a_{i+1} \in V(H^0)$. Therefore $\{a_i, a_{i+1}\} \subseteq V(H^0)$.

Towards a contradiction, suppose that H^0 does not contain a path between a_i and a_{i+1} with all vertices in W_i , and let D be the component of $H^0[W_i \cap V(H^0)]$ which contains a_i but not a_{i+1} . In such case we obtain a nontrivial 1-separation $((W_i \cup W_{i+1} \setminus V(D)) \cap V(H^0), (W_{i+2} \cup U \cup V(D)) \cap V(H^0))$ of H^0 , which is a contradiction. Hence (ii) holds. \diamond

Claim 7. *Among any $(2m+2)(4k+1)+3$ consecutive indices in $\{i_0, \dots, i_0 + \lambda_0\}$ we can find an index i such that some component of H has all its vertices contained in W_i .*

Proof. Suppose to the contrary that there is an index α with $i_0 \leq \alpha$ and $\alpha + (2m+2)(4k+1) + 2 \leq i_0 + \lambda_0$ such that for every $i \in \{\alpha, \dots, \alpha + (2m+2)(4k+1) + 2\}$ there is no $j \in \{1, \dots, N\}$ with $V(H_j) \subseteq W_i$.

By Claim 6 applied to all indices $i \in \{\alpha, \dots, \alpha + (2m+2)(4k+1)\}$, we can fix a component H^0 of H and a path P contained in H^0 with endpoints in a_α and $a_{\alpha+(2m+2)(4k+1)+1}$ which traverses the vertices $a_\alpha, \dots, a_{\alpha+(2m+2)(4k+1)+1}$ in that order and the subpath of P from a_i to a_{i+1} has all its vertices in W_i for $i \in \{\alpha, \dots, \alpha + (2m+2)(4k+1)\}$.

The graph H^0 contains $2k$ vertices which are of degree 3 or belong to X . Such vertices lie in at most $4k$ of the sets $W_\alpha, \dots, W_{\alpha+(2m+2)(4k+1)}$. Hence there exists an index α' with $\alpha \leq \alpha'$ and $\alpha' + (2m+1) \leq \alpha + (2m+2)(4k+1)$ such that $W_{\alpha'} \cup \dots \cup W_{\alpha'+(2m+1)}$ does not contain a vertex of degree 3 in H^0 nor a vertex from $V(H^0) \cap X$. This in particular implies that the subgraph of H^0 induced by those vertices which lie in W_i is the subpath $a_i P a_{i+1}$ for $i \in \{\alpha', \dots, \alpha' + 2m + 1\}$. Let us fix such an index α' .

We claim that actually $a_i P a_{i+1}$ contains all vertices in W_i for $i \in \{\alpha', \dots, \alpha' + 2m + 1\}$. Suppose to the contrary that for some index i there is a vertex $v \in W_i \setminus V(a_i P a_{i+1})$. We consider two cases: when $v \in V(H)$ and when $v \notin V(H)$. In the former case, the component of H containing v must contain all its vertices in W_i as W_i has no marked vertices and $\{a_i, a_{i+1}\} \subseteq V(P) \subseteq V(H^0)$, and this is a contradiction. In the latter case, by Claim 1, every neighbor of v lies in X . But v cannot have neighbors in $X \cap Z$ because W_i does not contain marked vertices, and $X \cap W_i = \emptyset$ by our choice of α' . Hence v must be an isolated vertex in G , contradicting its 3-connectedness. This completes the proof that $a_i P a_{i+1}$ contains all vertices in W_i .

Note finally that $a_i P a_{i+1}$ is an induced path in G , as otherwise we could replace the component H^0 of H with a subdivision of L_k whose vertex set is a proper subset of $V(H^0)$ thus contradicting the minimality of $|V(H)|$.

Hence $P' := a_{\alpha'} P a_{\alpha'+2m+2}$ is an induced path in G of length at least $2m+2 \geq 2|Z|+3$ such that each internal vertex of P' has all its neighbors in $V(P') \cup Z$. By Lemma 19, there is an edge e of P' whose contraction preserves the 3-connectivity of G . As H^0/e is still isomorphic to a subdivision of L_k , we get a contradiction to the minimality of $|V(H)|$. The lemma follows. \diamond

By Claim 7 and the choice of λ_0 , there exists a set of indices I with $|I| \geq m^2 + 2k^3(m-1) + 1$ with the property that for every $i \in I$, there exists $j \in \{1, \dots, N\}$ such that $H_j \subseteq W_i$, and for distinct $i, i' \in I$, the sets W_i and $W_{i'}$ are disjoint, that is, $|i' - i| \geq 2$.

For each $i \in I$, let H^i be one of the graphs H_1, \dots, H_N that is contained in W_i . For every $i \in I$, fix a linkage Q_1^i, Q_2^i, Q_3^i from $\{a_i, a_{i+1}\} \cup Z$ to $V(H^i)$ in which the number of paths having an endpoint in $\{a_i, a_{i+1}\}$ is largest possible. Note that $V(Q_j^i) \subseteq W_i \cup Z$ for $j = 1, 2, 3$. We can classify the linkages Q_1^i, Q_2^i, Q_3^i by whether zero, one or two of the paths have an endpoint in $\{a_i, a_{i+1}\}$. For $j \in \{0, 1, 2\}$, define $I_j \subseteq I$ to be the subset of indices $i \in I$ such that exactly j of the paths Q_1^i, Q_2^i, Q_3^i have an endpoint in $\{a_i, a_{i+1}\}$.

Claim 8. $I_0 = \emptyset$.

Proof. Suppose to the contrary that there exists $i \in I_0$. Let R be a path from $\{a_i, a_{i+1}\}$ to $V(H^i \cup Q_1^i \cup Q_2^i \cup Q_3^i)$ in W_i . The path R intersects at most one of the paths Q_1^i, Q_2^i, Q_3^i , so without loss of generality we may assume that R does not intersect $Q_1^i \cup Q_2^i$. In $Q_3^i \cup R$ we can find a path Q' from $\{a_i, a_{i+1}\}$ to H^i . The path Q' does not intersect Q_1^i nor Q_2^i , so after replacing Q_3^i with Q' we obtain a linkage with one more path with an endpoint in $\{a_i, a_{i+1}\}$, contradiction. \diamond

Claim 9. $|I_1| \leq m^2$.

Proof. Without loss of generality, assume that for every $i \in I_1$, the path Q_1^i has an endpoint in $\{a_i, a_{i+1}\}$. For every $i \in I_1$, fix R_i to be a path from $\{a_i, a_{i+1}\} \setminus V(Q_1^i)$ to $V(H^i \cup Q_1^i \cup Q_2^i \cup Q_3^i)$ in W_i . Since the linkage Q_1^i, Q_2^i, Q_3^i maximizes the number of endpoints in $\{a_i, a_{i+1}\}$, the path R_i must have an endpoint in Q_1^i , as otherwise we would reroute one of the paths Q_2^i or Q_3^i to $\{a_i, a_{i+1}\} \setminus V(Q_1^i)$. Thus, $Q_1^i \cup R_i$ is a tree, which contains a_i, a_{i+1} and a vertex of H^i . Let P be an a_1 - a_λ path in $G - Z$ such that for every $i \in I_1$, we have $P[W_i \cap V(P)] \subseteq Q_1^i \cup R_i$.

Let F be the union of P and all paths Q_j^i with $i \in I_1$ and $j \in \{1, 2, 3\}$. Every component of F is either a union of paths having a common endpoint in a vertex from Z but otherwise disjoint, or the union of P and paths Q_1^i with $i \in I_1$. Thus F has at most $|Z| + 1$ components, and each H^i with $i \in I_1$ intersects three components of F , each in one vertex. The graph G does not have an L_{k+1} minor, so by Lemma 23 applied to the graphs H_i with $i \in I_1$, we have $|I_1| \leq (|Z| + 1)^2 \leq m^2$. \diamond

By Claims 8 and 9 and the bound on $|I|$, we see that $|I_2| > 2k^3(m-1)$. Without loss of generality, for all $i \in I_2$, assume that Q_1^i has a_i as

an endpoint and Q_2^i has a_{i+1} as an endpoint. It follows that Q_3^i has an endpoint in Z . For every $i \in I_2$, fix an L_k -model ϕ^i of L_k in H^i , and for each $j \in \{1, 2, 3\}$, let $t(i, j)$ be the index such that Q_j^i has an endpoint in $\phi^i((1, t(i, j))) \cup \phi^i((2, t(i, j)))$. Thus, $t(i, j) \in \{1, \dots, k\}$ for all i and j .

As $|I_2| > 2k^3(m-1)$, there exist indices i, i' and i'' in I_2 with $i < i' < i''$ and a vertex $z \in Z$ such that $t(i, j) = t(i', j) = t(i'', j)$ for all $j \in \{1, 2, 3\}$, and the paths $Q_3^i, Q_3^{i'}$ and $Q_3^{i''}$ all have z as an endpoint. Using the symmetries of a ladder, we may assume that $t(i, 1) \leq t(i, 2)$, and thus $t(i', 1) \leq t(i', 2)$ and $t(i'', 1) \leq t(i'', 2)$.

There are now three cases to consider: $t(i, 3) \leq t(i, 1)$, $t(i, 1) < t(i, 3) < t(i, 2)$, and $t(i, 2) \leq t(i, 3)$. In each case, we find two paths R_1 and R_2 linking H^i and $H^{i''}$ such that by joining a rooted ladder minor in H^i to a rooted ladder minor in $H^{i''}$, we show that G has an L_{k+1} minor, yielding a contradiction. The rooted ladder minors will be obtained by applying Lemma 21.




FIGURE 8. Illustration of Case 1. The paths R_1 and R_2 are depicted in bold.

Case 1: $t(i, 3) \leq t(i, 1)$. See Figure 8 for an illustration of this case. Let R_1 be a $V(H^i)$ – $V(H^{i''})$ path contained in $V(Q_3^i \cup Q_3^{i'} \cup H^{i'} \cup Q_2^{i'}) \cup W_{i'+1} \cup \dots \cup W_{i''-1} \cup V(Q_1^{i''})$. Thus, R_1 links a vertex of $\phi^i((1, t(i, 3))) \cup \phi^i((2, t(i, 3)))$ to a vertex of $\phi^{i''}((1, t(i'', 1))) \cup \phi^{i''}((2, t(i'', 1)))$ and is internally disjoint from $H^i \cup H^{i''}$. The path R_1 has only one vertex not contained in C , namely z . Hence by our choice of i_0 , there exists a path P between $W_0 \cup \dots \cup W_{i_0-1}$ and $W_{i_0+\lambda_0+1} \cup \dots \cup W_\lambda$ that is internally disjoint from C and does not contain the vertex z . Let R_2 be a $V(H^i)$ – $V(H^{i''})$ path contained in $V(Q_1^i) \cup W_0 \cup \dots \cup W_{i-1} \cup V(P) \cup W_{i''+1} \cup \dots \cup W_\lambda \cup V(Q_2^{i''})$. Thus, the path R_2 links a vertex of $\phi^i((1, t(i, 1))) \cup \phi^i((2, t(i, 1)))$ to a vertex of $\phi^{i''}((1, t(i'', 2))) \cup \phi^{i''}((2, t(i'', 2)))$ and is internally disjoint from $H^i \cup H^{i''}$ and completely disjoint from R_1 . By Lemma 21, H^i contains an $L_{k-t(i,1)+1}$ -model rooted on the endpoints of R_1 and R_2 , and $H^{i''}$ has an $L_{t(i'',1)}$ -model rooted on the endpoints of R_1 and R_2 . Together, we see that G contains an L_{k+1} minor.

Case 2: $t(i, 1) < t(i, 3) < t(i, 2)$. Let R_1 be the path formed by the union of Q_3^i and $Q_3^{i''}$. Thus, R_1 links a vertex of $\phi^i((1, t(i, 3))) \cup \phi^i((2, t(i, 3)))$ to a vertex of $\phi^{i''}((1, t(i'', 3))) \cup \phi^{i''}((2, t(i'', 3)))$ and is internally disjoint from $H^i \cup H^{i''}$. Let R_2 be a $V(H^i)$ – $V(H^{i''})$ path contained in

$V(Q_2^i) \cup W_{i+1} \cup \dots \cup W_{i''-1} \cup V(Q_1^{i''})$. The path R_2 links a vertex of $\phi^i((1, t(i, 2))) \cup \phi^i((2, t(i, 2)))$ to a vertex of $\phi^{i''}((1, t(i'', 1))) \cup \phi^{i''}((2, t(i'', 1)))$. By Lemma 21, H_i contains an $L_{t(i,3)}$ -model rooted on the endpoints of R_1 and R_2 , and $H_{i''}$ contains an $L_{k-t(i'',3)+1}$ -model rooted on the endpoints of R_1 and R_2 , implying that G contains an L_{k+1} minor.

Case 3: $t(i, 2) \leq t(i, 3)$. Let R_1 be a $V(H^i)$ – $V(H^{i''})$ path contained in $V(Q_2^i) \cup W_{i+1} \cup \dots \cup W_{i'-1} \cup V(Q_1^{i'} \cup H^{i'} \cup Q_3^{i'} \cup Q_3^{i''})$. Thus, R_1 links a vertex of $\phi^i((1, t(i, 2))) \cup \phi^i((2, t(i, 2)))$ to a vertex of $\phi^{i''}((1, t(i'', 3))) \cup \phi^{i''}((2, t(i'', 3)))$ and is internally disjoint from $H^i \cup H^{i''}$. The path R_1 has only one vertex not contained in C , namely z . Hence by our choice of i_0 , there exists a path P between $W_0 \cup \dots \cup W_{i_0-1}$ and $W_{i_0+\lambda_0+1} \cup \dots \cup W_\lambda$ which is internally disjoint from C and does not contain the vertex z . Let R_2 be a $V(H^i)$ – $V(H^{i''})$ path contained in $V(Q_1^i) \cup W_0 \cup \dots \cup W_{i-1} \cup V(P) \cup W_{i''+1} \cup \dots \cup W_\lambda \cup V(Q_2^{i''})$. Thus, the path R_2 links a vertex of $\phi^i((1, t(i, 1))) \cup \phi^i((2, t(i, 1)))$ to a vertex of $\phi^{i''}((1, t(i'', 2))) \cup \phi^{i''}((2, t(i'', 2)))$ and is internally disjoint from $H^i \cup H^{i''}$ and completely disjoint from R_1 . By Lemma 21, H^i contains an $L_{k-t(i,2)+1}$ -model of rooted on the endpoints of R_1 and R_2 , and $H^{i''}$ has an $L_{t(i'',2)+1}$ -model rooted on the endpoints of R_1 and R_2 . Together, we see that G contains an L_{k+1} minor.

In each case, we showed that G contains an L_{k+1} minor, and thus, in each case we arrive at a contradiction to our assumptions on the graph G , completing the proof of the theorem. \square

ACKNOWLEDGEMENTS

We are much grateful to the three anonymous referees for their careful reading and very helpful comments.

REFERENCES

- [1] Kiyoshi Ando, Hikoe Enomoto, and Akira Saito. Contractible edges in 3-connected graphs. *Journal of Combinatorial Theory, Series B*, 42(1):87–93, 1987.
- [2] Reinhard Diestel. *Graph Theory*. Springer Publishing Company, Incorporated, 5th edition, 2017.
- [3] P. Erdős and G. Szekeres. A combinatorial problem in geometry. *Compositio Math.*, 2:463–470, 1935.
- [4] Maximilian Gorsky and Michał T. Seweryn. Posets with k -outerplanar cover graphs have bounded dimension. [arXiv:2103.15920](https://arxiv.org/abs/2103.15920).
- [5] R. Halin. Untersuchungen über minimale n -fach zusammenhängende graphen. *Mathematische Annalen*, 182:175–188, 1969.
- [6] R. Halin. Zur theorie der n -fach zusammenhängenden graphen. *Abh. Math. Sem. Univ. Hamburg*, 33:133–164, 1969.
- [7] David M. Howard, Noah Streib, William T. Trotter, Bartosz Walczak, and Ruidong Wang. Dimension of posets with planar cover graphs excluding two long incomparable chains. *Journal of Combinatorial Theory, Series A*, 164:1–23, 2019.
- [8] Gwenaél Joret, Piotr Micek, Patrice Ossona de Mendez, and Veit Wiechert. Nowhere dense graph classes and dimension. *Combinatorica*, 39(5):1055–1079, 2019. [arXiv:1708.05424](https://arxiv.org/abs/1708.05424).
- [9] Gwenaél Joret, Piotr Micek, Kevin G. Milans, William T. Trotter, Bartosz Walczak, and Ruidong Wang. Tree-width and dimension. *Combinatorica*, 36(4):431–450, 2016. [arXiv:1301.5271](https://arxiv.org/abs/1301.5271).

- [10] Gwenaël Joret, Piotr Micek, William T. Trotter, Ruidong Wang, and Veit Wiechert. On the dimension of posets with cover graphs of treewidth 2. *Order*, 34(2):185–234, 2017. [arXiv:1406.3397](#).
- [11] Gwenaël Joret, Piotr Micek, and Veit Wiechert. Planar posets have dimension at most linear in their height. *SIAM J. Discrete Math.*, 31(4):2754–2790, 2018. [arXiv:1612.07540](#).
- [12] Gwenaël Joret, Piotr Micek, and Veit Wiechert. Sparsity and dimension. *Combinatorica*, 38(5):1129–1148, 2018. [arXiv:1507.01120](#).
- [13] David Kelly. On the dimension of partially ordered sets. *Discrete Mathematics*, 35:135–156, 1981.
- [14] Jakub Kozik, Piotr Micek, and William T. Trotter. Dimension is polynomial in height for posets with planar cover graphs. [arXiv:1907.00380](#).
- [15] Piotr Micek and Veit Wiechert. Topological minors of cover graphs and dimension. *Journal of Graph Theory*, 86(3):295–314, 2017. [arXiv:1504.07388](#).
- [16] Jaroslav Nešetřil and Patrice Ossona de Mendez. *Sparsity*, volume 28 of *Algorithms and Combinatorics*. Springer, Heidelberg, 2012. Graphs, structures, and algorithms.
- [17] Neil Robertson and Paul D. Seymour. Graph minors. V. Excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 41(1):92–114, 1986.
- [18] Michał T. Seweryn. Improved bound for the dimension of posets of treewidth two. *Discrete Mathematics*, 343(1):111605, 2020. [arXiv:1902.01189](#).
- [19] Noah Streib and William T. Trotter. Dimension and height for posets with planar cover graphs. *European J. Combin.*, 35:474–489, 2014.
- [20] William T. Trotter, Bartosz Walczak, and Ruidong Wang. Dimension and cut vertices: an application of Ramsey theory. In *Connections in Discrete Mathematics*, pages 187–199. Cambridge Univ. Press, Cambridge, 2018. [arXiv:1505.08162](#).
- [21] Bartosz Walczak. Minors and dimension. *J. Combin. Theory Ser. B*, 122:668–689, 2017. [arXiv:1407.4066](#).