Limit Theorems for Branching Processes with Immigration in a Random Environment

Bojan Basrak* and Péter Kevei[†] January 10, 2022

Abstract

We investigate subcritical Galton–Watson branching processes with immigration in a random environment. Using Goldie's implicit renewal theory we show that under general Cramér condition the stationary distribution has a power law tail. We determine the tail process of the stationary Markov chain, prove point process convergence, and convergence of the partial sums. The original motivation comes from Kesten, Kozlov and Spitzer seminal 1975 paper, which connects a random walk in a random environment model to a special Galton–Watson process with immigration in a random environment. We obtain new results even in this very special setting.

Keywords: branching process in random environment, regularly varying stationary sequences, tail process, implicit renewal theory $MSC2010:\ 60J80,\ 60F05$

1 Introduction and notation

Kesten et al. in their article [25] on random walks in a random environment discovered close connections of such walks with the so-called stochastic recurrence equations, and with a class of branching processes with immigration in a random environment. In this article, we aim to give a precise description of the long term behavior of such branching processes. It is known that in an i.i.d. random environment, a branching process with immigration, (X_n) say, has a Markovian structure i.e. it satisfies recursion $X_n = \psi(X_{n-1}, Z_n)$ for

^{*}Department of Mathematics, Faculty of Science, University of Zagreb, Bijenička 30, 10000 Zagreb, Croatia; e-mail: bbasrak@math.hr

[†]Bolyai Institute, University of Szeged, Aradi vértanúk tere 1, 6720 Szeged, Hungary; e-mail: kevei@math.u-szeged.hu

an i.i.d. sequence of random elements (Z_n) . Although, its evolution seems somewhat more involved than the one described by the standard stochastic recurrence equations (as studied in [11] for instance), we show in Section 2 that Goldie's implicit renewal theory of [19] can be readjusted to characterize the tails of the corresponding stationary distribution. Next, in Sections 3 and 4 respectively, we give detailed description of the asymptotic limits of properly normalized values (X_i) for $i = 1, \ldots, n$, and of their partial sums $S_n = X_1 + \cdots + X_n$, showing that the latter can exhibit both Gaussian and non-Gaussian limits under appropriate conditions. Note that Roitershtein [34] showed Gaussian asymptotics for partial sums of certain multitype branching processes with immigration in a more general random environment. As an application of our main results, in Section 5 we reconsider Kesten et al.'s [25] random walk in a random environment model with positive drift and present an alternative analysis of the asymptotic behavior of such a walk. We explain how their conditions yield various limiting distributions of such walks with the emphasis on the arguably more interesting non-Gaussian case. Moreover, we show that one can relax original conditions in [25] somewhat, and also write out the characteristic function of the limiting distributions. The precise form of the characteristic function which follows from our analysis seems to be new. Finally, our method additionally yields the long term behavior of the worst traps such a random walk encounters when moving to the right. It is already understood that it is exactly those traps, that is the edges visited over and over again as the walk moves to the right, which give rise to the non-Gaussian limits for random walks in random environment.

Throughout the article, the random environment is modeled by an i.i.d. sequence ξ , $(\xi_t)_{t\in\mathbb{Z}}$ with values in a measurable space \mathbb{X} . It may help in the sequel to specify $\mathbb{X}=\Delta^2$, where Δ denotes the space of probability measures on $\mathbb{N}=\{0,1,\ldots\}$, in that case we may write $\xi=(\nu_\xi,\nu_\xi^\circ)$. Alternatively, we assume that there exist a measurable function which maps each ξ to a pair $(\nu_\xi,\nu_\xi^\circ)\in\Delta^2$. The two components of the pair (ν_ξ,ν_ξ°) are called the offspring and the immigration distribution, respectively. The Galton–Watson process with immigration in the random environment $\mathcal{E}=\sigma\{\xi_t:t\in\mathbb{Z}\}$ evolves as follows. Let $X_0=x\in\mathbb{N}$, and then set

$$X_{n+1} = \sum_{i=1}^{X_n} A_i^{(n+1)} + B_{n+1} =: \theta_{n+1} \circ X_n + B_{n+1}, \quad n \ge 0,$$
 (1)

where we assume that, conditioned on the environment \mathcal{E} , the variables $\{A_i^{(n)}, B_n : n \in \mathbb{Z}, i \geq 1\}$ are independent, moreover, for n fixed, $(A_i^{(n)})_{i \geq 1}$

are i.i.d. with distribution ν_{ξ_n} , and B_n has distribution $\nu_{\xi_n}^{\circ}$. We interpret the variable $A_i^{(n)}$ as the number of offspring of the *i*th element in the (n-1)st generation, and B_n as the number of immigrants in the *n*th generation. Hence, θ_n as in (1) denote random operators on nonnegative integers which act as follows

$$\theta_n \circ x = \sum_{i=1}^x A_i^{(n)} .$$

In the sequel, we abuse the notation, by writing $\theta_n \circ (x+y) = \theta_n \circ x + \theta_n \circ y$, keeping in mind that the two random operators on the right hand side are not really the same.

For an X-valued random element ξ denote by

$$m(\xi) = \sum_{i=1}^{\infty} i\nu_{\xi}(\{i\}), \quad m^{\circ}(\xi) = \sum_{i=1}^{\infty} i\nu_{\xi}^{\circ}(\{i\}),$$

the expectation of its offspring and immigration distribution. This is clearly a nonnegative random variable, potentially equal to $+\infty$. We assume in the sequel that the process is subcritical, i.e.

$$\mathbf{E}\log m(\xi) < 0,\tag{2}$$

and that the following Cramér's condition holds

$$\mathbf{E}m(\xi)^{\kappa} = 1 \quad \text{for some } \kappa > 0. \tag{3}$$

Finally, to avoid trivial situations, we assume that ν_{ξ}° is not concentrated at 0.

Note that by our assumptions the sequence of random elements $Z_n = (\xi_n, B_n, A_1^{(n)}, A_2^{(n)}, \ldots), \ n \in \mathbb{Z}$ with values in $\mathbb{X} \times \mathbb{N}^{\mathbb{N}}$ are iid. Therefore, one can represent the evolution of the process (X_n) using a measurable mapping ψ and i.i.d. sequence (Z_n) as

$$X_{n+1} = \psi(X_n, Z_{n+1}) = \Psi_{n+1}(X_n) = \theta_{n+1} \circ X_n + B_{n+1}, \tag{4}$$

emphasizing the Markovian character of the process (X_n) . Here Ψ_n denotes a random map $x \mapsto \psi(x, Z_n)$.

By iterating (1) backward one may expect that the stationary distribution of the process can be found as

$$X_{\infty} = B_0 + \theta_0 \circ B_{-1} + \theta_0 \circ \theta_{-1} \circ B_{-2} + \dots \stackrel{d}{=} \sum_{i=0}^{\infty} \theta_0 \circ \theta_1 \circ \dots \circ \theta_{i-1} \circ B_i, \quad (5)$$

provided that the r.h.s converges a.s. to a finite limit (cf. Lemma 1). Clearly, if well-defined, such a distribution satisfies the distributional fixed point equation

$$X \stackrel{d}{=} \psi(X, Z) = \sum_{i=1}^{X} A_i + B = \theta \circ X + B, \tag{6}$$

where $Z = (\xi, B, A_1, A_2, ...)$ and X on the right-hand side are independent.

2 Implicit renewal theory

2.1 Moments of the stationary distribution

One of the main steps in the analysis of the process (X_n) is to determine whether the random variable X_{∞} in (5) has finite moments of order $\alpha > 0$ say. If this holds, X_{∞} would be clearly finite with probability one. On the other hand, by the conditional Jensen inequality $\mathbf{E}m(\xi)^t \leq \mathbf{E}A^t$ for $t \geq 1$, while $\mathbf{E}m(\xi)^t \geq \mathbf{E}A^t$ for $t \leq 1$. In particular, for any fixed t > 0, $\mathbf{E}A^t = \infty$ is possible with the assumptions (2) and (3) still being satisfied for some $\kappa > 0$.

In deterministic environment for multitype processes, the existence and explicit expression for the moments of order α were subject of Quine [32] for $\alpha=1,2$, and of Barczy et al. [2, Lemma 1] for $\alpha=3$. Under additional ergodicity conditions, the corresponding result for multitype processes for general $\alpha>0$ was proved by Szűcs [36]. In our case, under the condition that $\alpha>0$ satisfies

$$\mathbf{E}m(\xi)^{\alpha} < 1$$
, $\mathbf{E}A^{\alpha} < \infty$, and $\mathbf{E}B^{\alpha} < \infty$, (7)

by Lemma 3.1 in Buraczewski and Dyszewski [12] one can find constants c>0 and $0<\varrho<1$ such that

$$\mathbf{E}(\theta_0 \circ \theta_1 \circ \dots \theta_{i-1} \circ B_i)^{\alpha} \le c\varrho^i.$$

Therefore, for $\alpha \geq 1$ by Minkowski's inequality

$$(\mathbf{E}X_{\infty}^{\alpha})^{1/\alpha} \leq \sum_{i=0}^{\infty} (\mathbf{E}(\theta_0 \circ \theta_1 \circ \dots \theta_{i-1} \circ B_i)^{\alpha})^{1/\alpha} \leq c^{1/\alpha} \sum_{i=0}^{\infty} \varrho^{i/\alpha} < \infty,$$

while for $\alpha < 1$, simply by subadditivity

$$\mathbf{E} X_{\infty}^{\alpha} \leq \sum_{i=0}^{\infty} \mathbf{E} (\theta_0 \circ \theta_1 \circ \dots \theta_{i-1} \circ B_i)^{\alpha} \leq c \sum_{i=0}^{\infty} \varrho^i < \infty,$$

which immediately yields the following useful result.

Lemma 1. If (7) holds for some $\alpha > 0$, then the random variable X_{∞} in (5) satisfies $\mathbf{E}X_{\infty}^{\alpha} < \infty$.

Lemma 1 implies in particular that (5) represents a solution of the distributional equation in (6). The next statement can be viewed as a slight extension of Theorem 1.5.1 in Gut [20]. Put $a \lor b = \max\{a, b\}$.

Lemma 2. Let Y be a nonnegative integer-valued random variable, A, A_1, A_2, \ldots are identically distributed random variables, independent of Y, such that they are i.i.d. given the environment ξ , $\mathbf{E}[A|\xi] = 0$, furthermore Y is independent of ξ .

(i) Let $\alpha \geq 2$. Assume that $\mathbf{E}Y^{\alpha/2} < \infty$, and $\mathbf{E}|A|^{\alpha} < \infty$. Then there is $a \ c = c(\alpha) > 0$ depending only on α such that

$$\mathbf{E} \left| \sum_{i=1}^{Y} A_i \right|^{\alpha} < c \, \mathbf{E} Y^{\alpha/2} \, \mathbf{E} |A|^{\alpha}.$$

(ii) Let $\alpha \in (0,2]$, $\eta \in [0,1]$ be such that $2\eta \leq \alpha \leq 1+\eta$, and $\mathbf{E}Y^{\alpha-\eta} < \infty$, and

$$\mathbf{E}\left(\mathbf{E}\left[|A|^{\frac{\alpha}{\alpha-\eta}}\Big|\xi\right]\right)^{\alpha-\eta}<\infty.$$

Then there is a $c = c(\alpha, \eta) > 0$ depending only on α and η such that

$$\mathbf{E} \left| \sum_{i=1}^{Y} A_i \right|^{\alpha} < c \, \mathbf{E} Y^{\alpha - \eta} \, \mathbf{E} \left(\mathbf{E} \left[|A|^{\frac{\alpha}{\alpha - \eta}} \Big| \xi \right] \right)^{\alpha - \eta}.$$

(iii) For $\alpha \in [1,2]$ assume that $\mathbf{E}Y < \infty$ and $\mathbf{E}|A|^{\alpha} < \infty$. Then there is a $c = c(\alpha) > 0$ such that

$$\mathbf{E} \left| \sum_{i=1}^{Y} A_i \right|^{\alpha} \le c \mathbf{E} Y \mathbf{E} |A|^{\alpha}.$$

Proof. (i) By Rosenthal's inequality [31, Theorem 2.10]

$$\mathbf{E} \Big| \sum_{i=1}^{n} A_i \Big|^{\alpha} = \mathbf{E} \left(\mathbf{E} \left[\Big| \sum_{i=1}^{n} A_i \Big|^{\alpha} \Big| \xi \right] \right) \le c \, n^{\alpha/2} \, \mathbf{E} |A|^{\alpha}. \tag{8}$$

From inequality (8) the statement follows easily, as

$$\mathbf{E} \Big| \sum_{i=1}^{Y} A_i \Big|^{\alpha} = \sum_{n=1}^{\infty} \mathbf{P}(Y=n) \, \mathbf{E} \Big| \sum_{i=1}^{n} A_i \Big|^{\alpha}$$

$$\leq c \, \mathbf{E} |A|^{\alpha} \sum_{n=1}^{\infty} \mathbf{P}(Y=n) \, n^{\alpha/2}$$

$$\leq c \, \mathbf{E} |A|^{\alpha} \, \mathbf{E} Y^{\alpha/2} < \infty.$$

(ii) First using the conditional Jensen inequality $(\alpha - \eta \le 1)$, then the Marcinkiewicz–Zygmund inequality [31, 2.6.18], and finally the subadditivity $(\alpha/[2(\alpha - \eta)] \le 1)$ we obtain

$$\begin{split} \mathbf{E} \Big| \sum_{i=1}^{Y} A_i \Big|^{\alpha} &= \mathbf{E} \bigg(\mathbf{E} \bigg[\Big| \sum_{i=1}^{Y} A_i \Big|^{\alpha} \Big| \, \xi, Y \bigg] \bigg) \\ &\leq \mathbf{E} \bigg(\mathbf{E} \bigg[\Big| \sum_{i=1}^{Y} A_i \Big|^{\frac{\alpha}{\alpha - \eta}} \Big| \, \xi, Y \bigg] \bigg)^{\alpha - \eta} \\ &\leq c \, \mathbf{E} \bigg(\mathbf{E} \bigg[\bigg(\sum_{i=1}^{Y} A_i^2 \bigg)^{\frac{\alpha}{2(\alpha - \eta)}} \Big| \, \xi, Y \bigg] \bigg)^{\alpha - \eta} \\ &\leq c \, \mathbf{E} Y^{\alpha - \eta} \, \mathbf{E} \left(\mathbf{E} \left[|A|^{\frac{\alpha}{\alpha - \eta}} \Big| \xi \right] \right)^{\alpha - \eta}. \end{split}$$

(iii) This follows from (ii) choosing $\eta = \alpha - 1$.

2.2 Goldie's condition

In (4) we described the evolution of the Markov chain (X_n) using an i.i.d. sequence of random functions (Ψ_n) , $\Psi_n: \Omega \times \mathbb{N} \to \mathbb{N}$, $n \in \mathbb{Z}$, having the following general form

$$\Psi_n(k) = \sum_{i=1}^k A_i^{(n)} + B_n, \quad k \in \mathbb{N}.$$

Clearly, distributional fixed point equation in (6) can be written as

$$X \stackrel{d}{=} \Psi(X), \tag{9}$$

with Ψ and X independent on the right-hand side.

Lemma 3. Assume that there exist $\kappa > 0$ such that $\mathbf{E}m(\xi)^{\kappa} = 1$, $\mathbf{E}A^{\kappa} < \infty$, $\mathbf{E}B^{\kappa} < \infty$. Then the law in (5) represents the unique stationary distribution for the Markov chain (X_n) . Suppose further that at least one of the following three conditions holds

- (i) $\kappa > 1$ and $\mathbf{E}(m(\xi)^{\kappa-1}(\mathbf{E}[A^2|\xi])^{1/2}) < \infty$;
- (ii) $\kappa > 1$ and there exists $\delta > 0$ such that $\mathbf{E} A^{\kappa + \delta} < \infty$;
- (iii) $\kappa \leq 1$ and there exists $\eta \in (0, \kappa/2]$ such that

$$\mathbf{E}\left(\mathbf{E}\left[A^{\frac{\kappa}{\kappa-\eta}}\,\Big|\,\xi\right]\right)^{\kappa-\eta}<\infty.$$

Then the random variable $X = X_{\infty}$ further satisfies

$$\mathbf{E} |\Psi(X)^{\kappa} - (m(\xi)X)^{\kappa}| < \infty.$$

Proof. It is straightforward to see that the assumptions of the lemma imply (7) for any $\alpha \in (0, \kappa)$, which, by Lemma 1 further proves that the Markov chain in (1) has a stationary distribution. Denote by

$$d_0 = \min\{k : \mathbf{P}(A_1 = 0, B_1 = k) > 0\}.$$

The nonnegative integer d_0 is well defined since $\mathbf{P}(A_1 = 0) > 0$ by the subcriticality assumption (2). Moreover, it represents an accessible atom for the Markov chain (X_n) , this makes the chain irreducible, and the stationary distribution unique, see Douc et al. [16, Theorem 7.2.1] for instance.

First consider the $\kappa>1$ case. We use that for any $\alpha\geq 1$ for some $c=c_{\alpha}>0$

$$|x^{\alpha} - y^{\alpha}| \le c|x - y| (y^{\alpha - 1} + |x - y|^{\alpha - 1}).$$

Therefore

$$\mathbf{E} |\Psi(X)^{\kappa} - (m(\xi)X)^{\kappa}|$$

$$\leq c \left(\mathbf{E} |\Psi(X) - m(\xi)X| (m(\xi)X)^{\kappa-1} + \mathbf{E} |\Psi(X) - m(\xi)X|^{\kappa} \right).$$
(10)

For the second term in (10) by Minkowski's inequality

$$(\mathbf{E}|\Psi(X) - m(\xi)X|^{\kappa})^{1/\kappa} \le \left(\mathbf{E}\Big|\sum_{i=1}^{X} (A_i - m(\xi))\Big|^{\kappa}\right)^{1/\kappa} + (\mathbf{E}B^{\kappa})^{1/\kappa}. \quad (11)$$

The second term is finite according to our assumptions. The finiteness of the first term in (11) follows from Lemmas 1 and 2. Indeed, for $\kappa \geq 2$ Lemma 2 (i), for $\kappa \in (1,2)$ Lemma 2 (iii) applies.

For the first term in (10) we have

$$\mathbf{E}\left(|\Psi(X) - m(\xi)X|(m(\xi)X)^{\kappa-1}\right)$$

$$\leq \mathbf{E}\left(\left|\sum_{i=1}^{X} (A_i - m(\xi))\right| (m(\xi)X)^{\kappa-1}\right) + \mathbf{E}B(m(\xi)X)^{\kappa-1}.$$
(12)

By independence and Hölder's inequality

$$\mathbf{E}B(m(\xi)X)^{\kappa-1} = \mathbf{E}X^{\kappa-1}\mathbf{E}Bm(\xi)^{\kappa-1}$$

$$\leq \mathbf{E}X^{\kappa-1}(\mathbf{E}B^{\kappa})^{1/\kappa}(\mathbf{E}m(\xi)^{\kappa})^{(\kappa-1)/\kappa} < \infty,$$

so the second term in (12) is finite. Therefore, it only remains to show the finiteness of the first term in (12).

Assume (i). Applying first the Marcinkiewicz–Zygmund inequality and then Jensen's inequality

$$\mathbf{E}\left[\left|\sum_{i=1}^{n}(A_{i}-m(\xi))\right|\left|\xi\right] \leq c\mathbf{E}\left[\left(\sum_{i=1}^{n}(A_{i}-m(\xi))^{2}\right)^{1/2}\left|\xi\right]\right]$$

$$\leq c\left(\mathbf{E}\left[\sum_{i=1}^{n}(A_{i}-m(\xi))^{2}\left|\xi\right|\right)^{1/2}$$

$$\leq c n^{1/2}\left(\mathbf{E}[A^{2}|\xi]\right)^{1/2}.$$

Substituting back into the first term in (12)

$$\mathbf{E}\left(\left|\sum_{i=1}^{X} (A_i - m(\xi))\right| (m(\xi)X)^{\kappa - 1}\right)$$

$$\leq c\mathbf{E}X^{\kappa - 1/2}\mathbf{E}\left(m(\xi)^{\kappa - 1}(\mathbf{E}[A^2|\xi])^{1/2}\right),$$

which is finite whenever (i) holds.

Assume (ii). For the first term in (12) by from Hölder's inequality we have

$$\mathbf{E}\left(\left|\sum_{i=1}^{X} (A_i - m(\xi))\right| (m(\xi)X)^{\kappa - 1}\right) \\
\leq \left(\mathbf{E}\left|\sum_{i=1}^{X} (A_i - m(\xi))\right|^p\right)^{1/p} \left(\mathbf{E}(m(\xi)X)^{q(\kappa - 1)}\right)^{1/q}, \tag{13}$$

with 1/p + 1/q = 1. Choose $p = \kappa + \varepsilon$, for some $0 < \varepsilon < \delta$, with $\delta > 0$ given in the condition (ii). Then easy computation shows that $q = \kappa/(\kappa - 1) - \varepsilon'$,

where $\varepsilon' \downarrow 0$ as $\varepsilon \downarrow 0$. Since $q(\kappa - 1) < \kappa$, the second factor is finite by the independence of X and ξ , and by Lemma 1. The finiteness of the first factor in (13) follows from Lemmas 1 and 2. Indeed, for $\kappa \geq 2$ this is immediate. For $\kappa \in (1,2)$ choose $p = \kappa + \varepsilon \leq 2$ and apply Lemma 2 (iii).

The case $\kappa \leq 1$ is simpler. By the inequality

$$|x^{\alpha} - y^{\alpha}| \le |x - y|^{\alpha},$$

we have

$$|\Psi(X)^{\kappa} - (m(\xi)X)^{\kappa}| \le |\Psi(X) - m(\xi)X|^{\kappa}.$$

Thus by subadditivity

$$\mathbf{E} \left| \Psi(X)^{\kappa} - (m(\xi)X)^{\kappa} \right| \le \mathbf{E} \left| \sum_{i=1}^{X} (A_i - m(\xi)) \right|^{\kappa} + \mathbf{E} B^{\kappa}.$$

Since the second term is finite by assumption, it is enough to show that

$$\mathbf{E} \left| \sum_{i=1}^{X} (A_i - m(\xi)) \right|^{\kappa} < \infty.$$

This follows from Lemma 2 (ii), with η given in condition (iii).

Remark 1. For special classes of offspring distributions the conditions of the lemma can be simplified. If,

$$\mathbf{E}[A^2|\xi] \le c(m(\xi)^2 + 1)$$
 a.s. for some $c > 1$, (14)

then both the condition for $\kappa \leq 1$ and condition (i) reduces to $\mathbf{E}m(\xi)^{\kappa} < \infty$, which holds since $\mathbf{E}m(\xi)^{\kappa} = 1$.

In particular if, conditionally on ξ , A has Poisson distribution with parameter $\lambda(\xi) > 0$, then $m(\xi) = \mathbf{E}[A|\xi] = \lambda(\xi)$ and $\mathbf{E}[A^2|\xi] = \lambda(\xi)^2 + \lambda(\xi) \le 2(m(\xi)^2 + 1)$. While if A, conditionally on ξ , has geometric distribution with parameter $p(\xi) \in (0,1)$, i.e. $\mathbf{P}(A=k\mid\xi) = (1-p(\xi))^k p(\xi)$, $k \ge 0$, then $m(\xi) = \mathbf{E}[A|\xi] = (1-p(\xi))/p(\xi)$ and $\mathbf{E}[A^2|\xi] = (3-2p(\xi)+p(\xi)^2)/p(\xi)^2 \le 3(1-p(\xi))^2/p(\xi)^2 + 3 = 3m(\xi)^2 + 3$. Thus, in both cases (14) holds.

From Goldie's Corollary 2.4 [19] we obtain the following.

Theorem 1. Assume the conditions of Lemma 3, $\mathbf{E}m(\xi)^{\kappa} \log m(\xi) < \infty$, and that the law of $\log m(\xi)$ given $m(\xi) > 0$ is nonarithmetic. Then the law

of X_{∞} in (5) represents the unique stationary distribution for the Markov chain (X_n) and

$$\mathbf{P}(X_{\infty} > x) \sim Cx^{-\kappa} \quad as \ x \to \infty, \tag{15}$$

where

$$C = \frac{1}{\kappa \mathbf{E} m(\xi)^{\kappa} \log m(\xi)} \mathbf{E} \left[\Psi(X_{\infty})^{\kappa} - m(\xi)^{\kappa} X_{\infty}^{\kappa} \right] > 0.$$

Proof. According to Lemma 3, the Markov chain (X_n) has unique stationary distribution given in (5). Because the law of the immigrant distribution is not concentrated at 0, this distribution is not trivial and therefore there exists $d = \min\{k > 0 : \mathbf{P}(X_{\infty} = k) > 0\} > 0$. The state d is necessarily positive recurrent, i.e. $\mathbf{E}_d(\tau_d) < \infty$ where $\tau_d = \min\{n \geq 1 : X_n = d\}$ denotes the return time to the state d.

From Lemma 3, we can conclude that the conditions of Corollary 2.4 in [19] hold which yields the tail asymptotics in (15). It remains to show the strict positivity of the constant C above. One can deduce this from Afanasyev's Theorem 1 [1] as follows.

To the original process X_n , one can couple a process Y_n , starting at $Y_0 = 1$, and satisfying

$$Y_{n+1} = \sum_{i=1}^{Y_n} A_i^{(n+1)}, \quad n \ge 0,$$

where the sequence of environments $(\xi_1, \xi_2, ...)$ are the same as in (1). Thus, (Y_n) is a subcritical branching process in random environment, such that by the construction $X_n \geq Y_n$ a.s. for every $n \geq 0$ provided that we start (X_n) at any state different from 0. Theorem 1 in [1] states that

$$\lim_{x \to \infty} x^{\kappa} \mathbf{P}(\sup_{n \ge 1} Y_n > x) = c > 0.$$
 (16)

Let $\tau_Y = \min\{n \geq 1 : Y_n = 0\}$, clearly $\tau_Y \leq \tau_d$. In particular, the process Y_n dies out almost surely. By the standard theory of Markov chains, see Theorem 7.2.1 in [16] or Theorem 10.4.9 in [30], we have

$$\mathbf{P}(X_{\infty} > x) = \frac{1}{\mathbf{E}_{d}\tau_{d}} \mathbf{E}_{d} \left[\sum_{i=0}^{\tau_{d}-1} \mathbb{1}\{X_{i} > x\} \right]$$

$$\geq \frac{1}{\mathbf{E}_{d}\tau_{d}} \mathbf{E} \left[\sum_{i=0}^{\tau_{Y}-1} \mathbb{1}\{Y_{i} > x\} \right] \geq \frac{1}{\mathbf{E}_{d}\tau_{d}} \mathbf{P}(\sup_{n \geq 1} Y_{n} > x),$$

which by (16) implies

$$\liminf_{x \to \infty} x^{\kappa} \mathbf{P}(X_{\infty} > x) > 0,$$

as claimed. \Box

Remark 2. As a consequence of the applied machinery, apart from some special cases, the constant C in the theorem is merely implicit; the formula contains the limit law itself. However, for $\kappa=1$ we simply have $\mathbf{E}[\Psi(X_{\infty})-m(\xi)X_{\infty}]=\mathbf{E}B$, so C can be computed.

Remark 3. Analogous results hold in the arithmetic case. Using Theorem 2 in [27] (see also Theorem 3.7 by Jelenković and Olvera-Cravioto [22]) one can show the following. If the conditions of Lemma 3 hold, and the law of $\log m(\xi)$ given $m(\xi) > 0$ is arithmetic with span h > 0, then there exists a function q such that

$$\lim_{n \to \infty} x^{\kappa} e^{\kappa nh} \mathbf{P}(X_{\infty} > x e^{nh}) = q(x),$$

whenever x is a continuity point of q. The function $x^{-\kappa}q(x)$ is nonincreasing, and $q(xe^h) = q(x)$ for all x > 0.

2.3 Relaxing Cramér's condition

In what follows, we weaken condition (3) in two ways, such that the tail of the stationary distribution is still regularly varying. We use a slight extension of Goldie's renewal theory by Kevei [26].

First we consider weakening the assumption $\mathbf{E}m(\xi)^{\kappa}\log m(\xi) < \infty$. The condition $\mathbf{E}m(\xi)^{\kappa} = 1$ ensures that

$$F_{\kappa}(x) = \mathbf{E}(\mathbb{1}\{\log m(\xi) \le x\} m(\xi)^{\kappa})$$

is a distribution function. The additional logarithmic moment condition in (3) is equivalent to the finiteness of the expectation of the distribution F_{κ} . This condition is needed to use the standard key renewal theorem. However, strong renewal theorems in the infinite mean case have been known since the 1962 paper by Garsia and Lamperti [18]. They showed that the strong renewal theorem holds if the underlying distribution belongs to the domain of attraction of an α -stable law with $\alpha \in (1/2,1]$, while for $\alpha \leq 1/2$ extra conditions are needed. Recently, Caravenna and Doney [13] obtained necessary and sufficient conditions for the strong renewal theorem to hold, solving a 50-year old open problem.

Assume that F_{κ} belongs to the domain of attraction of a stable law of index $\alpha \in (0,1]$, i.e., for some $\kappa > 0$ and $\alpha \in (0,1]$, for a slowly varying function ℓ

$$\mathbf{E}m(\xi)^{\kappa} = 1, \quad 1 - F_{\kappa}(x) = \frac{\ell(x)}{x^{\alpha}}.$$
 (17)

Define the truncated expectation as

$$M(x) = \int_0^x (1 - F_{\kappa}(x)) dx \sim \frac{\ell(x)x^{1-\alpha}}{1-\alpha},$$
 (18)

where the asymptotic equality holds for $\alpha < 1$. If $\alpha \in (0,1/2)$ further assume the Caravenna–Doney condition

$$\lim_{\delta \to 0} \limsup_{x \to \infty} x [1 - F_{\kappa}(x)] \int_{1}^{\delta x} \frac{1}{y [1 - F_{\kappa}(y)]^{2}} F_{\kappa}(x - dy) = 0.$$
 (19)

For our second extension, assume now that $\mathbf{E}m(\xi)^{\kappa} = \varphi < 1$ for some $\kappa > 0$. If $\mathbf{E}m(\xi)^t < \infty$ for some $t > \kappa$, then by Lemma 1 the tail of X_{∞} cannot be regularly varying with index κ . Therefore we assume F_{κ} is heavy-tailed, i.e. $\mathbf{E}m(\xi)^t = \infty$ for any $t > \kappa$. Define now the distribution function

$$F_{\kappa}(x) = \varphi^{-1} \mathbf{E}(\mathbb{1}\{\log m(\xi) \le x\} m(\xi)^{\kappa}). \tag{20}$$

The analysis of the stochastic fixed point equation (6) leads to a defective renewal equation. To understand the asymptotic behavior of the solution of these equations we need to introduce *locally subexponential distributions*.

For $T \in (0, \infty]$ let $\nabla_T = (0, T]$ and for a distribution function (df) H we put $H(x + \nabla_T) = H(x + T) - H(x)$. Let * denote the usual convolution operator. A df H is locally subexponential, $H \in \mathcal{S}_{loc}$, if for each $T \in (0, \infty]$ we have (i) $H(x+t+\nabla_T) \sim H(x+\nabla_T)$ as $x \to \infty$ uniformly in $t \in [0,1]$, (ii) $H(x+\nabla_T) > 0$ for x large enough, and (iii) $H*H(x+\nabla_T) \sim 2H(x+\nabla_T)$ as $x \to \infty$. For more details see Foss, Korshunov and Zachary [17, Section 4.7]. Informally, a locally subexponential distribution is a subexponential distribution with well-behaved density function.

Our assumptions on $m(\xi)$ are the following:

$$\mathbf{E}m(\xi)^{\kappa} = \varphi < 1, \ \kappa > 0, \quad F_{\kappa} \in \mathcal{S}_{loc},$$
 for each $T \in (0, \infty] \sup_{y>x} F_{\kappa}(y + \nabla_T) = O(F_{\kappa}(x + \nabla_T))$ as $x \to \infty$. (21)

Theorem 2. Assume that condition (i) or (iii) in Lemma 3 holds, the law of $\log m(\xi)$ given $m(\xi) > 0$ is nonarithmetic, $\mathbf{E}B^{\nu} < \infty$ for some $\nu > \kappa$, and one of the following two conditions is satisfied:

- (i) condition (17) holds, and if $\alpha \in (0, 1/2)$ also (19) holds;
- (ii) condition (21) holds.

Then the law of X_{∞} in (5) represents the unique stationary distribution for the Markov chain (X_n) and

$$\mathbf{P}(X_{\infty} > x) \sim Cx^{-\kappa}L(x)$$
 as $x \to \infty$,

where

$$L(x) = \begin{cases} (\Gamma(\alpha)\Gamma(2-\alpha)M(\log x))^{-1}, & \text{in case (i),} \\ (F_{\kappa}(1+\log x) - F_{\kappa}(\log x))\varphi/(1-\varphi)^{2}, & \text{in case (ii),} \end{cases}$$

is a slowly varying function, and

$$C = \frac{1}{\kappa} \mathbf{E} \left[\Psi(X_{\infty})^{\kappa} - m(\xi)^{\kappa} X_{\infty}^{\kappa} \right] \ge 0,$$

with C > 0 for $\kappa \geq 1$.

Proof. The result follows from Theorems 2.1 and 2.3 in [26]. We only have to check that

$$\mathbf{E} \left| \left(\sum_{i=1}^{X_{\infty}} A_i + B \right)^{\kappa + \delta} - \left(m(\xi) X_{\infty} \right)^{\kappa + \delta} \right| < \infty$$

for some $\delta > 0$. This can be done exactly the same way as in the proof of Lemma 3 case (i) and (iii).

The strict positivity of the constant C in the theorem can be deduced directly from its form. Indeed, since X_{∞} is independent of Ψ and ξ

$$\mathbf{E}\left[\Psi(X_{\infty})^{\kappa} - m(\xi)^{\kappa} X_{\infty}^{\kappa}\right] = \sum_{n=1}^{\infty} \mathbf{P}(X_{\infty} = n) \mathbf{E}\left[\left(\sum_{i=1}^{n} A_{i} + B\right)^{\kappa} - m(\xi)^{\kappa} n^{\kappa}\right]$$

$$> \sum_{n=1}^{\infty} \mathbf{P}(X_{\infty} = n) \mathbf{E}\left[\left(\sum_{i=1}^{n} A_{i}\right)^{\kappa} - m(\xi)^{\kappa} n^{\kappa}\right],$$
(22)

where in the last step we used that B is not identically 0. By Jensen's inequality

$$\mathbf{E}\left[\left(\sum_{i=1}^{n} A_{i}\right)^{\kappa} - m(\xi)^{\kappa} n^{\kappa}\right] = \mathbf{E}\left(\mathbf{E}\left[\left(\sum_{i=1}^{n} A_{i}\right)^{\kappa} \middle| \xi\right] - m(\xi)^{\kappa} n^{\kappa}\right)$$

$$\geq \mathbf{E}\left(n^{\kappa} m(\xi)^{\kappa} - m(\xi)^{\kappa} n^{\kappa}\right) = 0,$$

from which the strict positivity follows.

Remark 4. Note that in general we only proved the strict positivity of C above for $\kappa \geq 1$. However, in some special cases it is possible to show that C > 0 for a general $\kappa > 0$. In particular, if the expectation in the infinite sum in the first line of (22) is strictly positive for each n, then clearly C > 0.

Remark 5. Note that both remarks after Theorem 1 apply in this setup as well.

Remark 6. The condition $F_{\kappa} \in \mathcal{S}_{loc}$ is stronger than the regular variation condition, in particular Pareto distribution is locally subexponential. It is known that lognormal and Weibull distributions belong to the class \mathcal{S}_{loc} as well. In the Pareto case, i.e. if for large enough x we have $1 - F_{\kappa}(x) = c x^{-\beta}$, for some $c > 0, \beta > 0$, then $\mathbf{P}(X_{\infty} > x) \sim c' x^{-\kappa} (\log x)^{-\beta - 1}$. In the lognormal case, when $F_{\kappa}(x) = \Phi(\log x)$ for x large enough, with Φ being the standard normal df, we have $\mathbf{P}(X_{\infty} > x) \sim c x^{-\kappa} e^{-(\log \log x)^2/2}/\log x$, c > 0. While, for Weibull tails $1 - F_{\kappa}(x) = e^{-x^{\beta}}$, $\beta \in (0, 1)$, we obtain $\mathbf{P}(X_{\infty} > x) \sim c x^{-\kappa} (\log x)^{\beta - 1} e^{-(\log x)^{\beta}}$, c > 0.

3 Asymptotic behavior of the process (X_t)

3.1 Dependence structure of the process

In the sequel we only need that our process has a stationary distribution with regularly varying tail. In the previous section we derived several conditions which ensures regularly varying tail. Since in Theorem 2 we only have the strict positivity of the underlying constant C for $\kappa \geq 1$, we assume that one of the following holds:

conditions of Theorem 1, or conditions of Theorem 2 and C > 0. (23)

Then in particular, there exist a strictly stationary Galton–Watson process with immigration in random environment $(X_n)_{n\in\mathbb{Z}}$ which satisfies

$$X_{n+1} = \sum_{i=1}^{X_n} A_i^{(n+1)} + B_{n+1}, \quad n \in \mathbb{Z},$$
 (24)

with the same interpretation of the random variables $\{A_i^{(n)}, B_n : n \in \mathbb{Z}, i \geq 1\}$. Here again by ξ , $(\xi_i)_{i\in\mathbb{Z}}$ we denote i.i.d. random variables representing the environment. The offspring and immigration distributions are governed by the environment \mathcal{E} in the same way as before.

It is useful in the sequel to introduce a deterministic sequence $(a_n)_{n\in\mathbb{N}}$ such that

$$n\mathbf{P}(X_{\infty} > a_n) \longrightarrow 1 \quad \text{as } n \to \infty.$$
 (25)

Observe that by Theorem 1 and 2, (a_n) is a regularly varying sequence with index $1/\kappa$, i.e. $a_n = \tilde{\ell}(n)n^{1/\kappa}$, with an appropriate slowly varying function $\tilde{\ell}$. In particular, if the conditions of Theorem 1 hold we may set $a_n = (Cn)^{1/\kappa}$ and that any other sequence (a_n) in (25) necessarily satisfies $a_n \sim (Cn)^{1/\kappa}$.

Once we have shown that the marginal stationary distribution of the X_t 's is regularly varying, it is relatively easy to prove that all the finite dimensional distributions of (X_t) in (24) have multivariate regular variation property, cf. [33], i.e. $(X_n)_{n\in\mathbb{Z}}$ is regularly varying sequence in the sense of [7]. According to Theorem 2.1 in [7], this is equivalent to the existence of the so-called tail sequence, which is the content of the first theorem below.

Observe first that under Cramér's condition (3), one can construct a tilted distribution of the following form

$$\mathbf{P}(\xi^* \in \cdot) = \mathbf{E} \left(\mathbb{1} \{ \xi \in \cdot \} m(\xi)^{\kappa} \right) .$$

Similar change of measure appeared already in Afanasyev [1, Lemma 1], see also [12, Lemma 3.1]. By the convexity of the function $\lambda(\alpha) = \mathbf{E}m(\xi)^{\alpha}$

$$\mathbf{E}\log m(\xi^*) = \mathbf{E}m(\xi)^{\kappa}\log m(\xi) = \lambda'(\kappa) > 0,$$

possibly infinite, that is, the branching after the change of measure becomes supercritical.

To simplify notation, denote $m=m(\xi), m_i=m(\xi_i), i \geq 1$, under their original distribution. Introduce further an auxiliary i.i.d. sequence $m^*=m(\xi^*), m_i^*=m(\xi_i^*), i \geq 1$, with the common distribution $\mathbf{P}(m^* \in A) = \mathbf{E}[\mathbb{1}\{m(\xi) \in A\}m(\xi)^{\kappa}]$ and independent of (m_i) .

On the other hand, if (21) holds, i.e. $\mathbf{E}m(\xi)^{\kappa} = \varphi < 1$, then m_i^* , $i \geq 1$, is a sequence of extended random variables with common distribution $\mathbf{P}(m^* \in A) = \mathbf{E}[\mathbb{1}\{m(\xi) \in A\}m(\xi)^{\kappa}]$, for $A \subset \mathbb{R}$, and $\mathbf{P}(m^* = \infty) = 1 - \varphi$, and independent of (m_i) . In the following result we use the usual convention $1/\infty = 0$.

Theorem 3. Let $(X_n)_{n\in\mathbb{Z}}$ be a strictly stationary sequence satisfying (24). Assume (23), and let $(m_n)_{n\geq 1}$, $(m_n^*)_{n\geq 1}$ be independent i.i.d. sequences introduced above and independent of Y_0 with Pareto distribution $\mathbf{P}(Y_0 > u) = u^{-\kappa}$, $u \geq 1$. Then, for any integers $k, \ell \geq 0$

$$\mathcal{L}\left(\frac{X_{-k}}{x}, \dots, \frac{X_0}{x}, \dots, \frac{X_\ell}{x} \mid X_0 > x\right)$$

$$\xrightarrow{d} Y_0\left((m_k^* \cdots m_1^*)^{-1}, \dots, m_1^{*-1}, 1, m_1, \dots, m_1 \cdots m_\ell\right).$$

Writing the random vector on the r.h.s. above as $(Y_t, t = -k, ..., l)$, note that, in the language of Basrak and Segers [7], the sequence $(Y_t)_{t \in \mathbb{Z}}$ represents the tail process of the sequence (X_t) . Hence, in this case, both the forward and backward tail processes are multiplicative random walks.

Proof. We will first show that for arbitrary $\ell \geq 0$ and $\varepsilon > 0$

$$\mathbf{P}\left(\left|\frac{X_{\ell}}{X_0} - m_1 \cdots m_{\ell}\right| > \varepsilon \,\middle|\, X_0 > x\right) \longrightarrow 0, \text{ as } x \to \infty.$$
 (26)

Indeed, consider first $\ell = 1$, by the independence of $A_j^{(1)}$'s and B_1 from X_0 ,

$$\mathbf{P}\left(\left|\frac{X_{1}}{X_{0}}-m_{1}\right|>\varepsilon\left|X_{0}>x\right)\right)$$

$$=\sum_{k>x}\mathbf{P}\left(\left|\frac{\sum_{j=1}^{X_{0}}A_{j}^{(1)}+B_{1}}{X_{0}}-m_{1}\right|>\varepsilon;X_{0}=k\right)\frac{1}{\mathbf{P}\left(X_{0}>x\right)}.$$

$$=\sum_{k>x}\mathbf{P}\left(\left|\frac{\sum_{j=1}^{k}A_{j}^{(1)}+B_{1}}{k}-m_{1}\right|>\varepsilon\right)\frac{\mathbf{P}\left(X_{0}=k\right)}{\mathbf{P}\left(X_{0}>x\right)}.$$

By the ergodic theorem $n^{-1}\sum_{j=1}^n A_j^{(1)} \to m_1$ a.s. Hence the r.h.s. above tends to 0 as $x \to \infty$. Instead of general $\ell \geq 1$, consider for simplicity $\ell = 2$. One can write $X_2 = \sum_{j=1}^{X_0} \tilde{A}_j^{(2)} + \sum_{j=1}^{B_1} A_j^{(2)} + B_2$. Given the environment \mathcal{E} , the three terms on the r.h.s. are independent and $\tilde{A}_j^{(2)}$'s are i.i.d. with the following distribution

$$\tilde{A}_1^{(2)} \stackrel{d}{=} \sum_{j=1}^{A_1^{(1)}} A_j^{(2)}$$
.

In particular, $\mathbf{E}[\tilde{A}_1^{(2)} | \mathcal{E}] = m_1 m_2$, the ergodic theorem applied on the sequence $(\tilde{A}_j^{(2)})$ again yields

$$\mathbf{P}\left(\left|\frac{X_2}{X_0} - m_1 m_2\right| > \varepsilon \mid X_0 > x\right)$$

$$= \sum_{k>x} \mathbf{P}\left(\left|\frac{\sum_{j=1}^k \tilde{A}_j^{(2)} + \sum_{j=1}^{B_1} A_j^{(2)} + B_2}{k} - m_1 m_2\right| > \varepsilon\right) \frac{\mathbf{P}\left(X_0 = k\right)}{\mathbf{P}\left(X_0 > x\right)} \to 0$$

as $x \to \infty$. The same argument works for any $\ell \ge 1$, which implies (26). Observing that $\mathcal{L}(X_0/x \mid X_0 > x) \stackrel{d}{\longrightarrow} Y_0$, we can conclude that for an

arbitrary ℓ

$$\mathcal{L}\left(\frac{X_0}{x}, \frac{X_1}{X_0}, \dots, \frac{X_\ell}{X_0} \mid X_0 > x\right) \stackrel{d}{\longrightarrow} (Y_0, m_1, \dots, m_1 \cdots m_\ell).$$

This proves the statement of the theorem for k = 0. In particular, the sequence (X_t) is regularly varying, and the multiplicative random walk

$$Y_i = Y_0 \Theta_i = Y_0 m_1 \cdots m_i \,, \qquad i \ge 0 \tag{27}$$

represents the forward part of its tail process. By Theorem 3.1 in [7], this uniquely determines the distribution of the whole tail process, including the negative indices. The past distribution of the tail process has been determined already (see Theorem 5.2 and Example 3.3 in Segers [35] for instance). It turns out that the backward part of the tail process has the representation

$$Y_{-k} = Y_0 \Theta_{-k} = Y_0 / (m_k^* \cdots m_1^*), \qquad k > 0$$
 (28)

for an i.i.d. sequence (m_n^*) as in the statement of the theorem.

Alternatively, the theorem above can be obtained using the general results by Janssen and Segers [21, Theorem 2.1].

Next we prove that the large values in the sequence (X_n) cannot linger for "too long", i.e. the so called anticlustering condition from [15] holds.

Lemma 4. Let r_n be a sequence such that $r_n = o(n)$. Then for any u > 0

$$\lim_{k \to \infty} \limsup_{n \to \infty} \mathbf{P} \left(\max_{k \le |t| \le r_n} X_t > a_n u \, \middle| \, X_0 > a_n u \right) = 0.$$

Proof. The proof is similar to the proof of Lemma 3.2 in [4]. By the strict stationarity

$$\mathbf{P}\left(\max_{k \le |t| \le r_n} X_t > a_n u \mid X_0 > a_n u\right) \le 2 \sum_{t=k}^{r_n} \mathbf{P}(X_t > a_n u \mid X_0 > a_n u).$$

Consider the decomposition

$$X_{t} = \sum_{i=1}^{X_{0}} \widetilde{A}_{i}^{(t)} + \sum_{k=0}^{t-1} C_{t,k},$$
(29)

where the first sum stands for the descendants of generation 0, while the second sum stands for the descendants of the immigrants arrived after generation 0, i.e. $C_{t,k} = \theta_t \circ \cdots \circ \theta_{t-k+1} \circ B_{t-k}$ denotes the number of descendants in generation t from immigrants arrived in generation t-k. Then X_0 and $(C_{t,k})_{k=0,1,\ldots,t-1}$ are independent, and $(\widetilde{A}_i^{(t)})_{i\in\mathbb{N}}$ are independent given the environments ξ_1,\ldots,ξ_t . Then

$$\mathbf{P}(X_t > a_n u | X_0 > a_n u) \le \mathbf{P}\left(\sum_{k=0}^{t-1} C_{t,k} > a_n u / 2 \, \big| \, X_0 > a_n u\right) + \mathbf{P}\left(\sum_{i=1}^{X_0} \widetilde{A}_i^{(t)} > a_n u / 2 \, \big| \, X_0 > a_n u\right).$$

Using the independence of X_0 and $(C_{t,k})_{k=0,1,\dots,t-1}$

$$\mathbf{P}\left(\sum_{k=0}^{t-1} C_{t,k} > a_n u/2 \,\middle|\, X_0 > a_n u\right) = \mathbf{P}\left(\sum_{k=0}^{t-1} C_{t,k} > a_n u/2\right)$$

$$\leq \mathbf{P}\left(\sum_{k=0}^{\infty} C_{t,k} > a_n u/2\right) = \mathbf{P}\left(X_0 > a_n u/2\right).$$

By (25)

$$\lim_{n\to\infty} \sup r_n \mathbf{P}\left(X_0 > a_n u/2\right) = 0,$$

so it remains to show that

$$\lim_{k\to\infty}\limsup_{n\to\infty}\sum_{t=k}^{r_n}\mathbf{P}\left(\sum_{i=1}^{X_0}\widetilde{A}_i^{(t)}>a_nu/2\,\Big|\,X_0>a_nu\right)=0.$$

Choose $\alpha < \min\{\kappa, 1\}$. Then, by the convexity of the function $u \mapsto \mathbf{E} m(\xi)^u$ we obtain $\mathbf{E} m(\xi)^{\alpha} < 1$. Note that

$$\mathbf{E}[\widetilde{A}_i^{(t)}|\xi_1,\ldots,\xi_t] = m(\xi_1)\ldots m(\xi_t).$$

Therefore,

$$\begin{split} &\mathbf{E}\left[\left(\frac{\sum_{i=1}^{k}\widetilde{A}_{i}^{(t)}}{km(\xi_{1})\dots m(\xi_{t})}\right)^{\alpha}\Big|\xi_{1},\dots,\xi_{t}\right]\\ &\leq &\mathbf{E}\left[\frac{\sum_{i=1}^{k}\widetilde{A}_{i}^{(t)}}{km(\xi_{1})\dots m(\xi_{t})}\mathbb{1}\left\{\sum_{i=1}^{k}\widetilde{A}_{i}^{(t)}>km(\xi_{1})\dots m(\xi_{t})\right\}+1\Big|\xi_{1},\dots,\xi_{t}\right]\leq 2, \end{split}$$

which implies that

$$\mathbf{E}\left(\sum_{i=1}^k \widetilde{A}_i^{(t)}\right)^{\alpha} \leq 2k^{\alpha}\mathbf{E}\left(m(\xi_1)^{\alpha}\dots m(\xi_t)^{\alpha}\right) = 2k^{\alpha}\left[\mathbf{E}m(\xi)^{\alpha}\right]^t.$$

Note that the upper bound is summable in t, thus, by Markov's inequality

$$\sum_{t=k}^{r_n} \mathbf{P} \left(\sum_{i=1}^{X_0} \widetilde{A}_i^{(t)} > a_n u/2 \, \middle| \, X_0 > a_n u \right)$$

$$= \frac{1}{\mathbf{P}(X_0 > a_n u)} \sum_{t=k}^{r_n} \sum_{\ell > a_n u} \mathbf{P}(X_0 = \ell) \, \mathbf{P} \left(\sum_{i=1}^{\ell} \widetilde{A}_i^{(t)} > a_n u/2 \right)$$

$$\leq \frac{1}{\mathbf{P}(X_0 > a_n u)} \sum_{t=k}^{r_n} \sum_{\ell > a_n u} \frac{2^{1+\alpha}}{(a_n u)^{\alpha}} \ell^{\alpha} \left[\mathbf{E} m(\xi)^{\alpha} \right]^t$$

$$\leq 2^{1+\alpha} \frac{\mathbf{E} X_0^{\alpha} \mathbb{1} \{ X_0 > a_n u \}}{(a_n u)^{\alpha} \mathbf{P}(X_0 > a_n u)} \sum_{t=k}^{\infty} \left[\mathbf{E} m(\xi)^{\alpha} \right]^t.$$

As $n \to \infty$ the second factor here tends to $\kappa/(\kappa-\alpha)$ by Karamata's theorem, while the third factor vanishes as $k \to \infty$. Thus the result follows.

Recall that by our assumptions on progeny and immigrant distributions, there exists α , $0 < \alpha < \kappa$, $\alpha \le 1$, such that

$$\mathbf{E}m(\xi)^{\alpha} < 1$$
 and $\mathbf{E}m^{\circ}(\xi)^{\alpha} < \infty$.

Denote the Markov transition kernel of the sequence (X_n) by $P(x,\cdot)$ and consequently, by $P^n(x,\cdot)$, $n \geq 1$ denote the n-step Markov transition kernel corresponding to P. Next we apply the standard drift method to show that the Markov chain (X_n) is uniformly V-geometrically ergodic. First we introduce some notation. For a function $V: \mathbb{N} \to [1, \infty)$, and any two probability measures $\nu_1, \nu_2 \in \Delta$ put

$$\|\nu_1 - \nu_2\|_V = \sup_{g: |g| \le V} \Big| \sum_{n=0}^{\infty} g(n) \left(\nu_1(\{n\}) - \nu_2(\{n\}) \right) \Big|.$$

We use the notation \mathbf{E}_{ζ} when the initial distribution of X_0 is ζ , in particular, \mathbf{E}_x means that $X_0 = x$. The stationary distribution is denoted by π .

Lemma 5. Let $V(x) = x^{\alpha} + 1$, with $\alpha < \kappa$, $\alpha \le 1$. The Markov chain $(X_n)_n$ is uniformly V-geometrically ergodic, that is, there exists $\rho \in (0,1)$ and C > 0 such that for each $x \in \mathbb{N}$

$$||P^n(x,\cdot) - \pi||_V \le CV(x)\rho^n$$
,

where π denotes the unique stationary distribution of the Markov chain.

Proof. Applying Theorem 16.0.1 in [30] (equivalence of (ii) and (iv)), it is enough to prove that for some $\beta \in (0,1)$, b > 0, and a petite set C

$$\mathbf{E}_x[V(X_1)] = \mathbf{E}[V(X_1)|X_0 = x] \le \beta V(x) + b\mathbb{1}\{x \in C\}.$$

Using Jensen's inequality and that $(u+v)^{\alpha} \leq u^{\alpha} + v^{\alpha}$, u, v > 0, we have

$$\mathbf{E}\left[\left(\sum_{i=1}^{x} A_i + B\right)^{\alpha}\right] \leq \mathbf{E}\left[\left(\mathbf{E}\left[\sum_{i=1}^{x} A_i + B \mid \mathcal{E}\right]\right)^{\alpha}\right] \leq x^{\alpha}\mathbf{E}m(\xi)^{\alpha} + \mathbf{E}m^{\circ}(\xi)^{\alpha}.$$

Therefore, there exist $\beta \in (0,1)$, b > 0, and $x_0 \in \mathbb{N}$ such that for all $x \in \mathbb{N}$

$$\mathbf{E}_x[V(X_1)] = \mathbf{E}[V(X_1)|X_0 = x] \le \beta V(x) + b\mathbb{1}\{x \le x_0\}.$$

Moreover, the level set $M = \{x : x \leq x_0\}$ is small in terminology of Meyn and Tweedie [30]. Indeed, for any $x \leq x_0$

$$P(x,C) = \mathbf{P}\left(\sum_{i=1}^{x} A_i + B \in C\right) \ge \mathbf{P}\left(\sum_{i=1}^{x_0} A_i = 0; B \in C\right) =: \mu(C).$$

Since we assumed that the process is subcritical, measure μ is not trivial, M is small, and therefore petite as well. Thus the result follows.

Geometric ergodicity implies that $(X_n)_n$ is strongly mixing (Meyn and Tweedie [30] or Jones [23]), which further implies by Proposition 1.34 in [28] the mixing condition $\mathcal{A}'(a_n)$ (see Condition 2.2 in [3]). That is, there is a sequence $r_n \to \infty$, $r_n = o(n)$, such that for any $f: [0,1] \times [0,\infty) \to [0,\infty)$ for which there exists a $\delta > 0$ such that f(x,y) = 0 whenever $y \le \delta$, we have

$$\mathbf{E} \exp \left\{ -\sum_{i=1}^{n} f\left(\frac{i}{n}, \frac{X_i}{a_n}\right) \right\} - \prod_{k=1}^{k_n} \left(\mathbf{E} \exp \left\{ -\sum_{i=1}^{r_n} f\left(\frac{kr_n}{n}, \frac{X_i}{a_n}\right) \right\} \right) \to 0,$$

where $k_n = [n/r_n]$.

Roughly speaking, the last condition ensures that the sequence (X_t) can be split into blocks of consecutive observations

$$C_i = C_i(n) = (X_{(i-1)r_n+1}, \dots, X_{ir_n}), \qquad i = 1, 2, \dots, k_n,$$

which are asymptotically independent. Individual blocks could be considered as random elements of the space

$$l_0 = \{(x_i)_{i \in \mathbb{Z}} : \lim_{|i| \to \infty} |x_i| = 0\},$$

see also [6]. This embedding boils down to concatenating infinitely many zeros before and after a given block. We equip the space l_0 with the supnorm $\|(x_i)_i\| := \sup_{i \in \mathbb{Z}} |x_i|$ and with the corresponding Borel σ -field.

Consider now stationary sequence (X_t) , and recall that

$$\mathcal{L}\left(\left(\frac{X_t}{x}\right)_t \middle| X_0 > x\right) \stackrel{d}{\longrightarrow} (Y_t)_t , \qquad (30)$$

where the convergence in distribution is to be understood here with respect to the product topology (which simply corresponds to the convergence of finite-dimensional distributions). Recall further that the tail process has the form $Y_k = Y_0\Theta_k$, $k \in \mathbb{Z}$, where (Θ_k) is two sided multiplicative random walk introduced in (27) and (28). Since the walk has negative drift, $Y_t \to 0$ a.s. for $|t| \to \infty$, that is $(Y_t) \in l_0$ with probability 1. Moreover, since the anticlustering condition holds by Lemma 4, Proposition 4.2 in [7] (see also the proof of Theorem 4.3 in [7]) implies that

$$\theta = \mathbf{P}\left(\sup_{t>0} Y_t < 1\right) = \mathbf{P}\left(\sup_{t>0} Y_t \le 1\right) = \mathbf{P}\left(\max_{t>0} Y_0 m_1 \cdots m_t \le 1\right)$$
(31)

is strictly positive. By Theorem 4.3 and Remark 4.6 in [7] there exist two distributions of random elements, (Z_t) and $(Q_t)_t$ say, in l_0 such that

$$(Z_t)_t \stackrel{d}{=} (Y_t)_t \mid \sup_{t \le 0} Y_t \le 1$$
, and $(Q_t)_t \stackrel{d}{=} (Z_t)_t / \max\{Z_t : t \in \mathbb{Z}\}$. (32)

3.2 Point process convergence and partial maximum

Consider now a branching process with immigration in random environment started from an arbitrary initial distribution ζ . Recall that $P(x,\cdot)$ denotes the Markov transition kernel of the sequence (X_n) . By ζP^n we represent the distribution of the random variable X_n . In the following theorem we use the notion of convergence in distribution for point process. Following

Kallenberg [24], we endow the space of point measures on the the state space $[0,1] \times (0,\infty)$ by the vague topology. Recall, that (deterministic) measures ν_n converge vaguely to ν in such a topology if $\int f d\nu_n \to \int f d\nu$ for any continuous bounded function f with a support in some set of the form $[0,1] \times (x,\infty)$, x > 0.

Theorem 4. Let $(X_t)_{t\geq 0}$ be a branching process with immigration in random environment – b.p.i.r.e. satisfying (1) with an arbitrary initial distribution. Assume (23), then

$$N_n = \sum_{i=0}^n \delta_{(i/n, X_i/a_n)} \xrightarrow{d} N = \sum_i \sum_j \delta_{(T_i, P_i Q_{ij})}, \tag{33}$$

where

- i) $\sum_{i} \delta_{(T_i, P_i)}$ is a Poisson process on $[0, 1] \times (0, \infty)$ with intensity measure $Leb \times \nu$ where $\nu(\mathrm{d}y) = \theta \kappa y^{-\kappa 1} \mathrm{d}y$ for y > 0.
- ii) $((Q_{ij})_j)$, i = 1, 2, ..., is an i.i.d. sequence of elements in l_0 independent of $\sum_i \delta_{(T_i, P_i)}$ with common distribution equal to the distribution of (Q_i) in (32).

Remark 7. As we have seen in (31), one can characterize the key constant θ in the theorem using the tail process of (30) as

$$\theta = \mathbf{P} \Big(\sup_{t>0} Y_0 \Theta_t \le 1 \Big) = \mathbf{P} \Big(E_0 + \sup_{t>0} S_t \le 0 \Big),$$

where random variable E_0 is independent of two sided random walk $S_t = \log \Theta_t$, $t \in \mathbb{Z}$, and has exponential distribution with parameter κ .

To describe the distribution of the (Q_j) in the theorem, consider the quotient space \tilde{l}_0 of elements in l_0 which are shift-equivalent (elements $(x_i)_i, (y_i)_i \in l_0$ are shift-equivalent if for some $j \in \mathbb{Z}$, $(x_{i+j})_i = (y_i)_i$), cf. [6, Section 2]. It is shown in Basrak and Planinić [5] that in \tilde{l}_0 , (Q_t) has the same distribution as (Θ_t) under the condition that $\Theta_t < 1$ for t < 0 and $\Theta_t \le 1$ for t > 0. In other words, (Q_t) has the same distribution as $(\exp S_t)$, with random walk (S_t) conditioned on staying strictly negative for t < 0 and non positive for t > 0. We refer to Biggins [9] for more about random walks conditioned in this way.

Proof. Assume first that $(X_t)_{t\geq 0}$ is a stationary branching process with immigration in random environment – b.p.i.r.e. satisfying (24). The statement

of the theorem follows immediately from Theorem 3.1 in Basrak and Tafro [8] together with Lemmas 4 and 5 and discussion following the second lemma.

It remains to prove the convergence of point processes in (33) in the case when X_0 has an arbitrary initial distribution ζ . Observe that by the proof of Lemma 5, the function V is superharmonic for the Markov transition kernel $P(x,\cdot)$ outside of the level set M. Moreover, by the last argument in the proof of Lemma 5 each level set $\{x: x \leq r\}$ is petite. By Theorem 10.2.13 in Douc et al. [16] the Markov transition kernel P is Harris recurrent. This, together with Theorem 11.3.1 in [16] shows that for any initial distribution ζ of X_0 ,

$$\|\zeta P^m - \pi\|_{TV} \longrightarrow 0,$$

as $m \to \infty$, where ζP^m denotes the distribution of X_m .

Take an arbitrary continuous nonnegative function f with support in $[0,1] \times (\varepsilon,\infty) > 0$ for some $\varepsilon > 0$. Consider Laplace functional (see Kallenberg [24, Chapter 4]) of the point process N_n in (33) under initial distribution $X_0 \sim \zeta$

$$\mathbf{E}_{\zeta} \exp\left(-\sum_{i=1}^{n} f(i/n, X_i/a_n)\right) = \mathbf{E}_{\zeta} \exp\left(-\sum_{i=m+1}^{n} f(i/n, X_i/a_n)\right) + r_{n,m},$$

where for m fixed, $r_{n,m} \to 0$ as $n \to \infty$, because $X_i/a_n \to 0$ a.s. for $i = 1, 2, \ldots, m$. Thus

$$\begin{split} \mathbf{E}_{\zeta} \exp \left(& - \sum_{i=m+1}^{n} f(i/n, X_i/a_n) \right) \\ & = \mathbf{E}_{\zeta P^m} \exp \left(- \sum_{i=1}^{n-m} f((i+m)/n, X_i/a_n) \right) =: \int_{\mathbb{N}} H_n \, \mathrm{d}\zeta P^m \,, \end{split}$$

for a suitably chosen function H_n , which is nonnegative and bounded by 1. Similarly, for the stationary Markov chain (X_n)

$$\mathbf{E}_{\pi} \exp\left(-\sum_{i=1}^{n} f(i/n, X_i/a_n)\right) = \int_{\mathbb{N}} H_n \, \mathrm{d}\pi + r'_{n,m}$$
$$\longrightarrow \mathbf{E} \exp\left(-\sum_{i} \sum_{j} f(T_i, P_i Q_{ij})\right).$$

Observing that $\left| \int_{\mathbb{N}} H_n \, \mathrm{d}\zeta P^m - \int_{\mathbb{N}} H_n \, \mathrm{d}\pi \right| \leq \|\zeta P^m - \pi\|_{TV}$, uniformly over $n \in \mathbb{N}$, and $r'_{n,m} \to 0$ for fixed m as $n \to \infty$, yields the statement. \square

We consider next partial maxima of the sequence $(X_t)_{t\in\mathbb{N}}$, namely we define $M_n = \max\{X_1, \ldots, X_n\}$, for any $n \geq 1$. Observe that event $\{M_n/a_n \leq x\}$ corresponds to the event $\{N_n([0,1]\times(x,\infty))=0\}$. Moreover, for any x>0, the limit point process $N=\sum_i\sum_j\delta_{T_i,P_iQ_{ij}}$ has probability 0 of hitting the boundary of the set $[0,1]\times(x,\infty)$, thus $\mathbf{P}(N_n([0,1]\times(x,\infty))=0)\to \mathbf{P}(N([0,1]\times(x,\infty))=0)$. However, by (32), (Q_{ij}) in (33) satisfy $Q_{ij}\leq 1$ with at least one point exactly equal to 1. Thus $\mathbf{P}(N([0,1]\times(x,\infty))=0)=\mathbf{P}(\sum_i\delta_{T_i,P_i}([0,1]\times(x,\infty))=0)$. Therefore, for any initial distribution of X_0 , partial maxima converge to a rescaled Fréchet distribution.

Corollary 1. Let $(X_t)_{t\geq 0}$ be a branching process with immigration in random environment – b.p.i.r.e. satisfying (1) with an arbitrary initial distribution. Suppose that (23) holds. Then for any $x \geq 0$

$$\mathbf{P}\left(\frac{M_n}{a_n} \le x\right) \to e^{-\theta x^{-\kappa}} \quad as \quad n \to \infty.$$

4 Partial sums

Denote by (X_t) a stationary branching process with immigration in a random environment and assume that at least one of the conditions in (23) holds. To derive limit theorem for the partial sums from the point process convergence is immediate for $\alpha \in (0,1)$, but needs an extra condition for $\alpha \in [1,2)$.

Lemma 6 (Vanishing small values). Assume that $\kappa \in [1, 2)$. Then for any $\varepsilon > 0$

$$\lim_{\gamma \downarrow 0} \limsup_{n \to \infty} \mathbf{P} \left(\left| \sum_{i=1}^{n} \left[X_i \mathbb{1} \{ X_i \le a_n \gamma \} - \mathbf{E} [X_0 \mathbb{1} \{ X_0 \le a_n \gamma \}] \right] \right| > a_n \varepsilon \right) = 0.$$

Proof. Choose $\alpha < 1$ such that $1 + \alpha > \kappa$. (For $\kappa > 1$ we may choose $\alpha = 1$.) Lemma 5 holds with $V(x) = 1 + x^{\alpha}$. Let $\varepsilon = 1 - \alpha$. Then,

$$h(x) := \frac{x}{(a_n \gamma)^{\varepsilon}} \mathbb{1}\{x \le a_n \gamma\} \le x^{\alpha}, \text{ for } x \in \mathbb{N}.$$

Thus $|h| \leq V$, therefore by Lemma 5, for some $\rho \in (0,1)$ and C > 0 we have

$$\left| \mathbf{E} \left[\frac{X_i}{(a_n \gamma)^{\varepsilon}} \mathbb{1} \{ X_i \le a_n \gamma \} \, \middle| X_0 = m \right] - \mathbf{E} \frac{X_0}{(a_n \gamma)^{\varepsilon}} \mathbb{1} \{ X_0 \le a_n \gamma \} \right| \le C(1 + m^{\alpha}) \rho^i.$$

Thus

$$\mathbf{E}\left[X_{i}\mathbb{1}\left\{X_{i} \leq a_{n}\gamma\right\}X_{0}\mathbb{1}\left\{X_{0} \leq a_{n}\gamma\right\}\right]$$

$$= \sum_{m \leq a_{n}\gamma} m\mathbf{P}(X_{0} = m)(a_{n}\gamma)^{\varepsilon}\mathbf{E}\left[\frac{X_{i}}{(a_{n}\gamma)^{\varepsilon}}\mathbb{1}\left\{X_{i} \leq a_{n}\gamma\right\} \mid X_{0} = m\right]$$

$$\leq \sum_{m \leq a_{n}\gamma} m\mathbf{P}(X_{0} = m)(a_{n}\gamma)^{\varepsilon}\left(\mathbf{E}\frac{X_{0}}{(a_{n}\gamma)^{\varepsilon}}\mathbb{1}\left\{X_{0} \leq a_{n}\gamma\right\} + C(1 + m^{\alpha})\rho^{i}\right)$$

$$\leq \left[\mathbf{E}X_{0}\mathbb{1}\left\{X_{0} \leq a_{n}\gamma\right\}\right]^{2} + 2C\rho^{i}(a_{n}\gamma)^{\varepsilon}\mathbf{E}X_{0}^{2-\varepsilon}\mathbb{1}\left\{X_{0} \leq a_{n}\gamma\right\}.$$
(34)

For $\beta > \kappa$ by the regular variation and Karamata's theorem as $u \to \infty$

$$\mathbf{E}X_0^{\beta} \mathbb{1}\{X_0 \le u\} \sim \frac{\beta}{\beta - \kappa} u^{\beta} \overline{F}(u). \tag{35}$$

Substituting back into (34), and using that $\overline{F}(a_n\gamma) \sim \gamma^{-\kappa} n^{-1}$ we have

$$\operatorname{Cov}\left(X_0 \mathbb{1}\left\{X_0 \le a_n \gamma\right\}, X_i \mathbb{1}\left\{X_i \le a_n \gamma\right\}\right) \le C \rho^i \frac{a_n^2}{n} \gamma^{2-\kappa}.$$

Therefore, using the stationarity, (34), and (35), we obtain

$$\mathbf{E} \left(\sum_{i=1}^{n} [X_{i} \mathbb{1}\{X_{i} \leq a_{n}\gamma\} - \mathbf{E}[X_{0} \mathbb{1}\{X_{0} \leq a_{n}\gamma\}]] \right)^{2}$$

$$= n \operatorname{Var} (X_{0} \mathbb{1}\{X_{0} \leq a_{n}\gamma\})$$

$$+ \sum_{i=1}^{n-1} 2(n-i) \operatorname{Cov} (X_{0} \mathbb{1}\{X_{0} \leq a_{n}\gamma\}, X_{i} \mathbb{1}\{X_{i} \leq a_{n}\gamma\})$$

$$\leq n \operatorname{Var} (X_{0} \mathbb{1}\{X_{0} \leq a_{n}\gamma\}) + C \sum_{i=1}^{n-1} (n-i) \rho^{i} \frac{a_{n}^{2}}{n} \gamma^{2-\kappa}$$

$$\leq C a_{n}^{2} \gamma^{2-\kappa}.$$

Therefore, the claim follows from Chebyshev's inequality.

Remark 8. In fact, we proved condition (9) in Davis [14].

Theorem 5. Let $(X_t)_{t\geq 0}$ be a branching process with immigration in random environment – b.p.i.r.e. satisfying (1) with an arbitrary initial distribution. Assume (23) and denote by (b_n) a sequence of real numbers given by

$$b_n = 0, \quad \kappa < 1, \quad and \quad b_n = n \mathbf{E} \left(\frac{X_{\infty}}{a_n} \mathbb{1} \left\{ \frac{X_{\infty}}{a_n} \le 1 \right\} \right), \quad for \ \kappa \in [1, 2).$$

Then

$$V_n = \sum_{k=1}^n \frac{X_k}{a_n} - b_n \xrightarrow{d} V, \qquad n \to \infty, \tag{36}$$

where V is a κ -stable random variable. For $\kappa > 2$, as $n \to \infty$

$$\frac{1}{\sqrt{n}\sigma}\sum_{i=1}^{n}(X_i - \mathbf{E}X_{\infty}) \stackrel{d}{\longrightarrow} Z,$$

where Z is a standard normal random variable, and $\sigma^2 = \frac{1 + \mathbf{E}A}{1 - \mathbf{E}A} \operatorname{Var}(X_{\infty}) > 0$.

Remark 9. Recall that under the conditions of Theorem 1 for the normalizing sequence we may choose $a_n = (Cn)^{1/\kappa}$, in general we have $a_n = \tilde{\ell}(n)n^{1/\kappa}$, for some slowly varying $\tilde{\ell}$, as observed after (25). Concerning the centering constants (b_n) , recall that the the law of X_{∞} represents the unique stationary distribution for b.p.i.r.e. satisfying (1). For $\kappa \in (1,2)$, the mean of X_{∞} is finite, so one could substitute centering constants (b_n) by $n\mathbf{E}X_{\infty}/a_n$ to show

$$\frac{1}{a_n} \sum_{i=1}^n (X_i - \mathbf{E} X_\infty) \xrightarrow{d} V - \frac{\kappa}{\kappa - 1}.$$
 (37)

Thus, we again get a κ -stable limit with different location parameter, cf. Remark 3.1 in [15]. Moreover, under conditions of Theorem 1, in the case $\kappa = 1$, $b_n \sim C^{-1} \log n$.

Remark 10. One can also describe the limiting stable distribution in (36) in terms of parameters κ , θ and the distribution of (Q_j) , see Remark 3.2 in [15] for details. In the case $\kappa < 1$ for instance, the limiting random variable V has a characteristic function of relatively simple form

$$\mathbf{E}e^{itV} = \exp\left(-d|t|^{\kappa} \left(1 - i\operatorname{sgn}(t)\tan\frac{\pi\kappa}{2}\right)\right), \qquad (38)$$

where $d = \theta \Gamma(1-\kappa) \mathbf{E}(\sum_j Q_j)^{\kappa} \cos(\pi \kappa/2)$, which is known to be finite since for $\kappa \leq 1$, $\mathbf{E}(\sum_j Q_j)^{\kappa} \leq \mathbf{E}(\sum_j Q_j^{\kappa}) < \infty$, see [15, Remark 3.2, Theorem 3.2]. For $\kappa \in (1,2)$, multiplicative random walk (Θ_j) given by (27) and (28) satisfies $\mathbf{E}(\sum_j \Theta_j)^{\kappa-1} \leq \mathbf{E}(\sum_j \Theta_j^{\kappa-1}) < \infty$. By Theorem 5.5.1 in Kulik and Soulier [29], this yields $\mathbf{E}(\sum_j Q_j)^{\kappa} < \infty$. Applying Proposition 8.3.2 and Theorem 8.4.3 in [29], and observing that they use $n\mathbf{E}X_{\infty}/a_n$ as centering constants, one can deduce that

$$\mathbf{E}e^{itV} = \exp\left(-d|t|^{\kappa} \left(1 - i\operatorname{sgn}(t)\tan\frac{\pi\kappa}{2}\right) + ict\right),\,$$

with the same expression for the scale parameter d and with $c = \kappa/(\kappa - 1)$, cf. (37).

For $\kappa = 1$, (Θ_j) satisfies $\mathbf{E}[\log(\sum_j \Theta_j)] \leq \mathbf{E}(\sum_j \Theta_j)^{1/2} < \infty$. By Chapter 5 in Kulik and Soulier [29] (see Problem 5.31 therein), this implies $\mathbf{E}(\sum_j Q_j \log Q_j^{-1}) < \infty$. This time, Theorem 8.4.3 in [29] yields

$$\mathbf{E}e^{itV} = \exp\left(-d|t|\left(1 + i\frac{2}{\pi}\operatorname{sgn}(t)\log|t|\right) + ict\right),\,$$

where $d = \theta \frac{\pi}{2} \mathbf{E}(\sum_{j} Q_{j})$ while an expression for the location parameter c can be found in Proposition 8.3.3 of [29].

Note finally, that for $\kappa > 2$ one can easily check that the conditions of Theorem 19.1 in Billingsley [10] hold, therefore the functional version of the CLT also holds.

Proof. Assume first that $\kappa < 2$. As in the proof of Theorem 4, we first assume that (X_t) is a stationary b.p.i.r.e. process satisfying the conditions of the theorem. The claim then follows directly from Theorem 4 and Theorem 3.1 in Davis and Hsing [15]. Observe that the condition (3.2) therein follows from Lemma 6.

To prove the theorem when X_0 has an arbitrary initial distribution we can use a similar argument as in the proof of Theorem 4. Observe first, that

$$V_n = \sum_{k=1}^n \frac{X_k}{a_n} - b_n = \sum_{k=1}^{n-m} \frac{X_{k+m}}{a_n} - b_n + r_{n,m}$$

with $r_{n,m} \to 0$ in probability as $n \to \infty$ for any fixed m. Denote by ζP^m the distribution of X_m , and note that for any $s \in \mathbb{R}$

$$\mathbf{E}_{\zeta} \left[\exp \left(is \left(\sum_{k=1}^{n-m} \frac{X_{k+m}}{a_n} - b_n \right) \right) \right] = \mathbf{E}_{\zeta P^m} \left[\exp \left(is \left(\sum_{k=1}^{n-m} \frac{X_k}{a_n} - b_n \right) \right) \right]$$

$$= \mathbf{E}_{\pi} \left[\exp \left(is \left(\sum_{k=1}^{n-m} \frac{X_k}{a_n} - b_n \right) \right) \right] + u_{n,m},$$

where for all n

$$u_{n,m} \le \|\zeta P^m - \pi\|_{TV} \,,$$

with the right hand side tending to 0 as $m \to \infty$. Observe now that a stationary b.p.i.r.e. (X_t) satisfies $r'_{n,m} = \sum_{k=n-m+1}^n \frac{X_k}{a_n} \to 0$ in probability as $n \to \infty$. Therefore, it also satisfies

$$\sum_{k=1}^{n-m} \frac{X_k}{a_n} - b_n = \sum_{k=1}^{n} \frac{X_k}{a_n} - b_n - r'_{n,m} \stackrel{d}{\longrightarrow} V.$$

Let now $\kappa > 2$. Then $\mathrm{Var}(X_\infty) < \infty$, and the result follows from the standard Markov chain theory. To apply Theorem 1 (i) in Jones [23] we have to prove the drift condition as in Lemma 5 with $V(x) = 1 + x^2$. Simple calculation gives

$$\mathbf{E}\left(\sum_{i=1}^{x} A_i + B\right)^2 = x^2 \mathbf{E} m(\xi)^2 + x \left(\mathbf{E} A^2 - \mathbf{E} m(\xi)^2 + 2\mathbf{E} A B\right) + \mathbf{E} B^2.$$

Since $\mathbf{E}m(\xi)^2 < 1$, there exist $\beta \in (0,1)$, b > 0, and $x_0 \in \mathbb{N}$ such that for all $x \in \mathbb{N}$

$$\mathbf{E}_x[V(X_1)] = \mathbf{E}[V(X_1)|X_0 = x] \le \beta V(x) + b\mathbb{1}\{x \le x_0\}.$$

Moreover, the level set $M = \{x : x \leq x_0\}$ is small, so the conditions in Theorem 1 (i) in Jones [23] hold.

We only have to check that $\sigma^2 = \operatorname{Var}_{\pi}(X_0) + 2 \sum_{t=1}^{\infty} \operatorname{Cov}_{\pi}(X_0, X_t) > 0$. Using decomposition (29) from the proof of Lemma 4 we see that

$$\operatorname{Cov}_{\pi}(X_0, X_t) = \operatorname{Cov}_{\pi}\left(X_0, \sum_{i=1}^{X_0} \widetilde{A}_i^{(t)}\right),\,$$

where $\widetilde{A}_i^{(t)}$ stands for the number of descendants in generation t of a single individual in generation 0. Thus

$$\mathbf{E}\left(X_0 \sum_{i=1}^{X_0} \widetilde{A}_i^{(t)}\right) = (\mathbf{E}A)^t \mathbf{E} X_0^2.$$

Therefore

$$\operatorname{Cov}_{\pi}(X_0, X_t) = \operatorname{Var}_{\pi}(X_0) (\mathbf{E}A)^t,$$

and the formula $\sigma^2 = \operatorname{Var}_{\pi}(X_0)(1 + \mathbf{E}A)/(1 - \mathbf{E}A)$ follows.

To show that $\sigma > 0$ we have to prove that the stationary solution cannot be deterministic. Indeed, the fixed point equation (6) (or (9)) has no deterministic solution, since Cramér's condition (3) implies $\mathbf{P}(A > 1) > 0$, thus $\mathbf{P}(\sum_{i=1}^{m} A_i + B > m) > 0$ for any positive integer m.

5 Random walks in a random environment

Connection between random walks in a random environment and branching processes with immigration was made already in the seminal paper by

Kesten et al. [25]. To describe the setup of their model, we can use again a sequence of i.i.d. random variables ξ , $(\xi_t)_{t\in\mathbb{Z}}$ with values in the interval (0,1). Consider the process $W_0 = 0$ and

$$W_t = W_{t-1} + \eta_{W_{t-1},t} \tag{39}$$

where conditioned on the environment \mathcal{E} , $\eta_{x,t}$'s are independent random variables taking value +1 with probability ξ_x and value -1 with probability $\xi_x' = 1 - \xi_x$. The process (W_t) is called random walk in random environment – r.w.r.e.

Following [25] we assume throughout that

$$\mathbf{E}\log\frac{\xi'}{\xi} < 0\tag{40}$$

and

$$\mathbf{E} \left(\frac{\xi'}{\xi} \right)^{\kappa} = 1 \quad \text{and} \quad \mathbf{E} \left(\frac{\xi'}{\xi} \right)^{\kappa} \log \frac{\xi'}{\xi} < \infty, \tag{41}$$

for some $\kappa > 0$. The first of these conditions ensures that (W_t) drifts to $+\infty$ and the second one corresponds to (3). As observed in [25], for the asymptotic analysis of (W_t) , it is crucial to understand asymptotic behavior of the random variables

$$T_n = \min\{t : W_t = n\}, \quad n > 1,$$

which are all finite a.s. Observe that on the way to the state n, process (W_t) visits each state $k = 0, 1, \ldots, n-1$ at least once. For $i = 1, \ldots, n$ denote

$$L_i^n = 1 + U_i^n = 1 + \#\{\text{visits to the state } n - i \text{ from the right before } T_n\}.$$

Note, until time T_n each of the visits to n-i from the right is canceled by one movement back to the right. Note U_i^n are well defined and possibly different from 0 also for i>n. Observe next that the total number of visits by the process (W_t) to the left of 0, say $2U_{\infty}$, is a.s. finite. Thus, $R_n=2\sum_{i>n}U_i^n\leq 2U_{\infty}$. Thus in

$$T_n = n + 2\sum_{i \ge 1} U_i^n = n + 2\sum_{i=1}^n U_i^n + R_n,$$

the term R_n remains bounded as $n \to \infty$.

Observe next that $U_1^n=0$, so that $L_1^n=1$ and the sequence L_i^n evolves as follows

$$L_2^n = G_{n,1,1} + 1,$$

where $G_{n,1,1}$ represents the number of right visits to n-2 from n-1 before eventual move to the right. Conditioned on \mathcal{E} , $G_{n,1,1}$ has a geometric distribution with mean ξ'_{n-1}/ξ_{n-1} . Similarly, for each $i \geq 2$

$$L_i^n = \sum_{j=1}^{L_{i-1}^n} G_{n,i-1,j} + 1, \qquad (42)$$

where conditionally on \mathcal{E} , $G_{n,i-1,j}$, $j=1,2,\ldots$, are i.i.d. with a geometric distribution with mean ξ'_{n-i+1}/ξ_{n-i+1} .

The finite sequences

$$L_1^n, L_2^n, \ldots, L_k^n, \ldots, L_n^n$$

represent a special case of branching process in random environment. Moreover, the initial part $L_1^n, L_2^n, \ldots, L_k^n$ has the same distribution for each $n \geq k$, hence if we want to understand the limiting distribution of $T_n = 2\sum_{i=1}^n L_i^n$ $n+R_n$, we can simply skip the index n in (42) and analyze $\sum_{i=1}^n L_i$, which is exactly the content of Theorem 5. Denote in the sequel by L_∞ the random variable which has the stationary distribution of the Markov chain in (42). The following theorem corresponds to the part of the main theorem in [25] concerning the case $\kappa < 2$. The corresponding statement for $\kappa > 2$ follows from Theorem 5 as well, in that case T_n after centering and normalization with \sqrt{n} converges to a normal distribution. Observe that in the case $\kappa = 2$, T_n converges to a normal limit again, but additional care has to be taken about normalizing and centering constants, see [25].

Theorem 6. Let $(W_t)_{t\geq 0}$ be a random walk in random environment satisfying (39). Suppose that the conditions of (40) and (41) hold for some $\kappa > 0$. Suppose further that the law of $\log(\xi_i'/\xi_i)$ is nonarithmetic.

(i) For $\kappa \in (0,1)$ as $n \to \infty$

$$\frac{1}{n^{1/\kappa}}T_n \stackrel{d}{\longrightarrow} \widetilde{V},$$

where \widetilde{V} has a strictly positive κ -stable distribution, while, as $t \to \infty$

$$\frac{1}{t^{\kappa}}W_t \stackrel{d}{\longrightarrow} \widetilde{V}^{-\kappa}.$$

(ii) For
$$\kappa = 1$$

$$\frac{1}{n}T_n - 2Cb_n \stackrel{d}{\longrightarrow} \widetilde{V},$$

where $b_n \sim \log n$ and \widetilde{V} has a 1-stable distribution. Moreover, as $t \to \infty$

$$\frac{(\log t)^2}{t}(W_t - \delta(t)) \stackrel{d}{\longrightarrow} \frac{-\widetilde{V}}{(2C)^2},$$

with $\delta(t) \sim t/(2C \log t)$.

iii) For
$$\kappa \in (1,2)$$

$$\frac{1}{n^{1/\kappa}} (T_n - n(2\mathbf{E}L_{\infty} - 1)) \stackrel{d}{\longrightarrow} \widetilde{V}.$$

Moreover, as $t \to \infty$

$$\frac{1}{t^{1/\kappa}} \left(W_t - \frac{t}{2\mathbf{E}L_{\infty} - 1} \right) \xrightarrow{d} \frac{-\widetilde{V}}{(2\mathbf{E}L_{\infty} - 1)^{1 + 1/\kappa}}.$$

Proof. Recall that $T_n \stackrel{d}{=} \sum_{i=1}^n (2L_i - 1) + R_n$, with random variables R_n a.s. bounded and where (L_n) represents a special b.p.i.r.e. which is initialized at one, with progeny conditionally geometric given the environmental variables ξ_j 's and with the immigration identically equal to 1. Recall further that the conditions of Theorems 1 and 5 are met by assumptions (40), (41) and Remark 1.

Define now

$$c_n = \begin{cases} 0 & \text{for } \kappa \in (0,1), \\ 2\mathbf{E}L_{\infty} \mathbb{1}\left\{L_{\infty} \le (Cn)^{1/\kappa}\right\} - 1 & \text{for } \kappa = 1, \\ 2\mathbf{E}L_{\infty} - 1 & \text{for } \kappa \in (1,2). \end{cases}$$

Theorem 5 and Remark 9 imply that $2^{-1}(Cn)^{-1/\kappa} \sum_{i=1}^{n} (2L_i - 1 - c_n)$ converges in distribution to a κ -stable random variable for all $\kappa \in (0,2)$. After multiplication by $2C^{1/\kappa}$, this yields

$$\frac{1}{n^{1/\kappa}}(T_n - nc_n) \stackrel{d}{\longrightarrow} \widetilde{V} ,$$

for a certain κ -stable random variable \widetilde{V} . Using the particular form of constants c_n gives the centering sequence in each of the three cases. The convergence in distribution of the r.w.r.e. W_t now follows as in Kesten et al. [25].

Remark 11. Observe that the characteristic function, and the exact distribution therefore, of the limit \tilde{V} in all three statements of the theorem can now be read out from Remark 10. In particular, location and scale parameters of the stable law \tilde{V} are now determined in terms of the values κ , θ and the conditional multiplicative random walk (Q_j) .

The analogous results hold under the second condition in (23). We translate this to the RWRE setup in the following theorem. Recall first that normalizing sequence (a_n) from (25) satisfies $a_n = n^{1/\kappa} \tilde{\ell}(n)$ for some slowly varying function $\tilde{\ell}$, and extend this to a function $a(t) = t^{1/\kappa} \tilde{\ell}(t)$ on $(0, \infty)$.

Theorem 7. Let $(W_t)_{t\geq 0}$ be a random walk in random environment satisfying either condition (i) or (ii) in Theorem 2 for $\kappa \in (1,2)$ with $m(\xi) = \xi'/\xi$. Suppose further that the law of $\log(\xi'_i/\xi_i)$ is nonarithmetic. Then

$$\frac{1}{a_n}(T_n - n(2\mathbf{E}L_{\infty} - 1)) \stackrel{d}{\longrightarrow} \widetilde{V} \quad as \ n \to \infty.$$

Moreover, as $t \to \infty$

$$\frac{1}{a(t)} \left(W_t - \frac{t}{2\mathbf{E}L_{\infty} - 1} \right) \xrightarrow{d} \frac{-\widetilde{V}}{(2\mathbf{E}L_{\infty} - 1)^{1 + 1/\kappa}}.$$

Proof. The first part follows again from Theorem 5, while the second part follows from the inverse relation between T_n and W_t ; see (2.38) in [25]. We omit the details.

Example. Assume that the random variable ξ satisfies $\mathbf{E}(\xi'/\xi)^{\kappa} = 1$, and has a density function f such that for all u > 0 small enough

$$f(u) = u^{\kappa - 1} \alpha \frac{\log \log u^{-1}}{(\log u^{-1})^{\alpha + 1}},$$

where $\kappa \geq 1$, $\alpha \in (1/2,1)$. Then straightforward calculation shows that

$$\mathbf{E}\mathbb{1}\left\{\log\frac{1-\xi}{\xi} > x\right\} \left(\frac{1-\xi}{\xi}\right)^{\kappa} = \int_0^{(1+e^x)^{-1}} \left(\frac{1-u}{u}\right)^{\kappa} f(u) du \sim \frac{\log x}{x^{\alpha}}.$$

Thus, the conditions of the theorem above holds, and for M in (18)

$$M(x) \sim \frac{x^{1-\alpha} \log x}{1-\alpha}.$$

Therefore by Theorem 2, the stationary distribution L_{∞} satisfies

$$\mathbf{P}(L_{\infty} > x) \sim \frac{C(1-\alpha)}{x^{\kappa}(\log x)^{1-\alpha}\log\log x}.$$

Then

$$a_n \sim [\kappa^{1-\alpha}C(1-\alpha)]^{1/\kappa} \frac{n^{1/\kappa}}{(\log n)^{(1-\alpha)/\kappa}(\log\log n)^{1/\kappa}}.$$

According to Kesten et al. [25] even before they gave the proof of the theorem, it was conjectured by A.N. Kolmogorov and F. Spitzer that T_n might exhibit the behavior described above. The intuitive reason behind this observation may be the existence of so-called traps between 0 and n, i.e. sites $j \in \{0, ..., n\}$ where corresponding ξ_j is atypically small, which makes it very difficult for the random walk (W_t) to cross over to the right. It is interesting that our other main result, Theorem 4, provides a very simple argument characterizing the asymptotic distribution of the worst of such traps. Denote now for k < n by

$$V_k^n = \#\{\text{crossings over the edge } (k, k+1) \text{ before } T_n\}.$$

Clearly, for k = 0, ..., n-1, we have $V_k = 1 + 2U_{n-k}^n$ and while for k < 0 $V_k = 2U_{n-k}^n$. Observing that $\max_{k < 0} V_k^n$ remains bounded a.s. again, from Corollary 1 we can deduce the following result concerning the most visited edge until time T_n .

Corollary 2. Under the assumptions of Theorem 6 or Theorem 7

$$\mathbf{P}\left(\frac{\max_{k < n} V_k^n}{2a_n} \le x\right) \to e^{-\theta x^{-\kappa}},\,$$

as $n \to \infty$.

Recall that by Remark 7, θ in either case can be obtained as

$$\theta = \mathbf{P}\left(E_0 + \sup_{t>0} \sum_{i=1}^t \log(\xi_i'/\xi_i) \le 0\right),\,$$

where E_0 stands for an exponential random variable with parameter κ independent of the environment sequence (ξ_i) .

Acknowledgement. We are grateful to Dariusz Buraczewski for discussion on the problem and pointing out reference [1]. This work is in part financed within the Croatian-Swiss Research Program of the Croatian Science Foundation and the Swiss National Science Foundation – grant CSRP 018-01-180549. Péter Kevei is supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, by the NKFIH grant FK124141, and by the EU-funded Hungarian grant EFOP-3.6.1-16-2016-00008.

References

- [1] V. I. Afanasyev. On the maximum of a subcritical branching process in a random environment. *Stochastic Process. Appl.*, 93(1):87–107, 2001.
- [2] M. Barczy, F. K. Nedényi, and G. Pap. On aggregation of multitype Galton-Watson branching processes with immigration. *Mod. Stoch. Theory Appl.*, 5(1):53–79, 2018.
- [3] B. Basrak, D. Krizmanić, and J. Segers. A functional limit theorem for dependent sequences with infinite variance stable limits. *Ann. Probab.*, 40(5):2008–2033, 2012.
- [4] B. Basrak, R. Kulik, and Z. Palmowski. Heavy-tailed branching process with immigration. *Stoch. Models*, 29(4):413–434, 2013.
- [5] B. Basrak and H. Planinić. Compound Poisson approximation for random fields with application to sequence alignment, 2019.
- [6] B. Basrak, H. Planinić, and P. Soulier. An invariance principle for sums and record times of regularly varying stationary sequences. *Probab.* Theory Related Fields, 172(3-4):869–914, 2018.
- [7] B. Basrak and J. Segers. Regularly varying multivariate time series. Stochastic Process. Appl., 119(4):1055–1080, 2009.
- [8] B. Basrak and A. Tafro. A complete convergence theorem for stationary regularly varying multivariate time series. *Extremes*, 19(3):549–560, 2016.
- [9] J. D. Biggins. Random walk conditioned to stay positive. *J. London Math. Soc.* (2), 67(1):259–272, 2003.
- [10] P. Billingsley. Convergence of probability measures. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999.
- [11] D. Buraczewski, E. Damek, and T. Mikosch. Stochastic Models with Power-Law Tails. The Equation X=AX+B. Springer Series in Operations Research and Financial Engineering. Springer, 2016.
- [12] D. Buraczewski and P. Dyszewski. Precise large deviation estimates for branching process in random environment. https://arxiv.org/abs/1706.03874v1, 2019.

- [13] F. Caravenna and R. Doney. Local large deviations and the strong renewal theorem. *Electron. J. Probab.*, 24:Paper No. 72, 48, 2019.
- [14] R. A. Davis. Stable limits for partial sums of dependent random variables. *Ann. Probab.*, 11(2):262–269, 1983.
- [15] R. A. Davis and T. Hsing. Point process and partial sum convergence for weakly dependent random variables with infinite variance. Ann. Probab., 23(2):879–917, 1995.
- [16] R. Douc, E. Moulines, P. Priouret, and P. Soulier. *Markov chains*. Springer Series in Operations Research and Financial Engineering. Springer, Cham, 2018.
- [17] S. Foss, D. Korshunov, and S. Zachary. An introduction to heavy-tailed and subexponential distributions. Springer Series in Operations Research and Financial Engineering. Springer, New York, second edition, 2013.
- [18] A. Garsia and J. Lamperti. A discrete renewal theorem with infinite mean. *Comment. Math. Helv.*, 37:221–234, 1962/63.
- [19] C. M. Goldie. Implicit renewal theory and tails of solutions of random equations. *Ann. Appl. Probab.*, 1(1):126–166, 1991.
- [20] A. Gut. Stopped random walks. Springer Series in Operations Research and Financial Engineering. Springer, New York, second edition, 2009. Limit theorems and applications.
- [21] A. Janssen and J. Segers. Markov tail chains. *J. Appl. Probab.*, 51(4):1133–1153, 2014.
- [22] P. R. Jelenković and M. Olvera-Cravioto. Implicit renewal theorem for trees with general weights. Stochastic Process. Appl., 122(9):3209–3238, 2012.
- [23] G. L. Jones. On the Markov chain central limit theorem. *Probab. Surv.*, 1:299–320, 2004.
- [24] O. Kallenberg. Random measures, theory and applications, volume 77 of Probability Theory and Stochastic Modelling. Springer, Cham, 2017.
- [25] H. Kesten, M. V. Kozlov, and F. Spitzer. A limit law for random walk in a random environment. *Compositio Math.*, 30:145–168, 1975.

- [26] P. Kevei. A note on the Kesten–Grincevičius–Goldie theorem. *Electron. Commun. Probab.*, 21(51):1–12, 2016.
- [27] P. Kevei. Implicit renewal theory in the arithmetic case. *J. Appl. Probab.*, 54(3):732–749, 2017.
- [28] D. Krizmanić. Functional limit theorems for weakly dependent regularly varying time series. PhD thesis, University of Zagreb, 2010.
- [29] R. Kulik and P. Soulier. Heavy tailed time series. Manuscript, 2019.
- [30] S. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Cambridge University Press, Cambridge, second edition, 2009.
- [31] V. V. Petrov. Limit theorems of probability theory, volume 4 of Oxford Studies in Probability. The Clarendon Press, Oxford University Press, New York, 1995.
- [32] M. P. Quine. The multi-type Galton-Watson process with immigration. J. Appl. Probability, 7:411–422, 1970.
- [33] S. I. Resnick. Heavy-tail phenomena: probabilistic and statistical modeling. Springer Verlag, 2007.
- [34] A. Roitershtein. A note on multitype branching processes with immigration in a random environment. *Ann. Probab.*, 35(4):1573–1592, 2007.
- [35] J. Segers. Multivariate regular variation of heavy-tailed Markov chains. Available on arXiv: https://arxiv.org/abs/math/0701411, 2007.
- [36] G. Szűcs. Ergodic properties of subcritical multitype Galton-Watson processes. Available on arXiv: https://arxiv.org/abs/1402.5539.