

WELL-POSEDNESS AND QUALITATIVE BEHAVIOUR OF A TWO-PHASE NAVIER-STOKES/MULLINS-SEKERKA SYSTEM WITH BOUNDARY CONTACT

MAXIMILIAN RAUCHECKER AND MATHIAS WILKE

ABSTRACT. We consider a coupled two-phase Navier-Stokes/Mullins-Sekerka system describing the motion of two immiscible, incompressible fluids inside a bounded container. The moving interface separating the liquids meets the boundary of the container at a constant ninety degree angle. This common interface is unknown and has to be determined as a part of the problem.

We show well-posedness and investigate the long-time behaviour of solutions starting close to certain equilibria. We prove that for equal densities these solutions exist globally in time, are stable, and converge to an equilibrium solution at an exponential rate.

1. INTRODUCTION

In this article we study the two-phase Navier-Stokes equations with surface tension coupled to the Mullins-Sekerka problem inside a bounded domain in two or three space dimensions. In our model, the interface separating the two fluids meets the boundary of the domain at a constant ninety degree angle. This leads to a free boundary problem for the interface involving a contact angle problem at the boundary as well.

We assume that the domain $\Omega \subset \mathbb{R}^n$, $n = 2, 3$, can be decomposed as $\Omega = \Omega^+(t) \dot{\cup} \dot{\Gamma}(t) \dot{\cup} \Omega^-(t)$, where $\dot{\Gamma}(t)$ denotes the interior of $\Gamma(t)$, an $(n-1)$ -dimensional submanifold with boundary. We interpret $\Gamma(t)$ to be the interface separating the two phases, $\Omega^+(t)$ and $\Omega^-(t)$, which will both be assumed to be connected. The boundary of $\Gamma(t)$ will be denoted by $\partial\Gamma(t)$. Furthermore we assume $\Gamma(t)$ to be orientable, the unit normal vector field on $\Gamma(t)$ pointing from $\Omega^-(t)$ into $\Omega^+(t)$ will be denoted by $\nu_{\Gamma(t)}$.

Let us introduce some notation. Let $V_{\Gamma(t)}$ denote the normal velocity and $H_{\Gamma(t)}$ the mean curvature of the free interface $\Gamma(t)$. By $[[\cdot]]$ we denote the jump of a quantity across $\Gamma(t)$ in direction of $\nu_{\Gamma(t)}$, that is,

$$[[f]](x) := \lim_{\varepsilon \rightarrow 0+} [f(x + \varepsilon \nu_{\Gamma(t)}) - f(x - \varepsilon \nu_{\Gamma(t)})], \quad x \in \Gamma(t).$$

Furthermore, $a \otimes b$ is defined by $[a \otimes b]_{ij} := a_i b_j$ for vectors $a, b \in \mathbb{R}^n$ and A^\top denotes the transposed matrix of A .

MAXIMILIAN RAUCHECKER, FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, 93053 REGENSBURG, GERMANY

MATHIAS WILKE, INSTITUT FÜR MATHEMATIK, MARTIN-LUTHER-UNIVERSITÄT HALLE-WITTENBERG, 06099 HALLE, GERMANY

We assume that Ω is filled by two immiscible, incompressible fluids with respective constant densities $\rho^\pm > 0$ in the two phases. Their respective constant viscosities are denoted by $\mu^\pm > 0$ and $\sigma > 0$ is a given surface tension constant. To economize our notation, we let $\rho := \rho^+ \chi_{\Omega^+(t)} + \rho^- \chi_{\Omega^-(t)}$ and $\mu := \mu^+ \chi_{\Omega^+(t)} + \mu^- \chi_{\Omega^-(t)}$, where χ_M is the indicator function of a set M . In our model, u is the velocity of the fluids, p the pressure, η the chemical potential and $\Gamma(t)$ the free interface at time $t \geq 0$.

Let us consider the case where the domain is a cylindrical container $\Omega = \Sigma \times (L_1, L_2)$, where $-\infty < L_1 < 0 < L_2 < \infty$ and $\Sigma \subset \mathbb{R}^2$ is bounded and has smooth boundary. By a standard localization method however we can also show well-posedness for smooth, bounded domains. In a forthcoming paper discussing effects of gravity and Rayleigh-Taylor instability, cf. [26], this simpler geometry is useful. We denote the lateral walls of the cylinder Ω by $S_1 := \partial\Sigma \times (L_1, L_2)$ and bottom and top by $S_2 := \Sigma \times \{L_1, L_2\}$. As usual, $\nu_{\partial\Omega}$ denotes the unit normal vector field pointing outwards of Ω and $\nu_{S_1} = \nu_{\partial\Omega}$ on the walls S_1 . The projection to the tangent space of S_1 is defined by $P_{S_1} := I - \nu_{S_1} \otimes \nu_{S_1}$.

In a cylindrical domain the full problem for two possibly different, constant densities and viscosities reads as

$$\begin{aligned}
\rho \partial_t u - \mu \Delta u + \operatorname{div}[(\rho u + \llbracket \rho \rrbracket \nabla \eta) \otimes u] + \nabla p &= 0, & \text{in } \Omega \setminus \Gamma(t), \\
\operatorname{div} u &= 0, & \text{in } \Omega \setminus \Gamma(t), \\
-\llbracket \mu(Du + Du^\top) \rrbracket \nu_{\Gamma(t)} + \llbracket p \rrbracket \nu_{\Gamma(t)} &= \sigma H_{\Gamma(t)} \nu_{\Gamma(t)}, & \text{on } \Gamma(t), \\
\llbracket u \rrbracket &= 0, & \text{on } \Gamma(t), \\
V_{\Gamma(t)} - u|_{\Gamma(t)} \cdot \nu_{\Gamma(t)} &= -\llbracket \nu_{\Gamma(t)} \cdot \nabla \eta \rrbracket, & \text{on } \Gamma(t), \\
\nu_{\Gamma(t)} \cdot \nu_{S_1} &= 0, & \text{on } \partial\Gamma(t), \\
\Delta \eta &= 0, & \text{in } \Omega \setminus \Gamma(t), \\
\eta|_{\Gamma(t)} &= \sigma H_{\Gamma(t)}, & \text{on } \Gamma(t), \\
\nu_{\partial\Omega} \cdot \nabla \eta|_{\partial\Omega} &= 0, & \text{on } \partial\Omega \setminus \Gamma(t), \\
P_{S_1}(\mu(Du + Du^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial\Gamma(t), \\
u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial\Gamma(t), \\
u &= 0, & \text{on } S_2, \\
u(0) &= u_0, & \text{on } \Omega \setminus \Gamma(0), \\
\Gamma(0) &= \Gamma_0.
\end{aligned} \tag{1.1}$$

Here we want to mention that we implicitly impose that $\Gamma(t) \subset \Omega$ and $\partial\Gamma(t) \subset S_1$ for all $t \geq 0$, that is, the interface stays inside the domain for positive times and the boundary of the interface is contained in the boundary of the domain as well. This only makes sense from a physical standpoint.

Note that in this model proposed by Abels, Garcke, and Grün in [2] the momentum balance (1.1)₁ contains an extra term involving the chemical potential η since the densities in the two phases are different. This term however is needed to get an energy structure for the system, cf. Section 5 in [2]. It is shown there that the energy

$$E(t) := \int_{\Gamma(t)} \sigma d\mathcal{H}^{n-1} + \frac{1}{2} \int_{\Omega} \rho(t) u(t)^2 dx \tag{1.2}$$




FIGURE 1. The cylindrical capillary Ω with lateral walls S_1 and bottom and top S_2 . $G := \Sigma \times \{0\}$ is the reference surface and $\Gamma(t)$ the time-dependent free interface.

satisfies the energy-dissipation relation

$$\frac{d}{dt} E(t) = -D(t) := - \int_{\Omega} \mu |\mathbb{D}u(t)|^2 dx - \int_{\Omega} |\nabla \eta(t)|^2 dx. \quad (1.3)$$

Hereby, $\mathbb{D}u$ is the symmetric part of the gradient Du . There is a remark in order regarding this extra term in (1.1)₁. Since $\operatorname{div} u = 0$ and $\Delta \eta = 0$ in the bulk phases $\Omega \setminus \Gamma(t)$, we obtain

$$\operatorname{div}[(\rho u + (\rho^+ - \rho^-)\nabla \eta) \otimes u] = \rho(u \cdot \nabla)u + (\rho^+ - \rho^-)(\nabla \eta \cdot \nabla)u, \quad \text{in } \Omega \setminus \Gamma(t).$$

In the case of equivalent densities, say for simplicity $\rho = 1$, the extra term $\operatorname{div}[(\rho^+ - \rho^-)\nabla \mu \otimes u]$ vanishes and the system reduces to

$$\begin{aligned} \partial_t u - \mu^{\pm} \Delta u + (u \cdot \nabla)u + \nabla p &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ \operatorname{div} u &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ -\llbracket \mu^{\pm} (Du + Du^{\top}) \rrbracket \nu_{\Gamma(t)} + \llbracket p \rrbracket \nu_{\Gamma(t)} &= \sigma H_{\Gamma(t)} \nu_{\Gamma(t)}, & \text{on } \Gamma(t), \\ \llbracket u \rrbracket &= 0, & \text{on } \Gamma(t), \\ V_{\Gamma(t)} - u|_{\Gamma(t)} \cdot \nu_{\Gamma(t)} &= -\llbracket \nu_{\Gamma(t)} \cdot \nabla \eta \rrbracket, & \text{on } \Gamma(t), \\ \nu_{\Gamma(t)} \cdot \nu_{S_1} &= 0, & \text{on } \partial \Gamma(t), \\ \Delta \eta &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ \eta|_{\Gamma(t)} &= \sigma H_{\Gamma(t)}, & \text{on } \Gamma(t), \\ \nu_{\partial \Omega} \cdot \nabla \eta|_{\partial \Omega} &= 0, & \text{on } \partial \Omega \setminus \Gamma(t), \\ P_{S_1} (\mu^{\pm} (Du + Du^{\top}) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Gamma(t), \\ u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Gamma(t), \\ u &= 0, & \text{on } S_2, \\ u(0) &= u_0, & \text{on } \Omega \setminus \Gamma(0), \\ \Gamma(0) &= \Gamma_0. \end{aligned} \quad (1.4)$$

Note that in both cases, (1.1) and (1.4), individual masses are conserved,

$$\frac{d}{dt}|\Omega^\pm(t)| = 0, \quad t \in \mathbb{R}_+, \quad (1.5)$$

since $\Delta\eta = 0$ and $\operatorname{div} u = 0$ in the bulk phases $\Omega \setminus \Gamma(t)$.

Outline of this paper. In Section 2 we briefly introduce techniques and function spaces we work with. In Section 3 we rewrite the free boundary problem of the moving interface as a nonlinear problem for the height function parametrizing the interface. Section 4 deals with an analysis of the underlying linear problem proving maximal regularity in an $L_p - L_q$ scale for the distance function and an L_r scale for the velocity. Section 5 renders that the nonlinear problem is also well-posed, whereas Section 6 deals with qualitative behaviour, stability properties, and convergence to equilibrium solutions.

2. PRELIMINARIES AND FUNCTION SPACES

We now introduce function spaces and techniques we work with. For a more detailed discussion we refer the reader to the books of Triebel [28] and Prüss and Simonett [24].

2.1. Bessel-Potential, Besov and Triebel-Lizorkin Spaces. As usual, we will denote the classical L_p -Sobolev spaces on \mathbb{R}^n by $W_p^k(\mathbb{R}^n)$, where k is a natural number and $1 \leq p \leq \infty$. The Bessel-potential spaces will be denoted by $H_p^s(\mathbb{R}^n)$ for $s \in \mathbb{R}$ and the Sobolev-Slobodeckij spaces by $W_p^s(\mathbb{R}^n)$. We will also denote the usual Besov spaces by $B_{pr}^s(\mathbb{R}^n)$, where $s \in \mathbb{R}, 1 \leq p, r \leq \infty$. Lastly, the Triebel-Lizorkin spaces are denoted by $F_{pr}^s(\mathbb{R}^n)$.

These function spaces on a domain $\Omega \subset \mathbb{R}^n$ are defined in a usual way by restriction. The Banach space-valued versions of these spaces are denoted by $L_p(\Omega; X)$, $W_p^k(\Omega; X)$, $H_p^s(\Omega; X)$, $W_p^s(\Omega; X)$, $B_{pr}^s(\Omega; X)$, $F_{pr}^s(\Omega; X)$, respectively. For precise definitions we refer to [19].

For results on embeddings, traces, interpolation and extension operators we refer to [1], [8], [17], [18], [24], [27], [28].

2.2. Maximal Regularity. Let us recall the property of maximal L_p -regularity, as is e.g. done in Definition 3.5.1 in [24].

Definition 2.1. *Let X be a Banach space, $J = (0, T), 0 < T < \infty$ or $J = \mathbb{R}_+$, and A a closed, densely defined operator on X with domain $D(A) \subset X$. Then the operator A is said to have maximal L_p -regularity on J , if and only if for every $f \in L_p(J; X)$ there is a unique $u \in W_p^1(J; X) \cap L_p(J; D(A))$ solving*

$$\frac{d}{dt}u(t) + Au(t) = f(t), \quad t \in J, \quad u|_{t=0} = 0,$$

in an almost-everywhere sense in $L_p(J; X)$.

There is a wide class of results on operators having maximal regularity, we refer to [5], [6], [9], [10], [11], [20], [21], and [24], for further discussion.

3. REDUCTION TO A FLAT INTERFACE

In this section we transform the equations defined on the time-dependent domain $\Omega \setminus \Gamma(t)$ and the moving interface $\Gamma(t)$ to a fixed reference frame. We follow the ideas of [29], see also [15], [23], [3], [14], [25], [22]. To simplify notation let $n = 3$, the modifications for $n = 2$ are obvious.

We now assume that the interface at time t is given as a graph over the fixed reference surface $\Sigma := \Omega \cap \{x_3 = 0\}$. More precisely, we assume that there is a height function $h : \Sigma \times [0, \infty) \rightarrow (L_1, L_2)$, such that

$$\Gamma(t) = \Gamma_h(t) := \{x \in \Sigma \times (L_1, L_2) : x_3 = h(x', t), \ x' = (x_1, x_2) \in \Sigma\}, \quad t \geq 0.$$

We will now construct a Hanzawa-type transformation, which is an isomorphism on Ω and maps the moving interface $\Gamma(t)$ to the reference surface Σ for every $t \geq 0$. To this end pick some smooth bump function $\chi \in C_0^\infty(\mathbb{R}; [0, 1])$ such that $\chi(s) = 1$ for $|s| \leq \delta/2$ and $\chi(s) = 0$ for $|s| \geq \delta$, where $0 < \delta \leq \min\{-L_1, L_2\}/3$. Define a mapping

$$\Theta_h : \Omega \times \mathbb{R}_+ \rightarrow \Omega, \quad \Theta_h(x, t) := x + \chi(x_3)h(x', t)e_3 =: x + \theta_h(x, t),$$

where $x = (x', x_3)$. Then

$$D\Theta_h = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \partial_1 h \chi & \partial_2 h \chi & 1 + h \chi' \end{pmatrix}.$$

It clearly follows that $D\Theta_h$ is a regular matrix and Θ_h is invertible, provided $h\chi'$ is sufficiently small. For instance, this is the case whenever

$$|h|_{L_\infty((0, T) \times \Sigma)} \leq \frac{1}{2|\chi'|_{L_\infty(\mathbb{R})}}.$$

Note that $|\chi'|_\infty$ can be bounded by a constant depending on δ only. For the sequel we will fix the bump function χ and choose $0 < d_0 < 1/(2|\chi'|_\infty)$ sufficiently small and assume that $|h|_\infty \leq d_0$. This way we ensure that the inverse $\Theta_h^{-1} : \Omega \rightarrow \Omega$ is well defined and maps the free interface $\Gamma(t)$ to the fixed reference surface Σ .

We will now calculate how the equations behave under this transformation. Define the transformed quantities

$$w(x, t) := u(\Theta_h(x, t), t), \quad q(x, t) := p(\Theta_h(x, t), t), \quad \vartheta(x, t) := \eta(\Theta_h(x, t), t),$$

for $x \in \Omega$, $t \in \mathbb{R}_+$. We now determine the equations which (w, q, ϑ) solve. Define $D\Theta_h^{-\top} := ((D\Theta_h)^{-1})^\top$, as well as the transformed quantities

$$\nabla_h := D\Theta_h^{-\top} \nabla, \quad \nabla_h u := (\nabla_h u_k^\top)_{k=1}^3, \quad \operatorname{div}_h := \operatorname{Tr}(\nabla_h), \quad \Delta_h := \operatorname{div}_h \nabla_h.$$

With this it is straightforward to check that

$$\begin{aligned} \nabla u(\Theta_h(x, t), t) &= \nabla_h w(x, t), \quad [(u \cdot \nabla)u](\Theta_h(x, t), t) = [(w \cdot \nabla_h)w](x, t), \\ \Delta u(\Theta_h(x, t), t) &= \Delta_h w(x, t), \quad \operatorname{div} u(\Theta_h(x, t), t) = \operatorname{div}_h w(x, t), \quad x \in \Omega, t \in \mathbb{R}_+. \end{aligned}$$

Furthermore,

$$\partial_t u(\Theta_h(x, t), t) = \partial_t w(x, t) + Dw(x, t) \partial_t \Theta_h^{-1}(\Theta_h(x, t), t), \quad x \in \Omega, t \in \mathbb{R}_+.$$

The upper unit normal at the free interface $\Gamma(t)$ and the normal velocity of which can both be expressed in terms of h by

$$\nu_{\Gamma(t)} = \frac{(-\nabla h, 1)^\top}{\sqrt{1 + |\nabla h|^2}}, \quad V_{\Gamma(t)} = \frac{\partial_t h}{\sqrt{1 + |\nabla h|^2}}, \quad x \in \Sigma, t \in \mathbb{R}_+.$$

We are now able to transform the Two-phase Navier-Stokes/Mullins-Sekerka system (1.1) to the fixed reference frame, the transformed system reads as

$$\begin{aligned} \rho^\pm \partial_t w - \mu^\pm \Delta w + \nabla q &= a^\pm(h; D_x, D_x^2)(w, q) + \bar{a}(h, w), & \text{in } \Omega \setminus \Sigma, \\ \operatorname{div} w &= G_d(h, w), & \text{in } \Omega \setminus \Sigma, \\ -\llbracket \mu^\pm (Dw + Dw^\top) - qI \rrbracket \nu_\Sigma &= \sigma \Delta_{x'} h \nu_\Sigma + G_S(h, w, q), & \text{on } \Sigma, \\ \llbracket w \rrbracket &= 0, & \text{on } \Sigma, \\ \partial_t h &= w \cdot \nu_\Sigma - \llbracket \partial_3 \vartheta \rrbracket + G_\Sigma(h, w, \vartheta), & \text{on } \Sigma, \\ (-\nabla_{x'} h, 1)^\top \cdot \nu_{S_1} &= 0, & \text{on } \partial \Sigma, \\ \Delta \vartheta &= G_c(h, \vartheta), & \text{in } \Omega \setminus \Sigma, \\ \vartheta|_\Sigma - \sigma \Delta_{x'} h &= G_\kappa(h), & \text{on } \Sigma, \\ \nu_{\partial \Omega} \cdot \nabla \vartheta|_{\partial \Omega} &= G_N(h, \vartheta), & \text{on } \partial \Omega \setminus \Sigma, \\ P_{S_1}(\mu^\pm (Dw + Dw^\top) \nu_{S_1}) &= G_P^\pm(h, w), & \text{on } S_1 \setminus \partial \Sigma, \\ w \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\ w &= 0, & \text{on } S_2, \\ w(0) &= w_0, & \text{on } \Omega \setminus \Sigma, \\ h(0) &= h_0, & \text{on } \Sigma, \end{aligned} \tag{3.1}$$

where $\nu_\Sigma = e_3$, and

$$\begin{aligned} a^\pm(h; D_x, D_x^2)(w, q) &:= \mu^\pm (\Delta_h - \Delta)w + (\nabla - \nabla_h)q, \\ \bar{a}(h, w, \vartheta) &:= Dw \cdot \partial_t \Theta_h^{-1} - (w \cdot \nabla_h)w - (\rho^+ - \rho^-)(\nabla_h \vartheta \cdot \nabla_h)w, \\ G_d(h, w) &:= (\operatorname{div} - \operatorname{div}_h)w, \\ G_S(h, w, q) &:= \llbracket \mu^\pm ((D\Theta_h - I)Dw + Dw^\top (D\Theta_h - I)^\top) \rrbracket \nu_{\Gamma_h} + \\ &\quad + \llbracket (\mu^\pm (Dw + Dw^\top) - qI) (e_3 - \nu_{\Gamma_h}) \rrbracket + \sigma(K(h)\nu_{\Gamma_h} - \Delta_{x'} h e_3), \\ G_\Sigma(h, w, \vartheta) &:= w \cdot (-\nabla_{x'} h, 0)^\top - \llbracket e_3 \cdot (\nabla - \nabla_h)\vartheta \rrbracket - \llbracket (-\nabla_{x'} h, 0)^\top \cdot \nabla_h \vartheta \rrbracket, \\ G_c(h, \vartheta) &:= (\Delta - \Delta_h)\vartheta, \\ G_\kappa(h) &:= \sigma(K(h) - \Delta_{x'} h), \\ G_N(h, \vartheta) &:= \nu_{\partial \Omega} \cdot (\nabla - \nabla_h)\vartheta, \\ G_P^\pm(h, w) &:= P_{S_1}(\mu^\pm ((D\Theta_h - I)Dw + Dw^\top (D\Theta_h - I)^\top) \nu_{S_1}). \end{aligned}$$

Here, cf. [13], the mean curvature is given in terms of h by

$$K(h) = H(\Gamma_h) = \operatorname{div} \left(\frac{\nabla h}{\sqrt{1 + |\nabla h|^2}} \right), \quad x \in \Sigma, t \in \mathbb{R}_+.$$

Furthermore, we want to point out that we used the fact that the normal ν_{S_1} is independent of x_3 and that the transformation Θ_h leaves the Dirichlet-boundary S_2 invariant.

Since $\nu_\Sigma = e_3$, one can also easily decompose the stress tensor condition $(3.1)_3$ into tangential and horizontal parts, cf. [29]. Then, $(3.1)_3$ reads as

$$\begin{aligned} -\llbracket \mu^\pm \partial_3(w_1, w_2) \rrbracket - \llbracket \mu^\pm \nabla_{x'} w_3 \rrbracket &= (G_S(h, w, q))_{1,2}, & \text{on } \Sigma, \\ -2\llbracket \mu^\pm \partial_3 w_3 \rrbracket + \llbracket q \rrbracket - \sigma \Delta_{x'} h &= (G_S(h, w, q))_3, & \text{on } \Sigma. \end{aligned}$$

To economize notation, we define

$$G_S^\parallel(h, w, q) := (G_S(h, w, q))_{1,2}, \quad G_S^\perp(h, w, q) := (G_S(h, w, q))_3.$$

Hereby we understand $a_{1,2}$ to be (a_1, a_2) for a given vector $a = (a_1, a_2, a_3) \in \mathbb{R}^3$.

4. MAXIMAL REGULARITY OF THE LINEAR PROBLEM

The main goal of this section is to derive a maximal regularity result for the linearization of (1.1).

4.1. Linearization, regularity and compatibility conditions. In this section we consider the linear part of the Two-phase Navier-Stokes/Mullins-Sekerka system, which reads as

$$\begin{aligned} \rho^\pm \partial_t u - \mu^\pm \Delta u + \nabla \pi &= g_1, & \text{in } \Omega \setminus \Sigma, \\ \operatorname{div} u &= g_2, & \text{in } \Omega \setminus \Sigma, \\ -\llbracket \mu^\pm \partial_3(u_1, u_2) \rrbracket - \llbracket \mu^\pm \nabla_{x'} u_3 \rrbracket &= g_3, & \text{on } \Sigma, \\ -2\llbracket \mu^\pm \partial_3 u_3 \rrbracket + \llbracket \pi \rrbracket - \sigma \Delta_{x'} h &= g_4, & \text{on } \Sigma, \\ \llbracket u \rrbracket &= g_5, & \text{on } \Sigma, \\ \partial_t h - (u_3^+ + u_3^-)/2 + \llbracket \partial_3 \eta \rrbracket &= g_6, & \text{on } \Sigma, \\ (-\nabla_{x'} h, 0)^\top \cdot \nu_{S_1} &= g_7, & \text{on } \partial \Sigma, \\ \Delta \eta &= g_8, & \text{in } \Omega \setminus \Sigma, \\ \eta|_\Sigma - \sigma \Delta_{x'} h &= g_9, & \text{on } \Sigma, \\ \nu_{\partial \Omega} \cdot \nabla \eta|_{\partial \Omega} &= g_{10}, & \text{on } \partial \Omega \setminus \Sigma, \\ P_{S_1}(\mu^\pm (Du + Du^\top) \nu_{S_1}) &= P_{S_1} g_{11}, & \text{on } S_1 \setminus \partial \Sigma, \\ u \cdot \nu_{S_1} &= g_{12}, & \text{on } S_1 \setminus \partial \Sigma, \\ u &= g_{13}, & \text{on } S_2, \\ u(0) &= u_0, & \text{on } \Omega \setminus \Sigma, \\ h(0) &= h_0, & \text{on } \Sigma. \end{aligned} \tag{4.1}$$

Here we take $(u_3^+ + u_3^-)/2$ instead of the trace of u in equation $(4.1)_6$ since u is allowed to have a jump across Σ . Hereby u_3^\pm denote the directional traces of u_3 with respect to $\{x_3 \gtrless 0\}$.

4.2. Regularity of the solution. The question of function spaces is now a very delicate matter. The main idea already used by Abels and Wilke in the case of no boundary contact [4] is to treat the Navier-Stokes part of the evolution as lower order compared to the Mullins-Sekerka part. They consider some height function h as given and solve the two-phase Navier-Stokes equations in dependence of h by a function $u = u(h)$. Afterwards plugging in the solution $u(h)$ in the evolution equation for h they obtain a problem only dependent on h . If now u is sufficiently more regular as the other terms in the evolution equation for $\partial_t h$, the Navier-Stokes equations can be seen as a lower order perturbation. By choosing the time interval sufficiently small one gets well-posedness also for the coupled system, stemming from the unique solvability of the pure Mullins-Sekerka evolution of h .

Let us begin by recalling the maximal regularity class for h of the pure Mullins-Sekerka system with boundary contact, cf. [3]. For $6 < p < \infty$ and $q \in (5/3, 2) \cap (2p/(p+1), 2)$, we obtained a unique local in time strong solution

$$h \in W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma)),$$

and $\eta \in L_p(0, T; W_q^2(\Omega \setminus \Sigma))$ of the linearized Mullins-Sekerka with boundary contact for some $T > 0$, cf. Theorem 5.1 in [3].

Two things are important for the analysis: to be later able to treat the Navier-Stokes part as lower order, we need to know that $u|_\Sigma$ has better time regularity and at least as much space regularity as the other terms in $(4.1)_6$, namely $L_p(0, T; W_q^{1-1/q}(\Sigma))$. On the other hand, the linearized curvature term $\Delta_{x'} h$ has to be at least of the same regularity as $Du|_\Sigma$, cf. $(4.1)_4$. By choosing a setting where u is too regular, $\Delta_{x'} h$ fails to be admissible data, and by choosing u not regular enough, $u|_\Sigma$ may not be treated as a lower order perturbation. In the following lines we want to explain a setting of function spaces, in which the coupling is of lower order and u is still regular enough to control the nonlinear terms.

The first possibility is to choose an $L_p - L_p$ ansatz, where p as above is large. The vector field u would then be very regular, hence making the nonlinearities easy to handle since in particular $p > 5$. In this ansatz we search for

$$u \in W_p^1(0, T; L_p(\Omega)) \cap L_p(0, T; W_p^2(\Omega \setminus \Sigma)),$$

whence by classical theory, $u \in BUC([0, T]; W_p^{2-2/p}(\Omega \setminus \Sigma))$. Taking traces yields $u|_\Sigma \in BUC([0, T]; W_p^{2-3/p}(\Sigma))$, and hence it can be seen as a lower order perturbation in $L_p(0, T; W_q^{1-1/q}(\Sigma))$. However,

$$\Delta_{x'} h \in W_p^{2/3-1/(3q)}(0, T; L_q(\Sigma)) \cap L_p(0, T; W_q^{2-1/q}(\Sigma)), \quad (4.2)$$

on the other hand,

$$Du|_\Sigma \in W_p^{1/2-1/(2p)}(0, T; L_p(\Sigma)) \cap L_p(0, T; W_p^{1-1/p}(\Sigma)).$$

It is now a consequence of Sobolev-type embedding theorems to see that $W_q^{2-1/q}(\Sigma)$ does not embed into $W_p^{1-1/p}(\Sigma)$ in general, due to $5/3 < q < 2$ and $p > 6$. Hence this $L_p - L_p$ ansatz with large p does not work.

Alternatively, one can make an $L_q - L_q$ ansatz, searching for some

$$u \in W_q^1(0, T; L_q(\Omega)) \cap L_q(0, T; W_q^2(\Omega \setminus \Sigma)),$$

where $5/3 < q < 2$. Clearly, the function u possesses way less regularity in this ansatz. It is then easy to check that $\Delta_{x'}h$ is admissible data by comparing the regularity classes of $\Delta_{x'}h$ and $Du|_{\Sigma}$. Also,

$$u|_{\Sigma} \in L_{2q/(2-q)}(0, T; W_q^{1-1/q}(\Sigma)).$$

Note that as $q \rightarrow 2$, the time regularity index $2q/(2-q)$ tends to $+\infty$. Hence the Stokes part may be treated as lower order whenever $q < 2$ is close to 2. However we want to point out that handling the nonlinearities may be more difficult since certain Sobolev embeddings fail since $q < 2$.

By choosing an $L_p - L_q$ approach one may get better regularity for u , however if one takes any trace of u on the boundary, for instance in the simplest case of the Dirichlet conditions on top and bottom of the container, one ends up with Triebel-Lizorkin spaces in time. It is well known that the optimal regularity for the trace of a function

$$u \in W_p^1(0, T; L_q(\Omega)) \cap L_p(0, T; W_q^2(\Omega \setminus \Sigma))$$

on the boundary, e.g. S_2 , is

$$u|_{S_2} \in F_{pq}^{1-1/(2q)}(0, T; L_q(S_2)) \cap L_p(0, T; W_q^{2-1/q}(S_2)).$$

It is particularly hard to treat this problem in a mixed $L_p - L_q$ setting, since even in the model problems it is not clear how to generalize for instance the results of Prüss and Simonett in [23] regarding the Dirichlet-to-Neumann operator. This operator is well understood in an $L_p - L_p$ setting, however the proof of Proposition 3.3 given in [23] is not easily generalizable to a mixed setting where $p \neq q$. The proof heavily relies on real interpolation method and Triebel-Lizorkin spaces do not naturally arise as real interpolation spaces.

The explanations above motivate our introduction of a third integration scale. We will show that for given $q < 2$ sufficiently close to 2 and $6 < p < \infty$ finite but large, there is some exponent $3 < r = r(q) < 7/2 < \infty$ such that the following is true: $\Delta_{x'}h$ is admissible data in the Stokes part, and $u|_{\Sigma}$ is lower order in the evolution equation for h . This $L_r - L_r$ approach with $r > 3$ circumvents the problem of Triebel-Lizorkin data spaces in the Stokes part completely and hence makes the problem a lot easier to tackle. Also it allows to make use of known results of Prüss and Simonett in [23] and makes the nonlinearities easier to handle in the contraction estimates. We will give the precise choice of r below in Theorem 4.1 and prove the above assertions rigorously.

Theorem 4.1. *Let $n = 3$, that is, $\dim \Sigma = n - 1 = 2$. Let $5/3 < q < 2$ and $6 < p < \infty$. Furthermore, let $0 < T \leq T_0$ for some fixed $T_0 < \infty$. Let*

$$2 \leq r < \frac{7}{6/q - 1}.$$

Then, for any $h \in W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma))$, we have that

$$\Delta_{x'}h \in W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma)).$$

Furthermore, there is some $C = C(T) > 0$, such that

$$\begin{aligned} & |\Delta_{x'}h|_{W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma))} \\ & \leq C(T)|h|_{W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma))}. \end{aligned} \tag{4.3}$$

Furthermore, if $2 > q > 9/5$, we can choose r to satisfy $3 < r < 7/2$. If $3 < r < 7/2$, we have

$$\begin{aligned} W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma)) &\hookrightarrow \\ &\hookrightarrow L_\infty(0, T; L_\infty(\Omega)) \cap L_\infty(0, T; W_r^1(\Omega \setminus \Sigma)) \cap L_r(0, T; W_\infty^1(\Omega \setminus \Sigma)). \end{aligned} \quad (4.4)$$

Moreover,

$$\text{tr}_\Sigma : W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma)) \rightarrow L_\infty(0, T; W_q^{1-1/q}(\Sigma)), \quad (4.5)$$

is bounded provided the trace on Σ is well defined, for instance if $\llbracket u \rrbracket = 0$. Otherwise the statement is true for the restrictions on Ω^\pm , that is, $\text{tr}_\Sigma^\pm : u \mapsto u^\pm|_\Sigma$, where $u^\pm := u|_{\Omega^\pm}$.

By restricting to height functions h with initial trace zero, $h(0) = 0$, the embedding constant in (4.3) can be chosen to be independent of T and only depending on T_0 . In particular, the embedding does not degenerate and the embedding constant stays bounded as $T \downarrow 0$.

Restricting to vanishing traces at $t = 0$ in (4.4), the embedding constant is also independent of $T > 0$.

Proof. Let $2 \leq r \leq p$ and $0 < T \leq T_0$. Note that due to $p \geq r$, we have that $L_p(0, T) \hookrightarrow L_r(0, T)$. The embedding constant here only depends on T_0 , which stems from Hölder's inequality,

$$|f|_{L_r(0, T)} \leq T^{(p-r)/(pr)} |f|_{L_p(0, T)} \leq T_0^{(p-r)/(pr)} |f|_{L_p(0, T)}, \quad f \in L_p(0, T).$$

Now, due to Sobolev's embedding theorem, $W_q^{2-1/q}(\Sigma) \hookrightarrow W_r^{1-1/r}(\Sigma)$, provided that $2 - 3/q > 1 - 3/r$, which gives an upper restriction on r reading as

$$r < \frac{3q}{3-q}, \quad (4.6)$$

cf. [1], [28]. Summing up, $L_p(0, T; W_q^{2-1/q}(\Sigma)) \hookrightarrow L_r(0, T; W_r^{1-1/r}(\Sigma))$, provided $r \leq p$ and (4.6) holds.

Since we want to use the results of [16] on the half line, we now consider some

$$h \in W_p^1(\mathbb{R}_+; W_q^{1-1/q}(\Sigma)) \cap L_p(\mathbb{R}_+; W_q^{4-1/q}(\Sigma)).$$

Firstly, using Proposition 5.37 in [16] on the half line, $h \in H_p^\theta(\mathbb{R}_+; W_q^{1-1/q+3(1-\theta)}(\Sigma))$, whenever $\theta \in (0, 1)$. It follows that

$$\Delta_{x'} h \in H_p^\theta(\mathbb{R}_+; W_q^{2-1/q-3\theta}(\Sigma)), \quad \theta \in (0, 1).$$

Let $\epsilon > 0$ small. By choosing $\theta := 2/3 - 1/q + 2/(3r) - \epsilon \in (0, 1)$, we obtain

$$\Delta_{x'} h \in H_p^{2/3-1/q+2/(3r)-\epsilon}(\mathbb{R}_+; W_q^{2/q-2/r+3\epsilon}(\Sigma)).$$

By Sobolev embeddings for Besov spaces,

$$\Delta_{x'} h \in H_p^{2/3-1/q+2/(3r)-\epsilon}(0, T; L_r(\Sigma)),$$

for any small $\epsilon > 0$. Assume for a moment that

$$2/3 - 1/q + 2/(3r) > 1/2 - 1/(2r). \quad (4.7)$$

Then we may choose $\epsilon > 0$ so small, such that

$$2/3 - 1/q + 2/(3r) - \epsilon > 1/2 - 1/(2r). \quad (4.8)$$

Then $\Delta_{x'} h \in W_p^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \hookrightarrow W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma))$. Inequality (4.7) however is equivalent to $r < 7/(6/q - 1)$ since $q < 2$. Estimate (4.3) is a direct consequence of these considerations. Furthermore, whenever h has vanishing trace at $t = 0$, a standard extension argument allows to see that the estimate does not degenerate as $T \rightarrow 0$, that is, $C(T)$ stays bounded as $T \rightarrow 0$ since it only depends on T_0 .

Choosing $q < 2$ close enough to 2 we may assume that $r > 3$, since $7/(6/q - 1) \rightarrow 7/2$ as $q \rightarrow 2$. Now let $u \in W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma))$ for $r > 3$. We may use the embedding

$$W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma)) \hookrightarrow BUC([0, T]; W_r^{2-2/r}(\Omega \setminus \Sigma)), \quad (4.9)$$

cf. [5], to see that $Du \in BUC([0, T]; W_r^{1-2/r}(\Omega \setminus \Sigma))$, which then in turn yields $Du \in L_\infty(0, T; L_r(\Omega))$. It also follows that $Du \in L_r(0, T; L_\infty(\Omega))$. Regarding the trace operator, we note that

$$BUC([0, T]; W_r^{2-2/r}(\Omega \setminus \Sigma)) \hookrightarrow L_\infty(0, T; W_q^1(\Omega \setminus \Sigma)), \quad (4.10)$$

whenever $r \geq 5q/(q + 3)$, which is surely satisfied since $r > 3$ and $q < 2$. The proof is complete. \square

Remark 4.2. *Let us comment on the regularity of solutions.*

- (1) *Note that we can choose from now on $p \in (6, \infty)$, $q \in (9/5, 2) \cap (2p/(p+1), 2)$, and $r \in (3, 7/2)$. In particular, the set of admissible indices is not empty.*
- (2) *Note that if $r > 5/2$, it holds that $u \in BUC([0, T]; C^0(\Omega))$.*
- (3) *There is still room for improvement in these embeddings. For instance, u is $L_\infty(0, T; L_\infty(\Omega))$ whenever $r \geq 5/2$. Furthermore, it can be shown that*

$$Du \in L_{2r/(5-r)-\varepsilon}(0, T; L_\infty(\Omega)) \cap L_\infty(0, T; L_{3r/(5-r)-\varepsilon}(\Omega)),$$

for any small $\varepsilon > 0$. This may be used to lower the index r and consider initial data with lower regularity.

This motivates to choose the following setting for the solutions to the Two-phase Navier-Stokes/Mullins-Sekerka system and its linearization (4.1).

Let $T \in (0, \infty)$, $p \in (6, \infty)$, $q \in (9/5, 2) \cap (2p/(p+1), 2)$, and $r \in (3, 7/2)$ as in Theorem 4.2. From now on, we will fix the integration scales p, q and r . We are looking for solutions (u, π, h, μ) of (4.1) with

$$\begin{aligned} u &\in W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma)), \quad \pi \in L_r(0, T; \dot{H}_r^1(\Omega)), \\ \llbracket \pi \rrbracket &\in W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma)), \\ h &\in W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma)), \quad \mu \in L_p(0, T; W_q^2(\Omega \setminus \Sigma)). \end{aligned}$$

4.3. Regularity of the data. To be able to derive a maximal regularity result, we will now deduce optimal regularity classes for the data in problem (4.1). Given a solution $(u, \pi, \llbracket \pi \rrbracket, h, \mu)$ in the classes of (4.2), we derive by standard trace theory

the following necessary conditions for the data,

$$\begin{aligned}
g_1 &\in L_r(0, T; L_r(\Omega)), \quad g_2 \in L_r(0, T; W_r^1(\Omega \setminus \Sigma)), \\
g_3, g_4 &\in W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma)), \\
g_5 &\in W_r^{1-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{2-1/r}(\Sigma)), \\
g_6 &\in L_p(0, T; W_q^{1-1/q}(\Sigma)), \quad g_8 \in L_p(0, T; L_q(\Omega)), \\
g_7 &\in F_{pq}^{1-2/(3q)}(0, T; L_q(\partial\Sigma)) \cap L_p(0, T; W_q^{3-2/q}(\partial\Sigma)), \\
g_9 &\in L_p(0, T; W_q^{2-1/q}(\Sigma)), \quad g_{10} \in L_p(0, T; W_q^{1-1/q}(\partial\Omega)), \\
P_{S_1} g_{11} &\in W_r^{1/2-1/(2r)}(0, T; L_r(S_1)) \cap L_r(0, T; W_r^{1-1/r}(S_1)), \\
g_{12} &\in W_r^{1-1/(2r)}(0, T; L_r(S_1)) \cap L_r(0, T; W_r^{2-1/r}(S_1)), \\
g_{13} &\in W_r^{1-1/(2r)}(0, T; L_r(S_2)) \cap L_r(0, T; W_r^{2-1/r}(S_2)), \\
u_0 &\in W_r^{2-2/r}(\Omega \setminus \Sigma), \quad h_0 \in B_{qp}^{4-3/p-1/q}(\Sigma).
\end{aligned} \tag{4.11}$$

For the regularity of g_7 we refer to Appendix A in [3]. At this point we note that in (4.11) the function g_2 does not have to have the time regularity of Du in $\Omega \setminus \Sigma$. This is due to the fact that there is some compatibility condition hidden in the system stemming from the divergence equation, which inherits a certain time regularity for $(g_2, g_5, g_{12}, g_{13})$. This will be discussed in the next section regarding compatibility conditions. However we clearly want to point out that g_2 being $L_r(0, T; W_r^1(\Omega \setminus \Sigma))$ alone is a necessary but not a sufficient condition.

4.4. Compatibility conditions. We now shall discuss all the compatibility conditions for the data $(g_j)_{j=1}^{13}, u_0, h_0$ of system (4.1). In Lemma 4.3 below we rigorously show these conditions all occur and are well-defined. The following observations have already been made in [3] and [29].

At the starting point of the evolution at time $t = 0$ we have to have that

$$\begin{aligned}
\operatorname{div} u_0 &= g_2|_{t=0}, \quad -\llbracket \mu^\pm \partial_3(u_0)_{1,2} \rrbracket - \llbracket \mu^\pm \nabla_{x'}(u_0)_3 \rrbracket = g_3|_{t=0}, \\
\llbracket u_0 \rrbracket &= g_5|_{t=0}, \quad u_0 \cdot \nu_{S_1} = g_{12}|_{t=0}, \quad u_0|_{S_2} = g_{13}|_{t=0}, \\
(-\nabla_{x'} h_0, 1)^\top \cdot \nu_{S_1} &= g_7|_{t=0}, \quad P_{S_1}(\mu^\pm(Du_0 + Du_0^\top)\nu_{S_1}) = P_{S_1} g_{11}|_{t=0},
\end{aligned} \tag{4.12}$$

by evaluating the respective equations at time zero. Here, $(u_0)_{1,2}$ denotes the vector in \mathbb{R}^2 with the first two entries of u_0 , similarly $(u_0)_3$ denotes the last entry of u_0 .

Since $\partial\Sigma \subseteq S_1 \neq \emptyset$ and bottom, top and walls of the container have a common boundary, $\partial S_1 \cap \partial S_2 \neq \emptyset$, there are additional compatibility conditions. Simply by

comparing equations we get

$$\begin{aligned}
\llbracket g_{12} \rrbracket &= g_5 \cdot \nu_{S_1}, & \text{on } \partial\Sigma, \\
\llbracket (g_{11} \cdot e_3)/\mu^\pm - \partial_3 g_{12} \rrbracket &= \partial_{\nu_{S_1}}(g_5 \cdot e_3), & \text{on } \partial\Sigma, \\
P_{\partial\Sigma}[(D_x \Pi g_5 + (D_x \Pi g_5)^\top) \nu_{\partial\Sigma}] &= \llbracket P_{\partial\Sigma} \Pi g_{11} / \mu^\pm \rrbracket, & \text{on } \partial\Sigma, \\
g_3 \cdot (\nu_{S_1})_{1,2} &= -\llbracket g_{11} \cdot e_3 \rrbracket, & \text{on } \partial\Sigma, \\
g_{13} \cdot \nu_{S_1} &= g_{12}, & \text{on } \partial S_2, \\
P_{\partial\Sigma}[\mu^\pm (D_x \Pi g_{13} + (D_x \Pi g_{13})^\top) \nu_{\partial\Sigma}] &= P_{\partial\Sigma} \Pi g_{11}, & \text{on } \partial S_2, \\
\mu^\pm \partial_{\nu_{S_1}}(g_{13} \cdot e_3) + \mu^\pm \partial_3 g_{12} &= g_{11} \cdot e_3, & \text{on } \partial S_2.
\end{aligned} \tag{4.13}$$

Here, $\Pi v := (v_1, v_2) \in \mathbb{R}^2$ for $v = (v_1, v_2, v_3) \in \mathbb{R}^3$ and $\nu_{\partial\Sigma} := \Pi \nu_{S_1}$. The projection then is given by $P_{\partial\Sigma} := I - \nu_{\partial\Sigma} \otimes \nu_{\partial\Sigma}$. For further discussion we refer to [29].

We want to point out that there is no additional compatibility condition for $\partial_t g_7$ on $\partial\Sigma$ as there is in [29], since g_7 does not have a well defined time derivative on $\partial\Sigma$ in our regularity class. This is due to the fact that we have a different maximal regularity class for h as in [29].

Finally we turn to the divergence equation and want to point out that there is another compatibility and regularity condition hidden in the system, which has already been investigated in [29]. For completeness we explain it here briefly.

Consider the divergence equation $\operatorname{div} u = g_2$ and multiply this equation with a testfunction $\varphi \in W_{r'}^1(\Omega)$, where $r' = r/(r-1)$ is the conjugate exponent. An integration by parts on the two Lipschitz domains $\Omega \cap \{x_3 \geq 0\}$ and using the equations entails that

$$\begin{aligned}
&\int_{\Omega \setminus \Sigma} g_2 \varphi dx - \int_{S_1} g_{12} \varphi|_{S_1} dS_1 - \int_{S_2} (g_{13} \cdot \nu_{S_2}) \varphi|_{S_2} dS_2 \\
&\quad + \int_{\Sigma} (g_5 \cdot \nu_{\Sigma}) \varphi|_{\Sigma} d\Sigma = - \int_{\Omega \setminus \Sigma} u \cdot \nabla \varphi dx,
\end{aligned} \tag{4.14}$$

see also Proposition A.14 in [29]. Hence the functional $\varphi \mapsto \langle (g_2, g_5, g_{12}, g_{13}), \varphi \rangle$ defined by the left hand side of (4.14) is continuous on $W_{r'}^1(\Omega)$ with respect to the seminorm $|\nabla \cdot|_{L_{r'}(\Omega)}$. Since $C_0^\infty(\Omega) \subseteq W_{r'}^1(\Omega)$ is dense in the homogeneous space $\dot{H}_{r'}^1(\Omega)$ with respect to this seminorm, it follows that $\varphi \mapsto \langle (g_2, g_5, g_{12}, g_{13}), \varphi \rangle$ defines a functional on $\dot{H}_{r'}^1(\Omega)$. In other words, $(g_2, g_5, g_{12}, g_{13}) \in \hat{H}_r^{-1}(\Omega) := (\dot{H}_{r'}^1(\Omega))'$. The norm of $(g_2, g_5, g_{12}, g_{13})$ in $\hat{H}_r^{-1}(\Omega)$ is then given by

$$|(g_2, g_5, g_{12}, g_{13})|_{\hat{H}_r^{-1}(\Omega)} := \sup\{ \langle (g_2, g_5, g_{12}, g_{13}), \varphi \rangle / |\nabla \varphi|_{L_{r'}(\Omega)} : \varphi \in W_{r'}^1(\Omega) \}.$$

We now turn again to the equations. Since $u \in W_r^1(0, T; L_r(\Omega))$, it follows from (4.14) that $\frac{d}{dt}(g_2, g_5, g_{12}, g_{13})$ is well defined and is in $L_r(0, T; \hat{H}_r^{-1}(\Omega))$. Consequently,

$$(g_2, g_5, g_{12}, g_{13}) \in W_r^1(0, T; \hat{H}_r^{-1}(\Omega)) \tag{4.15}$$

is another necessary compatibility and regularity condition. We close this subsection by showing that the compatibility conditions we have deduced above are all well-defined conditions.

Lemma 4.3. *Let $r > 3$. Then all appearing traces and hence the compatibility conditions are all well-defined.*

Proof. Firstly, g_j , $j = 2, 3, 5, 7, 12, 13$, and $P_{S_1}g_{11}$ all have a well-defined trace at $t = 0$ since $r > 3$. Indeed, the condition for g_7 is independent of r (and fulfilled by choice of p and q) and the rest easily follow by trace theory. Pick for instance g_3 . Then g_3 surely has a trace at $t = 0$ whenever $1/2 - 1/(2r) - 1/r > 0$. This is however equivalent to $r > 3$. By taking traces in the spatial variables one easily sees that all the other traces are well-defined. \square

4.5. Maximal regularity. Let us consider the linear problem

$$\begin{aligned}
\rho^\pm \partial_t u - \mu^\pm \Delta u + \nabla \pi &= g_1, & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} u &= g_2, & \text{in } \Omega \setminus \Sigma, \\
-[\mu^\pm \partial_3(u_1, u_2)] - [\mu^\pm \nabla_{x'} u_3] &= g_3, & \text{on } \Sigma, \\
-2[\mu^\pm \partial_3 u_3] + [\pi] - \sigma \Delta_{x'} h &= g_4, & \text{on } \Sigma, \\
[u] &= g_5, & \text{on } \Sigma, \\
\partial_t h - (u_3^+ + u_3^-)/2 + [\partial_3 \mu] &= g_6, & \text{on } \Sigma, \\
(-\nabla_{x'} h, 1)^\top \cdot \nu_{S_1} &= g_7, & \text{on } \partial \Sigma, \\
\Delta \mu &= g_8, & \text{in } \Omega \setminus \Sigma, \\
\mu|_\Sigma - \sigma \Delta_{x'} h &= g_9, & \text{on } \Sigma, \\
\nu_{\partial \Omega} \cdot \nabla \mu|_{\partial \Omega} &= g_{10}, & \text{on } \partial \Omega \setminus \Sigma, \\
P_{S_1}(\mu^\pm(Du + Du^\top)\nu_{S_1}) &= P_{S_1}g_{11}, & \text{on } S_1 \setminus \partial \Sigma, \\
u \cdot \nu_{S_1} &= g_{12}, & \text{on } S_1 \setminus \partial \Sigma, \\
u &= g_{13}, & \text{on } S_2, \\
u(0) &= u_0, & \text{on } \Omega \setminus \Sigma, \\
h(0) &= h_0, & \text{on } \Sigma.
\end{aligned} \tag{4.16}$$

The main result on maximal regularity for (4.16) is the following.

Theorem 4.4. *Let $\mu^\pm, \rho^\pm, \sigma > 0$ be constant, $-\infty < L_1 < 0 < L_2 < \infty$, (p, q, r) as in Theorem 4.1 and $\Sigma \subset \mathbb{R}^2$ be a bounded, smooth domain. Let $\Omega := \Sigma \times (L_1, L_2)$, $S_1 := \partial \Sigma \times (L_1, L_2)$, and $S_2 := \Sigma \times \{L_1, L_2\}$. Let $0 < T < \infty$. The coupled linear system (4.16) then admits a unique solution $(u, \pi, [\pi], h, \mu)$ with regularity (4.2), if and only if the data satisfy the regularity and compatibility conditions (4.11), (4.12), (4.13), and (4.15). Furthermore, the solution map $[(g_j)_{j=1, \dots, 13}, u_0, h_0] \mapsto (u, \pi, [\pi], h, \mu)$ between the above spaces is continuous.*

Proof. First we reduce to trivial initial data by solving an auxiliary ninety degree angle linear Mullins-Sekerka problem of type

$$\begin{aligned}
\partial_t \bar{h} + \llbracket \partial_3 \bar{\mu} \rrbracket &= g_6, & \text{on } \Sigma, \\
(-\nabla_{x'} \bar{h}, 1)^\top \cdot \nu_{S_1} &= g_7, & \text{on } \partial\Sigma, \\
\Delta \bar{\mu} &= g_8, & \text{in } \Omega \setminus \Sigma, \\
\bar{\mu}|_\Sigma - \sigma \Delta_{x'} \bar{h} &= g_9, & \text{on } \Sigma, \\
\nu_{\partial\Omega} \cdot \nabla \bar{\mu}|_{\partial\Omega} &= g_{10}, & \text{on } \partial\Omega \setminus \Sigma, \\
\bar{h}(0) &= h_0, & \text{on } \Sigma,
\end{aligned}$$

by functions

$$\bar{h} \in W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma)), \quad \bar{\mu} \in L_p(0, T; W_q^2(\Omega \setminus \Sigma)).$$

Then we solve an auxiliary two-phase Stokes problem

$$\begin{aligned}
\rho^\pm \partial_t \bar{u} - \mu^\pm \Delta \bar{u} + \nabla \bar{\pi} &= g_1, & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} \bar{u} &= g_2, & \text{in } \Omega \setminus \Sigma, \\
-\llbracket \mu^\pm \partial_3(\bar{u}_1, \bar{u}_2) \rrbracket - \llbracket \mu^\pm \nabla_{x'} \bar{u}_3 \rrbracket &= g_3, & \text{on } \Sigma, \\
-2\llbracket \mu^\pm \partial_3 \bar{u}_3 \rrbracket + \llbracket \bar{\pi} \rrbracket &= g_4 - \sigma \Delta_{x'} \bar{h}, & \text{on } \Sigma, \\
\llbracket \bar{u} \rrbracket &= g_5, & \text{on } \Sigma, \\
P_{S_1}(\mu^\pm (D\bar{u} + D\bar{u}^\top) \nu_{S_1}) &= P_{S_1} g_{11}, & \text{on } S_1 \setminus \partial\Sigma, \\
\bar{u} \cdot \nu_{S_1} &= g_{12}, & \text{on } S_1 \setminus \partial\Sigma, \\
\bar{u} &= g_{13}, & \text{on } S_2, \\
\bar{u}(0) &= u_0, & \text{on } \Omega \setminus \Sigma,
\end{aligned} \tag{4.17}$$

using Theorem A.11 in [29] by functions

$$\bar{u} \in W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma)), \quad \bar{\pi} \in L_r(0, T; \dot{H}_r^1(\Omega \setminus \Sigma)), \tag{4.18}$$

with $\llbracket \bar{\pi} \rrbracket \in W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma))$. Here we want to point out two things: $\Delta_{x'} \bar{h}$ has sufficient regularity to be admissible data and that there is no compatibility condition stemming from $(4.17)_4$. Hence $g_4 - \sigma \Delta_{x'} \bar{h}$ is admissible

data for the problem. Having now $(\bar{u}, \bar{\pi}, \bar{h}, \bar{\mu})$ at hand, we are left to solve

$$\begin{aligned}
\rho^\pm \partial_t u - \mu^\pm \Delta u + \nabla \pi &= 0, & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} u &= 0, & \text{in } \Omega \setminus \Sigma, \\
-[\mu^\pm \partial_3(u_1, u_2)] - [\mu^\pm \nabla_{x'} u_3] &= 0, & \text{on } \Sigma, \\
-2[\mu^\pm \partial_3 u_3] + [\pi] - \sigma \Delta_{x'} h &= 0, & \text{on } \Sigma, \\
[u] &= 0, & \text{on } \Sigma, \\
\partial_t h - u_3|_\Sigma + [\partial_3 \mu] &= (\bar{u}_3^+ + \bar{u}_3^-)/2, & \text{on } \Sigma, \\
(-\nabla_{x'} h, 1)^\top \cdot \nu_{S_1} &= 0, & \text{on } \partial \Sigma, \\
\Delta \mu &= 0, & \text{in } \Omega \setminus \Sigma, \\
\mu|_\Sigma - \sigma \Delta_{x'} h &= -\sigma \Delta_{x'} \bar{h}, & \text{on } \Sigma, \\
\nu_{\partial \Omega} \cdot \nabla \mu|_{\partial \Omega} &= 0, & \text{on } \partial \Omega \setminus \Sigma, \\
P_{S_1}(\mu^\pm (Du + Du^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u &= 0, & \text{on } S_2, \\
u(0) &= 0, & \text{on } \Omega \setminus \Sigma, \\
h(0) &= 0, & \text{on } \Sigma.
\end{aligned}$$

We do this as follows. Define $L_{MS} : {}_0\mathbb{E}_{MS,T} \rightarrow {}_0\mathbb{F}_{MS,T}$ by

$$L_{MS} : (h, \mu) \mapsto \begin{pmatrix} \partial_t h - [\partial_3 \mu] \\ \Delta \mu \\ \mu|_\Sigma - \sigma \Delta_{x'} h \\ n_{\partial \Omega} \cdot \nabla \mu|_{\partial \Omega} \\ (-\nabla_{x'} h|_{\partial \Sigma}, 1)^\top \cdot \nu_{S_1} \end{pmatrix}$$

where

$${}_0\mathbb{E}_{MS,T} := [{}_0W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma))] \times L_p(0, T; W_q^2(\Omega \setminus \Sigma)),$$

and

$$\begin{aligned}
{}_0\mathbb{F}_{MS,T} &:= L_p(0, T; W_q^{1-1/q}(\Sigma)) \times L_p(0, T; L_q(\Omega)) \times L_p(0, T; W_q^{1-1/q}(\Sigma)) \\
&\quad \times L_p(0, T; W_q^{1-1/q}(\partial \Omega)) \times [{}_0F_{pq}^{1-2/(3q)}(0, T; L_q(\partial \Sigma)) \cap L_p(0, T; W_q^{3-2/q}(\partial \Sigma))].
\end{aligned}$$

In [3] we have shown that $L_{MS} : {}_0\mathbb{E}_{MS,T} \rightarrow {}_0\mathbb{F}_{MS,T}$ is boundedly invertible. Define $L_S : {}_0\mathbb{E}_{MS,T} \rightarrow [{}_0W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma))]$ by $L_S(h) := u$, where (u, π)

is the unique solution of

$$\begin{aligned}
\rho^\pm \partial_t u - \mu^\pm \Delta u + \nabla \pi &= 0, & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} u &= 0, & \text{in } \Omega \setminus \Sigma, \\
-[\mu^\pm \partial_3(u_1, u_2)] - [\mu^\pm \nabla_{x'} u_3] &= 0, & \text{on } \Sigma, \\
-2[\mu^\pm \partial_3 u_3] + [\pi] &= \sigma \Delta_{x'} h, & \text{on } \Sigma, \\
[u] &= 0, & \text{on } \Sigma, \\
P_{S_1}(\mu^\pm (Du + Du^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u &= 0, & \text{on } S_2, \\
u(0) &= 0, & \text{on } \Omega \setminus \Sigma,
\end{aligned}$$

cf. Theorem A.11 in [29]. It then stems from Theorem 4.1 that L_S is well defined, linear and bounded. Define $B : {}_0\mathbb{E}_{MS,T} \rightarrow {}_0\mathbb{F}_{MS,T}$ and $G(\bar{u}, \bar{h}) \in {}_0\mathbb{F}_{MS,T}$ by

$$B(h) := (-L_S(h)|_\Sigma, 0, 0, 0, 0)^\top, \quad G(\bar{u}, \bar{h}) := ((\bar{u}_3^+ + \bar{u}_3^-)/2, 0, -\sigma \Delta_{x'} \bar{h}, 0, 0)^\top.$$

We can hence rewrite the problem as

$$L_{MS}(h, \mu) = -B(h) + G(\bar{u}, \bar{h}), \quad \text{in } {}_0\mathbb{F}_{MS,T}.$$

We now solve this equation by a Neumann series argument. Clearly this equation is equivalent to

$$(I + L_{MS}^{-1}B)(h, \mu) = L_{MS}^{-1}G(\bar{u}, \bar{h}), \quad \text{in } {}_0\mathbb{F}_{MS,T},$$

hence it remains to show that $|L_{MS}^{-1}B|_{\mathcal{B}({}_0\mathbb{E}_{MS,T})} \leq \frac{1}{2}$, if $T > 0$ is small enough. Then by a Neumann series argument, $(I + L_{MS}^{-1}B)$ is invertible and the theorem is shown. Since we now have that L_{MS} is boundedly invertible and the norm of the inverse is independent of T - recall we only consider functions with vanishing time trace at $t = 0$ - the claim follows from Theorem 4.1. Indeed,

$$\begin{aligned}
|B(h)|_{{}_0\mathbb{F}_{MS,\tau}} &= |L_S(h)|_{L_p(0,\tau;W_q^{1-1/q}(\Sigma))} \leq \tau^{1/p} |L_S(h)|_{L_\infty(0,\tau;W_q^{1-1/q}(\Sigma))} \\
&\leq \tau^{1/p} |h|_{{}_0\mathbb{E}_{MS,\tau}}, \quad \tau > 0.
\end{aligned}$$

Note that again since h has vanishing time trace, all embeddings in Theorem 4.1 are time-independent. In particular, by choosing $\tau > 0$ sufficiently small, we get a unique solution (h, μ) in the proper regularity class on $(0, \tau)$. Solving then the two-phase Stokes system for this particular h gives a proper (u, π) in the L_r -regularity scale, again on $(0, \tau)$.

Shifting back the equations via $\tilde{u}(t) := u(t - \tau)$, $\tilde{\pi}(t) := \pi(t - \tau)$, $\tilde{h}(t) := h(t - \tau)$ and $\tilde{\mu} := \mu(t - \tau)$ we can again apply this argument and solve again on the same length time interval $(0, \tau)$, which in turn gives us now a solution on $(0, 2\tau)$ in fact. Repeating the steps we can solve then the problem on $(0, T)$, cf. Section 2.3 in [29]. \square

5. NONLINEAR WELL-POSEDNESS

In this section we show local well-posedness for the full nonlinear problem (3.1). The main result is the following.

Theorem 5.1. *Let $\mu^\pm, \rho^\pm, \sigma > 0$ be constant, $-\infty < L_1 < 0 < L_2 < \infty$, $p \in (6, \infty)$, $q \in (2p/(p+1), 2) \cap (19/10, 2)$, $3 < r < 7/2$, $\Sigma \subset \mathbb{R}^2$ be a bounded, smooth domain. Let $\Omega := \Sigma \times (L_1, L_2)$, $S_1 := \partial\Sigma \times (L_1, L_2)$ be the walls and $S_2 := \Sigma \times \{L_1, L_2\}$ bottom and top of the container. Furthermore let $(u_0, h_0) \in W_r^{2-2/r}(\Omega \setminus \Sigma) \times B_{qp}^{4-1/q-3/p}(\Sigma)$ be admissible by the compatibility conditions*

$$\begin{aligned} \operatorname{div} u_0 &= G_d(h_0, u_0), \quad \text{in } \Omega \setminus \Sigma, \\ -\llbracket \mu^\pm \partial_3(u_0)_{1,2} \rrbracket - \llbracket \mu^\pm \nabla_{x'}(u_0)_3 \rrbracket &= G_S^\parallel(u_0, h_0), \quad \llbracket u_0 \rrbracket = 0, \quad \text{on } \Sigma, \\ P_{S_1}(\mu^\pm(Du_0 + Du_0^\top)\nu_{S_1}) &= 0, \quad u_0 \cdot \nu_{S_1} = 0, \quad \text{on } S_1, \\ u_0|_{S_2} &= 0, \quad \text{on } S_2, \quad (-\nabla_{x'} h_0, 1)^\top \cdot \nu_{S_1} = 0, \quad \text{on } \partial\Sigma. \end{aligned} \tag{5.1}$$

Then the full nonlinear (transformed) problem (3.1) admits a unique local-in-time strong solution, that is, there is some $T_0 > 0$, such that for every $0 < T \leq T_0$ there is some $\varepsilon = \varepsilon(T) > 0$, such that whenever the smallness condition

$$|u_0|_{W_r^{2-2/r}(\Omega \setminus \Sigma)} + |h_0|_{B_{qp}^{4-1/q-3/p}(\Sigma)} \leq \varepsilon \tag{5.2}$$

is satisfied there is a unique strong solution $(u, \pi, \llbracket \pi \rrbracket, h, \mu)$ of (3.1) on $(0, T)$ with regularity (4.2).

Proof. We first again reduce the problem to $(u_0, h_0) = 0$. This can be done by solving an auxiliary problem first by functions (u_*, π_*, h_*, μ_*) in the proper regularity classes, cf. Section 3.2 in [29].

Let us now introduce notation. Let

$$\begin{aligned} {}_0\mathbb{E}_u(T) &:= {}_0W_r^1(0, T; L_r(\Omega)) \cap L_r(0, T; W_r^2(\Omega \setminus \Sigma)), \quad \mathbb{E}_\pi(T) := L_r(0, T; \dot{H}_r^1(\Omega \setminus \Sigma)), \\ {}_0\mathbb{E}_q(T) &:= {}_0W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma)), \\ {}_0\mathbb{E}_h(T) &:= {}_0W_p^1(0, T; W_q^{1-1/q}(\Sigma)) \cap L_p(0, T; W_q^{4-1/q}(\Sigma)), \end{aligned}$$

and $\mathbb{E}_\mu(T) := L_p(0, T; W_q^2(\Omega \setminus \Sigma))$. Furthermore, let

$${}_0\mathbb{E}(T) := {}_0\mathbb{E}_u(T) \times \mathbb{E}_\pi(T) \times {}_0\mathbb{E}_q(T) \times {}_0\mathbb{E}_h(T) \times \mathbb{E}_\mu(T) \cap \{(u, \pi, q, h, \mu) : q = \llbracket \pi \rrbracket\}.$$

Moreover, let

$$\begin{aligned}
\mathbb{F}_1(T) &:= L_r(0, T; L_r(\Omega)), \quad \mathbb{F}_2(T) := L_r(0, T; W_r^1(\Omega \setminus \Sigma)), \\
\mathbb{F}_3(T) &:= {}_0W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma)), \\
\mathbb{F}_4(T) &:= {}_0W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma)), \\
\mathbb{F}_5(T) &:= {}_0W_r^{1-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{2-1/r}(\Sigma)), \\
\mathbb{F}_6(T) &:= L_p(0, T; W_q^{1-1/q}(\Sigma)), \\
\mathbb{F}_7(T) &:= {}_0F_{pq}^{1-2/(3q)}(0, T; L_q(\partial\Sigma)) \cap L_p(0, T; W_q^{3-2/q}(\partial\Sigma)), \\
\mathbb{F}_8(T) &:= L_p(0, T; L_q(\Omega)), \quad \mathbb{F}_9(T) := L_p(0, T; W_q^{2-1/q}(\Sigma)), \\
\mathbb{F}_{10}(T) &:= L_p(0, T; W_q^{1-1/q}(\partial\Omega)), \\
\mathbb{F}_{11}(T) &:= {}_0W_r^{1/2-1/(2r)}(0, T; L_r(S_1)) \cap L_r(0, T; W_r^{1-1/r}(S_1)), \\
\mathbb{F}_{12}(T) &:= {}_0W_r^{1-1/(2r)}(0, T; L_r(S_1)) \cap L_r(0, T; W_r^{2-1/r}(S_1)), \\
\mathbb{F}_{13}(T) &:= {}_0W_r^{1-1/(2r)}(0, T; L_r(S_2)) \cap L_r(0, T; W_r^{2-1/r}(S_2)).
\end{aligned}$$

Let

$${}_0\mathbb{F}(T) := \times_{j=1}^{13} \mathbb{F}_j(T) \cap \{(g_2, g_5, g_{12}, g_{13}) \in W_r^1(\mathbb{R}_+; \hat{H}_r^{-1}(\Omega))\}. \quad (5.3)$$

Define a linear operator by the left hand side of (3.1), that is, define $\mathbb{L} : {}_0\mathbb{E}(T) \rightarrow {}_0\mathbb{F}(T)$ via

$$\mathbb{L}(u, \pi, q, h, \mu) := \begin{pmatrix} \rho^\pm \partial_t u - \mu^\pm \Delta u + \nabla \pi \\ \operatorname{div} u \\ -\llbracket \mu^\pm \partial_3(u_1, u_2) \rrbracket - \llbracket \mu^\pm \nabla_{x'} u_3 \rrbracket \\ -2\llbracket \mu^\pm \partial_3 u_3 \rrbracket + q - \sigma \Delta_{x'} h \\ \llbracket u \rrbracket \\ \partial_t h - u_3|_\Sigma + \llbracket \partial_3 \mu \rrbracket \\ (-\nabla_{x'} h, 1)^\top|_{\partial\Sigma} \cdot \nu_{S_1} \\ \Delta \mu \\ \mu|_\Sigma - \sigma \Delta_{x'} h \\ \nu_{\partial\Omega} \cdot \nabla \mu|_{\partial\Omega} \\ P_{S_1}(\mu^\pm (Du + Du^\top) \nu_{S_1}) \\ u|_{S_1} \cdot \nu_{S_1} \\ u|_{S_2} \end{pmatrix}.$$

We collect the right hand side in the operator $R : \mathbb{E}(T) \rightarrow \mathbb{F}(T)$ defined by

$$R(u, \pi, q, h, \mu) := \begin{pmatrix} a^\pm(h; D_x^2)(u, \pi) + \bar{a}(h, u) \\ G_d(u, h) \\ G_S(u, \pi, h)_{1,2} \\ G_S(u, \pi, h)_3 \\ 0 \\ G_\Sigma(u, h, \mu) \\ 0 \\ G_c(h, \mu) \\ G_\kappa(h) \\ G_N(h, \mu) \\ G_P^\pm(u, h) \\ 0 \\ 0 \end{pmatrix}.$$

Hereby $\mathbb{E}(T)$ and $\mathbb{F}(T)$ are defined similarly but without the trace properties at $t = 0$.

It is now clear that for $h \in {}_0\mathbb{E}(T)$ (which is a function having vanishing time trace) the compatibility condition $(-\nabla_x h(t=0), 1)^\top|_{\partial\Sigma} \cdot \nu_{S_1} = 0$ is satisfied. Regarding the compatibility conditions for the Stokes system we refer to Section 3.1 in [29]. Therefore both operators are well defined.

Let $z := (u, \pi, q, h, \mu)$ and $z_* := (u_*, \pi_*, \llbracket \pi_* \rrbracket, h_*, \mu_*)$ the reference solution as above. We can now rewrite the problem abstractly as

$$L(z + z_*) = R(z + z_*), \quad z \in {}_0\mathbb{E}(T).$$

Note that we already know that L is invertible from ${}_0\mathbb{E}(T)$ to ${}_0\mathbb{F}(T)$ and the norms are independent of T . This renders the fixed point equation

$$z = L^{-1}(R(z + z_*) - Lz_*), \quad \text{in } {}_0\mathbb{E}(T).$$

Define $K : {}_0\mathbb{E}(T) \rightarrow {}_0\mathbb{E}(T)$ by means of $[z \mapsto L^{-1}(R(z + z_*) - Lz_*)]$. We now need to establish contraction estimates for R .

Lemma 5.2. *We have*

$$\begin{aligned} & |R(z_1 + z_*) - R(z_2 + z_*)|_{{}_0\mathbb{F}(T)} \\ & \leq C(T^\alpha + |z_*|_{\mathbb{E}(T)} + |z_1|_{{}_0\mathbb{E}(T)} + |z_2|_{{}_0\mathbb{E}(T)}) |z_1 - z_2|_{{}_0\mathbb{E}(T)}, \end{aligned} \quad (5.4)$$

for some $\alpha > 0$ and for all $z_1, z_2 \in \mathbb{B}(r, 0) \subset {}_0\mathbb{E}(T)$, if $r > 0$ and $T = T(r) > 0$ are sufficiently small.

Having these estimates at hand we proceed as in the proof of Theorem 5.1 in [3] to obtain a fixed point of K by Banach's contraction mapping principle by choosing $\varepsilon(T) > 0$ in (5.2) small enough. This finishes the proof. \square

Proof of Lemma 5.2. Let us first note that

$$[h \mapsto \Delta_h] \in C^1(U; \mathcal{B}(W_r^2(\Omega \setminus \Sigma); L_r(\Omega))), \quad (5.5)$$

$$[h \mapsto \nabla_h] \in C^1(U; \mathcal{B}(W_r^k(\Omega \setminus \Sigma); W_r^{k-1}(\Omega \setminus \Sigma))), \quad k = 1, 2, \quad (5.6)$$

where $U \subset B_{qp}^{4-1/q-3/p}(\Sigma)$ is a sufficiently small neighbourhood of zero. This can be shown as in Lemma 3.4 in [3].

We estimate every nonlinearity separately. We recall that $a^\pm(h; D_x)(u, \pi) = \mu^\pm(\Delta_h - \Delta)u - (\nabla - \nabla_h)\pi$. Clearly,

$$|(\Delta_h - \Delta)u|_{L_r(0,T;L_r(\Omega))} \leq |\Delta_h - \Delta|_{L_\infty(0,T;\mathcal{B}(W_r^2(\Omega \setminus \Sigma); L_r(\Omega)))} |u|_{L_r(0,T;W_r^2(\Omega \setminus \Sigma))}.$$

Using (5.5) this gives

$$|(\Delta_h - \Delta)u|_{L_r(0,T;L_r(\Omega))} \leq C|h|_{0\mathbb{E}(T)} |u|_{L_r(0,T;W_r^2(\Omega \setminus \Sigma))}.$$

The same arguments give

$$|(\nabla_h - \nabla)\pi|_{L_r(0,T;L_r(\Omega))} \leq C|h|_{0\mathbb{E}(T)} |\pi|_{L_r(0,T;\dot{H}_r^1(\Omega \setminus \Sigma))},$$

since (5.6) is also true for the homogeneous counterparts \dot{H}_r^k replacing W_r^k . Note that these estimates and the C^1 -dependence of h and the bilinear structure in (u, π) of $a^\pm(h)(u, \pi)$ then automatically give rise to a contraction estimate of form

$$\begin{aligned} & |a^\pm(h_1)(u_1, \pi_1) - a^\pm(h_2)(u_2, \pi_2)|_{L_r(0,T;L_r(\Omega))} \leq \\ & \leq C|h_1 - h_2|_{0\mathbb{E}(T)} \left(|u_1 - u_2|_{L_r(0,T;W_r^2(\Omega \setminus \Sigma))} + |\pi_1 - \pi_2|_{L_r(0,T;\dot{H}_r^1(\Omega \setminus \Sigma))} \right), \end{aligned} \quad (5.7)$$

valid for all $h_1, h_2 \in U, u_1, u_2 \in L_r(0, T; W_r^2(\Omega \setminus \Sigma)), \pi_1, \pi_2 \in L_r(0, T; \dot{H}_r^1(\Omega \setminus \Sigma))$. Indeed, we have that $a^\pm \in C^2(\mathbb{E}_h(T) \times \mathbb{E}_u(T) \times \mathbb{E}_\pi(T); L_r(0, T; L_r(\Omega)))$ and $a^\pm(0) = 0, Da^\pm(0) = 0$. Alternatively, one can explicitly estimate the difference and end up with (5.7). Before we estimate $\bar{a}(u, h, \eta) := Du \cdot \partial_t \Theta_h^{-1} + (u \cdot \nabla_h)u + (\rho^+ - \rho^-)(\nabla_h \eta \cdot \nabla_h)u$, some remarks are in order. Firstly, $Du \cdot \partial_t \Theta_h^{-1} = -\chi \partial_t h (1 + h\chi')^{-1} \partial_3 u$, see [24].

Contracting the (transformed) convection term $(u \cdot \nabla_h)u$ is easy due to the fact that $\mathbb{E}_u(T) \hookrightarrow L_\infty(0, T; L_\infty(\Omega)) \cap L_\infty(0, T; W_r^1(\Omega \setminus \Sigma))$. More precisely,

$$\begin{aligned} & |(u \cdot \nabla_h)u|_{L_r(0,T;L_r(\Omega))} \leq \\ & T^{1/r} |u|_{L_\infty(0,T;L_\infty(\Omega))} |\nabla_h|_{L_\infty(0,T;\mathcal{B}(W_r^1(\Omega \setminus \Sigma); L_r(\Omega)))} |u|_{L_\infty(0,T;W_r^1(\Omega \setminus \Sigma))}. \end{aligned}$$

Regarding the other terms we recall that $Du \in L_{2r}(0, T; L_\infty(\Omega))$ due to $r > 3$. We then get by Hölder inequality that

$$\begin{aligned} & |Du \cdot \partial_t \Theta_h^{-1}|_{L_r(0,T;L_r(\Omega))} \leq \\ & |\chi(1 + h\chi')^{-1}|_{L_\infty(0,T;L_\infty(\Omega))} |Du|_{L_{p_1}(0,T;L_\infty(\Omega))} |\partial_t h|_{L_p(0,T;L_r(\Sigma))} |1|_{L_{p_0}(0,T;L_\infty(\Omega))}, \end{aligned}$$

where $1 < p_0, p_1 < \infty$ are such that

$$\frac{1}{r} = \frac{1}{p_1} + \frac{1}{p} + \frac{1}{p_0}. \quad (5.8)$$

By choice of $q < 2$ and $3 < r < 7/2$, Sobolev's embedding theorem gives

$$W_q^{1-1/q}(\Sigma) \hookrightarrow L_r(\Sigma).$$

Choosing $p_1 = 2r$ and recalling $p > 2r$ gives that there is some $1 < p_0 < \infty$ such that (5.8) is fulfilled. Note that these estimates are not optimal but sufficient in our case. We then obtain that there is some $\varepsilon = \varepsilon(p, q, r) > 0$ such that

$$|Du \cdot \partial_t \Theta_h^{-1}|_{L_r(0,T;L_r(\Omega))} \leq CT^\varepsilon |u|_{\mathbb{E}_u(T)} |h|_{\mathbb{E}_h(T)}.$$

Furthermore,

$$|(\nabla_h \eta \cdot \nabla_h)u|_{L_r(0,T;L_r(\Omega))} \leq |\nabla_h \eta|_{L_p(0,T;L_r(\Omega))} |\nabla_h u|_{L_{p_1}(0,T;L_\infty(\Omega))} |1|_{L_{p_0}(0,T;L_\infty(\Omega))},$$

where p_0, p_1 are as above. Again by Sobolev embedding, $W_q^1(\Omega \setminus \Sigma) \hookrightarrow L_r(\Omega)$, whence

$$|(\nabla_h \eta \cdot \nabla_h)u|_{L_r(0,T;L_r(\Omega))} \leq |\nabla_h \eta|_{L_p(0,T;W_q^1(\Omega \setminus \Sigma))} |\nabla_h u|_{L_{p_1}(0,T;L_\infty(\Omega))} \|1\|_{L_{p_0}(0,T;L_\infty(\Omega))}.$$

In view of (5.6), these estimates together with the smooth dependence

$$\bar{a} \in C^\infty(\mathbb{E}_u(T) \times \mathbb{E}_h(T); \mathbb{F}_1(T))$$

as well as $\bar{a}(0, 0, 0) = 0$ and $D\bar{a}(0, 0, 0) = 0$ give rise to contraction estimates for \bar{a} .

For $G_d(u, h) := (\operatorname{div} - \operatorname{div}_h)u$, the estimate in $\mathbb{F}_2(T)$ is straightforward,

$$|G_d(u, h)|_{L_r(0,T;W_r^1(\Omega \setminus \Sigma))} \leq |\nabla - \nabla_h|_{L_\infty(0,T;\mathcal{B}(W_r^1(\Omega \setminus \Sigma); L_r(\Omega)))} \|u\|_{L_r(0,T;W_r^2(\Omega \setminus \Sigma))},$$

where we used that $G_d(u, h) = \operatorname{Tr}(\nabla - \nabla_h)u$.

The contraction estimates for $G_c(h, \eta) := (\Delta - \Delta_h)\eta = (\operatorname{div} \nabla - \operatorname{div}_h \nabla_h)\eta$ and $G_N(h, \eta) := \nu_{\partial\Omega} \cdot (\nabla - \nabla_h)\eta$ easily stem from (5.5)-(5.6) with q replacing r , see also [3]. Note that there the contraction estimates for $G_\kappa(h) := \sigma(K(h) - \Delta_{x'}h)$ are already proven in a far more general setting. In this graph situation case we can give a much easier proof. Recall that in this case $K(h) = \operatorname{div}_{x'}(\nabla_{x'}h(1 + |\nabla_{x'}h|^2)^{-1/2})$, whence

$$G_\kappa(h) = \left(1 - \frac{1}{\sqrt{1 + |\nabla_{x'}h|^2}}\right) \Delta_{x'}h + \nabla_{x'}h \cdot \nabla_{x'} \left(\frac{1}{\sqrt{1 + |\nabla_{x'}h|^2}} \right). \quad (5.9)$$

Again using the product estimate

$$|\nabla h \cdot \nabla^2 h|_{\mathbb{F}_9(T)} \leq C |\nabla h|_{L_\infty(0,T;B_{qp}^{3-1/q-3/p}(\Sigma))} |\nabla^2 h|_{\mathbb{F}_9(T)} \leq C \|h\|_{\mathbb{E}_h(T)}^2$$

and the fact that $G_\kappa \in C^\infty({}_0\mathbb{E}_h(T); \mathbb{F}_9(T))$, $G_\kappa(0) = 0$, $DG_\kappa(0) = 0$ ensure the contraction property of G_κ .

Regarding $G_P^\pm(u, h)$ it is shown in Section 3.1 in [29], that

$$\begin{aligned} G_P^\pm(u, h) &= P_{S_1} \left[\frac{1}{1 + \chi'h} \left(\chi \partial_3 u ((-\nabla_{x'}h, 1)^\top \cdot \nu_{S_1}) + \left(\chi \frac{\nabla_{x'}h}{\chi'h} \right) \partial_3(u \cdot \nu_{S_1}) \right) \right]. \end{aligned} \quad (5.10)$$

Therefore, due to the fact that $u \cdot \nu_{S_1} = 0$ on $S_1 \setminus \partial\Sigma$ and $(-\nabla_{x'}h, 1)^\top \cdot \nu_{S_1} = 0$ on $\partial\Sigma \times (L_1, L_2)$, the nonlinearity $G_P^\pm(u, h)$ vanishes for the solution. Hence we may replace $G_P^\pm(u, h)$ by zero in the definition of \mathbf{R} .

Now, for $G_S(h, u, \pi)$ we split $G_S(h, u, \pi) = G_S^S(h, u, \pi) + G_S^\kappa(h)$, where

$$\begin{aligned} G_S^S(h, u, \pi) &:= \llbracket \mu^\pm ((D\Theta_h - I)Du + Du^\top(D\Theta_h - I)^\top) \rrbracket \nu_{\Sigma_h} + \\ &\quad + \llbracket (\mu^\pm(Du + Du^\top) - \pi I) (e_3 - \nu_{\Sigma_h}) \rrbracket, \\ G_S^\kappa(h) &:= \sigma(K(h)\nu_{\Sigma_h} - \Delta_{x'}he_3). \end{aligned}$$

Regarding the estimates of $G_S^S(h, u, \pi)$ we refer to [23]. Note that due to Remark 1.2. (c) in [23] we may use these results since $r > 3$ and $\mathbb{E}_h(T) \hookrightarrow BUC([0, T]; C^2(\Sigma))$.

Considering $G_S^\kappa(h)$ we may write $G_S^\kappa(h) = G_\kappa(h)e_3 + K(h)(\nu_{\Sigma_h} - e_3)$ and estimate each term separately. In particular, we have to control terms of the form $\nabla h \cdot \nabla^2 h$ in the norm of $\mathbb{F}_3(T) = {}_0W_r^{1/2-1/(2r)}(0, T; L_r(\Sigma)) \cap L_r(0, T; W_r^{1-1/r}(\Sigma))$. This stems from the observation in (5.9). Now, by Theorem 4.1 we already know that the space

in which second derivatives of h live in embeds into $\mathbb{F}_3(T)$. We may now use the product estimate of Proposition 5.7 in [19] to obtain

$$\begin{aligned} |\nabla h \cdot \nabla^2 h|_{W_r^{1/2-1/(2r)}(0,T;L_r(\Sigma))} &\lesssim |\nabla h|_{L_\infty(0,T;L_\infty(\Sigma))} |\nabla^2 h|_{W_r^{1/2-1/(2r)}(0,T;L_r(\Sigma))} + \\ &\quad + |\nabla h|_{W_r^{1/2-1/(2r)}(0,T;L_\infty(\Sigma))} |\nabla^2 h|_{L_\infty(0,T;L_r(\Sigma))}. \end{aligned}$$

Furthermore,

$$|\nabla h \cdot \nabla^2 h|_{L_r(0,T;W_r^{1-1/r}(\Sigma))} \lesssim |\nabla h|_{L_\infty(0,T;C^1(\Sigma))} |\nabla^2 h|_{L_r(0,T;W_r^{1-1/r}(\Sigma))}.$$

These estimates show that the product terms of form $\nabla h \cdot \nabla^2 h$ are well defined in $\mathbb{F}_3(T)$.

These observations allow us to conclude contraction estimates for G_S^κ since again $G_S^\kappa(0) = 0, DG_S^\kappa(0) = 0$.

Regarding $G_\Sigma(u, h, \mu) = u|_\Sigma \cdot (-\nabla_{x'} h, 0)^\top - \llbracket e_3 \cdot (\nabla - \nabla_h) \mu \rrbracket - \llbracket (-\nabla_{x'} h, 0)^\top \cdot \nabla_h \mu \rrbracket$, the last two terms can be controlled as before. Clearly the first term is smooth in (u, h) and quadratic and the bound

$$|u|_\Sigma \cdot (-\nabla_{x'} h, 0)^\top|_{L_p(0,T;W_q^{1-1/q}(\Sigma))} \leq T^{1/p} |u|_{L_\infty(0,T;W_q^1(\Omega \setminus \Sigma))} |\nabla h|_{L_\infty(0,T;C^1(\Sigma))}$$

renders contraction estimates also for G_Σ . This concludes the proof of the contraction estimates. \square

6. QUALITATIVE BEHAVIOUR

In this section we investigate the long-time behaviour of solutions starting close to equilibria. By a study of the spectrum of the linearization we will show that solutions starting close to certain equilibria converge to an equilibrium solution at an exponential rate.

Let us again consider the case of a cylindrical container $\Omega = \Sigma \times (L_1, L_2)$, where $-\infty < L_1 < 0 < L_2 < \infty$ and $\Sigma \subset \mathbb{R}^2$ is open, bounded and has smooth boundary. We want to study stability properties of

$$\begin{aligned} \rho \partial_t u - \mu \Delta u + \operatorname{div}[(\rho u + \llbracket \rho \rrbracket \nabla \eta) \otimes u] + \nabla p &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ \operatorname{div} u &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ -\llbracket \mu(Du + Du^\top) \rrbracket \nu_{\Gamma(t)} + \llbracket p \rrbracket \nu_{\Gamma(t)} &= \sigma H_{\Gamma(t)} \nu_{\Gamma(t)}, & \text{on } \Gamma(t), \\ \llbracket u \rrbracket &= 0, & \text{on } \Gamma(t), \\ V_{\Gamma(t)} - u|_{\Gamma(t)} \cdot \nu_{\Gamma(t)} &= -\llbracket \nu_{\Gamma(t)} \cdot \nabla \eta \rrbracket, & \text{on } \Gamma(t), \\ \nu_{\Gamma(t)} \cdot \nu_{S_1} &= 0, & \text{on } \partial \Gamma(t), \\ \Delta \eta &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ \eta|_{\Gamma(t)} &= \sigma H_{\Gamma(t)}, & \text{on } \Gamma(t), \\ \nu_{\partial \Omega} \cdot \nabla \eta|_{\partial \Omega} &= 0, & \text{on } \partial \Omega \setminus \Gamma(t), \\ P_{S_1}(\mu(Du + Du^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Gamma(t), \\ u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Gamma(t), \\ u &= 0, & \text{on } S_2, \\ u(0) &= u_0, & \text{on } \Omega \setminus \Gamma(0), \\ \Gamma(0) &= \Gamma_0. \end{aligned} \tag{6.1}$$

We recall that $\rho := \rho^+ \chi_{\Omega^+(t)} + \rho^- \chi_{\Omega^-(t)}$ and $\mu := \mu^+ \chi_{\Omega^+(t)} + \mu^- \chi_{\Omega^-(t)}$.

6.1. Equilibria and spectrum of the linearization. We note that the pressure p as well as the chemical potential μ may be reconstructed by the semiflow $(u(t), \Gamma(t))$ as follows. For given $\Gamma(t)$ we can solve the two-phase elliptic problem

$$\begin{aligned} \Delta \eta &= 0, & \text{in } \Omega \setminus \Gamma(t), \\ \eta|_{\Gamma(t)} &= \sigma H_{\Gamma(t)}, & \text{on } \Gamma(t), \\ n_{\partial\Omega} \cdot \nabla \eta|_{\partial\Omega} &= 0, & \text{on } \partial\Omega, \end{aligned}$$

and the weak transmission problem

$$\begin{aligned} (\nabla p / \rho | \nabla \varphi)_{L_2(\Omega)} &= ([\mu / \rho] \Delta u - u \cdot \nabla u | \nabla \varphi)_{L_2(\Omega)}, & \text{for all } \varphi \in W_{r'}^1(\Omega), \\ \llbracket p \rrbracket &= \llbracket \mu (Du + Du^\top) \nu_{\Gamma(t)} \cdot \nu_{\Gamma(t)} \rrbracket + \sigma H_{\Gamma(t)}, & \text{on } \Gamma(t), \end{aligned}$$

where $r' = r/(r-1)$, cf. Lemma A.7 in [29]. Therefore we may concentrate on the set of equilibria \mathcal{E} for the semiflow $(u(t), \Gamma(t))$. Note that the set of equilibria for (6.1) is given by $\mathcal{E} = \{(u, \Gamma) : u = 0, H_\Gamma = \text{const.}\}$. In particular, also μ is constant, p is constant in the two phases of $\Omega \setminus \Gamma$ and also the jump $\llbracket p \rrbracket$ is constant on Γ .

Remark 6.1. *We want to point out that in the special case when Γ is a C^2 -graph of a function h over Σ , we can even deduce that $H_\Gamma = 0$ and h is constant. A proof of this can be found in [3].*

We again now work in the graph situation, that is, we assume the free interface $\Gamma(t)$ is a graph of a height function h over Σ .

The linearization of the transformed Two-phase Navier-Stokes/Mullins-Sekerka problem (6.1) around the trivial equilibrium $(0, \Sigma) \in \mathcal{E}$ induces us to study the problem

$$\begin{aligned} \rho \partial_t u - \mu \Delta u + \nabla p &= f_u, & \text{in } \Omega \setminus \Sigma, \\ \operatorname{div} u &= 0, & \text{in } \Omega \setminus \Sigma, \\ -\llbracket \mu (Du + Du^\top) \rrbracket e_3 + \llbracket p \rrbracket e_3 + \sigma \Delta_{x'} h e_3 &= 0, & \text{on } \Sigma, \\ \llbracket u \rrbracket &= 0, & \text{on } \Sigma, \\ \partial_t h - u_3 + \llbracket \partial_3 \eta \rrbracket &= f_h, & \text{on } \Sigma, \\ (\nabla_{x'} h, -1)^\top \cdot \nu_{S_1} &= 0 & \text{on } \partial\Sigma, \\ \Delta \eta &= 0, & \text{in } \Omega \setminus \Sigma, \\ \eta|_\Sigma + \sigma \Delta_{x'} h &= 0, & \text{on } \Sigma, \\ \nu_{\partial\Omega} \cdot \nabla \mu|_{\partial\Omega} &= 0, & \text{on } \partial\Omega \setminus \Sigma, \\ P_{S_1} (\mu (Du + Du^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial\Sigma, \\ u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial\Sigma, \\ u &= 0, & \text{on } S_2, \\ u(0) &= u_0, & \text{in } \Omega \setminus \Sigma, \\ h(0) &= h_0, & \text{on } \Sigma, \end{aligned} \tag{6.2}$$

where f_h is assumed to be mean value free. Let us note the following observations. Integrating equation (6.2)₅ over Σ yields $\int_\Sigma h(t) dx = \int_\Sigma h_0 dx$ for all $t \in \mathbb{R}_+$. In

other words, whenever h_0 and f_h are mean value free, the solution h will stay mean value free for all times. Furthermore, applying $P_\Sigma = I - e_3 \otimes e_3$ to equation (6.2)₃ directly yields that $P_\Sigma(\llbracket \mu^\pm(Du + Du^\top) \rrbracket e_3) = 0$ on Σ .

We want to write system (6.2) as an abstract evolution equation. To this end let

$$X_0 := L_{r,\sigma}(\Omega) \times W_q^{1-1/q}(\Sigma), \quad X_1 := (L_{r,\sigma}(\Omega) \cap W_r^2(\Omega \setminus \Sigma)) \times W_q^{4-1/q}(\Sigma),$$

and define a linear operator $A : D(A) \subset X_1 \rightarrow X_0$ by

$$A(u, h) := (-[\mu/\rho]\Delta u + \nabla p, -u_3 + \llbracket \partial_3 \eta \rrbracket)$$

with domain

$$\begin{aligned} D(A) := \{ (u, h) \in X_1 : & \llbracket u \rrbracket = 0 \text{ on } \Sigma, \ P_{S_1}(\mu^\pm(Du + Du^\top)\nu_{S_1}) = 0 \text{ on } S_1 \setminus \partial\Sigma, \\ & u \cdot \nu_{S_1} = 0 \text{ on } S_1 \setminus \partial\Sigma, \ u = 0 \text{ on } S_2, \\ & P_\Sigma(\llbracket \mu^\pm(Du + Du^\top) \rrbracket e_3) = 0 \text{ on } \Sigma, \ (\nabla_{x'} h, -1)^\top \cdot \nu_{S_1} = 0 \text{ on } S_1 \}. \end{aligned}$$

Here, $p \in \dot{H}_r^1(\Omega \setminus \Sigma)$ solves the weak transmission problem

$$\begin{aligned} (\nabla p / \rho | \nabla \varphi)_{L_2(\Omega)} &= ([\mu/\rho]\Delta u | \nabla \varphi)_{L_2(\Omega)}, & \text{for all } \varphi \in W_r^1(\Omega), \\ \llbracket p \rrbracket &= \sigma \Delta_{x'} h + (\llbracket \mu^\pm(Du + Du^\top) \rrbracket e_3 | e_3)_{L_2(\Sigma)}, & \text{on } \Sigma, \end{aligned}$$

cf. Lemma A.7 in [29] and $\eta \in W_q^2(\Omega \setminus \Sigma)$ solves the elliptic problem

$$\begin{aligned} \Delta \eta &= 0, & \text{in } \Omega \setminus \Sigma, \\ \eta|_\Sigma + \sigma \Delta_{x'} h &= 0, & \text{on } \Sigma, \\ \partial_\nu \eta &= 0, & \text{on } \partial\Omega. \end{aligned}$$

As in [4] and [29], we will sometimes make use of the notation via solution operators, that is,

$$\nabla p / \rho = T_1[(\mu/\rho)\Delta u] + T_2[\sigma \Delta_{x'} h + (\llbracket \mu(Du + Du^\top) \rrbracket e_3 | e_3)_{L_2(\Sigma)}],$$

cf. Lemma A.7 in [29]. Note that $\Delta_{x'} h \in W_q^{2-1/q}(\Sigma) \hookrightarrow W_r^{1-1/r}(\Sigma)$ for $h \in W_q^{4-1/q}(\Sigma)$ by Sobolev embedding, since $r < 3q/(3-q)$.

We can then rewrite problem (6.2) in a more compact form as

$$\dot{z}(t) + Az(t) = f(t), \quad t \in \mathbb{R}_+, \quad z(0) = z_0, \quad (6.4)$$

where $z := (u, h)$, $f := (f_u, f_h)$ and $z_0 := (u_0, h_0)$. We can now show a similar result as in [4] about properties of the operator A .

Lemma 6.2. *Let $n = 2, 3$, (p, q, r) as in Theorem 4.1, $\rho^\pm, \mu^\pm, \sigma > 0$ constant and X_0 and A as above. Then the following statements are true.*

- (1) *The linear operator $-A$ generates an analytic C_0 -semigroup e^{-At} in X_0 .*
- (2) *The spectrum $\sigma(-A)$ consists of countably many eigenvalues with finite algebraic multiplicity.*
- (3) *$\lambda = 0$ is a semi-simple eigenvalue with multiplicity 1 and $X_0 = N(A) \oplus R(A)$.*
- (4) *$\sigma(-A) \setminus \{0\} \subset \mathbb{C}_- := \{z \in \mathbb{C} : \operatorname{Re} z < 0\}$.*
- (5) *The kernel $N(A)$ is isomorphic to the tangent space $T_{(0, \Sigma)} \mathcal{E}$ of \mathcal{E} at the trivial equilibrium $(0, \Sigma) \in \mathcal{E}$ and is given by $N(A) = \{(u, h) : u = 0, h = \text{const.}\}$.*
- (6) *The restriction of e^{-At} to $R(A)$ is exponentially stable.*

Proof. The first assertion follows from Theorem 4.4 and the proof of Proposition 1.2 in [21]. Since $D(A)$ compactly embeds into X_0 , the resolvent of A is compact and therefore the spectrum of A consists only of countably many eigenvalues with finite multiplicity. By classical results, it does not depend on q and r , cf. [7], [12]. So let $\lambda \in \sigma(-A)$ be an eigenvalue with eigenfunctions $(u, h) \in D(A)$. The corresponding eigenvalue problem reads as

$$\begin{aligned}
\lambda \rho u - \mu \Delta u + \nabla p &= 0, & \text{in } \Omega \setminus \Sigma, \\
\Delta \eta &= 0, \quad \operatorname{div} u = 0, & \text{in } \Omega \setminus \Sigma, \\
-\llbracket \mu(Du + Du^\top) \rrbracket e_3 + \llbracket p \rrbracket e_3 - \sigma \Delta_{x'} h e_3 &= 0, & \text{on } \Sigma, \\
\lambda h - u_3 + \llbracket \partial_3 \eta \rrbracket &= 0, & \text{on } \Sigma, \\
(\nabla_{x'} h, -0)^\top \cdot \nu_{S_1} &= 0 & \text{on } \partial \Sigma, \\
\llbracket u \rrbracket = 0, \quad \llbracket \eta \rrbracket = 0, \quad \eta|_\Sigma + \sigma \Delta_{x'} h &= 0, & \text{on } \Sigma, \\
\nu_{\partial \Omega} \cdot \nabla \mu|_{\partial \Omega} &= 0, & \text{on } \partial \Omega \setminus \Sigma, \\
u \cdot \nu_{S_1} = 0, \quad P_{S_1}(\mu(Du + Du^\top)\nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u &= 0, & \text{on } S_2.
\end{aligned} \tag{6.5}$$

Testing equation (6.5)₁ with u in $L_2(\Omega)$ and invoking boundary and transmission conditions yields

$$\lambda |\rho|^{1/2} u|_{L_2(\Omega)} + |\mu|^{1/2} (Du + Du^\top)|_{L_2(\Omega)} + \sigma \bar{\lambda} |\nabla_{x'} h|_{L_2(\Sigma)}^2 + |\nabla \eta|_{L_2(\Omega)} = 0. \tag{6.6}$$

Let $\lambda = 0$. Then $u = 0$ by Korn's inequality and $\eta = \text{const.}$, whence $\Delta_{x'} h$ is constant on Σ . An integration over Σ together with the boundary condition (6.5)₆ yields that $\Delta_{x'} h = 0$ on Σ . Hence h has to be constant. We obtain that the kernel $N(A)$ is one-dimensional and $N(A) = \{(u, h) : u = 0, h = \text{const.}\}$. Taking real parts in (6.6) yields $\operatorname{Re} \lambda \leq 0$. We also easily obtain that $\sigma(-A) \cap i\mathbb{R} = \{0\}$, hence $\sigma(-A) \setminus \{0\} \subset \mathbb{C}_-$. Next we show that the eigenvalue $\lambda = 0$ is semi-simple. Pick $z = (u, h) \in N(A^2)$. Then $z_1 := Az \in N(A)$, hence $z_1 = (0, h_1)$ and h_1 is constant. The problem for $z = (u, h)$ now reads as

$$\begin{aligned}
-\mu \Delta u + \nabla p &= 0, & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} u &= 0, & \text{in } \Omega \setminus \Sigma, \\
-\llbracket \mu(Du + Du^\top) \rrbracket e_3 + \llbracket p \rrbracket e_3 - \sigma \Delta_{x'} h e_3 &= 0, & \text{on } \Sigma, \\
\llbracket u \rrbracket &= 0, & \text{on } \Sigma, \\
-u_3 + \llbracket \partial_3 \eta \rrbracket &= h_1, & \text{on } \Sigma, \\
(\nabla_{x'} h, -1)^\top \cdot \nu_{S_1} &= 0 & \text{on } \partial \Sigma, \\
\Delta \eta &= 0, & \text{in } \Omega \setminus \Sigma, \\
\eta|_\Sigma + \sigma \Delta_{x'} h &= 0, & \text{on } \Sigma, \\
\nu_{\partial \Omega} \cdot \nabla \mu|_{\partial \Omega} &= 0, & \text{on } \partial \Omega \setminus \Sigma, \\
P_{S_1}(\mu(Du + Du^\top)\nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
u &= 0, & \text{on } S_2.
\end{aligned} \tag{6.7}$$

Integrating (6.7)₅ over Σ and using the fact that h_1 is constant yields that $h_1 = 0$, since the other terms are mean value free. This yields that $(u, h) \in N(A)$, whence $N(A^2) \subset N(A)$. Since A has compact resolvent, $R(A)$ is closed in X_0 and $\lambda = 0$ is a pole of $(\lambda - A)^{-1}$. Therefore $\lambda = 0$ is semi-simple, cf. [17], and $X_0 = N(A) \oplus R(A)$. Since also $\sigma(A|_{R(A)}) \subset \mathbb{C}_+$ we obtain that the restricted semigroup $e^{-At}|_{R(A)}$ is exponentially stable. \square

Define now a linear operator $L : D(L) \subset X_1 \rightarrow \tilde{X}_0$ by $L(u, h) := A(u, h)$, where

$$D(L) := D(A) \cap \{(u, h) \in X_1 : (h, 1)_{L_2(\Sigma)} = 0\},$$

and $\tilde{X}_0 := X_0 \cap \{(u, h) \in X_0 : P_0^\Sigma h = 0\}$. Hereby

$$P_0^\Sigma h := \frac{1}{|\Sigma|} \int_\Sigma h dx.$$

Then L is well-defined and $\sigma(-L) \subset \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \leq -\kappa < 0\}$ for some $\kappa > 0$, since we have a spectral gap.

6.2. Parametrization of the nonlinear phase manifold. We recall, cf. (3.1), that the transformed equations around the trivial equilibrium $(0, \Sigma) \in \mathcal{E}$ read as

$$\begin{aligned} \rho \partial_t u - \mu \Delta u + \nabla p &= F_u(h, u, p), & \text{in } \Omega \setminus \Sigma, \\ \operatorname{div} u &= G_d(h, u), & \text{in } \Omega \setminus \Sigma, \\ -\llbracket \mu(Du + Du^\top) - pI \rrbracket e_3 &= \sigma \Delta_{x'} h e_3 + G_S(h, u, p), & \text{on } \Sigma, \\ \llbracket u \rrbracket &= 0, & \text{on } \Sigma, \\ \partial_t h &= u_3 - \llbracket \partial_3 \eta \rrbracket + G_\Sigma(h, u, \eta), & \text{on } \Sigma, \\ (-\nabla_{x'} h, 0)^\top \cdot \nu_{S_1} &= 0, & \text{on } \partial \Sigma, \\ \Delta \eta &= G_c(h, \eta), & \text{in } \Omega \setminus \Sigma, \\ \eta|_\Sigma - \sigma \Delta_{x'} h &= G_\kappa(h), & \text{on } \Sigma, \\ \nu_{\partial \Omega} \cdot \nabla \eta|_{\partial \Omega} &= G_N(h, \eta), & \text{on } \partial \Omega \setminus \Sigma, \\ P_{S_1}(\mu(Du + Du^\top)\nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\ u \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\ u &= 0, & \text{on } S_2, \\ u(0) &= u_0, & \text{on } \Omega \setminus \Sigma, \\ h(0) &= h_0, & \text{on } \Sigma, \end{aligned} \tag{6.8}$$

where $F_u(h, u, p) := a^\pm(h; D_x)(u, p) + \bar{a}(h, u)$, cf. (3.1). The nonlinear phase manifold is given by

$$\begin{aligned} \text{PM} &:= \{(u, h) \in W_r^{2-2/r}(\Omega \setminus \Sigma) \cap B_{qp}^{4-1/q-3/p}(\Sigma) : \operatorname{div} u = G_d, \\ &\quad P_\Sigma(\mu^\pm(Du + Du^\top)e_3) = ((G_S)_{1,2}, 0), \llbracket u \rrbracket = 0, (\nabla_{x'} h|_{n_{\partial \Sigma}}) = 0, \\ &\quad (h|1)_{L_2(\Sigma)} = 0, P_{S_1}(\mu^\pm(Du + Du^\top)\nu_{S_1}) = 0, u \cdot \nu_{S_1} = 0, u|_{S_2} = 0\} \end{aligned}$$

as a subset of $X_\gamma := W_r^{2-2/r}(\Omega \setminus \Sigma) \cap B_{qp}^{4-1/q-3/p}(\Sigma)$. The linear phase manifold is given by

$$\begin{aligned} \text{PM}_0 := \{ & (u, h) \in W_r^{2-2/r}(\Omega \setminus \Sigma) \cap B_{qp}^{4-1/q-3/p}(\Sigma) : \operatorname{div} u = 0, \\ & P_\Sigma(\mu^\pm(Du + Du^\top)e_3) = 0, \llbracket u \rrbracket = 0, (\nabla_{x'} h|_{n_{\partial\Sigma}}) = 0, \\ & (h|_1)_{L_2(\Sigma)} = 0, P_{S_1}(\mu^\pm(Du + Du^\top)\nu_{S_1}) = 0, u \cdot \nu_{S_1} = 0, u|_{S_2} = 0 \}. \end{aligned}$$

We now refer to Section 4.2 in [29], where it is shown that there is a local parametrization of PM over PM_0 around zero. More precisely there is a small $r > 0$, such that for every $(u_0, h_0) \in B(r, 0) \subset \text{PM}$ there is a C^2 -function φ and a decomposition

$$(u_0, h_0) = (\tilde{u}_0, \tilde{h}_0) + (\varphi(\tilde{u}_0, \tilde{h}_0), 0), \quad (\tilde{u}_0, \tilde{h}_0) \in \text{PM}_0. \quad (6.9)$$

For details we refer to Proposition 4.3 and Section 4.2 in [29].

6.3. Convergence to equilibria. We now state and prove the main result.

Theorem 6.3. *The trivial equilibrium $(0, \Sigma) \in \mathcal{E}$ is stable in the following sense. For each $\varepsilon > 0$ there exists some $\delta = \delta(\varepsilon) > 0$ such that for all initial values $(u_0, h_0) \in X_\gamma \cap \text{PM}$ satisfying*

$$|u_0|_{W_r^{2-2/r}(\Omega \setminus \Sigma)} + |h_0|_{B_{qp}^{4-1/q-3/p}(\Sigma)} \leq \delta(\varepsilon), \quad (6.10)$$

there exists some global in time solution

$$\begin{aligned} u & \in W_r^1(\mathbb{R}_+; L_r(\Omega)) \cap L_r(\mathbb{R}_+; W_r^2(\Omega \setminus \Sigma)), \\ h & \in W_p^1(\mathbb{R}_+; W_q^{1-1/q}(\Sigma)) \cap L_p(\mathbb{R}_+; W_q^{4-1/q}(\Sigma)), \end{aligned}$$

such that

$$|u(t)|_{W_r^{2-2/r}(\Omega \setminus \Sigma)} + |h(t)|_{B_{qp}^{4-1/q-3/p}(\Sigma)} \leq \varepsilon, \quad t \in \mathbb{R}_+.$$

Moreover,

$$|u(t)|_{W_r^{2-2/r}(\Omega \setminus \Sigma)} + |h(t) - P_0^\Sigma h_0|_{B_{qp}^{4-1/q-3/p}(\Sigma)} \xrightarrow{t \rightarrow \infty} 0,$$

where $P_0^\Sigma h_0 = \frac{1}{|\Sigma|} \int_\Sigma h_0$ is the mean value of h_0 . The convergence is at an exponential rate.

Proof. We follow the lines of [4] and [29]. Let $\varepsilon > 0$ be given and $(u_0, h_0) \in X_\gamma \cap \text{PM}$ such that the smallness condition (6.10) holds for some $\delta > 0$ to be specified later. By (6.9), we can decompose the initial data

$$(u_0, h_0) = (0, P_0^\Sigma h_0) + (\tilde{u}_0, \tilde{h}_0) + (\varphi(\tilde{u}_0, \tilde{h}_0), 0),$$

where $(\tilde{u}_0, \tilde{h}_0) + (\varphi(\tilde{u}_0, \tilde{h}_0), 0) \in \text{PM}$ and $(\tilde{u}_0, \tilde{h}_0) \in \text{PM}_0$. We now want to decompose the solution $(u(t), h(t))$ suitably and write

$$(u(t), h(t)) = (0, P_0^\Sigma h_0) + (\tilde{u}(t), \tilde{h}(t)) + (\bar{u}(t), \bar{h}(t)), \quad t \in \mathbb{R}_+,$$

where $(\tilde{u}(t), \tilde{h}(t)) \in \mathbf{PM}_0$ for $t \in \mathbb{R}_+$, and estimate each term separately. We consider the two coupled systems

$$\begin{aligned}
\omega \rho \bar{u} + \rho \partial_t \bar{u} - \mu^\pm \Delta \bar{u} + \nabla \bar{\pi} &= F_u(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{u} + \bar{u}, \tilde{\pi} + \bar{\pi}), & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} \bar{u} &= G_d(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{u} + \bar{u}), & \text{in } \Omega \setminus \Sigma, \\
-P_\Sigma(\llbracket \mu^\pm (D\bar{u} + D\bar{u}^\top) e_3 \rrbracket) &= G_S^\parallel(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{u} + \bar{u}), & \text{on } \Sigma, \\
-2\llbracket \mu^\pm \partial_3 \bar{u}_3 \rrbracket + \llbracket \bar{\pi} \rrbracket - \sigma \Delta_{x'} \bar{h} &= G_S^\perp(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{u} + \bar{u}), & \text{on } \Sigma, \\
\llbracket \bar{u} \rrbracket &= 0, & \text{on } \Sigma, \\
\omega \bar{h} + \partial_t \bar{h} - \bar{u}_3 + \llbracket \partial_3 \bar{\eta} \rrbracket &= G_\Sigma(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{u} + \bar{u}, \tilde{\eta} + \bar{\eta}), & \text{on } \Sigma, \\
(-\nabla_{x'} \bar{h}, 0)^\top \cdot \nu_{S_1} &= 0, & \text{on } \partial \Sigma, \\
\Delta \bar{\eta} &= G_c(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{\eta} + \bar{\eta}), & \text{in } \Omega \setminus \Sigma, \\
\bar{\eta}|_\Sigma - \sigma \Delta_{x'} \bar{h} &= G_\kappa(P_0^\Sigma h_0 + \tilde{h} + \bar{h}), & \text{on } \Sigma, \\
\nu_{\partial \Omega} \cdot \nabla \bar{\eta}|_{\partial \Omega} &= G_N(P_0^\Sigma h_0 + \tilde{h} + \bar{h}, \tilde{\eta} + \bar{\eta}), & \text{on } \partial \Omega \setminus \Sigma, \\
P_{S_1}(\mu^\pm (D\bar{u} + D\bar{u}^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
\bar{u} \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
\bar{u} &= 0, & \text{on } S_2, \\
\bar{u}(0) &= \varphi(\tilde{u}_0, \tilde{h}_0), & \text{on } \Omega \setminus \Sigma, \\
\bar{h}(0) &= 0, & \text{on } \Sigma,
\end{aligned} \tag{6.11}$$

where $\omega > 0$, and

$$\begin{aligned}
\rho \partial_t \tilde{u} - \mu^\pm \Delta \tilde{u} + \nabla \tilde{\pi} &= \omega \rho (I - T_1) \tilde{u}, & \text{in } \Omega \setminus \Sigma, \\
\operatorname{div} \tilde{u} &= 0, & \text{in } \Omega \setminus \Sigma, \\
-P_\Sigma(\llbracket \mu^\pm (D\tilde{u} + D\tilde{u}^\top) e_3 \rrbracket) &= 0, & \text{on } \Sigma, \\
-2\llbracket \mu^\pm \partial_3 \tilde{u}_3 \rrbracket + \llbracket \tilde{\pi} \rrbracket - \sigma \Delta_{x'} \tilde{h} &= 0, & \text{on } \Sigma, \\
\llbracket \tilde{u} \rrbracket &= 0, & \text{on } \Sigma, \\
\partial_t \tilde{h} - \tilde{u}_3 + \llbracket \partial_3 \tilde{\eta} \rrbracket &= \omega (\bar{h} - P_0^\Sigma \tilde{h}), & \text{on } \Sigma, \\
(-\nabla_{x'} \tilde{h}, 0)^\top \cdot \nu_{S_1} &= 0, & \text{on } \partial \Sigma, \\
\Delta \tilde{\eta} &= 0, & \text{in } \Omega \setminus \Sigma, \\
\tilde{\eta}|_\Sigma - \sigma \Delta_{x'} \tilde{h} &= 0, & \text{on } \Sigma, \\
\nu_{\partial \Omega} \cdot \nabla \tilde{\eta}|_{\partial \Omega} &= 0, & \text{on } \partial \Omega \setminus \Sigma, \\
P_{S_1}(\mu^\pm (D\tilde{u} + D\tilde{u}^\top) \nu_{S_1}) &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
\tilde{u} \cdot \nu_{S_1} &= 0, & \text{on } S_1 \setminus \partial \Sigma, \\
\tilde{u} &= 0, & \text{on } S_2, \\
\tilde{u}(0) &= \tilde{u}_0, & \text{on } \Omega \setminus \Sigma, \\
\tilde{h}(0) &= \tilde{h}_0, & \text{on } \Sigma.
\end{aligned} \tag{6.12}$$

Let us note a few things here. The right hand side of (6.12)₁ can equivalently be written as $\omega\rho(I - T_1)\bar{u} = \omega\rho\bar{u} - \omega\rho\nabla\bar{q}$, where $\bar{q} \in \dot{H}_r^1(\Omega \setminus \Sigma)$ is the unique solution of the weak transmission problem

$$\begin{aligned} (\nabla\bar{q}|\nabla\psi)_{L_2(\Omega)} &= (\bar{u}|\nabla\psi)_{L_2(\Omega)}, & \text{for all } \psi \in W_{r'}^1(\Omega), \\ \llbracket\bar{q}\rrbracket &= 0, & \text{on } \Sigma. \end{aligned}$$

Furthermore, the initial value \tilde{h}_0 in (6.12) is mean value free. Note that the right hand side of (6.12)₆ is mean value free as well, hence an integration of (6.12)₆ over Σ yields that \tilde{h} stays mean value free for all times $t > 0$. In particular, we can equivalently rewrite (6.12) in the projected base space \tilde{X}_0 as

$$\frac{d}{dt}\tilde{z}(t) + L\tilde{z}(t) = R(\tilde{z})(t), \quad t > 0, \quad z(0) = \tilde{z}_0 := (\tilde{u}_0, \tilde{h}_0). \quad (6.13)$$

Here, $\tilde{z} := (\tilde{u}, \tilde{h})$, $\tilde{z} := (\tilde{u}, \tilde{h})$ and $R(\tilde{z}) := (\omega(I - T_1)\tilde{u}, (I - P_0^\Sigma)\tilde{h})$. Note that by Lemma 6.2, the spectral bound of $-L$ satisfies $s(-L) \leq -\kappa < 0$ and the restricted semigroup e^{-Lt} is exponentially stable on \tilde{X}_0 .

We now solve this evolution equation in exponentially time-weighted spaces to get suitable decay estimates, cf. [4] and [29]. Let us introduce notation. Let $\mathbb{E}_u(\mathbb{R}_+) := H_r^1(\mathbb{R}_+; L_r(\Omega)) \cap L_r(\mathbb{R}_+; H_r^2(\Omega \setminus \Sigma))$ and

$$\mathbb{E}_h(\mathbb{R}_+) := W_p^1(\mathbb{R}_+; W_q^{1-1/q}(\Sigma)) \cap L_p(\mathbb{R}_+; W_q^{4-1/q}(\Sigma)).$$

For $\beta \in [0, -s(-L))$ define

$$\begin{aligned} e^{-\beta t}\mathbb{E}_u(\mathbb{R}_+) &:= \{w \in L_r(\mathbb{R}_+; L_r(\Omega)) : e^{\beta t}w \in \mathbb{E}_u(\mathbb{R}_+)\}, \\ e^{-\beta t}\mathbb{E}_h(\mathbb{R}_+) &:= \{w \in L_p(\mathbb{R}_+; L_q(\Sigma)) : e^{\beta t}w \in \mathbb{E}_h(\mathbb{R}_+)\}. \end{aligned}$$

In a similar way we define $e^{-\beta t}L_r(\mathbb{R}_+; L_r(\Omega))$. Since $0 \leq \beta < -s(-L)$, we obtain that for every

$$(f_u, f_h) \in e^{-\beta t}[L_r(\mathbb{R}_+; L_r(\Omega)) \times L_p(\mathbb{R}_+; W_q^{1-1/q}(\Sigma))],$$

and $(\hat{u}_0, \hat{h}_0) \in X_\gamma$ there is a unique solution $(u, h) \in e^{-\beta t}[\mathbb{E}_u(\mathbb{R}_+) \times \mathbb{E}_h(\mathbb{R}_+)]$ of the linear evolution problem

$$\partial_t(u, h) + L(u, h) = (f_u, f_h), \quad t \in \mathbb{R}_+, \quad (u, h)|_{t=0} = (\hat{u}_0, \hat{h}_0),$$

by maximal regularity in exponentially time-weighted spaces. Furthermore, there is some $M > 0$ such that

$$\begin{aligned} |(u, h)|_{e^{-\beta t}[\mathbb{E}_u(\mathbb{R}_+) \times \mathbb{E}_h(\mathbb{R}_+)]} \\ \leq M|(f_u, f_h, \hat{u}_0, \hat{h}_0)|_{e^{-\beta t}[L_r(\mathbb{R}_+; L_r(\Omega)) \times L_p(\mathbb{R}_+; W_q^{1-1/q}(\Sigma))] \times X_\gamma}. \end{aligned}$$

In particular, we may then easily solve (6.13) in dependence of $\tilde{z} = (\tilde{u}, \tilde{h})$,

$$(\tilde{u}, \tilde{h}) = \left(\frac{d}{dt} + L, \text{tr}|_{t=0} \right)^{-1} (\omega(I - T_1)\tilde{u}, (I - P_0^\Sigma)\tilde{h}, \tilde{u}_0, \tilde{h}_0). \quad (6.14)$$

Let us now discuss problem (6.11). For given $\omega > 0$, let L_ω be given by the left hand side of (6.11) and N the collection of nonlinearities on the right hand side. Then we can rewrite problem (6.11) in the shorter form

$$L_\omega \bar{w} = N(w_\infty + \tilde{w} + \bar{w}), \quad (\bar{u}, \bar{h})(0) = (\varphi(\tilde{u}_0, \tilde{h}_0), 0),$$

where $\bar{w} := (\bar{u}, \bar{h}, \bar{\pi}, \bar{\eta})$, $\tilde{w} := (\tilde{u}, \tilde{h}, \tilde{\pi}, \tilde{\eta})$ and $w_\infty := (0, P_0^\Sigma h_0, 0, 0)$. Note at this point that w_∞ is constant and N does not explicitly depend on w_∞ . Furthermore, due to the first part of the proof, \tilde{w} depends only on $(\tilde{u}_0, \tilde{h}_0, \bar{u}, \bar{h})$, cf. (6.14).

In order to solve problem (6.11) we need to resolve the initial data and the compatibility conditions at $t = 0$ properly. By solving certain auxiliary problems in exponentially weighted spaces, we may construct an extension operator

$$\text{ext}_\beta : \bar{X}_\gamma \rightarrow e^{-\beta} [\mathbb{E}_u(\mathbb{R}_+) \times \mathbb{E}_h(\mathbb{R}_+)],$$

satisfying $\text{ext}_\beta(v, g)|_{t=0} = (v, g)$ for all $(v, g) \in \bar{X}_\gamma$, where

$$\begin{aligned} \bar{X}_\gamma := \{ & (u, h) \in X_\gamma : u|_{S_2} = 0, (u|_{S_1}) = 0, P_{S_1}(\mu^\pm(Du + Du^\top)\nu_{S_1}) = 0, \\ & \llbracket u \rrbracket = 0, (\nabla_{x'} h|_{\nu_{\partial\Sigma}}) = 0 \}, \end{aligned}$$

cf. [29]. Now define

$$M(\tilde{u}_0, \tilde{h}_0, \bar{w}) := N(w_\infty + \tilde{w} + \bar{w} + \text{ext}_\beta[(\varphi(\tilde{u}_0, \tilde{h}_0), 0) - (\bar{u}(0), \bar{h}(0))]).$$

By construction, $M(\tilde{u}_0, \tilde{h}_0, \bar{w})|_{t=0} = N(u_0, h_0, 0, 0)$. This allows us to solve the problem

$$L_\omega \bar{w} = M(\tilde{u}_0, \tilde{h}_0, \bar{w}), \quad (\bar{w}_1, \bar{w}_2)|_{t=0} = (\varphi(\tilde{u}_0, \tilde{h}_0), 0),$$

by the implicit function theorem, since all relevant compatibility conditions at $t = 0$ are satisfied. Following the lines of [4], we obtain that there is some small $\rho > 0$ and a ball $B(0, \rho) \subset X_\gamma \cap \text{PM}_0$, such that there is a $\Phi \in C^1(B(0, \rho); e^{-\beta t} [\mathbb{E}_u(\mathbb{R}_+) \times \mathbb{E}_h(\mathbb{R}_+) \times \mathbb{E}_\pi(\mathbb{R}_+) \times \mathbb{E}_\eta(\mathbb{R}_+)])$ satisfying $\bar{w} = \Phi(\tilde{u}_0, \tilde{h}_0)$. By construction, \bar{w} is the solution of (6.11). Here, $\mathbb{E}_\pi(\mathbb{R}_+) := L_r(\mathbb{R}_+; \dot{H}_r^1(\Omega \setminus \Sigma))$, $\mathbb{E}_\eta(\mathbb{R}_+) := L_p(\mathbb{R}_+; W_q^2(\Omega \setminus \Sigma))$.

We then obtain that the convergence $(u(t), h(t)) \rightarrow (0, P_0^\Sigma h_0)$ in X_γ is at an exponential rate. The proof is complete. \square

ACKNOWLEDGEMENTS

M.R. would like to thank Harald Garcke for pointing out existing work in [2] and Helmut Abels for inspiring discussions regarding the boundary conditions. The work of M.R. is financially supported by the DFG graduate school GRK 1692. The support is gratefully acknowledged.

REFERENCES

- [1] H. Abels. *Pseudodifferential and Singular Integral Operators*. De Gruyter, 2011.
- [2] H. Abels, G. Garcke, and G. Grün. Thermodynamically consistent, frame indifferent diffuse interface models for incompressible two-phase flows with different densities. *Mathematical Models and Methods in Applied Sciences*, 22(3), 2012.
- [3] H. Abels, M. Rauchecker, and M. Wilke. Well-Posedness and qualitative behaviour of the Mullins-Sekerka problem with ninety-degree angle boundary contact, 2019. <http://arxiv.org/abs/1902.03611>.
- [4] H. Abels and M. Wilke. Well-posedness and qualitative behaviour of solutions for a two-phase Navier-Stokes-Mullins-Sekerka system. *Interfaces and Free Boundaries*, 15:39–75, 2013.
- [5] H. Amann. Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. *Function Spaces, Differential Operators and Nonlinear Analysis*, pages 9–126, 1993.
- [6] H. Amann. *Linear and Quasilinear Parabolic problems, Volume I: Abstract linear theory*. Birkhäuser, 1995.
- [7] W. Arendt. Gaussian estimates and interpolation of the spectrum in L^p . *Differential and integral equations*, 7(5):1153–1168, 1994.

- [8] H. Bahouri, J. Chemin, and R. Danchin. *Fourier Analysis and nonlinear partial differential equations*. Springer, 2011.
- [9] D. Bothe and J. Prüss. Lp-Theory for a Class of Non-Newtonian Fluids. *SIAM J. Math. Analysis*, 39:379–421, 2007.
- [10] J. Bourgain. Extension of a result of Benedek, Calderon and Panzone. *Ark. Mat.*, 22(1-2):91–95, 12 1984.
- [11] G. Dore. Maximal regularity in L^p spaces for an abstract Cauchy problem. *Adv. Differential Equations*, 5(1-3):293–322, 2000.
- [12] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*. Springer, New York, 2000.
- [13] J. Escher and G. Simonett. A Center Manifold Analysis for the Mullins-Sekerka Model. *Journal of Differential Equations*, 143:267–292, 1998.
- [14] H. Garcke and M. Rauchecker. Stability analysis for stationary solutions of the Mullins-Sekerka flow with boundary contact, 2019. <https://arxiv.org/abs/1907.00833>.
- [15] E. Hanzawa. Classical solutions of the stefan problem. *Tohoku Math. J. (2)*, 33(3):297–335, 1981.
- [16] M. Kaip. *General parabolic mixed order systems in L^p and applications*. PhD thesis, Universität Konstanz, 2012.
- [17] A. Lunardi. *Analytic semigroups and Optimal Regularity in Parabolic Problems*. Springer, 1995.
- [18] A. Lunardi. *Interpolation Theory*. Springer, 2009.
- [19] A. Meyries and M. Veraar. Pointwise multiplication on vector-valued function spaces with power weights. *Journal of Fourier Analysis and Applications*, 1:95–136, February 2015.
- [20] J. Prüss. Maximal Regularity for abstract parabolic problems with inhomogeneous boundary data in L_p -spaces. *Mathematica Bohemica*, 127(2):311–327, 2002.
- [21] J. Prüss. Maximal Regularity for evolution equations in L_p -spaces. *Conf. Semin. Mat. Univ. Bari*, 2002.
- [22] J. Prüss and G. Simonett. On the Rayleigh-Taylor instability for the two-phase Navier-Stokes equations. *Indiana Univ. Math. J.*, 59(6):1853–1871, 2010.
- [23] J. Prüss and G. Simonett. On the two-phase Navier-Stokes equations with surface tension. *Interfaces Free Bound.*, 12(3):311–345, 2010.
- [24] J. Prüss and G. Simonett. *Moving interfaces and quasilinear parabolic evolution equations*. Birkhäuser Verlag, 2016.
- [25] M. Rauchecker. Strong solutions to the Stefan problem with Gibbs-Thomson correction and boundary contact. <https://arxiv.org/abs/2001.06438>.
- [26] M. Rauchecker and M. Wilke. Rayleigh-Taylor instability of the two-phase Navier-Stokes/Mullins-Sekerka equations with boundary contact. 2019.
- [27] T. Runst. Mapping properties of nonlinear operators in spaces of Triebel-Lizorkin and Besov type. *Analysis Mathematica*, 12(4):313–346, 1986.
- [28] H. Triebel. *Theory of Function Spaces*. Birkhäuser, 2000.
- [29] M. Wilke. Rayleigh-Taylor instability for the two-phase Navier-Stokes equations with surface tension in cylindrical domains, 2013. Habilitationsschrift, Universität Halle. Available online at <http://arxiv.org/abs/1703.05214>.