

# THE TOPOLOGICAL DIMENSION OF RADIAL JULIA SETS

DAVID S. LIPHAM

**ABSTRACT.** We prove that the meandering set for  $f_a(z) = e^z + a$  is homeomorphic to the space of irrational numbers whenever  $a$  belongs to the Fatou set of  $f_a$ . This extends recent results by Vasiliki Evdoridou and Lasse Rempe. It implies that the radial Julia set of  $f_a$  has topological dimension zero for all attracting and parabolic parameters, including all  $a \in (-\infty, -1]$ . Similar results are obtained for Fatou's function  $f(z) = z + 1 + e^{-z}$ .

## 1. INTRODUCTION

The focus of this paper is the class of exponential functions

$$f_a(z) = e^z + a$$

for attracting and parabolic parameters  $a \in \mathbb{C}$ . The Fatou set  $F(f_a)$  is the set of normality for the family of iterates  $\{f_a^n : n \in \mathbb{N}\}$ , and the Julia set  $J(f_a) = \mathbb{C} \setminus F(f_a)$  is its complement. The *escaping set*  $I(f_a)$  is the set of points  $z \in \mathbb{C}$  such that  $f_a^n(z) \rightarrow \infty$ , where  $\infty$  is the point at infinity on the Riemann sphere. It is known that  $I(f_a) \subset J(f_a)$  in general, and  $f_a$  has an attracting or parabolic cycle if and only if  $a \in F(f_a)$  [7, Proposition 2.1].

The *radial Julia set*  $J_r(f_a)$  is defined to be the set of points  $z \in J(f_a)$  with the following property: There exists  $\delta > 0$  and infinitely many  $n \in \mathbb{N}$  such that the spherical disc of radius  $\delta$  around  $f^n(z)$  can be pulled back univalently along the orbit of  $z$ . For the parameters of interest in this paper,  $J_r(f_a)$  is approximately equal to the set of non-escaping points in  $J(f_a)$ . Indeed,

$$J_r(f_a) \subset J(f_a) \setminus I(f_a)$$

for all  $a \in F(f_a)$ , and if  $f_a$  has an attracting periodic orbit then

$$J_r(f_a) = J(f_a) \setminus I(f_a);$$

see [7, Corollary 2.2]. The *meandering set*  $J_m(f_a)$  is the larger collection of points in  $J(f_a)$  whose orbits do not converge to  $\infty$  as quickly as possible. Precise definitions of  $J_m(f_a)$  and  $J_r(f_a)$  are provided in Section 2.

All of the sets mentioned above will be viewed as topological subspaces of the complex plane  $\mathbb{C}$ . A topological space  $X$  is

- *totally separated* if for every two points  $x, y \in X$  there is a clopen subset of  $X$  which contains  $x$  and misses  $y$ ; and
- *zero-dimensional* if  $X$  has a basis of clopen sets.

Note that every zero-dimensional space is totally separated, but the converse is false. In fact,  $J(f_a)$  contains totally separated spaces which are not zero-dimensional; see the Remark in Section 3.2.

2010 *Mathematics Subject Classification.* 37F10, 30D05, 54F45.

*Key words and phrases.* radial Julia set, complex exponential, meandering set, zero-dimensional.

Evdoridou and Rempe recently proved that if  $a \in F(f_a)$ , then  $J_m(f_a) \cup \{\infty\}$  (as a subspace of the Riemann sphere  $\mathbb{C} \cup \{\infty\}$ ) is totally separated [7, Theorem 1.3]. In this paper, we strengthen their result by proving that  $J_m(f_a)$  and  $J_m(f_a) \cup \{\infty\}$  are each homeomorphic to the zero-dimensional space of irrational numbers  $\mathbb{P}$ . Since  $J_r(f_a) \subset J_m(f_a)$  and zero-dimensionality is hereditary, we find that  $J_r(f_a)$  and  $J_r(f_a) \cup \{\infty\}$  are topologically zero-dimensional as well.

By contrast, the Hausdorff dimension of  $J_r(f_a)$  is always greater than 1; see [15, Theorem 2.1] and [9, Theorem 2]. The topological dimension when  $a \in J(f_a)$  can also differ. For instance, if  $f_a$  is postsingularly finite (meaning that  $\{f_a^n(a) : n \in \mathbb{N}\}$  is finite), then  $J_r(f_a) = J(f_a) \setminus I(f_a)$  contains an unbounded connected set; see [7, Corollary 2.2(c)], [13, Theorem 4.3] and [7, Proposition 2.4(e)]. In this case  $J_r(f_a)$  and  $J_m(f_a)$  must have positive topological dimension. A simple example of a postsingularly finite parameter is  $a = \ln(\pi) + \frac{\pi}{2}i$ .

At the end of the paper we will apply our work to Fatou's function

$$z \mapsto z + 1 + e^{-z}.$$

This will strengthen the results of [6] and [7, Section 5]. In particular, we will see that the entire non-escaping set for Fatou's function is zero-dimensional.

**Structure of the paper.** In Section 2 we will state the definitions of  $J_r(f_a)$ ,  $J_m(f_a)$  and related sets, including the fast escaping set  $A(f_a)$ . The endpoints and connected components of  $J(f_a)$  are analyzed in Section 3, and in Section 4 we prove two crucial lemmas regarding  $A(f_a)$ . All of that will be integrated in Section 5 to prove our main results. And in Section 6 we consider Fatou's function.

## 2. FORMAL DEFINITIONS

In the following definitions we assume that  $f$  is a transcendental entire function, and  $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$  is the Riemann sphere.

**2.1. Definition of  $J_r(f)$ .** The radial Julia set was introduced by Urbanski [14] and McMullen [10]. It is defined as follows.

For any  $z_0 \in \mathbb{C}$  let  $\mathbb{D}_\delta(z_0) = \{z \in \widehat{\mathbb{C}} : \text{dist}(z_0, z) < \delta\}$ , where the distance is with respect to the conformal metric on  $\widehat{\mathbb{C}}$  (the unique metric of constant curvature 1). If  $A \subset \widehat{\mathbb{C}}$  and  $z \in A$ , then let  $\text{Comp}(z, A)$  denote the connected component of  $z$  in  $A$ . Define  $J_r(f)$  to be the set of points  $z \in J(f)$  with the property: There exists  $\delta > 0$  and infinitely many  $n \in \mathbb{N}$  such that

$$f^n : \text{Comp}(z, f^{-n}(\mathbb{D}_\delta(f^n(z)))) \rightarrow \mathbb{D}_\delta(f^n(z))$$

is a conformal isomorphism.

The definition of  $J_r(f)$  will not be directly applied in this paper. We will only need the fact  $J_r(f) \subset J(f) \setminus I(f)$  for the functions studied here.

**2.2. Definitions of  $A(f)$  and  $J_m(f)$ .** The *fast escaping set*  $A(f)$  was introduced by Bergweiler and Hinkkanen [4], and was studied further by Rippon and Stallard [12]. It is defined as follows.

For each  $r > 0$  let  $M(r) = M(r, f) = \max\{|f(z)| : |z| = r\}$ . Since  $f$  is transcendental, we can choose  $R > 0$  sufficiently large so that  $M^n(R) \rightarrow \infty$  and  $M(r) > r$  for all  $r \geq R$ . Define

$$A(f) = \{z \in \mathbb{C} : \exists \ell \geq 0 \text{ such that } f^{\ell+n}(z) \geq M^n(R) \text{ for all } n \in \mathbb{N}\}.$$

This definition is independent of the choice of  $R$  by [12, Theorem 2.2]. Informally,  $A(f)$  is the set of points  $z \in \mathbb{C}$  such that  $f^n(z) \rightarrow \infty$  as quickly as possible. Note that  $A(f) \subset I(f)$ .

The set  $J_m(f)$  is defined to be the complement  $J(f) \setminus A(f)$ . As previously indicated,  $J_m(f_a) \supset J_r(f_a)$  for all  $a \in F(f_a)$ .

### 3. GEOMETRY OF JULIA SETS

In this section we examine some important aspects of  $J(f_a)$  and its connected components. *Throughout the section we assume  $a \in F(f_a)$ .*

**3.1. General structure of  $J(f_a)$ .** For certain attracting parameters, including all  $a \in (-\infty, -1)$ ,  $J(f_a)$  is homeomorphic to a particular subspace  $\bar{X}$  of  $\mathbb{Z}^{\mathbb{N}} \times [0, \infty)$ . The set  $\bar{X}$  is constructed in [11] and is of the form

$$\bar{X} = \bigcup_{\underline{s} \in \mathbb{Z}^{\mathbb{N}}} \{\underline{s}\} \times [t_{\underline{s}}, \infty)$$

where each  $t_{\underline{s}} \in [0, \infty]$  and  $[\infty, \infty) = \emptyset$ . More generally, by [11, Theorem 9.1], for any parameter  $a \in F(f_a)$  there is a closed continuous surjection

$$H : \bar{X} \rightarrow J(f_a).$$

If  $H$  is a homeomorphism (e.g. when  $a \in (-\infty, -1)$ ), then  $J(f_a)$  is a *Cantor bouquet* in the sense of [2] and [3, Definition 2.7]. Otherwise,  $J(f_a)$  is a *pinched Cantor bouquet*. Figures 1 and 2 show a Cantor bouquet Julia set for  $a = -2$ , and pinched Cantor bouquet Julia sets for three other parameters.

We now indicate some features of the escaping set  $I(f_a)$  and the pinching that may occur. There is a set  $X \subset \bar{X}$  (the same  $X$  defined in [11]) such that:

- (1)  $\{\underline{s}\} \times (t_{\underline{s}}, \infty) \subset X$  for all  $\underline{s} \in \mathbb{Z}^{\mathbb{N}}$  (see [11, Observation 3.1]),
- (2)  $H^{-1}(I(f_a)) = X$  (see [11, Corollary 5.3]), and
- (3)  $H \upharpoonright X : X \rightarrow I(f_a)$  is a homeomorphism (see [11, Theorem 9.1]).

Thus, if  $H^{-1}(z)$  is non-degenerate then  $z$  may be viewed as a point of  $J(f_a)$  where multiple rays of  $\bar{X}$  are “pinched together” at their endpoints. Observe also that a non-degenerate pre-image  $H^{-1}(z)$  cannot intersect  $X$ , so the pinched points in  $J(f_a)$  are all non-escaping (and thus belong to  $J_r(f_a)$  and  $J_m(f_a)$ ).



**3.2. Endpoints of  $J(f_a)$ .** The endpoints of rays of  $\bar{X}$  (the points  $\langle \underline{s}, t_{\underline{s}} \rangle \in \bar{X}$ ) are closely related to the endpoints of  $J(f_a)$  that are defined as follows.

A point  $z \in J(f_a)$  is on a *hair* if there exists an arc  $\alpha : [-1, 1] \hookrightarrow I(f_a)$  such that  $\alpha(0) = z$ . A point  $z_0 \in J(f_a)$  is an *endpoint* if  $z_0$  is not on a hair and there is an arc  $\alpha : [0, 1] \hookrightarrow J(f_a)$  such that  $\alpha(0) = z_0$  and  $\alpha(t) \in I(f_a)$  for all  $t > 0$ .

The set of all endpoints of  $J(f_a)$  is denoted  $E(f_a)$ .

**Observation 1.**  $E(f_a) = \{H(\langle \underline{s}, t_{\underline{s}} \rangle) : \underline{s} \in \mathbb{Z}^{\mathbb{N}} \text{ and } t_{\underline{s}} < \infty\}$ .

*Proof.* If  $t \in (t_{\underline{s}}, \infty)$  then  $H(\langle \underline{s}, t \rangle)$  is on a hair by items (1) and (3) above. Therefore  $E(f_a) \subset \{H(\langle \underline{s}, t_{\underline{s}} \rangle) : \underline{s} \in \mathbb{Z}^{\mathbb{N}} \text{ and } t_{\underline{s}} < \infty\}$ . We now prove the other inclusion. Let  $z_0 = H(\langle \underline{s}, t_{\underline{s}} \rangle)$  be given. Since  $\mathbb{Z}^{\mathbb{N}}$  is totally separated, every arc in  $\bar{X}$  containing  $\langle \underline{s}, t_{\underline{s}} \rangle$  is contained in  $\{\underline{s}\} \times [t_{\underline{s}}, \infty)$ . So by (2) and (3), every arc in  $I(f_a)$  containing  $z_0$  is contained in  $H(\{\underline{s}\} \times [t_{\underline{s}}, \infty))$ . Since  $H \upharpoonright \{\underline{s}\} \times [t_{\underline{s}}, \infty)$  is a homeomorphism, it follows that  $z_0$  is not on a hair. On the other hand,  $\alpha(t) = H(\langle \underline{s}, t_{\underline{s}} + t \rangle)$  defines

FIGURE 1. Julia sets of  $f_a$  for  $a = -2$  (left) and  $a = 5 + 3.14i$  (right).FIGURE 2. Julia sets of  $f_a$  for  $a = 2.06 + 1.57i$  (left) and  $a = 1.004 + 2.9i$  (right).

an arc  $\alpha : [0, 1] \hookrightarrow J(f_a)$  with  $\alpha(0) = z_0$  and  $\alpha(t) \in I(f_a)$  for all  $t > 0$ . Thus  $z_0 \in E(f_a)$ .  $\square$

**Remark.** The endpoint set  $E(f_a)$  is an example of a totally separated space which is not zero-dimensional. This is a consequence of [3, Theorem 1.7], which states that  $\infty$  is an explosion point for  $E(f_a)$  (when  $a \in F(f_a)$ ). This means that  $E(f_a)$  is totally separated and  $E(f_a) \cup \{\infty\}$  is connected. The latter implies that  $E(f_a)$  is not zero-dimensional, since every clopen neighborhood in  $E(f_a)$  must be unbounded.

**3.3. Connected components of  $J(f_a)$ .** The following is a simple consequence of Observation 1 and the properties of  $H$ .

**Proposition 2.** *Each connected component of  $J(f_a)$  contains a point of  $E(f_a)$ .*

*Proof.* Let  $C$  be a connected component of  $J(f_a)$ . Since  $H$  is surjective, there exists  $\langle \underline{s}, t \rangle \in H^{-1}(C)$ . By continuity of  $H$  the set  $H(\{\underline{s}\} \times [t_{\underline{s}}, \infty))$  is connected, so we have  $H(\{\underline{s}\} \times [t_{\underline{s}}, \infty)) \subset C$ . Then  $H(\langle \underline{s}, t_{\underline{s}} \rangle) \in C \cap E(f_a)$  by Observation 1.  $\square$

The next proposition shows that every connected component of  $J(f_a)$  can be written as an intersection of relatively clopen subsets of  $J(f_a)$ .

**Proposition 3.** *For every connected component  $C$  of  $J(f_a)$  there is a sequence of open subsets  $O_1 \supset O_2 \supset \dots$  of  $\mathbb{C}$  such that  $\partial O_n \subset F(f_a)$  and*

$$\bigcap_{n \in \mathbb{N}} \overline{O_n} \cap J(f_a) = C.$$

*Proof.* By [3, Proposition 6.15] there is a closed set  $\gamma \subset F(f_a)$  such that every two endpoints  $z_0, z_1 \in E(f_a)$  have different itineraries with respect to  $f_a^{-1}(\gamma)$ . This means that there exists  $n \geq 0$  such that  $f_a^n(z_0)$  and  $f_a^n(z_1)$  belong to different connected components of  $\mathbb{C} \setminus f_a^{-1}(\gamma)$ ; see [3, Definition 6.1]. For such  $n$ , note that  $z_0$  and  $z_1$  belong to different connected components of  $\mathbb{C} \setminus f_a^{-1-n}(\gamma)$ .

Now let  $C$  be a connected component of  $J(f_a)$ . Let  $z_0 \in C \cap E(f_a)$  be given by Proposition 2. For each  $n \in \mathbb{N}$  let  $O_n$  be the connected component of  $z_0$  in  $\mathbb{C} \setminus \bigcup_{k=1}^n f_a^{-k}(\gamma)$ . Clearly  $O_n$  is open,  $O_n \supset O_{n+1}$ , and  $\partial O_n \subset \bigcup_{k=1}^n f_a^{-k}(\gamma) \subset F(f_a)$  for every  $n \in \mathbb{N}$ . Also,  $\bigcap_{n \in \mathbb{N}} O_n \cap E(f_a) = \{z_0\}$  by the previous paragraph. Since  $O_n \cap J(f_a)$  is clopen in  $J(f_a)$ , we have that  $\bigcap_{n \in \mathbb{N}} O_n \cap J(f_a)$  is a union of connected components of  $J(f_a)$ . By Proposition 2 we therefore have  $\bigcap_{n \in \mathbb{N}} O_n \cap J(f_a) = C$ . Since  $O_n \cap J(f_a) = \overline{O_n} \cap J(f_a)$ , the proof is complete.  $\square$

The proof above shows that every two elements of  $E(f_a)$  belong to different connected components of  $J(f_a)$ . Thus Proposition 2 can be improved to:

**Proposition 4.** *Each connected component of  $J(f_a)$  contains exactly one endpoint.*

The final proposition implies, in particular, that each connected component of  $J(f_a)$  is bounded in the imaginary direction.

**Proposition 5.** *There exist  $c, \delta > 0$  and a closed set  $\mathcal{M} \subset F(f_a)$  such that:*

- (i)  $\{z : \operatorname{Re}(z) \leq -c\} \subset \mathcal{M}$ ; and
- (ii)  $\sup\{|\operatorname{Im}(z) - \operatorname{Im}(z')| : z, z' \in O\} \leq \delta$  for every connected component  $O$  of  $\mathbb{C} \setminus \mathcal{M}$ .

*Proof.* The proof is contained in the first part of the proof of [7, Theorem 3.1]. We include it here for completeness. Let  $U$  be the component of  $F(f_a)$  containing  $a$ . Let  $c > 0$  be large enough so that the closed disk  $D := \{z \in \mathbb{C} : |z - a| \leq e^{-c}\}$  is contained in  $U$ . By [7, Proposition 2.1] there is an arc  $\sigma \subset U$  along which the real part tends to infinity, and which intersects  $D$  at a single point. Then

$$\mathcal{M} := f_a^{-1}(D \cup \sigma) \subset F(f_a)$$

is a closed set consisting of the left half-plane  $\{z : \operatorname{Re}(z) \leq -c\}$  and countably many arcs connecting this half-plane to  $\infty$ . The arcs are all  $2\pi i\mathbb{Z}$ -translates of each other, and each arc is bounded in the imaginary direction. Thus there exists  $\varepsilon > 0$  such that the imaginary part of each arc varies by less than  $\varepsilon$ . Each connected component  $O$  of  $\mathbb{C} \setminus \mathcal{M}$  is an open region between consecutive arcs that are  $2\pi i$  apart, so  $\delta = \varepsilon + 2\pi$  is as desired.  $\square$

#### 4. KEY LEMMAS

Assume  $a \in F(f_a)$ . Let  $R > 0$  be large enough so that  $M^n(R) = M^n(R, f_a) \rightarrow \infty$  and  $M(r) > r$  for all  $r \geq R$ . For example, based on the calculations in [7, Lemma 2.5] we may take  $R = 3 + 2|a|$ .

The first lemma (below) concerns the set

$$A_R(f_a) := \{z \in \mathbb{C} : |f_a^n(z)| \geq M^n(R) \text{ for all } n \geq 0\},$$

and was mostly proved in [7].

**Lemma 6.** *There exists  $\lambda > 0$  such that for every  $z_0 \in J(f_a)$  there is an open set  $V \subset \mathbb{C}$  such that  $z_0 \in V$ ,  $\partial V \cap J(f_a) \subset A_R(f_a)$ , and  $\sup\{\operatorname{Re}(z) : z \in V\} \leq |z_0| + \lambda$ .*

*Proof.* Let  $c, \delta > 0$  and  $\mathcal{M} \subset F(f_a)$  be given by Proposition 5. Let

$$\lambda = \max\{R, c, \ln(1 + 2(|a| + \delta)), \ln(5 + |a|)\} + 6,$$

and let  $z_0 \in J(f_a)$  be given. Let  $R' = |z_0| + \lambda - 3$ . To construct  $V$  and verify its properties, we now follow the proof of [7, Theorem 3.1] with  $R'$  in the place of  $R$ .

For each  $k \geq 0$  let  $X_k = \{z \in \mathbb{C} : |f_a^k(z)| \geq M^k(R')\} \cup \bigcup_{j=0}^{k-1} f_a^{-j}(\mathcal{M})$ . Put  $X = \bigcap_{k \geq 0} X_k$ . Observe that  $X \subset A_{R'}(f_a) \cup F(f_a)$ . We know  $z_0 \notin F(f_a)$ , and  $z_0 \notin A_{R'}(f_a)$  because  $|z_0| < R' = M^0(R')$ . Hence  $z_0 \notin X$ . Let  $V$  be the connected component of  $z_0$  in  $\mathbb{C} \setminus X$ . Since  $\partial V \subset X$  and  $R' > R$ , we get

$$\partial V \cap J(f_a) \subset A_{R'}(f_a) \subset A_R(f_a).$$

Using (i) and (ii) from Proposition 5, and the fact

$$R' > \max\{|z_0|, c, 3, \ln(1 + 2(|a| + \delta))\},$$

the proof of [7, Theorem 3.1] proceeds to show that  $\sup\{\operatorname{Re}(z) : z \in V\} \leq K$  for a particular constant  $K$  that depends on  $R'$ . Tracing the definition of  $K$  back to [7, Corollary 2.7], we find that  $K = \max\{2 + \ln(5 + |a|), R' + 3\} = R' + 3 = |z_0| + \lambda$ .  $\square$

We will also need the next lemma which compares the orbits of fast escaping points and meandering points.

**Lemma 7.** *Let  $s \in A(f_a)$  and  $z_0 \in \mathbb{C} \setminus A(f_a)$ . Then for every  $\kappa > 0$  there exists  $n \in \mathbb{N}$  such that  $|f_a^n(s)| > 2|f_a^n(z_0)| + \kappa$  and  $\operatorname{Re}(f_a^n(s)) > 0$ .*

*Proof.* Let  $\kappa > 0$ . Let  $\ell \geq 0$  be such that  $f_a^{\ell+n}(s) \geq M^n(R)$  for all  $n \in \mathbb{N}$ .

By [7, Lemma 2.5],  $M(r) - r \geq e^{r-1} - r > r + \kappa$  when  $r$  is sufficiently large. Since  $M^n(R) \rightarrow \infty$ , we may substitute  $r = M^n(R)$  to see that there exists  $m \in \mathbb{N}$  such that  $M^{n+1}(R) - 2M^n(R) > \kappa$  for all  $n \geq m$ . By increasing  $m$ , we may assume  $|f_a^n(s)| > |a| + 1$  for all  $n \geq m$ . Since  $z_0 \notin A(f_a)$ , there exists  $N \in \mathbb{N}$  such that

$$|f_a^{\ell+m+1+N}(z_0)| < M^N(R).$$

Let  $n = \ell + m + N + 1$ . Then

$$2|f_a^n(z_0)| < 2M^N(R) \leq 2M^{m+N}(R) < M^{m+N+1}(R) - \kappa \leq |f_a^n(s)| - \kappa.$$

Therefore  $|f_a^n(s)| > 2|f_a^n(z_0)| + \kappa$ . Also,  $|f_a^{n+1}(s)| > |a| + 1$  implies  $\operatorname{Re}(f_a^n(s)) > 0$  since  $f_a$  maps each point with negative real part into the unit disc around  $a$ .  $\square$

## 5. MAIN RESULTS

We will now combine Propositions 3 through 5 with Lemmas 6 and 7 to compute the topological dimension of  $J_m(f_a)$ .

**Theorem 8.** *If  $a \in F(f_a)$  then  $J_m(f_a)$  is zero-dimensional.*

*Proof.* Let  $z_0 \in J_m(f_a)$ , and let  $U$  be any bounded open subset of  $\mathbb{C}$  with  $z_0 \in U$ . We will construct a relatively clopen subset of  $J_m(f_a)$  which contains  $z_0$  and is contained in  $U$ . To that end, let  $C$  be the connected component of  $z_0$  in  $J(f_a)$ . Let  $S = C \cap \partial U$ . By Proposition 4 and [7, Proposition 2.4(b),(c)] we have

$$S \subset J(f_a) \setminus E(f_a) \subset A(f_a).$$

Let  $\delta$  and  $\mathcal{M}$  be given by Proposition 5. Let  $\lambda > 0$  be given by Lemma 6, and for each  $n \in \mathbb{N}$  define

$$S_n = \{s \in S : |f_a^n(s)| > 2|f_a^n(z_0)| + \lambda + \delta \text{ and } \operatorname{Re}(f_a^n(s)) > 0\}.$$

By Lemma 7 and compactness of  $S$ , there is a finite  $\mathcal{F} \subset \mathbb{N}$  such that  $S \subset \bigcup\{S_n : n \in \mathcal{F}\}$ . For each  $n \in \mathcal{F}$ , Lemma 6 provides an open set  $V_n \subset \mathbb{C}$  such that:

- $f_a^n(z_0) \in V_n$ ,
- $\partial V_n \cap J(f_a) \subset A_R(f_a)$ , and
- $\sup\{\operatorname{Re}(z) : z \in V_n\} \leq |f_a^n(z_0)| + \lambda$ .

Notice that we can shrink  $V_n$  to obtain

$$\sup\{|\operatorname{Im}(z) - \operatorname{Im}(f_a^n(z_0))| : z \in V_n\} \leq \delta$$

in addition to the three properties above; simply intersect  $V_n$  with the connected component of  $f_a^n(z_0)$  in  $\mathbb{C} \setminus \mathcal{M}$ . Now if  $z \in V_n$  and  $\operatorname{Re}(z) > 0$ , then

$$|z| \leq \operatorname{Re}(z) + |\operatorname{Im}(z)| \leq 2|f_a^n(z_0)| + \lambda + \delta.$$

Therefore  $f_a^n(S_n) \cap \overline{V_n} = \emptyset$ . So  $S_n \cap \overline{f_a^{-n}(V_n)} = \emptyset$ .

Let  $V = \bigcap\{f_a^{-n}(V_n) : n \in \mathcal{F}\}$ . Then  $V$  is open, and

$$S \cap \overline{V} = \emptyset. \tag{5.1}$$

Further,  $\partial f_a^{-n}(V_n) \cap J(f_a) \subset f_a^{-n}[A_R(f_a)] \subset A(f_a)$  implies

$$\partial V \cap J(f_a) \subset A(f_a). \tag{5.2}$$

By Proposition 3, there is a collection of open sets  $\{O_n : n \in \mathbb{N}\}$  such that

$$\partial O_n \subset F(f_a), \tag{5.3}$$

$$\overline{O_n} \supset \overline{O_{n+1}}, \text{ and} \tag{5.4}$$

$$\bigcap_{n \in \mathbb{N}} \overline{O_n} \cap J(f_a) = C. \tag{5.5}$$

By (5.1), (5.4), (5.5), and compactness of  $\overline{O_n} \cap \overline{V} \cap \partial U \cap J(f_a)$ , there exists  $n \in \mathbb{N}$  such that

$$\overline{O_n} \cap \overline{V} \cap \partial U \cap J(f_a) = \emptyset. \tag{5.6}$$

Then  $O_n \cap V \cap U \cap J_m(f_a)$  is a relatively clopen subset of  $J_m(f_a)$  (it is closed in  $J_m(f_a)$  by (5.2), (5.3), and (5.6)). It contains  $z_0$  and is contained in  $U$ .  $\square$

Now the topological type of  $J_m(f_a)$  can be determined using the Alexandroff-Urysohn characterization of the irrationals. The Alexandroff-Urysohn characterization [16, Theorem 1.9.8] states that every topologically complete, nowhere compact, zero-dimensional space is homeomorphic to  $\mathbb{P}$ .

A space  $X$  is *topologically complete* if there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space that is homeomorphic to  $X$  in its original topology.  $G_\delta$ -subsets of  $\mathbb{C}$  are topologically complete by [16, Theorem A.6.3].

A space  $X$  is *nowhere compact* if every compact subset of  $X$  has empty interior in  $X$ . We observe that if  $X \subset Y \subset \mathbb{C}$ ,  $X$  is dense in  $Y$ , and  $Y \setminus X$  is dense in

$Y$ , then  $X$  is nowhere compact. For let  $K$  be any compact subset of  $X$  and let  $x \in K$ . If  $Y \setminus X$  is dense in  $Y$ , then there is a sequence of points  $y_1, y_2, \dots \in Y \setminus K$  converging to  $x$ . By compactness of  $K$  we have that  $Y \setminus K$  is open in  $Y$ . Thus, assuming  $X$  is dense in  $Y$ , there is a sequence of points  $x_1, x_2, \dots \in X \setminus K$  with  $d(x_n, y_n) < \frac{1}{n}$ . Then  $x_n \rightarrow x$ , proving that  $K$  is not a neighborhood of  $x$  in  $X$ . Since  $x$  was an arbitrary point of  $K$ , we conclude that  $K$  has empty interior in  $X$ .

**Corollary 9.** *If  $a \in F(f_a)$  then  $J_m(f_a) \simeq J_m(f_a) \cup \{\infty\} \simeq \mathbb{P}$ .*

*Proof.* The Julia set  $J(f_a)$  is closed in  $\mathbb{C}$ , and

$$A(f_a) = \bigcup_{\ell \in \mathbb{N}} f_a^{-\ell}[A_R(f_a)]$$

is  $F_\sigma$  in  $\mathbb{C}$ . So  $J_m(f_a)$  is a  $G_\delta$ -subset of  $\mathbb{C}$  and is therefore topologically complete. The sets  $J_m(f_a)$  and  $J(f_a) \setminus J_m(f_a) (= A(f_a))$  are infinite and completely invariant under  $f_a$ , so by Montel's theorem they are both dense in  $J(f_a)$ . By the observation above (with  $Y = J(f_a)$  and  $X = J_m(f_a)$ ),  $J_m(f_a)$  is nowhere compact. Finally,  $J_m(f_a)$  is zero-dimensional by Theorem 8. By [16, Theorem 1.9.8] we have  $J_m(f_a) \simeq \mathbb{P}$ . The one-point extension  $J_m(f_a) \cup \{\infty\}$  is also zero-dimensional by Theorem 8 and [1, Theorem 3.11]. Hence the characterization [16, Theorem 1.9.8] can be applied again to see that  $J_m(f_a) \cup \{\infty\} \simeq \mathbb{P}$ .  $\square$

**Corollary 10.** *If  $a \in F(f_a)$  then  $J_r(f_a)$  and  $J_r(f_a) \cup \{\infty\}$  are zero-dimensional.*

*Proof.* This follows immediately from Corollary 9 and the fact  $J_r(f_a) \subset J_m(f_a)$ .  $\square$

## 6. FATOU'S FUNCTION

The function defined by  $f(z) = z + 1 + e^{-z}$  is called *Fatou's function* [8]. Here it is known that  $J_r(f) = \mathbb{C} \setminus I(f) = J(f) \setminus I(f) \subset J_m(f)$  [citation to appear].

Previous work has shown that  $J_r(f) \cup \{\infty\}$  and  $J_m(f) \cup \{\infty\}$  are totally separated; see [6, Theorem 5.2] and [7, Theorem 5.1]. The corollaries below, together with [1, Theorem 3.11] (zero-dimensionality and one-point extensions), imply that both of these sets are in fact zero-dimensional.

**Corollary 11.**  *$J_m(f)$  is homeomorphic to  $\mathbb{P}$ .*

*Proof.* Define  $h : \mathbb{C} \rightarrow \mathbb{C}$  by  $h(z) = e^{-1}ze^{-z}$ . The proof of [7, Theorem 5.1] shows that  $J_m(f)$  is covered by a discrete collection of homeomorphic copies of  $J_m(h)$ . And  $J_m(h) \simeq J_m(f_{-2})$  by [7, Proposition 5.2]. We can now apply Theorem 8 with  $a = -2$  to see that  $J_m(f)$  is zero-dimensional. Further, the proof of Corollary 9 shows that  $J_m(f) \simeq \mathbb{P}$ .  $\square$

**Corollary 12.**  *$J_r(f)$  is zero-dimensional.*

*Proof.* This follows from Corollary 11 and the fact  $J_r(f) \subset J_m(f)$ .  $\square$

**Acknowledgements.** We thank the referee for their careful reading and suggestions which greatly improved the paper.



## REFERENCES

- [1] J. M. Aarts and T. Nishiura, *Dimension and Extensions*, North-Holland Mathematical Library, vol. 48, NorthHolland Publishing Co, Amsterdam, 1993.
- [2] J. M. Aarts and L. G. Oversteegen, The geometry of Julia sets, *Trans. Amer. Math. Soc.* 338 (1993), no. 2, 897–918.
- [3] N. Alhabib and L. Rempe-Gillen, Escaping Endpoints Explode. *Comput. Methods Funct. Theory* 17, 1 (2017), 65–100.
- [4] W. Bergweiler and A. Hinkkanen, On semiconjugation of entire functions, *Math. Proc. Cambridge Philos. Soc.* 126 (1999), no. 3, 565–574.
- [5] A. E. Eremenko and M. Y. Lyubich, Dynamical properties of some classes of entire functions, *Ann. Inst. Fourier (Grenoble)* 42 (1992), no. 4, 989–1020.
- [6] V. Evdoridou, Fatou’s web, *Proc. Amer. Math. Soc.* 144 (2016), no. 12, 5227–5240.
- [7] V. Evdoridou and L. Rempe-Gillen, Non-escaping endpoints do not explode. *Bulletin of the London Mathematical Society*, 50(5) (2018) pp. 916–932.
- [8] P. Fatou, Sur l’itération des fonctions transcendantes entières (French), *Acta Math.* 47 (1926), no. 4, 337–370.
- [9] B. Karpinska, Area and Hausdorff dimension of the set of accessible points of the Julia sets of  $\lambda e^z$  and  $\lambda \sin z$ , *Fund. Math.* 159 (1999), 269–287.
- [10] C. T. McMullen, Hausdorff dimension and conformal dynamics II. Geometrically finite rational maps, *Comment. Math. Helv.* 75 (2000), no. 4, 535–593.
- [11] L. Rempe, Topological dynamics of exponential maps on their escaping sets. *Ergod. Theory Dyn. Syst.* 26(6) (2006), 1939–1975.
- [12] P. J. Rippon and G. M. Stallard, Fast escaping points of entire functions, *Proc. London Math. Soc.*, 105 (2012), 787–820.
- [13] D. Schleicher and J. Zimmer, Periodic points and dynamic rays of exponential maps. *Ann. Acad. Scient. Fenn.* 28 2 (2003), 327–354.
- [14] M. Urbański. On some aspects of fractal dimensions in higher dimensional dynamics. In *Proceedings of Workshop: Problems in higher dimensional dynamics*, pages 18–25. Preprint SFB 170 Göttingen, 1995.
- [15] M. Urbański and A. Zdunik, The finer geometry and dynamics of the hyperbolic exponential family, *Michigan Math. J.* 51 (2003), no. 2, 227–250.
- [16] J. van Mill, *The Infinite-Dimensional Topology of Function Spaces*, North-Holland Publishing Co., Amsterdam, 2001.

DEPARTMENT OF MATHEMATICS, AUBURN UNIVERSITY AT MONTGOMERY, MONTGOMERY AL 36117, UNITED STATES OF AMERICA  
 Email address: ds10003@auburn.edu