

Energetics of Large Water Clusters [up to (H₂O)₂₀] by means of Explicitly Correlated, Localized Coupled Cluster Methods

Nitai Sylvetsky[§] and Jan M. L. Martin^{,§}*

Revised manuscript CPLETT-20-280

[§] Department of Organic Chemistry, Weizmann Institute of Science, 76100 Rehovot, Israel.

Email: gershom@weizmann.ac.il FAX: +972 8 934-3029

KEYWORDS water clusters • explicitly correlated methods • localized coupled cluster • canonical coupled cluster

ABSTRACT We consider the performance of combined PNO-F12 approaches for the interaction energies of water clusters as large as (H₂O)₂₀ by comparison to canonical CCSD(T)/CBS reference values obtained through n-body decomposition of post-MP2 corrections. We find that PNO-LCCSD(T)-F12b approaches with “Tight” cutoffs are generally capable of reproducing canonical CCSD(T) interaction energies to within ~0.25% and isomerization energies to ~1.5%, while requiring only a fraction of the canonical computational cost. However, basis set convergence patterns and effect of counterpoise corrections are more erratic than for canonical calculations, highlighting the need for canonical benchmarks on closely related systems.

Introduction

Noncovalent interactions (NCIs) are well-known to significantly influence various physical and chemical properties of many (supra)molecular systems.[1,2] In biochemistry, for instance, NCIs are known to play important roles in protein folding, protein-ligand interaction, and nucleobase packing and stacking.[3,4] They are also of great importance in various chemical contexts such as organocatalysis, supramolecular chemistry, and

conformational stability. That being said, individual NCIs (which may amount to as little as a few tenths of a kcal/mol; see, e.g., Ref.[5]) are extremely difficult to measure experimentally. For this reason, systematically-convergent wavefunction *ab initio* methods constitute a crucial, well-nigh exclusive source of information on such individual NCIs. As such, these methods are extensively used for calibration of approximate computational approaches, such as molecular mechanics force fields, semiempirical MO methods, and DFT functionals. Biochemically relevant NCIs are generally taking place in aqueous solutions. NCIs between biomolecules and water molecules, and between the latter among themselves, are therefore essential for understanding vital biochemical mechanisms, such as binding of substrates/inhibitors to their corresponding enzymes, maintaining 3-dimensional structures of organic complexes, and so forth. In this context, water clusters have long been the subject of basic scientific interest because they dictate water's singular bulk properties (including but not limited to a remarkably high boiling point, low thermal expansion coefficient, and unusual density behavior). Moreover, they have also been expected to play important roles in hydration patterns of small- and medium-sized molecules – as well as in hydration of small ions.[6,7] Of particular interest are cyclic clusters, which have been shown to provide important insights regarding the significance of many-body interactions in bulk water.[8–10] Thus, a great many experimental and theoretical studies have been carried out on the structure and properties of water in both the gas and liquid phases,[11–15] and wavefunction *ab initio* methods have already provided valuable insight into the structures and energetics of small water clusters.[16–20] A few representative reviews of diverse computational and experimental studies of water clusters can be found in Refs.[19,21–23]

One of our group's recent ventures[24] was the re-evaluation of the water cluster benchmarks BEGDB[25] and WATER27[26] using conventional and explicitly-correlated (F12) coupled cluster methods: the latter were effectively used to accelerate the notoriously slow 1-particle basis set convergence for the MP2 and CCSD energetic components (see Refs.[27–30] for reviews on F12 methodology). Canonical CCSD(T)-F12b calculations on larger water clusters such as (H₂O)₂₀ – which has been studied extensively[19,21–23,31,32] for representing a transition point between the four Wales-Hodges families of three-dimensional structures (i.e., single cage, box-kite, edge-sharing- and face-sharing-prisms)[33] – are, however, currently beyond the reach of commodity high-performance computing (HPC) hardware, except for woefully inadequate small basis sets like aVDZ.

For this reason, pair natural orbital (PNO) local correlation approaches[34–37] may constitute an attractive, affordable alternative. PNO-type approaches may additionally be

combined with the F12 ansatz in order to provide the best of both worlds –accelerated basis set convergence as well as nearly-linear CPU time scaling with system size.

Despite enabling *ab initio* methods to ever-larger systems of interest, practical implementations of both PNO-type and F12 approaches involve many cutoffs, screening thresholds, and other details ‘under the hood’; these are hidden from the end user as tuned collections invoked through keywords like, e.g. in ORCA,[38] LoosePNO, NormalPNO (the default), and TightPNO: for details see Table 1 of Ref.[39] (The MOLPRO equivalents are specified in Tables 1-4 of Ref.[37] and the accompanying discussion.) For this reason, and especially in chemical territories where PNO-type approaches have not yet proven to be consistent and robust, comparing calculated results with a canonical benchmark is necessary to ensure the latter’s reliability.

For water clusters, or indeed any cluster A_n for large enough n , such cutoffs put an additional fly in the ointment. While they do lead to substantial pruning in A_n , the monomer A is so small that nothing is screened out and the PNO calculation becomes functionally equivalent to a canonical one. This creates an intrinsic imbalance for a cluster interaction energy.

So how do we validate our PNO calculations for clusters as large as $(\text{H}_2\text{O})_{20}$? As stated above, canonical CCSD(T) calculations in adequate basis sets are not a practical option (or we need not resort to PNO in the first place). However, we learned from our earlier work[24] that in an n -body expansion (see Methods section below), the high-level correction, $\text{HLC} \equiv [\text{CCSD(T)} - \text{MP2}]$, converges fairly rapidly with n . Hence, a very good approximation to the canonical CCSD(T)/CBS energetics can be obtained from combining canonical whole-system MP2-F12/CBS with at most 4-body HLCs. Such a calculation involves thousands of single-point CCSD(T) energy calculations, but on at most tetramers, and of course it is embarrassingly parallel. This offers us a practical route toward nearly exact canonical benchmark results as a touchstone for the PNO approaches.

For smaller clusters like $(\text{H}_2\text{O})_6$ – which is of particular interest as $n=6$ represents the transition point between cyclic and three-dimensional structures[31] – we of course need not resort to such approximations. In the present paper, interaction energies of water clusters from the BEGDB[25] and WATER27[26] datasets will be considered in ascending order of size: (a) eight $(\text{H}_2\text{O})_6$ structures, for which we were able to obtain canonical CCSD (T)/CBS reference values; (b) one $(\text{H}_2\text{O})_{10}$ system (where we were able to improve on the previous reference value [24]); and (c) four $(\text{H}_2\text{O})_{20}$ clusters.

Computational Methods

All calculations were carried out on the Faculty of Chemistry's Linux cluster 'chemfarm' at the Weizmann Institute of Science. Single-point PNO-LCCSD(T)-F12b[37,40] calculations were carried out using default and "domopt=tight" options, as defined in Ref.[37] and implemented in the Molpro2019 program package.[41] (We note that L in the Werner group's notation refers to "localized" and not to linearized coupled cluster theory.) All PNO-based calculations discussed in this paper employ tight PNO domains unless explicitly stated otherwise.

CCSD(T)-F12b[42,43] and DF-MP2-F12[44] calculations were also used for the purpose of obtaining reliable canonical reference values.

For the explicitly correlated [i.e., CCSD(T)-F12b and PNO-LCCSD(T)-F12b] calculations, the correlation-consistent cc-pVnZ-F12 basis sets (a.k.a VnZ-F12) of Peterson et al.[45] were used in conjunction with the appropriate auxiliary basis sets for JKfit[46] (Coulomb and exchange), MP2fit[47,48] (density fitting in MP2), and OptRI[49,50] (complementary auxiliary basis set, CABS) basis sets. We also employed our own aug-cc-pVnZ-F12 (or aVnZ-F12) basis sets, introduced in Ref.[51]; the issue of the appropriate CABS basis set in such calculations is investigated in detail in Ref.[52] As recommended in Ref. [53], the geminal exponent (β) value was set to 1.0 for all basis sets used in explicitly-correlated calculations under consideration.

Conventional (i.e., orbital-based) *ab initio* CCSD(T)[54,55] calculations were performed using correlation-consistent[56] basis sets. In general, we used the combination of diffuse function-augmented basis sets aug-cc-pVnZ ($n = T, Q, 5$) on non-hydrogen atoms and regular cc-pVnZ basis sets on hydrogen – to be denoted haVnZ for short.

Basis set extrapolations were carried out using the two-point formula:

$$E_{\infty} = E(L) - [E(L) - E(L-1)] \left[\left(\frac{L}{L-1} \right)^{\alpha} - 1 \right] \quad (1)$$

where L is the highest angular momentum present in the basis set for elements B–Ne and Al–Ar and α an exponent specific to the level of theory and basis set pair. The basis set extrapolation exponents (α) were taken from Table 2 of Ref.[57].

Aside from Boys-Bernardi counterpoise corrections[58] and the uncorrected values, we also apply the average of both (so-called "half-CP"), as rationalized by Sherrill and coworkers[59] for orbital-based *ab initio* methods and by our group[60] for F12 calculations. In short, such practice exploits the balance between basis set superposition error (which causes overbinding) and intrinsic basis set incompleteness (which causes underbinding).[60]

As mentioned in the introduction, decomposing water cluster interaction energies (for which MP2 is a good approximation) into MP2 and HLC, and then applying an n-body expansion to the latter, may offer a route towards accurate energetics for large water clusters, where all-atom HLCs are computationally too costly.[24,61] (For brief overviews of the n-body decomposition scheme, see Refs.[62,63].) Indeed, multiple levels of theory may be nested, with the 2-body HLCs treated at the highest level, the 3-body HLCs at a lower one, etc. By way of illustration, our reference value for (H₂O)₁₀ below consists of:

$$\begin{aligned} \text{REF}_{\text{water}10} = & \text{half-CP-MP2-F12/a'V}\{T,Q\}\text{-F12} \{ \text{half-CP, whole system} \} \\ & + 2\text{-body}([\text{CCSD}(T) - \text{MP2}] / \text{AV}\{Q,5\}Z) + 3\text{-body}([\text{CCSD}(T) - \text{MP2}] / \text{AV}\{T,Q\}Z) \\ & + 4 \dots 10\text{-body}([\text{CCSD}(T) - \text{MP2}] / \text{AVTZ}) \end{aligned} \quad (2)$$

(We note that this approach is philosophically very similar to that of Ref.[64]) For the avoidance of doubt, all calculated interaction energies considered in this work are “vertical” – that is, the isolated monomer geometries are the same as those found within the cluster, and the interaction energy does not include monomer relaxation terms. To facilitate comparisons with earlier work, reference geometries were taken verbatim from the BEGDB[25] and WATER27[26] datasets and not optimized further.

Results and Discussion

a. (H₂O)₆ structures

For our reference level, we were able to obtain both CCSD(T)/A'V{Q,5}Z and [CCSD-F12b/a'V{Q,5}Z-F12 + (T)/A'V{Q,5}Z] half-CP energetics for the hexamers under consideration. However, as can be seen in Table S1 (found in the ESI), the latter turned out to be virtually indistinguishable from the former (± 0.01 kcal/mol RMSD) – which indicates that 1-particle basis set convergence has been achieved. The RMSD from the best values in the earlier benchmark paper[24] is just 0.08 kcal/mol; this is probably a conservative estimate for the residual uncertainty in our reference values.

Error statistics for the (H₂O)₆ structures (Table 1) reveal that PNO-based methods are capable of reproducing canonical results to within less than 0.1 kcal/mol (arguably within the uncertainty of the reference values). However, it can be seen that basis set convergence patterns are quite erratic. Using both VnZ-F12 and a'VnZ-F12 basis set families, neither CP corrections nor basis set extrapolations consistently reduce error. (Indeed, since basis set extrapolation is itself subject to a certain degree of empiricism, some might argue that PNO-

based methods require a dedicated set of extrapolation exponents for the procedure to be fruitful. On the flip side, strong sensitivity to extrapolation exponents is in our experience a sign that the basis sets being extrapolated from are still too far from the CBS limit.[65,66])

RMSD							
Raw							
n=	T	T*	Q	Q*	5	{T/Q}	{Q/5}
VnZ-F12	0.102	0.228	0.063	0.024	-	0.109	-
a'VnZ-F12	0.024	0.313	0.173	0.021	-	0.240	-
conv. haVnZ	0.099	-	0.030	-	0.150	0.032	0.249
CP							
n=	T	T*	Q	Q*	5	{T/Q}	{Q/5}
VnZ-F12	0.507	0.389	0.242	0.187	-	0.164	-
a'VnZ-F12	0.390	-	0.206	-	-	0.140	-
conv. haVnZ	3.152	-	1.188	-	0.609	0.247	0.139
half-CP							
n=	T	T*	Q	Q*	5	{T/Q}	{Q/5}
VnZ-F12	0.206	0.097	0.151	0.095	-	0.135	-
a'VnZ-F12	0.189	-	0.189	-	-	0.189	-
conv. haVnZ	1.615	-	0.608	-	0.379	0.130	0.194

MSD							
Raw							
n=	T	T*	Q	Q*	5	{T/Q}	{Q/5}
VnZ-F12	0.098	0.227	-0.057	0.000	-	-0.104	-
a'VnZ-F12	0.013	0.313	-0.171	0.019	-	-0.238	-
conv. haVnZ	-0.080	-	-0.027	-	-0.149	0.011	-0.247
CP							
n=	T	T*	Q	Q*	5	{T/Q}	{Q/5}
VnZ-F12	-0.506	-0.388	-0.187	-0.187	-	-0.163	-
a'VnZ-F12	-0.389	-	-0.206	-	-	-0.139	-
conv. haVnZ	-3.149	-	-1.186	-	-0.608	0.247	-0.139
half-CP							
n=	T	T*	Q	Q*	5	{T/Q}	{Q/5}
VnZ-F12	-0.204	-0.095	-0.150	-0.094	-	-0.133	-
a'VnZ-F12	-0.188	-	-0.188	-	-	-0.188	-
conv. haVnZ	-1.615	-	-0.607	-	-0.378	0.129	-0.193

* = Constant scaling of triples (Ts), cf. Table 3 of Ref.[67]

Table 1. Eight isomers of (H₂O)₆: Error statistics (kcal/mol) for PNO-LCCSD(T)-F12b interaction energies obtained using various basis sets; half-CP corrected, canonical CCSD(T)/A'V{Q,5}Z is used as a reference.

However, it should also be kept in mind that the (T) contributions — which we have seen in past work[24] are quite nontrivial — do not benefit at all from F12 as (T) is a 3-body, not a 2-body effect. Marchetti and Werner[68] in the past proposed scaling of the (T) contribution by the $E_{\text{corr}}[\text{MP2-F12}]/E_{\text{corr}}[\text{MP2}]$ ratio as a workaround. In Ref.[67] we considered both this approach, and scaling by the $E_{\text{corr}}[\text{CCSD-F12b}]/E_{\text{corr}}[\text{CCSD}]$ ratio, and found that constant scaling by a fixed ratio determined for a small training set [given in Table 3 of Ref.[67], denoted (Ts) there] yields better results for thermochemistry (besides being trivially size-consistent). For the problem at hand, triples scaling (denoted by an asterisk) does seem to offer an improvement over unscaled results – with the exception of raw triple-zeta options, which are arguably too far from the basis set limit anyway (thus, unscaled results obtained in this manner seem to benefit from a fortuitous error compensation). Indeed, that (Ts) scaling of the largest basis set is more effective than basis set extrapolations for PNO-F12 methods may be a useful finding with broader implications for thermochemistry of larger systems. We intend to explore this issue further in a future paper.

Ineffectiveness of CP corrections may be associated with the fundamental principles on which PNO methodology rests: while ‘strong pairs’ that exceed a certain energetic cutoff, fully benefit from the availability of added basis functions, ‘weak pairs’ are only treated at a lower level of theory, or are simply neglected, thus not making optimal use of the added functions and creating an imbalance for the overall calculated energy. Unlike for the canonical F12 case,[51] where a’VnZ-F12 offers considerably faster and smoother basis set convergence than VnZ-F12, we see no obvious advantage for PNO-F12 calculation on (H₂O)₆ isomers. This can, however, be an artifact of the smallness of these clusters (see below). Mean signed deviations for the methods under consideration indicate that PNO-based results are generally underbound relative to canonical reference values (except for VTZ-F12, a’VTZ-F12 and scaled VQZ-F12 and a’VQZ-F12 raw interaction energies).

b. (H₂O)₁₀

Unlike for the hexamers, fully canonical CCSD(T) energetics with sufficiently large basis sets were previously found to be computationally prohibitive.[24] Thus, we have employed the n-body decomposition scheme in Eq.(2). Our best calculated interaction energy, 98.50 kcal/mol, is somewhat lower than the 99.24 kcal/mol found in Ref.[24] where because of computational resource limitations, the (T) contribution was extrapolated from relatively small a’V{D,T}Z basis sets.

Results obtained for $(\text{H}_2\text{O})_{10}$ (Table 2) display a somewhat similar picture to $(\text{H}_2\text{O})_6$, except that this time around, a'VnZ-F12 seems to be generally preferable over VnZ-F12. Like for the hexamers, it can be seen that PNO-based methods can reproduce the canonical values to within ~ 0.1 kcal/mol (equivalent to $\sim 0.1\%$ percent of the overall interaction, which amounts to 98.5 kcal/mol). In this case, however, some of the errors – and specifically those for the raw interaction energies – are larger in magnitude. This fact illustrates the thermochemical imbalance in PNO association energies pointed out in the introduction: This problem is expected to be exacerbated for A_n clusters as n increases. Unlike for $(\text{H}_2\text{O})_6$, mean signed deviations testify that PNO-based results are, in fact, overbound in this case (except for the CP-corrected VTZ-F12 and a'VTZ-F12 options); unfortunately, this finding precludes using an *ad hoc*, *a posteriori* correction for PNO-based results – such as a constant scaling factor – in order to eliminate biases from corresponding canonical limits. Again, basis set extrapolations do not seem to bring us closer to the canonical reference, and while (Ts) scaling does offer an improvement for some CP-corrected results, it is not as useful since values of comparable quality may be obtained using CP-uncorrected (and therefore less computationally costly) alternatives. (For instance, raw a'VQZ-F12 is of similar quality to scaled, CP-corrected quadruple zeta options).

raw					
n=	T	T*	Q	Q*	{T/Q}
VnZ-F12	0.561	0.856	0.412	0.543	0.304
a'VnZ-F12	0.409	0.757	0.204	0.358	0.054

CP					
n=	T	T*	Q	Q*	{T/Q}
VnZ-F12	-0.505	-0.233	0.035	0.162	0.429
a'VnZ-F12	-0.273	0.053	0.094	0.245	0.362

half-CP					
n=	T	T*	Q	Q*	{T/Q}
VnZ-F12	0.028	0.311	0.224	0.353	0.367
a'VnZ-F12	0.068	0.405	0.149	0.302	0.208

* = Constant scaling of triples (Ts), cf. Table 3 of Ref.[67]

Table 2. $(\text{H}_2\text{O})_{10}$: Signed deviation (kcal/mol) for PNO-LCCSD(T)-F12b interaction energies obtained using various basis sets; $\text{REF}_{\text{water}10}$, eq. (2), is used as a reference.

c. $(\text{H}_2\text{O})_{20}$ structures

As for our n-body-based strategy used for $(\text{H}_2\text{O})_{10}$, we have managed to obtain benchmark, fully-canonical results using the following level of theory:

$$\begin{aligned}
\text{REF}_{\text{water20}} = & \text{MP2-F12/a}'V\{T,Q\}\text{-F12 } \{\text{raw, whole system}\} \\
& + 2\text{-body}([\text{CCSD-F12b} - \text{MP2-F12}]/\text{a}'VQZ\text{-F12} + (\text{T})/\text{AV5Z}) \\
& + 3\text{-body}([\text{CCSD-F12b} - \text{MP2-F12}]/\text{a}'VTZ\text{-F12} + (\text{T})/\text{a}'V\{D,T\}Z) \\
& + 4\text{-body}([\text{CCSD(T)} - \text{MP2}]/\text{a}'VTZ) \tag{3}
\end{aligned}$$

As can be seen in Table S2 in the ESI, this reference level represents an improvement over the one previously established in Ref.[24] (see ESI-2 of same paper), which corresponds to:

$$\begin{aligned}
\text{REFOLD}_{\text{water20}} = & \text{MP2-F12/V}\{T,Q\}\text{-F12 } \{\text{raw, whole system}\} \\
& + 2\text{-body}([\text{CCSD-F12b} - \text{MP2-F12}]/VQZ\text{-F12} + (\text{T})/\text{A}'V\{T,Q\}Z) \\
& + 3\text{-body}([\text{CCSD-F12b} - \text{MP2-F12}]/VTZ\text{-F12} + (\text{T})/\text{a}'V\{D,T\}Z) \tag{4}
\end{aligned}$$

which neglected 4-body HLC contributions entirely. Primarily because of the latter, the new total interaction energies differ by 0.37 kcal/mol RMS from those given in Ref.[24]. Our best values are: edge-sharing pentagonal 219.19, face-sharing cubes 215.98, face-sharing pentagonal 217.03, and dodecahedron 211.58 kcal/mol, respectively. We conservatively assign an uncertainty of about 0.4 kcal/mol to these new reference values.

Error statistics for the four structures of $(\text{H}_2\text{O})_{20}$ (Table 3) further confirm our hypothesis regarding accumulation of errors in PNO-based calculations, depending on chosen PNO cutoffs – as some of the errors for the raw calculations now approach 1 kcal/mol.

RMSD						MSD					
raw						raw					
n=	T	T*	Q	Q*	{T/Q}	n=	T	T*	Q	Q*	{T/Q}
VnZ-F12	0.309	0.430	0.504	0.225	0.533	VnZ-F12	-0.243	0.398	-0.494	-0.209	-0.531
a'VnZ-F12	0.511	0.271	0.942	0.605	1.092	a'VnZ-F12	-0.498	0.258	-0.940	-0.603	-1.091
CP						CP					
n=	T	T*	Q	Q*	{T/Q}	n=	T	T*	Q	Q*	{T/Q}
VnZ-F12	2.556	1.408	1.323	0.756	0.641	VnZ-F12	-2.553	-1.406	-1.322	-0.755	-0.641
a'VnZ-F12	1.938	0.764	1.190	0.619	0.753	a'VnZ-F12	-1.935	-0.76	-1.19	-0.62	-0.752
half-CP						half-CP					
n=	T	T*	Q	Q*	{T/Q}	n=	T	T*	Q	Q*	{T/Q}
VnZ-F12	1.407	0.518	0.911	0.485	0.586	VnZ-F12	-1.398	-0.504	-0.908	-0.482	-0.586
a'VnZ-F12	1.221	0.261	1.065	0.612	0.922	a'VnZ-F12	-1.216	-0.25	-1.06	-0.61	-0.922

* = Constant scaling of triples (Ts), cf. Table 3 of Ref.[67]

Table 3. $(\text{H}_2\text{O})_{20}$: Error statistics (kcal/mol) for PNO-LCCSD(T)-F12b interaction energies obtained using various basis sets; $\text{Ref}_{\text{water20}}$, Eq. (3), is used as a reference.

Nevertheless, PNO-based methods can still be used to reproduce canonical reference values to within 0.5 kcal/mol, which at about 0.25% of the cluster association energy might be deemed negligible in relative terms. Once again, CP corrections prove to be quite ineffectual, and do not justify the required additional computational cost (i.e., using the entire cluster's basis functions for calculating monomer energies). (Ts) scaling, on the other hand, does seem to improve most calculated results – even a'VTZ-F12, which is relatively compact and computationally economical, comes close to the reference. Thus, a PNO-LCCSD(T)-F12b calculation with tight PNO settings on the edge-sharing (H₂O)₂₀ structure took 6 days and 14 hours CPU time (with nearly perfect parallelization), whereas our n-body-based canonical reference calculation according to eq. (3) required more than a year (!) for the same system. This is indeed good news, as PNO-based methods can now be recognized as both remarkably economical and fairly accurate for the systems under consideration.

Would switching to default PNO domains come at a substantial cost in accuracy? As can be seen in Table 4, employing such settings leads to further under-binding: for the a'VnZ-F12 (n=T,Q) basis sets considered here, applying default PNO domains results in an error two to six times as large than that obtained using tight PNO settings. That being said, it should once more be noted that default PNO settings are more computationally economical, and may thus be preferable for larger systems for which tight PNOs are too demanding. Again, a PNO-LCCSD(T)-F12b calculation with tight PNO settings on the edge-sharing (H₂O)₂₀ structure required 6 days and 14 hours CPU time, compared to only 23 hours with default PNOs (6.843:1 ratio). This finding may indeed be useful for, say, still larger water clusters [such as (H₂O)₁₀₀]. However, the associated compromises on accuracy are not quite justified for the case under consideration.

RMSD

Raw

a'VnZ-F12	T	T*	Q	Q*	{T/Q}
tightDomain	0.361	0.191	0.666	0.428	0.772
default	1.689	1.203	1.716	1.494	1.641

* = Constant scaling of triples (Ts), cf. Table 3 of Ref.[67]

MSD

raw

a'VnZ-F12	T	T*	Q	Q*	{T/Q}
tightDomain	-0.498	0.258	-0.940	-0.603	-1.091
default	-2.381	-1.694	-2.421	-2.107	-2.316

Table 4. (H₂O)₂₀: Error statistics (kcal/mol) for PNO-LCCSD(T)-F12b interaction energies obtained using a'VnZ-F12 (n=T,Q) basis sets with tight and default PNO domains; Ref_{water20}, eq. (3), is used as a reference.

A reviewer wondered if perhaps the various truncation errors in the PNO result inject a random, rather than systematic, error component into *relative* energies of the different structural isomers. This would be a major downside of PNO methods if true, since for canonical approaches, relative energies of structures are well known to converge much faster and more smoothly with basis set and electron correlation approach than total interaction energies. As can be seen in Table 5 for the relative energies of the four (H₂O)₂₀ isomers, however, PNO-LCCSD(T)-F12b methods clearly reproduce canonical reference values without compromising much on accuracy: raw DLPNO-CCSD(T)-F12b/a'VTZ-F12 with default PNO domains underestimates the icos-ES difference by ~0.4 kcal/mol (or by ~5%), but makes significantly smaller errors for the FC-ES and FS-ES energetic gaps (0.05 and 0.02 kcal/mol, or ~1% and ~1.6% respectively). Switching to tight PNO domains and larger basis sets further reduces these errors – which amount to just 0.01 (icos-ES), 0.03 (FC-ES), and 0.09 (FS-ES) kcal/mol using raw DLPNO-CCSD(T)-F12b/a'V{T,Q}Z-F12 with tight PNO domains. Thus, we see that errors in relative energies obtained using PNO methods converge relatively smoothly to the canonical basis set limit – as previously observed for calculated interaction energies.

a'VnZ-F12	T	T*	Q	Q*	{T/Q}	Reference Values
tightDomain						
FC – ES	0.039	0.025	0.027	0.020	0.015	3.204
FS – ES	-0.056	-0.057	-0.037	-0.037	-0.032	2.151
icos – ES	-0.260	-0.187	-0.146	-0.113	-0.093	7.608
Default						
FC – ES	0.052	0.041	-0.043	-0.049	-0.096	
FS – ES	-0.021	-0.020	-0.089	-0.090	-0.126	
icos – ES	-0.422	-0.360	-0.414	-0.384	-0.378	

Table 5. (H₂O)₂₀: Signed deviations (kcal/mol) for PNO-LCCSD(T)-F12b relative isomer energies obtained using a'VnZ-F12 (n=T,Q) basis sets with tight and default PNO domains; canonical reference values are also given (upper right; Ref_{water20}, eq. (3), is used as the reference level).

Interaction energy for the edge-shared global minimum is 219.19 kcal/mol.

To sum up: we have seen that PNO-based methods can be used to reproduce accurate canonical interaction energies of water clusters up to (H₂O)₂₀. This further corroborates that such methods are a viable alternative to canonical calculations for calculating NCIs of biologically relevant systems. That being said, and since systematic convergence is perhaps *the* key benefit of wavefunction *ab initio* calculations, the somewhat erratic behavior of

PNO-based regarding CP corrections and basis set extrapolations is indeed somewhat troubling. Thus, we can again see that results obtained using such methods should be treated with some caution in cases where no calibration against canonical values is available.

Conclusions

In this paper, we have investigated the performance of combined PNO and F12 methodologies [i.e., PNO-LCCSD(T)-F12b] for reproducing high-quality, canonical CCSD(T)-based NCIs for water clusters as large as $(\text{H}_2\text{O})_{20}$. Based upon our analysis of all associated error statistics and computation times, we were able to draw the following conclusions:

- PNO-LCCSD(T)-F12b approaches are generally capable of reproducing canonical CCSD(T) results to within $\sim 0.25\%$. Deviations from the reference values are not systematically positive or negative in sign. This fact prevents using ad-hoc, *a posteriori* corrections such as employing constant scaling factors and the like.
- Basis set convergence patterns, however, are quite erratic. First, using both VnZ-F12 and a'VnZ-F12 families, neither CP corrections nor basis set extrapolations systematically improve results. Second, results of triple-zeta quality are sometimes fortuitously closer to the reference than quadruple-zeta ones – apparently due to different error compensations (e.g., intrinsic basis set insufficiency, F12 and PNO cutoffs, etc.). Third, constant (Ts) scaling of the triples contribution, as suggested in Ref. [67], is somewhat helpful when used with relatively-large basis sets (namely VQZ-F12 and a'VQZ-F12). Finally, unlike for canonical F12 methods alone, augmented basis sets (i.e., a'VnZ-F12) do not necessarily outperform their non-augmented counterparts (VnZ-F12) in PNO-F12 calculations. We tentatively attribute this to PNO domains being too small for using diffuse functions in an effective manner (the latter are designed to “capture” spatial regions found far from the nuclei, and are thus somewhat contradictory to the localized PNO strategy).
- As previously noted,[37,69] PNO-LCCSD(T)-F12b calculations require only a fraction of the computational cost associated with canonical CCSD(T). In our case, a PNO-LCCSD(T)-F12b calculation on the edge-sharing $(\text{H}_2\text{O})_{20}$ structure took 6 days and 14 hours CPU time, whereas the canonical reference calculation (MP2-F12/CBS + n-body-based HLC) required more than a year (!)

- Based on the error statistics for a VnZ-F12 ($n=T,Q$) $(\text{H}_2\text{O})_{20}$ energetics, employing default PNO domains leads to errors two to six times larger than those obtained using tight domains. Nevertheless, default domains are much more economical computationally, and may therefore be an option for larger systems where tight PNO domains are already impractical. (For instance, for the edge-sharing $(\text{H}_2\text{O})_{20}$ structure, tight PNO settings required ~ 7 times as much CPU time as the same calculation with default PNOs.)
- Despite what one might intuitively expect on account of potential PNO truncation imbalances, relative energies of larger water clusters such as the $(\text{H}_2\text{O})_{20}$ isomer do seem to be reliably reproduced.

The present paper shows that, despite balance issues between a cluster and its separate small monomers due to PNO truncation disproportionately affecting the former, PNO-LCCSD(T)-F12b nevertheless offers a path to reliable NCI energetics for larger water clusters (perhaps as large as $(\text{H}_2\text{O})_{100}$). The usefulness of combined PNO-F12 approaches for dimeric noncovalent interactions was previously shown in detail by Ma and Werner.[70]

In future work, we hope to apply PNO and F12 methodologies to large biochemical systems of interest, such as a 300-atom model of alcohol dehydrogenase.[71] Energy decomposition schemes, such as the recently-proposed LED (local energy decomposition[72]) will be applied for interpretative purposes.

Acknowledgments

Research at Weizmann was funded by the Israel Science Foundation (grant 1358/15) and by the Estate of Emile Mimran (Weizmann), while computational resources and services were provided by Chemfarm (the Weizmann Institute Faculty of Chemistry HPC facility). NS acknowledges a doctoral fellowship from the Feinberg Graduate School at the Weizmann Institute. The authors would like to thank Profs. Frank Neese and Hans-Joachim Werner for helpful discussions.

Supporting information

Our best estimates for all total interaction energies considered in this work are provided in a dedicated spreadsheet, alongside Tables S1-S2. In addition, water cluster geometries used in our calculations are given in .xyz format; example input files for PNO-

LCCSD(T)F12b/a'VQZ-F12 calculations, as well as those required for establishing our reference level, are included for the H₂O₂₀es structure.

References

- [1] K.E. Riley, M. Pitoňák, P. Jurečka, P. Hobza, Stabilization and Structure Calculations for Noncovalent Interactions in Extended Molecular Systems Based on Wave Function and Density Functional Theories, *Chem. Rev.* 110 (2010) 5023–5063. <https://doi.org/10.1021/cr1000173>.
- [2] J. Řezáč, P. Hobza, Benchmark Calculations of Interaction Energies in Noncovalent Complexes and Their Applications, *Chem. Rev.* 116 (2016) 5038–5071. <https://doi.org/10.1021/acs.chemrev.5b00526>.
- [3] K.E. Riley, P. Hobza, On the importance and origin of aromatic interactions in chemistry and biodisciplines., *Acc. Chem. Res.* 46 (2013) 927–36. <https://doi.org/10.1021/ar300083h>.
- [4] K.E. Riley, P. Hobza, Noncovalent interactions in biochemistry, *Wiley Interdiscip. Rev. Comput. Mol. Sci.* 1 (2011) 3–17. <https://doi.org/10.1002/wcms.8>.
- [5] M.K. Kesharwani, D. Manna, N. Sylvetsky, J.M.L. Martin, The X40×10 Halogen Bonding Benchmark Revisited: Surprising Importance of (n – 1)d Subvalence Correlation, *J. Phys. Chem. A.* 122 (2018) 2184–2197. <https://doi.org/10.1021/acs.jpca.7b10958>.
- [6] A.D. Kulkarni, K. Babu, S.R. Gadre, L.J. Bartolotti, Exploring Hydration Patterns of Aldehydes and Amides: Ab Initio Investigations, *J. Phys. Chem. A.* 108 (2004) 2492–2498. <https://doi.org/10.1021/jp0368886>.
- [7] A.D. Kulkarni, S.R. Gadre, S. Nagase, Quantum chemical and electrostatic studies of anionic water clusters, *J. Mol. Struct. THEOCHEM.* 851 (2008) 213–219. <https://doi.org/10.1016/j.theochem.2007.11.019>.
- [8] S. Maheshwary, N. Patel, N. Sathyamurthy, A.D. Kulkarni, S.R. Gadre, Structure and Stability of Water Clusters (H₂O)_n, n = 8–20: An Ab Initio Investigation, *J. Phys. Chem. A.* 105 (2001) 10525–10537. <https://doi.org/10.1021/jp013141b>.
- [9] Y. Shi, Z. Zhou, Density functional theory study of the hydrogen bonding interaction complexes of hydrogen peroxide with glycine, *J. Mol. Struct. THEOCHEM.* 674 (2004) 113–119. <https://doi.org/10.1016/j.theochem.2003.12.018>.
- [10] L. Gonzalez, O. Miquel, M. Yáñez, High-level ab initio versus DFT calculations on (H₂O)₂ and H₂O₂-H₂O complexes as prototypes of multiple hydrogen bond systems, *J. Comput. Chem.* 18 (1997) 1124–1135. [https://doi.org/10.1002/\(SICI\)1096-987X\(19970715\)18:9<1124::AID-JCC2>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1096-987X(19970715)18:9<1124::AID-JCC2>3.0.CO;2-T).
- [11] R. Ludwig, Water: From Clusters to the Bulk, *Angew. Chemie Int. Ed.* 40 (2001) 1808–1827. [https://doi.org/10.1002/1521-3773\(20010518\)40:10<1808::AID-ANIE1808>3.0.CO;2-1](https://doi.org/10.1002/1521-3773(20010518)40:10<1808::AID-ANIE1808>3.0.CO;2-1).
- [12] S.K. Reddy, S.C. Straight, P. Bajaj, C. Huy Pham, M. Riera, D.R. Moberg, M.A. Morales, C. Knight, A.W. Götz, F. Paesani, On the accuracy of the MB-pol many-body potential for water: Interaction energies, vibrational frequencies, and classical thermodynamic and dynamical properties from clusters to liquid water and ice, *J. Chem. Phys.* 145 (2016) 194504. <https://doi.org/10.1063/1.4967719>.
- [13] U. Göra, W. Cencek, R. Podszwa, A. van der Avoird, K. Szalewicz, Predictions for water clusters from a first-principles two- and three-body force field, *J. Chem. Phys.* 140 (2014) 194101. <https://doi.org/10.1063/1.4875097>.
- [14] P. Jankowski, G. Murdachaew, R. Bukowski, O. Akin-Ojo, C. Leforestier, K.

- Szalewicz, Ab Initio Water Pair Potential with Flexible Monomers, *J. Phys. Chem. A*. 119 (2015) 2940–2964. <https://doi.org/10.1021/jp512847z>.
- [15] S.Y. Willow, X.C. Zeng, S.S. Xantheas, K.S. Kim, S. Hirata, Why Is MP2-Water “Cooler” and “Denser” than DFT-Water?, *J. Phys. Chem. Lett.* 7 (2016) 680–684. <https://doi.org/10.1021/acs.jpcclett.5b02430>.
- [16] R.M. Shields, B. Temelso, K.A. Archer, T.E. Morrell, G.C. Shields, Accurate Predictions of Water Cluster Formation, $(\text{H}_2\text{O})_n$, $n=2-10$, *J. Phys. Chem. A*. 114 (2010) 11725–11737. <https://doi.org/10.1021/jp104865w>.
- [17] B. Temelso, C.R. Renner, G.C. Shields, Importance and Reliability of Small Basis Set CCSD(T) Corrections to MP2 Binding and Relative Energies of Water Clusters, *J. Chem. Theory Comput.* 11 (2015) 1439–1448. <https://doi.org/10.1021/ct500944v>.
- [18] E. Miliordos, S.S. Xantheas, An accurate and efficient computational protocol for obtaining the complete basis set limits of the binding energies of water clusters at the MP2 and CCSD(T) levels of theory: Application to $(\text{H}_2\text{O})_m$, $m = 2-6, 8, 11, 16$, and 17, *J. Chem. Phys.* 142 (2015) 234303. <https://doi.org/10.1063/1.4922262>.
- [19] J.C. Howard, G.S. Tschumper, Wavefunction methods for the accurate characterization of water clusters, *Wiley Interdiscip. Rev. Comput. Mol. Sci.* 4 (2014) 199–224. <https://doi.org/10.1002/wcms.1168>.
- [20] N. Sahu, G. Singh, A. Nandi, S.R. Gadre, Toward an Accurate and Inexpensive Estimation of CCSD(T)/CBS Binding Energies of Large Water Clusters, *J. Phys. Chem. A*. 120 (2016) 5706–5714. <https://doi.org/10.1021/acs.jpca.6b04519>.
- [21] G.A. Cisneros, K.T. Wikfeldt, L. Ojamäe, J. Lu, Y. Xu, H. Torabifard, A.P. Bartók, G. Csányi, V. Molinero, F. Paesani, Modeling Molecular Interactions in Water: From Pairwise to Many-Body Potential Energy Functions, *Chem. Rev.* 116 (2016) 7501–7528. <https://doi.org/10.1021/acs.chemrev.5b00644>.
- [22] F.N. Keutsch, R.J. Saykally, Water clusters: Untangling the mysteries of the liquid, one molecule at a time, *Proc. Natl. Acad. Sci.* 98 (2001) 10533–10540. <https://doi.org/10.1073/pnas.191266498>.
- [23] S.Y. Lo, W.C. Li, S.H. Huang, Water clusters in life, *Med. Hypotheses*. 54 (2000) 948–953. <https://doi.org/10.1054/mehy.1999.0991>.
- [24] D. Manna, M.K. Kesharwani, N. Sylvetsky, J.M.L. Martin, Conventional and Explicitly Correlated ab Initio Benchmark Study on Water Clusters: Revision of the BEGDB and WATER27 Data Sets, *J. Chem. Theory Comput.* 13 (2017) 3136–3152. <https://doi.org/10.1021/acs.jctc.6b01046>.
- [25] J. Řezáč, P. Jurečka, K.E. Riley, J. Černý, H. Valdes, K. Pluháčková, K. Berka, T. Řezáč, M. Pitoňák, J. Vondrášek, P. Hobza, Quantum Chemical Benchmark Energy and Geometry Database for Molecular Clusters and Complex Molecular Systems (www.begdb.com): A Users Manual and Examples, *Collect. Czechoslov. Chem. Commun.* 73 (2008) 1261–1270. <https://doi.org/10.1135/cccc20081261>.
- [26] V.S. Bryantsev, M.S. Diallo, A.C.T. van Duin, W.A. Goddard, Evaluation of B3LYP, X3LYP, and M06-Class Density Functionals for Predicting the Binding Energies of Neutral, Protonated, and Deprotonated Water Clusters, *J. Chem. Theory Comput.* 5 (2009) 1016–1026. <https://doi.org/10.1021/ct800549f>.
- [27] S. Ten-no, Explicitly correlated wave functions: summary and perspective, *Theor. Chem. Acc.* 131 (2012) 1070. <https://doi.org/10.1007/s00214-011-1070-1>.
- [28] S. Ten-no, J. Noga, Explicitly correlated electronic structure theory from R12/F12 ansätze, *Wiley Interdiscip. Rev. Comput. Mol. Sci.* 2 (2012) 114–125. <https://doi.org/10.1002/wcms.68>.
- [29] L. Kong, F.A. Bischoff, E.F. Valeev, Explicitly correlated R12/F12 methods for electronic structure., *Chem. Rev.* 112 (2012) 75–107.

- <https://doi.org/10.1021/cr200204r>.
- [30] C. Hättig, W. Klopper, A. Köhn, D.P. Tew, Explicitly correlated electrons in molecules., *Chem. Rev.* 112 (2012) 4–74. <https://doi.org/10.1021/cr200168z>.
- [31] G.S. Fanourgakis, E. Aprà, S.S. Xantheas, High-level ab initio calculations for the four low-lying families of minima of (H₂O)₂₀. I. Estimates of MP2/CBS binding energies and comparison with empirical potentials, *J. Chem. Phys.* 121 (2004) 2655–2663. <https://doi.org/10.1063/1.1767519>.
- [32] G.S. Fanourgakis, E. Aprà, W.A. de Jong, S.S. Xantheas, High-level ab initio calculations for the four low-lying families of minima of (H₂O)₂₀. II. Spectroscopic signatures of the dodecahedron, fused cubes, face-sharing pentagonal prisms, and edge-sharing pentagonal prisms hydrogen bonding networks, *J. Chem. Phys.* 122 (2005) 134304. <https://doi.org/10.1063/1.1864892>.
- [33] D.J. Wales, M.P. Hodges, Global minima of water clusters (H₂O)_n, n ≤ 21, described by an empirical potential, *Chem. Phys. Lett.* 286 (1998) 65–72. [https://doi.org/10.1016/S0009-2614\(98\)00065-7](https://doi.org/10.1016/S0009-2614(98)00065-7).
- [34] F. Neese, F. Wennmohs, A. Hansen, Efficient and accurate local approximations to coupled-electron pair approaches: An attempt to revive the pair natural orbital method, *J. Chem. Phys.* 130 (2009) 114108. [citeulike-article-id:4208818](https://doi.org/10.1063/1.3173827).
- [35] C. Riplinger, F. Neese, An efficient and near linear scaling pair natural orbital based local coupled cluster method, *J. Chem. Phys.* 138 (2013) 034106. <https://doi.org/10.1063/1.4773581>.
- [36] F. Neese, A. Hansen, D.G. Liakos, Efficient and accurate approximations to the local coupled cluster singles doubles method using a truncated pair natural orbital basis., *J. Chem. Phys.* 131 (2009) 064103. <https://doi.org/10.1063/1.3173827>.
- [37] Q. Ma, H. Werner, Explicitly correlated local coupled-cluster methods using pair natural orbitals, *Wiley Interdiscip. Rev. Comput. Mol. Sci.* 8 (2018) e1371. <https://doi.org/10.1002/wcms.1371>.
- [38] F. Neese, Software update: the ORCA program system, version 4.0, *WIREs Comput. Mol. Sci.* 8 (2018). <https://doi.org/10.1002/wcms.1327>.
- [39] D.G. Liakos, M. Sparta, M.K. Kesharwani, J.M.L. Martin, F. Neese, Exploring the Accuracy Limits of Local Pair Natural Orbital Coupled-Cluster Theory, *J. Chem. Theory Comput.* 11 (2015) 1525–1539. <https://doi.org/10.1021/ct501129s>.
- [40] Q. Ma, H.-J. Werner, Scalable Electron Correlation Methods. 5. Parallel Perturbative Triples Correction for Explicitly Correlated Local Coupled Cluster with Pair Natural Orbitals, *J. Chem. Theory Comput.* 14 (2018) 198–215. <https://doi.org/10.1021/acs.jctc.7b01141>.
- [41] H.-J. Werner, P.J. Knowles, G. Knizia, F.R. Manby, M. Schütz, P. Celani, W. Györffy, D. Kats, T. Korona, R. Lindh, A. Mitrushenkov, G. Rauhut, K.R. Shamasundar, T.B. Adler, R.D. Amos, S.J. Bennie, A. Bernhardsson, A. Berning, D.L. Cooper, M.J.O. Deegan, A.J. Dobbyn, F. Eckert, E. Goll, C. Hampel, A. Hesselmann, G. Hetzer, T. Hrenar, G. Jansen, C. Köppl, S.J.R. Lee, Y. Liu, A.W. Lloyd, Q. Ma, R.A. Mata, A.J. May, S.J. McNicholas, W. Meyer, T.F. Miller III, M.E. Mura, A. Nicklaß, D.P. O’Neill, P. Palmieri, D. Peng, K. Pflüger, R. Pitzer, M. Reiher, T. Shiozaki, H. Stoll, A.J. Stone, R. Tarroni, T. Thorsteinsson, M. Wang, M. Welborn, MOLPRO, A Package of Ab Initio Programs, version 2019.2, Institute for Theoretical Chemistry, University of Stuttgart, 2019.
- [42] T.B. Adler, G. Knizia, H.-J. Werner, A simple and efficient CCSD(T)-F12 approximation., *J. Chem. Phys.* 127 (2007) 221106. <https://doi.org/10.1063/1.2817618>.
- [43] G. Knizia, T.B. Adler, H.-J. Werner, Simplified CCSD(T)-F12 methods: theory and benchmarks., *J. Chem. Phys.* 130 (2009) 054104. <https://doi.org/10.1063/1.3054300>.

- [44] H.-J. Werner, T.B. Adler, F.R. Manby, General orbital invariant MP2-F12 theory., *J. Chem. Phys.* 126 (2007) 164102. <https://doi.org/10.1063/1.2712434>.
- [45] K.A. Peterson, T.B. Adler, H.-J. Werner, Systematically convergent basis sets for explicitly correlated wavefunctions: the atoms H, He, B-Ne, and Al-Ar., *J. Chem. Phys.* 128 (2008) 084102. <https://doi.org/10.1063/1.2831537>.
- [46] F. Weigend, A fully direct RI-HF algorithm: Implementation, optimised auxiliary basis sets, demonstration of accuracy and efficiency, *Phys. Chem. Chem. Phys.* 4 (2002) 4285–4291. <https://doi.org/10.1039/b204199p>.
- [47] F. Weigend, A. Köhn, C. Hättig, Efficient use of the correlation consistent basis sets in resolution of the identity MP2 calculations, *J. Chem. Phys.* 116 (2002) 3175–3183. <https://doi.org/10.1063/1.1445115>.
- [48] C. Hättig, Optimization of auxiliary basis sets for RI-MP2 and RI-CC2 calculations: Core–valence and quintuple- ζ basis sets for H to Ar and QZVPP basis sets for Li to Kr, *Phys. Chem. Chem. Phys.* 7 (2005) 59–66. <https://doi.org/10.1039/b415208e>.
- [49] K.E. Yousaf, K.A. Peterson, Optimized auxiliary basis sets for explicitly correlated methods., *J. Chem. Phys.* 129 (2008) 184108. <https://doi.org/10.1063/1.3009271>.
- [50] K.E. Yousaf, K.A. Peterson, Optimized complementary auxiliary basis sets for explicitly correlated methods: aug-cc-pVnZ orbital basis sets, *Chem. Phys. Lett.* 476 (2009) 303–307. <https://doi.org/10.1016/j.cplett.2009.06.003>.
- [51] N. Sylvetsky, M.K. Kesharwani, J.M.L. Martin, The aug-cc-pVnZ-F12 basis set family: Correlation consistent basis sets for explicitly correlated benchmark calculations on anions and noncovalent complexes, *J. Chem. Phys.* 147 (2017) 134106. <https://doi.org/10.1063/1.4998332>.
- [52] N. Sylvetsky, M.K. Kesharwani, J.M.L. Martin, MP2-F12 basis set convergence for the S66 noncovalent interactions benchmark: Transferability of the complementary auxiliary basis set (CABS), in: *Proc. ICCMSE-2017, Thessaloniki, Greece, April 21-26, 2017*, 2017: p. 030006. <https://doi.org/10.1063/1.5012285>.
- [53] J.G. Hill, K.A. Peterson, G. Knizia, H.-J. Werner, Extrapolating MP2 and CCSD explicitly correlated correlation energies to the complete basis set limit with first and second row correlation consistent basis sets., *J. Chem. Phys.* 131 (2009) 194105. <https://doi.org/10.1063/1.3265857>.
- [54] K. Raghavachari, G.W. Trucks, J.A. Pople, M. Head-Gordon, A fifth-order perturbation comparison of electron correlation theories, *Chem. Phys. Lett.* 157 (1989) 479–483. [https://doi.org/10.1016/S0009-2614\(89\)87395-6](https://doi.org/10.1016/S0009-2614(89)87395-6).
- [55] J.D. Watts, J. Gauss, R.J. Bartlett, Coupled-cluster methods with noniterative triple excitations for restricted open-shell Hartree–Fock and other general single determinant reference functions. Energies and analytical gradients, *J. Chem. Phys.* 98 (1993) 8718–8733. <https://doi.org/10.1063/1.464480>.
- [56] K.A. Peterson, Chapter 11 Gaussian Basis Sets Exhibiting Systematic Convergence to the Complete Basis Set Limit, *Annu. Rep. Comput. Chem.* 3 (2007) 195–206. [https://doi.org/10.1016/S1574-1400\(07\)03011-3](https://doi.org/10.1016/S1574-1400(07)03011-3).
- [57] B. Brauer, M.K. Kesharwani, S. Kozuch, J.M.L. Martin, The S66x8 benchmark for noncovalent interactions revisited: explicitly correlated ab initio methods and density functional theory, *Phys. Chem. Chem. Phys.* 18 (2016) 20905–20925. <https://doi.org/10.1039/C6CP00688D>.
- [58] S.F. Boys, F. Bernardi, The calculation of small molecular interactions by the differences of separate total energies. Some procedures with reduced errors, *Mol. Phys.* 19 (1970) 553–566. <https://doi.org/10.1080/00268977000101561>.
- [59] L.A. Burns, M.S. Marshall, C.D. Sherrill, Comparing Counterpoise-Corrected, Uncorrected, and Averaged Binding Energies for Benchmarking Noncovalent

- Interactions, *J. Chem. Theory Comput.* 10 (2014) 49–57.
<https://doi.org/10.1021/ct400149j>.
- [60] B. Brauer, M.K. Kesharwani, J.M.L. Martin, Some observations on counterpoise corrections for explicitly correlated calculations on noncovalent interactions, *J. Chem. Theory Comput.* 10 (2014) 3791–3799. <https://doi.org/10.1021/ct500513b>.
- [61] S.S. Xantheas, Ab initio studies of cyclic water clusters (H₂O)_n, n=1–6. II. Analysis of many-body interactions, *J. Chem. Phys.* 100 (1994) 7523.
<https://doi.org/10.1063/1.466846>.
- [62] J.C. Howard, G.S. Tschumper, N -body:Many-body QM:QM vibrational frequencies: Application to small hydrogen-bonded clusters, *J. Chem. Phys.* 139 (2013) 184113.
<https://doi.org/10.1063/1.4829463>.
- [63] D.M. Bates, J.R. Smith, T. Janowski, G.S. Tschumper, Development of a 3-body:many-body integrated fragmentation method for weakly bound clusters and application to water clusters (H₂O)_n = 3 – 10, 16, 17, *J. Chem. Phys.* 135 (2011) 044123. <https://doi.org/10.1063/1.3609922>.
- [64] U. Góra, R. Podeszwa, W. Cencek, K. Szalewicz, Interaction energies of large clusters from many-body expansion, *J. Chem. Phys.* 135 (2011) 224102.
<https://doi.org/10.1063/1.3664730>.
- [65] J.M.L. Martin, A simple ‘range extender’ for basis set extrapolation methods for MP2 and coupled cluster correlation energies, *AIP Conf. Proc.* 2040 (2018) 020008.
<https://doi.org/10.1063/1.5079050>.
- [66] M.K. Kesharwani, N. Sylvetsky, A. Köhn, D.P. Tew, J.M.L. Martin, Do CCSD and approximate CCSD-F12 variants converge to the same basis set limits? The case of atomization energies, *J. Chem. Phys.* 149 (2018) 154109.
<https://doi.org/10.1063/1.5048665>.
- [67] K.A. Peterson, M.K. Kesharwani, J.M.L. Martin, The cc-pV5Z-F12 basis set: reaching the basis set limit in explicitly correlated calculations, *Mol. Phys.* 113 (2015) 1551–1558. <https://doi.org/10.1080/00268976.2014.985755>.
- [68] O. Marchetti, H.-J. Werner, Accurate calculations of intermolecular interaction energies using explicitly correlated wave functions., *Phys. Chem. Chem. Phys.* 10 (2008) 3400–9. <https://doi.org/10.1039/b804334e>.
- [69] C. Riplinger, B. Sandhoefer, A. Hansen, F. Neese, Natural triple excitations in local coupled cluster calculations with pair natural orbitals, *J. Chem. Phys.* 139 (2013) 134101. <https://doi.org/10.1063/1.4821834>.
- [70] Q. Ma, H.-J. Werner, Accurate Intermolecular Interaction Energies Using Explicitly Correlated Local Coupled Cluster Methods [PNO-LCCSD(T)-F12], *J. Chem. Theory Comput.* 15 (2019) 1044–1052. <https://doi.org/10.1021/acs.jctc.8b01098>.
- [71] S. Moa, F. Himo, Quantum chemical study of mechanism and stereoselectivity of secondary alcohol dehydrogenase, *J. Inorg. Biochem.* 175 (2017) 259–266.
<https://doi.org/10.1016/j.jinorgbio.2017.07.022>.
- [72] A. Altun, F. Neese, G. Bistoni, Effect of Electron Correlation on Intermolecular Interactions: A Pair Natural Orbitals Coupled Cluster Based Local Energy Decomposition Study, *J. Chem. Theory Comput.* 15 (2019) 215–228.
<https://doi.org/10.1021/acs.jctc.8b00915>.