

Dynamical Algebras in the 1+1 Dirac Oscillator and the Jaynes–Cummings Model*

Wen-Ya Song and Fu-Lin Zhang[†]

Department of Physics, School of Science, Tianjin University, Tianjin 300072, China

(Dated: January 17, 2022)

We study the algebraic structure of the one-dimensional Dirac oscillator by extending the concept of spin symmetry to a noncommutative case. An $SO(4)$ algebra is found connecting the eigenstates of the Dirac oscillator, in which the two elements of Cartan subalgebra are conserved quantities. Similar results are obtained in the Jaynes–Cummings model.

The Dirac equation is a cornerstone of modern physics [1]. It is consistent with both the principles of quantum mechanics and the theory of special relativity, and it plays an essential role in the field of high energy physics [2]. The Dirac oscillator (DO), which is obtained by the substitution $\mathbf{p} \rightarrow \mathbf{p} - i\beta m\omega \mathbf{x}$ into the free Dirac equation [3, 4], has become the paradigm for the construction of covariant quantum models with some well determined nonrelativistic limits [5]. It is used in various branches of physics, such as nuclear physics [6] and subnuclear physics [7], and it has recently been simulated in quantum optics [8–11] and classical microwave setups [5, 12]. What is particularly noteworthy is that the simulation in quantum optics is based on the equivalence between the DO and the Jaynes–Cummings (JC) model, which is the simplest soluble model in light-matter interaction within the frame of completely quantized theory [13].

The concepts of dynamical symmetry and algebraic structure are essential and prevalent in both classical and quantum mechanics [14, 15]. In two simple examples, the Hydrogen atom and harmonic oscillator in nonrelativistic quantum mechanics [14, 15], the energy levels of a quantum system may be derived based on its algebraic structure. Algebraic properties of the DO has been studied from different perspectives [16, 17] shortly after the original work of Moshinsky [3]. Recently, Zhou *et al.* [18] constructed shift operators of the JC model¹ using the matrix-diagonalizing technique [19].

In this work, we re-examine the algebraic structures of the one-dimensional (1D) DO and the JC model. The motivation for this research is that the eigenstates of the JC model (and similarly, the DO) are determined by two good quantum numbers, but only one pair of raising and lowering operators were obtained in Ref. [18]. Therefore, our aim is to derive the two conserved quantities accompanied by their shift operators, which describe the full algebraic structure of the two models.

Our work begins with the 1D DO, which can be obtained by linearizing the quadratic form $E^2 = m^2 + p^2 +$

$m^2\omega^2x^2 - \beta m\omega$ because original approach to derive the Dirac Hamiltonian [1], where ω and β are the frequency and one of the Dirac matrices. We choose $\hbar = c = 1$ in this study. This implies that the 1D DO and the two-dimensional (2D) free Dirac system may share similar features, whereas difference comes from the commutator $[x, p] = i$.

The Dirac Hamiltonian, with scalar and vector potentials of equal magnitude, certainly including the free Dirac equation, has the spin or pseudospin symmetry corresponding to the same or opposite sign [20]. These systems are shown to have the same dynamical symmetries with their nonrelativistic counterparts [21–25]. The key step to reveal the symmetries is to derive the conserved quantities with the aid of a unitary operator [21–23]. Zhang *et al.* proposed a 2D version of the unitary operator [24], and showed that it can be used to deal with noncentral systems [25].

In this work, we extend the unitary for the 2D Dirac equation to a noncommutative case. Namely, one of the momentums in the unitary is replaced by a coordinate operator. Using this unitary, one can derive a conserved number operator and a spin of the 1D DO, i.e., the shift operators which can be constructed simultaneously. With these conserved quantities and shift operators, an $SO(4)$ algebraic structure of the system is revealed, which is said to be a dynamical symmetry in the broad sense [26]. Similar results are obtained in the JC model by a similar procedure.

Let us begin with the conserved angular momentum of the 2D free Dirac equation. The Hamiltonian reads

$$H = \boldsymbol{\alpha} \cdot \mathbf{p} + \beta m, \quad (1)$$

where the Dirac matrices are conveniently defined in terms of the Pauli matrices $\boldsymbol{\alpha} = (\sigma_1, \sigma_2)$ and $\beta = \sigma_3$. In the matrix form, it reads

$$H = \begin{pmatrix} m & B^\dagger \\ B & -m \end{pmatrix}, \quad (2)$$

with $B = p_1 - ip_2$ and $B^\dagger = p_1 + ip_2$. The spin-orbit coupling leads to the fact that the usual orbital angular momentum $l = x_1p_2 - x_2p_1$ does not commute with the Hamiltonian (1). A deformed orbit momentum defined

*Chin. Phys. Lett. 37(5) 050301 (2020).

[†]Corresponding author: flzhang@tju.edu.cn

¹ The words 'of the JC model' is missed out in the published version.

in Ref. [24]

$$L = U^\dagger l U \quad (3)$$

can be easily proved to be conserved, when the unitary

$$U = \begin{pmatrix} 1 & 0 \\ 0 & \frac{B}{p} \end{pmatrix} \quad (4)$$

with $p = \sqrt{p_1^2 + p_2^2} = \sqrt{B^\dagger B} = \sqrt{B B^\dagger}$. It equals

$$L = U l U^\dagger = \begin{pmatrix} l & 0 \\ 0 & l - 1 \end{pmatrix}. \quad (5)$$

and commutes with the 2D Dirac Hamiltonian with equal scalar and vector radial potentials, which is the key reason why the dynamical symmetries in these systems studied in Ref.[24] are not broken by the spin-orbit coupling.

Now, we turn to the conserved number operator of the 1D DO. The Hamiltonian of the 1D version of the DO is given by

$$\mathcal{H} = \alpha(p - i\beta m\omega x) + \beta m, \quad (6)$$

where the Dirac matrices $\alpha = \sigma_2$ and $\beta = \sigma_3$. In the matrix form, it reads

$$\mathcal{H} = \begin{pmatrix} m & \sqrt{2m\omega} a^\dagger \\ \sqrt{2m\omega} a & -m \end{pmatrix}, \quad (7)$$

where $a^\dagger = \sqrt{\frac{m\omega}{2}}(x - \frac{i}{m\omega}p)$ and $a = \sqrt{\frac{m\omega}{2}}(x + \frac{i}{m\omega}p)$ are the raising and lowering operators of a nonrelativistic harmonic oscillator. The 1D DO can be regarded as a coupled quantum system composed by the harmonic oscillator and a spin, and the coupling vanishes in the nonrelativistic limit [27]. This interaction leads to the case that the number operator $N = a^\dagger a$, which is a good quantum number of the harmonic oscillator, is no longer conserved. It is noteworthy that the matrix (7) can be obtained by replacing the first momentum with the coordinate corresponding to the second momentum. This similarity between the Hamiltonians enlightens us to construct a deformed number operator which commutes with the DO Hamiltonian. Our solution is to extend the unitary (4) to the case of the DO. An intuitive approach is to replace B and B^\dagger in (4) with the shift operators a and a^\dagger . However, there are several possible alternatives because of the uncommutation between a and a^\dagger , or equivalently x and p . Among them, one can directly find the unitary

$$\mathcal{U} = \begin{pmatrix} 1 & 0 \\ 0 & a \frac{1}{\sqrt{N}} \end{pmatrix} \quad (8)$$

transforming the number operator into a conserved one

as

$$\mathcal{N} = \mathcal{U} N \mathcal{U}^\dagger = \begin{pmatrix} N & 0 \\ 0 & N + 1 \end{pmatrix}. \quad (9)$$

It commutes with the Hamiltonian \mathcal{H} . In addition, the spin operator is invariant under the unitary; that is,

$$s = \mathcal{U} s \mathcal{U}^\dagger, \quad (10)$$

where $s = \frac{\sigma_3}{2}$.

The spectrum of the 1D DO can be derived in the following two steps. Under the inverse transformation of the unitary (8), the Hamiltonian becomes

$$\mathcal{U}^\dagger \mathcal{H} \mathcal{U} = \begin{pmatrix} m & \sqrt{2m\omega N} \\ \sqrt{2m\omega N} & -m \end{pmatrix}. \quad (11)$$

It can be diagonalized in the two-dimensional subspace with the same quantum number N ; that is,

$$e^{i\frac{\sigma_2}{2}\theta_N} \mathcal{U}^\dagger \mathcal{H} \mathcal{U} e^{-i\frac{\sigma_2}{2}\theta_N} = \sqrt{2m\omega N + m^2} \sigma_3, \quad (12)$$

where θ_N is an operator function defined by $\tan \theta_N = \sqrt{2\omega N}/m$. Here, we remark that the above form in (12) is different from the Foldy–Wouthuysen transformation [28].

The eigenstates of the DO can be obtained directly as

$$|\varphi_n^\pm\rangle = \mathcal{U} e^{-i\frac{\sigma_2}{2}\theta_N} |\pm\rangle \otimes |n\rangle, \quad (13)$$

where $|\pm\rangle$ and $|n\rangle$ are eigenstates of the spin and the harmonic oscillator, respectively, satisfying $\sigma_3|\pm\rangle = \pm|\pm\rangle$ and $N|n\rangle = n|n\rangle$. Here, the quantum number $n = 0, 1, 2, \dots$ for the plus sign (+), but $n = 1, 2, \dots$ for the minus sign (-). They satisfy

$$\mathcal{H}|\varphi_n^\pm\rangle = \mathcal{E}_n^\pm |\varphi_n^\pm\rangle, \quad (14)$$

and the corresponding eigenenergies are

$$\mathcal{E}_n^\pm = \pm \sqrt{2m\omega n + m^2}. \quad (15)$$

The eigenstates in the matrix form are given by

$$|\varphi_n^+\rangle = \begin{pmatrix} \cos \frac{\theta_n}{2} |n\rangle \\ \sin \frac{\theta_n}{2} |n-1\rangle \end{pmatrix}, \quad |\varphi_n^-\rangle = \begin{pmatrix} -\sin \frac{\theta_n}{2} |n\rangle \\ \cos \frac{\theta_n}{2} |n-1\rangle \end{pmatrix}, \quad (16)$$

where θ_n is the eigenvalue of θ_N satisfying $\tan \theta_n = \sqrt{2\omega n}/m$.

Now we investigate the full algebraic structure of the 1D DO. The diagonalized form of the Hamiltonian (12) indicates that there is a conserved quantity in addition to the deformed number operator, which is given by

$$\Sigma_3 = \mathcal{U} e^{-i\frac{\sigma_2}{2}\theta_N} \sigma_3 e^{i\frac{\sigma_2}{2}\theta_N} \mathcal{U}^\dagger. \quad (17)$$

The deformed number operator (9) equals

$$\mathcal{N} = \mathcal{U}e^{-i\frac{\sigma_2}{2}\theta_N} N e^{i\frac{\sigma_2}{2}\theta_N} \mathcal{U}^\dagger, \quad (18)$$

as $[N, \theta_N] = 0$. The shift operators of the two conserved quantities can be obtained as

$$b = \mathcal{U}e^{-i\frac{\sigma_2}{2}\theta_N} a e^{i\frac{\sigma_2}{2}\theta_N} \mathcal{U}^\dagger, \quad b^\dagger = \mathcal{U}e^{-i\frac{\sigma_2}{2}\theta_N} a^\dagger e^{i\frac{\sigma_2}{2}\theta_N} \mathcal{U}^\dagger, \quad (19)$$

$$\Sigma_\pm = \mathcal{U}e^{-i\frac{\sigma_2}{2}\theta_N} \sigma_\pm e^{i\frac{\sigma_2}{2}\theta_N} \mathcal{U}^\dagger, \quad (20)$$

where $\sigma_\pm = (\sigma_1 \pm i\sigma_2)/2$. They satisfy the commutation relations of the harmonic oscillator and the Pauli matrices as

$$[\mathcal{N}, b] = -b, \quad [\mathcal{N}, b^\dagger] = b^\dagger, \quad [b, b^\dagger] = 1, \quad (21)$$

$$[\Sigma_3, \Sigma_\pm] = \pm 2\Sigma_\pm, \quad [\Sigma_+, \Sigma_-] = \Sigma_3. \quad (22)$$

The two conserved quantities and their shift operators constitute two independent Lie algebras. The harmonic oscillator operators construct an $SU(1, 1)$ algebra [18, 19] as

$$K_3 = \mathcal{N} + \frac{1}{2}, \quad K_+ = b^\dagger \xi(K_3), \quad K_- = \xi(K_3) b, \quad (23)$$

with $\xi(K_3) = \sqrt{\mathcal{N} + 1}$, and they satisfy the commutation relations

$$[K_3, K_\pm] = \pm K_\pm, \quad [K_+, K_-] = -2K_3. \quad (24)$$

The deformed Pauli matrices construct an $SU(2)$ algebra naturally as

$$S_3 = \frac{\Sigma_3}{2}, \quad S_\pm = \Sigma_\pm, \quad (25)$$

satisfying

$$[S_3, S_\pm] = \pm S_\pm, \quad [S_+, S_-] = 2S_3. \quad (26)$$

The non-hermitian generators in the two Lie algebras can be rewritten as the Hermitian ones as follows

$$K_1 = \frac{1}{2}(K_+ + K_-), \quad K_2 = \frac{1}{2i}(K_+ - K_-), \quad (27)$$

$$S_1 = \frac{1}{2}(S_+ + S_-), \quad S_2 = \frac{1}{2i}(S_+ - S_-). \quad (28)$$

The two algebras are decoupled, as

$$[S_i, K_j] = 0, \quad (29)$$

with $i, j = 1, 2, 3$.

These results show that the 1D DO has an $SO(4)$ algebraic structure, and its six generators are defined as

$$I_i = \tilde{K}_i + S_i, \quad R_i = \tilde{K}_i - S_i, \quad \text{with } i = 1, 2, 3, \quad (30)$$

where $\tilde{K}_1 = iK_1$, $\tilde{K}_2 = iK_2$, $\tilde{K}_3 = K_3$. The commuta-

tion relations of the $SO(4)$ algebra are

$$\begin{aligned} [I_i, I_j] &= i\epsilon_{ijk} I_k, \\ [I_i, R_j] &= i\epsilon_{ijk} R_k, \\ [R_i, R_j] &= i\epsilon_{ijk} I_k, \end{aligned} \quad (31)$$

with $i, j, k = 1, 2, 3$. The two conserved Hermitian operators I_3 and R_3 , satisfying $[I_3, R_3] = 0$, form the Catalan subalgebra. The Hilbert space of the 1D DO provides a representation of the $SO(4)$ Lie algebra, in which the I_3 and R_3 are diagonal. One can derive their matrix elements directly as

$$\begin{aligned} A_1|\varphi_n^\pm\rangle &= \frac{i}{2}[(n+1)|\varphi_{n+1}^\pm\rangle + n|\varphi_{n-1}^\pm\rangle] + (-1)^\alpha \frac{1}{2}|\varphi_n^\mp\rangle, \\ A_2|\varphi_n^\pm\rangle &= \frac{1}{2}[(n+1)|\varphi_{n+1}^\pm\rangle - n|\varphi_{n-1}^\pm\rangle] \pm (-1)^\alpha \frac{i}{2}|\varphi_n^\mp\rangle, \\ A_3|\varphi_n^\pm\rangle &= [n+1 \pm (-1)^\alpha \frac{1}{2}]|\varphi_n^\pm\rangle, \end{aligned} \quad (32)$$

where $A = I$ with $\alpha = 0$, and $A = R$ with $\alpha = 1$.

We now turn to the JC model, the simplest completely quantized model in light-matter interaction, in which a two-level atom (matter) is coupled with a quantized mode (light) of an optical cavity. This has many applications, not only in the field of quantum optics [29] but also in the field of solidstate quantum information circuits [30], both experimentally and theoretically. The Hamiltonian reads

$$\mathcal{H}_{JC} = \omega a^\dagger a + \frac{\Omega}{2} \sigma_z + J(a^\dagger \sigma^- + a \sigma^+), \quad (33)$$

where Ω is the level splitting of the two-level system, a (a^\dagger) is the destruction (creation) operator of a single bosonic mode with frequency ω , J is the coupled coefficient. In the matrix form it is given by

$$\mathcal{H}_{JC} = \begin{pmatrix} \omega a^\dagger a + \frac{\Omega}{2} & Ja \\ Ja^\dagger & \omega a^\dagger a - \frac{\Omega}{2} \end{pmatrix}. \quad (34)$$

Following a similar procedure for the DO, one can define the conserved number operator

$$\mathcal{N}_{JC} = \mathcal{V} N \mathcal{V}^\dagger = \begin{pmatrix} N & 0 \\ 0 & N - 1 \end{pmatrix} \quad (35)$$

by the unitary

$$\mathcal{V} = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sqrt{N}} a^\dagger \end{pmatrix}. \quad (36)$$

The Hamiltonian can be diagonalized in two steps. First, by the unitary, it is transformed into

$$\mathcal{V}^\dagger \mathcal{H}_{JC} \mathcal{V} = \begin{pmatrix} \omega N + \frac{\Omega}{2} & J\sqrt{N+1} \\ J\sqrt{N+1} & \omega(N+1) - \frac{1}{2}\Omega \end{pmatrix}. \quad (37)$$

Second, one can choose a 2×2 operator in the subspace

with the same quantum number as

$$e^{i\frac{\sigma_2}{2}\phi_N}\mathcal{V}^\dagger\mathcal{H}_{JC}\mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N} \\ = \omega(N + \frac{1}{2}) + \sqrt{J^2(N + 1) + \frac{(\Omega - \omega)^2}{4}}\sigma_3, \quad (38)$$

where ϕ_N is an operator function defined by $\tan \phi_N = 2J\sqrt{N + 1}/(\Omega - \omega)$. Hence, its eigenvalues are

$$\epsilon_n^\pm = \omega(n + \frac{1}{2}) \pm \sqrt{J^2(n + 1) + \frac{(\Omega - \omega)^2}{4}}, \quad (39)$$

corresponding to the eigenstates

$$|\psi_n^\pm\rangle = \mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N}|\pm\rangle \otimes |n\rangle, \quad (40)$$

where $n = 0, 1, 2, \dots$. Two conserved quantities and their shift operators can be constructed as

$$\begin{aligned} \mathcal{N}_{JC} &= \mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N}Ne^{i\frac{\sigma_2}{2}\phi_N}\mathcal{V}^\dagger, \\ b_{JC} &= \mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N}ae^{i\frac{\sigma_2}{2}\phi_N}\mathcal{V}^\dagger, \\ b_{JC}^\dagger &= \mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N}a^\dagger e^{i\frac{\sigma_2}{2}\phi_N}\mathcal{V}^\dagger, \\ \Sigma_{JC}^{(3)} &= \mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N}\sigma_3e^{i\frac{\sigma_2}{2}\phi_N}\mathcal{V}^\dagger, \\ \Sigma_{JC}^{(\pm)} &= \mathcal{V}e^{-i\frac{\sigma_2}{2}\phi_N}\sigma_\pm e^{i\frac{\sigma_2}{2}\phi_N}\mathcal{V}^\dagger. \end{aligned} \quad (41)$$

By following the same steps to study the DO, one can show an $SO(4)$ algebraic structure in the JC model.

In summary, by extending the approach to study the 2D Dirac system with a spin symmetry, we present a unitary which transforms the number operator to a conserved quantity of the 1D DO. Based on this result, one can diagonalize the DO Hamiltonian, and obtain the two conserved quantities together with their shift operators. These operators show an $SO(4)$ algebra connecting the eigenstates of the Dirac oscillator. By a similar procedure, we also show the $SO(4)$ algebra in the JC model.

Further research on this topic in the two following directions would be interesting. First, whether one can extend the conserved number operator to the two- or three-dimensional Dirac oscillators, and derive a deformed isotropic harmonic oscillator, is a natural question. Second, it is fascinating to consider the algebraic of the multiphoton JC model [31]. More specifically, we look forward to obtaining a polynomial generalization of the $SO(4)$ algebra [25] in it.

Acknowledgments. This work was supported by the National Natural Science Foundation of China (Grant Nos. 11675119, 11575125, and 11105097).

- [2] Zee A 2010 *Quantum Field Theory in a Nutshell* (Princeton: Princeton University Press)
- [3] Moshinsky M and Szczepaniak A 1989 J. Phys. A: Math. Gen. **98**, 817
- [4] Sadurní E 2010 AIP Conf. Proc. **1334**, 249
- [5] Franco-Villafañe J A, Sadurní E, Barkhofen S, Kuhl U, Mortessagne F and Seligman T H 2013 Phys. Rev. Lett. **111**, 170405
- [6] Grineviciute J and Halderson D 2012 Phys. Rev. C **85**, 054617
- [7] Romera E 2011 Phys. Rev. A **84**, 052102
- [8] Bermudez A, Martin-Delgado M A and Solano E 2007 Phys. Rev. A **76**, 041801
- [9] Bermudez A, Martin-Delgado M A and Solano E 2007 Phys. Rev. Lett. **99**, 123602
- [10] Lamata L, León J, Schätz T and Solano E 2007 Phys. Rev. Lett. **98**, 253005
- [11] Torres J M, Sadurn E. and Seligman T H 2010 AIP Conf. Proc. **1323**, 301
- [12] Sadurní E, Seligman T H and Mortessagne F 2010 New J. Phys. **12**, 053014
- [13] Jaynes E T and Cummings F W 1963 Proceedings of the IEEE **51**, 89
- [14] Greiner W and Müller B 1994 *Quantum Mechanics (Symmetries)* (New York: Springer)
- [15] De Lange O L and Raab R E 1991 *Operator Methods in Quantum Mechanics* (Oxford: Clarendon Press)
- [16] De Lange O L 1991 J. Phys. A: Math. Gen. **24**, 667
- [17] Benitez J 1990 Phys. Rev. Lett. **64**, 1643
- [18] Zhou J, Su H Y, Zhang F L, Zhang H B and Chen J L 2018 Chin. Phys. Lett. **35**, 010302
- [19] Ge M L, Kwek L C, Liu Y, Oh C H and Wang X B 2000 Phys. Rev. A **62**, 052110
- [20] Ginocchio J N 2005 Phys. Rep. **414**, 165
- [21] Ginocchio J N 2004 Phys. Rev. C **69**, 034318
- [22] Ginocchio J N 2005 Phys. Rev. Lett. **95**, 252501
- [23] Zhang F L, Fu B and Chen J L 2008 Phys. Rev. A **78**, 040101
- [24] Zhang F L, Song C and Chen J L 2009 Ann. Phys. **324**, 173
- [25] Zhang F L, Fu B and Chen J L 2009 Phys. Rev. A **80**, 054102
- [26] Chen J L, Liu Y and Ge M L 1998 J. Phys. A: Math. Gen. **31**, 6473
- [27] Cao B X and Zhang F L 2019 arXiv:1908.09352
- [28] Moreno M and Zentella A 1989 Journal of Physics A: Mathematical and General **22**, L821
- [29] Vedral V 2005 *Modern Foundations of Quantum Optics* (London: World Scientific Publishing Company)
- [30] Irish E 2007 Phys. Rev. Lett. **99**, 173601
- [31] Lu H X and Wang X Q 2000 Chin. Phys. **9**, 568

[1] Dirac P A M 1982 *The Principles of Quantum Mechanics* (Oxford: Oxford University Press)