$ext{DISIM}_b(\mathbf{2})$ Local Relativistic Symmetry and Finslerian Extension of the Theory of Relativity

George Yu. Bogoslovsky

Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia bogoslov@theory.sinp.msu.ru

Finslerian extension of the theory of relativity implies that space-time can be not only in an amorphous state which is described by Riemann geometry but also in ordered, i.e. crystalline states which are described by Finsler geometry. Transitions between various metric states of space-time have the meaning of phase transitions in its geometric structure. These transitions together with the evolution of each of the possible metric states make up the general picture of space-time manifold dynamics. It is shown that there are only two types of curved Finslerian spaces endowed with local relativistic symmetry. However the metric of only one of them satisfies the correspondence principle with Riemannian metric of the general theory of relativity and therefore underlies viable Finslerian extension of the GR. Since the existing purely geometric approaches to a Finslerian generalization of Einstein's equations do not allow one to obtain such generalized equations which would provide a local relativistic symmetry of their solutions, special attention is paid to the property of the specific invariance of viable Finslerian metric under local conformal transformations of those fields on which it explicitly depends. It is this property that makes it possible to use the well-known methods of conventional field theory and thereby to circumvent the above-mentioned difficulties arising within the framework of purely geometric approaches to a Finslerian generalization of Einstein's equations.

Keywords: Finsler geometry; DISIM $_b(2)$ relativistic symmetry; Finslerian extension of GR

1. Introduction

The impressive successes of the contemporary cosmology are based to a large extent on the use of the Riemannian space-time model and the Einstein equations with Λ term

$$R_{ik} - \frac{1}{2}Rg_{ik} = \Lambda g_{ik} + T_{ik} \,. \tag{1}$$

In connection with the Λ term let us direct our attention to the Weyl theory [1, 2]. Trying to construct a unified theory of gravitation and electromagnetism, Weyl proceeded from the requirement of invariance of the theory with respect to the local conformal transformations of the Riemannian metric tensor $g_{ik} \to \exp[2\sigma(x)] g_{ik}$. The appearing Abelian gauge field, denoted below by A_i , was identified by Weyl with an electromagnetic field. The basis for such identification was the gradient character of the corresponding gauge transformations $A_i \to A_i + a\sigma_{;i}$, where a is a constant with the dimension of length. It is easy, however, to show [3] that in reality in the Weyl theory the A_i is not an

electromagnetic field but acquires mass due to the violation of local conformal invariance. This result is essentially contained in the Weyl variational principle

$$\delta \left[\frac{1}{16\Lambda} \int \left(R + \frac{6}{a} A^{i}_{;i} - \frac{6}{a^{2}} A_{i} A^{i} \right)^{2} \sqrt{-g} d^{4}x - \frac{1}{4} \int F_{ik} F^{ik} \sqrt{-g} d^{4}x \right] = 0, \qquad (2)$$

where $F_{ik} = A_{k;i} - A_{i;k}$ and Λ is a constant. In fact, using the Weyl gauge condition

$$R + \frac{6}{a}A^{i}_{;i} - \frac{6}{a^{2}}A_{i}A^{i} = -4\Lambda \tag{3}$$

and his variational principle (2), we get the following system of equations

$$F^{ik}_{;k} - \frac{6}{a^2} A^i = 0, (4)$$

$$R_{ik} - \frac{1}{2}Rg_{ik} - \Lambda g_{ik} = -F_{il}F^{l}_{k} + \frac{1}{4}F_{lm}F^{lm}g_{ik} + \frac{6}{a^{2}}(A_{i}A_{k} - \frac{1}{2}A_{l}A^{l}g_{ik}).$$
 (5)

Since, by virtue of (4), the equality $A_{;i}^i = 0$ holds and, by virtue of (5), the equality $R - \frac{6}{a^2}A_iA^i = -4\Lambda$ holds, it is clear that the Weyl gauge condition (3) is fulfilled for the solutions of the system of equations (4) and (5). Thus, a system of fourth-order field equations, following from the Weyl gauge invariant variational principle (2), together with the condition (3) fixing the gauge, is equivalent to the system of equations (4) and (5). In other words, the fixing of the gauge, which implies in the Weyl theory the incorporation of length and mass standards, converts the Abelian vector field A_i into a massive vector field and reduces the equations for gravitational field to the Einstein equations with Λ term.

Let us return to the Einstein equations (1). The vacuum Lorentz-invariant energy-momentum tensor Λg_{ik} can be written in the form $\Lambda g_{ik} = diag(\rho_v, -\rho_v, -\rho_v, -\rho_v)$, where the so-called dark energy density ρ_v is responsible for the accelerated expansion of the Universe. Thus, the corresponding vacuum space turns out to be filled with specific medium (scalar condensate, or quintessence), the pressure in which is negative and determined by the equation of state $p = -\rho_v$.

Apart from scalar condensate, the dynamic rearrangement of vacuum of quantized fields under spontaneous violation of initial gauge symmetry may give rise to anisotropic condensates. The flat space-time keeps being the Minkowski space in the only case of scalar condensate. But, if an anisotropic condensate arises, the corresponding anisotropy also appears in space-time [4]. Speaking of local anisotropy of space-time we have in mind its Finslerian geometric structure. In contrast to Riemann metrics, Finsler ones do not reduce to a quadratic form in the differentials of the coordinates. Being homogeneous functions of the differentials of the coordinates of the second degree of homogeneity, such metrics serve as a natural generalization of Riemann metrics.

In order to arrive naturally at the flat relativistically invariant Finslerian space-time we first confine ourselves to a two-dimensional space and show that it is possible to generalize the Lorentz transformations

$$\begin{cases} x'_0 = x_0 \cosh \alpha - x \sinh \alpha \\ x' = -x_0 \sinh \alpha + x \cosh \alpha; \qquad \tanh \alpha = v/c \end{cases}$$
 (6)

so that the new linear transformations will also form a group with a single parameter α and will keep invariance of the wave equation. Guided by the conformal invariance of the electrodynamic equations, we insert the additional scale transformations $e^{-b\alpha}$ into the standard Lorentz ones (6). As a result we obtain the generalized Lorentz transformations in the form

$$\begin{cases} x_0' = e^{-b\alpha} \left(x_0 \cosh \alpha - x \sinh \alpha \right) \\ x' = e^{-b\alpha} \left(-x_0 \sinh \alpha + x \cosh \alpha \right), \end{cases}$$
 (7)

where b is a dimensionless parameter of the scale transformations. Since the relation of the group parameter α to the velocity v of the primed frame remains unchanged, i.e. $\tanh \alpha = v/c$, the generalized Lorentz transformations can be rewritten as follows

$$\begin{cases} x'_0 = \left(\frac{1 - v/c}{1 + v/c}\right)^{b/2} \frac{x_0 - (v/c)x}{\sqrt{1 - v^2/c^2}} \\ x' = \left(\frac{1 - v/c}{1 + v/c}\right)^{b/2} \frac{x - (v/c)x_0}{\sqrt{1 - v^2/c^2}} \end{cases}$$

Obviously, in contrast to the standard Lorentz transformations (6), the generalized ones (7) do not leave invariant the Minkowski metric $ds^2 = dx_0^2 - dx^2$ but conformally modify it. Therefore, the question arises as to what the metric of an event space invariant under such generalized Lorentz transformations is. A rigorous solution of this problem leads to the following metric

$$ds^{2} = \left[\frac{(dx_{0} - dx)^{2}}{dx_{0}^{2} - dx^{2}}\right]^{b} (dx_{0}^{2} - dx^{2}).$$
(8)

Not being a quadratic form but a homogeneous function of the coordinate differentials of degree two, this metric falls into the category of Finslerian metrics. It describes a flat but anisotropic event space.

The observation made serves as a starting point for a viable Finslerian extension of both Special and General Relativity. To this end, we should first generalize twodimensional metric (8) to the case of a four-dimensional Finslerian event space. As it turned out, there are two independent approaches to such a generalization. The first approach leads to a flat Finslerian event space with partially broken 3D isotropy. The corresponding Finslerian metric, within the framework of the correspondence principle, generalizes the pseudo-Euclidean metric of the Minkowski space and describes another possible vacuum state of a flat space-time, namely, its axially symmetric crystalline state. Accordingly, the axially symmetric event space underlies the viable Finslerian extension of Special Relativity. The second approach leads to a flat Finslerian event space with entirely broken 3D isotropy. The corresponding Finslerian metric describes one more possible vacuum state of a flat space-time, namely, its entirely anisotropic crystalline state. Such a state is considered in detail in Section 4. However, the main purpose of this article is to describe the physically meaningful Finslerian extension of General Relativity (see Section 5). At the same time, there is no need to say that the corresponding Finslerian extension of Special Relativity should be considered first. Therefore, the following Sections 2 and 3 are devoted specifically to the Finslerian Special Relativity. Despite a preliminary publication [5], these Sections are presented here in their original form.

Although this article has a review character, it contains (with few exceptions) only those results on Finslerian extensions of the relativity theory, that were obtained by the author or with his participation. As for the results obtained in the framework of the traditional, purely geometric approach to a Finslerian modification of General Relativity, they are considered in a recent review [6].

2. Metric and group of relativistic symmetry of the flat Finslerian space-time with partially broken 3D isotropy

Proceeding from the Finslerian metric of the two-dimensional space of events (8) and using the substitutions $(dx_0^2 - dx^2) \rightarrow (dx_0^2 - dx^2)$; $(dx_0 - dx) \rightarrow (dx_0 - ndx)$, we arrive at the corresponding Finslerian metric [7] of the four-dimensional space of events

$$ds^{2} = \left[\frac{(dx_{0} - \mathbf{n}d\mathbf{x})^{2}}{dx_{0}^{2} - d\mathbf{x}^{2}}\right]^{b} (dx_{0}^{2} - d\mathbf{x}^{2}).$$
(9)

Here the unit vector \boldsymbol{n} indicates a preferred direction in 3D space while the parameter b determines the magnitude of space anisotropy and characterizes the degree of deviation of the metric (9) from the Minkowski metric. Thus, the flat Finslerian space-time (9) proves to be a generalization of the isotropic Minkowski space of conventional special relativity. Instead of the 3-parameter rotation group, the flat anisotropic event space (9) admits only 1-parameter group of rotations around the unit vector \boldsymbol{n} . This is why one can speak of a partially broken 3D isotropy. As to relativistic symmetry, it is realized by means of 3-parameter group of the generalized Lorentz transformations (of the generalized Lorentz boosts) which link the physically equivalent inertial reference frames in the flat anisotropic space-time and leave its metric (9) invariant. The generalized Lorentz boosts have the form

$$x^{\prime i} = D(\boldsymbol{v}, \boldsymbol{n}) R_j^i(\boldsymbol{v}, \boldsymbol{n}) L_k^j(\boldsymbol{v}) x^k,$$
(10)

where \boldsymbol{v} denotes the velocities of moving (primed) inertial reference frames, the matrices $L_k^j(\boldsymbol{v})$ represent the ordinary Lorentz boosts, the matrices $R_j^i(\boldsymbol{v},\boldsymbol{n})$ represent additional rotations of the spatial axes of the moving frames around the vectors $[\boldsymbol{v}\boldsymbol{n}]$ through the angles

$$\varphi = \arccos \left\{ 1 - \frac{(1 - \sqrt{1 - \boldsymbol{v}^2/c^2})[\boldsymbol{v}\boldsymbol{n}]^2}{(1 - \boldsymbol{v}\boldsymbol{n}/c)\boldsymbol{v}^2} \right\}$$

of relativistic aberration of n, and the diagonal matrices

$$D(\boldsymbol{v}, \boldsymbol{n}) = \left(\frac{1 - \boldsymbol{v} \boldsymbol{n}/c}{\sqrt{1 - \boldsymbol{v}^2/c^2}}\right)^b I$$

stand for the additional dilatational transformations of the event coordinates.

It should be noted that at b = 0 the Finslerian metric (9) reduces to the Minkowski metric, but the generalized Lorentz boosts, i.e. transformations (10) do not reduce to

ordinary Lorentz boosts and take the form

$$x^{\prime i} = R_j^i(\boldsymbol{v}, \boldsymbol{n}) L_k^j(\boldsymbol{v}) x^k. \tag{11}$$

At b=0, i.e. in the case of Minkowski space where all directions in 3D space are equivalent, \boldsymbol{n} has no physical meaning. In this case, each of the transformations (11) is differed from the respective Lorentz boost $x'^i=L_k^i(\boldsymbol{v})x^k$ by the additional rotation $x'^i=R_k^i(\boldsymbol{v},\boldsymbol{n})x^k$ of the spatial axes. These additional rotations are adjusted in such a way that if a ray of light has the the direction \boldsymbol{n} in one frame, then it will have the same direction in all frames under consideration. Thus, at b=0, i.e. within the framework of conventional special relativity, the transformations (11) represent an alternative to the Lorentz boosts, however, in contrast to the boosts, they constitute a 3-parameter noncompact group which turns out to be a subgroup of the 6-parameter homogeneous Lorentz group. Physically such noncompact transformations are realized as follows. First choose as \boldsymbol{n} a direction towards a preselected star and then perform an arbitrary Lorentz boost by complementing it with such a turn of the spatial axes that in a new reference frame the direction towards the star remains unchanged. The set of the transformations described has just been given by the above-displayed relation (11).

Now let us consider an inhomogeneous group of relativistic symmetry or, in other words, inhomogeneous group of isometries of the flat Finslerian space-time (9). Since the respective homogeneous noncompact group (10) is 3-parametric, with inclusion of 1-parameter group of rotations around the preferred direction \boldsymbol{n} and 4-parameter translation group, the inhomogeneous group of relativistic symmetry of the space-time (9) appears to have eight parameters. To obtain the simplest representation for its generators, it is enough to send the third spatial axis along \boldsymbol{n} and rewrite the homogeneous transformations (10) in the infinitesimal form. As a result, we come to the following eight generators

$$\begin{split} X_1 &= -(x^1p_0 + x^0p_1) - (x^1p_3 - x^3p_1) \,, \\ X_2 &= -(x^2p_0 + x^0p_2) + (x^3p_2 - x^2p_3) \,, \\ X_3 &= -bx^ip_i - (x^3p_0 + x^0p_3) \,, \\ R_3 &= x^2p_1 - x^1p_2 \,; \\ p_i &= \partial/\partial x^i \,. \end{split}$$

These generators satisfy the commutation relations

$$\begin{split} [X_1X_2] &= 0 \,, & [R_3X_3] &= 0 \,, \\ [X_3X_1] &= X_1 \,, & [R_3X_1] &= X_2 \,, \\ [X_3X_2] &= X_2 \,, & [R_3X_2] &= -X_1 \,; \\ \\ [p_ip_j] &= 0 \,; & \\ [X_1p_0] &= p_1 \,, & [X_2p_0] &= p_2 \,, & [X_3p_0] &= bp_0 + p_3 \,, & [R_3p_0] &= 0 \,, \\ [X_1p_1] &= p_0 + p_3 \,, & [X_2p_1] &= 0 \,, & [X_3p_1] &= bp_1 \,, & [R_3p_1] &= p_2 \,, \\ [X_1p_2] &= 0 \,, & [X_2p_2] &= p_0 + p_3 \,, & [X_3p_2] &= bp_2 \,, & [R_3p_2] &= -p_1 \,, \\ [X_1p_3] &= -p_1 \,, & [X_2p_3] &= -p_2 \,, & [X_3p_3] &= bp_3 + p_0 \,, & [R_3p_3] &= 0 \,. \end{split}$$

It is clear that the homogeneous isometry group of the flat Finslerian space-time (9) contains four parameters (generators X_1 , X_2 , X_3 and R_3). This group is a subgroup of

the 11-parametric Similitude (Weyl) group [8], and it is isomorphic to the corresponding 4-parametric subgroup (with generators $X_1, X_2, X_3|_{b=0}$ and R_3) of the homogeneous Lorentz group. Since the 6-parametric homogeneous Lorentz group does not have any 5-parametric subgroup [9], while its 4-parametric subgroup is unique up to isomorphisms, the passage from the Minkowski space-time to the Finslerian space-time (9) implies a minimum possible violation of Lorentz symmetry. With this, the relativistic symmetry represented now by generalized Lorentz boosts (10) remains valid [10]. Besides, in spite of a new geometry of event space, the Lobachevsky geometry of 3-velocity space and, respectively, the relativistic law of addition of 3-velocities remain unchanged. Finally, it should be noted that the above-described 8-parameter inhomogeneous group of relativistic symmetry (the group of motions) of the anisotropic event space (9) was first scrutinized in [11]-[13]. At the present time, thanks to [14], this group is known Deformed Inhomogeneous subgroup of SIMilitude group that as $DISIM_b(2)$, i.e. includes 2-parameter Abelian homogeneous noncompact subgroup. As for the group SIM(2), its generators X_1 , X_2 , $X_3|_{b=0}$, R_3 and their Lie algebra have already been described above. The Lie algebra of the corresponding Hermitian operators is also displayed in [15], where a nontrivial physical implementation of the SIM(2) symmetry is proposed.

3. Relativistic point mechanics and the rest momentum as a new observable in the flat Finslerian space-time with partially broken 3D isotropy

A remarkable property of the partially anisotropic space of events (9) is the fact that it keeps the conformal structure (light cone) of the Minkowski space, i.e. light propagates according to the equation $dx_0^2 - dx^2 = 0$. Therefore, the speed of light is independent of the direction of its propagation and is equal to c. It thus appears that the square of the distance dl between adjacent points of 3D space, determined by means of exchange of light signals, is expressed by the formula $dl^2 = dx^2$. Thus, although in the 3D space there is a preferred direction n, its geometry remains Euclidean. But, what does the anisotropy of space physically manifest itself in? First of all, the space anisotropy affects the dependence of proper time of a moving clock on the direction of clock's velocity: according to (9), the interval $d\tau$ of proper time read by the clock moving with a velocity v is related to the time interval $d\tau$ read by a clock at rest by the relation $d\tau = (d\tau/dt) dt$, where

$$\frac{d\tau}{dt} = \left(\frac{1 - \boldsymbol{v}\boldsymbol{n}/c}{\sqrt{1 - \boldsymbol{v}^2/c^2}}\right)^b \sqrt{1 - \boldsymbol{v}^2/c^2} \ .$$

Thus, in contrast to Minkowski space (for which: b=0, $(d\tau/dt)|_{b=0} = \sqrt{1-v^2/c^2} \le 1$ and, hence, the moving clock is always slow in comparison with the clock at rest), in the anisotropic space the time dilatation factor $(d\tau/dt)|_{b>0}$ can take on values greater than unity.

In order to generalize conventional relativistic point mechanics for the case of the flat partially anisotropic space-time, we replace the Minkowski line element $ds = \sqrt{dx_0^2 - dx^2}$ with the Finslerian element in the usual action integral. As a result, the action integral

¹In connection with the $DISIM_b(2)$ relativistic symmetry, note here a recent interesting paper [23].




Figure 1: Plots illustrating the dependence of time dilatation factor $(d\tau/dt)|_{b\equiv r=0.6}$ on the magnitude of clock's velocity at angles α between \boldsymbol{v} and \boldsymbol{n} equal to 0, $\pi/4$ and $\pi/2$. The semi-circumference corresponds to the isotropic Minkowski space for which $b\equiv r=0$.

for a free relativistic particle in the flat partially anisotropic space-time takes, according to (9), the form

$$S = -mc \int_{i}^{f} ds, \qquad (12)$$

where $ds = \left[(dx_0 - ndx) / \sqrt{dx_0^2 - dx^2} \right]^b \sqrt{dx_0^2 - dx^2}$. Now we have to make sure that formula (12) correctly determines the action for a free particle. To this end, consider, along with the clock at rest, another clock which starts to move from the clock at rest and, having executed nonuniform motion along a closed path, returns to the initial point. Since at certain values of the velocity v the time dilatation factor in the anisotropic space is greater than unity (see Figure 1), the moving clock is fast (on the corresponding parts of its path) in comparison with the clock at rest. Yet, on returning to the clock at rest, the moving clock will necessarily be behind. The validity of this assertion was rigorously proved in [16]. Thus in the flat anisotropic space-time (9) the integral $\int_i^f ds$ is at a maximum while the action (12) is at a minimum on the straight world line connecting points i and f. Therefore, (12) really correctly determines the action for a free particle in the flat anisotropic space-time (9).

The action (12) leads to the following relativistic Lagrange function

$$L = -mc^{2} \left(\frac{1 - v n/c}{\sqrt{1 - v^{2}/c^{2}}} \right)^{b} \sqrt{1 - v^{2}/c^{2}}.$$
 (13)

It follows from (13) that

$$E = \frac{mc^2}{\sqrt{1 - \mathbf{v}^2/c^2}} \left(\frac{1 - \mathbf{v} \mathbf{n}/c}{\sqrt{1 - \mathbf{v}^2/c^2}} \right)^b \left[1 - b + b \frac{1 - \mathbf{v}^2/c^2}{1 - \mathbf{v} \mathbf{n}/c} \right]. \tag{14}$$

Thus, the energy E of a free particle in the anisotropic space depends both on the magnitude and on the direction of its velocity \mathbf{v} . At $\mathbf{v}=0$ the energy reaches its absolute minimum, i.e the rest energy $E|_{\mathbf{v}=0}=mc^2$. As regards the momentum

$$\boldsymbol{p} = \frac{m}{\sqrt{1 - \boldsymbol{v}^2/c^2}} \left(\frac{1 - \boldsymbol{v} \boldsymbol{n}/c}{\sqrt{1 - \boldsymbol{v}^2/c^2}} \right)^b \left[(1 - b) \boldsymbol{v} + bc \boldsymbol{n} \frac{1 - \boldsymbol{v}^2/c^2}{1 - \boldsymbol{v} \boldsymbol{n}/c} \right], \tag{15}$$

Even in the case $\mathbf{v} = 0$ the momentum of the particle does not vanish; there remains its rest momentum. Thus, in the anisotropic space (9), any massive particle besides its rest energy $E|_{\mathbf{v}=0} = mc^2$ also has the rest momentum $\mathbf{p}|_{\mathbf{v}=0} = bmc\mathbf{n}$. Massless particles have no such property; for them, as in conventional special relativity, $\mathbf{v} = c$ and $E^2/c^2 - \mathbf{p}^2 = 0$.

The energy (14) and the momentum (15) are related by the following modified dispersion relation (modified mass-shell equation)

$$\left[\frac{(p_0 - \boldsymbol{p}\boldsymbol{n})^2}{p_0^2 - \boldsymbol{p}^2}\right]^{-b} (p_0^2 - \boldsymbol{p}^2) = (mc)^2 (1 - b)^{(1-b)} (1 + b)^{(1+b)},$$
(16)

where $p_0 = p^0 = E/c$. This modified relation describes a deformed two-sheeted hyperboloid. Such a hyperboloid is relativistically invariant owing to the fact that the generalized Lorentz transformations (10) of the event coordinates x^i , i.e. the transformations $x^i = DR^i_j L^j_k x^k$ are accompanied by the corresponding transformations $p^i = D^{-1}R^i_j L^j_k p^k$ of 4-momenta $p^i = (p^0, \mathbf{p})$. It is obvious that the deformation of a two-sheeted hyperboloid depends on the magnitude b of space anisotropy. The corresponding two-sheeted hyperboloids at $b \equiv r = 0, 0.3, 0.6$ and 0.8 are shown in Figure 2.

The Lagrange function (13) takes on the following form in the nonrelativistic limit

$$L = -mc^{2} + bmc(\mathbf{v}\mathbf{n}) + (1-b)m\mathbf{v}^{2}/2 + b(1-b)m(\mathbf{v}\mathbf{n})^{2}/2.$$
 (17)

Since in (17) the expression $-mc^2 + bmc(\boldsymbol{v}\boldsymbol{n})$ is a total derivative with respect to time, it can be omitted. As a result, we see that in the case of a nonrelativistic particle in the anisotropic space its kinetic energy $T = m_{\alpha\beta} v^{\alpha} v^{\beta}/2$ and momentum $p_{\alpha} = m_{\alpha\beta} v^{\beta}$ are determined by the inertial mass tensor

$$m_{\alpha\beta} = m(1-b)(\delta_{\alpha\beta} + b \, n_{\alpha} n_{\beta}). \tag{18}$$

Let us now rewrite Finslerian metric (9) so that it is expressed through the fourdimensional entities:

$$ds = \left[\frac{(dx_0 - \mathbf{n}d\mathbf{x})^2}{dx_0^2 - d\mathbf{x}^2} \right]^{b/2} \sqrt{dx_0^2 - d\mathbf{x}^2} = \left[\frac{(n_i dx^i)^2}{\eta_{ik} dx^i dx^k} \right]^{b/2} \sqrt{\eta_{ik} dx^i dx^k}.$$
(19)




Figure 2: Parametric 3D plots of the mass shells in a space of kinematic 4-momenta. Any of the deformed two-sheeted hyperboloids remains inscribed into a light cone and like a light cone it is an invariant of the generalized Lorentz transformations.

Since $n^2 = 1$, it is clear that here we have

$$n_i = \{1, -\mathbf{n}\}, \quad \eta_{ik} = diag\{1, -1, -1, -1\}, \quad n^i = \{1, \mathbf{n}\}, \quad n_i n^i = 0.$$

In accordance with (12) and (19), a constant field b defines the specific inseparable interaction of a constant null-vector field n_i with massive particles. Effect of this interaction is that the particles acquire - according to (18), the properties of quasi-particles in an axially symmetric crystalline medium, the role of which is played by axially symmetric quintessence. The latter is, thus, the physical carrier of the anisotropy of the flat space of events (19) and it consists of Weyl fermion-antifermion condensate. In this regard, it is worth noting that, in contrast to the standard Dirac equation, the corresponding constant spinors are non-trivial solutions of the $DISIM_b(2)$ -invariant generalized massive Dirac equation [17], whose Lagrangian has the form

$$\mathcal{L} = \frac{i}{2} \left(\bar{\psi} \gamma^j \partial_j \psi - \partial_j \bar{\psi} \gamma^j \psi \right) - m \left[\left(\frac{n_j \bar{\psi} \gamma^j \psi}{\bar{\psi} \psi} \right)^2 \right]^{b/2} \bar{\psi} \psi. \tag{20}$$

4. Metric and group of relativistic symmetry of the flat Finslerian space-time with entirely broken 3D isotropy

In [18] the metric describing relativistically invariant Finslerian spaces with entirely broken 3D isotropy has been obtained in the following most general form

$$ds = (dx_0 - dx_1 - dx_2 - dx_3)^{(1+b_1+b_2+b_3)/4} (dx_0 - dx_1 + dx_2 + dx_3)^{(1+b_1-b_2-b_3)/4} \times (dx_0 + dx_1 - dx_2 + dx_3)^{(1-b_1+b_2-b_3)/4} (dx_0 + dx_1 + dx_2 - dx_3)^{(1-b_1-b_2+b_3)/4}.$$
 (21)

The parameters characterizing the anisotropy of the spaces (21) are limited by the conditions

$$1 + b_1 + b_2 + b_3 \ge 0$$
, $1 + b_1 - b_2 - b_3 \ge 0$, $1 - b_1 + b_2 - b_3 \ge 0$, $1 - b_1 - b_2 + b_3 \ge 0$

It should be noted that if $b_1 = b_2 = b_3 = 0$, then the metric (21) takes the form

$$ds_{B-M} = \left[(dx_0 - dx_1 - dx_2 - dx_3)(dx_0 - dx_1 + dx_2 + dx_3) \times (dx_0 + dx_1 - dx_2 + dx_3)(dx_0 + dx_1 + dx_2 - dx_3) \right]^{1/4}.$$

Thus, in this particular case, we obtain the well-known Berwald-Moor metric, but written in the basis, which was introduced in [18].

Now consider the group of isometries of flat Finslerian spaces (21). The homogeneous 3-parametric non-compact group of isometries, i.e. the group of the relativistic symmetry of space-time (21) appears to be Abelian, and the transformations belonging to such a group have the same meaning as the ordinary Lorentz boosts. The explicit form of these transformations is

$$x_i' = D L_{ik} x_k, (22)$$

where

$$D = e^{-(b_1 \alpha_1 + b_2 \alpha_2 + b_3 \alpha_3)}$$

 L_{ik} are the unimodular matrices that are given by the formulas

$$L_{ik} = \begin{pmatrix} \mathcal{A} & -\mathcal{B} & -\mathcal{C} & -\mathcal{D} \\ -\mathcal{B} & \mathcal{A} & \mathcal{D} & \mathcal{C} \\ -\mathcal{C} & \mathcal{D} & \mathcal{A} & \mathcal{B} \\ -\mathcal{D} & \mathcal{C} & \mathcal{B} & \mathcal{A} \end{pmatrix} , \qquad (23)$$

 $\mathcal{A} = \cosh \alpha_1 \cosh \alpha_2 \cosh \alpha_3 + \sinh \alpha_1 \sinh \alpha_2 \sinh \alpha_3,$

 $\mathcal{B} = \cosh \alpha_1 \sinh \alpha_2 \sinh \alpha_3 + \sinh \alpha_1 \cosh \alpha_2 \cosh \alpha_3$

 $\mathcal{C} = \cosh \alpha_1 \sinh \alpha_2 \cosh \alpha_3 + \sinh \alpha_1 \cosh \alpha_2 \sinh \alpha_3$

 $\mathcal{D} = \cosh \alpha_1 \cosh \alpha_2 \sinh \alpha_3 + \sinh \alpha_1 \sinh \alpha_2 \cosh \alpha_3$

 α_1 , α_2 , α_3 are the parameters of the group. Along with the parameters α_i , the components $v_i = dx_i/dx_0$ of the coordinate velocity of the primed reference frame can also be used as group parameters. The parameters v_i and α_i are related by

$$v_{1} = (\tanh \alpha_{1} - \tanh \alpha_{2} \tanh \alpha_{3})/(1 - \tanh \alpha_{1} \tanh \alpha_{2} \tanh \alpha_{3}),$$

$$v_{2} = (\tanh \alpha_{2} - \tanh \alpha_{1} \tanh \alpha_{3})/(1 - \tanh \alpha_{1} \tanh \alpha_{2} \tanh \alpha_{3}),$$

$$v_{3} = (\tanh \alpha_{3} - \tanh \alpha_{1} \tanh \alpha_{2})/(1 - \tanh \alpha_{1} \tanh \alpha_{2} \tanh \alpha_{3}).$$
(24)

The reverse relations have the form

$$\alpha_1 = \frac{1}{4} \ln \frac{(1+v_1-v_2+v_3)(1+v_1+v_2-v_3)}{(1-v_1-v_2-v_3)(1-v_1+v_2+v_3)} ,$$

$$\alpha_2 = \frac{1}{4} \ln \frac{(1-v_1+v_2+v_3)(1+v_1+v_2-v_3)}{(1-v_1-v_2-v_3)(1+v_1-v_2+v_3)} ,$$

$$\alpha_3 = \frac{1}{4} \ln \frac{(1-v_1+v_2+v_3)(1+v_1-v_2+v_3)}{(1-v_1-v_2-v_3)(1+v_1+v_2-v_3)} .$$

As for the generators X_i of the homogeneous 3-parametric group of isometries (22) of the space-time (21), they can be represented as follows

$$X_1 = -b_1 x_{\alpha} p_{\alpha} - (x_1 p_0 + x_0 p_1) + (x_2 p_3 + x_3 p_2),$$

$$X_2 = -b_2 x_{\alpha} p_{\alpha} - (x_2 p_0 + x_0 p_2) + (x_1 p_3 + x_3 p_1),$$

$$X_3 = -b_3 x_{\alpha} p_{\alpha} - (x_3 p_0 + x_0 p_3) + (x_1 p_2 + x_2 p_1),$$

where $p_{\alpha} = \partial/\partial x_{\alpha}$ are the generators of the 4-parametric group of translations. Thus, with inclusion of the latter, a inhomogeneous group of isometries of the entirely anisotropic Finsler space of events (21) is a 7-parametric group. As to its generators, they satisfy the commutation relation

$$\begin{split} [X_i X_j] &= 0, & [p_\alpha p_\beta] &= 0, \\ [X_1 p_0] &= b_1 p_0 + p_1, & [X_2 p_0] &= b_2 p_0 + p_2, & [X_3 p_0] &= b_3 p_0 + p_3, \\ [X_1 p_1] &= b_1 p_1 + p_0, & [X_2 p_1] &= b_2 p_1 - p_3, & [X_3 p_1] &= b_3 p_1 - p_2, \\ [X_1 p_2] &= b_1 p_2 - p_3, & [X_2 p_2] &= b_2 p_2 + p_0, & [X_3 p_2] &= b_3 p_2 - p_1, \\ [X_1 p_3] &= b_1 p_3 - p_2, & [X_2 p_3] &= b_2 p_3 - p_1, & [X_3 p_3] &= b_3 p_3 + p_0. \end{split}$$

Determining the distance dl between adjacent points by means of light signals leads to the conclusion that the flat entirely anisotropic 3D space is endowed with the discrete symmetry of a regular rhombic dodecahedron shown in Figure 3. Accordingly, the dynamic properties of a nonrelativistic massive particle residing in such an anisotropic space are determined by the inertial mass tensor

$$m_{\alpha\beta} = m \begin{pmatrix} (1 - b_1^2) & (b_3 - b_1 b_2) & (b_2 - b_1 b_3) \\ (b_3 - b_1 b_2) & (1 - b_2^2) & (b_1 - b_2 b_3) \\ (b_2 - b_1 b_3) & (b_1 - b_2 b_3) & (1 - b_3^2) \end{pmatrix}.$$
(25)

5. A curved space-time with $DISIM_b(2)$ local relativistic symmetry and local gauge invariance of its Finslerian metric

Currently, the problem of Lorentz symmetry violation is widely debated in the literature, in which case the Finslerian approach to the problem is becoming more and more popular. It is based on the assumption of the Finsler geometric structure of space-time. As for Finslerian generalization of the Einstein equations, the corresponding first attempts




Figure 3: A regular rhombic dodecahedron as an Euclidean image of 3D sphere of radius dl, prescribed in the flat entirely anisotropic Finslerian space.

were made in the early seventies. At that time, equations for the metric function of Finslerian space-time were proposed by analogy with the Einstein equations, but using the Finsler curvature tensors. Unfortunately, the physical interpretation of such equations turned out to be obscure. Although recent studies are more advanced from a physical point of view (see, in particular, an interesting paper [24]), it should again be noted that, in all likelihood, it is impossible to build physically meaningful Finslerian generalization of the Einstein theory of relativity without going beyond the framework of the purely geometric Finslerian analogies.

The fact is that the transition from Riemannian to Finslerian space-time necessarily leads to a violation of the local Lorentz symmetry. However, the violation of the local Lorentz symmetry of space-time does not necessarily mean a violation of the local relativistic symmetry. In the previous sections of this paper, it is shown that there are only two types of flat Finslerian event spaces, transitions to which from the Minkowski space violate the Lorentz symmetry of the event space without breaking relativistic symmetry.²

The flat Finslerian event spaces of the first type are relativistically symmetric spaces with partially broken 3D isotropy, i.e. with axial 3D symmetry. Their $DISIM_b(2)$ invariant Finslerian metric has the form (9) and describes axially symmetric crystalline states of space-time. In addition, such a metric satisfies the correspondence principle with the pseudo-Euclidean metric of the Minkowski space and therefore underlies viable

²Since the Lorentz symmetry is usually identified with relativistic symmetry (although, strictly speaking, these symmetries do not coincide), we clarify that relativistic symmetry of a flat event space means that it has at least a 3-parameter homogeneous non-compact group of isometries, while the Lorentz symmetry means that an homogeneous group of isometries is represented by the 6-parameter Lorentz group, which, in addition to 3-parameter non-compact transformations, contains the 3-parameter subgroup of 3D rotations.

Finslerian extension of the SR.

The flat Finslerian event spaces of the second type are relativistically symmetric spaces with entirely broken 3D isotropy. They are described by relativistically invariant Finslerian metric (21) and their relativistic symmetry is realized by means of 3-parameter homogeneous non-compact Abelian group of isometries. As for the respective non-homogeneous group, it is a 7-parameter group. It is also worth noting that flat Finslerian metric (21) describes an entirely anisotropic crystalline state of space-time with a discrete symmetry of a regular rhombic dodecahedron.

It is now clear that Finslerian metric describing a curved locally anisotropic space-time with $DISIM_b(2)$ local relativistic symmetry can be obtained from the Finslerian metric (19) of a flat space-time by means of the following replacement:

$$\eta_{ik} \to g_{ik}(x)$$
, $n_i \to n_i(x)$, $b \to b(x)$

. As a result, we arrive at the only possible viable Finslerian space-time model [19] with the metric

$$ds = \left[\frac{(n_i dx^i)^2}{g_{ik} dx^i dx^k} \right]^{b/2} \sqrt{g_{ik} dx^i dx^k}, \tag{26}$$

where $g_{ik} = g_{ik}(x)$ is the Riemannian metric tensor related to the gravity field; b = b(x) is the scalar field characterizing the magnitude of local space anisotropy, and $n_i = n_i(x)$ is the null-vector field indicating the locally preferred directions in space-time. So, the correct Finslerian extension of the general theory of relativity (under which the physical meaning of local relativistic space-time symmetry is preserved) must be based on the Finslerian space-time model (26). At each point of a curved Finslerian space time (26), the flat tangent Finslerian spaces have their own values of the parameters b and n_i . These values are none other than the value of the fields b(x) and $n_i(x)$ at the corresponding space-time points.

Obviously, the dynamics of a curved Finslerian spacetime (26) is completely determined by the dynamics of interacting fields $g_{ik}(x), b(x), n_i(x)$, and these fields form, along with the matter fields, a unified dynamic system. Therefore, in contrast to the existing purely geometric approaches to a Finslerian generalization of the Einstein equations, our approach to the same problem is based on the use of the methods of conventional field theory.

One can readily check by direct substitution that the metric (26), depending on the fields $g_{ik}(x)$, b(x), $n_i(x)$ and thereby determining a curved partially anisotropic Finslerian space-time, is invariant under the following local transformations of these fields

$$\begin{cases}
g_{ik}(x) \to \tilde{g}_{ik}(x) = \exp\{2\sigma(x)\} g_{ik}(x), \\
n_i(x) \to \tilde{n}_i(x) = \exp\{\sigma(x)[b(x)-1]/b(x)\} n_i(x), \\
b(x) \to \tilde{b}(x) = b(x),
\end{cases} (27)$$

where $\sigma(x)$ is an arbitrary function.

From the invariance of metric (26) under the local transformations (27) it follows that the Lagrangian and the action

$$S = -mc \int_{a}^{b} \left[\frac{(n_i(x)\dot{x}^i)^2}{g_{ik}(x)\dot{x}^i\dot{x}^k} \right]^{b(x)/2} \sqrt{g_{ik}(x)\dot{x}^i\dot{x}^k} d\lambda$$
 (28)

(here \dot{x}^i is the generalized velocity, λ is a parameter on a world line) for a particle in the external fields $g_{ik}(x)$, b(x) and $n_i(x)$, are also invariant under the above transformations, which entails the invariance of Hamilton's equations of motion

$$\frac{du_i}{ds} = \frac{1}{2} \left[\frac{\partial b}{\partial x^i} \ln \frac{(1+b)(n^k u_k)^2}{(1-b)g^{lm} u_l u_m} + \frac{\partial n^p}{\partial x^i} \frac{2b u_p}{n^k u_k} - \frac{\partial g^{nl}}{\partial x^i} \frac{(1+b)u_n u_l}{g^{jk} u_j u_k} \right], \tag{29}$$

$$\frac{dx^{i}}{ds} = \frac{(1+b)g^{ik}u_{k}}{q^{lm}u_{l}u_{m}} - b\frac{n^{i}}{n^{k}u_{k}},$$
(30)

where

$$u_l = (1-b)\frac{v_l}{v_k v^k} + b\frac{n_l}{n_k v^k}; \qquad v^k = dx^k/ds.$$

Moreover, it can readily be seen that the observable quantities of proper time

$$cd\tau = \left[\frac{n_0^2}{g_{00}}\right]^{b/2} \sqrt{g_{00}} dx^0 \tag{31}$$

and 3-space distances

$$dl^{2} = \left[\frac{n_{0}^{2}}{g_{00}}\right]^{b} \left(-g_{\alpha\beta} + \frac{g_{0\alpha}g_{0\beta}}{g_{00}}\right) dx^{\alpha} dx^{\beta}, \qquad (32)$$

as well as the action

$$S = -\frac{1}{c} \int \mu^* \left(\frac{n_i \, v^i}{\sqrt{g_{ik} \, v^i \, v^k}} \right)^{4b} \sqrt{-g} \, d^4 x \,,$$

for a compressible fluid are invariant under (27). (In this formula μ^* is the invariant fluid energy density, $v^i = dx^i/ds$, and ds is Finslerian metric (26))

All this suggests that the fields coupled by the transformations (27) describe the same physical situation and that the transformations (27) themselves have the meaning of local gauge transformations.

In connection with the mentioned local gauge invariance, the dynamical system consisting of the fields g_{ik} , b, n^i and a compressible fluid must be supplemented by two vector gauge fields A_i and B_i , that under local transformations (27) are transformed in the corresponding gradient manner. The A_i field for a certain class of problems is a pure gauge field, and the B_i field, whose gauge transformation has the form

$$B_i \to B_i + l[(b-1)\sigma(x)/b]_{;i}$$
,

where l is a constant with the dimensionality of length, interacts with the conserved rest mass current j^i , adding the term proportional to $B_i j^i$ to the full gauge invariant Lagrangian.

6. Conclusion

Alongside with the phenomena, for explanation of which the concept of dark matter or dark energy is commonly used, there are the data of astrophysical observations and particle physics, indirectly indicating the existence of a local space-time anisotropy in the Universe. In this respect, we should mention first of all the anisotropy of the accelerated expansion of the Universe and the possible anisotropy of the fine-structure constant, a unified Finslerian description of which was proposed in [20]. Note also the Greisen-Zatsepin-Kuzmin effect in the physics of ultra-high energy cosmic rays and other effects considered in [21]. In particular, the absence of the GZK cutoff has yet not been explained convincingly and still remains the main empirical fact which indirectly speaks in favor of violation of Lorentz symmetry and, therefore, in favor of local space-time anisotropy.

Actually, the first direct evidence of the existence of local space-time anisotropy was obtained by the CMS collaboration at LHC within the framework of the so-called Ridge/CMS effect [22]. Let us note in passing that the Ridge/CMS effect is characteristic only to the high multiplicity events. Such events take place in case of the central collision of the initial protons where the energy density at the moment of the collision is comparable to the energy density shortly after the Big Bang, when instead of hadrons there was quark-gluon plasma. Since the initial total momentum within the CMS experiment is equal to zero, the appearance of a preferred direction, which coincides with the direction of an elongated ridge of measured correlation function and with the protons collision axis, speaks of vacuum rearrangement with the appearance of anisotropic axially-symmetric condensate. On the one hand, quantum-field vacuum filled with the anisotropic condensate is the physical carrier of the local anisotropy of space-time and it can be regarded as an anisotropic quintessence, on the other – it imparts all the particles the properties of quasi-particles in the crystalline environment. In particular, in addition to the rest energies, all massive particles acquire the respective rest momenta $p|_{v=0} = bmcn$ and the anisotropy (18) of their inertial properties.

Generally speaking, in contrast to the Minkowski space with its constant quintessence, the flat Finslerian space (9) and its anisotropic quintessence are able to change their anisotropy if b is changeable. It easy to see that, at $b \to 1$, the Finslerian metric $ds = \left[(dx_0 - ndx) / \sqrt{dx_0^2 - dx^2} \right]^b \sqrt{dx_0^2 - dx^2}$ degenerates into the total differential $ds = dx_0 - ndx$. This means that the notion of spatial extent disappears and in the space-time there remains the single physical characteristic, namely, time duration and it should be regarded as an interval of absolute time. Besides at $b \to 1$ in accordance with (18) there also disappears the inertial mass of any particle. All this suggests that absolute time, where the very notions of spatial extent and inertial mass together with unobservable primordial quintessence become meaningless, is not a stable degenerate state of space-time. As a result of geometric phase transition, which accompanies a spontaneous breaking of the original gauge symmetry, such a primordial space-time may turn into anisotropic space-time, the anisotropy of which and, accordingly, the anisotropy of the quintessence decrease in the course of the accelerated expansion of the Universe.

In this regard, it is important to bear in mind that due to the directly proportional dependence of the rest momentum on the magnitude of local space-time anisotropy, as well as on mass of a body and speed of light, the fact that in our time the magnitude of the local space-time anisotropy is extremely small does not at all mean the impossibility of detecting anisotropy in astrophysical processes with a giant energy release and, accordingly, with a giant mass defect. Obviously, in the processes with giant mass

defects, one can also expect noticeable momentum defects. With allowance for the local law of conservation of momentum, this implies a possible self-acceleration of matter and local isotropization of the Universe. If this scenario proves to be correct, then the question posed by the authors of work [14], namely, why are the magnitude of the local space anisotropy b and the Λ -term simultaneously so small, may be answered.

References

- [1] H. Weyl, Gravitation und Elektrizität, Sitzber. preuss Akad. Wiss., Phys.–Math. Kl. (1918) 465
- [2] H. Weyl, Eine neue Erweiterung der Relativitätstheorie, Ann. Phys., Lpz. 59 (1919) 101
- [3] G.Yu. Bogoslovsky, From the Weyl theory to a theory of locally anisotropic space-time, Class. Quant. Grav. 9 (1992) 569
- [4] G.Yu. Bogoslovsky, Dynamic rearrangement of vacuum and the phase transitions in the geometric structure of space-time, *Int. J. Geom. Meth. Mod. Phys.* **09** (2012) 1250007
- [5] G. Bogoslovsky, The rest momentum as an additional property of a massive particle in Finsler space-time, J. Phys.: Conf. Ser. 1051 (2018) 012007.
- [6] C. Lämmerzahl and V. Perlick, Finsler geometry as a model for relativistic gravity, Int. J. Geom. Meth. Mod. Phys. 15 (2018) 1850166
- [7] G.Yu. Bogoslovsky, On a special relativistic theory of anisotropic space-time, *Dokl. Akad. Nauk SSSR* **213** (1973) 1055
- [8] J. Patera, P. Winternitz and H. Zassenhaus, Continuous subgroups of the fundamental groups of physics, *J. Math. Phys.* **16** (1975) 1597
- [9] P. Winternitz and I. Fris, Invariant expansions of relativistic amplitudes and subgroups of the proper Lorentz group, *Yadern. Fiz.* **1** (1965) 889
- [10] G.Yu. Bogoslovsky, Lorentz symmetry violation without violation of relativistic symmetry, *Phys. Lett. A*, **350** (2006) 5
- [11] G.Yu. Bogoslovsky, A special relativistic theory of the locally anisotropic space-time. Part 1. The metric and group of motions of the anisotropic space of events, *Nuovo Cimento B* **40** (1977) 99
- [12] G.Yu. Bogoslovsky, A special relativistic theory of the locally anisotropic space-time. Part 2. Mechanics and electrodynamics in the anisotropic space, *Nuovo Cimento B* 40 (1977) 116
- [13] G.Yu. Bogoslovsky, Subgroups of the group of generalized Lorentz transformations and their geometric invariants, Symmetry, Integrability and Geometry: Methods and Applications 1 (2005) 017
- [14] G.W. Gibbons, J. Gomis and C.N. Pope, General very special relativity is Finsler geometry, *Phys. Rev. D* **76** (2007) 081701(R)
- [15] D.V. Ahluwalia and S.P. Horvath, Very special relativity as relativity of dark matter: the Elko connection, *Journal of High Energy Physics* **11** (2010) 078
- [16] G.Yu. Bogoslovsky, On the local anisotropy of space-time, inertia and force fields, Nuovo Cimento B 77 (1983) 181
- [17] G.Yu. Bogoslovsky and H.F. Goenner, Concerning the generalized Lorentz symmetry and the generalization of the Dirac equation, *Phys. Lett. A* **323** (2004) 40
- [18] G.Yu. Bogoslovsky and H.F. Goenner, Finslerian spaces possessing local relativistic symmetry, Gen. Relativ. Gravit. 31 (1999) 1565

- [19] G.Yu. Bogoslovsky, A curved spacetime with $DISIM_b(2)$ local relativistic symmetry and local gauge invariance of its Finslerian metric, In: Physical Interpretations of Relativity Theory, Int. Conf. (Moscow, 1-5 July, 2019), Abstracts, Moscow: BMSTU (2019) p.23; ISBN 978-5-7038-5226-2
- [20] X. Li, H-N. Lin, S. Wang and Z. Chang, A unified description for dipoles of the fine-structure constant and SnIa Hubble diagram in Finslerian universe, *Eur. Phys. J. C* **75** (2015) 181
- [21] V. Balan *et al*, Geometrical models of the locally anisotropic space-time, *J. Mod. Phys.* **3** (2012) 1314
- [22] The CMS collaboration, V. Khachatryan, A.M. Sirunyan *et al*, Observation of long-range near-side angular correlations in proton-proton collisions at the LHC, *Journal of High Energy Physics* **9** (2010) 1
- [23] S. Dhasmana, Z.K. Silagadze, Finsler spacetime in light of Segal's principle, *Mod. Phys. Lett. A* (2019) doi.org/10.1142/S0217732320500194
- [24] M. Hohmann, C. Pfeifer, and N. Voicu, Finsler gravity action from variational completion, Phys. Rev. D 100 (2019) 064035