

Quantum scrambling and the growth of mutual information

Akram Touil¹ and Sebastian Deffner²

Department of Physics, University of Maryland, Baltimore County, Baltimore, MD 21250, USA

E-mail: ¹akramt1@umbc.edu

E-mail: ²deffner@umbc.edu

Abstract. Quantum information scrambling refers to the loss of local recoverability of quantum information, which has found widespread attention from high energy physics to quantum computing. In the present analysis we propose a possible starting point for the development of a comprehensive framework for the thermodynamics of scrambling. To this end, we prove that the growth of entanglement as quantified by the mutual information is lower bounded by the time-dependent change of Out-Of-Time-Ordered Correlator. We further show that the rate of increase of the mutual information can be upper bounded by the sum of local entropy productions, and the exchange entropy arising from the flow of information between separate partitions of a quantum system. Our results are illustrated for the ion trap system, that was recently used to verify information scrambling in an experiment, and for the Sachdev-Ye-Kitaev model.

1. Introduction

One of the most intriguing problems in theoretical physics is the information paradox [1–3], which suggests that physical information crossing the event horizon could permanently disappear in a black hole. In essence, the paradox originates in the unresolved incompatibility of current formulations of quantum mechanics and general relativity. To date, many possible solutions have been proposed, some as esoteric as the many-worlds interpretation of reality [2, 4], whereas others are rooted in quantum information theory [5, 6].

A particularly fruitful concept has been dubbed *quantum information scrambling* [7]. Within this paradigm, information that passes the event horizon is quickly and chaotically “scrambled” across the entirety of the horizon. Thus, the information only appears lost, as no *local* measurement allows to fully reconstruct the original quantum state [8–10]. In recent years, the study of quantum information scrambling has led to new physical concepts, such as the black hole complementarity and the holographical principle [11, 12]. In addition, information scrambling has found attention in high energy physics [13, 14], quantum information [15, 16], condensed matter physics [17, 18], and quantum thermodynamics [19–21].

Remarkably, it has also been recognized that exploiting the AdS/CFT duality [22] and the “ER=EPR-conjecture” [23], the information scrambling dynamics of black holes can be studied with analog quantum systems. Loosely speaking, the dynamics of two black holes connected through an Einstein-Rosen bridge can be mathematically

mapped onto the dynamics of entangled quantum systems. In fact, this idea led to the first verification of quantum information scrambling [24] in an ion trap experiment.

This ubiquity of information scrambling poses the question which underlying physical principles determine, if, when, and how information is distributed. Since information is physical [25], and its processing requires thermodynamic resources [26–31], it is only natural to realize that the second law of (quantum) thermodynamics [32] must hold the answer. The task is then to uniquely quantify the thermodynamics resources (such as heat or work) that are consumed while information is scrambled.

In the literature, a plethora of quantifiers have been proposed that can track and characterize quantum information scrambling, such as, for instance, Out-of-Time-Ordered Correlators (OTOCs) [33–36], the Loschmidt Echo [37], and versions of the mutual information [38–40]. Out of these variety of measures, the OTOC has probably gained the most attention. This is due to the fact that the properties of the OTOC characterize the dynamical emergence of “non conventional” quantum chaos [38, 41]. To the very best of our knowledge, however, a concise, transparent, and practically relevant relationship between the OTOC and a thermodynamic observable appears to be lacking ‡.

Therefore, in the present analysis we prove upper and lower bounds on the quantum mutual information. As main results, we find (i) that the time-dependent mutual information is lower bounded by the change of the OTOC, and (ii) that the rate of change of the mutual information is upper bounded by the sum of the stochastic entropy productions [42] in the separate partitions of a quantum system. Our findings are illustrated for the experimental system described in Ref. [24] and an example of a quantum chaotic system, the SYK model.

2. Time-dependent mutual information and the change of the OTOC

Imagine a quantum system S that can be separated into two partitions, A and B . The total system S evolves under unitary dynamics, and the quantum state is initially prepared as a product, $\rho_S(0) = \rho_A(0) \otimes \rho_B(0)$. Typically, quantum information scrambling then occurs in situations, in which $\rho_S(0)$ is chosen to be pure, and the unitary dynamics of S yields the growth of entanglement between A and B . This means, in particular, state tomography on only A is not longer sufficient to reconstruct $\rho_A(0)$ for any time $t > 0$.

The loss of local recoverability is conveniently characterized by the Out-of-Time-Ordered-Correlator (OTOC) [7–10], which can be written as,

$$\mathcal{O}(t) = \left\langle O_A^\dagger O_B^\dagger(t) O_A O_B(t) \right\rangle. \quad (1)$$

Here, O_A and O_B are *local* operators acting only on A and B , respectively. More specifically, we have $O_A \equiv o_A \otimes I_B$, $O_B \equiv I_A \otimes o_B$, and $O_B(t) = U^\dagger(t) O_B(0) U(t)$, where $U(t)$ denotes the unitary time evolution operator of S . It has been argued, that $\mathcal{O}(t)$ characterizes the spread of the operator $O_B(t)$ as it evolves in time, which tracks how information is scrambled from A to B [7–10]. Note that the average in Eq. (1) is often taken over a thermal state in S [7, 37], which is, however, not necessarily an instrumental choice [37].

‡ For obvious reasons, the projective measurements considered in Ref. [19] are neither feasible nor practical in complex many body systems




Figure 1. Sketch of a black hole scrambling quantum information. A_1 represents the information falling into the black hole, A_2 is the initial state of the black hole, A_4 is the Hawking radiation, and A_3 is the remaining black hole.

Note, however, that (to the best of our knowledge) there is no rigorous proof of the existence of operators correctly tracking information scrambling in any physical scenario. For instance, taking operators that commute with the Hamiltonian of S results in a time-invariant OTOC. Thus, one often takes an average over operators, rather than working with $\mathcal{O}(t)$ directly [43, 44].

2.1. Lower bound on quantum mutual information

For the following analysis, we will be motivated by the conceptual framework that maps notions from high energy physics onto a quantum information theoretic language [23, 45]. Now, imagine a situation in which some quantum information falls across the event horizon of a black hole, and we can describe the dynamics of that black hole by a scrambling (entangling) unitary map, $U(t)$. The set-up is depicted in Fig. 1. Quantum information is initially encoded at A_1 , and over sufficiently long time-scales the scrambled information will be re-emitted as Hawking radiation denoted by A_4 . Further, the initial state of the black hole is encoded in A_2 . For such scenarios, it has been shown [45] that

$$\langle O_{A_1} O_{A_4}(t) O_{A_1} O_{A_4}(t) \rangle_{\text{avg}} = 2^{-\mathcal{I}_{A_1, A_2 A_4}^{(2)}}, \quad (2)$$

where $\mathcal{I}_{i,j}^{(2)}$ is the Rényi-2 mutual information between the partitions i and j . For initially pure states[§], we can write

$$\mathcal{I}_{i,j}^{(2)} = \mathcal{S}_i^{(2)} + \mathcal{S}_j^{(2)} = -\ln(\text{tr} \{\rho_i^2\}) - \ln(\text{tr} \{\rho_j^2\}). \quad (3)$$

[§] For the sake of simplicity we only consider initially pure states. If the composite quantum state was initially mixed, we would obtain the same results up to additive constants

Furthermore, the average in Eq. (2) is taken over the Haar measure on the unitary group $U(d)$ with

$$\int_{\text{Haar}} dU = 1, \quad (4)$$

and where we have for an arbitrary function f and $\forall V \in U(d)$,

$$\int_{\text{Haar}} dU f(U) = \int_{\text{Haar}} dU f(VU) = \int_{\text{Haar}} dU f(UV). \quad (5)$$

It is interesting to note that for the quantum systems comprised of qubits, such as the experimental system analyzed in Ref. [24], the Haar average is equivalent to an average over the Pauli group for each operator [44].

Equation (2) can now be used to relate the OTOC with a thermodynamically relevant quantity, the quantum mutual information. To this end, consider

$$1 - \langle O_A O_B(t) O_A O_B(t) \rangle_{\text{avg}} = 1 - 2^{-\mathcal{I}_{A,B}^{(2)}} \leq \mathcal{I}_{A,B}^{(2)}, \quad (6)$$

where we identified $A_1 \equiv A$ and $A_2 A_4 \equiv B$. To justify this identification, we note that knowledge about the degrees of freedom A_2 and A_4 is enough to infer the information encoded in A_1 . The same argument holds for any closed quantum system and any unitary evolution $U(t)$. In this case, the analog of ‘‘Hawking radiation’’, are a subset of the degrees of freedom that is enough to reconstruct the initial information.

Note that the Rényi-2 mutual information is upper bounded by the quantum mutual information \mathcal{I} [46, 47]. This is a direct consequence of the strong subadditivity of the von Neumann entropy [46, 47]. Thus, we have

$$\mathcal{I}(t) = \mathcal{S}_A(t) + \mathcal{S}_B(t) - \mathcal{S}_S(t), \quad (7)$$

where $\mathcal{S}_i = -\text{tr} \{ \rho_i \ln(\rho_i) \}$ is the von Neumann entropy of system i with density matrix ρ_i . Note that $\mathcal{S}_S(t) = \mathcal{S}_S(0)$ for unitary dynamics. Thus, we immediately obtain

$$1 - \langle O_A O_B(t) O_A O_B(t) \rangle_{\text{avg}} \leq \mathcal{I}(t). \quad (8)$$

Now introducing the notation $\bar{\mathcal{O}}(t) \equiv \langle O_A O_B(t) O_A O_B(t) \rangle_{\text{avg}}$ and noticing that by definition $\bar{\mathcal{O}}(0) = 1$, we can write

$$\mathcal{I}(t) \geq \bar{\mathcal{O}}(0) - \bar{\mathcal{O}}(t), \quad (9)$$

which is true for all times $t > 0$, and which constitutes our first main result.

Equation (9) is a rigorous relationship between the mutual information and the change of the OTOC. In scrambling dynamics, $\bar{\mathcal{O}}(0) - \bar{\mathcal{O}}(t)$ is a monotonically growing function, and Eq. (9) asserts that also $\mathcal{I}(t)$ has to be growing. This is consistent with intuitive understanding of ‘‘scrambling’’, which should be equivalent to the growth of entanglement between A and B . We will now continue the analysis by illustrating Eq. (9) with two important examples, before we discuss the thermodynamic significance of $\mathcal{I}(t)$.

2.2. Information scrambling in experimentally relevant systems

2.2.1. Verified quantum information scrambling in ion traps

The experimental verification of quantum information scrambling [24] was conducted with a 7-qubit




Figure 2. Mutual information \mathcal{I} (blue, top line), change of the OTOC $\Delta\mathcal{O}(t)$ (red, middle line), and change of the modified OTOC $\Delta\mathcal{MO}$ (orange, bottom line) as function of time.

fully-connected quantum computer with a family of 3-qubit entangling unitaries $U(t)$. These entangling (scrambling) unitaries were constructed from a combination of 1-qubit and 2-qubit gates. Due to the experimental specifics, the observables O_A and O_B had to be of special form. Therefore, Ref. [24] considered a modified version of the OTOC, namely

$$\mathcal{MO}(t) = \sum_{\phi, O_P} \langle O_1^\dagger O_P^\dagger(t) O_1 O_P(t) \rangle, \quad (10)$$

where $O_1 \equiv |\psi\rangle\langle\phi|$ acts on the first qubit, $|\psi\rangle$ denotes the state of the qubit, and $|\phi\rangle$ is the teleported state (the last qubit of the experiment). Moreover, $O_P(t)$ are Pauli matrices evolved by a scrambling unitary in the Heisenberg picture, and the average is taken over all Pauli matrices and state vectors $|\phi\rangle$.

It is relatively easy to see that $\Delta\mathcal{MO} \simeq \Delta\mathcal{O}$ at times close to zero, since $O_P(t)$ has a simple form (in terms of operator complexity). However, in general we have $\Delta\mathcal{MO} \leq \Delta\mathcal{O}$, since the specific average taken in Ref. [24] to compute \mathcal{MO} is only an approximate 1-design, which does not capture the complex dynamics of $O_P(t)$ as the operator spreads to the other support (becomes non-local) with time.

In Fig. 2 we plot the mutual information (7), together with $\Delta\mathcal{O}$ (9) and $\Delta\mathcal{MO}$ (10) for the scrambling dynamics of Ref. [24]. As in Ref. [24], $B = \{2, 3, 4, 5, 6, 7\}$ refers to the qubits in the experiment, and $A = \{1\}$ is the first qubit. We observe that all quantities are monotonically increasing functions of time t .

Furthermore, $\mathcal{I}(0) = 0$ indicates the absence of any scrambling in the system, while $\mathcal{I}(t) = 2 \ln(2)$ indicates maximal scrambling: the maximum value of the information that B can know about A (or that A can know about B) is reached. This accentuates an important advantage of \mathcal{I} as a measure of scrambling over \mathcal{MO} . In general, $\mathcal{I}_{\max}(t_*) = \min\{d_A, d_B\} \ln(2)$ is the maximum value of \mathcal{I} at the scrambling time, t_* , whereas the maximal value of \mathcal{MO} depends on the specifics of the performed experiment.

|| The exact and rather lengthy expressions for $U(t)$ can be found in the methods section of Ref. [24].




Figure 3. Mutual information $\mathcal{I}(t)$ (blue, upper line) and $\Delta\mathcal{O}$ (red, lower line) for the SYK model with $N = 10$ Majorana fermions, $q = 4$, $J^2=2$, and averaged over 300 realizations, where in each realization we generate a new Hamiltonian with different random interaction terms.

2.2.2. Quantum chaotic dynamics – the SYK model As a second example to illustrate Eq. (9) we choose the Sachdev-Ye-Kitaev (SYK) model [48–50]. This is an exactly solvable, chaotic many-body system consisting of N interacting Majorana fermions with random interactions between q of these fermions (q taken as an even number). The SYK model has found important applications, for instance, as a quantum gravity model of a 1 + 1-dimensional black hole [50] in the limit of large N .

The Hamiltonian can be written as

$$H = (i)^{\frac{q}{2}} \sum_{1 \leq i_1 < i_2 < \dots < i_q \leq N} J_{i_1 i_2 \dots i_q} \psi_{i_1} \psi_{i_2} \dots \psi_{i_q}, \quad (11)$$

where $J_{i_1 i_2 \dots i_q}$ are real independent random variables with values drawn from a Gaussian distribution with mean $\langle J_{i_1 \dots i_q} \rangle = 0$ and variance $\langle J_{i_1 \dots i_q}^2 \rangle = J^2 (q - 1)! / N^{q-1}$, the parameter J (in the variance) sets the scale of the Hamiltonian. Further, ψ_i are Majorana field operators for $i \in \{1, \dots, N\}$.

Remarkably, the problem can be mapped from interacting Majorana fermions to interacting qubits with random interaction terms by using the Jordan-Wigner transformation [51, 52]. For the exact mapping between interacting Majorana fermions and interacting spins, we follow the notation of Ref. [52], to get

$$\psi_{2j} = \frac{1}{\sqrt{2}} \left(\prod_{i=1}^{N/2-1} \sigma_i^z \right) \sigma_{N/2}^y, \quad \psi_{2j-1} = \frac{1}{\sqrt{2}} \left(\prod_{i=1}^{N/2-1} \sigma_i^z \right) \sigma_{N/2}^x, \quad (12)$$

such that $\forall i, j \in \{1, \dots, N\}$ we have $\{\psi_i, \psi_j\} = \delta_{ij}$. Notice that Eq. (12) demonstrates that N Majorana fermions can be represented by a string of $N/2$ Pauli operators.

For the present purposes, we choose the first qubit to be subsystem A and the complement of A constitutes subsystem B . In Fig. 3 we plot of the mutual information and together with $\Delta\mathcal{O}$, where for $N = 10$ Majorana fermions and $q = 4$ interacting




Figure 4. Sketch of the two subsystems A and B as represented in continuous variables.

Majorana fermions, which gives a total of 210 terms in Eq. (11). We observe that both $\Delta\mathcal{O}$ and \mathcal{I} are monotonically increasing functions with time. Moreover, we see that $\Delta\mathcal{O}$ is always upper bounded by the mutual information, and \mathcal{I} reaches a maximum value slightly lower than $2\ln(2)$ given that in each realization we have small oscillations due to finite size effects. Note that for large N the small oscillations (recurrences) will die out as the system becomes strongly chaotic.

In conclusion, Eq. (9) establishes a rigorous lower bound on the quantum mutual information and the Out-of-Time-Order Correlator. Thus, from a theoretical point of view the quantum mutual information has the same appealing properties as the OTOC in characterizing information scrambling. However, the mutual information has the added benefit that it is also a well-studied quantity in quantum thermodynamics [32], and it can be closely related to irreversible entropy production.

3. Stochastic entropy production in quantum scrambling

Quantum information scrambling is an inherently *dynamical* phenomenon. Especially in chaotic quantum systems it is, thus, important to understand the rate with which information is lost to local observation [41]. Moreover, from a thermodynamic point of view, it appears appealing to relate the rate with which the quantum mutual information, $\mathcal{I}(t)$, grows to the *local* entropy production in subsystems A and B . Therefore, motivated by analyses of the rate of information production [53–55], we now seek to upper bound the rate of change of $\mathcal{I}(t)$ in terms of the thermodynamic resources (locally) consumed while scrambling information.

3.1. Continuous quantum systems

We start by considering the continuity equation for S as expressed in continuous variables

$$\partial_t \rho_S(x, y; t) = -\nabla \cdot \mathbf{j}_S(x, y; t). \quad (13)$$

Here, $\rho_S(x, y; t) = \langle xy | \rho_S(t) | xy \rangle$ is the density function of the system evaluated in variables x and y . Without loss of generality and for the sake of simplicity, we choose x and y as the coordinates in which ρ_A and ρ_B are diagonal[¶], respectively, cf. Fig. 4. Finally, $\mathbf{j}_S(x, y; t)$ denotes the probability current.

The corresponding, *local* continuity equations are obtained by tracing out the corresponding other subsystem. In particular, we have $\int dx \rho_S(x, y; t) = \rho_B(x; t)$ and $\int dy \rho_S(x, y; t) = \rho_A(y; t)$. Therefore, we can write

$$\partial_t \rho_A(x; t) = -\nabla_x \cdot \mathbf{j}_A(x; t) + j_S(x, 0; t). \quad (14)$$

[¶] Choosing more general coordinates does not lead to any additional physical insight, and only makes the mathematical expressions messier.

and

$$\partial_t \rho_B(y; t) = -\nabla_y \cdot \mathbf{j}_B(y; t) - j_S(0, y; t), \quad (15)$$

where $j_S(x, 0; t)$ is a boundary term that describes the influx of information from A to B , and $j_S(0, y; t)$ is the flow from B to A .

Now, again using that at $t = 0$ subsystems A and B are prepared in a product state, we can write by simply taking the derivative of Eq. (7)

$$\dot{\mathcal{I}} = - \int dx (\partial_t \rho_A) \ln(\rho_A) - \int dy (\partial_t \rho_B) \ln(\rho_B). \quad (16)$$

Employing the local continuity equations (14) and (15), we thus have

$$\begin{aligned} \dot{\mathcal{I}} &= \int dx (\nabla_x \cdot \mathbf{j}_A) \ln(\rho_A) - \int dx j_S(x, 0; t) \ln(\rho_A) \\ &\quad + \int dy (\nabla_y \cdot \mathbf{j}_B) \ln(\rho_B) + \int dy j_S(0, y; t) \ln(\rho_B). \end{aligned} \quad (17)$$

The latter can be further simplified by partial integration, and we obtain

$$\begin{aligned} \dot{\mathcal{I}} &= - \int dx \mathbf{j}_A \cdot \nabla_x \ln(\rho_A) - \int dx j_S(x, 0; t) \ln(\rho_A) \\ &\quad - \int dy \mathbf{j}_B \cdot \nabla_y \ln(\rho_B) + \int dy j_S(0, y; t) \ln(\rho_B), \end{aligned} \quad (18)$$

for which it is now easy to find upper bounds.

Using the trivial inequality, $\dot{\mathcal{I}} \leq |\dot{\mathcal{I}}|$, and then bounding the absolute value with the Cauchy-Schwarz inequality we can write [54]

$$\begin{aligned} \dot{\mathcal{I}} &\leq \alpha \left(\int dx \frac{\mathbf{j}_A^2}{\rho_A} \right)^{1/2} + \gamma_1 \left(\int dx \frac{j_S(x, 0; t)}{\rho_S} \right)^{1/2} \\ &\quad + \beta \left(\int dy \frac{\mathbf{j}_B^2}{\rho_B} \right)^{1/2} + \gamma_2 \left(\int dy \frac{j_S(0, y; t)}{\rho_S} \right)^{1/2}, \end{aligned} \quad (19)$$

where we introduced the Frieden integrals [53, 54],

$$\alpha = \int dx \rho_A (\nabla_x \ln(\rho_A))^2 \quad \text{and} \quad \beta = \int dy \rho_B (\nabla_y \ln(\rho_B))^2. \quad (20)$$

The Frieden integral is related to the Fisher information [54], but generally depends on the choice of variables, x and y , and the geometry of the quantum system S . Similarly, we have

$$\gamma_1 = \int dx \rho_S(x, 0; t) (\ln(\rho_A))^2 \quad \text{and} \quad \gamma_2 = \int dy \rho_S(0, y; t) (\ln(\rho_B))^2, \quad (21)$$

which are geometric terms corresponding to the flow of information across the boundary separating A and B .

However, we also immediately recognize the stochastic entropy production [56, 57] in subsystems A and B , which reads,

$$\dot{S}_A = \int dx \frac{\mathbf{j}_A^2}{\rho_A} \quad \text{and} \quad \dot{S}_B = \int dy \frac{\mathbf{j}_B^2}{\rho_B}. \quad (22)$$

In conclusion, we obtain

$$\dot{\mathcal{I}} \leq \alpha \left(\dot{\mathcal{S}}_A \right)^{1/2} + \beta \left(\dot{\mathcal{S}}_B \right)^{1/2} + \gamma |\dot{\mathcal{S}}_E|, \quad (23)$$

where we introduced $\gamma |\dot{\mathcal{S}}_E|$ to denote to the *exchange entropy* due to flow of information between A and B .

Equation (23) provides an intuitive way to think about information scrambling, or information flow between any arbitrary partitions A and B . The mutual information achieves a maximum if and only if the stochastic irreversible entropy productions within A and B as well as the entropy flow between A and B vanish. So far we have only considered scenarios where the dynamics of the quantum system is driven by information flow. A fully thermodynamic formalism, where a system can be in contact with an information reservoir as well as the usual heat and work reservoirs [29], will require a thoroughly developed conceptual framework which is beyond the scope of the present analysis.

3.2. Discrete quantum systems

The above analysis can be generalized to discrete representations of S . To this end, we consider the von Neumann equation describing the unitary dynamics of ρ_S

$$\partial_t \rho_S = -\frac{i}{\hbar} [H, \rho_S]. \quad (24)$$

with which the rate of change of $\mathcal{I}(t)$ (7) can be written as

$$\dot{\mathcal{I}} = \frac{i}{\hbar} [\text{tr} \{ [H, \rho_S] (\ln(\rho_A) \otimes I_B) \} + \text{tr} \{ [H, \rho_S] (I_A \otimes \ln(\rho_B)) \}], \quad (25)$$

which is mathematical a little more tedious than the continuous case. Therefore, we relegate the technical details of the derivation to the appendix.

Expressing the quantum states in Fock-Liouville space and after straight forward manipulations we again find

$$\dot{\mathcal{I}} \leq \mathcal{A} \dot{\mathcal{S}}_A + \mathcal{B} \dot{\mathcal{S}}_B + \mathcal{C} |\dot{\mathcal{S}}_E|, \quad (26)$$

where as before \mathcal{A} , \mathcal{B} , and \mathcal{C} are discrete versions of the Frieden integral [53, 54], that depend only on the geometry of the problem. Furthermore, $\dot{\mathcal{S}}_A$ and $\dot{\mathcal{S}}_B$ are the stochastic irreversible entropy production [42] in A and B , respectively. Finally, $\dot{\mathcal{S}}_E$ is the entropy (or information) flow between A and B .

4. Concluding remarks

4.1. The thermodynamic limit

We conclude the analysis with a few remarks on thermodynamic implications. To this end, note that both quantum systems, A and B , can be considered as open systems, for which the respective other system plays the role of an “environment”. Imagine now that B is much larger than A in the sense that B becomes a heat reservoir for A . For such a scenario it was shown in Ref. [57] that the thermodynamic entropy, $\mathcal{S}_S(t)$, is directly related to the *correlation entropy*, \mathcal{S}_{cor} , and we have

$$\mathcal{S}_S(t) = \mathcal{S}_A(t) + \mathcal{S}_B(t) + \mathcal{S}_{cor}(t). \quad (27)$$

Comparing the latter with the definition of the quantum mutual information (7), we immediately conclude $\mathcal{S}_{cor}(t) = -\mathcal{I}(t)$ for unitary dynamics.

This observation indicates that the chaotic spread of quantum information is intimately related to thermalization in quantum systems – a conclusion that can hardly be substantiated by looking only at the OTOC.

4.2. Summary

The present analysis provided a possible starting point for the development of a comprehensive framework for the thermodynamics of information scrambling. In particular, we related the OTOC with thermodynamically relevant quantities by proving that the change of the OTOC sets a lower bound on the time-dependent quantum mutual information. This bound was demonstrated for two experimentally relevant scenarios, namely for a system of trapped ions and the SYK model. We further showed that the rate of increase of the mutual information is upper bounded by the sum of local stochastic entropy productions, and the flow of entropy between separate partitions of the quantum system.

Possible applications of our work may be sought in the study of the thermodynamics of quantum chaotic systems, and the dynamical emergence of classicality from quantum mechanics. For each of these cases, our results provide a possible route to quantify the thermodynamics resources, such as work and heat, that are consumed while information is scrambled.

Acknowledgments

Fruitful discussions with Nathan M. Myers are gratefully acknowledged. This research was supported by grant number FQXi-RFP-1808 from the Foundational Questions Institute and Fetzer Franklin Fund, a donor advised fund of Silicon Valley Community Foundation (S.D).

7. Appendix: Entropy production in Fock-Liouville space

This appendix is dedicated to the technical details that lead to Eq. (26). To this end, we express all quantum states in Fock-Liouville space [58]. In this formalism operators are represented as vectors and superoperators as operators, which is convenient to numerically simplify computations in Hilbert space.

We introduce the notation [58], $\rho_S \equiv |\rho_S\rangle$ and $\langle\rho|\sigma\rangle \equiv \text{tr}\{\rho^\dagger\sigma\}$, which defines a pre-Hilbert space and completeness is guaranteed by definition. Thus, the von Neumann equation becomes [59],

$$|\dot{\rho}_S\rangle = W |\rho_S\rangle \quad \text{and} \quad W = \frac{1}{i\hbar} (H \otimes I - I \otimes H^\top). \quad (\text{B.1})$$

Note, that in contrast to classical stochastic dynamics that are described by rate matrices, the dynamics in Fock-Liouville space is determined by a skew hermitian matrix, W .

Thus, the rate of change of $\mathcal{I}(t)$ (7) can be expressed as

$$\dot{\mathcal{I}} = - \sum_{m,m'} W_{m,m'} \rho_{m'} \ln(\rho'_{Am}) - \sum_{m,m'} W_{m,m'} \rho_{m'} \ln(\rho'_{Bm}), \quad (\text{B.2})$$

where $\rho'_A \equiv \rho_A \otimes I_B$ and $\rho'_B \equiv I_A \otimes \rho_B$. Using standard tricks from stochastic thermodynamics of adding and subtracting terms we write

$$\begin{aligned} \dot{\mathcal{I}} &= \sum_{m,m'} \rho_{m'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Am'}}{W_{m',m} \rho'_{Am}} \right) + \sum_{m,m'} \rho_{m'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Bm'}}{W_{m',m} \rho'_{Bm}} \right) \\ &+ 2 \sum_{m,m'} \rho_{m'} W_{m,m'} \ln \left(\frac{W_{m',m}}{W_{m,m'}} \right) \\ &- \sum_{m'} \ln(\rho'_{Am'}) \cdot \left(\sum_m W_{m,m'} \right) \cdot \rho_{m'} - \sum_{m'} \ln(\rho'_{Bm'}) \cdot \left(\sum_m W_{m,m'} \right) \cdot \rho_{m'}, \end{aligned} \quad (\text{B.3})$$

which is not as involved as it looks. In particular, note that we have independent of the choice of basis, $\sum_{m,m'} W_{m,m'} = 0$, and hence

$$\begin{aligned} \dot{\mathcal{I}} &= \sum_{m,m'} \rho_{m'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Am'}}{W_{m',m} \rho'_{Am}} \right) + \sum_{m,m'} \rho_{m'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Bm'}}{W_{m',m} \rho'_{Bm}} \right) \\ &+ 2 \sum_{m,m'} \rho_{m'} W_{m,m'} \ln \left(\frac{W_{m',m}}{W_{m,m'}} \right). \end{aligned} \quad (\text{B.4})$$

In complete analogy to the continuous case (18) we can now upper bound $\dot{\mathcal{I}}$ as

$$\begin{aligned} \dot{\mathcal{I}} &\leq \sum_{m,m'} \left| \rho_{m'} \rho'^{-1}_{Am'} \cdot \rho'_{Am'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Am'}}{W_{m',m} \rho'_{Am}} \right) \right| \\ &+ \sum_{m,m'} \left| \rho_{m'} \rho'^{-1}_{Bm'} \cdot \rho'_{Bm'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Bm'}}{W_{m',m} \rho'_{Bm}} \right) \right| + \mathcal{C} |\dot{S}_E|, \end{aligned} \quad (\text{B.5})$$

where we already introduced the exchange entropy production

$$\mathcal{C} |\dot{S}_E| = \mathcal{A} \sum_{m,m'} \left| \rho'_{Am'} W_{m,m'} \ln \left(\frac{W_{m,m'}}{W_{m',m}} \right) \right| + \mathcal{B} \sum_{m,m'} \left| \rho'_{Bm'} W_{m,m'} \ln \left(\frac{W_{m,m'}}{W_{m',m}} \right) \right|, \quad (\text{B.6})$$

with $\mathcal{A} = \sum_{m,m'} |\rho_{m'} \rho'^{-1}_{Am'}|$ and $\mathcal{B} = \sum_{m,m'} |\rho_{m'} \rho'^{-1}_{Bm'}|$. Note that this is an identification only by analogy, as W is not a proper rate matrix. Since S is closed and evolves under unitary dynamics, the exchange entropy describes the flow of information between A and B .

The first two terms in Eq. (B.5) can be further simplified to read

$$\dot{\mathcal{I}} \leq \mathcal{A} \dot{S}_A + \mathcal{B} \dot{S}_B + \mathcal{C} |\dot{S}_E|, \quad (\text{B.7})$$

where we finally introduced the *local*, stochastic entropy production [42]

$$\dot{S}_A = \sum_{m,m'} \left| \rho'_{Am'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Am'}}{W_{m',m} \rho'_{Am}} \right) \right| \quad (\text{B.8})$$

and

$$\dot{S}_B = \sum_{m,m'} \left| \rho'_{Bm'} W_{m,m'} \ln \left(\frac{W_{m,m'} \rho'_{Bm'}}{W_{m',m} \rho'_{Bm}} \right) \right|. \quad (\text{B.9})$$

References

- [1] S. W. Hawking. Black holes and thermodynamics. *Phys. Rev. D*, 13:191, 1976.
- [2] J. Preskill. Do black holes destroy information? In *Proceedings of the International Symposium on Black Holes, Membranes, Wormholes and Superstrings*, S. Kalara and DV Nanopoulos, eds. (World Scientific, Singapore, 1993) pp, page 22. World Scientific, 1992.
- [3] S. D. Mathur. The information paradox: a pedagogical introduction. *Class. Quantum Grav.*, 26:224001, 2009.
- [4] N. J. Popławski. Cosmology with torsion: An alternative to cosmic inflation. *Phys. Lett. B*, 694:181, 2010.
- [5] R. Kallosh and A. Linde. Strings, black holes, and quantum information. *Phys. Rev. D*, 73:104033, 2006.
- [6] S. B. Giddings. Black holes, quantum information, and unitary evolution. *Phys. Rev. D*, 85:124063, 2012.
- [7] B. Swingle. Unscrambling the physics of out-of-time-order correlators. *Nat. Phys.*, 14:988, 2018.
- [8] K. Hashimoto, K. Murata, and R. Yoshii. Out-of-time-order correlators in quantum mechanics. *JHEP*, 2017:138, 2017.
- [9] Y. Sekino and L. Susskind. Fast scramblers. *JHEP*, 2008:065, 2008.
- [10] P. Hayden and J. Preskill. Black holes as mirrors: quantum information in random subsystems. *JHEP*, 2007:120, 2007.
- [11] L. Susskind, L. Thorlacius, and J. Uglum. The stretched horizon and black hole complementarity. *Phys. Rev. D*, 48:3743, 1993.
- [12] L. Susskind. The world as a hologram. *J. Math. Phys.*, 36:6377, 1995.
- [13] Y. Ling, P. Liu, and J.-P. Wu. Holographic butterfly effect at quantum critical points. *JHEP*, 2017:25, 2017.
- [14] S. Grozdanov, K. Schalm, and V. Scopelliti. Black hole scrambling from hydrodynamics. *Phys. Rev. Lett.*, 120:231601, 2018.
- [15] J. Couch, S. Eccles, P. Nguyen, B. Swingle, and S. Xu. The speed of quantum information spreading in chaotic systems. *arXiv preprint arXiv:1908.06993*, 2019.
- [16] B. Yoshida and N. Y. Yao. Disentangling scrambling and decoherence via quantum teleportation. *Phys. Rev. X*, 9:011006, 2019.
- [17] M. Blake, R. A. Davison, S. Grozdanov, and H. Liu. Many-body chaos and energy dynamics in holography. *JHEP*, 2018:35, 2018.
- [18] E. Iyoda and T. Sagawa. Scrambling of quantum information in quantum many-body systems. *Phys. Rev. A*, 97:042330, 2018.
- [19] M. Campisi and J. Goold. Thermodynamics of quantum information scrambling. *Phys. Rev. E*, 95:062127, 2017.
- [20] N. Yunger Halpern. Jarzynski-like equality for the out-of-time-ordered correlator. *Phys. Rev. A*, 95:012120, 2017.
- [21] A. Chenu, J. Molina-Vilaplana, and A. del Campo. Work Statistics, Loschmidt Echo and Information Scrambling in Chaotic Quantum Systems. *Quantum*, 3:127, 2019.
- [22] J. Maldacena. The large-N limit of superconformal field theories and supergravity. *Int. J. Theo. Phys.*, 38:1113, 1999.
- [23] J. Maldacena and L. Susskind. Cool horizons for entangled black holes. *Fort. Phys.*, 61:781, 2013.
- [24] K. A. Landsman, C. Figgatt, T. Schuster, N. M. Linke, B. Yoshida, N. Y. Yao, and C. Monroe. Verified quantum information scrambling. *Nature*, 567:61, 2019.
- [25] R. Landauer. Irreversibility and heat generation in the computing process. *IBM J. Res. Dev.*, 5:183, 1961.
- [26] C. H. Bennett. Logical reversibility of computation. *IBM J. Res. Dev.*, 17:525, 1973.
- [27] C. H. Bennett. The thermodynamics of computation: a review. *Int. J. Theo. Phys.*, 21:905, 1982.
- [28] W. H Zurek. Reversibility and stability of information processing systems. *Phys. Rev. Lett.*, 53:391, 1984.
- [29] S. Deffner and C. Jarzynski. Information processing and the second law of thermodynamics: An inclusive, Hamiltonian approach. *Phys. Rev. X*, 3:041003, 2013.
- [30] J. M. R. Parrondo, J. M. Horowitz, and T. Sagawa. Thermodynamics of information. *Nat. Phys.*, 11:131, 2015.
- [31] A. B. Boyd and J. P. Crutchfield. Maxwell Demon Dynamics: Deterministic Chaos, the Szilard Map, and the Intelligence of Thermodynamic Systems. *Phys. Rev. Lett.*, 116:190601, 2016.
- [32] S. Deffner and S. Campbell. *Quantum Thermodynamics*. Morgan & Claypool Publishers, 2019.
- [33] J. Maldacena, S. H. Shenker, and D. Stanford. A bound on chaos. *JHEP*, 2016:106, 2016.

- [34] B. Swingle, G. Bentsen, M. Schleier-Smith, and P. Hayden. Measuring the scrambling of quantum information. *Phys. Rev. A*, 94:040302, 2016.
- [35] S. Nakamura, E. Iyoda, T. Deguchi, and T. Sagawa. Universal scrambling in gapless quantum spin chains. *Phys. Rev. B*, 99:224305, 2019.
- [36] P. D. Bergamasco, G. G. Carlo, and A. M. F. Rivas. OTOC, complexity and entropy in bi-partite systems. *arXiv preprint arXiv:1904.12830*, 2019.
- [37] B. Yan, L. Cincio, and W. H Zurek. Information scrambling and Loschmidt echo. *arXiv preprint arXiv:1903.02651*, 2019.
- [38] E. Iyoda and T. Sagawa. Scrambling of quantum information in quantum many-body systems. *Phys. Rev. A*, 97:042330, 2018.
- [39] A. Seshadri, V. Madhok, and A. Lakshminarayan. Tripartite mutual information, entanglement, and scrambling in permutation symmetric systems with an application to quantum chaos. *Phys. Rev. E*, 98:052205, 2018.
- [40] V. Alba and P. Calabrese. Quantum information scrambling after a quantum quench. *Phys. Rev. B*, 100:115150, 2019.
- [41] D. A. Roberts and B. Swingle. Lieb-Robinson bound and the butterfly effect in quantum field theories. *Phys. Rev. Lett.*, 117:091602, 2016.
- [42] M. Esposito and C. Van den Broeck. Three faces of the second law. I. Master equation formulation. *Phys. Rev. E*, 82:011143, 2010.
- [43] J. Cotler, N. Hunter-Jones, J. Liu, and B. Yoshida. Chaos, complexity, and random matrices. *JHEP*, 2017:48, 2017.
- [44] D. A Roberts and B. Yoshida. Chaos and complexity by design. *JHEP*, 2017:121, 2017.
- [45] N. R. Hunter-Jones. *Chaos and Randomness in Strongly-Interacting Quantum Systems*. PhD thesis, [California Institute of Technology](https://www.caltech.edu/content/caltech/library/publication/thesis/2018), 2018.
- [46] E. H. Lieb and M. B. Ruskai. Proof of the strong subadditivity of quantummechanical entropy. *J. Math. Phys.*, 14:267, 1973.
- [47] S. Verd. α -mutual information. In *2015 Information Theory and Applications Workshop (ITA)*, pages 1–6, 2015.
- [48] S. Sachdev and J. Ye. Gapless spin-fluid ground state in a random quantum heisenberg magnet. *Phys. Rev. Lett.*, 70:3339, 1993.
- [49] J. Maldacena and D. Stanford. Remarks on the Sachdev-Ye-Kitaev model. *Phys. Rev. D*, 94:106002, 2016.
- [50] V. Rosenhaus. An introduction to the SYK model. *J. Phys. A: Math. Theor.*, 52:323001, 2019.
- [51] C. D. Batista and G. Ortiz. Generalized Jordan-Wigner transformations. *Phys. Rev. Lett.*, 86:1082, 2001.
- [52] J. Cotler and N. Hunter-Jones. Spectral decoupling in many-body quantum chaos. *arXiv preprint arXiv:1911.02026*, 2019.
- [53] B. Nikolov and B. R. Frieden. Limitation on entropy increase imposed by Fisher information. *Phys. Rev. E*, 49:4815, 1994.
- [54] D. Brody and B. Meister. An upper bound for entropy production. *Phys. Lett. A*, 204:93, 1995.
- [55] S. Deffner. Quantum speed limits and the maximal rate of information production. *Phys. Rev. Research*, (in press) [arXiv:1910.06811], 2020.
- [56] U. Seifert. Stochastic thermodynamics: principles and perspectives. *Euro. Phys. J. B*, 64:423, 2008.
- [57] M. Esposito, K. Lindenberg, and C. Van den Broeck. Entropy production as correlation between system and reservoir. *New J. Phys.*, 12:013013, 2010.
- [58] D. Manzano. A short introduction to the Lindblad master equation. *AIP Advances*, 10:025106, 2020.
- [59] A. Royer. Wigner function in Liouville space: a canonical formalism. *Phys. Rev. A*, 43:44, 1991.