arXiv:2002.02947v3 [quant-ph] 26 Nov 2020

Adiabatic theorem for closed quantum systems initialized at finite temperature

Nikolai Il'in¹, Anastasia Aristova^{1,3}, and Oleg Lychkovskiy^{1,2,3}

¹ Skolkovo Institute of Science and Technology, Bolshoy Boulevard 30, bld. 1, Moscow 121205, Russia

² Department of Mathematical Methods for Quantum Technologies,

Steklov Mathematical Institute of Russian Academy of Sciences, 8 Gubkina St., Moscow 119991, Russia and

³ Laboratory for the Physics of Complex Quantum Systems, Moscow Institute of Physics and Technology,

Institutsky per. 9, Dolgoprudny, Moscow region, 141700, Russia

(Dated: January 4, 2022)

The evolution of a driven quantum system is said to be adiabatic whenever the state of the system stays close to an instantaneous eigenstate of its time-dependent Hamiltonian. The celebrated quantum adiabatic theorem ensures that such *pure state adiabaticity* can be maintained with arbitrary accuracy, provided one chooses a small enough driving rate. Here, we extend the notion of quantum adiabaticity to closed quantum systems initially prepared at finite temperature. In this case adiabaticity implies that the (mixed) state of the system stays close to a quasi-Gibbs state diagonal in the basis of the instantaneous eigenstates of the Hamiltonian. We prove a sufficient condition for the finite temperature adiabaticity. Remarkably, it implies that the finite temperature adiabaticity can be more robust than the pure state adiabaticity, particularly in many-body systems. We present an example of a many-body system where, in the thermodynamic limit, the finite temperature adiabaticity is maintained, while the pure state adiabaticity breaks down.

A concept of quantum adiabatic evolution Introduction. was introduced by Born and Fock in the early days of quantum mechanics [1, 2]. The concept pertains to a driven closed quantum system described by a time-dependent Hamiltonian. The evolution of the system is called adiabatic as long as the state of the system stays close to the time-dependent instantaneous eigenstate of the Hamiltonian. The celebrated adiabatic theorem [2, 3] states that adiabaticity can be maintained with any prescribed accuracy, provided the driving rate (i.e. the rate of change of the Hamiltonian) is chosen small enough. The adiabatic theorem enjoys a glorious history and a wide range of theoretical and practical applications, including dynamics of chemical reactions [4], population transfer between molecular vibrational levels [5, 6], theory of quantum topological order [7], quantized charge transport [8], quantum memory [9] and quantum adiabatic computation [10–12].

Nowadays there is a wealth of experimental techniques available to manipulate large quantum systems consisting of cold atoms in optical lattices, ions in ion traps, arrays of superconducting qubits and quantum dots *etc* [13]. However, these systems are rarely prepared in pure states. Rather, they are typically initialized at some finite temperature determined by the preparation protocol. Therefore the conventional concept of adiabaticity [1–3], which we refer to as *pure state adiabaticity* (PSA) in what follows, calls for extension to the case of finite temperature.

Here we define the *finite temperature adiabaticity* as the property by which the state of a system initially prepared at finite temperature stays close to the quasi-Gibbs state in the course of the unitary quantum evolution. The time-dependent quasi-Gibbs state, defined by eq. (12) below, is diagonal in the instantaneous eigenbasis of the Hamiltonian and has the same spectrum as the initial thermal state.

Clearly, if the driving rate is so low that the conditions

for PSA for any eigenstate are met, then the finite temperature adiabaticity is also present, irrespectively of the temperature. It turns out that, in fact, the finite temperature adiabaticity can be present at much higher driving rates. This follows from the finite temperature adiabatic condition proven in the present paper. Remarkably, the energy gaps do not enter this conditions directly, in contrast to the case of PSA. Instead, the role of the energy gaps is played by the temperature. This can be of particular importance for many-body systems, where energy gaps vanish in the thermodynamic limit, and the pure state adiabaticity typically breaks down whenever the driving rate is kept finite but the system size is increased [14, 15]. We provide a particular example of a many-body system where the finite temperature adiabaticity survives the thermodynamic limit, despite the pure state adiabaticity being broken.

The rest paper is organised as follows. We start from introducing required definitions and notions (most importantly, the notion of the quasi-Gibbs state). Then we state the adiabatic theorem for closed quantum systems prepared in thermal states and discuss its scope and implications. After that we illustrate the theorem by applying it to a particular many-body system. We conclude the paper by the summary and outlook. Technical details are relegated to the Supplementary material [16].

Preliminaries We describe an isolated driven quantum system by means of a time-dependent Hamiltonian. To introduce time dependence in a way convenient for our purposes, we consider a Hamiltonian H_s dependent on a parameter s and assume that s varies in time. Without loss of generality, we can assume that s is a linear function of time,

$$s = \omega t,$$
 (1)

where ω is the driving rate. The adiabatic limit is defined

$$\omega \to 0, \qquad t \to \infty, \qquad \omega t = \text{const} > 0.$$
 (2)

Let E_s^n and Φ_s^n be respectively eigenenergies and eigenvectors of H_s ,

$$H_s \Phi_s^n = E_s^n \Phi_s^n, \quad n = 1, 2, ..., d,$$
 (3)

where d is the dimension of the Hilbert space. We assume that E_s^n and Φ_s^n are continuously differentiable in s.

Importantly, H_s can be represented as

$$H_s = U_s H_s U_s^{\dagger}, \tag{4}$$

where U_s is a continuously differentiable unitary operator,¹ $U_0 = 1$, and \tilde{H}_s is an auxiliary operator with the same eigenvalues as H_s and the same eigenvectors as H_0 ,

$$\widetilde{H}_{s} = \sum_{n} E_{s}^{n} |n\rangle \langle n|, \qquad (5)$$

where $|n\rangle \equiv \Phi_0^n$. Note that time dependence enters H_s only through E_s^n . An important object in our study is the operator

$$V_s \equiv -iU_s^{\dagger} \,\partial_s U_s. \tag{6}$$

To characterize the spectrum, we define

$$\frac{1}{\mu_s} = \max_n \left| \frac{E_0^{n+1} - E_0^n}{E_s^{n+1} - E_s^n} \right| \tag{7}$$

and

$$\nu_s = \max_n |\partial_s \ln(E_s^{n+1} - E_s^n)|. \tag{8}$$

Often the spectrum of the driven Hamiltonian does not change with time, which we refer to as *isospectral driving*. In this case $\tilde{H}_s = H_0$, μ_s do not actually depend on s, and ν_s is identically zero. A particular simple instance of the isospectral driving is the *uniform* isospectral driving with

$$H_s = e^{isV} H_0 e^{-isV}.$$
 (9)

Here V coincides with V_s defined by eq. (6).

The state of the system ρ_t satisfies the von Neumann equation

$$i\partial_t \rho_t = [H_{\omega t}, \rho_t]. \tag{10}$$

We assume that at t = 0 the system is initialized in a thermal state,

$$p_0 = e^{-\beta H_0} / Z_0, \qquad Z_0 \equiv \operatorname{tr} e^{-\beta H_0}, \qquad (11)$$

 β being the inverse temperature.

If the system were prepared in an eigenstate (in particular, in the ground state, i.e. "at zero temperature"), the adiabatic theorem [2, 3, 11] would imply that for any given s one can choose sufficiently small ω so that the state of the

system at a (large) time $t = s/\omega$ is close (within a given error margin) to the corresponding instantaneous eigenstate. This is what we refer to as pure state adiabaticity (PSA).

When we turn to the case of finite temperatures, the first question we have to address is what state one should compare the dynamical state ρ_t with. If the conditions for PSA are met for any eigenstate, then ρ_t stays close to the *quasi-Gibbs* state given by (see also a recent ref. [17])

$$\theta_t^{\beta} \equiv Z_0^{-1} \sum_n e^{-\beta E_0^n} |\Phi_{\omega t}^n\rangle \langle \Phi_{\omega t}^n|.$$
(12)

We will prove that, in fact, this is also the case under different (and, generally, less stringent) conditions that those for PSA.

It should be emphasized that the quasi-Gibbs state (12) is diagonal in the time-dependent instantaneous eigenbasis of the Hamiltonian, but its spectrum does not change with time and coincides with the spectrum of the initial Gibbs state. The latter feature emerges because the spectrum of the density matrix ρ_t cannot be changed by the unitary evolution (10). For this reason the quasi-Gibbs state (12) is, in general, different from the instantaneous Gibbs state $\rho_t^{\beta} \equiv e^{-\beta H_{\omega t}}/\text{tr } e^{-\beta H_{\omega t}}$, whose spectrum varies with time.

In what follows we will need to quantify the difference between two mixed quantum states. To this end, we employ the trace distance

$$D_{\rm tr}(\rho_1, \rho_2) \equiv (1/2) \,{\rm tr}|\rho_2 - \rho_1|,$$
 (13)

which is known to have a straightforward operational meaning [18–21].

Adiabatic theorem for finite temperatures. Now we are in a position to state the following

Theorem: The trace distance between the dynamical state of the system ρ_t (initialized in the Gibbs state (11) and evolving according to the von Neumann equation (10)) and the quasi-Gibbs state θ_t^{β} (defined by eq. (12)) is bounded from above by

$$D_{tr}\left(\rho_{t},\theta_{t}^{\beta}\right) \leq \sqrt{\sqrt{2}\omega\beta} \left(\frac{1}{\mu_{\omega t}} \|V_{\omega t}\| + \int_{0}^{\omega t} \frac{1}{\mu_{s'}} \|\partial_{s'}V_{s'}\| ds' + \int_{0}^{\omega t} \frac{\nu_{s'}}{\mu_{s'}} \|V_{s'}\| ds' + \sqrt{2} \int_{0}^{\omega t} \frac{1}{\mu_{s'}} \|V_{s'}\|^{2} ds'\right)^{1/2}.$$
(14)

Here V_s , μ_s and ν_s are defined according to eqs. (6), (7) and (8), respectively, and $\| \dots \|$ refers to the operator norm.²

¹ Note that U_s is *not* an evolution operator.

 $^{^2}$ For our purposes, the operator norm $\| \dots \|$ can be defined as the maximum among absolute values of eigenvalues of the corresponding operator.

This theorem implies that ρ_t converges to θ_t^{β} in the adiabatic limit (2), provided the term in brackets remains finite. The proof of the theorem can be found in the Supplement [16].

Observe that the r.h.s. of the bound (14) vanishes in the limit of infinite temperature, $\beta = 0$. This is consistent with the simple fact that at the infinite temperature $\rho_t = \theta_t^{\beta=0} = 1/d$, and the evolution is adiabatic at any driving rate.

The theorem admits a particularly simple form in the case of the uniform isospectral driving (9):

Corollary: For the isospectrally and uniformly driven Hamiltonian (9) the bound (14) reads

$$D_{\rm tr}\left(\rho_t, \theta_t^\beta\right) \le \sqrt{\sqrt{2}\omega\beta} \, \|V\| \, \left(1 + \sqrt{2}\,\omega t \, \|V\|\right). \tag{15}$$

The corollary immediately implies that ρ_t converges to θ_t^{β} in the adiabatic limit (2) whenever ||V|| is finite.

Remarkably, energy gaps do not directly enter the bounds (14) and (15), in contrast to typical sufficient conditions for PSA [11] (see, however, [22, 23]). This is crucial for the robustness of adiabaticity in the thermodynamic limit, since the energy gaps vanish with increasing the system size. The system size may also enter the bounds (14) and (15) through $||V_s||$, $||\partial_s V_s||$ and (for the bound (14)) through μ_s , ν_s . When the above quantities are finite in the thermodynamic limit, the finite temperature adiabaticity survives in this limit even if the PSA fails. Below we consider a many-body system exhibiting such behavior.

Example. Consider a thin straight wire with N electrons and a quantum sensor which can be moved around the wire, see Fig. 1. We consider a toy model of the sensor consisting of a single quantum spin S with a magnetic moment μ_{magn} (not to be confused with μ_s defined in eq. (7)). The interaction between the spin and the electrons is mediated by the magnetic field produced by the electron motion.³ We consider the case of zero net current of electrons. Still, the interaction persists even in this case due to fluctuations of the current, both classical and quantum. The Hamiltonian of the system reads

$$H_{\alpha} = H^e + H_{\alpha}^{Se}, \tag{16}$$

where H_e is the Hamiltonian of electrons (we do not need its explicit form here), and

$$H_{\alpha}^{Se} = -\frac{\mu_{\text{magn}}}{2\pi r} \mathcal{J}\left(-\sin\alpha S_x + \cos\alpha S_y\right) \quad (17)$$

is the Hamiltonian of the magnetic field-mediated interaction between electrons and the spin. Here (S_x, S_y, S_z) are the components of the spin operator, \mathcal{J} is the operator of the electron current, r is the distance from the sensor to the



FIG. 1. (Color online) A quantum sensor with a single spin possessing a magnetic moment is moved around a wire along a circular trajectory. The net current through the wire is zero, however the electrons in the wire are still magnetically coupled to the spin due to fluctuations of the current, see eqs. (16), (17). The manybody adiabaticity of the electron-spin system at finite temperature is robust with respect to increasing the system size (i.e. the length of the wire). In contract, the pure state adiabaticity breaks down in the thermodynamic limit at any finite driving rate.

wire and α is the polar angle determining the position of the sensor, see Fig. 1.

We further assume that the sensor is moved along a circular trajectory around the wire with r = const and $\alpha = \omega t$. Then the Hamiltonian (16) can be cast in the form (9), $H_{\alpha} = e^{-i\alpha S_z} H_{\alpha=0} e^{i\alpha S_z}$, therefore the bound (15) with $V = -S_z$ applies. This bound implies that it suffices to choose

$$\omega \le \frac{\varepsilon^2}{\sqrt{2}\,\beta\,S\,(1+\sqrt{2}\,\alpha\,S)}\tag{18}$$

to move the sensor up to the angle α along the circular trajectory while maintaining adiabaticity with precision ε , $D_{\mathrm{tr}}\left(\rho_t, \theta_t^\beta\right)\Big|_{t=\alpha/\omega} \leq \varepsilon$.

Remarkably, the sufficient adiabatic condition (18) does not depend on the number of electrons. Thus the finite temperature adiabaticity is robust in the thermodynamic limit $N \to \infty, L \to \infty, A = \text{const}, \rho \equiv N/(LA) = \text{const},$ where L and A are, respectively, the length and the cross section of the wire, and ρ is the number density of electrons in the wire.

In contract, the pure state adiabaticity breaks down in the thermodynamic limit. This can be easily seen if periodic boundary conditions along the z direction are imposed on the electron wave functions. In this case the Hamiltonian (16) commutes with the current operator, the latter be-

³ We disregard the magnetic fields of the magnetic moments of electrons.

ing related to the total momentum of electrons, P_e ,

$$\mathcal{J} = \frac{e\rho A}{m_e N} P_e, \tag{19}$$

where e and m_e are the charge and the mass of the electron. As a result, the dynamics of the spin is governed by the effective Hamiltonian (17), where \mathcal{J} now refers to the eigenvalue of the current operator (19) in the eigenstate the system is initialized in. Since in the typical eigenstate from the Gibbs ensemble this eigenvalue is $O(1/\sqrt{N})$, the driving rate necessary to maintain the pure state adiabaticity also scales as $1/\sqrt{N}$ and vanishes in the thermodynamic limit (see a detailed analysis in the Supplement [16]).

Summary and outlook

To summarize, we have introduced the notion of finite temperature adiabaticity of an isolated quantum system and proved the finite temperature adiabatic theorem (14). The sufficient adiabatic condition which follows from this theorem does not contain energy gaps, in contrast to most of the adiabatic conditions for pure state adiabaticity. This indicates that the finite temperature adiabaticity can be more robust in the thermodynamic limit then the pure state adiabaticity. We confirm this expectation for the specific model (16). It should be noted that this robustness is consistent with earlier numerical observations that microcanonical mixed states are more robust to adiabaticity breaking than pure states [24].

It should be emphasized that our notion of adiabaticity refers to the many-body state of the system and is different from the notion of local adiabaticity [25–31]. The latter notion applies to the reduced density matrix of a subsystem coupled to a reservoir. The many-body adiabaticity implies the local adiabaticity, but not vice versa.

A considerable limitation of the bounds (14), (15) is that they contain the operator norms. For continuous systems operator norms of certain physically relevant operators (e.g. momentum) are infinite, which renders the bounds void. In fact, the operator norm can be replaced by the better behaved thermal averages in *some* of the terms in eqs. (14), (15), as we discuss in the Supplement [16]. However, at the moment we are not able to avoid the operator norms altogether, and leave the improvement of the bounds (14), (15) in this direction for further work.

Acknowledgements. We are grateful to V. Dobrovitski for a useful discussion. The work was supported by the Russian Science Foundation under the grant N^o 17-71-20158.

- Max Born, "Das adiabatenprinzip in der quantenmeehanik," Zeitschrift für Physik 40, 167 (1926).
- [2] Max Born and Vladimir Fock, "Beweis des adiabatensatzes," Zeitschrift für Physik 51, 165–180 (1928).
- [3] Tosio Kato, "On the adiabatic theorem of quantum mechanics," Journal of the Physical Society of Japan 5, 435–439 (1950).

- [4] Joel M Bowman, "Reduced dimensionality theory of quantum reactive scattering," The Journal of Physical Chemistry 95, 4960–4968 (1991).
- [5] U Gaubatz, P Rudecki, S Schiemann, and K Bergmann, "Population transfer between molecular vibrational levels by stimulated raman scattering with partially overlapping laser fields. a new concept and experimental results," The Journal of Chemical Physics **92**, 5363–5376 (1990).
- [6] Klaas Bergmann, Nikolay V Vitanov, and Bruce W Shore, "Perspective: Stimulated raman adiabatic passage: The status after 25 years," The Journal of chemical physics 142, 170901 (2015).
- [7] Jan Carl Budich and Björn Trauzettel, "From the adiabatic theorem of quantum mechanics to topological states of matter," physica status solidi (RRL)-Rapid Research Letters 7, 109–129 (2013).
- [8] DJ Thouless, "Quantization of particle transport," Physical Review B 27, 6083 (1983).
- [9] M. Fleischhauer and M. D. Lukin, "Quantum memory for photons: Dark-state polaritons," Phys. Rev. A 65, 022314 (2002).
- [10] E. Farhi, J. Goldstone, S. Gutmann, J. Lapan, A. Lundgren, and D. Preda, "A quantum adiabatic evolution algorithm applied to random instances of an np-complete problem," Science 292, 472 (2001).
- [11] Tameem Albash and Daniel A. Lidar, "Adiabatic quantum computation," Rev. Mod. Phys. **90**, 015002 (2018).
- [12] E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, "Quantum computation by adiabatic evolution," arXiv:quantph/0001106.
- [13] 2D Quantum Metamaterials: Proceedings of the 2018 NIST Workshop).
- [14] John Dirk Walecka Alexander L. Fetter, *Quantum Theory* of Many-Particle Systems, Dover Books on Physics (Dover Publications, 2003).
- [15] Oleg Lychkovskiy, Oleksandr Gamayun, and Vadim Cheianov, "Time scale for adiabaticity breakdown in driven many-body systems and orthogonality catastrophe," Phys. Rev. Lett. **119**, 200401 (2017).
- [16] See the supplementary material to this article for the proof of the finite temperature adiabatic theorem and analysis of the pure state adiabaticity in the spin-electron model.
- [17] AH Skelt and I D'Amico, "Characterizing adiabaticity in quantum many-body systems at finite temperature," arXiv:2004.05842 (2020).
- [18] Carl W Helstrom, "Quantum detection and estimation theory," Journal of Statistical Physics **1**, 231–252 (1969).
- [19] A S Holevo, "On quasiequivalence of locally normal states," Theor. Math. Phys. 13, 1071–1082 (1972).
- [20] A S Holevo, "Statistical decision theory for quantum systems," Journal of multivariate analysis 3, 337–394 (1973).
- [21] Mark M Wilde, *Quantum information theory* (Cambridge University Press, 2013).
- [22] Joseph E Avron and Alexander Elgart, "Adiabatic theorem without a gap condition," Communications in mathematical physics **203**, 445–463 (1999).
- [23] Stefan Teufel, "A note on the adiabatic theorem without gap condition," Letters in Mathematical Physics 58, 261–266 (2001).
- [24] R. Bürkle and J. R. Anglin, "Probabilistic hysteresis in an isolated quantum system: The microscopic onset of irre-

versibility from a quantum perspective," Phys. Rev. A **101**, 042110 (2020).

- [25] Walid K. Abou-Salem and Jürg Fröhlich, "Adiabatic theorems and reversible isothermal processes," Letters in Mathematical Physics 72, 153–163 (2005).
- [26] Walid K. Abou-Salem and Jürg Fröhlich, "Status of the fundamental laws of thermodynamics," Journal of Statistical Physics 126, 1045–1068 (2007).
- [27] Vojkan Jaksic and Claude-Alain Pillet, "A note on the landauer principle in quantum statistical mechanics," Journal of Mathematical Physics **55**, 075210 (2014).
- [28] Sven Bachmann, Wojciech De Roeck, and Martin Fraas, "The adiabatic theorem for many-body quantum systems," Phys. Rev. Lett. **119**, 060201 (2017).
- [29] Lorenzo Campos Venuti, Tameem Albash, Daniel A. Lidar, and Paolo Zanardi, "Adiabaticity in open quantum systems," Phys. Rev. A 93, 032118 (2016).
- [30] Tristan Benoist, Martin Fraas, Vojkan Jaksic, and Claude-Alain Pillet, "Full statistics of erasure processes: Isothermal adiabatic theory and a statistical landauer principle," arXiv:1602.00051 (2016).
- [31] Stefan Teufel, "Non-equilibrium almost-stationary states and linear response for gapped quantum systems," Communications in Mathematical Physics , 1–33 (2019).

SUPPLEMENTARY MATERIAL

Properties of μ_s and ν_s

Here we prove a Lemma about μ_s and ν_s required for the proof of the finite temperature adiabatic theorem. We introduce a shorthand notation

$$\Delta_{mn}(s) \equiv E_s^m - E_s^n. \tag{S1}$$

We assume that at a given s the spectrum is ordered:

$$\Delta_{mn}(s) \ge 0 \quad \text{for} \quad m > n. \tag{S2}$$

Let us show that μ_s and ν_s defined respectively by eqs. (7) and (8) of the main text, satisfy the following

Lemma:

$$\frac{1}{\mu(s)} = \max_{1 \le n < m \le d} \left| \frac{\Delta_{mn}(0)}{\Delta_{mn}(s)} \right|$$
(S3)

and

$$\nu(s) = \max_{1 \le n < m \le d} \left| \frac{\partial_s \Delta_{mn}(s)}{\Delta_{mn}(s)} \right|.$$
(S4)

Proof: Consider an arbitrary set of real numbers A_n , n = 1, 2, ..., d and introduce

$$q = \max_{m,n} \left| \frac{A_m - A_n}{E_s^m - E_s^n} \right|.$$
(S5)

We are going to prove that in fact

$$q = \max_{k} \left| \frac{A_{k+1} - A_{k}}{E_{s}^{k+1} - E_{s}^{k}} \right|.$$
 (S6)

This equality entails eq. (S3) for $A_n = E_0^n$ and eq. (S4) for $A_n = \partial_s E_s^n$.

To prove eq. (S6), we start from an obvious observation that

$$q \ge \max_{k} \left| \frac{A_{k+1} - A_{k}}{E_{s}^{k+1} - E_{s}^{k}} \right|.$$
(S7)

Let us show that, in fact, the strict inequality is impossible. To this end we assume the opposite, i.e. that

$$\left|\frac{A_{k+1} - A_k}{E_s^{k+1} - E_s^k}\right| < q \qquad \forall k.$$
(S8)

Then for any m > n we obtain

$$|A_m - A_n| = \left|\sum_{k=n}^{m-1} (A_{k+1} - A_k)\right| \le \sum_{k=n}^{m-1} |A_{k+1} - A_k| < q \sum_{k=n}^{m-1} (E_s^{k+1} - E_s^k) = q(E_s^m - E_s^n),$$
(S9)

where the ordering of energies, (S2), is used to get rid of the modulus. Eq. (S9) is inconsistent with eq. (S8). Thus the equality (S6) is true, q.e.d.

Proof of the finite temperature adiabatic theorem

Here we prove the bound (14) from the main text. We introduce

$$\sigma_t = U_{\omega t}^{\dagger} \rho_t U_{\omega t}, \tag{S10}$$

$$\partial_t \sigma_t = -i[\widetilde{H}_{\omega t} + \omega V_{\omega t}, \sigma_t], \qquad \sigma_0 = e^{-\beta H_0} / Z_0.$$
(S11)

We denote by dot the derivative of a function with respect to its argument $s = \omega t$. For example, $\dot{U}_s = \partial_s U_s$, $\dot{U}_{\omega t} =$ $\partial_s U_s|_{s=\omega t}$ but $\partial_t U_{\omega t} = \omega \dot{U}_{\omega t}$. We first estimate the quantity

$$D \equiv 1 - \operatorname{tr}(\sqrt{\theta_t^\beta}\sqrt{\rho_t}) = 1 - \operatorname{tr}(\sqrt{\rho_0}\sqrt{\sigma_t}).$$
(S12)

Note that

$$\partial_t \sqrt{\sigma_t} = -i [\tilde{H}_{\omega t} + \omega V_{\omega t}, \sqrt{\sigma_t}].$$
(S13)

Therefore

$$\partial_{t}D = -\operatorname{tr}(\sqrt{\rho_{0}}\,\partial_{t}\sqrt{\sigma_{t}}) = i\frac{1}{\sqrt{Z_{0}}}\operatorname{tr}(e^{-\beta H_{0}/2}\left[\widetilde{H}_{\omega t} + \omega V_{\omega t}, \sqrt{\sigma_{t}}\right]) = i\omega\frac{1}{\sqrt{Z_{0}}}\operatorname{tr}(\left[e^{-\beta H_{0}/2}, V_{\omega t}\right]\sqrt{\sigma_{t}})$$
$$= \frac{i\omega}{\sqrt{Z_{0}}}\sum_{n,k} \left(e^{-\frac{\beta}{2}E_{0}^{n}} - e^{-\frac{\beta}{2}E_{0}^{k}}\right) \langle n|V_{\omega t}|k\rangle \langle k|\sqrt{\sigma_{t}}|n\rangle$$
$$= \frac{i\beta\omega}{2\sqrt{Z_{0}}}\sum_{n,k} f_{nk}(\omega t)\Delta_{nk}(\omega t) \langle n|V_{\omega t}|k\rangle \langle k|\sqrt{\sigma_{t}}|n\rangle, \tag{S14}$$

where $|n\rangle = |\Phi_0^n\rangle$, $|k\rangle = |\Phi_0^k\rangle$ are eigenstates of H_0 and, consequently, of $\tilde{H}_{\omega t}$ for arbitrary t, E_0^n, E_0^k are eigenenergies of H_0 , and

$$f_{nk}(\omega t) \equiv \frac{e^{-\frac{\beta}{2}E_0^n} - e^{-\frac{\beta}{2}E_0^k}}{\beta \Delta_{nk}(\omega t)/2} \text{ for } n \neq k, \quad f_{nn} = 0.$$
(S15)

Note that we will occasionally drop an argument of the function f_{nk} when this does not lead to ambiguities.

We notice that

$$(E_{\omega t}^{n} - E_{\omega t}^{k}) \langle k | \sqrt{\sigma_{t}} | n \rangle = -\langle k | [\widetilde{H}_{\omega t}, \sqrt{\sigma_{t}}] | n \rangle = -i \langle k | \partial_{t} \sqrt{\sigma_{t}} | n \rangle + \omega \langle k | [V_{\omega t}, \sqrt{\sigma_{t}}] | n \rangle.$$
(S16)

Substituting this expression to eq. (S14) and integrating it over time one obtains

$$D = \frac{\beta\omega}{2\sqrt{Z_0}} \sum_{n,k} \int_0^t f_{nk}(\omega t') \langle n | V_{\omega t'} | k \rangle \left(\langle k | \partial_{t'} \sqrt{\sigma_{t'}} | n \rangle + i\omega \langle k | [V_{\omega t'}, \sqrt{\sigma_{t'}}] | n \rangle \right) dt'.$$
(S17)

Integrating (S17) by parts one gets

$$D = \frac{\beta\omega}{2\sqrt{Z_0}} \sum_{n,k} \left(f_{nk}(\omega t) \langle n | V_{\omega t} | k \rangle \langle k | \sqrt{\sigma_t} | n \rangle - \omega \int_0^t f_{nk}(\omega t') \langle n | \dot{V}_{\omega t'} | k \rangle \langle k | \sqrt{\sigma_{t'}} | n \rangle dt' \right. \\ \left. + \omega \int_0^t f_{nk}(\omega t') \langle n | V_{\omega t'} | k \rangle \frac{\dot{\Delta}_{nk}(\omega t')}{\Delta_{nk}(\omega t')} \langle k | \sqrt{\sigma_{t'}} | n \rangle dt' + i\omega \int_0^t f_{nk}(\omega t') \langle n | V_{\omega t'} | k \rangle \langle k | [V_{\omega t'}, \sqrt{\sigma_{t'}}] | n \rangle dt' \right) \\ \left. = \frac{\beta\omega}{2\sqrt{Z_0}} (K_1 + K_2 + K_3 + K_4), \right.$$
(S18)

where we us $\dot{f}_{nk}(\omega t) = -f_{nk}(\omega t)\dot{\Delta}_{nk}(\omega t)/\Delta_{nk}(\omega t)$ and

$$K_{1} = \sum_{n,k} f_{nk}(\omega t) \langle n | V_{\omega t} | k \rangle \langle k | \sqrt{\sigma_{t}} | n \rangle$$
(S19)

$$K_{2} = -\omega \sum_{n,k} \int_{0}^{t} f_{nk}(\omega t') \langle n | \dot{V}_{\omega t'} | k \rangle \langle k | \sqrt{\sigma_{t'}} | n \rangle dt'$$
(S20)

$$K_{3} = \omega \sum_{n,k} \int_{0}^{t} f_{nk}(\omega t') \langle n | V_{\omega t'} | k \rangle \frac{\dot{\Delta}_{nk}(\omega t')}{\Delta_{nk}(\omega t')} \langle k | \sqrt{\sigma_{t'}} | n \rangle dt'$$
(S21)

$$K_4 = i\omega \sum_{n,k} \int_0^t f_{nk}(\omega t') \langle n | V_{\omega t'} | k \rangle \langle k | [V_{\omega t'}, \sqrt{\sigma_{t'}}] | n \rangle dt'$$
(S22)

Obviously,

$$|D| \le \frac{\beta\omega}{2\sqrt{Z_0}} (|K_1| + |K_2| + |K_3| + |K_4|).$$
(S23)

Let us estimate $|K_1|$:

$$|K_1| \le \left(\sum_{n,k} f_{nk}^2 \langle n|V_{\omega t}|k\rangle \langle k|V_{\omega t}|n\rangle\right)^{1/2} \left(\sum_{n,k} \langle k|\sqrt{\sigma_t}|n\rangle \langle n|\sqrt{\sigma_t}|k\rangle\right)^{1/2}.$$
(S24)

The term in the second bracket reads

$$\sum_{n,k} \langle k | \sqrt{\sigma_t} | n \rangle \langle n | \sqrt{\sigma_t} | k \rangle = \operatorname{tr} \sigma_t = \operatorname{tr} \rho_0 = 1.$$
(S25)

To estimate the first term we need to estimate f_{nk}^2 :

$$f_{nk}^{2}(\omega t) = \left(\frac{e^{-\frac{\beta}{2}E_{0}^{n}} - e^{-\frac{\beta}{2}E_{0}^{k}}}{\beta(E_{0}^{n} - E_{0}^{k})/2} \frac{\Delta_{nk}(0)}{\Delta_{nk}(\omega t')}\right)^{2} \le \left(\frac{e^{-\frac{\beta}{2}E_{0}^{n}} - e^{-\frac{\beta}{2}E_{0}^{k}}}{\beta(E_{0}^{n} - E_{0}^{k})/2}\right)^{2} \frac{1}{\mu^{2}(\omega t)},$$
(S26)

where eq. (S3) is used to establish the inequality. Further, by the Lagrange's Mean Value Theorem there exists $a \in (0, 1)$ such that

$$\left(\frac{e^{-\frac{\beta}{2}E_0^n} - e^{-\frac{\beta}{2}E_0^k}}{\beta(E_0^n - E_0^k)/2}\right)^2 = e^{-\beta(aE_0^n + (1-a)E_0^k)} \le e^{-\beta\min\{E_0^n, E_0^k\}} \le e^{-\beta E_0^n} + e^{-\beta E_0^k}, \quad n \ne k.$$
(S27)

Combining inequalities (S26) and (S27) and extending them to the trivial case n = k (where $f_{nn} = 0$ by definition (S15)) we get

$$f_{nk}^{2}(\omega t) \leq \frac{1}{\mu^{2}(\omega t)} \left(e^{-\beta E_{0}^{n}} + e^{-\beta E_{0}^{k}} \right).$$
 (S28)

We use this bound to proceed further:

$$\sum_{n,k} f_{nk}^{2} \langle n | V_{\omega t} | k \rangle \langle k | V_{\omega t} | n \rangle$$

$$\leq \frac{1}{\mu^{2}(\omega t)} \sum_{n,k} \left(e^{-\beta E_{0}^{n}} \langle n | V_{\omega t} | k \rangle \langle k | V_{\omega t} | n \rangle + e^{-\beta E_{0}^{k}} \langle n | V_{\omega t} | k \rangle \langle k | V_{\omega t} | n \rangle \right)$$

$$= \frac{1}{\mu^{2}(\omega t)} \sum_{n,k} \left(\langle n | V_{\omega t} | k \rangle \langle k | V_{\omega t} e^{-\beta H_{0}} | n \rangle + \langle n | V_{\omega t} | k \rangle \langle k | e^{-\beta H_{0}} V_{\omega t} | n \rangle \right)$$

$$= 2 \frac{1}{\mu^{2}(\omega t)} \operatorname{tr}(V_{\omega t}^{2} e^{-\beta H_{0}})$$
(S29)

Next use the inequality

$$|\mathrm{tr}AB| \leqslant ||A||\mathrm{tr}B,\tag{S30}$$

valid for any B > 0 and diagonalisable A (be reminded that $\|...\|$ stands for the operator norm), to obtain

t

$$rV_{\omega t}^2 e^{-\beta H_0} \le \|V_{\omega t}^2\| \operatorname{tr} e^{-\beta H_0}.$$
(S31)

Finally

$$K_1 \le \sqrt{2Z_0} \frac{1}{\mu_{\omega t}} \| V_{\omega t} \|.$$
 (S32)

 K_2 can be bounded in an analogous way:

$$\left|\sum_{n,k} f_{nk}(\omega t') \langle n | \dot{V}_{\omega t'} | k \rangle \langle k | \sqrt{\sigma_{t'}} | n \rangle \right| \le \sqrt{2Z_0} \frac{1}{\mu_{\omega t'}} \| \dot{V}_{\omega t'} \|$$
(S33)

and

$$|K_2| \le \omega \sqrt{2Z_0} \int_0^t \frac{1}{\mu_{\omega t'}} \|\dot{V}_{\omega t'}\| dt' = \sqrt{2Z_0} \int_0^{\omega t} \frac{1}{\mu_{s'}} \|\dot{V}_{s'}\| ds'.$$
(S34)

Let us estimate $|K_3|$:

$$|K_3| \leq \int_0^t \left(\sum_{n,k} f_{nk}^2 \left(\frac{\dot{\Delta}_{nk}(\omega t')}{\Delta_{nk}(\omega t')} \right)^2 \langle n | V_{\omega t'} | k \rangle \langle k | V_{\omega t} | n \rangle \right)^{1/2} \left(\sum_{n,k} \langle k | \sqrt{\sigma_{t'}} | n \rangle \langle n | \sqrt{\sigma_{t'}} | k \rangle \right)^{1/2} dt', \tag{S35}$$

$$f_{nk}^{2}(\omega t') \left(\frac{\dot{\Delta}_{nk}(\omega t')}{\Delta_{nk}(\omega t')}\right)^{2} = \left(\frac{e^{-\frac{\beta}{2}E_{0}^{n}} - e^{-\frac{\beta}{2}E_{0}^{k}}}{\beta(E_{0}^{n} - E_{0}^{k})/2} \frac{\Delta_{nk}(0)\dot{\Delta}_{nk}(\omega t')}{\Delta_{nk}^{2}(\omega t')}\right)^{2} \le \frac{\nu^{2}(\omega t')}{\mu^{2}(\omega t')} (e^{-\beta E_{0}^{n}} + e^{-\beta E_{0}^{k}}), \tag{S36}$$

where eqs. (S3), (S4) are used to establish the inequality. Thus we obtain

$$|K_3| \le \sqrt{2Z_0} \int_0^{\omega t} \frac{\nu_{s'}}{\mu_{s'}} ||V_{s'}|| ds'$$
(S37)

Finally, let us estimate K_4 :

$$|K_4| \le \omega \int_0^t dt' \left(\sum_{n,k} f_{nk}^2 \langle n | V_{\omega t'} | k \rangle \langle k | V_{\omega t'} | n \rangle \right)^{1/2} \left(\sum_{n,k} \langle k | [V_{\omega t'}, \sqrt{\sigma_{t'}}] | n \rangle \langle n | [\sqrt{\sigma_{t'}}, V_{\omega t'}] | k \rangle \right)^{1/2}.$$
(S38)

The term in the first bracket has been already bounded, see eq. (S29). The term in the second bracket reads

$$\operatorname{tr}[V_{\omega t'}, \sqrt{\sigma_{t'}}]\left[\sqrt{\sigma_{t'}}, V_{\omega t'}\right] = 2\operatorname{tr}V_{\omega t'}^2 \sigma_{t'} - 2\operatorname{tr}\left(\sigma_{t'}^{1/4} V_{\omega t'} \sigma_{t'}^{1/4}\right)^2 \le 2\|V_{\omega t'}^2\|\operatorname{tr}\sigma_t = 2\|V_{\omega t'}\|^2,$$
(S39)

and we get

$$|K_4| \le 2\sqrt{Z_0} \,\omega \int_0^t \frac{1}{\mu_{\omega t'}} \|V_{\omega t'}\|^2 dt' = 2\sqrt{Z_0} \int_0^{\omega t} \frac{1}{\mu_{s'}} \|V_{s'}\|^2 ds'.$$
(S40)

Finally we collect all pieces (S32)–(S40) together and bound D according to eq. (S23):

$$D \leq \frac{\omega\beta}{\sqrt{2}} \left(\frac{1}{\mu_{\omega t}} \|V_{\omega t}\| + \int_0^{\omega t} \frac{1}{\mu(t')} \|\partial_{t'} V_{t'}\| dt' + \int_0^{\omega t} \frac{\nu(t')}{\mu(t')} \|V_{t'}\| dt' + \sqrt{2} \int_0^{\omega t} \frac{1}{\mu(t')} \|V_{t'}\|^2 dt' \right).$$
(S41)

The last thing we need to do is to connect D with the trace distance $D_{tr}\left(\rho_t, \theta_t^{\beta}\right)$. This can be done thanks to the inequality proven in [19] which reads

$$D_{\rm tr}\left(\rho_1,\rho_2\right) \le \sqrt{1 - \left({\rm tr}\sqrt{\rho_1}\sqrt{\rho_2}\right)^2} \le \sqrt{2D}.$$
(S42)

Eq. (14) of the main text follows from eqs. (S41) and (S42), q.e.d.

As was mentioned in the main text, the presence of operator norms in the final result makes the bound inapplicable in the cases where V or \dot{V} are unbounded operators. In fact, the operator norms are superficial for estimating K_1 , K_2 and K_3 above and can be substituted by thermal averages with respect to the initial Gibbs state. For K_1 this can be seen from the eq. (S29), and analogously for K_2 and K_3 . However, we were not able to avoid the operator norm when estimating K_4 .

Spin moved around a wire: pure state adiabaticity

Here we derive a condition for pure state adiabaticity in the electron-spin system (see (16),(17) of the main text) under the assumption of periodic boundary conditions for electron wave functions in the z-direction. To be specific, we choose

$$S = \frac{1}{2}.$$
(S43)

Using eq. (19) of the main text, we rewrite the Hamiltonian as

$$H_{\alpha} = H_e + \gamma \, e^{-i\alpha S_z} S_y \, e^{i\alpha S_z},\tag{S44}$$

where

$$\gamma = -\frac{\mu_{\text{magn}}}{2\pi r} \frac{e\rho A}{m_e N} P_e.$$
(S45)

The total momentum of electrons, P_e , commutes with H_{α} , therefore we treat it as a *c*-number. We initialize the system in an eigenstate of H_{α} ,

$$\Psi_0 = |\text{electrons}\rangle \otimes \psi_0, \tag{S46}$$

where $|\text{electrons}\rangle$ is an eigenstate of H_e and P_e , while ψ_0 is an eigenstate of S_y ,

$$S_y \psi_0 = \frac{1}{2} \psi_0. \tag{S47}$$

The time-dependent many-body wave function Ψ_t satisfies the Schrödinger equation

$$i\partial_t \Psi_t = H_{\omega t} \Psi_t. \tag{S48}$$

It is easy to see that Ψ_t factors as follows:

$$\Psi_t = \left(e^{-iH_e t} | \text{electrons} \right) \otimes \psi_t, \tag{S49}$$

where ψ_t satisfies the Schrödinger equation with the effective spin Hamiltonian $H^{Se}_{\omega t}$,

$$i\partial_t \psi_t = H^{Se}_{\omega t} \psi_t, \qquad H^{Se}_{\omega t} = \gamma \, e^{-i\omega t S_z} S_y \, e^{i\omega t S_z}.$$
(S50)

The figure of merit of the pure state adiabaticity is the adiabatic fidelity between the dynamical wave function Ψ_t and the instantaneous eigenfunction Φ_{α} of the Hamiltonian (S44):

$$F_t \equiv |\langle \Phi_{\alpha=\omega t} | \Psi_t \rangle|^2.$$
(S51)

From (S49) one immediately obtains that \mathcal{F}_t is given by

$$\mathcal{F}_t = |\langle \varphi_{\alpha = \omega t} | \psi_t \rangle|^2, \tag{S52}$$

where φ_{α} is the eigenstate of H_{α}^{Se} satisfying $\varphi_0 = \psi_0$. The dynamics of ψ_t can be easily inferred from eq. (S50) by transformation to the rotating frame. As a result one obtains

$$1 - \mathcal{F}_t = \frac{\omega^2}{\omega^2 + \gamma^2} \sin^2 \left(\frac{\alpha}{2} \sqrt{1 + \gamma^2 / \omega^2} \right) \Big|_{\alpha = \omega t}.$$
(S53)

We say that the adiabaticity is maintained up to some target α with the accuracy ε if for $t \leq \alpha/\omega$

$$1 - \mathcal{F}_t \le \varepsilon. \tag{S54}$$

Let us assume that the target α is greater than π . Then the sine squared in eq. (S53) will become equal to unity somewhere on the way to the target α . Taking this into account, we conclude from eq. (S53) that the maximal driving rate ω_{ε} that allows to maintain adiabaticity with the accuracy ε is given by

$$\omega_{\varepsilon} = \gamma \sqrt{\frac{\varepsilon}{1 - \varepsilon}}.$$
(S55)

Since $P_e \sim p_F \sqrt{N}$ in the majority of states in the Gibbs ensemble, it follows from eqs. (S45) and (S55) that for these states

$$\omega_{\varepsilon} \sim 1/\sqrt{N}$$
 (S56)

in the thermodynamic limit $N \to \infty$, $\rho = \text{const.}$

It should be stressed that the explicit solution of the dynamical problem presented here works only in the case when the total momentum of electrons in the wire is conserved, i.e. for periodic boundary conditions imposed on electron wave functions. If this is not the case, e.g. for a long piece of wire with open ends, the dynamics of electrons and the spin are coupled. However, we see no reasons to expect that different boundary conditions can make the scaling of the driving rate with the system size more favorable for the pure state adiabaticity.