

Five open problems in quantum information

Paweł Horodecki,^{1,2,3} Łukasz Rudnicki,^{1,2,4,*} and Karol Życzkowski^{1,5,4}

¹*National Quantum Information Centre (KCIK), University of Gdańsk, 81-824 Sopot, Poland*

²*International Centre for Theory of Quantum Technologies (ICTQT), University of Gdańsk, 80-308 Gdańsk, Poland*

³*Faculty of Applied Physics and Mathematics, Technical University of Gdańsk, 80-952 Gdańsk, Poland*

⁴*Center for Theoretical Physics, Polish Academy of Sciences, Aleja Lotników 32/46, 02-668 Warsaw, Poland*

⁵*Faculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University, 30-348 Kraków, Poland*

(Dated: February 8, 2020)

Five selected problems in the theory of quantum information are presented. The first four concern existence of certain objects relevant for quantum information, namely mutually unbiased bases in dimension six, an infinite family of symmetric informationally complete generalized measurements, absolutely maximally entangled states for four subsystems with six levels each and bound entangled states with negative partial transpose. The last problem requires checking whether a certain state of a two-ququart system is 2-copy distillable. Finding a correct answer to any of them will be rewarded by the Golden KCIK Award established by the National Quantum Information Centre (KCIK) in Poland. A detailed description of the problems in question, the motivation to analyze them, as well as the rules for the open competition are provided.

I. INTRODUCTION

KCIK was established in 2007 as a joint research unit gathering the researchers from the field of quantum information working in Poland. The mission of the Centre, since its foundation, has been to create an integrated basis for interdisciplinary research within the fields of quantum information processing and foundations of quantum physics [1]. As a new initiative, the Centre announces an open competition (further called the **Golden KCIK Award**) aiming at encouraging the quantum information community worldwide to work on a few not easy, but at

the same time well-motivated research problems extensively covered by the topical literature. The problems in question pertain to the research which was always in the scope of the scientists working within KCIK.

In the next section the general rules applying to the award are explained. Then, in Sec. III we provide a comprehensive description of the three problems (KCIK Problem 1.–3.) associated with symmetric structures in discrete Hilbert spaces, while Sec. IV covers the description of two problems (No. 4 and 5) relevant for entanglement distillability.

II. GENERAL RULES

Every scientist worldwide is eligible to participate in the competition. The Golden KCIK Award will be conferred for solving one of five KCIK problems on quantum information listed below. To participate, an author should send before January 31 of each year (next deadline in 2021), to the email address: kcikaward@ug.edu.pl:

- a) a single page summarizing the solution (in pdf format),
- b) a link to an arXiv preprint posted within the year previous to the deadline date, in which a solution of the problem is provided.

The Golden KCIK Award conferred for the year 2020 is set to 2020 EUR. If a given problem is not solved during the year $XXXX$, the competition will automatically be extended to the next Year $XXXX + 1$ with the same rules. The KCIK Award will be upgraded *linearly*¹ to $XXXX + 1$ EUR. More importantly, each recipient of the **KCIK Award** will be invited to present her/his work during the forthcoming *Symposium on Quantum Information in Gdańsk* (Poland). The local costs will be covered by KCIK.

Each year up to two prizes can be awarded. The Golden Award can be conferred to a group of authors who will share the prize. In such a case the awarded group will select a single representative to present the work during the Symposium.

The Competition will be closed if all five problems are solved. Then, as always happens, new problems will come into play...

¹ A reader educated in financial mathematics will know that during the last decade the Euro experienced an average inflation rate of 1.27% per year. Due to a positive risk-less interest rate the purchasing power of a single euro decreases exponentially in time, so the strategy to wait with a ready solution of a problem to the year $XXXX + k$ is suboptimal.

Detailed description of the problems

III. DISCRETE STRUCTURES IN THE HILBERT SPACE

The space of pure quantum states of a fixed dimension² N is isotropic – no quantum state is "more equal" than others. However, it is legitimate to ask, for which dimensions certain particular *constellations* of quantum states with prescribed properties do exist. Although it is widely believed that symmetric informationally complete generalized quantum measurements do exist for any N , they do so in different ways – see Problem 1 and references therein.

Furthermore, if one looks for other quantum states' configurations, the answer depends on the prime decomposition of the dimension. Two of the problems formulated below concern the dimension $N = 6$, the smallest *perfect number*, equal to the sum of its positive divisors. More importantly, six is the smallest natural number which is neither a prime nor a power of a prime. And from the quantum perspective, this is the smallest dimension, for which the problems concerning the number of mutually unbiased bases and the number of quantum orthogonal Latin squares remain open.

The list of open problems formulated for the KCIK Award and presented here consists of five items only. Why five? In view of problems 2 and 3, specified below, it looks like our understanding of the number six is still not satisfactory...

KCIK Problem 1. Construct an infinite sequence of SIC POVM in different dimensions, N_1, N_2, N_3, \dots

Setup. A symmetric informationally complete positive operator valued measure (SIC POVM) [2, 3] associated with N -dimensional Hilbert space is given by a set of N^2 vectors $|\psi_j\rangle \in \mathcal{H}_N$ satisfying relation $|\langle\psi_j|\psi_k\rangle|^2 = \frac{N\delta_{ij}+1}{N+1}$. This set defines a generalized quantum measurement capable to extract complete information concerning the measured density matrix, described by $N^2 - 1$ real parameters. Such a constellation of N^2 pure states forms a simplex inscribed in the set of one dimensional projectors, the convex hull of which forms the entire set of $N^2 - 1$ dimensional density matrices. For an accessible guide to the SIC problem in low dimensions consult [4].

Motivation. From a mathematical point of view, we ask about the maximal set of *complex equiangular lines* in a given dimension N . From a physical perspective one looks for an optimal scheme of a quantum measurement

of an arbitrary size N . Solving the SIC existence problem for any dimension will significantly contribute to our understanding of the set-of-quantum-states's geometry [5].

According to the 1999 dated conjecture by Zauner [2], for any dimension N there exists a *fiducial vector*, such that all remaining $N^2 - 1$ elements of the desired SIC can be obtained by acting on it with unitary matrices representing elements of the Weyl–Heisenberg group. For $N \leq 28$ and several other dimensions (e.g. $N = 30, 31, 35, 37, 39, 48, 53, 124, 195, 323, 1299$) the solutions are known analytically [6–10]. Numerical solutions are known for all $N \leq 189$ and for several much larger dimensions [6–8, 11], including $N = 844, 2208$. However, in spite of the recent research effort [12–15], the general conjecture of Zauner remains unproven. Finding an infinite family of SICs, as requested in KCIK Problem 1, could become decisive step in this direction. Furthermore, let us emphasize inspiring connections to some major open questions in algebraic number theory, including the 12th problem of Hilbert [10, 12, 16, 17].

KCIK Problem 2. Construct a set of at least 4 mutually unbiased bases (MUBs) of order six or prove that there are no 7 MUBs in \mathcal{H}_6 .

Setup. Consider a set of K bases $\{|\psi_i^{(m)}\rangle\}$ ($1 \leq m \leq K$, $1 \leq i \leq N$) in N -dimensional complex Hilbert space \mathcal{H}_N , so that all vectors in each basis are orthogonal, (i) $\langle\psi_i^{(m)}|\psi_j^{(m)}\rangle = \delta_{ij}$. These bases are called *mutually unbiased* (MUB), if any two bases are unbiased, (ii) $|\langle\psi_i^{(m)}|\psi_j^{(n)}\rangle|^2 = 1/N$ for $m \neq n$.

It is easy to show that there exist no more than $(N + 1)$ MUBs in \mathcal{H}_N . For any $N \geq 2$, there exist at least three MUBs. If the dimension N is a prime number or a power of a prime, $N = p^k$, there exists a complete set of $(N + 1)$ MUBs [18, 19]. Several methods to construct MUBs are known [20–22] and all solutions for dimensions 2 – 5 are classified [23]. If N is a power of prime, various properties of a complete set of $(N + 1)$ MUBs are already understood [24–29], but otherwise the number of existing MUBs remains unknown [30, 31]. In particular, for $N = 6$ a complete set would consist of seven MUBs, but up till now only solutions containing three bases were found [32–40]. If a complete set of seven MUB exists, it cannot contain a triple of product bases [41, 42].

A matrix U which relates two unbiased bases of order N , belongs to the set of complex Hadamard matrices, which includes unitary matrices, such that all its entries have the same squared modulus $|U_{ij}|^2 = 1/N$. Interestingly, the set of complex Hadamard matrices is fully characterized [43, 44] for $N \leq 5$, while for higher dimensions further connections between Hadamard matrices and MUBs were found [45].

Motivation. On the one hand, finding the complete set of MUBs in dimension 6 would yield an optimal scheme

² While presenting the problems we used notation common in each subfield. Thus the reader is advised that the notation is not entirely consistent throughout the entire document. Consequently, in different problems the symbols n, d, k and N have different meaning.

of orthogonal quantum measurement in this dimension. More importantly, deciding whether such a configuration exists has significant implications for foundations of quantum theory, as up till now our understanding of basic properties of finite dimensional Hilbert spaces is not complete. On the other hand, a possible non-existence result is of a considerable mathematical interest, as it would show that the number 6 is indeed very special and ‘less equal than others’. Research on the MUB problem reveals further intricate links between foundations of quantum theory and several fields of mathematics, including Galois rings, group theory, combinatorics, finite fields and projective geometry [46–53].

KCIK Problem 3. Determine whether there exist two quantum orthogonal Latin squares [54, 55] of order six. In other words, find a solution of the problem of 36 entangled officers of Euler or demonstrate that it does not exist.

Setup. A Latin square (LS) of size N consists of N sets of N symbols arranged in such a way that no row or column of the square contains the same number twice. The name refers to papers of Leonhard Euler [56], who used Latin characters as symbols to be arranged.

Two orthogonal Latin squares (also called Graeco-Latin squares) of size N consist of N^2 cells arranged in a square with a pair of ordered symbols in each cell, for instance one Greek character and one Latin. Every row and every column of the square contains each element of the pair exactly once, and no two cells contain the same ordered pair. It is easy to show that for a given dimension N there exist no more than $(N - 1)$ mutually orthogonal Latin squares (MOLS). This bound is saturated if N is a prime or a power of prime [57].

Euler analyzed the problem of 36 officers from six regiments, each containing 6 officers of 6 different ranks. They should be arranged before a parade into a square 6×6 such that each row and each column holds only one officer from each regiment and only one officer from each rank. Euler wrote in 1782 that this problem has no solution [56] without providing a formal proof, established only in 1901 by Gaston Tarry [58]. This result implies that there is no pair of orthogonal Latin squares of size 6, so that the upper bound for the number of MOLS, in this case $N - 1 = 5$, is not saturated. For any $N \geq 7$ there exist at least two MOLS, in particular also for $N = 2 \times 5 = 10$ – consult a novel by Georges Perec [59]. In general, the problem of finding the number of MOLS for an arbitrary value of N remains open [60].

As a rule of thumb, for any interesting classical notion one can find a quantum analogue. A quantum Latin square is an $N \times N$ table of N^2 vectors from N dimensional Hilbert space \mathcal{H}_N arranged in such a way that every row and every column of the table forms an orthonormal basis in the space [61]. Two quantum orthogonal Latin squares (QOLS) are defined [54] as collection of N^2 normalized vectors from a composite space $\mathcal{H}_N \otimes \mathcal{H}_N$, which are mutually orthogonal so they form an orthonor-

mal basis. They are arranged in an $N \times N$ table such that for every row (column) the sum of all states in each row (column) is proportional to the maximally entangled state, $|\psi_+\rangle = \frac{1}{\sqrt{N}} \sum_{j=1}^N |j\rangle \otimes |j\rangle$. Any Graeco-Latin square leads to such a design, since it suffices to treat the pair of classical objects (α, B) as a product state, $|\alpha\rangle \otimes |B\rangle$.

An n -partite pure state is called *absolutely maximally entangled* (AME) state if it is maximally entangled with respect all possible bipartitions [62], so that all its reductions consisting of k subsystems, with arbitrary $k \leq n/2$, are maximally mixed. A density matrix ρ on a given M -dimensional Hilbert space is *maximally mixed*, if it is proportional to the identity operator on this space, $\rho = \mathbb{1}_M/M$. It is known that there are no AME states of 4 qubits [63], and equivalently, a pair of QOLS does not exist for $N = 2$.

As there are no two orthogonal Latin squares (OLS) of size six, the famous classical problem of 36 officers of Euler has no solution [64]. An analogous quantum problem, which involves 36 entangled officers, remains open.

Motivation. This problem can be reformulated in several other settings. Establishing a negative result is equivalent to proving that a) there is no AME state of four subsystems with six levels each [62, 65, 66], thus the corresponding quantum error correction code [67], written $((4, 1, 3))_6$, does not exist; b) there is no 2-unitary matrix $U \in U(36)$ – see [68] – which saturates the absolute bound for *entangling power* [69]; c) there are no *perfect tensors* [70] with four indices, each running from 1 to 6. Furthermore, a negative result would directly imply the famous Euler conjecture that there are no two orthogonal Latin squares of size 6. On the other hand, rather unlikely, but still possible positive result, could become an important step towards development of *quantum combinatorics*: a search for particular constellations of discrete quantum objects, with special properties of symmetry and balance, hidden in the continuous Hilbert space.

As problems 2 and 3 refer to the same dimension, $N = 6$, it is natural to speculate that they might be somehow related. It seems, however, that a connection between problems of finding the maximal number of MOLSs and MUBs for a given dimension is not a direct one [51, 52].

IV. QUANTUM ENTANGLEMENT AND ITS DISTILLABILITY

Any bipartite product state is called separable, while all other pure states are entangled. A density matrix (mixed state) is called *entangled* if it cannot be represented as a convex combination of product states [71].

Entanglement forms a crucial resource used in the theory of quantum information processing. Therefore, one of the major problems in this field is to decide, whether

a given quantum state of a composite system is separable or entangled [72]. Up till now, such a problem is solved only for 2×2 and 2×3 systems, as in these cases the single positive partial transpose (PPT) criterion provides a constructive answer [73]. Already for the 3×3 system neither a finiter number of positive maps based separability criteria [74] nor a techique based on finite size semidefinite programming [75] can always allow us to settle, whether a given quantum state is entangled or not. The known procedure deciding separability of a given bi-partite quantum state in a finite number of steps [76] can not be applied in practice due to its high complexity.

KCIK Problem 4. Establish whether there exist bound entangled states with negative partial transpose (NPT).

Setup. A state is called *bound entangled* if it is entangled but not distillable [77–79]. A bipartite state ρ defined on a bipartite Hilbert space $\mathcal{H}_d \otimes \mathcal{H}_d$ is called *distillable*, if it is *n-copy distillable* for some finite n . The property of n -copy distillability means that there exist two-dimensional (i.e. of rank two) projectors P and Q such that the matrix $(P \otimes Q)[\rho^\Gamma]^{\otimes n}(P \otimes Q)$ has a negative eigenvalue [78, 80]. Here ρ^Γ stands for the *partial transpose* of the state ρ defined by its matrix elements as $\langle ij|\rho^\Gamma|kl\rangle = \langle il|\rho|kj\rangle$. One says that the state has *negative partial transpose* (NPT) iff its partial transpose ρ^Γ has some of its eigenvalues negative. It should be stressed that the projectors P and Q act on the product $(\mathcal{C}^d)^{\otimes n}$ of all n Hilbert spaces associated with left and right subsystems of copies of the considered bipartite system, respectively — see [81, 82].

The question of NPT bound entanglement is closely related to a mathematical problem concerning 2-co-positive maps [81],[83]. A linear map $\Lambda : M_d(\mathcal{C}) \rightarrow M_d(\mathcal{C})$ acting on \mathcal{H}_d is called *positive* iff it transforms any matrix with non-negative eigenvalues into a matrix with the same property. Furthermore, a linear map Λ is called *k-positive* if and only if the following extension $\mathbb{1}_k \otimes \Lambda : M_k(\mathcal{C}) \otimes M_d(\mathcal{C}) \rightarrow M_k(\mathcal{C}) \otimes M_d(\mathcal{C})$ is positive, where $\mathbb{1}_k$ stands for the identity map (i.e. the one that maps any complex matrix from $M_k(\mathcal{C})$ to itself). The map is called *completely positive* iff it is k-positive for any k. For a finite dimension d , to ensure complete positivity it is enough to check only k-positivity for $k = d$. A map Λ is called *k-co-positive* if and only if the composition $S = T \circ \Lambda$ is k-positive, where T stands for transposition.

Motivation. This is one of the long-standing open questions of quantum information theory [81, 84]. Its positive solution would have several consequences. If NPT bound entangled states exist then the set of non-distillable entangled states is neither closed under the tensor product nor under mixing – see [83]. The latter means that there would exist two non-distillable entangled states such that their mixture were distillable. This would imply that one of the central measures of entanglement theory, namely distillable entanglement (which

describes asymptotic amount of entanglement that can be distilled from many copies of a given state by local operations and classical communication [72]) is neither additive nor convex [83].

Therefore positive solution of the present problem would lead to an extremal example of *superadditivity*. Namely it has been proven that for any NPT state there exists PPT bound entangled state such that the product of the two is distillable [85]. If the NPT state were bound entangled, then we would have the pair of two bound entangled state (with distillable entanglement measure zero) such that their tensor product would be distillable (i.e. having the measure strictly positive). As already pointed out, such a scenario is an extremal case of superadditivity: two objects containing no resource of a given type, if put together constitute a single object that, surprisingly, turns out to contain some amount of the resource (for this type of effect on the ground of quantum channel capacities see [86]).

One can show [81] that the existence of n-copy non-distillable state is equivalent to the existence of a completely positive map Λ such that it is 2-co-positive and its n-th tensor power $\Lambda^{\otimes n} = \Lambda \otimes \dots \otimes \Lambda$ is also co-positive. Consequently, existence of NPT bound entanglement is equivalent to existence of a completely positive map that is 2-co-positive and its arbitrary high tensor power is also co-positive.

Note that for any n there exists an n -copy nondistillable state that is $(n + 1)$ -copy distillable (see [87]) which may be considered as an indication that the present problem of existence of NPT bound entanglement is hard.

KCIK Problem 5. Show that the two-ququart **Werner state** $\rho(4, -1/2)$ is not 2-copy distillable. This state, defined below, is the only two-ququart Werner state, for which its partial transpose ρ^Γ is proportional to a unitary matrix.

Setup. Consider the family of Werner states, defined on the Hilbert space $\mathcal{C}^d \otimes \mathcal{C}^d$ as $\rho(d, \alpha) = \frac{\mathbb{1} \otimes \mathbb{1} + \alpha V}{d^2 + \alpha d}$ with the general range of the parameter $\alpha \in [-1, 1]$. The matrix V stands for the *Swap operator*, defined by its matrix elements, $\langle ij|V|kl\rangle = \delta_{il}\delta_{jk}$. The Werner states are invariant with respect to twirling with local unitaries [71], so they are also called $U \otimes U$ -invariant. It is conjectured [81, 83, 88] that Werner states which are not 1-copy distillable are also 2-copy nondistillable.

Motivation. It would be a first step for a possible proof of existence of NPT bound entanglement. Equivalently, it would provide a very elegant completely positive map that is 2-co-positive, such that a tensor product of its two copies also has that property. Interestingly, this question is equivalent to the following mathematical problem [82]: show that sum of squares of the two largest singular values is bounded by $\frac{1}{2}$ for any matrix X of the form $X = A \otimes \mathbb{1} + \mathbb{1} \otimes B$, where A and B denote traceless matrices of size 4 satisfying $\text{Tr}(A^\dagger A) + \text{Tr}(B^\dagger B) = \frac{1}{4}$. So far, the bound equal to $\frac{1}{2}$ has been proven [82] under the

additional assumption that A and B are normal, so they commute with their hermitian conjugates.

Parameters $d = 4$ and $\alpha = -\frac{1}{2}$ in the problem are chosen since they correspond to the case of:

(i) the minimal dimension for which the very special Werner state is 1-copy nondistillable, namely the state $\rho(d, -\frac{2}{d})$ which has its partial transpose proportional to the dichotomic unitary operator $U = I - 2|\psi_+\rangle\langle\psi_+|$, where $|\psi_+\rangle = \frac{1}{\sqrt{d}} \sum_{j=1}^d |j\rangle \otimes |j\rangle$ denotes the maximally entangled state. A dichotomic unitary operator has eigenvalues ± 1 .

(ii) the unique dimension, for which the Werner state with the parameter $\alpha = -\frac{2}{d}$ is located just *on the boundary of a 1-copy nondistillability* (i.e. in case of $d = 4$ all the states with $\alpha < -\frac{1}{2}$ are already 1-copy distillable, which is not true for $d > 4$).

The choice of the state with its partial transpose proportional to the dichotomic unitary operator is motivated by the fact that checking its n -copy distillability seems to be easier than in the general case. In particular, proportionality to the dichotomic unitary operation is conserved under taking the tensor product.

Acknowledgments

We are pleased to thank Ingemar Bengtsson, Adam Burhardt, Jakub Czartowski, Dardo Goyeneche, Markus Grassl, Michał Horodecki, Felix Huber, Jędrzej Kaniewski, Arul Lakshminarayan, and Oliver Reardon-Smith for helpful comments and remarks.

-
- [1] KCIK webpage: <https://kcik.ug.edu.pl/kcik.php>
- [2] G. Zauner, Quantendesigns – Grundzüge einer nichtkommutativen Designtheorie. Dissertation, Universität Wien, 1999; English translation, *Int. J. Quantum Inf.* **9**, 445 (2004).
- [3] J. M. Renes, R. Blume-Kohout, A. J. Scott, and C. M. Caves, Symmetric Informationally Complete Quantum Measurements, *J. Math. Phys.* **45**, 2171 (2004).
- [4] D. Andersson, An Enthusiast’s Guide to SICs in Low Dimensions, Master Thesis, Stockholm University, 2015
- [5] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States. An Introduction to Quantum Entanglement*, II edition, Cambridge University Press, Cambridge 2017.
- [6] A. J. Scott and M. Grassl, SIC-POVMs: A new computer study, *J. Math. Phys.* **51**, 042203 (2010).
- [7] A. J. Scott, SICs: Extending the list of solutions, preprint arXiv:1703.03993
- [8] M. Grassl and A. J. Scott, Fibonacci-Lucas SIC-POVMs, *J. Math. Phys.* **58**, 122201 (2017).
- [9] M. Appleby, I. Bengtsson, I. Dumitru, and S. Flammia, Dimension towers of SICs. I. Aligned SICs and embedded tight frames, *J. Math. Phys.* **58**, 112201 (2017).
- [10] M. Appleby, T.-Y. Chien, S. Flammia, and S. Waldron, Constructing exact symmetric informationally complete measurements from numerical solutions, *J. Phys. A* **51**, 165302 (2018).
- [11] C. A. Fuchs, M.C. Hoang, and B.C. Stacey, The SIC Question: History and State of Play, *Axioms* **6**, 21 (2017).
- [12] G. S. Kopp, SIC-POVMs and the Stark conjectures, preprint arXiv:1807.05877
- [13] M. Appleby and I. Bengtsson, Simplified exact SICs, *J. Math. Phys.* **60**, 062203 (2019).
- [14] O. Andersson and I. Dumitru, Aligned SICs and embedded tight frames in even dimensions, preprint arXiv:1905.09737
- [15] I. Bengtsson, Algebraic units, anti-unitary symmetries, and a small catalogue of SICs, preprint arXiv:2001.08487
- [16] D. M. Appleby, H. Yadsan-Appleby, and G. Zauner, Galois automorphisms of a symmetric measurement, *Quantum Inf. Comput.* **13**, 672 (2013).
- [17] M. Appleby, S. Flammia, G. McConnell, and J. Yard, SICs and Algebraic Number Theory, *Found. Phys.* **47**, 1042 (2017).
- [18] I. D. Ivanović, Geometrical description of quantal state determination, *J. Phys. A* **14**, 3241 (1981).
- [19] W. K. Wootters and B. D. Fields, Optimal state-determination by mutually unbiased measurements, *Ann. Phys.* **191**, 363 (1989).
- [20] S. Bandyopadhyay, P. O. Boykin, V. Roychowdhury, and F. Vatan, A new proof for the existence of MUBs, *Algorithmica* **34**, 512 (2002).
- [21] A. Klappenecker and M. Rötteler, Construction of MUBs, *Lect. Notes Comp. Sci.* **2948**, 137 (2004).
- [22] C. Archer, There is no generalization of known formulas for MUBs, *J. Math. Phys.* **46**, 022106 (2005).
- [23] S. Brierley, S. Weigert, and I. Bengtsson, All mutually unbiased bases in dimensions two to five, *Quantum Info. Comp.* **10**, 0803 (2010).
- [24] J. Lawrence, Mutually unbiased bases and trinary operator sets for N qutrits, *Phys. Rev. A* **70**, 012302 (2004).
- [25] J. L. Romero, G. Björk, A. B. Klimov, and L. L. Sánchez-Soto, Structure of the sets of mutually unbiased bases for N qubits, *Phys. Rev. A* **72**, 062310 (2005).
- [26] P. Wocjan and T. Beth, New construction of MUBs in square dimensions, *Quant. Inf. Comput.* **5**, 181 (2005).
- [27] M. Combescure, Block-circulant matrices with circulant blocks, Weil sums, and mutually unbiased bases. II. The prime power case *J. Math. Phys.* **50**, 032104 (2009).
- [28] M. Wieśniak, T. Paterek, and A. Zeilinger, Entanglement in mutually unbiased bases, *New J. Phys.* **13**, 053047 (2011).
- [29] D. McNulty, B. Pammer, and S. Weigert, Mutually Unbiased Product Bases for Multiple Qudits, *J. Math. Phys.* **57**, 032202 (2016).
- [30] T. Durt, B.-G. Englert, I. Bengtsson, and K. Życzkowski, On mutually unbiased bases, *Int. J. Quant. Inf.* **8**, 535 (2010).
- [31] G. Alber and C. Charney, Mutually unbiased bases: a group and graph theoretical approach, *Phys. Scripta* **94**, 014007 (2018).
- [32] I. Bengtsson, W. Bruzda, A. Ericsson, J.-A. Larsson, W. Tadej, and K. Życzkowski, MUBs and Hadamards of order six, *J. Math. Phys.* **48**, 052106 (2007).
- [33] P. Butterley and W. Hall, Numerical evidence for the maximum number of mutually unbiased bases in dimension six, *Phys. Lett. A* **369**, 5 (2007).
- [34] S. Brierley and S. Weigert, Constructing mutually unbiased bases in dimension six, *Phys. Rev. A* **79**, 052316 (2009).
- [35] M. Grassl, On SIC-POVMs and MUBs in Dimension 6,

- preprint arxiv/quant-ph/0406175 (v.2; 2009).
- [36] P. Jaming, M. Matolcsi, P. Móra, F. Szőllösi, and M. Weiner, A generalized Pauli problem and an infinite family of MUB-triplets in dimension 6. *J. Phys. A* **42**, 245305 (2009).
- [37] P. Jaming, M. Matolcsi, and P. Móra, The problem of mutually unbiased bases in dimension 6, *Cryptography and Communications* **2**, (2010).
- [38] P. Raynal, X. Lü, and B.-G. Englert, Mutually unbiased bases in dimension six: The four most distant bases, *Phys. Rev. A* **83**, 062303 (2011).
- [39] J. Batle, A. Farouk, M. Naseri, and M. Elhoseny, New approach to finding the maximum number of Mutually Unbiased Bases in C^6 , *Appl. Math. Inf. Sci.* **10**, 1-6 (2016),
- [40] L. Chen and L. Yu, Mutually unbiased bases in dimension six containing a product-vector basis, *Quantum Inform. Proc.* **17**, 198 (2018).
- [41] D. McNulty and S. Weigert, All mutually unbiased product bases in dimension 6, *J. Phys. A* **45**, 135307 (2012).
- [42] D. McNulty and S. Weigert, On the impossibility to extend triples of mutually unbiased product bases in dimension six, *Int. J. Quant. Inf.* **10**, 1250056 (2012).
- [43] U. Haagerup, Orthogonal maximal abelian $*$ -subalgebras of the $n \times n$ matrices and cyclic n -roots, *Operator Algebras and Quantum Field Theory (Rome)*, Cambridge, MA: International Press, pp. 296-322, 1996.
- [44] W. Tadej and K. Życzkowski, A concise guide to complex Hadamard matrices, *Open Syst. Inf. Dyn.* **13**, 133-177 (2006) and online version at <http://chaos.if.uj.edu.pl/~karol/hadamard>
- [45] D. McNulty and S. Weigert, Isolated Hadamard matrices from mutually unbiased product bases, *J. Math. Phys.* **53**, 122202 (2012).
- [46] M. Saniga, M. Planat, and H. Rosu, MUBs and finite projective planes, *J. Opt.* **B 6**, L19 (2004).
- [47] I. Bengtsson and A. Ericsson, MUBs and the complementarity polytope, *Open Syst. Inf. Dyn.* **12**, 107 (2005).
- [48] M. R. Kibler and M. Planat, A $SU(2)$ recipe for MUBs, *Int. J. Mod. Phys. B* **20**, 1802 (2006).
- [49] M. Planat, H. C. Rosu, and S. Perrine, A survey of finite algebraic geometrical structures underlying mutually unbiased quantum measurements, *Found. Phys.* **36**, 1662 (2006).
- [50] P. O. Boykin, M. Sitharam, P. H. Tiep, and P. Wojan, MUBs and orthogonal decompositions of Lie algebras, *Quant. Inform. Comput.* **7**, 371 (2007).
- [51] T. Paterek, B. Dakić, and Č. Brukner, Mutually unbiased bases, orthogonal Latin squares, and hidden-variable models *Phys. Rev. A* **79**, 012109 (2009).
- [52] T. Paterek, M. Pawłowski, M. Grassl, C. Brukner, On the connection between mutually unbiased bases and orthogonal Latin squares, *Phys. Scr.* **T140**, 014031 (2010).
- [53] M. N. Kolountzakis, M. Matolcsi and M. Weiner, An application of positive definite functions to the problem of MUBs *Proc. Amer. Math. Soc.* **146**, 1143 (2018).
- [54] D. Goyeneche, Z. Raissi, S. Di Martino, and K. Życzkowski, Entanglement and quantum combinatorial designs, *Phys. Rev. A* **97**, 062326 (2018).
- [55] B. Musto and J. Vicary, Orthogonality for Quantum Latin Isometry Squares, *EPTCS* **287**, 253 (2019).
- [56] L. Euler, Recherches sur une nouvelle espece de quarres magiques, E530, (1782).
- [57] J. Dénes and A. D. Keedwell (eds.), *Latin Squares: New Developments in the Theory and Applications*, North-Holland, Amsterdam, 1991.
- [58] G. Tarry, Le Problème des 36 Officiers, *Compte Rendu de l'Association Française pour l'Avancement des Sciences* **2**, 170 (1901).
- [59] G. Perc, *La Vie mode d'emploi*, Hachette, 1978; English translation, *Life: A User's Manual*, Godline 1987
- [60] C. J. Colbourn and J. H. Dinitz, Mutually orthogonal latin squares: a brief survey of constructions, *J. Stat. Plan. Infer.* **95**, 9 (2001).
- [61] B. Musto and J. Vicary, Quantum Latin squares and unitary error bases, *Quantum Inf. Comput.* **16**, 1318 (2016).
- [62] W. Helwig, W. Cui, J. I. Latorre, A. Riera, and H. K. Lo, Absolute maximal entanglement and quantum secret sharing, *Phys. Rev. A* **86**, 052335 (2012).
- [63] A. Higuchi and A. Sudbery, How entangled can two couples get?, *Phys. Lett. A* **273**, 213 (2000).
- [64] L. Clarisse, S. Ghosh, S. Severini, and A. Sudbery, Entangling power of permutations, *Phys. Rev. A* **72**, 012314 (2005).
- [65] F. Huber, C. Eltschka, J. Siewert, and O. Gühne, Bounds on absolutely maximally entangled states from shadow inequalities, and the quantum MacWilliams identity, *J. Phys. A* **51**, 175301 (2018).
- [66] F. Huber and N. Wyderka, Table of AME states, Online available at <http://www.tp.nt.uni-siegen.de/~fhuber/ame.html>
- [67] M. Grassl, T. Beth, and M. Rötteler, On optimal quantum codes, *Int. J. Quant. Inform.* **2**, 55 (2004).
- [68] D. Goyeneche, D. Alsina, J. I. Latorre, A. Riera, and K. Życzkowski, Absolutely Maximally Entangled states, combinatorial designs and multi-unitary matrices, *Phys. Rev. A* **92**, 032316 (2015).
- [69] P. Zanardi, Ch. Zalka, and L. Faoro, Entangling power of quantum evolutions, *Phys. Rev. A* **62**, 030301 (2000).
- [70] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence, *J. High Energy Phys.* **06**, 149 (2015).
- [71] R. F. Werner, Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model, *Phys. Rev. A* **40**, 4277 (1989).
- [72] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, *Rev. Mod. Phys.* **81**, 865 (2009).
- [73] M. Horodecki, P. Horodecki, and R. Horodecki, Separability of mixed states: necessary and sufficient conditions, *Phys. Lett. A* **223**, 1 (1996).
- [74] L. Skowronek, There is no direct generalization of positive partial transpose criterion to the three-by-three case, *J. Math. Phys.* **57**, 112201 (2016).
- [75] H. Fawzi, The set of separable states has no finite semidefinite representation except in dimension 3×2 , preprint arXiv:1905.02575
- [76] P. Badziąg, P. Horodecki, R. Horodecki, and R. Augusiak, Separability in terms of a single entanglement witness, *Phys. Rev. A* **88**, 010301 (2013).
- [77] M. Horodecki, P. Horodecki, and R. Horodecki, Mixed-State Entanglement and Distillation: Is there a 'Bound' Entanglement in Nature?, *Phys. Rev. Lett.* **80**, 5239 (1998).
- [78] M. Horodecki and P. Horodecki, Reduction criterion of separability and limits for a class of protocols of entanglement distillation, *Phys. Rev. A* **59**, 4206 (1999).
- [79] P. Horodecki, Bound entanglement, in: D. Bruss, G. Leuchs (Editors) *Form Quantum Information: From Foundations to Quantum Technology Applications*, Wiley-VCH, 2019.
- [80] L. Clarisse, The distillability problem revisited, *Quant. Inform. Comput.* **6**, 539 (2006).
- [81] D. P. DiVincenzo, P. W. Shor, J. A. Smolin, B. M. Terhal, and A. V. Thapliyal, Evidence for bound entangled states with negative partial transpose, *Phys. Rev. A* **61**, 062312 (2000).
- [82] L. Pankowski, M. Piani, M. Horodecki, P. Horodecki, A few steps more towards NPT bound entanglement, *IEEE*

- Trans. Inf. Theory* **56**, 4085 (2010).
- [83] L. Clarisse, Entanglement Distillation. A Discourse on Bound Entanglement in Quantum Information Theory, arXiv:quant-ph/0612072v1 (2006).
- [84] W. Dür, J. I. Cirac, M. Lewenstein, D. Bruss, Distillability and partial transposition in bipartite systems, *Phys. Rev. A* **61**, 062313 (2000).
- [85] T. Eggeling, K. G. H. Vollbrecht, R. F. Werner, and M. M. Wolf, Distillability via protocols respecting the positivity of partial transpose, *Phys. Rev. Lett.* **87**, 257902 (2001).
- [86] G. Smith and J. Yard, Quantum communication with zero-capacity channels, *Science* **321**, 1812 (2010).
- [87] J. Watrous, Many copies may be required for entanglement distillation, *Phys. Rev. Lett.* **93**, 010502 (2004).
- [88] D. Doković, On two-distillable Werner states, *Entropy* **18**, 216 (2016).