

# Hartmann potential with a minimal length and generalized recurrence relations for matrix elements

Lamine Khodja<sup>a,b,1</sup>, Mohamed Achour<sup>b</sup> and Slimane Zaim<sup>c</sup>

<sup>a</sup>Département de Physique, Faculté des Mathématiques et des Sciences de la Matière, Université Kasdi Merbah – Ouargla, Algeria.

<sup>b</sup>Laboratoire de Physique Théorique, Faculté des Sciences Exactes, Université de Bejaia, 06000 Bejaia, Algeria.

<sup>c</sup>Département de Physique, Faculté des Sciences de la Matière, Université de Batna-1 Hadj Lakhdar, Algeria.

## Abstract

In this work we study the Schrödinger equation in the presence of the Hartmann potential with a generalized uncertainty principle. We perturbatively obtain the matrix elements of the hamiltonian at first order in the parameter of deformation  $\beta$  and show that some degenerate states are removed. We give analytic expressions for the solutions of the diagonal matrix elements. Finally, we derive a generalized recurrence formula for the angular average values.

**KEYWORDS:** Hartmann potential; Schrödinger equation; Generalized uncertainty principle (GUP).

**PACS NUMBERS:** 02.30.Gp, 31.15.-p, 31.15.xp

---

<sup>1</sup>Email: lamine.khodja@yahoo.fr, khodja.lamine@univ-ouagla.dz

# 1 Introduction

In recent years many arguments have been suggested to motivate a modified Heisenberg algebra in quantum mechanics such as theories of quantum gravity and string theory, which lead to the existence of a minimal observable length expected to be of order of the Planck length [1–11]. The minimal length can be obtained from the deformed canonical commutation relation between position and momentum operators [12–17]:

$$[X, P] = i\hbar(1 + \beta P^2), \quad (1)$$

where  $\beta$  is a positive parameter of deformation. This commutation relation implies the following generalized uncertainty principle (GUP):

$$\Delta X \Delta P \geq \frac{\hbar}{2} (1 + \beta (\Delta P)^2), \quad (2)$$

which corresponds to a minimal length  $(\Delta X)_{\min} = \hbar\sqrt{\beta}$ .

There has recently been a lot of interest in the study of quantum mechanics problems in the presence of a minimal length with various potentials such as the harmonic oscillator, the Coulomb and Yukawa potentials, the Woods-Saxon potential, the Kratzer potential, ... [18–28]. The first author to use the standard perturbation theory to solve the Schrödinger equation of the central potentials in the presence of a minimal length was Brau in Ref. [21], where energy-level corrections were calculated and splitting of degenerate levels were found to occur. On the other hand the solutions of the Schrödinger equation with Hartmann potential  $V(r, \theta) = \eta\sigma^2 \left( \frac{e^2}{r} + \frac{q\hbar^2}{2\mu r^2 \sin^2 \theta} \right)$ , where  $\eta$  and  $\sigma$  are positive real numbers with values ranging from about 1 to 10, and  $q$  is a real parameter, is well known [29–35]. This potential has been introduced in [36–38] to describe ring-shaped molecules. When  $q = 0$  and  $\eta\sigma^2 = Z$  the Hartmann potential reduces to the Coulomb potential.

The purpose of this paper is to study the extension of the Schrödinger equation with the Hartmann potential in the presence of a minimal length. In section 2 we use the first-order perturbation theory to give the general form of the hamiltonian matrix elements and, for a particular case, we show that the degeneracy is completely removed. In section 3 we give an explicit analytical expression of the diagonal matrix elements and show that the splitting of the degenerate energy levels also occurs. In section 4 we provide a general recurrence formula for the angular part. Finally, in the last section, we draw our conclusion.

## 2 Hamiltonian matrix elements

To calculate the hamiltonian matrix elements for a Hartmann potential in the presence of a minimal length, we solve the corresponding Schrödinger equation:

$$\left[ \frac{\hat{P}^2}{2\mu} + V(\hat{r}, \hat{\theta}) \right] \psi(\vec{r}) = E^{(\beta)} \psi(\vec{r}). \quad (3)$$

We choose to work with the following representation that verifies the relation (1) to first order in  $\beta$ :

$$\hat{X}_i \psi(\vec{r}) = x_i \psi(\vec{r}), \quad (4)$$

$$\hat{P}_i \psi(\vec{r}) = p_i (1 + \beta p^2) \psi(\vec{r}), \quad p_i = i\hbar \frac{\partial}{\partial x^i}. \quad (5)$$

To first order in  $\beta$  the Schrödinger equation (3) can be written as:

$$\left[ \frac{p^2}{2\mu} + \frac{\beta p^4}{\mu} + V(r, \theta) \right] \hat{\psi}(r, \theta, \varphi) = E^{(\beta)} \hat{\psi}(r, \theta, \varphi). \quad (6)$$

In this equation the Hartmann potential in the presence of a minimal length appears within the perturbation term:

$$\frac{\beta p^4}{\mu}$$

To investigate the correlations we use the first-order perturbation theory. For  $\beta = 0$  the spectrum of equation (6) and the corresponding wave functions are well-known and are given by [29, 39, 40]:

$$\psi(r, \theta, \varphi) = \frac{1}{r} R(r) \Theta(\theta) \Phi(\phi), \quad (7)$$

where

$$\begin{aligned} \Phi(\phi) &= \frac{1}{\sqrt{2\pi}} \exp(im\phi), \quad m = 0, \pm 1, \pm 2, \pm 3, \dots \\ \Theta_{nm}(x) &= \frac{\Gamma(2k+1)}{\Gamma(k+1)} \sqrt{\frac{(2n+2k+1)}{2^{2k+1}\Gamma(n+2k+1)}} (1-x^2)^{k/2} C_n^{(k+1/2)}(x), \quad x = \cos\theta \\ R_{Nnm}(r) &= \left( \frac{\mu\eta\sigma^2 e^2}{\hbar^2 n'} \right)^{1/2} \left[ \frac{(n'-l-1)!}{n'\Gamma(n'+l+1)} \right]^{1/2} \left( \frac{2\mu\eta\sigma^2 e^2}{\hbar^2 n'} r \right)^{l+1} \\ &\times \exp \left( -\frac{\mu\eta\sigma^2 e^2}{\hbar^2 n'} r \right) L_N^{(2l+1)} \left( \frac{2\mu\eta\sigma^2 e^2}{\hbar^2 n'} r \right), \\ k &= \sqrt{m^2 + \frac{q\eta\hbar^2\sigma^2}{2\mu}}, \quad n' = N + l + 1, \quad l = n + k, \quad N, n = 0, 1, 2, 3, \dots \quad (8) \end{aligned}$$

with  $N$  being the radial quantum number,  $L_n^{(\nu)}(x)$  and  $C_n^{(\nu)}(x)$  respectively stand for the associated Laguerre and the Gegenbauer (ultraspherical) polynomials. The orthogonality conditions for these functions are:

$$\begin{aligned} \int_{-1}^1 dx (1-x^2)^{\nu-1/2} [C_n^{(\nu)}(x)]^2 &= \frac{\pi 2^{1-2\nu} \Gamma(n+2\nu)}{n! (n+\nu) [\Gamma(\nu)]^2}, \\ \int_{-1}^1 dx e^{-x} x^\nu [L_n^{(\nu)}(x)]^2 &= \frac{\Gamma(\nu+n+1)}{n!}. \end{aligned} \quad (9)$$

The energy eigenvalues are:

$$E_{Nnm}^{(0)} = -\frac{\mu(\eta\sigma^2)^2 e^4}{2\hbar^2} \left[ N + n + \sqrt{m^2 + \frac{q\eta\sigma^2\hbar^2}{2\mu}} + 1 \right]^{-2}. \quad (10)$$

The first-order perturbation theory gives the matrix element of the hamiltonian operator (6) up to first order in  $\beta$  as follows [21]:

$$\begin{aligned} \frac{\beta}{\mu} \langle N_1 n_1 m_1 | p^4 | N_2 n_2 m_2 \rangle &= 4\mu\beta \left[ \left( E_{N_1 n_1 m_1}^{(0)} \right)^2 \delta_{N_1 N_2} \delta_{n_1 n_2} \delta_{m_1 m_2} \right. \\ &\quad \left. - 2E_{N_2 n_2 m_2}^{(0)} \langle N_1 n_1 m_1 | V(r, \theta) | N_2 n_2 m_2 \rangle + \langle N_1 n_1 m_1 | (V(r, \theta))^2 | N_2 n_2 m_2 \rangle \right], \end{aligned} \quad (11)$$

where

$$V(r, \theta) = \eta\sigma^2 \left( \frac{e^2}{r} + \frac{q\hbar^2}{2\mu r^2 \sin^2 \theta} \right). \quad (12)$$

Each of these terms can be written as

$$\begin{aligned} \langle N_1 n_1 m_1 | \frac{1}{r^s \sin^{2t} \theta} | N_2 n_2 m_2 \rangle &= \int_0^\infty R_{N_1 n_1 m_1}(r) R_{N_2 n_2 m_2}(r) r^{-s} dr \\ &\quad \times \int_{-1}^1 dx (1-x^2)^{-t} \Theta_{n_1 m_1}(x) \Theta_{n_2 m_2}(x) \end{aligned} \quad (13)$$

For the radial part, the first integral has been evaluated in [30] and its expression is given by:

$$\begin{aligned} \langle N_1 n_1 m_1 | r^s | N_2 n_2 m_2 \rangle &= \int_0^\infty R_{N_1 n_1 m_1}(r) R_{N_2 n_2 m_2}(r) r^s dr \\ &= \frac{\eta\sigma^2}{n_1 n_2} \sqrt{\frac{N_1! N_2!}{\Gamma(2l_1 + N + 2) \Gamma(2l_2 + N + 2)}} \left( \frac{2\eta\sigma^2}{n'_1} \right)^{l_1+1} \left( \frac{2\eta\sigma^2}{n'_2} \right)^{l_2+1} \\ &\quad \times [\eta\sigma^2 (1/n'_1 + 1/n'_2)]^{-s-l_1-l_2-3} \sum_{m_1}^N \sum_{m_2}^N \frac{(-1)^{2N+m_2}}{m_1! m_2!} \left( \frac{1/n'_1 - 1/n'_2}{1/n'_1 + 1/n'_2} \right)^{m_1+m_2} \\ &\quad \times \Gamma(l_1 + l_2 + s + m_1 + m_2 + 3) \sum_{m_3=0}^t \binom{l_1 + l_2 + s + m_1 + m_2 + 1}{N - m_1 - m_2 - m_3} \\ &\quad \times \binom{l_1 + l_2 + s + m_1 + 1}{N - m_2 - m_3} \binom{l_1 + l_2 + s + m_1 + m_2 + m_3 + 2}{m_3}, \end{aligned} \quad (14)$$

with  $t = \min(N - m_1, N - m_2)$  and  $s < l_1 + l_2 + 3$ . The only values of  $s$  which contribute in the calculation of the matrix elements given in the expression (11) are  $s = \{-1, -2, -3, -4\}$ .

For the angular part we use the following integral of the product of two Gegenbauer polynomials [41] with  $\theta > -1/2$ ,  $\mu > -1/2$ , and  $\lambda > -1/2$ :

$$\begin{aligned} \int_{-1}^1 C_l^\theta(x) C_m^\mu(x) (1-x^2)^{\lambda-1/2} dx &= \frac{\pi 2^{1-2\lambda}}{\Gamma(\mu)\Gamma(\theta)} \sum_{k=0}^{[l/2]} \left[ \frac{(l-2k+\lambda)}{(l-2k)!k!s!} \right. \\ &\times \left. \frac{\Gamma(l-\theta-k)}{\Gamma(l+\lambda-k+1)} \frac{\Gamma(l-2k+2\lambda)\Gamma(m+\mu-s)}{\Gamma(m+\lambda-s+1)} (\theta-\lambda)_k (\mu-\lambda)_s \right], \quad (15) \end{aligned}$$

where  $m = l - 2k + 2s$ , with  $[m/2] \geq s \in N$ , and  $(z)_n$  is the Pochhammer symbol [43]:

$$(z)_n = z(z+1)\dots(z+n-1) = \frac{\Gamma(z+n)}{\Gamma(z)}. \quad (16)$$

The integral (15) vanishes for odd values of  $l+m$ . We thus have the following special integrals:

$$\begin{aligned} \int_{-1}^1 dx (1-x^2)^{-1} \Theta_{n_1 m_1}(x) \Theta_{n_2 m_2}(x) &= \left| \cos \frac{\pi}{2} (n_1 + n_2) \right| \\ &\times \frac{\Gamma(2k_1+1)}{\Gamma(k_1+1)} \frac{\Gamma(2k_1+1)}{\Gamma(k_1+1)} \left[ \frac{(2n_1+2k_1+1)}{2^{2(k_1+k_2+1)} \Gamma(n_1+2k_1+1)} \frac{(2n_2+2k_2+1)}{\Gamma(n_2+2k_2+1)} \right]^{1/2} \\ &\times \frac{\pi 2^{2-k_1-k_2}}{\Gamma(k_1+1/2) \Gamma(k_2+1/2)} \sum_{p=0}^{[n_1/2]} \left[ \frac{[n_1-2p+(k_1+k_2-1)/2]}{(n_1-2p)!p!s!} \right. \\ &\times \frac{\Gamma(n_1-2p+k_1+k_2-1)\Gamma(n_2+k_1-s+1/2)}{\Gamma(n_2+\frac{k_1+k_2+1}{2}-s)} \\ &\times \left. \frac{\Gamma(n_1-k_1-p-1/2)}{\Gamma(n_1+\frac{k_1+k_2+1}{2}-p)} \left( \frac{k_1-k_2}{2} + 1 \right)_p \left( \frac{k_2-k_1}{2} + 1 \right)_s \right]. \quad (17) \end{aligned}$$

and

$$\begin{aligned} \int_{-1}^1 dx (1-x^2)^{-2} \Theta_{n_1 m_1}(x) \Theta_{n_2 m_2}(x) &= \left| \cos \frac{\pi}{2} (n_1 + n_2) \right| \times \\ &\times \frac{\Gamma(2k_1+1)}{\Gamma(k_1+1)} \frac{\Gamma(2k_1+1)}{\Gamma(k_1+1)} \left[ \frac{(2n_1+2k_1+1)}{2^{2(k_1+k_2+1)} \Gamma(n_1+2k_1+1)} \frac{(2n_2+2k_2+1)}{\Gamma(n_2+2k_2+1)} \right]^{1/2} \\ &\times \frac{\pi 2^{4-k_1-k_2}}{\Gamma(k_1+1/2) \Gamma(k_2+1/2)} \sum_{p=0}^{[n_1/2]} \left[ \frac{[n_1-2p+(k_1+k_2-3)/2]}{(n_1-2p)!p!s!} \right. \\ &\times \frac{\Gamma(n_1-2p+k_1+k_2-3)\Gamma(n_2+k_1-s+1/2)}{\Gamma(n_2+\frac{k_1+k_2-1}{2}-s)} \\ &\times \left. \frac{\Gamma(n_1-k_1-p-1/2)}{\Gamma(n_1+\frac{k_1+k_2-1}{2}-p)} \left( \frac{k_1-k_2}{2} + 2 \right)_p \left( \frac{k_2-k_1}{2} + 2 \right)_s \right]. \quad (18) \end{aligned}$$

Finally, the general form of the hamiltonian matrix elements in (11) is given by the expression:

$$\begin{aligned}
\frac{\beta}{4\mu} \langle N_1 n_1 m_1 | p^4 | N_2 n_2 m_2 \rangle &= \frac{\mu^2 (\eta\sigma^2)^4 e^8}{4\hbar^4} (n'_2)^{-4} \delta_{N_1 N_2} \delta_{n_1 n_2} \delta_{m_1 m_2} \\
&+ \frac{\mu (\eta\sigma^2)^4 e^6}{\hbar^2} (n'_2)^{-2} \langle N_1 n_1 m_1 | r^{-1} | N_2 n_2 m_2 \rangle \\
&+ (\eta\sigma^2)^2 e^4 \langle N n_1 m_1 | r^{-2} | N_2 n_2 m_2 \rangle \\
&+ \left[ \frac{(\eta\sigma^2)^4 e^6 q}{2} (n'_2)^{-2} \langle N_1 n_1 m_1 | r^{-2} | N_2 n_2 m_2 \rangle \right. \\
&\quad \left. + \frac{(\eta\sigma^2) e^2 q \hbar^2}{\mu} \langle N_1 n_1 m_1 | r^{-3} | N_2 n_2 m_2 \rangle \right] \\
&\times \frac{\Gamma(2k_1 + 1)}{\Gamma(k_1 + 1)} \frac{\Gamma(2k_1 + 1)}{\Gamma(k_1 + 1)} \left[ \frac{(2n_1 + 2k_1 + 1)}{2^{2(k_1+k_2+1)} \Gamma(n_1 + 2k_1 + 1)} \frac{(2n_2 + 2k_2 + 1)}{\Gamma(n_2 + 2k_2 + 1)} \right]^{1/2} \\
&\times \frac{\pi 2^{2-k_1-k_2}}{\Gamma(k_1 + 1/2) \Gamma(k_2 + 1/2)} \left| \cos \frac{\pi}{2} (n_1 + n_2) \right| \\
&\times \sum_{p=0}^{[n_1/2]} \left[ \frac{[n_1 - 2p + (k_1 + k_2 - 1)/2]}{(n_1 - 2p)! p! s!} \right. \\
&\times \frac{\Gamma(n_1 - 2p + k_1 + k_2 - 1) \Gamma(n_2 + k_1 - s + 1/2)}{\Gamma(n_2 + \frac{k_1+k_2+1}{2} - s)} \\
&\times \frac{\Gamma(n_1 - k_1 - p - 1/2)}{\Gamma(n_1 + \frac{k_1+k_2+1}{2} - p)} \left( \frac{k_1 - k_2}{2} + 1 \right)_p \left( \frac{k_2 - k_1}{2} + 1 \right)_s \\
&+ \left| \cos \frac{\pi}{2} (n_1 + n_2) \right| (\eta\sigma^2)^2 \frac{q^2 \hbar^4}{4\mu^2} \frac{\Gamma(2k_1 + 1)}{\Gamma(k_1 + 1)} \frac{\Gamma(2k_1 + 1)}{\Gamma(k_1 + 1)} \\
&\times \left[ \frac{(2n_1 + 2k_1 + 1)}{2^{2(k_1+k_2+1)} \Gamma(n_1 + 2k_1 + 1)} \frac{(2n_2 + 2k_2 + 1)}{\Gamma(n_2 + 2k_2 + 1)} \right]^{1/2} \\
&\times \frac{\pi 2^{4-k_1-k_2}}{\Gamma(k_1 + 1/2) \Gamma(k_2 + 1/2)} \sum_{p=0}^{[n_1/2]} \left[ \frac{[n_1 - 2p + (k_1 + k_2 - 3)/2]}{(n_1 - 2p)! p! s!} \right. \\
&\times \frac{\Gamma(n_1 - 2p + k_1 + k_2 - 3) \Gamma(n_2 + k_1 - s + 1/2)}{\Gamma(n_2 + \frac{k_1+k_2-1}{2} - s)} \\
&\times \frac{\Gamma(n_1 - k_1 - p - 1/2)}{\Gamma(n_1 + \frac{k_1+k_2-1}{2} - p)} \left( \frac{k_1 - k_2}{2} + 2 \right)_p \left( \frac{k_2 - k_1}{2} + 2 \right)_s \\
&\times \langle N_1 n_1 m_1 | r^{-4} | N_2 n_2 m_2 \rangle , \tag{19}
\end{aligned}$$

where  $\langle N_1 n_1 m_1 | r^s | N_2 n_2 m_2 \rangle$  are given by replacing  $s = \{-1, -2, -3, -4\}$  in (14).

### Example: The states $|010\rangle$ and $|100\rangle$

In the ordinary case (i.e.  $\beta = 0$ ),  $|010\rangle$  and  $|100\rangle$  are two degenerate states. It is clear that the matrix (11) is actually diagonal ( $\langle 100| p^4 |010\rangle = 0$ ) which can be seen by using the expressions:

$$L_0^{(\alpha)}(x) = 1, \quad L_1^{(\alpha)}(x) = -x + \alpha + 1 \quad (20)$$

$$C_0^{(\alpha)}(x) = 1, \quad C_1^{(\alpha)}(x) = 2\alpha x \quad (21)$$

From equation (11), a straightforward calculation gives the energy corrections at first order in the parameter  $\beta$  as follows:

$$\begin{aligned} \Delta E_{010} &= 4\mu^2 \left[ \left( E_{010}^{(0)} \right)^2 - \frac{abAE_{010}^{(0)}(2k_0+1)^2}{k_0+2} \left( \frac{2^{2k_0+1} [\Gamma(k_0+1)]^2}{\Gamma(2k_0+2)} - \frac{2^{2k_0+3} [\Gamma(k_0+2)]^2}{\Gamma(2k_0+4)} \right) \right. \\ &\quad - \frac{a^2bBE_{010}^{(0)}(2k_0+1)^2}{(k_0+2)(2k_0+3)} \left( \frac{2^{2k_0-1} [\Gamma(k_0)]^2}{\Gamma(2k_0)} - \frac{2^{2k_0+1} [\Gamma(k_0+1)]^2}{\Gamma(2k_0+2)} \right) \\ &\quad + \frac{a^2bA^2(2k_0+1)^2}{2(k_0+2)(2k_0+3)} \left( \frac{2^{2k_0+1} [\Gamma(k_0+1)]^2}{\Gamma(2k_0+2)} - \frac{2^{2k_0+3} [\Gamma(k_0+2)]^2}{\Gamma(2k_0+4)} \right) \\ &\quad + \frac{a^4bB^2(2k_0+1)}{2(k_0+2)(2k_0+3)(2k_0+2)} \left( \frac{2^{2k_0-1} [\Gamma(k_0-1)]^2}{\Gamma(2k_0-2)} - \frac{2^{2k_0+1} [\Gamma(k_0)]^2}{\Gamma(2k_0)} \right) \\ &\quad \left. + \frac{a^3bAB(2k_0+1)^2}{(k_0+2)(2k_0+3)(2k_0+2)} \left( \frac{2^{2k_0-1} [\Gamma(k_0)]^2}{\Gamma(2k_0)} - \frac{2^{2k_0+1} [\Gamma(k_0+1)]^2}{\Gamma(2k_0+2)} \right) \right] \\ \Delta E_{100} &= 4\mu^2 \left[ \left( E_{100}^{(0)} \right)^2 - \frac{abAE_{100}^{(0)}2^{2k_0+1}[\Gamma(k_0+1)]^2}{(2+k_0)\Gamma(2k_0+2)} \right. \\ &\quad - \frac{a^2bBE_{100}^{(0)}2^{2k_0-1}[\Gamma(k_0)]^2}{(2+k_0)(2k_0+1)\Gamma(2k_0)} + \frac{a^2bA^22^{2k_0}[\Gamma(k_0+1)]^2}{(2+k_0)(2k_0+1)\Gamma(2k_0+2)} \\ &\quad \left. + \frac{a^4bB^22^{2k_0-3}(2k_0-2)(4+k_0)[\Gamma(k_0-1)]^2}{(2+k_0)(2k_0+3)} + \frac{a^3bAB2^{2k_0}[\Gamma(k_0)]^2}{\Gamma(2k_0+3)} \right] \end{aligned} \quad (22)$$

where

$$E_{100}^{(0)} = E_{010}^{(0)} = -\frac{\mu(\eta\sigma^2)^2 e^4}{2\hbar^2} \left[ 2 + \sqrt{\frac{q\eta\sigma^2\hbar^2}{2\mu}} \right]^{-2}, \quad A = \eta\sigma^2 e^2, \quad B = \frac{\eta q\sigma^2\hbar^2}{2\mu}, \quad (23)$$

$$a = \frac{2\mu\eta\sigma^2 e^2}{\hbar^2 n'}, \quad b = \left[ \frac{\Gamma(2k+1)}{\Gamma(k+1)} \right]^2 \frac{(2n+2k+1)}{2^{2k+1}\Gamma(n+2k+1)}, \quad k_0 = \sqrt{\frac{q\eta\hbar^2\sigma^2}{2\mu}}. \quad (24)$$

Thus, at first order in the parameter  $\beta$ , the degeneracy of the two levels  $|010\rangle$  and  $|100\rangle$  is completely lifted.

### 3 Matrix elements for $\Delta N = 0$ , $\Delta n = 0$ and $\Delta m = 0$

Taking  $N_1 = N_2 = N$ ,  $n_1 = n_2 = n$  and  $m_1 = m_2 = m$ , we derive the explicit form of the energy corrections given in equation (11). Using the relation between the confluent hypergeometric function  $F(-n; l+1; z)$  and the associated Laguerre polynomials  $L_n^{(l)}(x)$ , namely:

$$L_n^{(l)}(z) = \frac{\Gamma(n+l+1)}{\Gamma(n+1)\Gamma(l+1)} F(-n; l+1; z), \quad (25)$$

where  $z = ar$  and  $a = \frac{2\mu\eta\sigma^2e^2}{\hbar^2n'}$ , and using the integral:

$$\begin{aligned} \int_0^\infty z^{l-1} e^{-z} [F(-n; \gamma; z)]^2 dx &= \frac{n! \Gamma(l)}{\gamma(\gamma+1) \cdots (\gamma+n-1)} \left\{ 1 + \right. \\ &\quad \frac{n(\gamma-l-1)(\gamma-l)}{1^2\gamma} + \frac{n(n-1)(\gamma-l-2)(\gamma-l-1)(\gamma-l)(\gamma-l+1)}{1^22^2\gamma(\gamma+1)} + \dots \\ &\quad \left. \dots + \frac{n(n-1) \cdots 1(\gamma-l-n) \cdots (\gamma-l+n-1)}{1^22^2 \cdots n^2\gamma(\gamma+1) \cdots (\gamma+n-1)} \right\}, \end{aligned} \quad (26)$$

we obtain the following average values for the radial part:

$$\langle Nnm | r^{-1} | Nnm \rangle = \int_0^\infty [R(r)]^2 r^{-1} dr = \frac{a}{2n'}, \quad (27)$$

$$\langle Nnm | r^{-2} | Nnm \rangle = \int_0^\infty [R(r)]^2 r^{-2} dr = \frac{1}{2l+1} \frac{a^2}{2n'}, \quad (28)$$

$$\langle Nnm | r^{-3} | Nnm \rangle = \int_0^\infty [R(r)]^2 r^{-3} dr = \frac{a^3}{2l(2l+1)(2l+2)} = \frac{a^3 \Gamma(2l)}{\Gamma(2l+3)}, \quad (29)$$

$$\begin{aligned} \langle Nnm | r^{-4} | Nnm' \rangle &= \int_0^\infty [R(r)]^2 r^{-4} dr \\ &= \frac{a^4}{n'} \left( \frac{3(n')^2 - l(l+1)}{(2l-1)2l(2l+1)(2l+2)(2l+3)} \right) \\ &= \frac{a^4 [3(n')^2 - l(l+1)]}{n'} \frac{\Gamma(2l-1)}{\Gamma(2l+4)} \end{aligned} \quad (30)$$

Now, we evaluate the following integral of the angular functions:

$$\begin{aligned} \int_{-1}^1 dx (1-x^2)^{-t} [\Theta(x)]^2 &= \frac{(2n+2k+1)}{2^{2k+1}\Gamma(n+2k+1)} \\ &\times \left[ \frac{\Gamma(2k+1)}{\Gamma(k+1)} \right]^2 \int_{-1}^1 dx (1-x^2)^{k-t} [C_n^{(k+1/2)}(x)]^2 \end{aligned} \quad (31)$$

We use the following expression given in Ref. [42]:

$$\int_0^\pi d\theta \sin^\nu \theta \sin(\gamma\theta) \left[ C_n^{(\lambda)} \left( \sqrt{1 + \rho \sin^2 \theta} \right) \right]^2 = \frac{2^{-\nu} \pi \Gamma(\nu + 1) (2\lambda)_n^2}{(n!)^2 \Gamma(\frac{\nu-\gamma}{2} + 1) \Gamma(\frac{\nu+\gamma}{2} + 1)} \\ \times \sin\left(\frac{\gamma\pi}{2}\right) {}_5F_4 \left( \begin{matrix} -n, \lambda, 2\lambda + n, \frac{\nu+1}{2}, 1 + \frac{\nu}{2}; -\rho \\ \lambda + \frac{1}{2}, 2\lambda, \frac{\nu-\gamma}{2} + 1, \frac{\nu+\gamma}{2} + 1 \end{matrix} \right), \quad (\text{Re}\nu > -1) \quad (32)$$

with  ${}_pF_q(a_1, a_2, \dots, a_p; b_1, b_2, \dots, b_q; x)$  being the hypergeometric function defined as:

$${}_pF_q \left( \begin{matrix} a_1, a_2, \dots, a_p \\ b_1, b_2, \dots, b_q \end{matrix} ; x \right) = {}_pF_q(a_1, a_2, \dots, a_p; b_1, b_2, \dots, b_q; x) \\ = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_p)_k}{(b_1)_k (b_2)_k \dots (b_q)_k} \frac{x^k}{k!} \quad (33)$$

We obtain:

$$\int_{-1}^1 dx (1-x^2)^{-1} [\Theta(x)]^2 = \frac{\pi}{2^{4k-3} (n!)^2} \frac{(2n+2k+1)}{[\Gamma(k+1)]^2} \frac{\Gamma(n+2k+1) \Gamma(2k-3)}{\Gamma(k-3/2) \Gamma(k-1/2)} \\ \times {}_5F_4 \left( \begin{matrix} -n, k+1/2, n+2k+1, k-1/2, k; 1 \\ k+1, 2k+1, k-1/2, k+1/2 \end{matrix} \right), \quad (34)$$

$$\int_{-1}^1 dx (1-x^2)^{-2} [\Theta(x)]^2 = \frac{\pi}{2^{4k} (n!)^2} \frac{(2n+2k+1)}{[\Gamma(k+1)]^2} \frac{\Gamma(n+2k+1) \Gamma(2k-1)}{\Gamma(k-1/2) \Gamma(k+1/2)} \\ \times {}_5F_4 \left( \begin{matrix} -n, k+\frac{1}{2}, n+2k+1, k-3/2, k-1; 1 \\ k+1, 2k+1, k-3/2, k-1/2 \end{matrix} \right). \quad (35)$$

Finally, the diagonal matrix elements up to first order in  $\beta$  take the form:

$$\frac{\beta}{4\mu} \langle Nnm | p^4 | Nnm \rangle = \beta \frac{a^4}{(n')^8} \left\{ \frac{\hbar^2}{8\mu} \left( \frac{\hbar^2}{2} + \frac{n'}{2l+1} \right) + \right. \\ + \frac{\eta q \sigma^2 \hbar^4}{\mu} \left( \frac{1}{2n' (2l+1)} + \frac{4n' \Gamma(2l)}{\Gamma(2l+3)} \right) \frac{\pi}{2^{4k-2} (n!)^2} \frac{(2n+2k+1)}{[\Gamma(k+1)]^2} \\ \times \frac{\Gamma(n+2k+1) \Gamma(2k-3)}{\Gamma(k-3/2) \Gamma(k-1/2)} {}_5F_4 \left( \begin{matrix} -n, k+1/2, n+2k+1, k-1/2, k; 1 \\ k+1, 2k+1, k-1/2, k+1/2 \end{matrix} \right) \\ \left. + \frac{\eta^2 q^2 \sigma^4 \hbar^4}{\mu} \frac{[3(n')^2 - l(l+1)] \Gamma(2l-1)}{\Gamma(2l+4) n'} \frac{\pi}{2^{4k+2} (n!)^2} \frac{(2n+2k+1)}{[\Gamma(k+1)]^2} \right. \\ \left. \times \frac{\Gamma(n+2k+1) \Gamma(2k-1)}{\Gamma(k-1/2) \Gamma(k+1/2)} {}_5F_4 \left( \begin{matrix} -n, k+\frac{1}{2}, n+2k+1, k-3/2, k-1; 1 \\ k+1, 2k+1, k-3/2, k-1/2 \end{matrix} \right) \right\} \quad (36)$$

This last expression depends on  $l$  ( $l = n+k$ ), which lifts the degeneracy.

## 4 Generalized Recurrence Relations

For the diagonal matrix elements, the radial part verifies the following recurrence relations given in Ref. [29](restoring  $\mu$ ,  $e$  and  $\hbar$ ):

$$\frac{\hbar^4 a^2}{4\mu^2 e^4} (s+1) \langle r^s \rangle = \frac{\hbar n' a}{2\mu e^2} \langle r^{s-1} \rangle - \frac{s \left[ (2l+1)^2 - s^2 \right]}{4} \langle r^{s-2} \rangle, \quad (37)$$

where the first average elements of  $r^s$  were evaluated in Ref. [29]. Next, we evaluate the recurrence formula for the angular part. For this we denote:

$$\begin{aligned} \langle \sin^{2t} \theta \rangle_{n,k} &= \langle Nnm | \sin^{2t} \theta | Nnm \rangle = \int_{-1}^1 dx (1-x^2)^t [\Theta(x)]^2 \\ &= \frac{(2n+2k+1)}{2^{2k+1} \Gamma(n+2k+1)} \left[ \frac{\Gamma(2k+1)}{\Gamma(k+1)} \right]^2 \int_{-1}^1 dx (1-x^2)^{k+t} \left[ C_n^{(k+1/2)}(x) \right]^2. \end{aligned} \quad (38)$$

We can write:

$$\begin{aligned} &\int_{-1}^1 dx (1-x^2)^{k+t+1} \left[ C_n^{(k+1/2)}(x) \right]^2 \\ &= \int_{-1}^1 dx (1-x^2)^{k+t} \left( \left[ C_n^{(k+1/2)}(x) \right]^2 - \left[ x C_n^{(k+1/2)}(x) \right]^2 \right). \end{aligned} \quad (39)$$

Using the following recurrence rule of the Gegenbauer polynomials [43]:

$$2\alpha (1-x^2) C_{n-1}^{(\alpha+1)}(x) = (2\alpha+n+1) C_{n-1}^{(\alpha)}(x) - nx C_n^{(\alpha)}(x), \quad (40)$$

we straightforwardly obtain:

$$\begin{aligned} \langle \sin^{2(t+1)} \theta \rangle_{n,k} &= \langle \sin^{2t} \theta \rangle_{n,k} - \frac{1}{4} (n+2k+1) \langle \sin^{2(t+1)} \theta \rangle_{n-1,k+1} \\ &\quad - \frac{(n+2k+2)(n+2k+1)}{n(2n+2k+1)} \langle \sin^{2t} \theta \rangle_{n-1,k+1} \\ &\quad - \frac{(n+2k+2)}{(2n+2k+1)(2n+2k-1)} \langle \sin^{2t} \theta \rangle_{n-1,k}. \end{aligned} \quad (41)$$

From equations (37) and (41) we can derive the general formula of the averages values of  $r^p \sin^{2s} \theta$ . The recurrence formula (41) requires the two initial values  $\langle \sin^{2t} \theta \rangle_{0,k}$  and  $\langle \sin^{2t} \theta \rangle_{1,k}$ . Then, taking the special cases (21) and using of the following integral [44]:

$$\int_0^\pi d\theta \left( z + \sqrt{z^2 - 1} \cos \theta \right)^\mu \sin^{2\nu-1} \theta = \frac{2^{2\nu-1} \Gamma(\mu+1) [\Gamma(\nu)]^2}{\Gamma(2\nu+\mu)} C_\mu^{(\nu)}(z), \quad (42)$$

with  $\text{Re}(\nu) > 0$ , we obtain the first matrix elements:

$$\langle \sin^{2t} \theta \rangle_{0,k} = \frac{2^{2t} \Gamma(2k+2)}{\Gamma[2(k+t+1)]} \left[ \frac{\Gamma(k+t+1)}{\Gamma(k+1)} \right]^2, \quad (43)$$

$$\langle \sin^{2t} \theta \rangle_{1,k} = \frac{2^{2t} (2k+3)}{2k+2t+3} \frac{[\Gamma(2k+1)]^4}{\Gamma(2k+2)\Gamma(2k+2t+1)} \left[ \frac{\Gamma(k+t+1)}{\Gamma(k+1)} \right]^2. \quad (44)$$

## 5 Conclusion

In this paper we studied the Schrödinger equation for the Hartmann potential with deformed Heisenberg algebra. Using perturbation theory at the first order in the parameter of deformation  $\beta$ , we obtained the general form of the hamiltonian matrix elements and, as an example, we showed that the degeneracy of the two states  $|010\rangle$  and  $|100\rangle$  is completely lifted. For the diagonal matrix elements, we derived an explicit analytical expression which depends on  $l$ . In this case, some degenerate states split into sub-levels, and new transitions appear. In addition to the recurrence formula for the radial average values given in [29], we derived the one for the angular part which leads to the general formula of the average values of  $r^p \sin^{2s} \theta$  for the non-relativistic Hartmann potential. These results are useful in the calculations of the bound-state transitions and, on the experimental side, the energy levels can be measured and an upper bound on the minimal length  $(\Delta X)_{\min}$  can be obtained.

## References

- [1] D. J. Gross and P. F. Mende, Nucl. Phys. B 303, 407 (1988)
- [2] M. Maggiore, Phys. Lett. B 304, 65 (1993)
- [3] E. Witten, Phys. Today 49, 24 (1996)
- [4] D. Amati, M. Ciafaloni, G. Veneziano, Phys. Lett. B 197, 81 (1987)
- [5] D. Amati, M. Ciafaloni, G. Veneziano, Phys. Lett. B 216, 41 (1989)
- [6] M. Maggiore, Phys. Lett. B 319, 83 (1993)
- [7] L.J. Garay, Int. J. Mod. Phys. A 10, 145 (1995)
- [8] S. Hossenfelder, Mod. Phys. Lett. A 19, 2727 (2004)
- [9] S. Hossenfelder, Phys. Rev. D 70, 105003 (2004)
- [10] S. Hossenfelder, Phys. Lett. B 598, 92 (2004)
- [11] M. Sprenger, P. Nicolini, M. Bleicher, Eur. J. Phys. 33, 853 (2012)
- [12] S. Haouat, Phys. Lett B 729, 33 (2014)

- [13] A. Kempf, J. Math. Phys. 35, 4483 (1994)
- [14] A. Kempf, G. Mangano and R. B. Mann, Phys. Rev. D 52, 1108 (1995)
- [15] H. Hinrichsen and A. Kempf, J. Math. Phys. 37, 2121 (1996)
- [16] A. Kempf, J. Math. Phys. 38, 1347 (1997)
- [17] A. Kempf, J. Phys. A 30, 2093 (1997)
- [18] B. Hamil and M. Merad, Few Body Syst. 60, 36 (2019)
- [19] Y. Chargui and A. Dahabi, Few Body Syst. 61, 2 (2020)
- [20] L.N. Chang, D. Minic, N. Okamura, T. Takeuchi, Phys. Rev. D 65, 125027 (2002)
- [21] F. Brau, J. Phys. A, Math. Gen. 32, 7691 (1999)
- [22] M.M. Stetsko, V.M. Tkachuk, Phys. Rev. A 74, 012101 (2006)
- [23] M.M. Stetsko, Phys. Rev. A 74, 062105 (2006)
- [24] D. Bouaziz, N. Ferkous, Phys. Rev. A 82, 022105 (2010)
- [25] P. Pedram, Europhys. Lett. 101, 30005 (2013)
- [26] D. Bouaziz, M. Bawin, Phys. Rev. A 78, 032110 (2008)
- [27] M.M. Stetsko, V.M. Tkachuk, Phys. Rev. A 76, 012707 (2007)
- [28] H. Hassanabadi, S. Zarrinkamar, E. Maghsoodi, Phys. Lett. B 718, 678 (2012)
- [29] Chang-Yuan Chen , Cheng-Lin Liu and Dong-Sheng Sun, Phys. Lett. A 305, 341 (2002)
- [30] Chang-Yuan Chen , Dong-Sheng Sun and Cheng-Lin Liu, Phys. Lett. A 317, 80 (2003)
- [31] C.C. Gerry, Phys. Lett. A 118, 445 (1986)
- [32] M. Kibler, T. Negadi, Int. J. Quantum Chem. 26, 405 (1984)
- [33] I. Sökmen, Phys. Lett. A 115, 249 (1986)
- [34] M. Kibler, T. Negadi, Theor. Chim. Acta 66, 31 (1984)
- [35] M. Kibler, P. Winternitz, J. Phys. A 20, 4097 (1987)
- [36] H. Hartmann, Theor. Chim. Acta 24, 201 (1972)
- [37] H. Hartmann, R. Schuck, J. Radtke, Theor. Chim. Acta 46, 1 (1976)
- [38] H. Hartmann, D. Schuck, Int. J. Quantum Chem. 18, 125 (1980)

- [39] F. Yasuk, C. Berkdemir and A. Berkdemir, J. Phys. A: Math. Gen. 38, 6579 (2005)
- [40] Sameer M. Ikhdair and Ramazan Sever, Int. J. Theor. Phys. 46, 2384 (2007)
- [41] F. El Wassouli, African Journal of Mathematical Physics 5, 51 (2007)
- [42] Yuri A. Brychkov, Handbook of special functions (CRC Press 2008)
- [43] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables, (Dover, (1965))
- [44] I. S. Gradshteyn and I. M. Ryzhik, Table of Integrals, Series and Products, 7th edition, (Acedemic Press, 2007)