

Optimal liquidation for a risk averse investor in a one-sided limit order book driven by a Lévy process

Arne Løkka*

Junwei Xu†

February 5, 2022

Abstract

In a one-sided limit order book, satisfying some realistic assumptions, where the unaffected price process follows a Lévy process, we consider a market agent that wants to liquidate a large position of shares. We assume that the agent has constant absolute risk aversion and aims at maximising the expected utility of the cash position at the end of time. The agent is then faced with the problem of balancing the market risk and the cost of a rapid execution. In particular we are interested in how the agent should go about optimally submitting orders. Since liquidation normally takes place within a short period of time, modelling the risk as a Lévy process should provide a realistic model with good statistical fit to observed market data, and thus the model should provide a realistic reflection of the agent's market risk. We reduce the optimisation problem to a deterministic two-dimensional singular problem, to which we are able to derive an explicit solution in terms of the model data. In particular we find an expression for the optimal intervention boundary, which completely characterise the optimal liquidation strategy.

1 Introduction

This paper is concerned with how a market agent should go about selling (or purchasing) a large position of shares. This kind of problem has attracted considerable interest over the past few years following the introduction of electronic trading platforms. In the model we consider,

*Department of Mathematics Columbia House London School of Economics Houghton Street, London WC2A 2AE United Kingdom (a.lokka@lse.ac.uk)

†Department of Mathematics Columbia House London School of Economics Houghton Street, London WC2A 2AE United Kingdom (j.xu19@lse.ac.uk)

we specify the limit order book and how this recovers over time. Thus the optimal liquidation strategy will explicitly specify the orders the agent submits to the market, as opposed to just specifying the optimal speed at which to trade that is the case for the popular impact models. We refer the reader to Lehalle and Laruelle (2013), Cartea et al. (2015) and Guéant (2016) for an introduction to optimal execution and common models.

More precisely, we consider a market agent with constant absolute risk aversion (CARA) that wants to maximise the expected utility of the cash position at the end of time. Thus the agent does not face any restrictions on the duration of the liquidation, but the rapidness is determined by the market risk and the agent's preference towards risk. Working with an infinite time-horizon also avoids the time dimension of the problem, and hence make the problem more tractable. We assume that the market risk of the stock price is modelled by a Lévy process, which is allowed to have a drift, but which we assume satisfy a certain exponential moment condition. A number of studies demonstrate that Lévy processes are able to capture the essential statistical properties of stock price movements over short time-horizons (Madan and Seneta, 1990), (Eberlein and Keller, 1995), (Barndorff-Nielsen, 1997) and (Cont and Tankov, 2004). Since the main bulk of the liquidation tends to finish within a short period of time, this model should provide a reasonable reflection of the market risk faced by the agent. For the same reason, a Lévy process model is a reasonable approximation of an exponential Lévy process model. Such a Bachelier-Lévy type model may seem simplistic, but this kind of modelling of the unaffected price process is widely used in the optimal liquidation literature (Almgren and Chriss, 2000), (Kissell and Malamut, 2005), (Schied and Schöneborn, 2009) and (Gatheral, 2010). In particular, Forsyth et al. (2012) demonstrated that the linear model provides an excellent approximation to models with prices modelled as a geometric Brownian motion and multiplicative impact in the Almgren-Chriss framework.

We consider a bid limit order book with general shape and with a general resilience function satisfying some reasonable conditions. In particular, some of these assumptions are crucial in order to solve the problem since the assumptions imply that a certain function is concave, which is needed for optimality of our strategy. So with an infinite time horizon, we solve the problem of maximising the expected utility of the agent's final cash position. We do this by showing that the problem can be reduced to a two-dimensional deterministic singular optimisation problem to which we can obtain an explicit solution in terms of the characteristics of the limit order book and the agent's risk aversion. With reference to Løkka (2014) and the nature of the problem, we guess that the optimal strategy consists initially of either a block

sale or a period of waiting, and that the agent thereafter continuously submit sales orders in such a way that the state process remains on the optimal intervention boundary. The state process here is the two-dimensional process consisting of the number of shares the agent currently hold and the current state of the order book. The optimal intervention boundary is associated with the Hamilton-Jacobi-Bellman (HJB) variational inequalities corresponding to the optimisation problem. This intervention boundary might have discontinuities as well as constant parts. The discontinuities corresponds to periods of waiting while the order book recovers, while the constant part corresponds to submitting sales orders at the same rate as the resilience rate. Following the idea in Løkka (2014), the value function in our problem can be expressed in an explicit way in terms of the problem's data, and we characterise the intervention boundary via the HJB variational inequalities. The strategy associated with this intervention boundary is shown to be optimal by a verification argument.

We also provide an example in order to illustrate the optimal strategies for various parameters of risk aversion for the case when the risk is modelled by a Brownian motion and for the case when the risk is modelled by a Lévy process with jumps. For the case of a Lévy process with jumps we choose the Lévy process approximation of the exponential variance-gamma process. We find that the two models produce similar optimal strategies if the agent is not very risk averse, but as the agent's stock position and the agent's risk aversion increase, the differences become more pronounced.

The model we use is a version of the model introduced in Obizhaeva and Wang (2013), which was later generalised in Alfonsi et al. (2010), and then further in Predoiu et al. (2011). However, these papers did not consider risk-aversion, but assume that the agent wants to maximise the expected value of the cash position. The problem we consider in this paper is an extension of Løkka (2014) in the sense that unaffected price process follows a general Lévy process which could have a drift, and not simply a Brownian motion. Moreover, in this paper we consider an order book with a general resilience function, not just exponential resilience.

This paper is structured as follows. In Section 2 we introduce the limit order book model and the agent's optimisation problem. We simplify the problem and show that it can be reduced to a deterministic optimisation problem in Section 3. The simplified deterministic optimisation problem is solved in Section 4. The proofs omitted in the previous sections are contained in Section 5.

2 Problem formulation

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a complete filtered probability space satisfying the usual conditions and supporting a one-dimensional non-trivial Lévy process L .

Assumption 2.1. We assume that there exists some $\delta > 0$ such that $\mathbb{E}[e^{\theta L_1}] < \infty$, for $|\theta| < \delta$.

Let κ denote the cumulant generating function of L_1 , i.e.

$$\kappa(\theta) = \ln(\mathbb{E}[e^{\theta L_1}]), \quad \theta \in \mathbb{R}.$$

Assumption 2.1 guarantees that the cumulant generating function κ is continuously differentiable in a neighbourhood of 0 and that the Lévy process L is square integrable. Hence L admits the representation

$$L_t = \mu t + \sigma W_t + \int_{\mathbb{R} \setminus \{0\}} z(N(t, dz) - t\nu(dz)), \quad t \geq 0,$$

where $\mu \in \mathbb{R}$, W is a standard Brownian motion, N is a Poisson random measure which is independent of W with compensator $\pi(t, dz) = t\nu(dz)$, where ν denotes the Lévy measure associated with L (Kyprianou, 2006). The cumulant generating function κ can then be expressed as

$$\kappa(\theta) = \mu\theta + \frac{1}{2}\sigma^2\theta^2 + \int_{\mathbb{R} \setminus \{0\}} (e^{\theta z} - 1 - \theta z) \nu(dz), \quad |\theta| < \delta. \quad (2.1)$$

In particular,

$$\kappa(0) = 0, \quad \kappa'(0) = \mu \quad \text{and} \quad \kappa''(0) = \sigma^2 + \int_{\mathbb{R} \setminus \{0\}} z^2 \nu(dz).$$

Moreover, κ is strictly convex and continuously differentiable on its effective domain.

Assumption 2.2. We assume that $\mu \leq 0$, i.e. the Lévy process L is a supermartingale.

The main reason behind this assumption is that the optimisation problem we consider in this paper does not have a solution when μ is positive. This is because when μ is positive, it is optimal for the agent to hold on to some amount of shares for as long as possible, and as a consequence our formulation of the liquidation problem does not have a solution when μ is

positive (see Remark 4.4). For the remainder of the paper, we therefore restrict ourselves to the case $\mu \leq 0$ for which our optimisation problem is well formulated. To simplify notation set

$$\mathbb{R}^+ = [0, \infty) \quad \text{and} \quad \mathbb{R}^- = (-\infty, 0].$$

We consider a market agent that aims to sell a large amount of shares of a single stock with no time restrictions. Let Y_t denote the number of shares held by the agent at time t . We refer to a process Y as a liquidation strategy if Y_t tends to 0 as t tends to infinity. We consider the following set of admissible liquidation strategies.

Definition 2.3. For $y \in \mathbb{R}^+$, let $\mathcal{A}(y)$ denote the set of all $(\mathcal{F}_t)_{t \geq 0}$ -adapted, predictable, decreasing, càdlàg processes Y , satisfying $Y_{0-} = y$ and

$$\int_0^\infty \kappa_A(\|Y_t\|_{L^\infty(\mathbb{P})}) dt < \infty. \quad (2.2)$$

Moreover, let $\mathcal{A}_D(y)$ denote the set of all deterministic strategies in $\mathcal{A}(y)$.

Remark 2.4. Condition (2.2) amounts to the following

$$\begin{aligned} \int_0^\infty \|Y_t\|_{L^\infty(\mathbb{P})} dt &< \infty, & \text{if } \mu < 0, \\ \int_0^\infty \|Y_t\|_{L^\infty(\mathbb{P})}^2 dt &< \infty, & \text{if } \mu = 0. \end{aligned}$$

When $\mu > 0$ (which we do not consider), condition (2.2) would have to be replaced with a condition like the one for $\mu < 0$ above. Conditions like these are not just technical conditions, but are necessary in order for the problem that we consider in this paper to be meaningful. If we do not impose any restrictions, then the value function is infinite. With reference to (2.11), the reason is that since there is no discounting, there is no penalty in waiting until the best bid price reach a certain level and then start liquidating. For instance if L is a Brownian motion, then the time it takes for the best bid price to reach a given level p is finite almost surely, for any level p . We skip the mathematical details, but intuitively a higher price level p results in a higher expected utility of the cash position. Consequently the value function will be infinite if such strategies are admissible.

To describe the agent's execution price, we explicitly model a bid limit order book. We assume that the unaffected bid price process B^0 , which is the process describing the best bid

prices in the market if the agent does not trade, is given by

$$B_t^0 = b + L_t, \quad t \geq 0,$$

where $b > 0$ is the best bid price at the initial time. This Bachelier-Lévy type price model may seem simplistic, but this kind of modelling of the unaffected price process is widely used in the optimal liquidation literature (Almgren and Chriss, 2000), (Kissell and Malamut, 2005), (Schied and Schöneborn, 2009) and (Gatheral, 2010). There are studies which show that liquidation models with linear price processes provide a good approximation to models with exponential price processes (Gatheral and Schied, 2011) and Forsyth et al. (2012).

In order to describe the bid limit order book, we consider a measure m defined on the Borel σ -algebra on \mathbb{R}^- , denoted by $\mathcal{B}(\mathbb{R}^-)$. If $\mathcal{S} \in \mathcal{B}(\mathbb{R}^-)$, then $m(\mathcal{S})$ represents the number of bid orders with prices in the set $B_t^0 + \mathcal{S} = \{B_t^0 + s \mid s \in \mathcal{S}\}$, provided that the agent did not make any trades before time t . Notice that the undisturbed bid order book described by m is relative to the unaffected bid prices in the sense that it shifts together with the movement of the unaffected price. We impose the following assumptions on m .

Assumption 2.5. We assume that

- (i) there exists some $\bar{x} \in (-\infty, 0)$ such that $m((\bar{x}, 0]) = m(\mathbb{R}^-) < \infty$,
- (ii) m is absolutely continuous with respect to Lebesgue measure, and is non-zero on any interval properly containing the origin,
- (iii) the function $x \mapsto m((x, 0])$ is concave in x , for $x \in \mathbb{R}^-$.

The concavity of $x \mapsto m((x, 0])$ means that there are less and less bid orders in the undisturbed limit order book the further we get from the best bid price. Set $\bar{z} = -m(\mathbb{R}^-)$, which represents the total amount of bid limit orders in the undisturbed limit order book, and introduce the functions $\phi : [-\infty, 0] \rightarrow \mathbb{R}^-$ and $\psi : \mathbb{R}^- \rightarrow [-\infty, 0]$ by

$$\phi(x) = -m((x, 0]) \quad \text{and} \quad \psi(z) = \phi^{-1}(z),$$

where $\phi(\psi(z)) = z$, for all $z \in [\bar{z}, 0]$, and $\psi(z) = -\infty$, for all $z < \bar{z}$. A consequence of Assumption 2.5 is that ϕ is convex, ψ is concave, and ϕ and ψ are both continuous and strictly increasing on their effective domain. They also satisfy

$$\phi(0) = \psi(0) = 0, \tag{2.3}$$

as well as

$$\int_0^{\bar{z}} \psi(u) du < \infty \quad \text{and} \quad \psi(\bar{z}) > -\infty. \quad (2.4)$$

In order to model the dynamic of the bid order book during trading, we need to introduce one more process that captures the state of the order book. For a given strategy Y , let Z^Y be an \mathbb{R}^- -valued process such that $-Z_t^Y$ represents the volume spread at time t . That is $-Z_t^Y$ is equal to the total number of bid orders which have already been executed subtracted by the total amount of limit orders which have arrived to refill the book up to time t . We call Z^Y the state process of the bid limit order book associated with a trading strategy Y . Let $Z_{0-}^Y = z$, where $z \geq \bar{z}$ is the initial state of our bid order book. Therefore, we have $\psi(Z_t^Y) = B_t^Y - B_t^0$, where B_t^Y is the best bid price at time t corresponding to Y , and $\psi(Z_t^Y)$ can be understood as the extra price spread at time t , caused by the investor who implements a strategy Y . Note that we have defined $\psi(z) = -\infty$, for all $z < \bar{z}$. This implies that the best bid price drops down to $-\infty$, if one sell more shares than available bids in the book. The rate at which bid orders are refilling the order book is described by a resilience function $h : \mathbb{R}^- \rightarrow \mathbb{R}^-$ which satisfies the following.

Assumption 2.6. We assume that the resilience function $h : \mathbb{R}^- \rightarrow \mathbb{R}^-$ is increasing, locally Lipschitz continuous, satisfies $h(0) = 0$ with $h(x) < 0$ for all $x < 0$, and that the function $x \mapsto 1/h(x)$ is a concave function for $x < 0$.

Note that the choice $h(x) = \lambda x$, for $\lambda > 0$, which corresponds to exponential resilience, satisfies Assumption 2.6. We then consider the state process Z^Y with dynamic

$$dZ_t^Y = -h(Z_{t-}^Y) dt + dY_t, \quad Z_{0-}^Y = z \in \mathbb{R}^-. \quad (2.5)$$

For any admissible strategy Y , we refer to Predoiu et al. (2011) Appendix A for the existence and uniqueness of a negative, càdlàg and adapted solution to this dynamic. From Assumption 2.6 and equation (2.5) we observe that the further the best bid price is away from the unaffected bid price, the larger the speed of resilience for the best bid price.

If the agent does not make any trades from time t_1 to t_2 , then $(Z_t^Y)_{t_1 < t < t_2}$ satisfies

$$dZ_t^Y = -h(Z_t^Y) dt. \quad (2.6)$$

Now define a strictly decreasing function $H : \mathbb{R}^- \rightarrow \mathbb{R} \cup \{-\infty\}$ by

$$H(x) = \int_{-1}^x \frac{1}{h(u)} du. \quad (2.7)$$

Let H^{-1} denote the inverse of H , which satisfies $H^{-1}(H(x)) = x$ for all $x \leq 0$ and $H^{-1}(u) = 0$ for $u \in (-\infty, \lim_{x \rightarrow 0-} H(x)]$. Then, it can be verified that the process Z given by

$$Z_t = H^{-1}(H(Z_0) - t) \quad (2.8)$$

has dynamic (2.6). Hence, for any t between time t_1 and t_2 , $Z_t^Y = H^{-1}(H(Z_{t_1}^Y) - t + t_1)$. Moreover, if $Z_{t_2}^Y < 0$, then

$$t_2 - t_1 = H(Z_{t_1}^Y) - H(Z_{t_2}^Y). \quad (2.9)$$

Suppose that the agent's initial cash position is c and that the agent implements a strategy $Y \in \mathcal{A}(y)$. Then the agent's cash position at time $T > 0$ is

$$C_T(Y) = c - \int_0^T B_{t-}^Y dY_t^c - \sum_{0 \leq t \leq T} \int_0^{\Delta Y_t} \{B_{t-}^0 + \psi(Z_{t-}^Y + x)\} dx, \quad (2.10)$$

which corresponds to the best bids offered at all times being executed first so as to match the agent's orders, where the first integral represents the cost from the continuous component of the liquidation strategy and the sum of integrals represents the total cost due to all block sales. We also suppose the agent has a constant absolute risk aversion (CARA). With initial cash position c , an initial share position y and infinite time-horizon, the agent wants to maximise the expected utility of the cash position at the end of time. Mathematically, the agent's optimal liquidation problem is

$$\sup_{Y \in \mathcal{A}(y)} \mathbb{E}[U(C_\infty(Y))], \quad (2.11)$$

where the utility function U is given by

$$U(c) = -e^{-Ac}, \quad A > 0.$$

Observe that if $Z_t^Y < \bar{z}$, then $B_t^Y = B_t^0 + \psi(Z_t^Y) = -\infty$. Clearly receiving the price $-\infty$

is unfavourable to the agent. Indeed, (2.10) shows that this brings the agent an infinite cost. Due to this consideration, we will from now on only focus on admissible strategies Y for which $Z_t^Y \geq \bar{z}$, for all $t \geq 0$. Define the function $\kappa_A : \mathbb{R}^+ \rightarrow [0, \infty]$ by

$$\kappa_A(y) = \kappa(-Ay), \quad y \geq 0,$$

and set

$$\bar{y}_A = \sup\{y \geq 0 \mid \kappa_A(y) < \infty\}.$$

Then κ_A is strictly increasing, strictly convex and continuously differentiable on $[0, \bar{y}_A)$, with $\kappa_A(0) = 0$.

3 Problem simplification

In this section, we show that the utility maximisation problem in (2.11) can be reduced to a deterministic optimisation problem. This kind of result was first derived in Schied et al. (2010), which proved that with a certain market structure and an agent with constant absolute risk aversion, the optimal liquidation strategy is deterministic.

Let $Y \in \mathcal{A}(y)$. Then it follows from (2.10) that

$$C_T(Y) = c + by - (b + L_T)Y_T + \int_0^T Y_{t-} dL_t + \sum_{0 \leq t \leq T} \Delta L_t \Delta Y_t - F_T(Y),$$

where F_T is given by

$$F_T(Y) = \int_0^T \psi(Z_{t-}^Y) dY_t^c + \sum_{0 \leq t \leq T} \int_0^{\Delta Y_t} \psi(Z_{t-}^Y + x) dx. \quad (3.1)$$

Since $t \mapsto \|Y_T\|_{L^\infty(\mathbb{P})}$ is decreasing, condition (2.2) implies that any admissible strategy $Y \in \mathcal{A}(y)$ satisfies

$$\lim_{t \rightarrow \infty} t \kappa_A(\|Y_t\|_{L^\infty(\mathbb{P})}) = 0. \quad (3.2)$$

Also observe that

$$\lim_{x \rightarrow 0} \frac{\kappa_A(x)}{x} = -A\mu.$$

Therefore, if $\mu < 0$, there exists an $\epsilon > 0$ and constants $C_1, C_2 > 0$ such that

$$C_1 x \leq \kappa_A(x) \leq C_2 x, \quad \text{for } x \in [0, \epsilon].$$

With reference to (3.2), It follows that for every $Y \in \mathcal{A}(y)$,

$$\lim_{t \rightarrow \infty} t \|Y_t\|_{L^\infty(\mathbb{P})} = 0, \quad \text{if } \mu < 0. \quad (3.3)$$

If $\mu = 0$, then

$$\lim_{x \rightarrow 0} \frac{\kappa_A(x)}{x^2} = K, \quad \text{for some } K > 0.$$

Therefore, if $\mu = 0$, there exists an $\epsilon > 0$ and constants $C_1, C_2 > 0$ such that

$$C_1 x^2 \leq \kappa_A(x) \leq C_2 x^2, \quad \text{for } x \in [0, \epsilon].$$

With reference to (3.2) it follows that for every $Y \in \mathcal{A}(y)$,

$$\lim_{t \rightarrow \infty} t \|Y_t\|_{L^\infty(\mathbb{P})}^2 = 0, \quad \text{if } \mu = 0. \quad (3.4)$$

Let Y be an admissible strategy in $\mathcal{A}(y)$. Then with reference to (3.3) and (3.4), we calculate

$$\lim_{T \rightarrow \infty} \mathbb{E}[|L_T Y_T|^2] \leq \lim_{T \rightarrow \infty} (\mu^2 T^2 + \kappa''(0)T) \|Y_T\|_{L^\infty(\mathbb{P})}^2 = 0.$$

We conclude that $B_T^0 Y_T$ tends to 0 in $L^2(\mathbb{P})$ as $T \rightarrow \infty$. Furthermore,

$$\begin{aligned} \mathbb{E} \left[\left(\int_0^\infty Y_{t-} dL_t \right)^2 \right]^{\frac{1}{2}} &\leq \mathbb{E} \left[\left(\int_0^\infty Y_{t-} \mu dt \right)^2 \right]^{\frac{1}{2}} + \mathbb{E} \left[\left(\int_0^\infty Y_{t-} d(L_t - \mu t) \right)^2 \right]^{\frac{1}{2}} \\ &\leq \mu \int_0^\infty \|Y_t\|_{L^\infty(\mathbb{P})} dt + \left[\kappa''(0) \int_0^\infty \|Y_t\|_{L^\infty(\mathbb{P})}^2 dt \right]^{\frac{1}{2}} < \infty. \end{aligned}$$

Hence, $\int_0^\infty Y_{t-} dL_t$ is well-defined in $L^2(\mathbb{P})$. Due to the predictability of Y , we also have that

$$\mathbb{E} \left[\left(\sum_{0 \leq t \leq T} \Delta L_t \Delta Y_t \right)^2 \right] = \mathbb{E} \left[\int_0^T (\Delta Y_t)^2 dt \right] \left(\int_{\mathbb{R} \setminus \{0\}} z^2 \nu(dz) \right) = 0,$$

for all $T > 0$, which shows that the quadratic covariation of the jumps of L and Y is almost surely 0. Moreover, note that $F_T(Y) \geq 0$ is an increasing function of T . Therefore, F_∞ is a well defined function from the set of càdlàg non-increasing functions into the extended positive real numbers. The final cash position is hence given by

$$C_\infty(Y) = c + by + \int_0^\infty Y_{t-} dL_t - F_\infty(Y), \quad (3.5)$$

where $c + by$ represents the mark-to-market value of the total wealth of the agent's position at the start of the liquidation, $\int_0^\infty Y_{t-} dL_t$ represents the profit or loss due to the market risk, and $F_\infty(Y)$ represents the cost due to the price impact.

Let $Y \in \mathcal{A}(y)$ and define the process M^Y by

$$M_t^Y = \exp \left(-A \int_0^t Y_{s-} dL_s - \int_0^t \kappa_A(Y_{s-}) ds \right), \quad t \geq 0.$$

Then it follows from Theorem 3.2 in Kallsen and Shiryaev (2002) that M^Y is a uniformly integrable martingale. We can therefore define a probability measure $\tilde{\mathbb{P}} = \mathbb{P}^Y$ by

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = M_\infty^Y.$$

Following the idea of the proof of Theorem 2.1 in Schied et al. (2010), we set

$$I = \inf_{Y \in \mathcal{A}_D(y)} \int_0^\infty \kappa_A(Y_{t-}) dt + AF_\infty(Y),$$

and note that $\kappa_A(\cdot)$ and $F_\infty(\cdot)$ are deterministic. Let $Y^\epsilon \in \mathcal{A}_D(y)$ be such that

$$\int_0^\infty \kappa_A(Y_{t-}^\epsilon) dt + AF_\infty(Y^\epsilon) \leq I + \epsilon.$$

For an arbitrary $Y \in \mathcal{A}(y)$, we calculate that

$$\begin{aligned}
\mathbb{E}[U(C_\infty(Y))] &= -e^{-A(c+by)} \mathbb{E} \left[\exp \left(-A \int_0^\infty Y_{t-} dL_t + AF_\infty(Y) \right) \right] \\
&= -e^{-A(c+by)} \mathbb{E} \left[M_\infty \exp \left(\int_0^\infty \kappa_A(Y_{t-}) dt + AF_\infty(Y) \right) \right] \\
&= -e^{-A(c+by)} \tilde{\mathbb{E}} \left[\exp \left(\int_0^\infty \kappa_A(Y_{t-}) dt + AF_\infty(Y) \right) \right] \\
&\leq -e^{-A(c+by)} e^{-\epsilon} \tilde{\mathbb{E}} \left[\exp \left(\int_0^\infty \kappa_A(Y_{t-}^\epsilon) dt + AF_\infty(Y^\epsilon) \right) \right] \\
&\leq -e^{-A(c+by)} e^{-\epsilon} \exp \left(\inf_{Y \in \mathcal{A}_D(y)} \left\{ \int_0^\infty \kappa_A(Y_{t-}) dt + AF_\infty(Y) \right\} \right),
\end{aligned}$$

since

$$\int_0^\infty \kappa_A(Y_{t-}) dt + AF_\infty(Y) \geq I \geq \int_0^\infty \kappa_A(Y_{t-}^\epsilon) dt + AF_\infty(Y^\epsilon) - \epsilon.$$

By letting ϵ tend to 0 and taking the supremum over all admissible strategies on the left-hand side of (3.6), we obtain

$$\sup_{Y \in \mathcal{A}(y)} \mathbb{E}[U(C_\infty(Y))] = -e^{-A(c+by)} \exp \left(\inf_{Y \in \mathcal{A}_D(y)} \left\{ \int_0^\infty \kappa_A(Y_{t-}) dt + AF_\infty(Y) \right\} \right) \quad (3.6)$$

Lemma 3.1. *Let F be given by (3.1). Then for every $Y \in \mathcal{A}_D(y)$ and $z \in [\bar{z}, 0]$,*

$$F_\infty(Y) = \int_z^0 \psi(s) ds + \int_0^\infty h(Z_{t-}^Y) \psi(Z_{t-}^Y) dt. \quad (3.7)$$

With reference to Lemma 3.1 and (3.6), the optimal liquidation problem amounts to solving

$$V(y, z) = \inf_{Y \in \mathcal{A}_D(y)} \int_0^\infty \left(\kappa_A(Y_{t-}) + Ah(Z_{t-}^Y) \psi(Z_{t-}^Y) \right) dt, \quad (3.8)$$

with $y = Y_{0-}$ and $z = Z_{0-}^Y$. Since h and ψ are both negative-valued and $\kappa_A \geq 0$, we have $V \geq 0$. Suppose $y > \bar{y}_A$, which is the upper bound for which κ_A is finite (\bar{y}_A might be $+\infty$). In this case, the market agent will want to make an immediate block sale to bring the number of shares less than \bar{y}_A , since otherwise Y does not satisfy (2.2) and $V(y, z) = \infty$. However, if

the agent sell more than $z - \bar{z}$ number of shares, the value function $V(y, z)$ will be infinite. We therefore define the solvency region to be

$$\mathcal{D} = \{ (y, z) \in \mathbb{R}^+ \times [\bar{z}, 0] \mid z > y - \bar{y}_A + \bar{z} \},$$

and for the remainder of the paper we focus on this region. For technical reasons, we do not consider $z = y - \bar{y}_A + \bar{z}$, as the value function may explode also along this line.

4 Solution to the problem

Our next aim is to derive a solution to the problem (3.8). The derivation will be based on applying a time-change, and the principle of dynamic programming. With reference to the results in Løkka (2014) and the general theory of optimal control (Fleming and Soner, 2006), it is natural to guess that there exists a decreasing¹ càglàd function $\beta = \beta^* : \mathbb{R}^+ \rightarrow [\bar{z}, 0]$ which separates the (y, z) domain into two different regions, a region where the agent makes an immediate block sale and another where the agent waits for the order book to recover. Let β_* denote the càdlàg version of β^* , and set

$$\begin{aligned} \bar{\mathcal{S}}^\beta &= \{ (y, z) \in \mathcal{D} \mid z \geq \beta_*(y) \}, \\ \bar{\mathcal{W}}^\beta &= \{ (y, z) \in \mathcal{D} \mid z \leq \beta^*(y) \} \cup \{ (y, z) \mid y = 0 \}, \\ \mathcal{G}^\beta &= \bar{\mathcal{S}}^\beta \cap \bar{\mathcal{W}}^\beta. \end{aligned}$$

$\bar{\mathcal{S}}^\beta$ represents the immediate sales region, $\bar{\mathcal{W}}^\beta$ the waiting region, and \mathcal{G}^β is the graph of the intervention boundary β and represents the continuous sales region. For $y > 0$, the Hamilton-Jacobi-Bellman equation corresponding to V given by (3.8) takes the form

$$D_y^- v(y, z) + v_z(y, z) = 0, \quad \text{for } (y, z) \in \bar{\mathcal{S}}^\beta, \quad (4.1)$$

$$h(z)v_z(y, z) - \kappa_A(y) - Ah(z)\psi(z) \leq 0, \quad \text{for } (y, z) \in \bar{\mathcal{S}}^\beta \setminus \mathcal{G}^\beta, \quad (4.2)$$

¹When the volume spread is small, but the stock position is large, it seems intuitive to sell rapidly. On the other hand, if the volume spread is large, but the stock position is small, it seems intuitive to wait for the order book to recover. This motivates us to guess that the optimal intervention boundary is decreasing and separates the (y, z) domain.

and

$$h(z)v_z(y, z) - \kappa_A(y) - Ah(z)\psi(z) = 0, \quad \text{for } (y, z) \in \overline{\mathcal{W}}^\beta, \quad (4.3)$$

$$D_y^- v(y, z) + v_z(y, z) \leq 0, \quad \text{for } (y, z) \in \overline{\mathcal{W}}^\beta \setminus \mathcal{G}^\beta, \quad (4.4)$$

with associated boundary condition $v(0, z) = A \int_0^z \psi(u) du$ for all $z \in [\bar{z}, 0]$, where²

$$D_y^- v(y, z) = \lim_{\epsilon \rightarrow 0^-} \frac{1}{\epsilon} \left(v(y + \epsilon, z) - v(y, z) \right).$$

The equations (4.1)–(4.4) can be motivated as follows. When the market agent is trying to optimise over deterministic strategies, the agent basically has two options. The agent can either sell a certain number $\Delta > 0$ of shares or wait. Given a state (y, z) , it may or may not be optimal to sell Δ amount of shares, thus

$$v(y, z) \leq v(y - \Delta, z - \Delta),$$

because the share position is decreased from y to $y - \Delta$, due to Δ number of shares being sold, while at the same time the state of the bid order book changes from z to $z - \Delta$. This inequality should hold for all $0 < \Delta \leq y$, therefore

$$\max_{0 < \Delta \leq y} \{v(y, z) - v(y - \Delta, z - \Delta)\} \leq 0. \quad (4.5)$$

On the other hand, during a period of time $\Delta t > 0$, it may or may not be optimal to wait, hence

$$\begin{aligned} v(y, z) &\leq v(y, Z_{\Delta t}) + \int_0^{\Delta t} \left(\kappa_A(y) + Ah(Z_{u-})\psi(Z_{u-}) \right) du \\ &= v(y, z) + \int_0^{\Delta t} \left(\kappa_A(y) + Ah(Z_{u-})(Z_{u-}) - v_z(y, Z_{u-})h(Z_{u-}) \right) du, \end{aligned}$$

²The value function turns out to be continuously differentiable in z , but only continuous with a one-sided derivative in y (see Proposition 4.7).

where $dZ_u = -h(Z_u) du$, for $0 \leq u \leq \Delta t$. Multiplying the above inequality by $(\Delta t)^{-1}$ and sending Δt to 0, we obtain

$$h(z)v_z(y, z) - \kappa_A(y) - Ah(z)\psi(z) \leq 0. \quad (4.6)$$

Since one of these strategies should be optimal, equality should hold in either (4.5) or (4.6). We therefore get

$$\max \left\{ \max_{0 < \Delta \leq y} \{v(y, z) - v(y - \Delta, z - \Delta)\}, h(z)v_z(y, z) - \kappa_A(y) - Ah(z)\psi(z) \right\} = 0,$$

from which (4.1)–(4.4) follow.

We define the liquidation strategy Y^β corresponding to an intervention boundary β as the càdlàg function with the following properties:

- (i) If $(y, z) \in \overline{\mathcal{S}}^\beta$, then the agent makes an immediate block trade of size Δ such that $(Y_0^\beta, Z_0^\beta) = (y - \Delta, z - \Delta) \in \mathcal{G}^\beta$, and set $t_w = 0$.
- (ii) If $(y, z) \in \overline{\mathcal{W}}^\beta$, then the agent waits until the time $t_w = \inf\{t \geq 0 \mid Z_t^{Y^\beta} = \beta(y)\}$, where

$$Z_t^{Y^\beta} = z - \int_0^t h(Z_u^{Y^\beta}) du, \quad 0 \leq t \leq t_w.$$

- (iii) For $t \geq t_w$, the agent continuously sell shares in such a way that $(Y_t^\beta, Z_t^{Y^\beta}) \in \mathcal{G}^\beta$, where

$$Z_t^{Y^\beta} = Z_{t_w}^{Y^\beta} - \int_{t_w}^t h(Z_u^{Y^\beta}) du + Y_t^\beta - Y_{t_w}^\beta, \quad t \geq t_w.$$

- (iv) The agent takes no further action once $Y_t^\beta = 0$.

Figure 1 provides an illustration of such a strategy. We will later characterise an optimal intervention boundary, and prove that the corresponding strategy exists, is admissible and optimal. The key to characterise the optimal intervention boundary is that we are able to obtain expressions for the performance of the strategy corresponding to a given intervention boundary. But first, let us examine in more detail the strategy corresponding to a given intervention boundary function β . We will consider any intervention boundary $\beta : \mathbb{R}^+ \rightarrow [\bar{z}, 0]$ which is decreasing, càglàd and satisfy $\beta(y) < 0$, for all $y > 0$, $\lim_{y \rightarrow \infty} \beta(y) = \bar{z}$ and $\beta(0) = 0$. Now, given an intervention boundary β , one may ask whether the corresponding liquidation




Figure 1: An illustration of the strategy Y^β corresponding to an intervention boundary β (the graph of β is \mathcal{G}^β). The solvency region \mathcal{D} is the region that is not shaded. For an initial state $(y, z) \in \bar{\mathcal{S}}^\beta$, the strategy Y^β consists of an initial block sale that brings the state onto \mathcal{G}^β . For an initial state $(y, z) \in \bar{\mathcal{W}}^\beta$, the strategy Y^β consists of waiting while the order book recovers until the state reach \mathcal{G}^β . Once the state is on \mathcal{G}^β , the strategy Y^β consists of continuously submitting sales orders in such a way that the state stays on \mathcal{G}^β .

strategy Y^β exists and is unique. In order to answer this, we need to introduce the following functions related to β .

$$\gamma_\beta(y) = \beta(y) - y, \quad \text{for } y \in \mathbb{R}^+, \quad (4.7)$$

$$\rho_\beta(z) = z - \beta^{-1}(z), \quad \text{for } z \in [\bar{z}, 0]. \quad (4.8)$$

We also introduce the inverse functions

$$\beta^{-1}(z) = \inf\{y \in \mathbb{R}^+ \mid \beta(y) \leq z\}, \quad \text{for } z \in [\bar{z}, 0]; \quad (4.9)$$

$$\gamma_\beta^{-1}(x) = \inf\{y \in \mathbb{R}^+ \mid \gamma_\beta(y) \leq x\}, \quad \text{for } x \in \mathbb{R}^-; \quad (4.10)$$

$$\rho_\beta^{-1}(x) = \inf\{z \in [\bar{z}, 0] \mid \rho_\beta(z) \geq x\}, \quad \text{for } x \in \mathbb{R}^-. \quad (4.11)$$

Note that β and γ_β are càglàd, β^{-1} and ρ_β are càdlàg, and γ_β^{-1} as well as ρ_β^{-1} are continuous³. Moreover, β , β^{-1} and γ_β^{-1} are decreasing, γ_β is strictly decreasing, ρ_β is strictly increasing, and ρ_β^{-1} is increasing. Furthermore, it follows directly from the definitions of β^{-1} , γ_β , γ_β^{-1} , ρ_β and ρ_β^{-1} that the following three identities hold.

$$\rho_\beta^{-1}(x) = x + \gamma_\beta^{-1}(x), \quad \text{for all } x \in \mathbb{R}^-, \quad (4.12)$$

$$\gamma_\beta^{-1}(\rho_\beta(z)) = \beta^{-1}(z), \quad \text{for all } z \in [\bar{z}, 0], \quad (4.13)$$

$$\rho_\beta^{-1}(\gamma_\beta(y)) = \beta(y), \quad \text{for all } y \in \mathbb{R}^+. \quad (4.14)$$

Also, by the definitions of \mathcal{G}^β , β and β^{-1} , we see that the set \mathcal{G}^β is the union of the graphs of the functions β and β^{-1} restricted to \mathcal{D} .

Observe that if $z > \beta(y)$, then the strategy Y^β corresponding to the intervention boundary described by β consists of an initial sale of Δ number of shares so that $(y - \Delta, z - \Delta)$ is in \mathcal{G}^β (see Figure 1). Let $Y_{0-}^\beta = y$ and $Y_0^\beta = y - \Delta$. Suppose $(y - \Delta, z - \Delta)$ is on the graph of β . Then $(y - \Delta, z - \Delta) = (y - \Delta, \beta(y - \Delta))$ and this equality is equivalent to

$$\gamma_\beta(Y_0^\beta) = \beta(Y_0^\beta) - Y_0^\beta = z - y,$$

from which it follows that $Y_0^\beta = \gamma_\beta^{-1}(z - y)$ and $\Delta = y - \gamma_\beta^{-1}(z - y)$. Now suppose $(y - \Delta, z - \Delta)$ is on the graph of β^{-1} , and let $Z_{0-}^{Y^\beta} = z$ and $Z_0^{Y^\beta} = z - \Delta$. Then $(y - \Delta, z - \Delta) =$

³It can be checked that for $x \in \mathbb{R}^-$, $\gamma_\beta^{-1}(x)$ and $\rho_\beta^{-1}(x)$ is respectively the y -coordinate and the z -coordinate of the intersection of the line $z = y + x$ and \mathcal{G}^β .

$(\beta^{-1}(z - \Delta), z - \Delta)$, which is equivalent to

$$\rho_\beta(Z_0^{Y^\beta}) = Z_0^{Y^\beta} - \beta^{-1}(Z_0^{Y^\beta}) = z - y,$$

and it follows that $Z_0^{Y^\beta} = \rho_\beta^{-1}(z - y)$ and $\Delta = z - \rho_\beta^{-1}(z - y)$. According (4.12), the number Δ of shares in both of the aforementioned two cases can be expressed by

$$\Delta = y - \gamma_\beta^{-1}(z - y) = z - \rho_\beta^{-1}(z - y).$$

On the other hand, if $z \leq \beta(y)$, then the strategy Y^β consists of an initial waiting period until $(Y_t^\beta, Z_t^{Y^\beta})$ is on the graph of β (see Figure 1). As long as no action is taken, we have $Y_t^\beta = y$, and with reference to (2.8) and (2.9), we obtain $Z_t^{Y^\beta} = H^{-1}(H(z) - t)$. The first time t_w that the state process is on the graph of β is therefore given by

$$t_w = H(z) - H(\beta(y)). \quad (4.15)$$

Once the state process (Y^β, Z^{Y^β}) is in the set \mathcal{G}^β , the strategy Y^β consists of selling shares in such a way that the state process remains in \mathcal{G}^β (see Figure 1). Therefore, $(Y_t^\beta, Z_t^{Y^\beta}) = (Y_t^\beta, \beta(Y_t^\beta))$ whenever $\beta(Y_t^\beta +) = \beta(Y_t^\beta)$. With reference to (2.5), this implies that Y_t^β should solve

$$d\beta(Y_t^\beta) = -h(\beta(Y_t^\beta)) dt + dY_t^\beta,$$

which is equivalent to

$$d\gamma_\beta(Y_t^\beta) = -h(\beta(Y_t^\beta)) dt.$$

If $\beta^{-1}(Z_t^{Y^\beta}) = \beta^{-1}(Z_t^{Y^\beta} -)$, then $(Y_t^\beta, Z_t^{Y^\beta}) = (\beta^{-1}(Z_t^{Y^\beta}), Z_t^{Y^\beta})$. According to (2.5) and the definition of β^{-1} , Z^{Y^β} should solve

$$dZ_t^{Y^\beta} = -h(Z_{t-}^{Y^\beta}) dt.$$

Set

$$t_w = \begin{cases} 0, & \text{if } z > \beta(y), \\ H(z) - H(\beta(y)), & \text{if } z \leq \beta(y), \end{cases} \quad (4.16)$$

and

$$\bar{t} = \inf\{t \geq 0 \mid Y_t^\beta = 0\}. \quad (4.17)$$

Denote by $\{y_n\}_{n \in \mathbb{I}}$ the set of discontinuity points of β . Then \mathbb{I} is countable since β is càglàd. Define $\{t_n\}_{n \in \mathbb{I}}$ by

$$t_n = \inf\{t \geq t_w \mid Y_t^\beta = y_n\}, \quad (4.18)$$

and $\{s_n\}_{n \in \mathbb{I}}$ by

$$s_n = \inf\{t \geq t_w \mid Y_t^\beta < y_n\}. \quad (4.19)$$

If $\{t \geq t_w \mid Y_t^\beta = y_n\} = \emptyset$, set $t_n = \infty$; and set $s_n = \infty$, if $\{t \geq t_w \mid Y_t^\beta < y_n\} = \emptyset$. The following result establish existence and uniqueness of such a strategy Y^β corresponding to a given intervention boundary β .

Lemma 4.1. *Let $(y, z) \in \mathcal{D}$ where $y > 0$, and let β be an intervention boundary function. Let $H, \beta^{-1}, \gamma_\beta, \gamma_\beta^{-1}, t_w, \bar{t}, y_n, t_n$ and s_n be given by (2.7), (4.9), (4.7), (4.10) and (4.16)–(4.19), respectively. Set $(Y_t^\beta)_{t \geq 0} = (Y_{t \wedge \bar{t}}^\beta)_{t \geq 0}$, with $Y_{0-}^\beta = y$, which denotes the decreasing càdlàg liquidation strategy corresponding to β , and let $(Z_t^{Y^\beta})_{t \geq 0}$, with $Z_{0-}^{Y^\beta} = z$, be the state process of the bid order book associated with Y^β . Suppose Y^β satisfies the following description:*

(i) *If $z > \beta(y)$,*

- (a) *when $y \in \cup_{n \in \mathbb{I}}(z - \beta(y_n) + y_n, z - \beta(y_{n+}) + y_n]$, immediately sell $y - \gamma_\beta^{-1}(z - y)$ number of shares. This block trade ensures $Y_0^\beta = \beta^{-1}(Z_0^{Y^\beta})$.*
- (b) *when $y \in (z, \infty) \setminus \cup_{n \in \mathbb{I}}(z - \beta(y_n) + y_n, z - \beta(y_{n+}) + y_n]$, immediately sell $y - \gamma_\beta^{-1}(z - y)$ number of shares. This block trade ensures $Z_0^{Y^\beta} = \beta(Y_0^\beta)$.*

Then continuously sell shares so that $(Y_t^\beta, Z_t^{Y^\beta}) \in \mathcal{G}^\beta$ for all $t \in [t_w, \bar{t}]$.

- (ii) *If $z \leq \beta(y)$, then wait until time t_w . The time t_w has the property that $Z_{t_w}^{Y^\beta} = \beta(y)$. Continuously sell shares so that $(Y_t^\beta, Z_t^{Y^\beta}) \in \mathcal{G}^\beta$ for all $t \in [t_w, \bar{t}]$.*

Such a strategy Y^β exists and is unique, and it is continuous for all $t > 0$. In particular,

$$Y_t^\beta = y_n \quad \text{for } t \in [t_w, \bar{t}] \cap \cup_{n \in \mathbb{I}}[t_n, s_n), \quad (4.20)$$

with corresponding $Z_t^{Y^\beta}$ being the unique solution to

$$dZ_t^{Y^\beta} = -h(Z_t^{Y^\beta}) dt, \quad (4.21)$$

where

$$Z_{t_w}^{Y^\beta} = \rho_\beta^{-1}(z - y) \text{ if } z > \beta(y), \quad \text{and} \quad Z_{t_n}^{Y^\beta} = \beta(Y_{t_n-}^\beta) \text{ for } t_n > t_w. \quad (4.22)$$

Moreover,

$$Z_t^{Y^\beta} = \beta(Y_t^\beta), \quad \text{for } t \in [t_w, \bar{t}] \setminus \cup_{n \in \mathbb{I}} [t_n, s_n), \quad (4.23)$$

where Y^β is the unique solution to

$$d\gamma_\beta(Y_t^\beta) = -h(\beta(Y_t^\beta)) dt, \quad (4.24)$$

with

$$Y_{t_w}^\beta = y \text{ if } z \leq \beta(y), \quad Y_{t_w}^\beta = \gamma_\beta^{-1}(z - y) \text{ if } z > \beta(y), \quad \text{and} \quad Y_{s_n}^\beta = y_n \text{ for } s_n > t_w. \quad (4.25)$$

If $t_w > 0$, then $Y_t^\beta = y$ and $Z_t^{Y^\beta} = H^{-1}(H(z) - t)$, for $0 \leq t \leq t_w$.

We can also describe $Z_t^{Y^\beta}$ for $t \in [\bar{t}, \infty)$ since it satisfies (4.21) with initial condition

$$Z_{\bar{t}}^{Y^\beta} = \begin{cases} Z_{t_w}^{Y^\beta}, & \text{if } \bar{t} = t_w, \\ z, & \text{if } \bar{t} < t_w, \\ \beta(0+), & \text{if } \bar{t} > t_w. \end{cases} \quad (4.26)$$

The value $\beta(0+)$ can then be used to determine whether the liquidation period is finite or not. More specifically, we have that $\beta(0+) < 0$ implies $\bar{t} < \infty$. To see this, it is enough to consider

$$\gamma_\beta(Y_{\bar{t}}^\beta) - \gamma_\beta(Y_t^\beta) = \int_t^{\bar{t}} -h(\beta(Y_u^\beta)) du$$

which follows from (4.24) when there is no waiting period between the times t and \bar{t} . To get a contradiction, suppose $\bar{t} = \infty$. Then it is clear that $\int_t^{\bar{t}} -h(\beta(Y_u^\beta)) du = \infty$, as $\beta(Y_u^\beta)$ is

bounded away from 0 on the interval (t, \bar{t}) . However, $\gamma_\beta(Y_{\bar{t}}^\beta) - \gamma_\beta(Y_t^\beta)$ is finite, so we get a contradiction.

It follows from the dynamics of $Z_t^{Y^\beta}$ that Z^{Y^β} is càdlàg and increasing to 0. Moreover, the continuity of Y_t^β for $t > 0$ implies that Z^{Y^β} is also continuous for all $t > 0$.

We now progress by deriving an explicit expression for the performance function associated with the strategy Y^β described by Lemma 4.1 for an arbitrary intervention boundary β . This expression can then later be used to derive an explicit expression for the value function of our problem. For the strategy Y^β with associated state process Z^{Y^β} , given an initial state (y, z) , and with reference to (3.8), we define the performance function J_β by

$$J_\beta(y, z) = \int_0^\infty \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt, \quad (4.27)$$

where $Y_{0-}^\beta = y$, $Z_{0-}^{Y^\beta} = z$ and $(y, z) \in \mathcal{D}$. Since $\kappa_A(0) = 0$, it follows that

$$\int_{\bar{t}}^\infty \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt = A \int_0^{Z_{\bar{t}}^{Y^\beta}} \psi(u) du. \quad (4.28)$$

Therefore,

$$J_\beta(0, z) = A \int_0^z \psi(u) du. \quad (4.29)$$

Lemma 4.2. *Let β , Y^β , Z^{Y^β} , t_w and \bar{t} be defined as in Lemma 4.1. If $t_w < \bar{t}$, then*

$$\begin{aligned} & \int_{t_w}^\infty \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt \\ &= \int_{\beta(0+)}^{Z_{t_w}^{Y^\beta} - Y_{t_w}^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du, \end{aligned}$$

where γ_β^{-1} and ρ_β^{-1} are defined by (4.10) and (4.11), respectively.

In case (i) (a) of Lemma 4.1, the strategy Y^β consists of an initial sale of $y - \gamma_\beta^{-1}(z - y) = z - \rho_\beta^{-1}(z - y)$ number of shares. The state after the block sale is $(Y_0^\beta, Z_0^{Y^\beta}) = (\beta^{-1}(\rho_\beta^{-1}(z -$

$y)) , \rho_\beta^{-1}(z - y))$, Hence, according to (4.27) and Lemma 4.2,

$$\begin{aligned} J_\beta(y, z) &= J_\beta\left(\beta^{-1}(\rho_\beta^{-1}(z - y)), \rho_\beta^{-1}(z - y)\right) \\ &= \int_{\beta(0+)}^{\rho_\beta(\rho_\beta^{-1}(z-y))} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du \\ &= \int_{\beta(0+)}^{z-y} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du. \end{aligned}$$

In case (i) (b), we immediately sell $y - \gamma_\beta^{-1}(z - y)$ number of shares at the beginning. The state after the block sale is $(Y_0^\beta, Z_0^{Y^\beta}) = (\gamma_\beta^{-1}(z - y), \beta(\gamma_\beta^{-1}(z - y)))$. Hence, similar to the above calculation, we have

$$\begin{aligned} J_\beta(y, z) &= J_\beta\left(\gamma_\beta^{-1}(z - y), \beta(\gamma_\beta^{-1}(z - y))\right) \\ &= \int_{\beta(0+)}^{z-y} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du. \end{aligned}$$

We therefore conclude that for case (i) of Lemma 4.1,

$$J_\beta(y, z) = \int_{\beta(0+)}^{z-y} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du. \quad (4.30)$$

Moreover, in case (ii), $z \leq \beta(y)$. So we need to wait until time $t_w > 0$ at which $Z_{t_w}^{Y^\beta} = \beta(y)$. With reference to (2.7) and (4.15), we have

$$t_w = H(z) - H(\beta(y)) = \int_{\beta(y)}^z \frac{1}{h(u)} du.$$

Also, observe that

$$\int_0^{t_w} h(Z_t^{Y^\beta}) \psi(Z_t^{Y^\beta}) dt = - \int_0^{t_w} \psi(Z_t^{Y^\beta}) dZ_t^{Y^\beta} = - \int_z^{\beta(y)} \psi(u) du.$$

Hence in case (ii), the performance function is given by

$$\begin{aligned}
J_\beta(y, z) &= \int_0^{t_w} \left(\kappa_A(y) + Ah(Z_t^{Y^\beta}) \psi(Z_t^{Y^\beta}) \right) dt + J_\beta(y, \beta(y)) \\
&= \kappa_A(y) \int_{\beta(y)}^z \frac{1}{h(u)} du - A \int_z^{\beta(y)} \psi(u) du \\
&\quad + \int_{\beta(0+)}^{\gamma_\beta(y)} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du. \quad (4.31)
\end{aligned}$$

Although this provides an explicit expression for $J_\beta(y, z)$, it is not entirely straightforward to conclude about properties of continuity and differentiability for $J_\beta(y, z)$ in y since β is only a càglàd function. However, we can further calculate that

$$\begin{aligned}
\int_{\beta(0+)}^{\gamma_\beta(y)} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du &= \int_0^y \left(\frac{\kappa_A(u)}{h(\beta(u))} + A\psi(\beta(u)) \right) d\gamma_\beta^c(u) \\
&\quad + \sum_{0 < u < y} \kappa_A(u) \int_{\beta(u)}^{\beta(u+)} \frac{1}{h(x)} dx \\
&\quad + A \sum_{0 < u < y} \int_{\beta(u)}^{\beta(u+)} \psi(s) ds.
\end{aligned}$$

From this expression, as well as

$$\int_{\beta(0+)}^{\beta(y)} \psi(u) du = \int_0^y \psi(\beta(u)) d\beta^c(u) + \sum_{0 < u < y} \int_{\beta(u)}^{\beta(u+)} \psi(s) ds,$$

and

$$\begin{aligned}
\kappa_A(y)H(\beta(y)) &= \kappa_A(0)H(\beta(0+)) + \int_0^y \kappa'_A(u)H(\beta(u)) du + \int_0^y \frac{\kappa_A(u)}{h(\beta(u))} d\beta^c(u) \\
&\quad + \sum_{0 < u < y} \kappa_A(u) \int_{\beta(u)}^{\beta(u+)} \frac{1}{h(x)} dx,
\end{aligned}$$

it follows from (4.31) that the performance function $J_\beta(y, z)$, for case (ii) of Lemma 4.1, admits

the expression

$$J_\beta(y, z) = \kappa_A(y)H(z) + A \int_0^z \psi(u) du - \int_0^y \left(\frac{\kappa_A(u)}{h(\beta(u))} + A\psi(\beta(u)) + \kappa'_A(u)H(\beta(u)) \right) du. \quad (4.32)$$

In the above calculations, we have assumed the existence and finiteness of $\lim_{u \rightarrow 0+} \frac{\kappa_A(u)}{h(\beta(u))}$ and $\lim_{u \rightarrow 0+} \kappa'_A(u)H(\beta(u))$. We have also used that $\lim_{u \rightarrow y-} \kappa_A(u) < \infty$ as well as $\lim_{u \rightarrow y-} \kappa'_A(u) < \infty$. The finiteness of $\lim_{u \rightarrow 0+} \kappa'_A(u)H(\beta(u))$ together with (3.3) and (3.4) imply that $\kappa_A(0)H(\beta(0+)) = 0$. For an optimal intervention boundary β , all of these properties will be demonstrated below in Lemma 4.6.

Suppose β is an intervention boundary such that Y^β is optimal, i.e. the value function v and the performance function J_β coincide. Then according to the Hamilton-Jacobi-Bellman equation as well as (4.32), we should have

$$D_y^- v(y, z) + v_z(y, z) = \Gamma(z; y) - \Gamma(\beta(y); y) \leq 0, \quad \text{for all } (y, z) \in \mathcal{D},$$

where

$$\Gamma(x; y) = A\psi(x) + \frac{\kappa_A(y)}{h(x)} + \kappa'_A(y)H(x).$$

Therefore, for any given y , $\beta(y)$ should be a maximiser of $\Gamma(x; y)$. The next lemma helps us characterise an intervention boundary β whose value maximises $\Gamma(x; y)$. We will later show that such a β indeed describes an optimal intervention boundary for our problem.

Lemma 4.3. *For $y \in (0, \bar{y}_A)$, define the function $\Gamma(\cdot; y) : [\bar{z}, 0] \rightarrow \overline{\mathbb{R}}$ by*

$$\Gamma(x; y) = A\psi(x) + \frac{\kappa_A(y)}{h(x)} + \kappa'_A(y)H(x), \quad \text{for } x \in (\bar{z}, 0), \quad (4.33)$$

and

$$\Gamma(0; y) = \lim_{x \rightarrow 0+} \Gamma(x; y), \quad \Gamma(\bar{z}; y) = \lim_{x \rightarrow \bar{z}} \Gamma(x; y).$$

Let $\beta^* = \beta^*(y)$ and $\beta_* = \beta_*(y)$ denote the functions defined as the largest and smallest

$\beta \in [\bar{z}, 0]$ satisfying

$$\max_{x \in [\bar{z}, 0]} \Gamma(x; y) = \Gamma(\beta; y), \quad (4.34)$$

respectively. Then for all $y \in (0, \bar{y}_A)$, we have $\bar{z} \leq \beta_*(y) \leq \beta^*(y) < 0$. Furthermore, if $\bar{y}_A < \infty$, write $\beta^*(y) = \beta_*(y) = \bar{z}$, for all $y > \bar{y}_A$. Set

$$\beta^*(0) = 0, \quad \beta_*(0) = \lim_{y \rightarrow 0^+} \beta_*(y),$$

and

$$\beta^*(\bar{y}_A) = \lim_{y \rightarrow \bar{y}_A^-} \beta^*(y), \quad \beta_*(\bar{y}_A) = \lim_{y \rightarrow \bar{y}_A^+} \beta_*(y).$$

This uniquely defines two decreasing functions $\beta^*, \beta_* : \mathbb{R}^+ \rightarrow [\bar{z}, 0]$ that are càglàd and càdlàg, respectively, and they are the left and the right-continuous version of each other.

Remark 4.4. We will see later that the optimal intervention boundary will be given by β^* , and we have previously assumed that $\mu \leq 0$. But let's examine some of the properties that β^* would have if $\mu > 0$. If $\mu > 0$, then κ_A is no longer strictly increasing, but will be strictly decreasing on $[0, \underline{y})$ and then strictly increasing on $(\underline{y}, \bar{y}_A)$, where \underline{y} denotes the unique point where κ_A attains its minimum. In particular $\kappa_A(\underline{y}) < 0$ and $\kappa'_A(\underline{y}) = 0$. Therefore

$$\Gamma(x; \underline{y}) = A\psi(x) + \frac{\kappa_A(\underline{y})}{h(x)},$$

which is maximised for $x = 0$. Thus $\beta^*(\underline{y}) = \beta_*(\underline{y}) = 0$ if $\mu > 0$. This means that if the optimal intervention boundary is given by β^* , then the optimal strategy would involve holding on to at least \underline{y} number of shares until eternity if the initial stock position is larger than \underline{y} . But such a strategy is not admissible as we want to restrict the set of admissible strategies to strategies which tend to zero sufficiently fast (we would also have to rephrase the admissibility condition if $\mu > 0$ since in this case κ_A is not strictly increasing). We conclude that if $\mu > 0$, then the optimal liquidation problem is not well posed.

Lemma 4.5. Let $\beta = \beta^*$ where β^* is as in Lemma 4.3. For $\bar{z} - \bar{y}_A < s < 0$ and $\rho_\beta^{-1}(s) < z < 0$

it holds that

$$\Gamma(\rho_\beta^{-1}(s); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) \geq 0.$$

Lemma 4.5 is needed for the proof of Proposition 4.7 (see in particular equation (5.20)). The result relies on the assumptions that $x \mapsto \nu([x, 0])$ and $x \mapsto 1/h(x)$ are concave functions. If we do not impose these conditions then one can show that the strategy Y^β with $\beta = \beta^*$ given by (4.34) may not be optimal (compare Theorem 4.8). Thus while these concavity conditions may seem realistic, they are also needed in order to solve the problem. If these conditions does not hold, we simply do not know what the solution to the liquidation problem looks like.

Lemma 4.6. *Let β^* be given by Lemma 4.3, it follows that if $\lim_{x \rightarrow y-} \kappa_A(x) = \infty$ or $\lim_{x \rightarrow y-} \kappa'_A(x) = \infty$, then $\lim_{x \rightarrow y-} \beta^*(x) = \bar{z}$. Furthermore, we have*

$$\lim_{y \rightarrow 0+} \frac{\kappa_A(y)}{h(\beta^*(y))} = 0 \quad \text{and} \quad \lim_{y \rightarrow 0+} \kappa'_A(y)H(\beta^*(y)) = 0. \quad (4.35)$$

Clearly, the function β^* given in Lemma 4.3 satisfies the properties we require of an intervention boundary. With this intervention boundary, the proposition below provides an explicit expression for the value function that solves (4.1)–(4.4) with associated boundary condition $v(0, z) = A \int_0^z \psi(u) du$, for all $z \in [\bar{z}, 0]$. As a consequence, the optimal liquidation strategy is characterised by this intervention boundary.

Proposition 4.7. *Let $\beta = \beta^*$ denote the largest solution to (4.34), and let γ_β^{-1} and ρ_β^{-1} be the corresponding functions defined by (4.10) and (4.11). Then the function $v : \mathcal{D} \rightarrow \mathbb{R}$ given by*

$$v(y, z) = \int_{\beta(0+)}^{z-y} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du + A \int_0^{\beta(0+)} \psi(u) du \quad (4.36)$$

for $z > \beta(y)$, and

$$v(y, z) = \kappa_A(y)H(z) + A \int_0^z \psi(u) du - \int_0^y \left(\frac{\kappa_A(u)}{h(\beta(u))} + A\psi(\beta(u)) + \kappa'_A(u)H(\beta(u)) \right) du, \quad (4.37)$$

for $z \leq \beta(y)$, is a $C^{0,1}(\mathcal{D})$ solution to (4.1)–(4.4) with the boundary condition

$$v(0, z) = A \int_0^z \psi(u) du, \quad \text{for all } z \in [\bar{z}, 0].$$

Moreover, $D_y^- v(y, z)$ is càglàd in y and continuous in z .

The following theorem verifies that the function v given by (4.36)–(4.37) is equal to the value function V given by (3.8), and that the strategy Y^β corresponding to $\beta = \beta^*$ in Lemma 4.3 is an optimal liquidation strategy. Hence, such a Y^β provides a solution to the utility maximization problem in (2.11).

Theorem 4.8. *Let β denote the largest solution to (4.34), let v be given by (4.36) and (4.37), and let V be given by (3.8). Then $v = V$ on \mathcal{D} and*

$$\sup_{Y \in \mathcal{A}(y)} \mathbb{E}[U(C_\infty(Y))] = -\exp\left(-A(c + by) + A \int_z^{z-y} \psi(s) ds\right) \exp(v(y, z)),$$

where A denotes the agent's risk aversion, b is the initial unaffected price, c is the agent's initial cash position, $z = Z_{0-}^Y$ is the initial state of the bid order book, and y is the agent's initial share position. The optimal strategy Y^* is equal to $Y^\beta \in \mathcal{A}_D(y)$, where Y^β is the strategy described in Lemma 4.1 corresponding to β with $Y_{0-}^\beta = y$.

Example 1. Suppose that the bid order book has an equal number n of orders at every price point to a level \bar{x} below the unaffected best bid price. This corresponds to $x \mapsto m([x, 0]) = -n(x \wedge \bar{x})$. Therefore $\psi(z) = \frac{1}{n}z$, for $\bar{z} = n\bar{x} < z \leq 0$, and $-\infty$, for $z \leq \bar{z}$. We calculate that

$$\Gamma'(x; y) = \frac{A}{n} - \frac{\kappa_A(y)}{\lambda x^2} + \frac{\kappa'_A(y)}{\lambda x}, \quad \text{for } n\bar{x} < x < 0.$$

In order to find the maximiser, we want to solve $\Gamma'(x; y) = 0$. This amounts to solving

$$\frac{\lambda A}{n} x^2 + \kappa'_A(y) x - \kappa_A(y) = 0. \tag{4.38}$$

The unique solution is

$$x = \frac{n}{2\lambda A} \left\{ -\kappa'_A(y) - \sqrt{(\kappa'_A(y))^2 + \frac{4\lambda A}{n} \kappa_A(y)} \right\}.$$

Since $\Gamma(x; y)$ is concave in x for every $0 \leq y < \bar{y}_A$ and equation (4.38) has a unique solution for $x \leq 0$, it follows that

$$\beta(y) = \frac{n}{2\lambda A} \left\{ -\kappa'_A(y) - \sqrt{(\kappa'_A(y))^2 + \frac{4\lambda A}{n} \kappa_A(y)} \right\}, \quad \text{for } 0 \leq y < \bar{y}_A, \quad (4.39)$$

provided that $\beta(y) > n\bar{x}$.

- (i) **(BM case)** If L is a Brownian motion with drift $\mu \leq 0$, then

$$\kappa_A(y) = -A\mu y + \frac{1}{2}A^2\sigma^2 y^2, \quad y \geq 0,$$

and hence,

$$\beta(y) = \frac{n}{2\lambda A} \left\{ \mu A - \sigma^2 A^2 y - \sqrt{\mu^2 A^2 - \left(\frac{4\lambda \mu A^2}{n} - 2\mu \sigma^2 A^3 \right) y + \left(\sigma^4 A^4 + \frac{2\lambda \sigma^2 A^3}{n} \right) y^2} \right\}.$$

If $\mu = 0$, then it is straightforward to check that the equation for the optimal intervention boundary coincides with the result obtained in the example in Løkka (2014).

- (ii) **(LVG case)** Consider a Lévy process L with drift $\mu \leq 0$ and a pure jump part with Lévy-measure

$$\nu(dx) = \begin{cases} \frac{-1}{\eta \ln(x+1)} (x+1)^{C+D-1} dx, & x \in (-1, 0), \\ \frac{1}{\eta \ln(x+1)} (x+1)^{C-D-1} dx, & x \in (0, \infty). \end{cases} \quad (4.40)$$

where

$$C = \frac{\theta}{\rho^2} \quad \text{and} \quad D = \frac{\sqrt{\theta^2 + \frac{2\rho^2}{\eta}}}{\rho^2}. \quad (4.41)$$

If the initial stock price is 1 (if the initial price is $s \neq 1$, one can deal with this case by replacing the risk aversion A with $\tilde{A} = sA$) and

$$\mu = -\frac{1}{\eta} \ln\left(1 - \frac{\rho^2 \eta}{2} - \theta \eta\right), \quad (4.42)$$

then this choice of Lévy process corresponds to the (linear) Lévy process approximation of the exponential variance-gamma process with parameters (ρ, η, θ) . We refer the reader

to Løkka and Xu (2020) for more details. The common Lévy processes considered as models for financial price data assume models of the exponential type, and taking L to simply be one of these Lévy processes will in general not do a particularly good job as a model for financial price data. We then have

$$\begin{aligned}\kappa_A(y) &= -A\mu y + \int_{(-1,\infty)\setminus\{0\}} \left(e^{-Ayz} - 1 + Ayz \right) \nu(dz), \\ \kappa'_A(y) &= -A\mu + A \int_{(-1,\infty)\setminus\{0\}} \left(1 - e^{-Ayz} \right) z \nu(dz),\end{aligned}$$

where ν is given by (4.40). With reference to (4.39) we can calculate the optimal intervention boundary β .

From Figure 2 we see that the difference between the BM case and the LVG case increase with the agent's share position. Moreover the difference becomes more pronounced the more risk averse the agent is. If the agent is not particularly risk averse, then the more heavy-tailed risk in the LVG model does not make much difference to the agent, but if the agent is risk averse then it does. However, from Figure 2 we see that while for $A = 10^{-2}$ the optimal intervention boundary for the LVG case and the BM case are quite different, in terms of the corresponding strategy Y^β , the initial block sale in the LVG case is about 200 shares and in the BM case about 150 shares, if the initial stock position is 10^4 . An initial block sale of 200 shares would eat into the order book and reduce the best bid price by $\frac{200}{1000}$, so the best bid immediately after the block sale would be 0.8 (remember that we assumed the initial stock price to be 1). In the case of an initial block sale of 150 shares the best bid would be $1 - \frac{150}{1000} = 0.85$ immediately after the block sale.




Figure 2: The optimal intervention boundary β for the two cases of L being a Brownian motion (BM case) and L being the Lévy process approximating the exponential variance-gamma (LVG case). We take $n = 1000$ and $\lambda = 5$. For the LVG case, we take $\theta = -0.002$, $\rho = 0.02$ and $\eta = 0.6$ (timescale in days), which according to Rathgeber et al. (2013) are realistic parameters for stock price data for the exponential variance-gamma model. We choose $\mu = -0.0018$, which is in accordance with (4.42). For the BM case, we also choose $\mu = -0.0018$, and take $\sigma^2 = 4.011 * 10^{-4}$, which match the variance of L in the LVG case. Solid graphs correspond to the LVG case and a dotted graphs to the BM case. When $A = 10^{-3}$ we see that the LVG and the BM graphs are pretty much identical.

5 Proofs

Proof of Lemma 3.1. With reference to the dynamic of Z^Y , we calculate that for $z \geq \bar{z}$,

$$\begin{aligned}
\int_0^{Z_T^Y} \psi(u) du &= \int_0^z \psi(u) du + \int_0^T \psi(Z_{t-}^Y) dY_t^c \\
&\quad - \int_0^T h(Z_{t-}^Y) \psi(Z_{t-}^Y) dt + \sum_{0 \leq t \leq T} \int_{Z_{t-}^Y}^{Z_{t-}^Y + \Delta Y_t} \psi(u) du \\
&= \int_0^z \psi(u) du + \int_0^T \psi(Z_{t-}^Y) dY_t^c \\
&\quad - \int_0^T h(Z_{t-}^Y) \psi(Z_{t-}^Y) dt + \sum_{0 \leq t \leq T} \int_0^{\Delta Y_t} \psi(Z_{t-}^Y + u) du.
\end{aligned}$$

Then,

$$\begin{aligned}
F_T(Y) &= \int_0^T \psi(Z_{t-}^Y) dY_t^c + \sum_{0 \leq t \leq T} \int_0^{\Delta Y_t} \psi(Z_{t-}^Y + x) dx \\
&= \int_0^{Z_T^Y} \psi(u) du - \int_0^z \psi(u) du + \int_0^T h(Z_{t-}^Y) \psi(Z_{t-}^Y) dt \\
&= \int_z^{Z_T^Y} \psi(u) du + \int_0^T h(Z_{t-}^Y) \psi(Z_{t-}^Y) dt.
\end{aligned}$$

Notice that for any admissible liquidation strategy Y , we have that either Y and Z^Y become 0 at the same time or Y becomes 0 at some time s while $Z_s^Y < 0$. In the second case, for all $t > s$, Z^Y satisfies

$$dZ_t^Y = -h(Z_t^Y) dt.$$

According to (2.8), we know that the solution to the above dynamic tends to 0, as $t \rightarrow \infty$. Therefore, $Z_t^Y \rightarrow 0$, as $t \rightarrow \infty$ in any case. It then follows from the above expression for $F_T(Y)$ that

$$F_\infty(Y) = \int_z^0 \psi(u) du + \int_0^\infty h(Z_{t-}^Y) \psi(Z_{t-}^Y) dt.$$

□

Proof of Lemma 4.1. We first prove that on any time interval I contained in $[t_w, \bar{t}] \setminus \cup_{n \in \mathbb{I}} [t_n, s_n)$, there exists a unique solution to the dynamic (4.24). On such an interval I , the process Y^β does not cross any jump of β . Thus, in terms of the function β , we shall only focus on those parts without jumps. Also, it is sufficient to consider Y starting from time 0 (rather than starting at any time in $[t_w, \bar{t}] \setminus \cup_{n \in \mathbb{I}} [t_n, s_n)$). Write $Y_t^0 = Y_0 > 0$ and

$$Y_t^{k+1} = \gamma_\beta^{-1} \left(\left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^k)) du \right\} \wedge \beta(0+) \right). \quad (5.1)$$

Let $T \in [0, \infty)$. Then

$$\begin{aligned} & \sup_{0 \leq t \leq T} |\beta(Y_t^{k+1}) - \beta(Y_t^k)| \\ &= \sup_{0 \leq t \leq T} \left| \left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^k)) du \right\} \wedge \beta(0+) - \left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^{k-1})) du \right\} \wedge \beta(0+) \right. \\ & \quad \left. + \gamma_\beta^{-1} \left(\left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^k)) du \right\} \wedge \beta(0+) \right) \right. \\ & \quad \left. - \gamma_\beta^{-1} \left(\left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^{k-1})) du \right\} \wedge \beta(0+) \right) \right| \\ &\leq 2 \sup_{0 \leq t \leq T} \left| \left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^k)) du \right\} \wedge \beta(0+) - \left\{ \gamma_\beta(Y_0) - \int_0^t h(\beta(Y_u^{k-1})) du \right\} \wedge \beta(0+) \right| \\ &\leq 2 \sup_{0 \leq t \leq T} \left| \int_0^t h(\beta(Y_u^k)) - h(\beta(Y_u^{k-1})) du \right| \\ &\leq 2L \int_0^T |\beta(Y_u^k) - \beta(Y_u^{k-1})| du \\ &\leq 2L \int_0^T \sup_{0 \leq t \leq u} |\beta(Y_t^k) - \beta(Y_t^{k-1})| du, \end{aligned} \quad (5.2)$$

where the first equality holds because when β has no jumps we have $\beta(\gamma_\beta^{-1}(x)) = x + \gamma_\beta^{-1}(x)$, the first inequality is due to the triangle inequality and $|\gamma_\beta^{-1}(x) - \gamma_\beta^{-1}(y)| \leq |x - y|$, and the third inequality follows from the boundedness of the processes $\beta(Y^k)$ and $\beta(Y^{k-1})$ and the local Lipschitz continuity of h with a Lipschitz constant L . By induction and with reference to (5.2), it can be shown that

$$\sup_{0 \leq t \leq T} |\beta(Y_t^{k+1}) - \beta(Y_t^k)| \leq \frac{(2LT)^k}{k!} 2|\beta(Y_0)|.$$

Taking k to infinity, we have that $\beta(Y_t^k)$ converges uniformly on $[0, T]$. Define $\beta_t = \lim_{k \rightarrow \infty} \beta(Y_t^k)$, for $t \in [0, T]$. Since $T \in [0, \infty)$ is arbitrary, it follows that $\beta_t = \lim_{k \rightarrow \infty} \beta(Y_t^k)$ for all $t \in [0, \infty)$. With reference to (5.1) and the dominated convergence theorem it follows that, for every $t \in [0, \infty)$, $(Y_t^k)_{k=0}^\infty$ is convergent. We define $Y_t^\beta = \lim_{k \rightarrow \infty} Y_t^k$. It can be checked that Y^β decreases to 0. Then since β is continuous, we obtain $\beta_t = \beta(Y_t^\beta)$, for all $t \in [0, \infty)$. Therefore, by sending k to infinity in (5.1), since we only consider Y_t^β before time \bar{t} , we have that

$$Y_t^\beta = \gamma_\beta^{-1} \left(\gamma_\beta(Y_0^\beta) - \int_0^t h(\beta(Y_u^\beta)) du \right), \quad \text{for } t \leq \bar{t}.$$

This proves the existence of a solution to the dynamic (4.24) on any time interval contained in $[t_w, \bar{t}] \setminus \cup_{n \in \mathbb{I}} [t_n, s_n)$. For uniqueness, let's assume that $Y^{(1)}$ and $Y^{(2)}$ satisfy (4.24), where $Y_t^{(1)} = Y_t^{(2)}$ for $0 \leq t \leq t_1$, and $Y_t^{(1)} < Y_t^{(2)}$ for $t_1 < t < t_2$. Then for $t_1 < t < t_2$,

$$\begin{aligned} Y_t^{(1)} &= \gamma_\beta^{-1} \left(\gamma_\beta(Y_0^{(1)}) - \int_0^t h(\beta(Y_u^{(1)})) du \right) \\ &\geq \gamma_\beta^{-1} \left(\gamma_\beta(Y_0^{(2)}) - \int_0^t h(\beta(Y_u^{(2)})) du \right) \\ &= Y_t^{(2)}, \end{aligned}$$

which contradicts the assumption that $Y_t^{(1)} < Y_t^{(2)}$ for $t_1 < t < t_2$. So uniqueness holds. The existence and uniqueness of a solution to the dynamic in (4.21) on any time interval contained in $[t_w, \bar{t}] \cap \cup_{n \in \mathbb{I}} [t_n, s_n)$ follow from the locally Lipschitz continuity of the function h .

Now let Y^β and Z^{Y^β} be processes satisfying (4.20)–(4.26) with $(Y_{0-}^\beta, Z_{0-}^{Y^\beta}) = (y, z) \in \mathcal{D}$. Note that $(Y_t^\beta, Z_t^{Y^\beta}) \in \mathcal{G}^\beta$ for all $t \in [t_w, \bar{t}]$. We need to show that (2.5) is satisfied. We first focus on the case when $t \leq t_w$. Suppose $z > \beta(y)$, i.e. $t_w = 0$. Then in case (i) (a),

$$\begin{aligned} Y_0^\beta - Y_{0-}^\beta &= \beta^{-1}(\rho_\beta^{-1}(z - y)) - y \\ &= \gamma_\beta^{-1}(z - y) - y \\ &= (z - y + \gamma_\beta^{-1}(z - y)) - z \\ &= Z_0^{Y^\beta} - Z_{0-}^{Y^\beta}, \end{aligned}$$

where we have used the identity $\beta^{-1}(\rho_\beta^{-1}(z - y)) = \gamma_\beta^{-1}(z - y)$, which follows from (4.13) and

is valid under the condition of (i) (a). In case (i) (b), we obtain

$$\begin{aligned}
Z_0^{Y^\beta} - Z_{0-}^{Y^\beta} &= \beta(\gamma_\beta^{-1}(z - y)) - z \\
&= z - y + \gamma_\beta^{-1}(z - y) - z \\
&= \gamma_\beta^{-1}(z - y) - y \\
&= Y_0^\beta - Y_{0-}^\beta,
\end{aligned}$$

where we have used that $\beta(\gamma_\beta^{-1}(z - y)) = \rho_\beta^{-1}(z - y)$. Suppose $z \leq \beta(y)$, i.e. $t_w > 0$. It can be checked that $Z_t^{Y^\beta} = H^{-1}(H(z) - t)$ has dynamic (4.21). Because Y_t^β is now constant, (2.5) is satisfied. In the case when $t > t_w$, Y_t^β and $Z_t^{Y^\beta}$ follow (4.20)–(4.26), which satisfy (2.5).

We next like to prove that Y^β is càdlàg and decreasing. Note that by the definitions of t_n , s_n , t_w and \bar{t} and (4.21), (4.24) and the first part of the proof, we have Y_t^β and $Z_t^{Y^\beta}$ are continuous when $(Y_t^\beta, Z_t^{Y^\beta})$ is in each continuous part of the graph of β or β^{-1} , for $t > 0$. Also, each initial condition associated with the dynamics (4.21) and (4.24) is chosen to make Y_t^β and $Z_t^{Y^\beta}$ to be continuous at t_n , s_n and t_w when $t_w > 0$. It can also be seen that Y^β and Z^{Y^β} are right continuous at $t = 0$. These together with the well-defined Y_{0-}^β and $Z_{0-}^{Y^\beta}$ imply that Y^β and Z^{Y^β} are continuous for $t > 0$ and they are right-continuous with left-limit at $t = 0$. That Y^β decreases to 0 follows from (4.20), (4.21), (4.24), and the first part of this proof. Finally, that $Z_t^{Y^\beta} = H^{-1}(H(z) - t)$, for $0 \leq t \leq t_w$, follows from (2.8). \square

Proof of Lemma 4.2. Let $\{y_n\}_{n \in \mathbb{I}}$ be the set of all points at which the intervention boundary β is discontinuous. Consider a time interval $[t, s] \subseteq [t_n, s_n)$, for some $n \in \mathbb{I}$, where t_n and s_n are given by (4.18) and (4.19). With reference to (2.5), we note that formally,

$$dt = -\frac{d\rho_\beta(Z_t^{Y^\beta})}{h(Z_t^{Y^\beta})} \quad \forall t \in [t_n, s_n),$$

and hence,

$$\begin{aligned}
& \int_t^s \left(\kappa_A(Y_r^\beta) + Ah(Z_r^{Y^\beta})\psi(Z_r^{Y^\beta}) \right) dr \\
&= \int_s^t \left(\frac{\kappa_A(\beta^{-1}(Z_r^{Y^\beta}))}{h(Z_r^{Y^\beta})} + A\psi(Z_r^{Y^\beta}) \right) d\rho_\beta(Z_r^{Y^\beta}) \\
&= \int_{\rho_\beta(Z_s^{Y^\beta})}^{\rho_\beta(Z_t^{Y^\beta})} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du \\
&= \int_{Z_s^{Y^\beta} - Y_{s_n}^\beta}^{Z_t^{Y^\beta} - Y_{t_n}^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du, \tag{5.3}
\end{aligned}$$

where we have used the identity in (4.13). Similarly, since

$$dt = -\frac{d\gamma_\beta(Y_t^\beta)}{h(\beta(Y_t^\beta))} \quad \forall t \in [t_w, \bar{t}] \setminus \cup_{n \in \mathbb{I}} [t_n, s_n),$$

applying (4.14), it can be calculated that on some time interval $[s, t] \subset [t_w, \bar{t}] \setminus \cup_{n \in \mathbb{I}} [t_n, s_n)$, for some $n \in \mathbb{I}$,

$$\begin{aligned}
& \int_s^t \left(\kappa_A(Y_r^\beta) + Ah(Z_r^{Y^\beta})\psi(Z_r^{Y^\beta}) \right) dr \\
&= \int_{Z_t^{Y^\beta} - Y_t^\beta}^{Z_s^{Y^\beta} - Y_s^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du. \tag{5.4}
\end{aligned}$$

Let $t_w < \bar{t}$. Suppose the number of t_n and s_n in the interval $[t_w, \bar{t}]$ is equal to $m < \infty$ (possibly $m = 0$). Consider $r_0 \leq r_1 < \dots < r_m < r_{m+1}$, where $r_0 = t_w$, $r_{m+1} = \bar{t}$ and for $k = 1, \dots, m$, r_k are equal to those $t_n, s_n \in [t_w, \bar{t}]$. We assume r_1, \dots, r_m are in an ascending order. Then it follows from (5.3), (5.4) and the continuity of Y_t^β and $Z_t^{Y^\beta}$ when $t > 0$ that

$$\begin{aligned}
& \int_{t_w}^{\bar{t}} \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt \\
&= \sum_{k=0}^m \int_{r_k}^{r_{k+1}} \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt \\
&= \int_{Z_t^{Y^\beta}}^{Z_{t_w}^{Y^\beta} - Y_{t_w}^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du.
\end{aligned}$$

Suppose there are infinitely many t_n and s_n in the interval $[t_w, \bar{t}]$. Let $r \in [t_w, \bar{t}]$ be an accumulation point of the sequence $\{t_n\}_{n \in \mathbb{I}}$. Then without loss of generality, consider a subsequence $\{t_{n_k}\}_{k=1}^\infty \subset [t_w, \bar{t}]$ increasing to r . Consider some time interval $[t, s]$ in which r is the only accumulation point of $\{t_n\}_{n \in \mathbb{I}}$. Then, it follows that

$$\begin{aligned}
& \int_t^s \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt \\
&= \lim_{n \rightarrow \infty} \int_t^{t_n} \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt + \int_r^s \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt \\
&= \lim_{n \rightarrow \infty} \int_{Z_{t_n}^{Y^\beta} - Y_{t_n}^\beta}^{Z_t^{Y^\beta} - Y_t^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du \\
&\quad + \int_{Z_s^{Y^\beta} - Y_s^\beta}^{Z_r^{Y^\beta} - Y_r^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du, \\
&= \int_{Z_s^{Y^\beta} - Y_s^\beta}^{Z_t^{Y^\beta} - Y_t^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du.
\end{aligned}$$

This implies that

$$\begin{aligned}
& \int_{t_w}^{\bar{t}} \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt \\
&= \int_{Z_{\bar{t}}^{Y^\beta} - Y_{\bar{t}}^\beta}^{Z_{t_w}^{Y^\beta} - Y_{t_w}^\beta} \left(\frac{\kappa_A(\gamma_\beta^{-1}(u))}{h(\rho_\beta^{-1}(u))} + A\psi(\rho_\beta^{-1}(u)) \right) du.
\end{aligned}$$

Therefore the result follows from the above equality as well as (4.26) and (4.28). \square

Proof of Lemma 4.3. First notice that, for any $y \in (0, \bar{y}_A)$, the function $\Gamma(x, y)$ is concave in x , but that this concavity may not be strict. Observe that for $y \in (0, \bar{y}_A)$,

$$\lim_{x \rightarrow 0^-} \Gamma(x; y) = -\infty.$$

Also, $\Gamma(x; y) \in \mathbb{R}$, for $x \in [\bar{z}, 0)$. These observations imply that $\bar{z} \leq \beta_*(y) \leq \beta^*(y) < 0$, for all $0 < y < \bar{y}_A$. The largest and smallest solution to (4.34) uniquely define the functions β^* and

β_* . For $0 < y < y + \Delta < \bar{y}_A$ and $x \in [\bar{z}, 0)$, we calculate that

$$\frac{d}{dx} \left[\Gamma(x; y + \Delta) - \Gamma(x; y) \right] = -\frac{(\kappa_A(y + \Delta) - \kappa_A(y))h'(x)}{h^2(x)} + \frac{\kappa'_A(y + \Delta) - \kappa'_A(y)}{h(x)} < 0, \quad (5.5)$$

since κ_A is convex and $\kappa'_A(u) > 0$, for $u > 0$. We want to show that β^* and β_* are decreasing functions. In order to get a contradiction, suppose that there exists $y \in (0, \bar{y}_A)$ and $\Delta > 0$ such that $\beta^*(y + \Delta) > \beta_*(y)$. With reference to (5.5), we obtain

$$\Gamma(\beta^*(y + \Delta); y + \Delta) - \Gamma(\beta^*(y + \Delta); y) < \Gamma(\beta_*(y); y + \Delta) - \Gamma(\beta_*(y); y).$$

However, this contradicts the definitions of β^* and β_* , which imply that

$$\Gamma(\beta^*(y + \Delta); y + \Delta) \geq \Gamma(\beta_*(y); y + \Delta) \quad \text{and} \quad \Gamma(\beta_*(y); y) \geq \Gamma(\beta^*(y + \Delta); y).$$

Therefore, for all $0 < y < \bar{y}_A$,

$$\beta_*(y + \Delta) \leq \beta^*(y + \Delta) \leq \beta_*(y) \leq \beta^*(y), \quad (5.6)$$

from which it follows that β^* and β_* are decreasing. By (4.33), we know that for $\bar{z} \leq x < 0$, $\Gamma(x; y)$ is continuous in y . Then for $y \in (0, \bar{y}_A)$, we have

$$\begin{aligned} \Gamma(\beta_*(y+); y+) &= \Gamma(\beta_*(y+); y) \leq \Gamma(\beta_*(y); y) = \Gamma(\beta_*(y); y+) \\ \Gamma(\beta^*(y-); y) &= \Gamma(\beta^*(y-); y-) \geq \Gamma(\beta^*(y); y-) = \Gamma(\beta^*(y); y). \end{aligned}$$

Since β^* and β_* are defined as respectively the largest and smallest maximiser to (4.34), and β^* and β_* are decreasing, it follows that $\beta_*(y+) = \beta_*(y)$ and $\beta^*(y-) = \beta^*(y)$. By monotonicity, the right limit of β^* and the left limit of β_* exist. Hence, we have proved that β^* is càglàd and β_* is càdlàg. The claim that β^* is the càglàd version of β_* and that β_* is the càdlàg version of β^* follows from (5.6). \square

Proof of Lemma 4.5. With reference to (4.14), we have that

$$\beta(\gamma_\beta^{-1}(s)) = \rho_\beta^{-1}(\gamma_\beta(\gamma_\beta^{-1}(s))) = \rho_\beta^{-1}(s) \quad \text{if } \gamma_\beta(\gamma_\beta^{-1}(s)) = s.$$

Moreover, $\gamma_\beta(\gamma_\beta^{-1}(s)) = s$, unless β has a jump at $\gamma_\beta^{-1}(s)$. Thus if β does not have a jump at

$\gamma_\beta^{-1}(s)$ then

$$\Gamma(\rho_\beta^{-1}(s); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) = \Gamma(\beta(\gamma_\beta^{-1}(s)); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) \geq 0, \quad (5.7)$$

by the definition of β . If on the other hand β has a jump at $\gamma_\beta^{-1}(s)$, then γ_β^{-1} is flat on the interval $[s_*, s^*]$, where $s_* = \beta(\gamma_\beta^{-1}(s)+) - \gamma_\beta^{-1}(s)$ and $s^* = \beta(\gamma_\beta^{-1}(s)) - \gamma_\beta^{-1}(s)$. Also

$$\beta(\gamma_\beta^{-1}(s_*)+) = \rho_\beta^{-1}(\gamma_\beta(\gamma_\beta^{-1}(s_*+))) = \rho_\beta^{-1}(s_*)$$

and

$$\beta(\gamma_\beta^{-1}(s^*)) = \rho_\beta^{-1}(\gamma_\beta(\gamma_\beta^{-1}(s^*))) = \rho_\beta^{-1}(s^*).$$

In particular,

$$\beta(\gamma_\beta^{-1}(s_*)+) = \rho_\beta^{-1}(s_*) \leq \rho_\beta^{-1}(s) \leq \rho_\beta^{-1}(s^*) = \beta(\gamma_\beta^{-1}(s^*)) \quad (5.8)$$

and so

$$\Gamma(\rho_\beta^{-1}(s^*), \gamma_\beta^{-1}(s)) = \Gamma(\rho_\beta^{-1}(s_*), \gamma_\beta^{-1}(s)) \geq \Gamma(z, \gamma_\beta^{-1}(s)),$$

by the definition of β . With reference to Assumption 2.5 and Assumption 2.6, it follows that $x \mapsto \Gamma(x; y)$ is concave. According to (5.8), there exists $\lambda \in [0, 1]$ such that $\rho_\beta^{-1}(s) = \lambda \rho_\beta^{-1}(s_*) + (1 - \lambda) \rho_\beta^{-1}(s^*)$. Hence,

$$\Gamma(\rho_\beta^{-1}(s); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) \geq \Gamma(\beta(\gamma_\beta^{-1}(s)); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) \geq 0. \quad (5.9)$$

□

Proof of Lemma 4.6. If $y > \bar{y}_A$, then by the definition of β^* , it holds that if $\lim_{x \rightarrow y-} \kappa_A(x) = \infty$ or $\lim_{x \rightarrow y-} \kappa'_A(x) = \infty$, then $\lim_{x \rightarrow y-} \beta^*(x) = \bar{z}$. The remaining case is when $y = \bar{y}_A$. We will prove this case by contradiction. Suppose $\beta^*(\bar{y}_A) > \bar{z}$. For any $x \in (\bar{z}, \beta^*(\bar{y}_A))$ and

$y \in (0, \bar{y}_A)$ such that $\beta^*(y) \geq \beta^*(\bar{y}_A)$, we have

$$\begin{aligned} A\psi(x) &\leq A\left(\psi(x) - \psi(\beta^*(y))\right) \\ &\leq \kappa_A(y)\left(\frac{1}{h(\beta^*(y))} - \frac{1}{h(x)}\right) + \kappa'_A(y)\left(H(\beta^*(y)) - H(x)\right) \\ &\leq \kappa_A(y)\left(\frac{1}{h(\beta^*(\bar{y}_A))} - \frac{1}{h(x)}\right) + \kappa'_A(y)\left(H(\beta^*(\bar{y}_A)) - H(x)\right). \end{aligned}$$

Taking y to be arbitrarily close to \bar{y}_A implies $\psi(x) = -\infty$. This means $x < \bar{z}$, which contradicts $x > \bar{z}$. Hence, we conclude that $\beta^*(\bar{y}_A) = -\infty$.

Next we prove (4.35). Observe that if $\beta^*(0+) < 0$, then (4.35) is true. However, if $\beta^*(0+) = 0$, then

$$\frac{\kappa_A(y)}{h(\beta^*(y))} \geq \Gamma(x; y) - A\psi(\beta^*(y)) - \kappa'_A(y)H(\beta^*(y)) \geq \Gamma(x; y) - \kappa'_A(y)H(\beta^*(y)),$$

from which it follows that for any $x \in (\bar{z}, 0)$,

$$0 \geq \liminf_{y \rightarrow 0+} \frac{\kappa_A(y)}{h(\beta^*(y))} \geq A\psi(x) - \limsup_{y \rightarrow 0+} \kappa'_A(y)H(\beta^*(y)), \quad (5.10)$$

$$0 \geq \limsup_{y \rightarrow 0+} \frac{\kappa_A(y)}{h(\beta^*(y))} \geq A\psi(x) - \liminf_{y \rightarrow 0+} \kappa'_A(y)H(\beta^*(y)). \quad (5.11)$$

Therefore,

$$\begin{aligned} 0 &\geq \limsup_{y \rightarrow 0+} \kappa'_A(y)H(\beta^*(y)) \geq A\psi(x), \\ 0 &\geq \liminf_{y \rightarrow 0+} \kappa'_A(y)H(\beta^*(y)) \geq A\psi(x). \end{aligned}$$

By letting x tend to 0, then with reference to (2.3), we get $\lim_{y \rightarrow 0+} \kappa'_A(y)H(\beta^*(y)) = 0$. Also, by letting x tend to 0 in (5.10) and (5.11), $\lim_{y \rightarrow 0+} \frac{\kappa_A(y)}{h(\beta^*(y))} = 0$ follows. \square

Proof of Proposition 4.7. To show that v is continuous, we first prove it is finite. With reference to (4.28)-(4.32), it is sufficient to show that the function J_β given by (4.27) is finite for β defined by Lemma 4.3. By the continuity of Y^β and Z^{Y^β} after time 0 and condition

(2.4), we have that there exists some $s > 0$ such that

$$\int_0^s \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt < \infty \quad (5.12)$$

and $Y_s^\beta < \bar{y}_A$. According to the condition in Lemma 4.6,

$$\lim_{y \rightarrow 0^+} \frac{\kappa_A(y)}{h(\beta(y))} = 0,$$

so it follows that there exists $C_1 > 0$ and $0 < \epsilon < \bar{y}_A$ such that

$$\kappa_A(y) \leq -C_1 h(\beta(y)), \quad \text{for all } y \in [0, \epsilon].$$

Since $\psi(Z_t^{Y^\beta})$ is bounded for all $t \geq s$ (it increases to 0), this and the above inequality imply that

$$\begin{aligned} \int_s^\infty \left(\kappa_A(Y_t^\beta) + Ah(Z_t^{Y^\beta})\psi(Z_t^{Y^\beta}) \right) dt &\leq \int_s^\infty \left(-C_1 h(\beta(Y_t^\beta)) - C_2 h(Z_t^{Y^\beta}) \right) dt \\ &\leq \int_s^\infty \left(-C_1 h(Z_t^{Y^\beta}) - C_2 h(Z_t^{Y^\beta}) \right) dt \\ &\leq (C_1 + C_2)(Y_s^\beta - Z_s^{Y^\beta}) < \infty, \end{aligned} \quad (5.13)$$

where $C_2 > 0$ is some constant. Therefore, (5.12) and (5.13) show that v is finite.

Note that both of the expressions (4.36) and (4.37) are continuous in y and z . It is therefore sufficient to prove that v is continuous across \mathcal{G}^β . Let $J_u(y, z)$ denote the expression for $v(y, z)$ given by (4.36), and let $J_l(y, z)$ denote the expression in (4.37). Suppose (y, z) is a point on the graph of β , i.e, $z = \beta(y)$. Consider a sequence of points $(y_n, z_n)_{n=1}^\infty$ contained in $\bar{\mathcal{S}}^\beta \setminus \mathcal{G}^\beta$, converging to (y, z) . With reference to (4.31) and (4.32), we calculate that

$$\lim_{n \rightarrow \infty} v(y_n, z_n) = J_u(y, \beta(y)) = J_l(y, \beta(y)) = v(y, \beta(y)). \quad (5.14)$$

If (y, z) lies on the graph of β^{-1} , i.e, $y = \beta^{-1}(z)$, then using the property that $\beta^{-1}(u) = \beta^{-1}(z)$, for $u \in (z, \beta(\beta^{-1}(z)))$, direct calculation results in (5.14). We therefore conclude

that v is a continuous function. Differentiating v gives

$$D_y^- v(y, z) = -\frac{\kappa_A(\gamma_\beta^{-1}(z - y))}{h(\rho_\beta^{-1}(z - y))} - A\psi(\rho_\beta^{-1}(z - y)), \quad z > \beta(y), \quad (5.15)$$

$$v_z(y, z) = \frac{\kappa_A(\gamma_\beta^{-1}(z - y))}{h(\rho_\beta^{-1}(z - y))} + A\psi(\rho_\beta^{-1}(z - y)), \quad z > \beta(y), \quad (5.16)$$

$$D_y^- v(y, z) = \kappa'_A(y)H(z) - \frac{\kappa_A(y)}{h(\beta(y))} - A\psi(\beta(y)) - \kappa'_A(y)H(\beta(y)), \quad z \leq \beta(y), \quad (5.17)$$

$$v_z(y, z) = \frac{\kappa_A(y)}{h(z)} + A\psi(z), \quad z \leq \beta(y). \quad (5.18)$$

These expressions are left-continuous with right limit in y and continuous in z (all of these expressions are continuous at $(0, 0)$, which is guaranteed by (4.35)). Also, we check that for any $(y_n, z_n)_{n=1}^\infty \subseteq \overline{\mathcal{G}}^\beta$, $(y, z) \in \mathcal{G}^\beta$ and $\lim_{n \rightarrow \infty} (y_n, z_n) = (y, z)$, we have $v_z(y_n, z_n) \rightarrow v_z(y, z)$, as $n \rightarrow \infty$. Further, $\lim_{z \rightarrow \beta(y)+} D_y^- v(y, z) = D_y^- v(y, \beta(y))$. Therefore, we conclude that $v_z(y, z)$ is continuous, and $D_y^- v(y, z)$ is càglàd in y and continuous in z .

Standard calculations show that v satisfies (4.1) and (4.3). When $z = 0$, (4.2) is clearly true. In order to verify (4.2) for $z \neq 0$, we compute that when $z > \beta(y)$,

$$\begin{aligned} & h(z)v_z(y, z) - \kappa_A(y) - Ah(z)\psi(z) \\ &= h(z) \left\{ \frac{\kappa_A(\gamma_\beta^{-1}(s))}{h(\rho_\beta^{-1}(s))} - \frac{\kappa_A(z - s)}{h(z)} + A\{\psi(\rho_\beta^{-1}(s)) - \psi(z)\} \right\}, \end{aligned} \quad (5.19)$$

where $s = z - y$. Observe that $h(\rho_\beta^{-1}(s)) = 0$ implies $y = 0$, but (4.1)–(4.4) are under the condition that $y > 0$. So $h(\rho_\beta^{-1}(s))$ is non-zero. By the definition of γ_β^{-1} , we must have $\gamma_\beta^{-1}(s) \in (0, \bar{y}_A)$ if $\beta(\bar{y}_A) = \bar{z}$, or $\gamma_\beta^{-1}(s) \in (0, \bar{y}_A]$ if $\beta(\bar{y}_A) > \bar{z}$. Then according to the limiting behaviour of β in Lemma 4.6, $\kappa_A(\gamma_\beta^{-1}(s))$ must be finite, and therefore also $\kappa'_A(\gamma_\beta^{-1}(s))$. However, $\kappa_A(z - s)$ may be infinite, but then it follows that (5.19) is negative. Otherwise, if $\kappa_A(y) < \infty$, write

$$G(s; z) = \frac{\kappa_A(\gamma_\beta^{-1}(s))}{h(\rho_\beta^{-1}(s))} - \frac{\kappa_A(z - s)}{h(z)} + A\{\psi(\rho_\beta^{-1}(s)) - \psi(z)\}.$$

Then in order to verify (4.2), it is sufficient to show that $G(s; z) \geq 0$, for all $\rho_\beta^{-1}(s) < z < 0$.

We calculate that $G(s; y)$ can be expressed by

$$\begin{aligned} G(s; z) &= \left[\Gamma(\rho_\beta^{-1}(s); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) \right] \\ &\quad - \kappa'_A(\gamma_\beta^{-1}(s)) \left[H(\rho_\beta^{-1}(s)) - H(z) \right] + \frac{1}{h(z)} \left[\kappa_A(\gamma_\beta^{-1}(s)) - \kappa_A(z - s) \right], \end{aligned} \quad (5.20)$$

According to Lemma 4.5,

$$\Gamma(\rho_\beta^{-1}(s); \gamma_\beta^{-1}(s)) - \Gamma(z; \gamma_\beta^{-1}(s)) \geq 0.$$

Furthermore, we calculate that

$$\begin{aligned} &\frac{1}{h(z)} \left[\kappa_A(\gamma_\beta^{-1}(s)) - \kappa_A(z - s) \right] - \kappa'_A(\gamma_\beta^{-1}(s)) \left[H(\rho_\beta^{-1}(s)) - H(z) \right] \\ &= \int_{\rho_\beta^{-1}(s)}^z \left(\frac{[\kappa_A(u - s) - \kappa_A(\rho_\beta^{-1}(s) - s)] h'(u)}{h^2(u)} + \frac{\kappa'_A(\rho_\beta^{-1}(s) - s) - \kappa'_A(u - s)}{h(u)} \right) du \\ &\geq 0. \end{aligned} \quad (5.21)$$

(4.2) then follows from (5.20)–(5.21). Moreover, from the definition of β we get

$$\begin{aligned} D_y^- v(y, z) + v_z(y, z) &= \kappa'_A(y) H(z) - \frac{\kappa_A(y)}{h(\beta(y))} + A\psi(z - y) - A\psi(\beta(y)) \\ &\quad - \kappa'_A(y) H(\beta(y)) + \frac{\kappa_A(y)}{h(z)} + A\psi(z) - A\psi(z - y) \\ &= \Gamma(z; y) - \Gamma(\beta(y); y) \\ &\leq 0, \end{aligned}$$

which verifies (4.4).

Finally, the expression in (4.36) satisfies the boundary condition since, for any $u \in [\beta(0+), z]$, we have $\gamma_\beta^{-1}(u) = 0$ and $\rho_\beta^{-1}(u) = u$. The expression in (4.37) clearly satisfies the boundary condition. \square

Proof of Theorem 4.8. Let δ be a positive-valued $C^\infty(\mathbb{R})$ function with support on $[0, 1]$ satisfying $\int_0^1 \delta(x) dx = 1$, and define a sequence of functions $\{\delta_n\}_{n=1}^\infty$ by

$$\delta_n(s) = n \delta(ns), \quad s \geq 0.$$

We mollify v to obtain a sequence of functions $\{v^{(n)}\}_{n=1}^\infty$ which are given by

$$v^{(n)}(y, z) = \int_0^1 v(y - s, z) \delta_n(s) ds.$$

(One may extend the lower bound of the domain of $v(\cdot, z)$ properly so that $v^{(n)}$ is well-defined at $y = 0$.) Then $v^{(n)} \in C^{1,1}(\mathcal{D})$, for all $n \in \mathbb{N}$, and

$$\begin{aligned} v(y, z) &= \lim_{n \rightarrow \infty} v^{(n)}(y, z), \\ v_z(y, z) &= \lim_{n \rightarrow \infty} v_z^{(n)}(y, z), \\ D_y^- v(y, z) &= \lim_{n \rightarrow \infty} v_y^{(n)}(y, z), \end{aligned}$$

where the last equality is due to $D_y^- v(y, z)$ being càglàd in y . Moreover, for every $(y_0, z_0) \in \mathcal{D}$ there exists a $K > 0$ such that on the set $\{(y, z) \in \mathcal{D} \mid z \geq y + z_0 - y_0\}$,

$$|v^{(n)}(y, z)| \leq K, \quad n \in \mathbb{N}, \quad (5.22)$$

$$|v_y^{(n)}(y, z)| \leq K, \quad n \in \mathbb{N}, \quad (5.23)$$

$$|v_z^{(n)}(y, z)| \leq K, \quad n \in \mathbb{N}. \quad (5.24)$$

(If Y is admissible and $(Y_{0-}, Z_{0-}^Y) = (y_0, z_0)$, then $(Y_t, Z_t^Y) \in \{(y, z) \in \mathcal{D} \mid z \geq y + z_0 - y_0\}$, for all $t \geq 0$.) By Itô's formula, we calculate that

$$\begin{aligned} &v^{(n)}(Y_T, Z_T^Y) + \int_0^T \left(\kappa_A(Y_{t-}) + Ah(Z_{t-}^Y) \psi(Z_{t-}^Y) \right) dt \\ &= v^{(n)}(y, z) + \int_0^T \left(v_y^{(n)}(Y_{t-}, Z_{t-}^Y) + v_z^{(n)}(Y_{t-}, Z_{t-}^Y) \right) dY_t^c \\ &\quad + \int_0^T \left(\kappa_A(Y_{t-}) + Ah(Z_{t-}^Y) \psi(Z_{t-}^Y) - v_z^{(n)}(Y_{t-}, Z_{t-}^Y) h(Z_{t-}^Y) \right) dt \\ &\quad + \sum_{0 \leq t \leq T} \left\{ v^{(n)}(Y_{t-} + \Delta Y_t, Z_{t-}^Y + \Delta Y_t) - v^{(n)}(Y_{t-}, Z_{t-}^Y) \right\}, \end{aligned} \quad (5.25)$$

for all $Y \in \mathcal{A}_D(y)$. Observe that for $t \geq 0$,

$$0 \leq - \int_0^t h(Z_u^Y) du = Z_t^Y - Y_t - Z_0^Y + Y_0 \leq y - z.$$

Then, with reference to (5.22)–(5.24), we have

$$\int_0^\infty \sup_{n \in \mathbb{N}} \left| v_z^{(n)}(Y_{t-}, Z_{t-}^Y) h(Z_{t-}^Y) \right| dt \leq K(y - z).$$

Similarly,

$$\int_0^\infty \sup_{n \in \mathbb{N}} \left| v_y^{(n)}(Y_{t-}, Z_{t-}^Y) + v_z^{(n)}(Y_{t-}, Z_{t-}^Y) \right| d(-Y_t^c) \leq 2Ky$$

and

$$\sum_{0 \leq t} \sup_{n \in \mathbb{N}} \left| v^{(n)}(Y_{t-} + \Delta Y_t, Z_{t-}^Y + \Delta Y_t) - v^{(n)}(Y_{t-}, Z_{t-}^Y) \right| \leq 2Ky.$$

Hence, by (5.25) and the boundary condition $v(0, z) = A \int_0^z \psi(u) du$, it follows from the dominated convergence theorem that for any $Y \in \mathcal{A}_D(y)$,

$$\begin{aligned} & \int_0^\infty \left(\kappa_A(Y_{t-}) + Ah(Z_{t-}^Y) \psi(Z_{t-}^Y) \right) dt \\ &= v(y, z) + \int_0^\infty \left(D_y^- v(Y_{t-}, Z_{t-}^Y) + v_z(Y_{t-}, Z_{t-}^Y) \right) dY_t^c \\ & \quad + \int_0^\infty \left(\kappa_A(Y_{t-}) + Ah(Z_{t-}^Y) \psi(Z_{t-}^Y) - v_z(Y_{t-}, Z_{t-}^Y) h(Z_{t-}^Y) \right) dt \\ & \quad + \sum_{t \geq 0} \left\{ v(Y_{t-} + \Delta Y_t, Z_{t-}^Y + \Delta Y_t) - v(Y_{t-}, Z_{t-}^Y) \right\}, \end{aligned} \tag{5.26}$$

as $n \rightarrow \infty$ and $T \rightarrow \infty$. According to Proposition 4.7, v satisfies (4.1)–(4.4), and therefore,

$$\int_0^\infty \left(\kappa_A(Y_{t-}) + Ah(Z_{t-}^Y) \psi(Z_{t-}^Y) \right) dt \geq v(y, z). \tag{5.27}$$

Hence, $V \geq v$.

From from (5.12)–(5.13), we know that with β being the largest solution to (4.34) and Y^β being the strategy described in Lemma 4.1 corresponding to β , Y^β is admissible, in particular (2.2) is satisfied. Therefore, with reference to (5.27), in order to complete the proof, we need to show that (5.27) holds with equality for Y^β . Observe that $\Delta Y^\beta < 0$ only if $t = 0$ and $z > \beta(y)$. But by (4.1) and Proposition 4.7, we have that $D_y^- v(y, z) + v_z(y, z) = 0$, for

$z > \beta(y)$. Therefore,

$$\sum_{t \geq 0} \left\{ v(Y_{t-}^{\beta} + \Delta Y_t^{\beta}, Z_{t-}^{Y^{\beta}} + \Delta Y_t^{\beta}) - v(Y_{t-}^{\beta}, Z_{t-}^{Y^{\beta}}) \right\} = 0.$$

For any $z \leq 0$, if $0 \leq t \leq t_w$, where t_w is defined by (4.16), then $d(Y_t^{\beta})^c = 0$, hence

$$\int_0^{t_w} \left(D_y^- v(Y_{t-}^{\beta}, Z_{t-}^{Y^{\beta}}) + v_z(Y_{t-}^{\beta}, Z_{t-}^{Y^{\beta}}) \right) d(Y_t^{\beta})^c = 0;$$

if $t > t_w$, then $(Y_t^{\beta}, Z_t^{Y^{\beta}}) \in \mathcal{G}^{\beta}$, which implies

$$\int_{t_w}^{\infty} \left(D_y^- v(Y_{t-}^{\beta}, Z_{t-}^{Y^{\beta}}) + v_z(Y_{t-}^{\beta}, Z_{t-}^{Y^{\beta}}) \right) d(Y_t^{\beta})^c = 0.$$

Finally we have

$$\int_0^{\infty} \left(\kappa_A(Y_{t-}^{\beta}) + Ah(Z_{t-}^{Y^{\beta}}) \psi(Z_{t-}^{Y^{\beta}}) - v_z(Y_{t-}^{\beta}, Z_{t-}^{Y^{\beta}}) h(Z_{t-}^{Y^{\beta}}) \right) dt = 0,$$

since the integrand is equal to 0, for all $(Y_t^{\beta}, Z_t^{Y^{\beta}}) \in \overline{\mathcal{W}}^{\beta}$, and the Lebesgue measure of the set of $t \geq 0$ for which $(Y_t^{\beta}, Z_t^{Y^{\beta}}) \in \overline{\mathcal{S}}_s^{\beta} \setminus \mathcal{G}^{\beta}$ is 0. With reference to (5.26), we therefore conclude that $v = V$ and that $Y^* = Y^{\beta} \in \mathcal{A}_D(y)$ is an admissible optimal liquidation strategy for the optimization problem (3.8), and the result follows from (3.6). \square

References

- Alfonsi, A., Fruth, A., and Schied, A. (2010). Optimal execution strategies in limit order books with general shape functions. *Quantitative Finance*, 10:143–157.
- Almgren, R. and Chriss, N. (2000). Optimal execution of portfolio transactions. *Risk*, 3(2):5–39.
- Barndorff-Nielsen, O. E. (1997). Normal inverse gaussian distribution and stochastic volatility modelling. *Scandinavian Journal of Statistics*, 24(1):1–13.
- Cartea, A., Jaimungal, S., and Penalva, J. (2015). *Algorithmic and High-Frequency Trading*. Cambridge University Press.

- Cont, R. and Tankov, P. (2004). *Financial modelling with jump processes*. Chapman & Hall/CRC.
- Eberlein, E. and Keller, U. (1995). Hyperbolic distributions in finance. *Bernoulli*, 1(3):281–299.
- Fleming, W. H. and Soner, H. (2006). *Controlled Markov Processes and Viscosity Solutions*. Springer.
- Forsyth, P. A., Kennedy, J. S., Tse, S. T., and Windcliff, H. (2012). Optimal trade execution: a mean-quadratic variation approach. *Journal of Economic Dynamics and Control*, 36(12):1971–1991.
- Gatheral, J. (2010). No-dynamic-arbitrage and market impact. *Quantitative Finance*, 10(7):749–759.
- Gatheral, J. and Schied, A. (2011). Optimal trade execution under geometric brownian motion in the Almgren and Chriss framework. *International Journal of Theoretical and Applied Finance*, 14:353–368.
- Guéant, O. (2016). *The Financial Mathematics of Market Liquidity*. Chapman and Hall.
- Kallsen, J. and Shiryaev, A. (2002). The cumulant process and Esscher’s change of measure. *Finance and Stochastics*, 6:397–428.
- Kissell, R. and Malamut, R. (2005). Understanding the profit and loss distribution of trading algorithms. *Institutional Investor*.
- Kyprianou, A. (2006). *Introductory lectures on fluctuations of Levy processes with applications*. Springer.
- Lehalle, C.-A. and Laruelle, S. (2013). *Market Microstructure in Practice*. World Scientific.
- Løkka, A. (2014). Optimal liquidation in a limit order book for a risk-averse investor. *Mathematical Finance*, 24(4):696–727.
- Løkka, A. and Xu, J. (2020). Optimal liquidation trajectories for the Almgren-Chriss model.
- Madan, D. and Seneta, E. (1990). The variance gamma (V.G.) model for share market returns. *The Journal of Business*, 63(4):511–524.

- Obizhaeva, A. and Wang, J. (2013). Optimal trading strategy and supply/demand dynamics. *Journal of Financial Markets*, 16:1–32.
- Predoiu, S., Shaikhet, G., and Shreve, S. (2011). Optimal execution in a general one-sided limit-order book. *SIAM Journal on Financial Mathematics*, 2:183–212.
- Rathgeber, A., Stadler, J., and Stöckl, S. (2013). Modeling share returns - an empirical study on the variance gamma model. *Proceedings of European Financial Management Association, Reading 2013*.
- Schied, A. and Schöneborn, T. (2009). Risk aversion and the dynamics of optimal liquidation strategies in illiquid markets. *Finance and Stochastics*, 13:181–204.
- Schied, A., Schöneborn, T., and Tehranchi, M. (2010). Optimal basket liquidation for CARA investors is deterministic. *Applied Mathematical Finance*, 17:471–489.