

# Feynman Propagator for a System of Interacting Scalar Particles in the Fokker Theory

Natalia Gorobey,<sup>1</sup> Alexander Lukyanenko,<sup>1,\*</sup> and A. V. Goltsev<sup>2</sup>

<sup>1</sup>*Peter the Great Saint Petersburg Polytechnic University,  
Polytekhnicheskaya 29, 195251, St. Petersburg, Russia*

<sup>2</sup>*Ioffe Physical- Technical Institute, Polytekhnicheskaya 26, 195251, St. Petersburg, Russia*

A generalized canonical formulation of the theory of the electromagnetic Fokker interaction for a system of two particles is proposed. The functional integral on the generalized phase space is defined as the initial one in quantum theory. After integration over the momenta, the measure of integration in the generalized configuration space of world particle lines is determined. The problem of dynamic interpretation of the Feynman propagator for a system of particles with independent proper time parameters is discussed. A modification of the propagator is proposed, in which the role of independent time parameters is taken by the time coordinates of the particles in Minkowski space.

## I. INTRODUCTION

Wheeler and Feynman [1, 2] formulated classical electrodynamics without the concept of electromagnetic field in terms of the Fokker principle of least action [3]. For the simplest system of two particles, the Fokker action has the form ( $c = 1$ ):

$$I_F = \frac{1}{2} \int_0^{S_1} ds_1 \left( \dot{x}_1^2 + m_1^2 \right) + \frac{1}{2} \int_0^{S_2} ds_2 \left( \dot{x}_2^2 + m_2^2 \right) + e_1 e_2 \int_0^{S_1} ds_1 \int_0^{S_2} ds_2 \delta(s_{12}^2) \dot{x}_1 x_2'. \quad (1)$$

Here, the world lines of particles are parametrized by the intrinsic time of each particle, the dot and dash denote the derivatives with respect to the corresponding time. For definiteness, we consider the scattering problem and assume that the boundary points of the particle trajectories lie in the asymptotic regions of free motion. The squares of the velocity vectors and their scalar product are implied in the Minkowski space with the signature  $(+, -, -, -)$ . The same applies to the squared distance between particles

$$s_{12}^2 = (x_1(s_1) - x_2(s_2))^2 \quad (2)$$

The Fokker action Eq.(1) contains two independent parameters of time, which excludes the usual Hamiltonian form of the theory and the use of the usual rules of canonical quantization. This was partly an incentive for the development of a new formulation of quantum mechanics in terms of the functional integral [4]. This formulation turned out to be useful, in particular, in the case of systems with a nonlocal in time interaction potential [5]. However, for the Fokker action Eq.(1), quantum theory in terms of the functional integral was not formulated. There are two reasons for this. The first is the uncertainty of the integration measure in the space of world particle lines. This issue is considered in this paper. In

the framework of the generalized canonical formulation of the Fokker theory, a representation of the Feynman propagator for a system of interacting particles will be obtained. The second reason is common to relativistic quantum mechanics: the Feynman propagator has no dynamic meaning, since the proper time of particles is not an observable parameter. This issue requires a separate consideration. For ordinary systems with a local interaction potential, there is a canonical representation, and it is the initial one in the definition of the functional integral in the phase space for the kernel of the evolution operator - the solution of the Schrödinger equation [6]. From it, after integration over the momenta, we obtain the Feynman integral in the configuration space. The canonical form of action Eq.(1) is also desirable for determining the functional integral. Despite the multi-temporal nature of dynamics in the Fokker theory, it admits a generalized canonical formulation in which the world lines of particles as a whole serve as canonical coordinates [7, 8].

In this paper, a variant of such a generalization is considered. Based on it, the functional integral on the generalized phase space is determined, which, after integration over the momenta, becomes the functional integral on the configuration space of world particle lines with a certain measure. The construction of the Fokker quantum theory is being completed by integrating particles in their own time. In the case of a single particle, the integral of the kernel of the evolution operator over its proper time is a representation of the Green function of the Dirac and Klein-Gordon operators, i.e. the Feynman propagator [9, 10]. The result of multiple Fock-Schwinger integration (FS) in the Fokker theory will also be called the Feynman propagator. However, the Feynman propagator itself does not describe the evolution of the system, but is a skeletal element of the diagram technique in perturbation theory [11]. To introduce the physical parameters of time and the dynamic interpretation of the Feynman propagator, a modification of the Fokker theory at the classical level is required, similar to that considered in the case of a single relativistic particle in [12].

The next section proposes a generalized canonical formalism for multitemporal dynamic theory and defines the

\* alex.lukyan@mail.ru

Feynman propagator in the Fokker theory. In the third section, a modification of the Feynman propagator is proposed, allowing its dynamic interpretation.

## II. GENERALIZED CANONICAL REPRESENTATION AND QUANTIZATION OF FOKKER THEORY

In the absence of the Lagrange function, we consider action Eq.(1) as the velocity  $\dot{x}_{1\mu}(s_1), \dot{x}'_{2\nu}(s_2)$  functional, which should now be considered as an infinite set of quantities, also numbered by the particle's own time, and we define canonical momenta as variational derivatives:

$$p_{1\mu}(s_1) = \frac{\delta I_F}{\delta \dot{x}_{1\mu}(s_1)} = \dot{x}_{1\mu}(s_1) + e_1 e_2 \int_0^{S_2} ds_2 \delta(s_{12}^2) \dot{x}'_{2\mu}(s_2), \quad (3)$$

$$p_{2\nu}(s_2) = \frac{\delta I_F}{\delta \dot{x}'_{2\nu}(s_2)} = \dot{x}'_{2\nu}(s_2) + e_1 e_2 \int_0^{S_1} ds_1 \delta(s_{12}^2) \dot{x}_{1\nu}(s_1). \quad (4)$$

Next, we act in a standard way. We introduce the generalized Hamiltonian functional  $H_F[p_1, x_1, p_2, x_2]$  by means of the generalized Legendre transform:

$$H_F[p_1, x_1, p_2, x_2] = \int_0^{S_1} ds_1 p_{1\mu}(s_1) \dot{x}_{1\mu}(s_1) + \int_0^{S_2} ds_2 p_{2\nu}(s_2) \dot{x}'_{2\nu}(s_2) - I_F[\dot{x}_1, x_1, \dot{x}'_2, x_2], \quad (5)$$

where velocities should also be excluded by solving Eqs.(3) and (4). The solution can be written in the form of series of perturbation theory:

$$\dot{x}_{1\mu}(s_1) = p_{1\mu}(s_1) - e_1 e_2 \int_0^{S_2} ds_2 \delta(s_{12}^2) p_{2\mu}(s_2) + \dots, \quad (6)$$

$$\dot{x}'_{2\nu}(s_2) = p_{2\nu}(s_2) - e_1 e_2 \int_0^{S_1} ds_1 \delta(s_{12}^2) p_{1\nu}(s_1) + \dots \quad (7)$$

Substituting Eqs. (6) and (7) into Eq.(5), we obtain the generalized Hamilton functional in the form of a quadratic form of generalized momenta:

$$H_F[p_1, x_1, p_2, x_2] = \sum_{\alpha, \beta=1,2} \int_0^{S_\alpha} ds_\alpha \int_0^{S_\beta} ds_\beta M_{\alpha\mu\beta\nu}(s_\alpha, s_\beta) \times p_{\alpha\mu}(s_\alpha) p_{\beta\nu}(s_\beta) - m_1^2 S_1 - m_2^2 S_2, \quad (8)$$

where  $M_{\alpha\mu\beta\nu}(s_\alpha, s_\beta)$  are the matrix entries that are series composed of multiple integrals of the products  $\delta(s_{\alpha\beta}^2)$ . They are explicitly contained in the equations for the generalized canonical momenta Eqs. (6) and (7) that follow from the generalized canonical form of the Fokker action,

$$\begin{aligned} \tilde{I}_F[p_1, x_1, p_2, x_2] &= \int_0^{S_1} ds_1 p_{1\mu}(s_1) \dot{x}_{1\mu}(s_1) \\ &+ \int_0^{S_2} ds_2 p_{2\nu}(s_2) \dot{x}'_{2\nu}(s_2) \\ &- H_F[p_1, x_1, p_2, x_2]. \end{aligned} \quad (9)$$

It is in this form the Fokker action is taken as the basis for constructing the functional integral on the generalized phase space, which determines the evolution of the system in terms of the particle's proper time:

$$\begin{aligned} K(S_1, S_2) &= \int \frac{dp_{1out}}{2\pi\hbar} \prod_{s_{1,\mu}} \frac{dp_{1\mu} dx_{1\mu}}{2\pi\hbar} \frac{dp_{2out}}{2\pi\hbar} \prod_{s_{2,\nu}} \frac{dp_{2\nu} dx_{2\nu}}{2\pi\hbar} \\ &\times \exp\left\{\frac{i}{\hbar} \tilde{I}_F[p_1, x_1, p_2, x_2]\right\}. \end{aligned} \quad (10)$$

Gaussian momentum integrals are easily calculated, and we arrive at a functional integral on the configuration space with the initial Fokker action:

$$\begin{aligned} K(S_1, S_2) &= \int \prod_{s_{1,\mu}} dx_{1\mu} \prod_{s_{2,\nu}} dx_{2\nu} \frac{1}{\sqrt{\det M}} \\ &\times \exp\left\{\frac{i}{\hbar} I_F[\dot{x}_1, x_1, \dot{x}'_2, x_2]\right\}. \end{aligned} \quad (11)$$

Thus, a measure of functional integration in the configuration space of particle trajectories is defined. This integral can be made convenient for calculations in the framework of perturbation theory by introducing the determinant of the inverse matrix  $A = M^{-1}$ ,

$$\sqrt{\det A} = \frac{1}{\sqrt{\det M}}, \quad (12)$$

and take into account that it is determined by the coefficients of Eqs.(3) and (4). Then, for the determinant, we obtain the integral representation using Grassmann variables [6]:

$$\begin{aligned} \det A &= \int \prod_{s_{1,\mu}} \frac{d\chi_{1\mu}^* d\chi_{1\mu}}{2\pi} \prod_{s_{2,\nu}} \frac{d\chi_{2\nu}^* d\chi_{2\nu}}{2\pi} \\ &\times \exp\left[\int_0^{S_1} ds_1 \chi_{1\mu}^* \chi_{1\mu} + \int_0^{S_2} ds_2 \chi_{2\nu}^* \chi_{2\nu}\right] \\ &+ e_1 e_2 \int_0^{S_1} ds_1 \int_0^{S_2} ds_2 \delta(s_{12}^2) \\ &\times (\chi_{1\mu}^* \chi_{2\nu} + \chi_{2\nu}^* \chi_{1\mu}). \end{aligned} \quad (13)$$

The functional integral Eq. (11) is a function of the boundary points of the world lines of particles at extreme times:  $K(S_1, x_{1out}, 0, x_{1in}, S_2, x_{2out}, 0, x_{2in})$ . Now

it is necessary to make the next step that will complicate the dynamic interpretation of the Fokker quantum theory. Since the internal time of particles in quantum theory is considered unobservable, we obtain the covariant Feynman propagator by integrating the FS along these parameters:

$$K(x_{1out}, x_{1in}; x_{2out}, x_{2in}) = \int_0^\infty dS_1 \int_0^\infty dS_2 K(S_1, S_2). \quad (14)$$

In the absence of interaction, propagator Eq.(14) is reduced to the product of ordinary Feynman propagators of scalar particles.

### III. MODIFIED FEYNMAN PROPAGATOR OF FOKKER THEORY

Having thus determined the covariant Feynman propagator in the Fokker quantum theory, one should think about its dynamic interpretation. This will require its modification. According to [12], the modification of the action is carried out at the classical level: its variation, generated by the infinitesimal shift of the proper time in the coordinates  $x_{10}(s_1), x_{20}(s_2)$  of Minkowski space,

$$\delta x_{10} = \dot{x}_{10}\varepsilon_1, \delta x_{20} = \dot{x}_{20}\varepsilon_2. \quad (15)$$

is added to it. One can show that, by analogy with [12], the modified Feynman propagator for interacting charges in the Fokker theory has the form:

$$\begin{aligned} & \tilde{K}(P_{1out}, x_{1out}, P_{1in}, x_{1in}; P_{2out}, x_{2out}, P_{2in}, x_{2in}) \\ &= \int_0^\infty dS_1 \int_0^\infty dS_2 \int \prod_{s_1, k} dx_{1k} \prod_{s_2, l} dx_{2l} \prod_{s_1} dP_1 \prod_{s_2} dP_2 \\ & \times \sqrt{\det \tilde{A}} \prod_{s_1} \delta(x_{10} - \sqrt{2P_1}) \prod_{s_2} \delta(x_{20} - \sqrt{2P_2}) \\ & \times \prod_{s_1} \delta\left(\dot{P}_1 + \sqrt{2P_1} \dot{x}_{1k} F_{1k}\right) \\ & \times \prod_{s_2} \delta\left(P_2' - \sqrt{2P_2} \dot{x}_{2l} F_{2l}\right) \\ & \times \exp\left\{\frac{i}{\hbar} \tilde{I}_F[x_1, x_1, P_1, x_2', x_2, P_2]\right\}. \quad (16) \end{aligned}$$

Here  $P_1, P_2$  are the self-energies of the particles, and

$$\begin{aligned} F_{1k} &= 2e_1 e_2 \int_0^\infty dS_2 \frac{d}{ds_{12}^2} \delta(s_{12}^2) \\ & \times [(x_{10} - x_{20}) \dot{x}'_{2k} - (x_{1k} - x_{2k}) \dot{x}'_{20}], \quad (17) \end{aligned}$$

$$\begin{aligned} F_{2l} &= 2e_1 e_2 \int_0^\infty dS_1 \frac{d}{ds_{12}^2} \delta(s_{12}^2) \\ & \times [(x_{10} - x_{20}) \dot{x}_{1l} - (x_{1k} - x_{2k}) \dot{x}_{10}] \quad (18) \end{aligned}$$

are forces acting on each particle,

$$\begin{aligned} & \tilde{I}_F[x_1, x_1, P_1, x_2', x_2, P_2] \\ &= \frac{1}{2} \int_0^{S_1} ds_1 \left(2P_1 - \dot{x}_{1k}^2 + m_1^2\right) \\ & + \frac{1}{2} \int_0^{S_2} ds_2 \left(2P_2 - \dot{x}_{2l}^2 + m_2^2\right) \\ & + e_1 e_2 \int_0^{S_1} ds_1 \int_0^{S_2} ds_2 \delta(s_{12}^2) \end{aligned}$$

is a modified Fokker action, and

$$\begin{aligned} \det \tilde{A} &= \int \prod_{s_1, k} \frac{d\chi_{1k}^* d\chi_{1k}}{2\pi} \prod_{s_2, l} \frac{d\chi_{2l}^* d\chi_{2l}}{2\pi} \\ & \times \exp\left[\int_0^{S_1} ds_1 \chi_{1k}^* \dot{\chi}_{1k} + \int_0^{S_2} ds_2 \chi_{2l}^* \dot{\chi}_{2l}\right] \\ & + e_1 e_2 \int_0^{S_1} ds_1 \int_0^{S_2} ds_2 \delta(s_{12}^2) \\ & \times (\chi_{1k}^* \chi_{2k} + \chi_{2k}^* \chi_{1k}) \quad (20) \end{aligned}$$

is a measure of integration in the space of particle trajectories. Here we turned to three-dimensional notation.

The presence of  $\delta$ -functions in the functional integral Eq. (16) makes it possible to remove the FS integration over the particle proper time after solving the classical equations of motion for  $(x_{10}, P_1)$  and  $(x_{20}, P_2)$ . In this case, the role of time for each particle, as one would expect, is assumed by their  $x_{10}, x_{20}$  - coordinates in Minkowski space, and the self-energies of particles  $P_1, P_2$  are additional observables.

### IV. CONCLUSIONS

Relativistic quantum mechanics of a long-range interaction particle system is formulated in terms of a functional integral on the generalized phase space of world lines parameterized by the particle's proper time. The covariance of quantum theory is ensured by integration over the parameters of proper time in the Feynman propagator. A dynamic interpretation of the covariant quantum theory can be achieved by a modification of Fokker theory at the classical level. This modification does not change the dynamic content of the classical theory, but allows to remove the integration over proper time of particles in the Feynman propagator. As a result, the role of independent time parameters is taken by the  $x_{10}, x_{20}$  coordinates of the particles in Minkowski space.

### V. ACKNOWLEDGEMENTS

We thank V.A. Franke for useful discussions.

- 
- [1] R. P. Feynman and J. A. Wheeler, *Rev. Mod. Phys.* **17**, 157 (1945).
- [2] R. P. Feynman and J. A. Wheeler, *Rev. Mod. Phys.* **21**, 425 (1949).
- [3] A. D. Fokker, *Z. Phys.* **58**, 386 (1929).
- [4] R. P. Feynman, *The Development of the Space-Time View of Quantum Electrodynamics*, Nobel Lecture, December 11, 1965. Preprint les Prix Nobel en 1965. The Nobel Foundation. Stockholm, 1966.
- [5] R.P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw-Hill, New-York, (1965).
- [6] L. D. Faddeev and A. A. Slavnov, *Gauge fields: An introduction to quantum theory*, Westview Press, 2d edition, 236 (1993).
- [7] X. Jaen, R. Jauregui, J. Llosa, and A. Molina, *Phys. Rev. D* **36**, 2385 (1987).
- [8] X. Jaen, R. Jauregui, a) J. Llosa, and A. Molina, *J. Math. Phys.* **30** (12) 2807-2814 (1989).
- [9] V.A. Fock, *Izv. Acad. Nauk SSSR*, 551 (1937).
- [10] J. Schwinger, *On gauge invariance and vacuum polarization*, *Phys. Rev.* **82**. N 5, 664–679 (1951).
- [11] James D. Bjorken, Sidney D. Drell, *Relativistic quantum fields*, Mc Graw-Hill Book Company (1978).
- [12] Natalia Gorobey<sup>1</sup> and Alexander Lukyanenko, arXiv: 2001.09003v1 [gr-qc] 22 Jan 2020.