

A PROBABILISTIC PROOF OF APRIORI L^p ESTIMATES FOR A CLASS OF DIVERGENCE FORM ELLIPTIC OPERATORS

TYMOTEUZ CHOJECKI AND TOMASZ KOMOROWSKI

ABSTRACT. Suppose that \mathcal{L} is a divergence form differential operator of the form $\mathcal{L}f := (1/2)e^U \nabla_x \cdot [e^{-U}(I + H)\nabla_x f]$, where U is scalar valued, I identity matrix and H an anti-symmetric matrix valued function. The coefficients are not assumed to be bounded, but are C^2 regular. We show that if $Z = \int_{\mathbb{R}^d} e^{-U(x)} dx < +\infty$ and the supremum of the numerical range of matrix $-\frac{1}{2}\nabla_x^2 U + \frac{1}{2}\nabla_x \{ \nabla_x \cdot H - [\nabla_x U]^T H \}$ satisfies some exponential integrability condition with respect to measure $d\mu = Z^{-1}e^{-U} dx$, then for any $1 \leq p < q < +\infty$ there exists a constant $C > 0$ such that $\|f\|_{W^{2,p}(\mu)} \leq C \left(\|\mathcal{L}f\|_{L^q(\mu)} + \|f\|_{L^q(\mu)} \right)$ for $f \in C_0^\infty(\mathbb{R}^d)$. Here $W^{2,p}(\mu)$ is the Sobolev space of functions that are $L^p(\mu)$ integrable with two derivatives. Our proof is probabilistic and relies on an application of the Malliavin calculus.

1. INTRODUCTION

In the present note we formulate some apriori bound for a divergence form differential operator of the form

$$(1.1) \quad \begin{aligned} \mathcal{L}f(x) &:= \frac{1}{2}e^{U(x)} \nabla_x \cdot \left[e^{-U(x)}(I + H(x))\nabla_x f(x) \right] \\ &= \sum_{i,j=1}^d \frac{1}{2}e^{U(x)} \partial_{x_i} \left\{ e^{-U(x)} [(\delta_{i,j} + H_{i,j}(x)) \partial_{x_j} f(x)] \right\}, \quad f \in C_0^\infty(\mathbb{R}^d), x \in \mathbb{R}^d. \end{aligned}$$

Here $U : \mathbb{R}^d \rightarrow \mathbb{R}$ and $H(x) = [H_{i,j}(x)]$ is a $d \times d$ antisymmetric matrix valued function, i.e.

$$(1.2) \quad H_{i,j}(x) = -H_{j,i}(x), \quad x \in \mathbb{R}^d, i, j = 1, \dots, d.$$

We do not assume that the potential $U(x)$ and antisymmetric part $H(x)$ are bounded. Instead we assume that

$$(1.3) \quad Z := \int_{\mathbb{R}^d} e^{-U(x)} dx < +\infty.$$

and furthermore they satisfy some exponential integrability condition with respect to the measure

$$(1.4) \quad \mu(dx) := \frac{1}{Z} e^{-U(x)} dx,$$

see (2.8) below.

In our main result, see Theorem 2.1 and Corollary 2.2 below, we show that for any $q > p > 1$, there exists constant $C > 0$, depending only on p, q , dimension d and exponential moments of the coefficients, with respect to μ , such that

$$(1.5) \quad \|f\|_{W^{m,p}(\mu)} \leq C(\|f\|_{L^q(\mu)} + \|\mathcal{L}f\|_{L^q(\mu)}), \quad f \in C_0^\infty(\mathbb{R}^d).$$

Here, for any integer $m \geq 1$ and $p \in [1, +\infty)$, we denote by $W^{m,p}(\mu)$ the respective Sobolev space, defined as the completion of $C_0^\infty(\mathbb{R}^d)$ in the norm

$$(1.6) \quad \|f\|_{W^{m,p}(\mu)} := \left\{ \|f\|_{L^p(\mu)}^p + \sum_{k=1}^m \|\nabla_x^k f\|_{L^p(\mu)}^p \right\}^{1/p}, \quad f \in C_0^\infty(\mathbb{R}^d).$$

These type of estimates have been used by the authors in proving homogenization of advection equations with non-stationary, locally ergodic coefficients, see [2].

A priori estimates for Poisson equation in case of Sobolev norms with respect to the flat Lebesgue measure, and the Laplacian, or a uniform elliptic operator with bounded coefficients, are classical, see e.g. [3]. For \mathcal{L} that is an Ornstein-Uhlenbeck operator (then U and H are quadratic polynomials, and μ is therefore Gaussian) one can in fact assume that $p = q$ in (1.5), see [6]. This result has been generalised in [7] to cover some generalisation of Ornstein-Uhlenbeck operators, quite different from the one considered in the present paper. Some further results concerning global L^p bounds for divergence form operators with lower order coefficients in an appropriate Morrey class can be found in [8].

Our proof of (1.5) is probabilistic and relies on an application of the Malliavin calculus. The gradient of function f can be represented using the Fréchet derivative of the stochastic flow corresponding to the diffusion with the generator \mathcal{L} , see (3.15) and (3.16) below. The principal novelty of the paper is to replace the Fréchet by the Malliavin derivative in the direction of a random path that is adapted with respect to the natural filtration of the flow and such that, after some initial boundary layer, the respective Fréchet and Malliavin derivatives coincide, see (3.26) and (3.28) below. Finding such a path requires to solve a simple linear control problem, see Section 3.3. The solution also yields an estimate of the moments of the random path, see (3.27). We mention here that an analogous argument has been used by D. Bell in [1] in the context of finding a representation of a certain class of vector fields, defined over the space of paths on a Riemannian manifold, in terms of an appropriately defined divergence operator of the field.

The organisation of the paper is as follows: the basic definitions and assumptions are stated in Section 2, where also the main results of the paper are rigorously stated, see Section 2.2. The proofs of the results are given throughout Section 3. Finally, in the appendix we give the proofs of the auxiliary results, mainly from the theory of diffusions that are used throughout the paper.

Acknowledgements. Both authors acknowledge the support of the National Science Centre: NCN grant 2016/23/B/ST1/00492. T.K. wishes to express his gratitude to Professors D. Bakry and K. Oleszkiewicz for useful discussions on the topic of the paper.

2. PRELIMINARIES AND FORMULATION OF THE MAIN RESULT

We assume that the coefficients of \mathcal{L} are sufficiently smooth, so there is no issue with the definition of the stochastic flow $(X(t, x))_{t \geq 0, x \in \mathbb{R}^d}$ and semigroup $(P_t)_{t \geq 0}$ corresponding to the operator, see Section 3.1 below.

2.1. Differential operator.

A1) We assume that $U \in C^2(\mathbb{R}^d)$ and $H = [H_{i,j}]$ is a $d \times d$ matrix valued function with C^2 smooth entries that satisfy (1.2).

We can rewrite \mathcal{L} in the following way

$$(2.1) \quad \mathcal{L}f = Lf + \mathcal{A}f, \quad f \in C_0^\infty(\mathbb{R}^d),$$

where operator L is defined as

$$(2.2) \quad Lf := \frac{1}{2}e^U \nabla_x \cdot (e^{-U} \nabla_x f) = \frac{1}{2} \Delta_x f - \frac{1}{2} \nabla_x U \cdot \nabla_x f, \quad f \in C_0^\infty(\mathbb{R}^d)$$

and

$$(2.3) \quad \mathcal{A}f := \frac{1}{2} b \cdot \nabla_x f, \quad f \in C_0^\infty(\mathbb{R}^d).$$

where $b^T = [b_1, \dots, b_d]$, is given by

$$(2.4) \quad b := \nabla_x \cdot H - [\nabla_x U]^T H,$$

or coordinatewise

$$(2.5) \quad b_j(x) := \sum_{i=1}^d (\partial_{x_i} H_{i,j}(x) - \partial_{x_i} U(x) H_{i,j}(x)), \quad x \in \mathbb{R}^d, \quad j = 1, \dots, d.$$

2.2. Main results. To ensure global existence of the stochastic flow corresponding to \mathcal{L} , see Section 3.1 below, we assume that:

A2) there exists $\rho > 0$ such that

$$(2.6) \quad \int_{\mathbb{R}^d} |x|^{2\rho} \mu(dx) + \int_{\mathbb{R}^d} (|x|^2 + 1)^{\rho-1} |x \cdot [-\nabla_x U(x) + b(x)]| \mu(dx) < +\infty.$$

Let \mathcal{U} be the matrix valued function, given by

$$(2.7) \quad \mathcal{U} := -\frac{1}{2} \nabla_x^2 U + \frac{1}{2} \nabla_x b = -\frac{1}{2} \nabla_x^2 U + \frac{1}{2} \nabla_x \{ \nabla_x \cdot H - [\nabla_x U]^T H \},$$

or entrywise $\mathcal{U} = [\mathcal{U}_{j,j'}]$, with

$$\mathcal{U}_{j,j'} := -\frac{1}{2} \partial_{x_j, x_{j'}}^2 U + \frac{1}{2} \sum_{i=1}^d \partial_{x_{j'}} \{ \partial_{x_i} H_{i,j} - H_{i,j} \partial_{x_i} U \}, \quad j, j' = 1, \dots, d.$$

We let

$$\mathbf{u} := \max_{|\ell|=1} \langle \mathcal{U} \ell, \ell \rangle_{\mathbb{R}^d}.$$

A3) Integrability condition. Assume that there exists $\gamma_0 > 0$ such that

$$(2.8) \quad E(\gamma_0) := \int_{\mathbb{R}^d} \exp \{ \gamma_0 \mathbf{u} \vee 0 \} d\mu < +\infty.$$

Here $a \vee b := \max\{a, b\}$.

Our main result gives an estimate of the $L^p(\mu)$ norm of the gradient of $f \in C_0^\infty(\mathbb{R}^d)$ in terms of the $L^q(\mu)$ norms of f and $\mathcal{L}f$, with $q > p$.

Theorem 2.1. *Suppose that (1.3) and assumptions A1) – A3) hold. Then, for any $p \in [1, +\infty)$ and $q > p$ there exists $C(d, r, \gamma_0) > 0$, depending only on the dimension d , γ_0 and r , given by $1/p = 1/q + 1/r$, for which*

$$(2.9) \quad \|\nabla_x f\|_{L^p(\mu)} \leq C(d, r, E(\gamma_0)) \left(\|\mathcal{L}f\|_{L^q(\mu)} + \|f\|_{L^q(\mu)} \right), \quad f \in C_0^\infty(\mathbb{R}^d).$$

The proof of this theorem is contained in Section 3.2.

Using an argument based on Calderon-Zygmund estimates we can also conclude the following.

Corollary 2.2. *Suppose that all the hypotheses of Theorem 2.1 remain in force. In addition, we assume that*

$$(2.10) \quad \ell_{2,*} := \sum_{j,j'=1}^d \|H_{j,j'}\|_{W^{1,r}(\mu)} + \|U\|_{W^{2,r}(\mu)} < +\infty, \quad \forall r \in [1, +\infty).$$

Then, for any $q > 1$ and $p \in [1, q)$ there exists $C(d, p, q, E(\gamma_0), \ell_{2,}) > 0$, depending only on the indicated parameters, such that*

$$(2.11) \quad \|\nabla_x^2 f\|_{L^p(\mu)} \leq C \left(\|\mathcal{L}f\|_{L^q(\mu)} + \|f\|_{L^q(\mu)} \right), \quad f \in C_0^\infty(\mathbb{R}^d).$$

The proof of the corollary is presented in Section 3.4.

3. PROOFS OF THE MAIN RESULTS

3.1. Diffusion corresponding to \mathcal{L} . It follows from the assumptions made about the coefficients of \mathcal{L} that they are locally Lipschitz, therefore we can define a diffusion $(X(t; x))_{0 \leq t < \tilde{\epsilon}_x}$ via an Itô stochastic differential equation

$$(3.1) \quad dX(t; x) = \left\{ -\frac{1}{2} \nabla_x U(X(t; x)) + \frac{1}{2} b(X(t; x)) \right\} dt + dw(t), \quad 0 \leq t < \tilde{\epsilon}_x \quad X(0; x) = x,$$

where $(w(t) = (w_1(t), \dots, w_d(t)))_{t \geq 0}$ is standard d -dimensional Brownian motion and $\tilde{\epsilon}_x$ is an explosion time, see (3.7) below. The solutions satisfy the flow property, i.e.

$$X(t; X(s, x)) = X(t + s; x), \quad t + s < \tilde{\epsilon}_x, \quad t, s \geq 0, \quad x \in \mathbb{R}^d$$

for a.s. realization of the field $(X(t; x))_{0 \leq t < \tilde{\epsilon}_x, x \in \mathbb{R}^d}$, see e.g. Theorem 4.2.5, p. 126 of [4].

Integrating by parts we conclude that L is symmetric and \mathcal{A} is anti-symmetric with respect to μ , i.e.

$$(3.2) \quad \int_{\mathbb{R}^d} Lfgd\mu = \int_{\mathbb{R}^d} fLgd\mu$$

and

$$(3.3) \quad \int_{\mathbb{R}^d} \mathcal{A}fgd\mu = - \int_{\mathbb{R}^d} f\mathcal{A}gd\mu, \quad f, g \in C_0^\infty(\mathbb{R}^d).$$

The processes $(X(t; x))_{t \geq 0}$ could, in principle, explode in finite time. This scenario is however precluded, thanks to condition A2). More precisely, for any path $\pi : [0, +\infty) \rightarrow \mathbb{R}^d$ and $R > 0$ define

$$(3.4) \quad \tau_R(\pi) := \inf_{t \geq 0} [|\pi(t)| \geq R],$$

with the convention that $\tau_R(\pi) = +\infty$, if the set over which we take infimum is empty.

The function $R \mapsto \tau_R$ is increasing. We can define therefore

$$\mathbf{e}(\pi) := \lim_{R \rightarrow +\infty} \tau_R(\pi).$$

Suppose that the random vector η is distributed according to μ and independent of $(w(t))_{t \geq 0}$. Consider the process

$$X(t) := X(t; \eta), \quad 0 \leq t < \tilde{\mathbf{e}} := \mathbf{e}(X(\cdot; \eta)).$$

Proposition 3.1. *Suppose that (1.3) and conditions A1) - A2) hold. Then,*

$$(3.5) \quad \mathbb{P}[\tilde{\mathbf{e}} < +\infty] = 0.$$

As a result of the above proposition, we can define the process $(X(t))_{t \geq 0}$ for all times.

Proposition 3.2. *The process $(X(t))_{t \geq 0}$ is stationary, i.e. for any $0 \leq t_1 < t_2 < \dots < t_n$, bounded measurable functions f_1, \dots, f_n and $h \geq 0$ we have*

$$(3.6) \quad \mathbb{E} \left[\prod_{i=1}^n f_i(X(t_i + h)) \right] = \mathbb{E} \left[\prod_{i=1}^n f_i(X(t_i)) \right].$$

The proofs of Propositions 3.1 and 3.2 are contained in Section A.1 of Appendix.

It turns out that we can define the diffusion starting from any x for all times. Let

$$(3.7) \quad \tilde{\mathbf{e}}_x := \mathbf{e}(X(\cdot; x)).$$

Proposition 3.3. *For any $x \in \mathbb{R}^d$ we have*

$$(3.8) \quad \tilde{\mathbf{e}}_x < +\infty, \quad a.s.$$

The proof of this result is contained in Section A.2 of Appendix. Thanks to the above result the trajectories $t \mapsto X(t; x)$ are defined for all $t \geq 0$, a.s. for any $x \in \mathbb{R}^d$.

Let $P_t(x, dy)$ be the transition probabilities corresponding to $(X(t; x))_{t \geq 0}$. Thanks to Proposition 3.2 we have

$$(3.9) \quad \int_{\mathbb{R}^d} \mu(dx) P_t(x, A) = \mu(A), \quad A \in \mathcal{B}(\mathbb{R}^d), t > 0.$$

Define by $(P_t)_{t \geq 0}$ the respective transition probability semigroup, given by $P_0 = I$,

$$(3.10) \quad P_t f(x) = \mathbb{E} f(X(t; x)) = \int_{\mathbb{R}^d} f(y) P_t(x, dy), \quad f \in B_b(\mathbb{R}^d), t > 0, x \in \mathbb{R}^d.$$

In consequence of the path continuity of diffusions we conclude the following.

Corollary 3.4. *The semigroup $(P_t)_{t \geq 0}$ extends to a C_0 semigroup on $L^p(\mu)$ for any $p \geq 1$.*

We use an abbreviated notation $P := P_1$.

Proposition 3.5. *There exists a Borel measurable, strictly positive function $p : \mathbb{R}^{2d} \rightarrow (0, +\infty)$ such that for every $f \in L^1(\mu)$*

$$(3.11) \quad P f(x) = \int_{\mathbb{R}^d} p(x, y) f(y) \mu(dy), \quad \mu \text{ a.e.}$$

The proof of the proposition is contained in Section A.3 of Appendix.

As a consequence of (3.11) and Theorem 6 of [11] we get

Corollary 3.6. *For any $f \in L^2(\mu)$ such that*

$$(3.12) \quad \int_{\mathbb{R}^d} f d\mu = 0$$

we have

$$(3.13) \quad \lim_{t \rightarrow +\infty} \|P_t f\|_{L^2(\mu)} = 0.$$

3.2. Proof of Theorem 2.1. We assume with no loss of generality that $f \in C_0^\infty(\mathbb{R}^d)$ satisfies (3.12). We have

$$(3.14) \quad f(x) = - \int_0^{+\infty} P_t \mathcal{L} f(x) dt = - \int_0^{+\infty} \mathbb{E}[\mathcal{L} f(X(t; x))] dt, \quad x \in \mathbb{R}^d.$$

Therefore,

$$(3.15) \quad f_{x_j}(x) = \int_0^{+\infty} v_j(t) dt,$$

where $v_j(t)$ is given by

$$(3.16) \quad \begin{aligned} v_j(t) &:= -\partial_{x_j} P_t \mathcal{L} f(x) = -\mathbb{E} [(\nabla_x \mathcal{L} f)(X(t; x)) \cdot \partial_{x_j} X(t; x)] \\ &= - \sum_{i=1}^d \mathbb{E} [(\mathcal{L} f)_{x_i}(X(t; x)) \xi_{i,j}(t)] \end{aligned}$$

and $\partial_{x_j} X_i(t; x) = \xi_{i,j}(t)$ is the Fréchet derivative of the stochastic flow $x \mapsto X(t; x)$.

Differentiating (3.1) with respect to the initial condition we conclude that

$$(3.17) \quad \begin{aligned} \frac{d\xi_{i,j}}{dt} &= \frac{1}{2} \sum_{i'=1}^d \left[-\partial_{x_i, x_{i'}}^2 U(X(t; x)) + \partial_{x_{i'}} b_i(X(t; x)) \right] \xi_{i',j}, \\ \xi_{i,j}(0) &= \delta_{i,j}, \quad i, j = 1, \dots, d, \end{aligned}$$

where $\delta_{i,j}$ is the Kronecker symbol, i.e. $\delta_{i,i} = 1$ and $\delta_{i,j} = 0$, if $i \neq j$.

For a given random matrix valued field

$$\mathbf{g}(t, x) := (\mathbf{g}_j(t, x))_{j=1, \dots, d} = [g_{i,j}(t, x)]_{i,j=1, \dots, d}, \quad t \geq 0, \quad x \in \mathbb{R}^d,$$

and $r \geq 1$ we let

$$(3.18) \quad \|\mathbf{g}\|_{r,t} := \sum_{i,j=1}^d \left\{ \int_{\mathbb{R}^d} \mu(dx) \mathbb{E} \left[\int_0^t |g_{i,j}(s, x)|^2 ds \right]^{r/2} \right\}^{1/r}.$$

Let also $\mathbf{h}_j(t, x) = (h_{1,j}(t, x), \dots, h_{d,j}(t, x))$, $j = 1, \dots, d$, where

$$(3.19) \quad h_{i,j}(t, x) := \int_0^t g_{i,j}(s, x) ds, \quad t \geq 0, \quad x \in \mathbb{R}^d.$$

Suppose that $(\mathbf{g}(t, \cdot))_{t \geq 0}$ is with respect to the natural filtration of $(w(t))_{t \geq 0}$. Treating the solution $X(t; x, w)$ of (3.1) as the functional of the Wiener process $w(t) = (w_i(t))_{i=1, \dots, d}$, we define the Malliavin derivative of $X(t; x, w)$ in the direction \mathbf{h}_j

$$\mathcal{D}_{\mathbf{h}_j} X(t; x, w) := \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \{X(t; x, w + \varepsilon \mathbf{h}_j) - X(t; x, w)\},$$

where the limit above is understood in the L^2 sense, see Def. 2.1, p. 35 of [5].

Denote

$$\zeta_{i,j}(t) := \mathcal{D}_{\mathbf{h}_j} X_i(t; x, w), \quad i, j = 1, \dots, d$$

the components of the Malliavin derivatives in the directions $\mathbf{h}_1, \dots, \mathbf{h}_d$. We can see, from (3.1), that they satisfy

$$(3.20) \quad \begin{aligned} \frac{d\zeta_{i,j}}{dt} &= \frac{1}{2} \sum_{i'=1}^d \left[-\partial_{x_i, x_{i'}}^2 U(X(t; x)) + \partial_{x_{i'}} b_i(X(t; x)) \right] \zeta_{i',j} + g_{i,j}, \\ \zeta_{i,j}(0) &= 0, \quad i, j = 1, \dots, d, \end{aligned}$$

From the chain rule for the Malliavin derivative, see Proposition 1.2.3, p. 28 of [9], we obtain

$$(3.21) \quad \mathbb{E} \left[\mathcal{D}_{\mathbf{h}_j} \mathcal{L}f(X(t, x)) \right] = \sum_{i=1}^d \mathbb{E} \left[\partial_{x_i} \mathcal{L}f(X(t, x)) \zeta_{i,j}(t) \right].$$

The difference of the Fréchet and Malliavin derivatives

$$(3.22) \quad \Theta(t, x) := \partial_x X(t; x) - \mathcal{D}_{\mathbf{h}} X(t; x) = (\Theta_{i,j}(t, x))_{i=1, \dots, d}$$

satisfies the following system of equations

$$(3.23) \quad \frac{d\Theta_{i,j}(t, x)}{dt} = \frac{1}{2} \sum_{i'=1}^d [-\partial_{x_i, x_{i'}} U(X(t; x)) + \partial_{x_{i'}} b_j(X(t; x))] \Theta_{i',j}(t, x) - g_{i,j}(t, x),$$

$$\Theta_{i,j}(0, x) = \delta_{i,j}, \quad i, j = 1, \dots, d.$$

Accordingly, using (3.21) together with (3.22), we get

$$(3.24) \quad v_j(t) = \tilde{v}_j(t) - \mathbb{E} [\mathcal{D}_{b_j} \mathcal{L}f(X(t; x))],$$

with $\tilde{v}_j(t)$ given by

$$(3.25) \quad \tilde{v}_j(t) := - \sum_{i=1}^d \mathbb{E} [(\mathcal{L}f)_{x_i}(X(t; x)) \Theta_{i,j}(t, x)].$$

Integrating by parts the second term on the right hand side of (3.24), see Theorem 2.1 p. 236 of [1] (and also Lemma 1.2.1 p. 25, of [9]), we conclude that

$$(3.26) \quad v_j(t) = \tilde{v}_j(t) - \sum_{i=1}^d \mathbb{E} \left[\mathcal{L}f(X(t; x)) \int_0^t g_{i,j}(s, x) dw_i(s) \right].$$

We shall look for $\mathbf{g}^T(t, t_0, x) = [\mathbf{g}_1(t, t_0, x), \dots, \mathbf{g}_d(t, t_0, x)]$ - a column vector - that satisfies the following conditions:

- i) it is adapted with respect to the natural filtration of $(w(t))_{t \geq 0}$,
- ii) given $t_0 \in (0, t_*]$, where $t_* := \gamma_0/r$ and parameter γ_0 as in (2.8), we have both $\Theta(t, x) \equiv 0$ and $\mathbf{g}(t, t_0, x) \equiv 0$ for $t \geq t_0$, $x \in \mathbb{R}^d$,
- iii) we have (cf (3.18))

$$(3.27) \quad \|\mathbf{g}(\cdot; t_0)\|_{r, t_0} < +\infty.$$

Suppose that we can construct such an object, that in what follows we call a control (we show how to do it in Section 3.3). In the remaining part of the present section, we demonstrate how to conclude with its help the proof of (2.9).

Assume first that $t \geq t_0$. Thanks to ii), we conclude that then $\tilde{v}_j(t) \equiv 0$. Using formula (3.26) and the Markov property of $(X(t; x))_{t \geq 0}$ we can write

$$(3.28) \quad v_j(t) = - \sum_{i=1}^d \mathbb{E} \left[P_{t-t_0} \mathcal{L}f(X(t_0; x)) \int_0^{t_0} g_{i,j}(s; t_0, x) dw_i(s) \right], \quad \text{for } t \geq t_0,$$

which, invoking Corollary 3.6, yields, upon integration from t_0 to $+\infty$,

$$(3.29) \quad (P_{t_0} f)_{x_j}(x) = \int_{t_0}^{+\infty} v_j(t) dt = \sum_{i=1}^d \mathbb{E} \left[f(X(t_0; x)) \int_0^{t_0} g_{i,j}(s; t_0, x) dw_i(s) \right].$$

First equality above comes from (3.14). Denote $\mathbf{g}(t; t_0) := \mathbf{g}(t; t_0, \eta)$, where η is the stationary initial data vector for (3.1).

For $1/p = 1/q + 1/r$, we can write, by the Burkholder-Davies-Gundy inequality

$$\begin{aligned}
(3.30) \quad & \|\nabla_x P_{t_0} f\|_{L^p(\mu)} \leq \|f\|_{L^q(\mu)} \left\{ \mathbb{E} \left| \int_0^{t_0} \mathfrak{g}(s; t_0) \cdot dw(s) \right|^r \right\}^{1/r} \\
& \leq C(d, r) \|f\|_{L^q(\mu)} \left\{ \mathbb{E} \left[\int_0^{t_0} \operatorname{tr} (\mathfrak{g}^T(s; t_0) \mathfrak{g}(s; t_0)) ds \right]^{r/2} \right\}^{1/r} \\
& = C(d, r) t_0^{1/2} \|f\|_{L^q(\mu)} \left\{ \mathbb{E} \left[\frac{1}{t_0} \int_0^{t_0} \operatorname{tr} (\mathfrak{g}^T(s; t_0) \mathfrak{g}(s; t_0)) ds \right]^{r/2} \right\}^{1/r} \\
& \leq C(d, r) t_0^{1/2} \|f\|_{L^q(\mu)} \left\{ \frac{1}{t_0} \int_0^{t_0} \mathbb{E} [\operatorname{tr} (\mathfrak{g}^T(s; t_0) \mathfrak{g}(s; t_0))]^{r/2} ds \right\}^{1/r}.
\end{aligned}$$

By virtue of estimate (3.45) below we conclude that

$$(3.31) \quad \|\nabla_x P_{t_0} f\|_{L^p(\mu)} \leq C(d, r) t_0^{-1/2} \|f\|_{L^q(\mu)} \left\{ \int_{\mathbb{R}^d} \mathfrak{E}(t_0 r \mathbf{u}(x)) \mu(dx) \right\}^{1/r},$$

with $\mathfrak{E}(\cdot)$ given by (3.44). Therefore (see (3.29)), we obtain the following bound

$$(3.32) \quad \left\| \int_{t_0}^{+\infty} v_j(t) dt \right\|_{L^p(\mu)} \leq \frac{C}{t_0^{1/2}} \|f\|_{L^q(\mu)}, \quad t_0 \in (0, t_*)$$

and

$$(3.33) \quad C := C(d, r) \left\{ \int_{\mathbb{R}^d} \exp \{ \gamma_0 \mathbf{u}(x) \vee 0 \} \mu(dx) \right\}^{1/r}$$

It remains to consider the case when $t \in (0, t_0)$. We will use estimate (3.31). Recall that t_0 appearing there can be chosen arbitrarily from $(0, t_*)$. Applying (3.31) with $\mathcal{L}f$ instead of f and $t \in (0, t_0]$ instead of t_0 we conclude that

$$\|v_j(t)\|_{L^p(\mu)} \leq \frac{C}{t^{1/2}} \|\mathcal{L}f\|_{L^q(\mu)}, \quad t \in (0, t_0],$$

with C given by (3.33). Therefore,

$$(3.34) \quad \left\| \int_0^{t_0} v_j(t) dt \right\|_{L^p(\mu)} \leq \int_0^{t_0} \|v_j(t)\|_{L^p(\mu)} dt \leq C \|\mathcal{L}f\|_{L^q(\mu)} \int_0^{t_0} \frac{dt}{\sqrt{t}} = C t_0^{1/2} \|\mathcal{L}f\|_{L^q(\mu)}.$$

From estimates (3.32) and (3.34) we conclude that there exists a constant $C > 0$ such that

$$(3.35) \quad \|\nabla_x f\|_{L^p(\mu)} \leq C \left(t_0^{1/2} \|\mathcal{L}f\|_{L^q(\mu)} + t_0^{-1/2} \|f\|_{L^q(\mu)} \right), \quad f \in C_0^\infty(\mathbb{R}^d), \quad t_0 \in (0, t_*)$$

and (2.9) follows. \square

3.3. Construction of a control $\mathbf{g}(t, t_0, x)$. We construct first $\mathbf{g}(t, 1, x)$. For a time being we suppress writing arguments 1 and x . Denote by $C(t, s) = [C_{i,i'}(t, s)]_{i,i'=1,\dots,d}$ the fundamental matrix of the system (3.17). It is a $d \times d$ -matrix, which is the solution of the equation

$$(3.36) \quad \frac{d}{dt}C(t, s) = A(t)C(t, s), \quad C(s, s) = I_d, \quad t, s \geq 0,$$

where I_d is the identity $d \times d$ -matrix and $A(t) = [A_{i,i'}(t)]_{i,i'=1,\dots,d}$, where

$$(3.37) \quad A_{i,i'}(t) := \frac{1}{2} \left[-\partial_{x_i x_{i'}}^2 U(X(t, x)) + \partial_{x_i} b_{i'}(X(t, x)) \right].$$

We have

$$C(u, t)C(t, s) = C(u, s), \quad u, t, s \in \mathbb{R}.$$

System (3.23) can be rewritten as follows

$$(3.38) \quad \frac{d\Theta(t)}{dt} = A(t)\Theta(t) - \mathbf{g}(t), \quad \Theta(0) = I_d,$$

where $\Theta(t) = [\Theta_{ij}(t)]$ was defined in (3.22) and I_d is a d -dimensional identity matrix.

The solution of equation (3.38) can be expressed by the Duhamel formula

$$(3.39) \quad \Theta(t) = - \int_0^t C(t, s)\mathbf{g}(s)ds + C(t, 0), \quad t \geq 0.$$

We wish to show that $\Theta(t) \equiv 0$ for $t \geq 1$, see condition ii). For that purpose let

$$(3.40) \quad \mathbf{g}(t) := C(t, 0), \quad t \in [0, 1],$$

and let $\mathbf{g}(t) := 0$ for $t \geq 1$. The process is adapted with respect to the natural filtration of $(w_t)_{t \geq 0}$, satisfying therefore condition i). Additionally, we have

$$\Theta(1) = - \int_0^1 C(1, s)\mathbf{g}(s)ds + C(1, 0) = - \int_0^1 C(1, 0)ds + C(1, 0) = 0.$$

Thus, $\Theta(t) \equiv 0$ for $t \geq 1$. Condition ii) is therefore fulfilled.

In the general case we let

$$(3.41) \quad \mathbf{g}(t; t_0, x) := \frac{1}{t_0}C(t, 0), \quad t \in (0, t_0) \quad \text{and} \quad \mathbf{g}(t; t_0, x) := 0, \quad t \geq t_0.$$

It remains to be checked that $(\mathbf{g}(t, t_0, x))_{t \geq 0}$, constructed above, satisfies estimate (3.27). From (3.36) we conclude that

$$(3.42) \quad \frac{d}{dt}\mathbf{g}_i(t, t_0, x) = A(t)\mathbf{g}_i(t, t_0, x), \quad t \in [0, t_0],$$

where $\mathbf{g}_i(t, t_0, x)$ is the i -th column of $\mathbf{g}(t, t_0, x)$. Multiplying scalarly both sides of (3.42) by $\mathbf{g}_i(t; t_0, x)$ we get

$$\frac{1}{2} \frac{d}{dt} |\mathbf{g}_i(t, t_0, x)|^2 = A(t)\mathbf{g}_i(t, t_0, x) \cdot \mathbf{g}_i(t, t_0, x) \leq \mathbf{u}(X(t, x)) |\mathbf{g}_i(t, t_0, x)|^2.$$

Thus, by the Gronwall inequality we obtain

$$(3.43) \quad |\mathbf{g}_i(t, t_0, x)|^2 \leq \frac{1}{t_0^2} \exp \left\{ 2 \int_0^t \mathbf{u}(X(s, x)) ds \right\}, \quad t \in [0, t_0]$$

Therefore, from (3.43) (recall $\mathbf{g}(s; t_0) = \mathbf{g}(s; t_0, \eta)$)

$$\begin{aligned} & \left\{ \frac{1}{t_0} \int_0^{t_0} \mathbb{E} [\text{tr} (\mathbf{g}^T(s; t_0) \mathbf{g}(s; t_0))]^{r/2} ds \right\}^{1/r} \\ & \leq \sqrt{d} \left\{ \frac{1}{t_0^{1+r}} \int_0^{t_0} \mathbb{E} \left[\exp \left\{ r \int_0^s \mathbf{u}(X(\tau)) d\tau \right\} \right] ds \right\}^{1/r}. \end{aligned}$$

Using the Jensen inequality and then, subsequently, the stationarity of $(X(t))_{t \geq 0}$, the right hand side estimates by

$$\begin{aligned} & \sqrt{d} \left\{ \frac{1}{t_0^{1+r}} \int_0^{t_0} \frac{ds}{s} \int_0^s \mathbb{E} [\exp \{rsu(X(\tau))\}] d\tau \right\}^{1/r} \\ & = \sqrt{d} \left\{ \frac{1}{t_0^{1+r}} \int_0^{t_0} \frac{ds}{s} \int_0^s d\tau \int_{\mathbb{R}^d} \exp \{rsu(x)\} \mu(dx) \right\}^{1/r} \\ & = \sqrt{d} \left\{ \frac{1}{t_0^{1+r}} \int_0^{t_0} ds \int_{\mathbb{R}^d} \exp \{rsu(x)\} \mu(dx) \right\}^{1/r}. \end{aligned}$$

Performing integration over the s variable, we get that the utmost the right hand side equals

$$\sqrt{d} \left\{ \frac{1}{t_0^r} \int_{\mathbb{R}^d} \mathfrak{E}(rt_0 \mathbf{u}(x)) \mu(dx) \right\}^{1/r},$$

where $\mathfrak{E}(0) := 1$ and

$$(3.44) \quad \mathfrak{E}(x) := \frac{e^x - 1}{x}, \quad x \neq 0.$$

Note that

$$0 < \mathfrak{E}(x) \leq \begin{cases} e^x & \text{for } x \geq 0 \\ 1 \wedge |x|^{-1} & \text{for } x < 0. \end{cases}$$

We have shown therefore that

$$(3.45) \quad \left\{ \frac{1}{t_0} \int_0^{t_0} \mathbb{E} [\text{tr} (\mathbf{g}^T(s; t_0) \mathbf{g}(s; t_0))]^{r/2} ds \right\}^{1/r} \leq \frac{\sqrt{d}}{t_0} \left\{ \int_{\mathbb{R}^d} \mathfrak{E}(rt_0 \mathbf{u}(x)) \mu(dx) \right\}^{1/r}.$$

This finishes the proof of (3.27).

3.4. Proof of Corollary 2.2. Let $F = fe^{-U/p}$. Note that

$$\begin{aligned} \partial_{x_i x_j}^2 F &= \partial_{x_i x_j} f e^{-U/p} - \frac{1}{p} \partial_{x_i} f \partial_{x_j} U e^{-U/p} \\ &\quad - \frac{1}{p} \partial_{x_j} f \partial_{x_i} U e^{-U/p} - \frac{1}{p} f \left(\partial_{x_i x_j} U - \frac{1}{p} \partial_{x_i} U \partial_{x_j} U \right) e^{-U/p}. \end{aligned}$$

Let $p' \in (p, q)$. Using the assumption (2.10) for U and an appropriate r , we can find $C > 0$ such that

$$\|\nabla_x^2 f\|_{L^p(\mu)} \leq \|\nabla_x^2 F\|_{L^p(\mathbb{R}^d)} + C\|\nabla_x f\|_{L^{p'}(\mu)} + C\|f\|_{L^q(\mu)}.$$

The classical Calderon-Zygmund L^p estimates (with respect to the Lebesgue measure) give

$$\|\nabla_x^2 F\|_{L^p(\mathbb{R}^d)} \leq C\|\Delta_x F\|_{L^p(\mathbb{R}^d)}.$$

Note that

$$\begin{aligned} \Delta_x F(x) &= \mathcal{L}f(x)e^{-U(x)/p} + a(U, \nabla_x U, H, \nabla_x H) \cdot \nabla_x f(x)e^{-U(x)/p} \\ &\quad + b(\nabla_x U, \Delta_x U, H, \nabla_x H)f(x)e^{-U(x)/p}, \end{aligned}$$

where $a(\cdot)$, $b(\cdot)$ are some polynomials in the indicated variables. Therefore

$$\|\Delta_x F\|_{L^p(\mathbb{R}^d)} \leq \|\mathcal{L}f\|_{L^p(\mu)} + C\|\nabla_x f\|_{L^{p'}(\mu)} + C\|f\|_{L^q(\mu)}.$$

Using the already proved estimated for $\|\nabla_x f\|_{L^{p'}(\mu)}$, in terms of $\|\mathcal{L}f\|_{L^q(\mu)} + \|f\|_{L^q(\mu)}$ for $p' < q$, we conclude the corollary. \square

APPENDIX A. PROOFS OF THE AUXILIARY FACTS ABOUT THE DIFFUSION

A.1. Proofs of Propositions 3.1 and 3.2. For $R > 0$, let $\chi_R \in C_0^\infty(\mathbb{R}^d)$ be such that $\chi_R(x) \equiv 1$ for $|x| \leq R + 1$, $\chi_R(x) \equiv 0$ for $|x| \geq R + 2$ and $\|\nabla_x \chi_R\|_\infty \leq 2$. Define $H^{(R)}(x) = [H_{i,j}^{(R)}(x)]$, where $H_{i,j}^{(R)}(x) := H_{i,j}(x)\chi_R(x)$, $i, j = 1, \dots, d$ and $U^{(R)}(\cdot)$ a C^2 -smooth potential, such that $U^{(R)}(x) = U(x)$, $|x| \leq R + 1$ and $U^{(R)}(x) = |x|$, $|x| \geq R + 2$.

Let $\mu_R(dx) = Z_R^{-1}e^{-U_R(x)}dx$ be a probability measure, with Z_R – the appropriate normalizing constant. We shall assume that the truncation is made in such a way that $\lim_{R \rightarrow +\infty} Z_R = Z$. As a result $\lim_{R \rightarrow +\infty} \mu_R = \mu$ in total variation.

Let L_R and \mathcal{A}_R be the differential operators defined by formulas analogous to (2.2) and (2.3), where the respective coefficients have been replaced by the truncated ones introduced above. Then $\mathcal{L}_R = L_R + \mathcal{A}_R$ is the generator of a family of diffusions $(X_R(t, x))_{t \geq 0}$ indexed by the starting point. The stationary $X_R(t) = X_R(t, \eta)$, where $\eta(x) = x$ is distributed according to μ_R . Let $\mathbf{e}_{R,x} := \tau_R(X_R(\cdot, x))$, see (3.4), be the exit time of diffusion $(X_R(t, x))_{t \geq 0}$ from the ball of radius $R > 0$, centered at 0. Note that

$$X_R(t, x) = X(t, x), \quad 0 \leq t \leq \mathbf{e}_{R,x} \quad \text{and} \quad \mathbf{e}_{R,x} = \tilde{\mathbf{e}}_{R,x} := \tau_R(X(\cdot, x)).$$

To prove (3.5) it suffices to show that for any $T > 0$

$$(A.1) \quad \lim_{R \rightarrow +\infty} \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[\tilde{\mathbf{e}}_{R,x} \leq T] = 0.$$

Since μ_R converges to μ in the total variation it suffices to show that

$$(A.2) \quad \lim_{R \rightarrow +\infty} \int_{\mathbb{R}^d} \mu_R(dx) \mathbb{P}[\mathbf{e}_{R,x} \leq T] = 0.$$

From Itô formula applied to $f(X_R(t, x))$, where

$$f(x) := (|x|^2 + 1)^\rho,$$

we conclude

$$\begin{aligned} f(X_R(T \wedge \mathbf{e}_{R,x}, x)) &= f(x) + \frac{1}{2} \int_0^{T \wedge \mathbf{e}_{R,x}} \nabla f(X_R(s, x)) \cdot [-\nabla_x U(X_R(s, x)) + b(X_R(s, x))] ds \\ &+ \frac{1}{2} \int_0^{T \wedge \mathbf{e}_{R,x}} \Delta_x f(X_R(s, x)) ds + \int_0^{T \wedge \mathbf{e}_{R,x}} \nabla f(X_R(s, x)) \cdot dw(s). \end{aligned}$$

Here, we assume that ρ is as in condition A3) and it belongs to $(0, 1)$. Applying the expectation to both sides of (A.2) and averaging over the initial data with respect to the stationary measure μ_R we conclude

$$(A.3) \quad \begin{aligned} \int_{\mathbb{R}^d} \mu_R(dx) \mathbb{E} f(X_R(T \wedge \mathbf{e}_{R,x}, x)) &\leq \int_{\mathbb{R}^d} f(x) \mu_R(dx) \\ &+ \frac{T}{2} \int_{\mathbb{R}} |\nabla_x f(x) \cdot [-\nabla_x U(x) + b(x)]| \mu_R(dx) + 2\rho T d. \end{aligned}$$

Here we have used the inequality

$$|\Delta_x f(x)| = 2\rho(|x|^2 + 1)^{\rho-2} |(d + 2(\rho - 1))|x|^2 + d| \leq 4\rho d, \quad x \in \mathbb{R}^d.$$

Hence, there exists some constant $C > 0$, independent of $R > 0$, such that

$$\int_{\mathbb{R}^d} \mu_R(dx) \mathbb{E} |X_R(T \wedge \mathbf{e}_{R,x}, x)|^{2\rho} \leq C + CT.$$

As a result

$$\int_{\mathbb{R}^d} \mu_R(dx) \mathbb{P}[\mathbf{e}_{R,x} \leq T] \leq \frac{C + CT}{R^{2\rho}}.$$

Passing with $R \rightarrow \infty$ we conclude (A.2), which ends the proof of Proposition 3.1. \square

To show Proposition 3.2, observe first that an analogue of (3.6) holds for $(X_R(t))_{t \geq 0}$ and functions f_1, \dots, f_n that are bounded and continuous. Letting $R \rightarrow +\infty$ we conclude the equality for stationary process $(X(t))_{t \geq 0}$ and bounded continuous functions. Using an approximation argument we can extend (3.6) to arbitrary bounded and measurable functions f_1, \dots, f_n that ends the proof of Proposition 3.2. \square

A.2. Proof of Proposition 3.3. From Proposition 3.1 it follows that

$$(A.4) \quad \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[\tilde{\mathbf{e}}_x < +\infty] = 0.$$

The above equality implies that there exists a Borel measurable set of null Lebesgue measure $\mathcal{Z} \subset \mathbb{R}^d$, such that for any $x \notin \mathcal{Z}$ we have $\tilde{\mathbf{e}}_x = +\infty$.

For an arbitrary $x \in \mathbb{R}^d$ there exists $R > 0$ such that the sphere $S_R(x) := [y : |y-x| = R]$ intersects with \mathcal{Z} on a set \mathcal{N} of a null surface Lebesgue measure $\omega_{d,R}$. Consider the harmonic measure $\omega_R^x(A) = \mathbb{P}[X_R(\mathbf{e}_{R,x}, x) \in A]$, where $A \in \mathcal{B}(S_R(x))$ – the Borel σ -algebra of subsets of the sphere. It is equivalent with respect to $\omega_{d,R}$, see Theorem 4.4, p. 311 of [10]. Hence $\omega_R^x(\mathcal{N}) = 0$. As a result

$$X_R(\mathbf{e}_{R,x}, x) = X(\tilde{\mathbf{e}}_{R,x}, x) \notin \mathcal{N}, \quad \text{a.s.}$$

and for such an event we can let

$$X(t, x) := X(t - \tilde{\mathbf{e}}_{R,x}, X(\tilde{\mathbf{e}}_{R,x})) \quad \text{for all } t \geq \tilde{\mathbf{e}}_{R,x},$$

which ends the proof of Proposition 3.3. \square

A.3. Proof of Proposition 3.5. We use the notation of Section A.1. Let ν be the law of the random vector $(X(0, \eta), X(1, \eta))$ in $(\mathbb{R}^{2d}, \mathcal{B}(\mathbb{R}^{2d}))$. We claim that it is equivalent with the Lebesgue measure m_{2d} , i.e. the families of null sets in both ν and m_{2d} measures are equal. First suppose that $A \in \mathcal{B}(\mathbb{R}^{2d})$ is a m_{2d} -null measure set. Given $R > 0$ let ν_R be the law of $(X_R(0, \eta), X_R(1, \eta))$, with η distributed according to μ_R . It is quite straightforward (e.g. using the Girsanov theorem) to conclude that ν_R is equivalent with m_{2d} . Let $A_R := A \cap (S_R(0) \times S_R(0))$ and let $\mathbf{e}_{R,x} := \tau_R(X_R(t, x)) = \tau_R(X(t, x))$. We have $\nu_R[A_R] = 0$ and, by the argument made in Section A.1, we know that

$$(A.5) \quad \lim_{R \rightarrow +\infty} \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[\mathbf{e}_{R,x} < 1] = 0.$$

Hence

$$\begin{aligned} \nu[A_R] &= \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[(x, X(1, x)) \in A_R] \\ &\leq \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[(x, X(1, x)) \in A_R, \mathbf{e}_{R,x} \geq 1] + \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[\mathbf{e}_{R,x} < 1] \\ &\leq \frac{Z_R}{Z} \nu_R[A_R] + \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[\mathbf{e}_{R,x} < 1] = \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[\mathbf{e}_{R,x} < 1]. \end{aligned}$$

Letting $R \rightarrow +\infty$, we conclude that $\nu[A] = \lim_{R \rightarrow +\infty} \nu[A_R] = 0$.

Now suppose that $m_{2d}(A) > 0$ and $\nu[A] = 0$. Then, for a sufficiently large $R > 0$ we would have $m_{2d}(A_R) > 0$ and

$$\begin{aligned} \nu[A] &\geq \nu[A_R] \geq \int_{\mathbb{R}^d} \mu(dx) \mathbb{P}[(x, X(1, x)) \in A_R, \mathbf{e}_{R,x} \geq 1] \\ &= \frac{Z_R}{Z} \int_{\mathbb{R}^d} \mu_R(dx) \mathbb{P}[(x, X_R(1, x)) \in A_R, \mathbf{e}_{R,x} \geq 1] > 0. \end{aligned}$$

The last strict inequality can be seen using, for example, the Girsanov theorem.

Using the Radon-Nikodym theorem we conclude therefore the existence of a strictly positive, Borel measurable density $p(x, y)$, such that

$$\mathbb{E}_\mu F(X(0), X(1)) = \int_{\mathbb{R}^{2d}} F(x, y) p(x, y) \mu(dx) \mu(dy)$$

for any Borel measurable and bounded function $F : \mathbb{R}^{2d} \rightarrow \mathbb{R}$. Here \mathbb{E}_μ is the expectation with respect to measure $\mu \otimes \mathbb{P}$. This in particular implies (3.10).

REFERENCES

- [1] Bell, D., *Divergence theorems in path space. III. Hypocoelliptic diffusions and beyond*. J. Funct. Anal. 251 (2007), 232-253.
- [2] Chojecki, T.; Komorowski, T. *Homogenization of an advection equation with locally stationary random coefficients*. J. Differential Equations 267 (2019), no. 3, 1972-2010.
- [3] Gilbarg, D., Trudinger, N., S., *Elliptic Partial Differential Equations of Second Order*, Springer (2001).
- [4] Kunita, H. *Stochastic Flows and Stochastic Differential Equations*, Cambridge Univ. Press 1990.
- [5] P. Malliavin, *Stochastic Analysis*, Grundlehren Math. Wiss., vol. 313, Springer-Verlag, Berlin, 1997.
- [6] Metafune, G.; Prüss, J.; Rhandi, A.; Schnaubelt, R. *The domain of the Ornstein-Uhlenbeck operator on an L^p -space with invariant measure*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 1 (2002), no. 2, 471-485.
- [7] Metafune, G.; Prüss, J.; Rhandi, A.; Schnaubelt, R. *L^p -regularity for elliptic operators with unbounded coefficients*. Adv. Differential Equations 10 (2005), no. 10, 1131-1164.
- [8] Monsurr, S., Transirico, M., *A priori bounds in L^p for solutions of elliptic equations in divergence form*, Bull. Sci. Math. 137 (2013) 851-866.
- [9] Nualart, D., *The Malliavin Calculus and Related Topics*, Springer New York 2000.
- [10] Ross G. Pinsky, *Positive Harmonic Functions and Diffusions*, Cambridge Univ. Press, 1995
- [11] Pichor, K.; Rudnicki, R.; Tyran-Kamiska, M. *Markov semigroups and their applications*. in Dynamics of Dissipation, P. Garbaczewski and R. Olkiewicz (eds.), Lecture Notes in Physics, vol. 597, Springer, Berlin 2002, 215-238.
- [12] Stein, E.,M., *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, New Jersey, 1970.

TYMOTEUSZ CHOJECKI: INSTITUTE OF MATHEMATICS, UMCS, PL. MARII CURIE-SKŁODOWSKIEJ 1, 20-031, LUBLIN, POLAND.

E-mail address: chojecki.tymoteusz@gmail.com

TOMASZ KOMOROWSKI: INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-636 WARSAW, POLAND.

E-mail address: komorow@hektor.umcs.lublin.pl