

Geometric Formulation for Discrete Points and its Applications

Yuuya Takayama
Nikon Corporation, Japan
Nikon Research Corporation of America, USA
yuuya.takayama@nikon.com

Abstract

We introduce a novel formulation for geometry on discrete points. It is based on *a universal differential calculus*, which gives a geometric description of a discrete set by the algebra of functions. We expand this mathematical framework so that it is consistent with differential geometry, and works on spectral graph theory and random walks. Consequently, our formulation comprehensively demonstrates many discrete frameworks in probability theory, physics, applied harmonic analysis, and machine learning. Our approach would suggest the existence of an intrinsic theory and a unified picture of those discrete frameworks.

1 Introduction

Mathematical approaches play an essential role in understanding of practical harmonic techniques. Though differential geometry has contributed to the theoretical studies of the Laplacian, it does not work on discrete points, such as data. In order to formulate its discrete analogue on a set of points, we focus on *a universal differential calculus* [14, 17], which has an advantage to define the exterior derivative without any additional assumption on points, likesuch as continuous models or graphs. Since it is also possible to extend it to define the (discrete) Laplacian, this framework is naturally expected to provide a unified view among Laplacian-based algorithms in applied harmonic analysis and machine learning. Therefore, in this paper, we aim to construct a general formulation to enable differential geometry to work on discrete points with the help of a universal differential calculus, and then study how it shows geometric relationship of frameworks in applied harmonic analysis, machine learning, and so on.

In order to build a general setting, we start from defining a differential 1-form, a measure on functions, an inner product on 1-forms, and the Dirichlet energy over a set of discrete points, which is regarded as a manifold. Then, the Laplacian is immediately given as the Laplace-Beltrami operator. It is worth emphasizing that this Laplacian is compatible with

Table 1.1: dictionary of formulation

manifold	discrete set V	detail
set of function	$\mathcal{A} = \{f: V \rightarrow \mathbb{R}\}$	§2.1
exterior derivative	$\partial: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$	Dfn. 2.3
differential 1-form	$\Omega_{\mathcal{A}}^1 = \partial\mathcal{A} \subset \mathcal{A} \otimes \mathcal{A}$	Dfn. 2.3
integral of function	$\int_V \cdot d\mu: \mathcal{A} \rightarrow \mathbb{R}$	(2.6)
inner product on functions	$\langle \cdot, \cdot \rangle_{\mathcal{A}}: \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{R}$	(2.7)
inner product on 1-forms	$\langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1}: \Omega_{\mathcal{A}}^1 \times \Omega_{\mathcal{A}}^1 \rightarrow \mathbb{R}$	(2.9)
Dirichlet energy	$\mathcal{E}(\cdot, \cdot): \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{R}$	(2.10)
Laplacian for function	$\mathcal{L} = \partial^* \circ \partial: \mathcal{A} \rightarrow \mathcal{A}$	(2.15)
Fourier transform	$\mathcal{F}: \mathcal{A} \rightarrow \mathbb{R}^{ V }$	(2.17)
embedding	$\vec{r}_d: V \rightarrow \mathbb{R}^d$	(2.19)
curvature vector	$\vec{H}_d = -\mathcal{L}\vec{r}_d \in \mathcal{A}^{\oplus d}$	(2.20)

that given in spectral graph theory [8, 27] and random walks [25, 1]. Finally, we define the Fourier transform and the curvature vector of an embedding, which characterize geometric aspects of points. In summary, our formulation for differential geometry on discrete points consists of those in Table 1.1.

To show advantages of this formulation, we demonstrate three types of applications. First, we study a graph based frameworks; spectral graph theory and random walks. There, we review useful techniques for other applications to verify compatibility between our setting and theirs. Second, we figure out geometric aspects of principal component analysis and classical many-body physics. Though these frameworks are usually not explained in geometric contexts, a covariance and a force are interpreted as the Dirichlet energy and the curvature vector respectively. Third, we understand practical applications, signal processing and manifold leaning, in applied harmonic analysis and machine learning by their relations with other frameworks.

This paper is organized as follows: In §2, we explain the way to construct discrete differential geometry as in Table 1.1. Since this section is discussed in an abstract manner, we summarize main concepts by matrix description in §3 for the sake of the reader. In §4 and §5, we review some results from spectral graph theory and random walks, in §6 and §7, we explain geometric viewpoints in principal component analysis and physics, and last we study signal processing and manifold learning in §8 and §9 respectively.

2 Differential geometry on discrete points

In this section, we review a universal differential calculus, and then define differential geometry on a set of discrete points.

In §2.1, we check algebraic aspects of a set of functions over discrete points. Then, we build a geometric setting in §2.2. The Laplacian, the Fourier transform, and the curvature vector are introduced in §2.3, §2.4 and 2.5 respectively. Their matrix description is explained in §3.

2.1 universal differential calculus

We recall algebraic structures on functions to make sure the definition of a universal differential calculus. See also [13, 17] for reference.

Let V be a finite set. Without loss of generality, we can assume $V = \{1, 2, \dots, n\}$. The set of functions $\{f: V \rightarrow \mathbb{R}\}$ is denoted by \mathcal{A} , which is an \mathbb{R} -vector space in a standard manner. It is useful to take its basis $\{e_i \in \mathcal{A}\}_{i \in V}$ as $e_i(x) = \delta_{ix}$, where δ_{ij} is Kronecker's delta. Define a product $\tilde{\sigma}: \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ as pointwise:

$$\tilde{\sigma}(f, g)(x) = (f \cdot g)(x) := f(x) \cdot g(x). \quad (2.1)$$

By bilinearity, $\tilde{\sigma}$ decomposes into two maps $\iota: \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ and $\sigma: \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ which satisfy $\tilde{\sigma} = \sigma \circ \iota$. Here, the tensor product \otimes is over \mathbb{R} , and $\mathcal{A} \otimes \mathcal{A}$ is regarded as functions on $V \times V$ by $(f \otimes g)(x, y) = f(x) \cdot g(y)$. It is easy to see these maps ι and σ are given as

$$\iota(e_i, e_j) = e_i \otimes e_j, \quad \sigma(e_i \otimes e_j) = \delta_{ij} e_i.$$

We set $1_{\mathcal{A}}$ as the constant function taking a value 1, which is written as $1_{\mathcal{A}} = \sum_{i \in V} e_i$. The equation $f \cdot 1_{\mathcal{A}} = f = 1_{\mathcal{A}} \cdot f$ follows from the definition (2.1), or is checked by $e_i \cdot e_j = \delta_{ij} e_i$. Then, we define left and right actions on $\mathcal{A} \otimes \mathcal{A}$ by $h \cdot (f \otimes g) := (h \cdot f) \otimes g$ and $(f \otimes g) \cdot h := f \otimes (g \cdot h)$ respectively. The next proposition follows:

Proposition 2.2. *\mathcal{A} is an \mathbb{R} -algebra with the product $\tilde{\sigma}$ and the unity $1_{\mathcal{A}}$. $\mathcal{A} \otimes \mathcal{A}$ is an \mathcal{A} -bimodule.*

Now, we introduce a universal differential calculus.

Definition 2.3 ([17]). *For $f = \sum_{i \in V} f_i e_i \in \mathcal{A}$, define a differential map $\partial: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ by*

$$\partial f := 1_{\mathcal{A}} \otimes f - f \otimes 1_{\mathcal{A}} = \sum_{i, j \in V} (f_j - f_i) e_i \otimes e_j,$$

and $\Omega_{\mathcal{A}}^1 \subset \mathcal{A} \otimes \mathcal{A}$ as the minimal left \mathcal{A} -submodule of $\mathcal{A} \otimes \mathcal{A}$ containing $\partial \mathcal{A}$. The pair $(\Omega_{\mathcal{A}}^1, \partial)$ is called the universal first order differential calculus on \mathcal{A} .

Lemma 2.4. *$\Omega_{\mathcal{A}}^1$ is an \mathcal{A} -bimodule.*

Proof. We can see the Leibniz rule holds:

$$\begin{aligned} \partial(f \cdot g) &= 1_{\mathcal{A}} \otimes (f \cdot g) - (f \cdot g) \otimes 1_{\mathcal{A}} \\ &= (1_{\mathcal{A}} \otimes f) \cdot g - f \otimes g + f \otimes g - (f \cdot g) \otimes 1_{\mathcal{A}} = \partial f \cdot g + f \cdot \partial g. \end{aligned}$$

Hence, the element $\partial f \cdot g$ produced by the right action belongs to $\Omega_{\mathcal{A}}^1$. □

The differential map ∂ can be defined on the higher tensor spaces in a similar way to the exterior derivative on manifolds [17, §2]. Hence, we refer to an element of $\Omega_{\mathcal{A}}^1$ as a 1-form.

Lemma 2.5. $\mathcal{A} \otimes \mathcal{A}$ is isomorphic to $\Omega_{\mathcal{A}}^1 \oplus \mathcal{A}$ as \mathcal{A} -bimodules.

Proof. Notice that $e_i \cdot \partial e_j = e_i \otimes e_j$ for $i \neq j$, $e_i \cdot \partial e_i = -\sum_{j \in V \setminus i} e_i \otimes e_j$. Thus, $\Omega_{\mathcal{A}}^1$ is spanned by a basis $\{e_i \otimes e_j \mid i \neq j\}$ in $\mathcal{A} \otimes \mathcal{A}$. The linear map $\sigma: e_i \otimes e_j \mapsto \delta_{ij} e_i$ means $\text{Ker } \sigma = \Omega_{\mathcal{A}}^1$, and then we have $\mathcal{A} \otimes \mathcal{A} \cong \text{Ker } \sigma \oplus \text{Im } \sigma = \Omega_{\mathcal{A}}^1 \oplus \mathcal{A}$. \square

2.2 measure and metric

Let μ be a measure on V , namely, $\mu \in \mathcal{A}$ and $\mu_x := \mu(x) > 0$ for any $x \in V$. We define an integral on \mathcal{A} with respect to the measure and an inner product $\langle \cdot, \cdot \rangle_{\mathcal{A}}: \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{R}$ for $f = \sum f_i e_i, g = \sum g_i e_i \in \mathcal{A}$:

$$\int_V f(x) d\mu_x := \sum_{x \in V} \sum_{i \in V} f_i e_i(x) \mu(x) = \sum_{i \in V} f_i \mu_i, \quad (2.6)$$

$$\langle f, g \rangle_{\mathcal{A}} := \int_V f(x) g(x) d\mu_x = \sum_{i \in V} f_i g_i \mu_i. \quad (2.7)$$

As usual, the corresponding norm $\langle f, f \rangle_{\mathcal{A}}^{1/2}$ is denoted by $\|f\|_{\mathcal{A}}$, and the volume of $A \subset V$ is given by $\text{vol}(A) := \int_A 1_{\mathcal{A}}(x) d\mu_x = \sum_{i \in A} \mu_i$. Put a mean of $f \in \mathcal{A}$ as $m_f := \text{vol}(V)^{-1} \int_V f(x) d\mu_x \in \mathbb{R}$. Note that the evaluation operator is represented in several ways:

$$f_i = f(i) = e_i \cdot f = \langle \mu_i^{-1} e_i, f \rangle_{\mathcal{A}}. \quad (2.8)$$

Let us consider an inner product on 1-forms given as a symmetric bilinear map $\langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1}: \Omega_{\mathcal{A}}^1 \times \Omega_{\mathcal{A}}^1 \rightarrow \mathbb{R}$. In this paper, we define it by

$$\langle e_i \otimes e_j, e_k \otimes e_l \rangle_{\Omega_{\mathcal{A}}^1} := \delta_{ik} \delta_{jl} w_{ij}, \quad (2.9)$$

with $w_{ij} \geq 0$ which satisfies $w_{ij} = w_{ji}$ for $i, j \in V, i \neq j$. For simplicity, we put $w_{ii} = 0$ for $i \in V$. This inner product satisfies the property; $\langle f u g, v \rangle_{\Omega_{\mathcal{A}}^1} = \langle u, f v g \rangle_{\Omega_{\mathcal{A}}^1}$ for $f, g \in \mathcal{A}$. For $u = \sum_{i \neq j} u_{ij} e_i \otimes e_j, v = \sum_{i \neq j} v_{ij} e_i \otimes e_j$, we have

$$\langle u, v \rangle_{\Omega_{\mathcal{A}}^1} = \sum_{i, j \in V, i \neq j} w_{ij} u_{ij} v_{ij}.$$

Define a degree of the inner product as $\text{deg}(i) := \|e_i \cdot \partial e_i\|_{\Omega_{\mathcal{A}}^1}^2$, which is often employed as a measure $\mu_i = \text{deg}(i)$. Since $e_i \cdot \partial e_i = e_i \otimes e_i - e_i \otimes 1_{\mathcal{A}}$, we get $\text{deg}(i) = \sum_{j \in V} w_{ij}$. For $f, g \in \mathcal{A}$,

$$\mathcal{E}(f, g) := \frac{1}{2} \langle \partial f, \partial g \rangle_{\Omega_{\mathcal{A}}^1} = \frac{1}{2} \sum_{i, j \in V} w_{ij} (f_i - f_j)(g_i - g_j) \quad (2.10)$$

is called *the Dirichlet energy* with respect to f and g . It is easy to check

$$\mathcal{E}(e_i, e_j) = \begin{cases} \deg(i) & \text{for } i = j \\ -w_{ij} & \text{for } i \neq j \end{cases} \quad (2.11)$$

Remark 2.12. *The above inner product can be defined through a metric $(\cdot, \cdot)_{\Omega_{\mathcal{A}}^1} : \Omega_{\mathcal{A}}^1 \times \Omega_{\mathcal{A}}^1 \rightarrow \mathcal{A} \otimes \mathcal{A}$:*

$$(e_i \otimes e_j, e_k \otimes e_l)_{\Omega_{\mathcal{A}}^1} := \delta_{ik} \delta_{jl} \frac{w_{ij}}{\mu_i \mu_j} e_i \otimes e_j,$$

which preserves the \mathcal{A} -bimodule structures $(fug, v)_{\Omega_{\mathcal{A}}^1} = f(u, v)_{\Omega_{\mathcal{A}}^1} g = (u, fvg)_{\Omega_{\mathcal{A}}^1}$. By integrating it over $V \times V$, we have the inner product. Moreover, its integration over V corresponds to a dual Riemann metric $g^ : \Omega_M^1 \times \Omega_M^1 \rightarrow \mathcal{C}(M)$ in differential geometry.*

Sometimes, it is useful to consider another basis $\{\tilde{e}_i\} := \{e_i/\sqrt{\mu_i}\}$, which is an orthonormal basis on $(\mathcal{A}, \langle \cdot, \cdot \rangle_{\mathcal{A}})$. By this basis, we can represent $f \in \mathcal{A}$ as

$$f = \sum_{i \in V} \tilde{f}_i \tilde{e}_i, \text{ where } \tilde{f}_i = \langle f, \tilde{e}_i \rangle_{\mathcal{A}} = f_i \sqrt{\mu_i}. \quad (2.13)$$

In general, V is regarded as a set of vertices in *an oriented graph* and w_{ij} as a weight on the oriented edge (i, j) . Here, we can ignore the orientation because of the condition $w_{ji} = w_{ij}$. In this sense, $(\mathcal{A}, \langle \cdot, \cdot \rangle_{\mathcal{A}})$ and $(\Omega_{\mathcal{A}}^1, \langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1})$ are Hilbert spaces on the vertices and the edges respectively [20]. When $w_{ij} = 0$, the edge (i, j) is viewed as disconnected. If there does not exist non-empty proper subset $V' \subset V$ which satisfies $w_{ij} = 0$ and $w_{ji} = 0$ for all $i \in V'$ and $j \in V \setminus V'$, the graph is called connected.

Remark 2.14. *The original universal differential calculus refers to a disconnected edge (i, j) as a non-allowed element $e_i \otimes e_j$, and then realizes a non complete graph as a quotient algebra of $\Omega_{\mathcal{A}}^1$ by the ideal generated by non-allowed elements [17, §4]. This construction seems to describe a topology of a graph, contrary, ours focus on its metric structure.*

According to this convention, we often refer to $\{w_{ij}\}$ as (graph) weights and $(\mathcal{A}, \langle \cdot, \cdot \rangle_{\mathcal{A}}, \Omega_{\mathcal{A}}^1, \langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1})$ as a graph.

2.3 Laplace operator

With the inner products given in 2.2, define a co-differential $\partial^* : \Omega_{\mathcal{A}}^1 \rightarrow \mathcal{A}$ to satisfy $\langle \partial^* u, f \rangle_{\mathcal{A}} = \langle u, \partial f \rangle_{\Omega_{\mathcal{A}}^1}$ for any $u \in \Omega_{\mathcal{A}}^1$ and $f \in \mathcal{A}$. Then, *the Laplacian* $\mathcal{L} : \mathcal{A} \rightarrow \mathcal{A}$ is defined by $\frac{1}{2} \partial^* \circ \partial$, in the same way as the Laplace-Beltrami operator in differential geometry. Since

$$\langle u, \partial f \rangle_{\Omega_{\mathcal{A}}^1} = \sum_{i, j \in V, i \neq j} w_{ij} u_{ij} (f_j - f_i) = \sum_{i, j \in V, i \neq j} w_{ij} (u_{ji} - u_{ij}) f_i,$$

we obtain

$$\partial^* u = \sum_{i \in V} \frac{1}{\mu_i} \left\{ \sum_{j \in V \setminus i} w_{ij} (u_{ji} - u_{ij}) \right\} e_i.$$

Thereby, the Laplacian $\mathcal{L} = \frac{1}{2} \partial^* \circ \partial$ is represented as

$$\mathcal{L}f = \sum_{i,j \in V} \frac{w_{ij}}{\mu_i} (f_i - f_j) e_i = \sum_{i \in V} \frac{\deg(i)}{\mu_i} f_i e_i - \sum_{i,j \in V} \frac{w_{ij}}{\mu_i} f_j e_i. \quad (2.15)$$

This is also known as *the graph Laplacian*, as explained in §3. By definition, we have $\mathcal{E}(f, g) = \langle f, \mathcal{L}g \rangle_{\mathcal{A}}$ and the above representation follows from (2.11) as well. When the corresponding graph is connected, the Dirichlet energy $\mathcal{E}(f, f)$ takes the minimum value 0 if and only if f is a constant function. Since \mathcal{L} is self-adjoint, we can take eigenfunctions $\{v_i\}$ as follows:

$$\mathcal{L}v_i = \rho_i v_i, \quad 0 = \rho_1 \leq \rho_2 \leq \dots \leq \rho_n, \quad \langle v_i, v_j \rangle_{\mathcal{A}} = \delta_{ij}. \quad (2.16)$$

Here, we see $v_1 = \text{vol}(V)^{-1/2} 1_{\mathcal{A}}$ and $v_i \perp 1_{\mathcal{A}}$ for $i \geq 2$.

2.4 Fourier analysis

In the continuous setting, the Fourier transform is given by $\mathcal{F}[f](\xi) = \int f(x) e^{\sqrt{-1}\xi x} dx$, and $e^{\sqrt{-1}\xi x}$ is an eigenfunction of the 1-dimensional Laplacian, $-\frac{d^2}{dx^2} e^{\sqrt{-1}\xi x} = \xi^2 e^{\sqrt{-1}\xi x}$.

On the analogy, in the graph setting, it is natural to use the eigenfunctions $\{v_i\}$ of the Laplacian \mathcal{L} , instead of $e^{\sqrt{-1}\xi x}$, and define $\mathcal{F}[f]_i := \langle f, v_i \rangle_{\mathcal{A}} \in \mathbb{R}$ for $f \in \mathcal{A}$. The transformation $\mathcal{F}: \mathcal{A} \ni f \mapsto \mathcal{F}[f] \in \mathbb{R}^n$ is known as *the graph Fourier transform* [18]. We call $\mathcal{F}[f]_i$ the i -th Fourier coefficient or the i -th frequency. The corresponding inverse Fourier transform is given by

$$f = \sum_{i=1}^n \mathcal{F}[f]_i v_i = \sum_{i=1}^n \langle f, v_i \rangle_{\mathcal{A}} v_i, \quad (2.17)$$

which is just the eigenfunction expansion by $\{v_i\}$. It is easy to see *Parseval's identity* holds:

$$\langle f, g \rangle_{\mathcal{A}} = \sum_{i=1}^n \langle f, v_i \rangle_{\mathcal{A}} \langle g, v_i \rangle_{\mathcal{A}} = \langle \mathcal{F}[f], \mathcal{F}[g] \rangle_{\mathbb{R}^n}.$$

This is valid for other expansions by orthogonal functions, such as (2.13). Sometimes, the convolution operator $*g: \mathcal{A} \rightarrow \mathcal{A}$ is defined so that $\mathcal{F}[f * g]_i = \mathcal{F}[f]_i \mathcal{F}[g]_i$ holds:

$$f * g := \sum_{i=1}^n \mathcal{F}[f]_i \mathcal{F}[g]_i v_i. \quad (2.18)$$

We also obtain relations $\|f\|_{\mathcal{A}} = \|\mathcal{F}[f]\|_{\mathbb{R}^n}$ and $\mathcal{F}[\mathcal{L}f]_i = \rho_i \mathcal{F}[f]_i$.

2.5 embedding and curvature

Our setting so far did not use a coordinate of points in V , just used their indexes. Herein, suppose that points are embedded in Euclidean space \mathbb{R}^d . Namely, we consider a map

$$V \ni i \mapsto (r_1(i), r_2(i), \dots, r_d(i)) =: \vec{r}_d(i) \in \mathbb{R}^d, \quad (2.19)$$

where $r_s \in \mathcal{A}$ and $\vec{r}_d \in \mathcal{A}^{\oplus d}$. This element $\vec{r}_d(i)$ is viewed as a coordinate for a point $i \in V$. The Euclidean group $E(d)$ acts on \mathbb{R}^d , hence it defines a coordinate transformation $\vec{r}_d(i) \mapsto \vec{r}'_d(i) = \vec{r}_d(i) \cdot \vec{R}_d + \vec{u}_d$ for $(\vec{R}_d, \vec{u}_d) \in E(d) \cong O(d) \times \mathbb{R}^d$ (as a set).

In differential geometry, an embedding $\vec{r}: \mathcal{M} \rightarrow \mathbb{E}^d$ induces a Riemann metric $g_{\vec{r}}$ on a manifold \mathcal{M} , and especially determines the Laplace-Beltrami operator $\mathcal{L}_{g_{\vec{r}}}$. The normal bundle is given on \mathcal{M} , and then the mean curvature vector $\vec{H}_{\vec{r}}$ is defined as the trace of the second fundamental form divided by $n = \dim \mathcal{M}$. Hence, the vector indicates the normal direction on each point of \mathcal{M} , and its length is called the mean curvature. Beltrami's formula relates those objects as

$$\mathcal{L}_{g_{\vec{r}}} \vec{r} = -n \cdot \vec{H}_{\vec{r}}.$$

One can refer to [7, 4] for mathematical details.

Motivated by this formula, we *define* a graph curvature vector $\vec{H}_d = (H_1, H_2, \dots, H_d) \in \mathcal{A}^{\oplus d}$ of an embedding \vec{r}_d by

$$\vec{H}_d := -\mathcal{L} \vec{r}_d, \quad H_s := -\mathcal{L} r_s, \quad \text{for } s = 1, 2, \dots, d. \quad (2.20)$$

Unlike differential geometry, this vector does not indicate the normal direction, because it is not defined for discrete points. Nevertheless, the vector has a special meaning in physics as explained in §7. The embedding energy $\mathcal{E}(\vec{r}_d, \vec{r}_d) := \sum_{s=1}^d \mathcal{E}(r_s, r_s)$ is given with the curvature vector, that is, $\mathcal{E}(\vec{r}_d, \vec{r}_d) = \int_V \langle \vec{r}_d(x), -\vec{H}_d(x) \rangle_{\mathbb{R}^d} d\mu_x$, and invariant by the Euclidean group action. In some cases, it is convenient to suppose a metric $\langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1}$ is induced by an embedding as in Figure 2.1. For example, we can define weights $\{w_{ij}\}$ by using the distance, such as

$$w_{ij} := C \cdot \exp \left(-\frac{\|\vec{r}_d(i) - \vec{r}_d(j)\|_{\mathbb{R}^d}^2}{2\sigma^2} \right).$$

Several researches show this type of weights converges into a heat kernel on a manifold in the limit $|V| \rightarrow \infty$ [21, 3, 9]. Instead, we study another type of weights in Theorem 7.4.

3 Matrix description of the geometric formulation

In this section, we give matrix description of the formulation discussed in §2.




Figure 2.1: Dependency of geometric objects. A dashed arrow means that its head object can be determined by its tail object, but not necessary.

First we remark that our formulation contains two types of parameters in a measure μ and an inner product $\langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1}$ independently. A measure μ is just given by a positive function on V , thus its degree of freedom is n . On the other hand, an inner product $\langle \cdot, \cdot \rangle_{\Omega_{\mathcal{A}}^1}$ is determined by weights $\{w_{ij}\}$ which satisfy $w_{ij} \geq 0, w_{ji} = w_{ij}$ and $w_{ii} = 0$, hence its degree of freedom is $n(n-1)/2$. When they are taken on a certain relation, well-known cases appear as follows.

Let $\mathbf{M} = \text{diag}_i(\mu_i)$, $\mathbf{W} = \{w_{ij}\}_{i,j}$, and $\mathbf{D} = \text{diag}_i(\text{deg}(i))$ be $n \times n$ -matrices.

By $\{e_i\}$ -basis, a function $f \in \mathcal{A}$ is represented as a numerical vector $\mathbf{f} = {}^t(f_1, \dots, f_n)$. Then, an inner product $\langle f, g \rangle_{\mathcal{A}}$ is written as ${}^t\mathbf{f}\mathbf{M}\mathbf{g}$, and the Dirichlet energy $\mathcal{E}(f, g) = 2\langle \partial f, \partial g \rangle_{\Omega_{\mathcal{A}}^1}$ is as ${}^t\mathbf{f}(\mathbf{D} - \mathbf{W})\mathbf{g}$, which does not depend on \mathbf{M} . The Laplacian \mathcal{L} is given as $\mathbf{M}^{-1}(\mathbf{D} - \mathbf{W})$, and its eigenvalue equation is

$$(\mathbf{D} - \mathbf{W})\mathbf{f} = \rho\mathbf{M}\mathbf{f}.$$

The corresponding eigenvectors, denoted by an $n \times n$ -matrix $\mathbf{V} = \{v_j(i)\}_{i,j}$, defines the Fourier transform $\mathcal{F}[\mathbf{f}] = {}^t\mathbf{V}\mathbf{M}\mathbf{f}$ and its inverse $\mathbf{f} = \mathbf{V}\mathcal{F}[\mathbf{f}]$, where ${}^t\mathbf{V}\mathbf{M}\mathbf{V} = \mathbf{I} = \mathbf{V}{}^t\mathbf{V}\mathbf{M}$. An embedding \vec{r}_d is described as an $n \times d$ -matrix $\mathbf{R} = \{r_s(i)\}_{i,s}$, then its curvature vector \vec{H}_d is as $-\mathbf{M}^{-1}(\mathbf{D} - \mathbf{W})\mathbf{R}$.

In the case of $\{\tilde{e}_i\}$ -basis, since $f = \sum_{i \in V} \tilde{f}_i \tilde{e}_i$ as mentioned in (2.13), we have $\langle f, g \rangle_{\mathcal{A}} = {}^t\tilde{\mathbf{f}}\tilde{\mathbf{g}}$, where $\tilde{\mathbf{f}} = {}^t(\tilde{f}_1, \dots, \tilde{f}_n)$. The Laplacian \mathcal{L} is described as $\mathbf{M}^{-1/2}(\mathbf{D} - \mathbf{W})\mathbf{M}^{-1/2}$, because

$$\mathcal{L}f = \sum_{i \in V} \frac{\text{deg}(i)}{\sqrt{\mu_i \mu_i}} \tilde{f}_i \tilde{e}_i - \sum_{i,j \in V} \frac{w_{ij}}{\sqrt{\mu_i \mu_j}} \tilde{f}_j \tilde{e}_i.$$

When assume $\mathbf{M} = \mathbf{I}$ or \mathbf{D} , we obtain the known Laplacians; the combinatorial Laplacian, the random walk Laplacian, and the normalized Laplacian [20]. However, we do not use this configuration for random walks in §5.

The above notations are summarized in Table 3.1.

Table 3.1: matrix description

	general by $\{e_i\}$	$M = I$	$M = D$ by $\{e_i\}$	$M = D$ by $\{\tilde{e}_i\}$
$\langle f, g \rangle_{\mathcal{A}}$	${}^t f M g$	${}^t f g$	${}^t f D g$	${}^t \tilde{f} \tilde{g}$
$\text{vol}(V)$	$\text{tr } M$	$ V = n$	$\text{tr } D$	$\text{tr } D$
$\mathcal{E}(f, g)$	${}^t f (D - W) g$	${}^t f (D - W) g$	${}^t f (D - W) g$	${}^t \tilde{f} D^{-\frac{1}{2}} (D - W) D^{-\frac{1}{2}} \tilde{g}$
\mathcal{L}	$M^{-1} (D - W)$	$D - W$	$I - D^{-1} W$	$I - D^{-\frac{1}{2}} W D^{-\frac{1}{2}}$
$\mathcal{F}[f]$	${}^t V M f$	${}^t V f$	${}^t V D f$	${}^t \tilde{V} \tilde{f}$
\overrightarrow{H}_d	$M^{-1} (W - D) R$	$(W - D) R$	$(D^{-1} W - I) R$	$(D^{-\frac{1}{2}} W D^{-\frac{1}{2}} - I) \tilde{R}$

4 Application I: spectral graph theory

We review basic results about eigenvalue estimation in spectral graph theory to check its compatibility with our formulation. These results are regarded as discrete analogues of spectral geometry and related to the graph cut problem in §9.1. See also [8, 27] for reference.

4.1 upper bound of eigenvalues

First, we estimate an upper bound for the largest eigenvalue ρ_n . Put $\delta := \max_{i \in V} \text{deg}(i) / \mu_i$.

Lemma 4.1. $\rho_n \leq 2\delta$.

Proof. We have

$$\begin{aligned} \rho_n &= \langle v_n, \mathcal{L} v_n \rangle_{\mathcal{A}} = \frac{1}{2} \sum_{i, j \in V} w_{ij} (v_n(i) - v_n(j))^2 \\ &\leq \sum_{i, j \in V} w_{ij} (v_n(i)^2 + v_n(j)^2) \leq 2 \max_{i \in V} \frac{\text{deg}(i)}{\mu_i} \end{aligned}$$

because $1 = \|v_n\|_{\mathcal{A}}^2 = \sum_{i \in V} v_n(i)^2 \mu_i$. □

Next, we give an upper bound for the second smallest eigenvalue ρ_2 , which is characterized as a minimum value of $\mathcal{E}(f, f) / \|f\|_{\mathcal{A}}^2$ in functions $\{f \in \mathcal{A} \mid f \perp 1_{\mathcal{A}}\}$. For this purpose, the *isoperimetric constant* β is useful, because it is defined in a similar way to the characterization of ρ_2 :

$$\beta = \min_{\emptyset \neq A \subsetneq V} \frac{\text{vol}(\partial A)}{\text{vol}(A)} := \min_{\emptyset \neq A \subsetneq V} \frac{\mathcal{E}(\chi_A, \chi_A)}{\|\chi_A\|_{\mathcal{A}}^2}, \quad (4.2)$$

where $\chi_A := \sum_{i \in V} e_i \in \mathcal{A}$ and A is taken over all subsets satisfying $\text{vol}(A) \leq \text{vol}(V)/2$. From (2.11), we can check $\mathcal{E}(\chi_A, \chi_A) = \sum_{i \in A, j \in A^c} w_{ij}$, then $\text{vol}(\partial A^c) = \text{vol}(\partial A)$ for a complement $A^c = V \setminus A$.

Lemma 4.3. $\rho_2 \leq 2\beta$.

Proof. For any $\emptyset \neq A \subsetneq V$ such that $\text{vol}(A) \leq \text{vol}(V)/2$, put $f_A = \text{vol}(A^c)\chi_A - \text{vol}(V)\chi_{A^c}$, which satisfies $f_A \perp 1_A$, then we have

$$\mathcal{E}(f_A, f_A) = \sum_{i \in A, j \in A^c} w_{ij} (\text{vol}(A) + \text{vol}(A^c))^2 = \text{vol}(V)^2 \text{vol}(\partial A),$$

$$\|f_A\|_{\mathcal{A}}^2 = \text{vol}(A) \text{vol}(A^c)^2 + \text{vol}(A)^2 \text{vol}(A^c) = \text{vol}(V) \text{vol}(A) \text{vol}(A^c).$$

Hence, we obtain $\rho_2 \leq \mathcal{E}(f_A, f_A)/\|f_A\|_{\mathcal{A}}^2 \leq 2 \text{vol}(\partial A)/\text{vol}(A)$ by $\text{vol}(V)/2 \leq \text{vol}(A^c)$. \square

4.2 lower bound of eigenvalues

Here, we estimate an lower bound for the second smallest eigenvalue ρ_2 .

Theorem 4.4. $\rho_2 \geq \beta^2/2\delta$.

Proof. First we claim

$$\beta \int_V f(x) d\mu_x \leq \frac{1}{2} \sum_{i, j \in V} w_{ij} |f_i - f_j|, \quad (4.5)$$

for a positive function $f \in \mathcal{A}$ such that $\text{vol}(\{i \in V \mid f(i) > 0\}) \leq \text{vol}(V)/2$. We take a sequence of subsets $\emptyset = A_0 \subsetneq A_1 \subsetneq \cdots \subsetneq A_l \subsetneq A_{l+1} = V$ so that f is represented as $\sum_{s=1}^l h_s \chi_{A_s}$ by $h_s = f|_{A_s \setminus A_{s-1}} - f|_{A_{s+1} \setminus A_s} \in \mathbb{R}_{>0}$. Then, we can see

$$\begin{aligned} \frac{1}{2} \sum_{i, j \in V} w_{ij} |f_i - f_j| &= \sum_{s=1}^l \sum_{i \in A_s, j \in A_s^c} w_{ij} h_s \\ &= \sum_{s=1}^l h_s \text{vol}(\partial A_s) \geq \beta \sum_{s=1}^l h_s \int_V \chi_{A_s}(x) d\mu_x, \end{aligned}$$

as required. Now, we can write $v_2 = g_+ - g_-$ by positive functions $g_+, g_- \in \mathcal{A}$ which satisfies $\text{vol}(\{i \in V \mid g_+(i) > 0\}) \leq \text{vol}(V)/2$ since $\int_V g_+(x) d\mu_x = \int_V g_-(x) d\mu_x$. It is easy to check $\mathcal{E}(g_+, g_+) \leq \mathcal{E}(v_2, g_+) = \rho_2 \|g_+\|_{\mathcal{A}}^2$, then applying (4.5) to $f = g_+$, we obtain

$$\begin{aligned} \beta^2 \|g_+\|_{\mathcal{A}}^4 &\leq \frac{1}{4} \left(\sum_{i, j \in V} w_{ij} |g_+(i) - g_+(j)| \cdot |g_+(i) + g_+(j)| \right)^2 \\ &\leq \frac{1}{4} \left(\sum_{i, j \in V} w_{ij} |g_+(i) - g_+(j)|^2 \right) \cdot \left(\sum_{i, j \in V} w_{ij} |g_+(i) + g_+(j)|^2 \right)^2 \\ &\leq \frac{1}{2} \mathcal{E}(g_+, g_+) \cdot 2 \sum_{i, j \in V} w_{ij} (g_+(i)^2 + g_+(j)^2) \leq 2\rho_2 \delta \|g_+\|_{\mathcal{A}}^4. \end{aligned}$$

In the second inequality, we used the Cauchy-Schwarz inequality: $\langle \omega_-, \omega_+ \rangle_{\Omega_{\mathcal{A}}^1}^2 \leq \|\omega_-\|_{\Omega_{\mathcal{A}}^1}^2 \cdot \|\omega_+\|_{\Omega_{\mathcal{A}}^1}^2$ for $\omega_{\pm} = \sum_{i, j \in V, i \neq j} |v_2(i) \pm v_2(j)| e_i \otimes e_j \in \Omega_{\mathcal{A}}^1$. \square

This theorem is called *Cheeger's inequality* and its continuous analogue is known in differential geometry [6].

5 Application II: random walks

In this section, we deduce some notations of random walks from our formulation. In §5.1, we review random walks briefly, and in §5.2, we consider their connection with a geometric distance.

5.1 heat equations

For $c > 0$, putting $S_c := \text{id} - c^{-1}\mathcal{L}: \mathcal{A} \rightarrow \mathcal{A}$, we have

$$S_c f = \sum_{i \in V} \frac{c\mu_i - \deg(i)}{c\mu_i} f_i e_i + \sum_{i,j \in V} \frac{w_{ij}}{c\mu_i} f_j e_i = \sum_{i,j \in V} \frac{\theta_{ij}}{c\mu_i} f_j e_i, \quad (5.1)$$

for $f \in \mathcal{A}$, where we put $\theta_{ii} = c\mu_i - \deg(i)$ and $\theta_{ij} = w_{ij}$ for $i, j \in V$. Needless to say, the eigenvalue decomposition of S_c is given as

$$S_c v_i = \lambda_i v_i, \quad \lambda_i = 1 - \frac{\rho_i}{c}, \quad 1 = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 1 - 2\frac{\delta}{c},$$

by (2.16) and Lemma 4.1. Besides, $S_c f$ is described as

$$S_c f = \sum_{i=1}^n \lambda_i \langle f, v_i \rangle_{\mathcal{A}} v_i. \quad (5.2)$$

Proposition 5.3. *For $k \in \mathbb{Z}_{\geq 0}$ and $t \in \mathbb{R}_{\geq 0}$, define operators*

$$P_k := S_c^k, \quad Q_t := \sum_{k=0}^{\infty} \frac{e^{-t} t^k}{k!} P_k.$$

In addition, put $p_k := P_k f_0$ and $q_t = Q_t f_0$ for any $f_0 \in \mathcal{A}$. Then, p_k and q_t satisfy the discrete and continuous time heat equations

$$p_{k+1} - p_k = -\frac{1}{c} \mathcal{L} p_k, \quad p_0 = f_0 \quad (5.4)$$

$$\frac{d}{dt} q_t = -\frac{1}{c} \mathcal{L} q_t, \quad q_0 = f_0 \quad (5.5)$$

respectively.

Proof. The discrete time heat equation follows from $p_{k+1} - p_k = (S_c - \text{id})p_k$. Besides, for q_t , we have,

$$\frac{d}{dt}q_t = \sum_{k=0}^{\infty} \frac{e^{-t}t^k}{k!} (-p_k + p_{k+1}) = (S_c - \text{id}) \sum_{k=0}^{\infty} \frac{e^{-t}t^k}{k!} p_k.$$

Hence, q_t is a solution of the continuous time heat equation.

The latter part is also checked by (5.2). We have $p_k = \sum_{i=1}^n \lambda_i^k \langle f_0, v_i \rangle_{\mathcal{A}} v_i$, thus,

$$q_t = \sum_{i=1}^n \sum_{k=0}^{\infty} e^{-t} \frac{t^k \lambda_i^k}{k!} \langle f_0, v_i \rangle_{\mathcal{A}} v_i = \sum_{i=1}^n e^{-t\rho_i/c} \langle f_0, v_i \rangle_{\mathcal{A}} v_i.$$

This gives $Q_t = \exp(-t\mathcal{L}/c)$, then the assertion immediately follows. See also §8.1. \square

In this sense, S_c is viewed as *an integral operator of a heat kernel*, which is given as $\sum_{i=1}^n \lambda_i v_i \otimes v_i \in \mathcal{A} \otimes \mathcal{A}$. If we impose the condition $c \geq \delta$, then we have $(S_c e_y)(x) = \theta_{xy} \geq 0$ for any $x, y \in V$. Thereby, a function $y \mapsto (S_c e_y)(x)$ defines a discrete probability distribution because of $S_c 1_{\mathcal{A}} = 1_{\mathcal{A}}$. In random walk settings,

$$P(x, y) := (S_c e_y)(x) = \frac{\theta_{xy}}{c\mu_x}, \quad p_k(x, y) := \frac{(P_k e_y)(x)}{\mu(y)}$$

is known as *the transition probability and transition density* respectively [25, 1]. Here, $(P_k e_y)(x) = \mathbb{P}^x(X_k = y)$ means the probability of transitioning from x to y in k steps.

5.2 commute time distance

Let $\tau_+ := \min\{k \geq 1 \mid X_k = y\}$ be the first hitting time and $m(x, y) := \mathbb{E}^x[\tau_+]$ be its expectation. It is easy to see

$$\begin{aligned} m(x, y) &= P(x, y) + \sum_{z \in V \setminus y} P(x, z)(1 + m(z, y)) \\ &= 1 + \sum_{z \in V} P(x, z)m(z, y) - P(x, y)m(y, y). \end{aligned}$$

Notice that we can rearrange the above equation as

$$m(\cdot, y) = 1_{\mathcal{A}} + S_c m(\cdot, y) - m(y, y) S_c e_y, \tag{5.6}$$

where $m(\cdot, y) \in \mathcal{A}$.

Theorem 5.7 ([15]). *Put $T(\cdot, y) := m(\cdot, y) - m(y, y)e_y \in \mathcal{A}$. Then, we have*

$$n(x, y) := \frac{T(x, y) + T(y, x)}{c \text{vol}(V)} = \sum_{i=2}^n \frac{1}{\rho_i} (v_i(y) - v_i(x))^2,$$

for any $x, y \in V$.

Proof. From the equation (5.6), we have

$$c^{-1}\mathcal{L}T(\cdot, y) = (\text{id} - S_c)(m(\cdot, y) - m(y, y)e_y) = 1_{\mathcal{A}} - m(y, y)e_y.$$

By taking the inner product with v_i and using (2.8), we get

$$\begin{aligned} 0 &= c^{-1}\rho_1\langle v_1, T(\cdot, y) \rangle_{\mathcal{A}} = \text{vol}(V)^{1/2} - m(y, y)\mu_y v_1(y), & \text{for } i = 1, \\ c^{-1}\rho_i\langle v_i, T(\cdot, y) \rangle_{\mathcal{A}} &= -m(y, y)\mu_y v_i(y), & \text{otherwise.} \end{aligned}$$

The first equation means $m(y, y)\mu_y = \text{vol}(V)$, and the second equation leads to the Fourier coefficients, hence we obtain

$$\begin{aligned} T(\cdot, y) &= \sum_{i=1}^n \mathcal{F}[T(\cdot, y)]_i v_i \\ &= a_0 1_{\mathcal{A}} - \sum_{i=2}^n \frac{c \text{vol}(V)}{\rho_i} v_i(y) v_i = \sum_{i=2}^n \frac{c \text{vol}(V)}{\rho_i} v_i(y) (v_i(y) 1_{\mathcal{A}} - v_i). \end{aligned}$$

At the last equality, we used the fact $T(y, y) = 0$ to determine a constant a_0 . \square

$n(x, y)$ is known as *the commute time distance* divided by $c \text{vol}(V)$, and equal to the Euclidean distance via the embedding

$$V \ni x \mapsto (\rho_2^{-1/2} v_2(x), \rho_3^{-1/2} v_3(x), \dots). \quad (5.8)$$

This is regarded as one of branches of *the Laplacian eigenmaps* [2],

$$V \ni x \mapsto (v_2(x), v_3(x), \dots), \quad (5.9)$$

and it links with PCA is studied by [15]. We also see these relations in the following section.

6 Application III: PCA

Here, we study a geometric aspect of the empirical covariance, and then consider a principal component analysis as a branch of manifold learning.

6.1 random variable and embedding

Let (Ω, \mathcal{F}, P) be a probability space, and we consider a random variable $X = (X_1, X_2, \dots, X_d): \Omega \rightarrow \mathbb{R}^d$. A mean $\mathbb{E}[X_s] \in \mathbb{R}$ and a covariance $C(X_s, X_t) \in \mathbb{R}$ are defined by

$$\mathbb{E}[X_s] := \int_{\Omega} X_s(\omega) dP_{\omega}, \quad C(X_s, X_t) := \mathbb{E}[X_s X_t] - \mathbb{E}[X_s] \mathbb{E}[X_t].$$

Now, we regard $(V, 2^V, \mu / \text{vol}(V))$ as a probability space, then it follows that $X_s \in \mathcal{A}$ and $m_{X_s} = \mathbb{E}[X_s]$. Moreover, we can characterize a covariance as well.

Theorem 6.1. Take $w_{ij} = \mu_i \mu_j / \text{vol}(V)^2$ for $i \neq j$. Then a covariance coincides with the Dirichlet energy: $C(X_s, X_t) = \mathcal{E}(X_s, X_t)$.

Proof. We have

$$\begin{aligned} \mathbb{E}[X_s X_t] - \mathbb{E}[X_s] \mathbb{E}[X_t] &= \frac{1}{2} \sum_{i,j \in V} \frac{\mu_i \mu_j}{\text{vol}(V)^2} (X_s(i) - X_s(j)) (X_t(i) - X_t(j)) \\ &= \mathcal{E}(X_s, X_t), \end{aligned}$$

as required. \square

Corollary 6.2. In the above setting, we have $\mathcal{L}X_s = (X_s - \mathbb{E}[X_s] \cdot 1_{\mathcal{A}}) / \text{vol}(V)$. In particular, $\rho_i = \text{vol}(V)^{-1}$ for $i \geq 2$ in (2.16).

Proof. From (2.15), we have

$$\mathcal{L}f = \sum_{i,j \in V} \frac{\mu_j}{\text{vol}(V)^2} (f_i - f_j) e_i = \frac{1}{\text{vol}(V)} (f - \mathbb{E}[f] \cdot 1_{\mathcal{A}}).$$

For $i \geq 2$, we get $\mathbb{E}[v_i] = 0$, then $\mathcal{L}v_i = \text{vol}(V)^{-1} v_i$. \square

Therefore, if X is a centered variable, then the corresponding curvature vector is 0, and besides if variables $\{X_s\}$ are independent, then they forms orthogonal eigenfunctions of the Laplacian.

6.2 principal component

For a random variable $X: \Omega \rightarrow \mathbb{R}^d$, put $\bar{X}(\omega) := X(\omega) - \mathbb{E}[X]$. Take eigenfunctions of the covariance matrix $C_{\bar{X}}$, that is, $C_{\bar{X}} u_s = \alpha_s u_s$, $u_s \in \mathbb{R}^d$ and $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_d$. This is viewed as a diagonalization by the Euclidean group $E(d)$ acting on \mathbb{R}^d . In this setting, the eigenfunction expansion of $\bar{X}(\omega)$ is given as

$$\bar{X}(\omega) = \sum_{s=1}^d \xi_s(\omega) u_s, \quad \text{where } \xi_s(\omega) = \langle \bar{X}(\omega), u_s \rangle_{\mathbb{R}^d} \in \mathbb{R}. \quad (6.3)$$

The coefficient ξ_s is a map $\Omega \rightarrow \mathbb{R}$, hence, a random variable. A straightforward calculation shows

$$\mathbb{E}[\xi_s] = 0, \quad \mathbb{E}[\xi_s \xi_t] = \alpha_s \delta_{st}, \quad (6.4)$$

for $1 \leq s, t \leq d$. This means the coefficients $\{\xi_s\}$ are not correlated each other, and their covariances decrease as the index becomes larger. Therefore, a mapping $\Omega \ni \omega \mapsto (\xi_1(\omega), \xi_2(\omega), \dots)$ is called a *principal component analysis* (PCA), where only the first few principal terms are usually taken.

Let us give geometric interpretations for PCA by Theorem 6.1. Since the random variables $\{\xi_s\}$ are centered and independent as (6.4), $\{\xi_s\}$ consist of as eigenfunctions of the Laplacian given in Corollary 6.2. Hence, PCA is regarded as a special case of the Laplacian eigenmaps (5.9) or the embedding given in (5.8).

In addition, PCA is reformulated as a problem to maximize the left hand side of

$$\langle C_{\bar{X}}a, a \rangle_{\mathbb{R}^d} = \sum_{s,t=1}^d a_s a_t \mathcal{E}(X_s, X_t), \quad (6.5)$$

for $a = (a_1, a_2, \dots, a_d) \in \mathbb{R}^d$ under the condition $\|a\|_{\mathbb{R}^d} = 1$. Then the first principal term is written as $\xi_1 = \sum_{s=1}^d a_s X_s \in \mathcal{A}$. By contrast, the right hand side means the embedding energy of one dimensional subspace in \mathbb{R}^d given by $r = \sum_{s=1}^d a_s X_s \in \mathcal{A}$. This shows a relation between PCA and *Locality Preserving Projections* (LPP), which provides another embedding by *minimizing* (6.5) under the condition $\|r\|_{\mathcal{A}} = 1$ [19].

7 Application IV: many-body physics

In this section, we regard embedded points in \mathbb{R}^d as point mass in a classical many-body system. We show the force coincides with the curvature vector by taking special weights.

7.1 Hooke's law of spring

Let $\{\vec{r}_d(i) \mid i \in V\}$ be n points in \mathbb{R}^d having masses $\{m_i > 0\}$. We assume all pair of points $(\vec{r}_d(i), \vec{r}_d(j))$ is connected by a zero-length spring with force constant $k_{ij} \geq 0$. For simplicity, put $k_{ii} = 0$ for $i \in V$. In this case, Hooke's law states the force $\vec{F} \in \Gamma(V; \mathbb{R}^d)$ and the potential $U \in \mathbb{R}$ are respectively given as

$$\vec{F}(i) = \sum_{j \in V} k_{ij} (\vec{r}_d(j) - \vec{r}_d(i)), \quad U = \frac{1}{2} \sum_{i,j \in V} k_{ij} \|\vec{r}_d(i) - \vec{r}_d(j)\|_{\mathbb{R}^d}^2.$$

Theorem 7.1. *Take $w_{ij} = k_{ij}$ and $\mu_i = m_i$ for $i, j \in V$. Then we have*

$$\vec{F}(i) = m_i \vec{H}_d(i), \quad U = \mathcal{E}(\vec{r}_d, \vec{r}_d).$$

Proof. The assertion immediately follows from (2.15), (2.20), and (2.10). □

Its equations of motion is represented as

$$m_i \frac{d^2}{dt^2} \vec{r}_d(i) = \vec{F}(i) = -m_i \mathcal{L} \vec{r}_d(i).$$

Since the eigenfunctions of \mathcal{L} does not depend of \vec{r}_d , we can see the i -th Fourier coefficient has frequency ρ_i :

$$\frac{d^2}{dt^2}\mathcal{F}[\vec{r}_d]_i = \frac{d^2}{dt^2}\langle \vec{r}_d, v_i \rangle_{\mathcal{A}} = -\rho_i \langle \vec{r}_d, v_i \rangle_{\mathcal{A}} = -\rho_i \mathcal{F}[\vec{r}_d]_i.$$

If we fix some points $A \subset V$, the stable positions of other free points A^c are calculated by solving the Dirichlet problem

$$\begin{cases} \mathcal{L}\vec{r}(i) = 0, \text{ for } i \in A^c, \\ \vec{r}|_A = \vec{r}_d|_A. \end{cases} \quad (7.2)$$

This idea is used to obtain a smooth surface in point cloud processing [33]. The Dirichlet problem for a function is considered in random walks and machine learning, and applied to image processing [16] and semi-supervised learning in [37] respectively.

7.2 Newton's law of gravitation

Now, we assume $d \geq 3$. Let $\{\vec{r}_d(i) \mid i \in V\}$ be n points in \mathbb{R}^d as before. The gravitational potential $\Phi \in \mathcal{C}(\mathbb{R}^d)$ is considered to follow the Poisson equation in \mathbb{R}^d with a boundary condition at infinity, and its general solution gives

$$\Phi_i(\vec{r}) = -C \sum_{j \in V \setminus i} \frac{Gm_j}{\|\vec{r} - \vec{r}_d(j)\|_{\mathbb{R}^d}^{d-2}}.$$

Here, Φ_i means the gravitational potential around $\vec{r} = \vec{r}_d(i)$ caused by $\{\vec{r}_d(j) \mid j \in V \setminus i\}$. The corresponding gravitational field $\vec{g} \in \Gamma(V; \mathbb{R}^d)$ are given by $\vec{g}(i) = -\vec{\nabla}_{\mathbb{R}^d} \Phi_i|_{\vec{r}=\vec{r}_d(i)}$, hence we have

$$\vec{g}(i) = -C(d-2) \sum_{j \in V \setminus i} \frac{Gm_j(\vec{r}_d(i) - \vec{r}_d(j))}{\|\vec{r}_d(i) - \vec{r}_d(j)\|_{\mathbb{R}^d}^d}. \quad (7.3)$$

The gravitational potential energy $U \in \mathbb{R}$ is described as

$$U = \frac{1}{2} \sum_{i \in V} m_i \Phi_i(\vec{r}_d(i)) = -\frac{C}{2} \sum_{i,j \in V, i \neq j} \frac{Gm_i m_j}{\|\vec{r}_d(i) - \vec{r}_d(j)\|_{\mathbb{R}^d}^{d-2}}.$$

These physical concepts defined on \mathbb{R}^d are directly described as those on V .

Theorem 7.4. *Take $w_{ij} = CGm_i m_j / \|\vec{r}_d(i) - \vec{r}_d(j)\|_{\mathbb{R}^d}^d$ for $i, j \in V, i \neq j$ otherwise 0, and $\mu_i = m_i$. We have*

$$\vec{g}(i) = (d-2)\vec{H}_d(i), \quad U = -\mathcal{E}(\vec{r}_d, \vec{r}_d).$$

Proof. Under the assumption, for $i \in V$ we have

$$-(d-2)\mathcal{L}\vec{r}_d(i) = -(d-2) \sum_{j \in V \setminus i} \frac{w_{ij}}{m_i} (\vec{r}_d(i) - \vec{r}_d(j)) = \vec{g}(i),$$

$$\mathcal{E}(\vec{r}_d, \vec{r}_d) = \frac{1}{2} \sum_{i,j \in V, i \neq j} w_{ij} \|\vec{r}_d(i) - \vec{r}_d(j)\|_{\mathbb{R}^d}^2 = -U,$$

as required. \square

Remark that the above weights depend on positions \vec{r}_d unlike the case of Hooke's law. Hence, to compute the stable positions, we cannot use direct calculation like (7.2). In this situation, we can use an iterative method like (8.2) with variable weights, which is viewed as *the mean shift algorithm* [10] in machine learning.

In these case, weights are given as physical constants defining the system, and the curvature vector (2.20) indicates the force (up to constant) in a totally discrete manner, like general relativity in a continuous setting.

Remark 7.5. *From a similar viewpoint, the Coulomb potential is studied as weights of a kernel by [24], and their exponent is different from that in Theorem 7.4.*

8 Application V: signal processing

In this section, we review several harmonic techniques used in signal processing. Here, the Fourier transform and the curvature vector play important roles.

8.1 filtering

The convolution operator given in (2.18) is viewed to weight the Fourier coefficients of f by those of g . This can be generalized as a filtering on a frequency domain. In other words, the eigenvalues $\{\rho_i\}$ are understood as the frequencies, thus, with a continuous function $g: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$, we define a filtering operator $T_g := g(\mathcal{L}): \mathcal{A} \rightarrow \mathcal{A}$ by

$$T_g f := \sum_{i=1}^n \mathcal{F}[f]_i g(\rho_i) v_i = \sum_{i=1}^n g(\rho_i) \mathcal{F}[f]_i v_i. \quad (8.1)$$

The operator $Q_t = \exp(-t\mathcal{L}/c)$ given in Proposition 5.3 is also understood in this sense. In particular, a low-pass filter is realized by taking a decreasing function g such that $g(0) = 1$ and $\lim_{x \rightarrow \infty} g(x) = 0$. Practically, it is convenient to approximate g with the Chebyshev polynomials to avoid calculating the eigenfunctions [30, 18]. The filtering operator for an embedding \vec{r}_d is proposed as *the manifold harmonic transform* [32] for analyzing point clouds, which is described as $\vec{r}_3 \in \mathcal{A}^{\oplus 3}$. In order to obtain useful functions on a graph, the filtering operators (8.1) is modified to add learnable parameters in the graph deep learning

[5, 23]. In another context, the filtering operator is extended to define the graph wavelet transform [18].

8.2 smoothing

In order to construct useful filtering operators, let us perturb an embedding \vec{r}_d to decrease the embedding energy $\mathcal{E}(\vec{r}_d, \vec{r}_d)$.

$$\vec{r}_d^{(k+1)} = \vec{r}_d^{(k)} - \epsilon \mathcal{L} \vec{r}_d^{(k)} = \vec{r}_d^{(k)} + \epsilon \vec{H}_d^{(k)}. \quad (8.2)$$

This is seen as the discrete time heat equation (5.4), or the explicit Euler-scheme for the continuous time heat equation (5.5). From a geometric viewpoint, it corresponds to the mean curvature flow [4]. Here, we remark that weights $\{w_{ij}\}$ are usually fixed during the iterations even if they depend on the initial $\vec{r}_d^{(0)} = \vec{r}_d$.

Since iterations of (8.2) asymptotically lead to a constant embedding, which means just one point, it is important to prevent from shrinking. Some improving methods are proposed, for example, using the second ordered Laplacian combining with a growing up process [31], considering the implicit Euler-scheme alternatively [22], or both [11];

$$\vec{r}_d^{(k+1)} = \begin{cases} (\text{id} - \epsilon' \mathcal{L}) (\text{id} - \epsilon \mathcal{L}) \vec{r}_d^{(k)}, \\ \vec{r}_d^{(k)} - \epsilon \mathcal{L} \vec{r}_d^{(k+1)}, \\ \vec{r}_d^{(k)} - \epsilon \mathcal{L}^2 \vec{r}_d^{(k+1)}, \end{cases}$$

where $\epsilon > 0$ and $\epsilon + \epsilon' < 0$. They are described respectively as $g(t) = (1 - \epsilon't)(1 - \epsilon t)$, $(1 - \epsilon t)^{-1}$ and $(1 - \epsilon t^2)^{-1}$ in terms of (8.1), and then behave like low-path filters. Other geometric flows are also well studied in [36].

Nowadays, image filtering is also understood in this context:

$$f' = f - \frac{1}{c} \mathcal{L} f = S_c f$$

as in (5.1). The weights $\{\theta_{ij}\}$ are usually decided by pixel's location and values [26].

9 Application VI: manifold learning

In machine learning, dimension reduction techniques based on graphs are sometimes called *manifold leaning*. In this section, we review some of them from viewpoints of the graph cut problem and the energy minimization.

9.1 graph cut problem

The graph cut problem aims to find clusters in graph by minimizing several cutting loss functions [34]. They have a similar form to the isoperimetric constant (4.2) and are closely related with the eigenvalue problem as discussed in §4.

In this paper, we set the problem as minimization of the following function

$$\text{Loss}_{\text{GC}} \left(\{\chi_{A_l}\}_{l=1}^k \right) := \sum_{l=1}^k \frac{\mathcal{E}(\chi_{A_l}, \chi_{A_l})}{\|\chi_{A_l}\|_{\mathcal{A}}^2},$$

where $\chi_{A_l} = \sum_{i \in A_l} e_i$ and $\{A_l\}_{l=1}^k$ is a k -partition of V , which satisfies $\bigsqcup_{l=1}^k A_l = V$ (disjoint union) and $A_l \neq \emptyset$ for any l . By (5.2), we have

$$\langle \chi_{A_l}, S_c \chi_{A_l} \rangle_{\mathcal{A}} = \sum_{i=1}^n \lambda_i \langle v_i, \chi_{A_l} \rangle_{\mathcal{A}}^2 = \sum_{i=1}^n \lambda_i \iint_{A_l \times A_l} v_i(x) v_i(y) d\mu_x d\mu_y,$$

hence $\mathcal{L} = c \text{id} - c S_c$ and $\|v_i\|_{\mathcal{A}} = 1$ lead to

$$\sum_{l=1}^k \frac{\mathcal{E}(\chi_{A_l}, \chi_{A_l})}{\|\chi_{A_l}\|_{\mathcal{A}}^2} = \sum_{i=1}^n \rho_i + \sum_{l=1}^k \sum_{i=1}^n \frac{c \lambda_i}{2 \text{vol}(A_l)} \iint_{A_l \times A_l} (v_i(x) - v_i(y))^2 d\mu_x d\mu_y.$$

Furthermore, in general, we get

$$\begin{aligned} \frac{1}{2} \iint_{A \times A} (f(x) - f(y))^2 d\mu_x d\mu_y &= \frac{1}{2} \iint_{A \times A} ((f(x) - m_{f,A}) - (f(y) - m_{f,A}))^2 d\mu_x d\mu_y \\ &\quad + \iint_{A \times A} (f(x) - m_{f,A}) (f(y) - m_{f,A}) d\mu_x d\mu_y \\ &= \text{vol}(A) \int_A (f(x) - m_{f,A})^2 d\mu_x, \end{aligned}$$

for any $f \in \mathcal{A}$ and $m_{f,A} := \text{vol}(A)^{-1} \int_A f(x) d\mu_x \in \mathbb{R}$. Therefore, by taking large c , we can obtain the minimum partition by the k -means algorithm for

$$\left\{ \left(\sqrt{\lambda_1} v_1(x), \sqrt{\lambda_2} v_2(x), \dots, \sqrt{\lambda_n} v_n(x) \right) \in \mathbb{R}^n \mid x \in V \right\},$$

which is nothing but *the kernel k -means algorithm* [29]. This fact is first shown by [12]. On the other hand, the graph cut problem is often translated into the k -means on the Laplacian eigenmaps (5.9), which is known as *spectral clustering* [34].

In both cases, mapping methods play an essential role, and such methods are researched as dimension reduction. We already explained its examples; the commute time distance embedding (5.8), PCA (6.3), and LPP (6.5). LLE we discuss in the following subsection is also one of examples.

9.2 weight learning

The algorithms in the previous subsection are highly dependent on choice of weights $\{w_{ij}\}$, which determines eigenvalues and eigenfunctions. In order to avoid its trial-and-error process, several researches propose methods to learn weights. Their basic idea is to minimize an energy function with respect to weights under some assumptions.

In the Locally Linear Embedding (LLE) [28], weights are determined to minimize the energy function $\sum_{i \in V} \|\vec{r}_d(i) - \sum_{j \in V} \tilde{w}_{ij} \vec{r}_d(j)\|_{\mathbb{R}^d}^2$ under the condition $\sum_{j \in V} \tilde{w}_{ij} = 1$. For these weights $\{\tilde{w}_{ij}\}$, we can take $\mu_i = \deg(i) = 1$ for all $i \in V$. Hence, the energy function is equivalent to the length of the curvature vector for the embedding:

$$\sum_{i=1}^n \left\| \vec{r}_d(i) - \sum_{j \in V} \tilde{w}_{ij} \vec{r}_d(j) \right\|_{\mathbb{R}^d}^2 = \sum_{s=1}^d \sum_{i=1}^n (-(\mathcal{L}r_s)_i)^2 = \sum_{s=1}^d \|H_s\|_{\mathcal{A}}^2.$$

Besides, the LLE requires minimizing the embedding cost function $\sum_{i \in V} \|\vec{y}_{d'}(i) - \sum_{j \in V} \tilde{w}_{ij} \vec{y}_{d'}(j)\|_{\mathbb{R}^{d'}}^2 = \sum_{l=1}^{d'} \|\mathcal{L}y_l\|_{\mathcal{A}}^2$ for $d' < d$ under the condition $\langle y_l, y_{l'} \rangle_{\mathcal{A}} = \delta_{ll'}$, which gives the Laplacian eigenmaps (5.9) again.

10 Conclusion

We introduced a formulation based on a universal differential calculus and differential geometry, and explained several frameworks to analyze discrete points. These demonstrations would show our formulation has a potential to understand various discrete frameworks and develop new harmonic techniques by combining graph theory, probability theory, spectral geometry, and topological techniques [35].

Acknowledgement

The author would like to thank Satoshi Takahashi, Tetsuya Koike, Yosuke Otsubo, Chikara Nakamura for useful discussion and constant encouragement. He is also grateful to Bausan Yuan, Ping-Wei Chang, Shruthi Kubatur, Henry Chau and Pranav Gundewar for their advices.

References

- [1] Martin T Barlow. *Random walks and heat kernels on graphs*, volume 438. Cambridge University Press, 2017.
- [2] Mikhail Belkin and Partha Niyogi. Laplacian eigenmaps and spectral techniques for embedding and clustering. In *Advances in neural information processing systems*, pages 585–591, 2002.

- [3] Mikhail Belkin and Partha Niyogi. Towards a theoretical foundation for laplacian-based manifold methods. In *International Conference on Computational Learning Theory*, pages 486–500. Springer, 2005.
- [4] Kenneth A Brakke. *The Motion of a Surface by Its Mean Curvature*. Princeton University Press, 1978.
- [5] Joan Bruna, Wojciech Zaremba, Arthur Szlam, and Yann LeCun. Spectral networks and locally connected networks on graphs. *arXiv preprint arXiv:1312.6203*, 2013.
- [6] Jeff Cheeger. A lower bound for the smallest eigenvalue of the laplacian. In *Proceedings of the Princeton conference in honor of Professor S. Bochner*, pages 195–199, 1969.
- [7] Bang-Yen Chen and Leopold Verstraelen. Laplace transformations of submanifolds. *arXiv preprint arXiv:1307.1515*, 2013.
- [8] Fan RK Chung and Fan Chung Graham. *Spectral graph theory*. Number 92. American Mathematical Soc., 1997.
- [9] Ronald R Coifman and Stéphane Lafon. Diffusion maps. *Applied and computational harmonic analysis*, 21(1):5–30, 2006.
- [10] Dorin Comaniciu and Peter Meer. Mean shift: A robust approach toward feature space analysis. *IEEE Transactions on Pattern Analysis & Machine Intelligence*, (5):603–619, 2002.
- [11] Mathieu Desbrun, Mark Meyer, Peter Schröder, and Alan H Barr. Implicit fairing of irregular meshes using diffusion and curvature flow. In *Proceedings of the 26th annual conference on Computer graphics and interactive techniques*, pages 317–324. Citeseer, 1999.
- [12] Inderjit S Dhillon, Yuqiang Guan, and Brian Kulis. Kernel k-means: spectral clustering and normalized cuts. In *Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 551–556. ACM, 2004.
- [13] Aristophanes Dimakis and Folkert Müller-Hoissen. Discrete differential calculus: Graphs, topologies, and gauge theory. *Journal of Mathematical Physics*, 35(12):6703–6735, 1994.
- [14] Aristophanes Dimakis and Folkert Müller-Hoissen. Discrete riemannian geometry. *Journal of Mathematical Physics*, 40(3):1518–1548, 1999.
- [15] Francois Fouss, Alain Pirotte, Jean-Michel Renders, and Marco Saerens. Random-walk computation of similarities between nodes of a graph with application to collaborative recommendation. *IEEE Transactions on knowledge and data engineering*, 19(3):355–369, 2007.
- [16] Leo Grady. Random walks for image segmentation. *IEEE Transactions on Pattern Analysis & Machine Intelligence*, (11):1768–1783, 2006.
- [17] Alexander Grigor’yan, Yong Lin, Yuri Muranov, and Shing-Tung Yau. Cohomology of digraphs and (undirected) graphs. *Asian J. Math*, 19(5):887–931, 2015.
- [18] David K Hammond, Pierre Vandergheynst, and Rémi Gribonval. Wavelets on graphs via spectral graph theory. *Applied and Computational Harmonic Analysis*, 30(2):129–150, 2011.
- [19] Xiaofei He and Partha Niyogi. Locality preserving projections. In *Advances in neural information processing systems*, pages 153–160, 2004.

- [20] Matthias Hein, Jean-Yves Audibert, and Ulrike von Luxburg. Graph laplacians and their convergence on random neighborhood graphs. *Journal of Machine Learning Research*, 8(Jun):1325–1368, 2007.
- [21] Matthias Hein, Jean-Yves Audibert, and Ulrike Von Luxburg. From graphs to manifolds—weak and strong pointwise consistency of graph laplacians. In *International Conference on Computational Learning Theory*, pages 470–485. Springer, 2005.
- [22] Matthias Hein and Markus Maier. Manifold denoising. In *Advances in neural information processing systems*, pages 561–568, 2007.
- [23] Mikael Henaff, Joan Bruna, and Yann LeCun. Deep convolutional networks on graph-structured data. *arXiv preprint arXiv:1506.05163*, 2015.
- [24] Sepp Hochreiter, Michael C Mozer, and Klaus Obermayer. Coulomb classifiers: Generalizing support vector machines via an analogy to electrostatic systems. In *Advances in neural information processing systems*, pages 561–568, 2003.
- [25] Takashi Kumagai. *Random walks on disordered media and their scaling limits*. Springer, 2014.
- [26] Peyman Milanfar. A tour of modern image filtering: New insights and methods, both practical and theoretical. *IEEE signal processing magazine*, 30(1):106–128, 2012.
- [27] Bogdan Nica. A brief introduction to spectral graph theory. *arXiv preprint arXiv:1609.08072*, 2016.
- [28] Sam T Roweis and Lawrence K Saul. Nonlinear dimensionality reduction by locally linear embedding. *science*, 290(5500):2323–2326, 2000.
- [29] Bernhard Schölkopf, Alexander Smola, and Klaus-Robert Müller. Nonlinear component analysis as a kernel eigenvalue problem. *Neural computation*, 10(5):1299–1319, 1998.
- [30] David I Shuman, Pierre Vandergheynst, and Pascal Frossard. Chebyshev polynomial approximation for distributed signal processing. In *2011 International Conference on Distributed Computing in Sensor Systems and Workshops (DCOSS)*, pages 1–8. IEEE, 2011.
- [31] Gabriel Taubin. A signal processing approach to fair surface design. In *Proceedings of the 22nd annual conference on Computer graphics and interactive techniques*, pages 351–358. ACM, 1995.
- [32] Bruno Vallet and Bruno Lévy. Spectral geometry processing with manifold harmonics. In *Computer Graphics Forum*, volume 27, pages 251–260. Wiley Online Library, 2008.
- [33] Jörg Vollmer, Robert Mencl, and Heinrich Mueller. Improved laplacian smoothing of noisy surface meshes. In *Computer graphics forum*, volume 18, pages 131–138. Wiley Online Library, 1999.
- [34] Ulrike Von Luxburg. A tutorial on spectral clustering. *Statistics and computing*, 17(4):395–416, 2007.
- [35] Larry Wasserman. Topological data analysis. *Annual Review of Statistics and Its Application*, 5:501–532, 2018.
- [36] Guoliang Xu, Qing Pan, and Chandrajit L Bajaj. Discrete surface modelling using partial differential equations. *Computer Aided Geometric Design*, 23(2):125–145, 2006.

- [37] Xiaojin Zhu, Zoubin Ghahramani, and John D Lafferty. Semi-supervised learning using gaussian fields and harmonic functions. In *Proceedings of the 20th International conference on Machine learning (ICML-03)*, pages 912–919, 2003.