

A CONTRACTIVE HARDY–LITTLEWOOD INEQUALITY

ALEKSEI KULIKOV

ABSTRACT. We prove a contractive Hardy–Littlewood type inequality for functions from $H^p(\mathbb{T})$, $0 < p \leq 2$ which is sharp in the first two Taylor coefficients and asymptotically at infinity.

1. INTRODUCTION

The classical Hardy–Littlewood inequality [1] says that for $f(z) = a_0 + a_1z + \dots \in H^p(\mathbb{T})$, $0 < p \leq 2$, we have

$$(1.1) \quad \sum_{n=0}^{\infty} \frac{|a_n|^2}{(n+1)^{2/p-1}} \leq C_p \|f\|_p^2.$$

In [2] the following more precise version of this inequality was conjectured.

Conjecture 1.1. *For the function $f(z) = a_0 + a_1z + a_2z^2 + \dots \in H^p(\mathbb{T})$, $0 < p \leq 2$ we have*

$$(1.2) \quad \sum_{n=0}^{\infty} \frac{|a_n|^2}{c_{2/p}(n)} \leq \|f\|_p^2,$$

where $c_{\alpha}(n) = \binom{n+\alpha-1}{n}$.

Despite vast numerical evidence this conjecture is currently proved only for $p = \frac{2}{k}$, $k \in \mathbb{N}$ by Burbea [3], the case $p = 1$ being the famous Carleman inequality (see e.g. [4] for a simple self-contained proof).

In [5] inequality (1.2) was proved for the first two coefficients. Namely for the function $f \in H^p(\mathbb{T})$, $0 < p \leq 2$ we have $|f(0)|^2 + \frac{p}{2}|f'(0)|^2 \leq \|f\|_p^2$. In [6], by means of Wiessler's inequality [7], the authors proved the following strengthening of this result.

Theorem 1.2. *For the function $f(z) = a_0 + a_1z + a_2z^2 + \dots \in H^p(\mathbb{T})$, $0 < p \leq 2$ we have*

$$(1.3) \quad \sum_{n=0}^{\infty} \frac{|a_n|^2}{\Phi_{2/p}(n)} \leq \|f\|_p^2,$$

where $\Phi_{\alpha}(n) = c_{[\alpha]}(n) \left(\frac{\alpha}{[\alpha]} \right)^n$.

Note that $\Phi_\alpha(0) = c_\alpha(0) = 1, \Phi_\alpha(1) = c_\alpha(1) = \alpha$ but for $\alpha \notin \mathbb{N}$ these coefficients grow exponentially when n goes to infinity.

In this paper we prove the following theorem which gives us an inequality that is also sharp in the first two terms but for $n \geq 2$ the weight decays as in the Hardy–Littlewood inequality (1.1).

Theorem 1.3. *For each $0 < p \leq 2$ there exists $\varepsilon_p > 0$ such that for all $f \in H^p(\mathbb{T})$, $f(z) = a_0 + a_1z + a_2z^2 + \dots$ we have*

$$(1.4) \quad |a_0|^2 + \frac{p}{2}|a_1|^2 + \varepsilon_p \sum_{n=2}^{\infty} \frac{|a_n|^2}{(n+1)^{2/p-1}} \leq \|f\|_p^2.$$

Note that the constant $\frac{p}{2}$ is optimal as can be seen from the function $f(z) = 1 + \varepsilon z, \varepsilon \rightarrow 0$.

The proof of this inequality is based on the following theorem which may be of independent interest.

Theorem 1.4. *For $0 < p \leq 2$ there exists $C'_p < \infty$ such that for all $f \in H^p(\mathbb{T})$ we have*

$$(1.5) \quad \|f(z) - f(0) - f'(0)z\|_p^2 \leq C'_p(\|f\|_p^2 - |a_0|^2 - \frac{p}{2}|a_1|^2).$$

Since this theorem is obviously true for $p = 2$ we will prove it only for $0 < p < 2$. Moreover, the constants C'_p will be uniformly bounded except possibly for $0 < p < \varepsilon$ and $2 - \varepsilon < p < 2$. It is easy to see that in the former case nonuniformity is unavoidable but we do not know what happens when p is close to 2.

2. WEAK FORM OF THEOREM 1.4

In this section we will prove the following lemma.

Lemma 2.1. *For every $0 < p \leq 2$ there exists a constant γ_p such that for all $f \in H^p(\mathbb{T})$ we have*

$$(2.1) \quad \|f - f(0)\|_p \leq \gamma_p \sqrt{\|f\|_p^2 - |f(0)|^2}.$$

In [8, Lemma 2.2] this is proved for $p \leq 1$ and in [9] this is proved for $1 < p \leq 2$ (in [9] this lemma is proved even for $f \in L^p$, but with $\gamma_p \rightarrow \infty$ as $p \rightarrow 1$). Nevertheless we present here a simple uniform proof of this lemma.

Proof. Without loss of generality we may assume that $\|f\|_p = 1$. Let $n = \left\lfloor \frac{2}{p} \right\rfloor$, $\frac{1}{q} + \frac{n}{2} = \frac{1}{p}$. We can decompose the function f as a product $f = f_0 f_1 \dots f_n$, $f_0 \in H^q(\mathbb{T})$, $f_1, \dots, f_n \in H^2(\mathbb{T})$ such that $\|f_0\|_q = 1, \|f_k\|_2 = 1, k = 1, \dots, n$.

Let $f_k(z) = a_k + g_k(z)$, $g_k(0) = 0$. Note that $|a_k| \leq 1$, $\prod_{k=0}^n |a_k| = |f(0)|$. Therefore $|a_k| \geq |f(0)|$. By orthogonality we have $\|g_k\|_2 \leq \sqrt{1 - |f(0)|^2}$ and this inequality is valid even for $k = 0$ since $\|f_0\|_2 \leq \|f_0\|_q$.

We have the following formula for $f - f(0)$:

$$(2.2) \quad f - f(0) = g_n \left(\prod_{k=0}^{n-1} f_k \right) + g_{n-1} a_n \left(\prod_{k=0}^{n-2} f_k \right) + \dots + g_1 \left(\prod_{k=2}^n a_k \right) f_0 + g_0 \left(\prod_{k=1}^n a_k \right).$$

For each of the first n summands, by the obvious estimate $|a_k| \leq 1$ and Hölder's inequality, we have H^p -norm is bounded by $\sqrt{1 - |f(0)|^2}$. For the last summand we have $\prod_{k=1}^n |a_k| \leq 1$ and $\|g_0\|_p \leq \|g_0\|_2 \leq \sqrt{1 - |f(0)|^2}$. Therefore by the triangle inequality (with the possible additional constant coming from the fact that $H^p(\mathbb{T})$ for $p < 1$ is not a Banach space) we get $\|f - f(0)\|_p \leq \gamma_p \sqrt{1 - |f(0)|^2}$. $\square$

3. PROOF OF THEOREM 1.4 FOR FUNCTIONS WITHOUT ZEROES

In this section we will prove the following theorem.

Theorem 3.1. *Let $0 < p < 2$ and $f \in H^p(\mathbb{T})$ has no zeroes in $\mathbb{D}$. Then the conclusion of Theorem 1.4 holds for this function f .*

For the proof of this theorem we will need the following result from [8].

Theorem 3.2. *For $f \in H^p(\mathbb{T})$ with $\|f\|_p = 1$ we have*

$$(3.1) \quad |f'(0)| \leq \kappa(p) = \begin{cases} 1, & p \geq 1, \\ \sqrt{\frac{2}{p}} (1 - \frac{p}{2})^{1/p-1/2}, & 0 < p < 1. \end{cases}$$

Note that for all $0 < p < 2$ we have $\frac{p}{2} \kappa(p)^2 < 1$.

Proof of Theorem 3.1. Without loss of generality we may assume that $\|f\|_p = 1$, $f(z) = a_0 + a_1 z + \tilde{f}$. Note that $\|\tilde{f}\|_p \leq A_p$ for some absolute constant $A_p < \infty$.

We fix $\delta_p > 0$ to be determined later and consider the following cases depending on the values of $|a_0|$ and $|a_1|$:

- (i) $|a_0| < \delta_p$,
- (ii) $\delta_p \leq |a_0| \leq 1 - \delta_p$, $|a_1| < \delta_p$,
- (iii) $\delta_p \leq |a_0| \leq 1 - \delta_p$, $|a_1| \geq \delta_p$,
- (iv) $1 - \delta_p < |a_0|$.

In the first three cases we will prove that $\|f\|_p^2 - |a_0|^2 - \frac{p}{2}|a_1|^2$ is greater than some absolute constant $\lambda_p > 0$ from which, by the inequality $\|\tilde{f}\|_p \leq A_p$, the desired result follows.

In the first case we have $\|f\|_p^2 - |a_0|^2 - \frac{p}{2}|a_1|^2 \geq 1 - \delta_p^2 - \frac{p}{2}\kappa(p)^2$ which is positive if δ_p is small enough.

In the second case we have $\|f\|_p^2 - |a_0|^2 - \frac{p}{2}|a_1|^2 \geq 1 - (1 - \delta_p)^2 - \delta_p^2 = 2(\delta_p - \delta_p^2) > 0$.

For the third case we will essentially repeat the proof of Lemma 1 from [5]. We have $U(z) = f^{p/2}(z) = a_0^{p/2} + \frac{p}{2}a_0^{p/2-1}a_1z + \dots$ with $\|U\|_2 = 1$. Therefore

$$(3.2) \quad |a_0|^p + \frac{p^2}{4}|a_0|^{p-2}|a_1|^2 \leq 1.$$

On the other hand we have

$$(3.3) \quad \left(|a_0|^p + \frac{p^2}{4}|a_0|^{p-2}|a_1|^2\right)^{2/p} = |a_0|^2 \left(1 + \left(\frac{p|a_1|}{2|a_0|}\right)^2\right)^{2/p} > |a_0|^2 \left(1 + \frac{p|a_1|^2}{2|a_0|^2}\right),$$

where the last inequality is a Bernoulli's inequality $(1+t)^r > 1+tr$ for $r > 1, t > 0$. Since we are on a compact set $\delta_p \leq |a_0| \leq 1 - \delta_p$, $\delta_p \leq |a_1| \leq \kappa(p)$ and the functions are continuous we actually have a nonzero loss in the Bernoulli's inequality

$$(3.4) \quad |a_0|^2 \left(1 + \left(\frac{p|a_1|}{2|a_0|}\right)^2\right)^{2/p} \geq |a_0|^2 \left(1 + \frac{p|a_1|^2}{2|a_0|^2}\right) + \lambda_p$$

for some $\lambda_p > 0$. Therefore $1 \geq |a_0|^2 \left(1 + \frac{p|a_1|^2}{2|a_0|^2}\right) + \lambda_p = |a_0|^2 + \frac{p}{2}|a_1|^2 + \lambda_p$ as desired.

Now we turn to the fourth case which requires some additional ideas. Put $U(z) = f^{p/2}(z) = a_0^{p/2} + \frac{p}{2}a_0^{p/2-1}a_1z + \tilde{U}(z) \in H^2(\mathbb{T})$, $\|U\|_2 = 1$ (here we used that f has no zeroes).

Denote $|a_0|^2 = 1 - \beta^2$, $\|\tilde{U}\|_2 = \varepsilon$. Our goal now is to prove that $\|\tilde{f}\|_p \lesssim (\beta^2 + \varepsilon)$.

Consider $V(z) = U(z)(1 - \frac{p}{2a_0}a_1z)$. We have

$$(3.5) \quad V(z) = a_0^{p/2} - \frac{p^2 a_0^{p/2-2}}{4} a_1^2 z^2 + \tilde{U} - \frac{p}{2a_0} a_1 \tilde{U} z = a_0^{p/2} + \tilde{V}.$$

Note also that by orthogonality it is easy to see from $\|U\|_2 = 1$ that $|a_1|, \varepsilon \lesssim \beta$. Therefore we can bound $\|\tilde{V}\|_2 \lesssim \varepsilon + \beta^2$. Thus, by Pythagoras's Theorem we have

$$(3.6) \quad \|V\|_2 = \sqrt{|a_0|^p + \|\tilde{V}\|_2^2} \leq \sqrt{|a_0|^p + O(\varepsilon^2 + \beta^4)} = |a_0|^{p/2} + O(\varepsilon^2 + \beta^4).$$

We will now apply Lemma 2.1 to the function $V^{2/p}$ (V has no zeroes for small enough $\frac{|a_1|}{|a_0|}$, that is for small enough δ_p):

$$(3.7) \quad \|V^{2/p} - a_0\|_p \lesssim \sqrt{\|V\|_2^{4/p} - |a_0|^2} \leq \sqrt{|a_0|^2 + O(\varepsilon^2 + \beta^4) - |a_0|^2} = O(\beta^2 + \varepsilon).$$

Now we are going to connect $V^{2/p} - a_0$ and $\tilde{f}$:

$$\begin{aligned} V^{2/p} - a_0 &= U^{2/p} \left(1 - \frac{p}{2a_0} a_1 z\right)^{2/p} - a_0 = (a_0 + a_1 z + \tilde{f}) \left(1 - \frac{a_1}{a_0} z + O(\beta^2)\right) - a_0 = \\ &O(\beta^2) + \tilde{f} + \tilde{f}(a_1 z + O(\beta^2)) = \tilde{f} + O(\beta^2) + O(\beta) \tilde{f}. \end{aligned}$$

Therefore $\|\tilde{f}\| = O(\beta^2 + \varepsilon)(1 + O(\beta))^{-1} = O(\beta^2 + \varepsilon)$, as required.

Since $\|U\|_2 = 1$ we have

$$(3.8) \quad |a_0|^p + \frac{p^2}{4} |a_0|^{p-2} |a_1|^2 + \varepsilon^2 = 1.$$

Recall that in the end we want to prove that

$$(3.9) \quad |a_0|^2 + \frac{p}{2} |a_1|^2 + \varepsilon_p \|\tilde{f}\|_p^2 \leq 1.$$

By our bound for $\|\tilde{f}\|_p$ it is enough to prove that

$$(3.10) \quad |a_0|^2 + \frac{p}{2} |a_1|^2 + c_p(\beta^4 + \varepsilon^2) \leq 1$$

holds for some $c_p > 0$. Substituting the value of $|a_1|^2$ from (3.8) we get

$$(3.11) \quad |a_0|^2 + \frac{2}{p} |a_0|^{2-p} (1 - \varepsilon^2 - |a_0|^p) + c_p(\beta^4 + \varepsilon^2) \leq 1.$$

Choosing $c_p \leq \frac{2}{p}(1 - \delta_p)^{2-p}$ we can neglect terms with ε and we are left with the inequality

$$(3.12) \quad (1 - \beta^2) + \frac{2}{p} (1 - \beta^2)^{1-p/2} (1 - (1 - \beta^2)^{p/2}) + c_p \beta^4 \leq 1.$$

Expanding the left-hand side via Taylor's formula we get

$$(3.13) \quad 1 + \frac{p-2}{4} \beta^4 + c_p \beta^4 + O(\beta^6),$$

and it is smaller than 1 for $c_p < \frac{2-p}{4}$ and small enough β (that is small enough δ_p) since the constant in front of β^4 is negative. $\square$

4. PROOF OF THEOREM 1.4

In this section we will finish the proof of Theorem 1.4 by taking into consideration the potential zeroes of the function f .

Let $f \in H^p(\mathbb{T})$, $\|f\|_p = 1$. Write it as $f = Bg$, $\|g\|_p = 1$, g has no zeroes, $B = \prod_{n=1}^N \frac{z-w_n}{1-\bar{w}_n z}$ (obviously, it is enough to consider finite Blaschke products). Let $g(z) = a_0 + a_1 z + \tilde{g}(z)$, $B(z) = b_0 + b_1 z + \tilde{B}(z)$. We know that $|a_0|^2 + \frac{p}{2}|a_1|^2 + \varepsilon_p \|\tilde{g}\|_p^2 \leq 1$ and we want to prove the same bound for f (with possibly smaller ε_p).

Note that if $|f(0)| < \delta_p$ then as in the proof of Theorem 3.1 we can prove the desired inequality. Therefore we can assume that $|f(0)| \geq \delta_p$. Since $|f(0)| \leq |w_n|$ for all n we have that $|w_n| \geq \delta_p$.

Put $f_k(z) = g(z) \prod_{n=1}^k \frac{z-w_n}{1-\bar{w}_n z}$. Note that $|f_k(0)| \geq |f_N(0)| = |f(0)| \geq \delta_p$.

Carefully reading the proof of Lemma 1 from [5] we can see that each factor $\frac{z-w_k}{1-\bar{w}_k z}$ decreases $|f(0)|^2 + \frac{p}{2}|f'(0)|^2$ by at least $c_p(1 - |w_k|)$ for some $c_p > 0$, that is

$$(4.1) \quad |f_{k-1}(0)|^2 + \frac{p}{2}|f'_{k-1}(0)|^2 \geq |f_k(0)|^2 + \frac{p}{2}|f'_k(0)|^2 + c_p(1 - |w_k|)$$

(note that $c_p \rightarrow 0$ as $p \rightarrow 2$).

We have

$$(4.2) \quad |b_0| = \prod_{n=1}^N |w_n| = \exp\left(\sum_{n=1}^N \log |w_n|\right) \geq \exp(-C_p \sum_{n=1}^N (1 - |w_n|)),$$

where $C_p < \infty$ since all w_n are bounded away from 0. By orthogonality we have

$$(4.3) \quad \|\tilde{B}\|_p \leq \|\tilde{B}\|_2 \leq \sqrt{1 - |b_0|^2} \leq \sqrt{1 - \exp(-C_p \sum_{n=1}^N (1 - |w_n|))} \leq \sqrt{C_p \sum_{n=1}^N (1 - |w_n|)}.$$

Let us now write $f(z) - f(0) - f'(0)z$ in terms of B and g :

$$(4.4) \quad f(z) - f(0) - f'(0)z = b_1 a_1 z^2 + B(z) \tilde{g}(z) + \tilde{B}(z)(a_0 + a_1 z).$$

Since Blaschke products are unimodular we have $\|B\tilde{g}\|_p = \|\tilde{g}\|_p$. Since $|a_0| \leq 1$, $|a_1| \leq \kappa(p)$, the last term has H^p -norm at most $\alpha_p \|\tilde{B}\|_p$ for some $\alpha_p < \infty$. Finally, for b_1 we have again by orthogonality

$$(4.5) \quad |b_1| \leq \sqrt{1 - |b_0|^2} \leq \sqrt{C_p \sum_{n=1}^N (1 - |w_n|)}.$$

Collecting everything we get

$$(4.6) \quad \|f(z) - f(0) - f'(0)z\|_p \leq A_p \left(\|\tilde{g}\|_p + \sqrt{\sum_{n=1}^N (1 - |w_n|)} \right).$$

On the other hand by (4.1)

$$(4.7) \quad |f(0)|^2 + \frac{p}{2}|f'(0)|^2 \leq |a_0|^2 + \frac{p}{2}|a_1|^2 - c_p \sum_{n=1}^N (1 - |w_n|)$$

and by Theorem 3.1

$$(4.8) \quad |a_0|^2 + \frac{p}{2}|a_1|^2 + \varepsilon_p \|\tilde{g}\|_p^2 \leq 1.$$

Now it is easy to see from (4.6), (4.7), (4.8) and the trivial inequality $(x+y)^2 \leq 2x^2 + 2y^2$ that for some $\varepsilon'_p > 0$ we have

$$(4.9) \quad |f(0)|^2 + \frac{p}{2}|a_1|^2 + \varepsilon'_p \|f(z) - f(0) - f'(0)z\|_p^2 \leq 1,$$

as required.

5. PROOF OF THEOREM 1.3

In this section we will deduce Theorem 1.3 from Theorem 1.4.

We can rewrite inequality (1.1) as

$$(5.1) \quad \frac{1}{C_p} \sum_{n=0}^{\infty} \frac{|a_n|^2}{(n+1)^{2/p-1}} \leq \|f\|_p^2.$$

Applying this to the function $\tilde{f}(z) = f(z) - f(0) - f'(0)z$ we get

$$(5.2) \quad \frac{1}{C_p} \sum_{n=2}^{\infty} \frac{|a_n|^2}{(n+1)^{2/p-1}} \leq \|\tilde{f}\|_p^2.$$

Combining it with the bound from Theorem 1.4 we get

$$(5.3) \quad |a_0|^2 + \frac{p}{2}|a_1|^2 + \frac{1}{C_p C'_p} \sum_{n=2}^{\infty} \frac{|a_n|^2}{(n+1)^{2/p-1}} \leq \|f\|_p^2.$$

Acknowledgments. I would like to thank my advisor Kristian Seip for introducing me to this problem and for his constant support and encouragement, as well as for carefully reading this note and spotting some inaccuracies in the first version of it. I also would like to thank Dmitriy Stolyarov and Pavel Zatitskiy for fruitful discussions. This work was supported by Grant 275113 of the Research Council of Norway and by «Native towns», a social investment program of PJSC «Gazprom Neft».

REFERENCES

- [1] G. H. Hardy and J. E. Littlewood, “Some new properties of fourier constants,” *Math. Ann.*, vol. 97, no. 1, pp. 159–209, 1927.
- [2] O. F. Brevig, J. Ortega-Cerdà, K. Seip, and J. Zhao, “Contractive inequalities for Hardy spaces,” *Funct. Approx. Comment. Math.*, vol. 59, no. 1, pp. 41–56, 2018.
- [3] J. Burbea, “Sharp inequalities for holomorphic functions,” *Illinois J. Math.*, vol. 31, no. 2, pp. 248–264, 1987.
- [4] H. Helson, “Hankel forms and sums of random variables,” *Studia Math.*, vol. 176, no. 1, pp. 85–92, 2006.
- [5] A. Bondarenko, W. Heap, and K. Seip, “An inequality of Hardy-Littlewood type for Dirichlet polynomials,” *J. Number Theory*, vol. 150, pp. 191–205, 2015.
- [6] A. Bondarenko, O. F. Brevig, E. Saksman, K. Seip, and J. Zhao, “Pseudomoments of the Riemann zeta function,” *Bull. Lond. Math. Soc.*, vol. 50, no. 4, pp. 709–724, 2018.
- [7] F. B. Weissler, “Logarithmic Sobolev inequalities and hypercontractive estimates on the circle,” *J. Functional Analysis*, vol. 37, no. 2, pp. 218–234, 1980.
- [8] A. Bondarenko, O. F. Brevig, E. Saksman, and K. Seip, “Linear space properties of H^p spaces of Dirichlet series,” *Trans. Amer. Math. Soc.*, vol. 372, no. 9, pp. 6677–6702, 2019.
- [9] H. Hedenmalm, D. M. Stolyarov, V. I. Vasyunin, and P. B. Zatitskiy, “Sharpening Hölder’s inequality,” *J. Funct. Anal.*, vol. 275, no. 5, pp. 1280–1319, 2018.

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NO-7491 TRONDHEIM, NORWAY
 lyosha.kulikov@mail.ru