Orbits of Free Cyclic Submodules over Rings of Lower Triangular Matrices

Edyta Bartnicka

Abstract

Given a ring $T_n, n \ge 2$, of lower triangular $n \times n$ matrices with entries from an arbitrary field F, a complete description is performed of the orbits of free cyclic submodules of 2T_n , under the action of the general linear group $GL_2(T_n)$. It is given its total number, which is equal to the Bell number B_n , and there are shown their representatives.

Mathematics Subject Classification: 15B33, 16D40 Keywords: orbits of submodules — free cyclic submodules — rings of lower triangular matrices

1 Introduction

The projective line over a ring R with unity is a generalization of the concept of the projective line over a field known from the algebraic geometry. It can be defined as a set of those free cyclic submodules of the two-dimensional left free module ${}^{2}R$ which are generated by admissible or unimodular pairs. In general, there exist free cyclic submodules generated by non-unimodular pairs. The natural question arises about orbits of action of the general linear group $GL_{2}(R)$ on the set of all free cyclic submodules of ${}^{2}R$. In this note we investigate the case of a ring T_{n} of lower triangular $n \times n$ matrices

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ a_{21} & a_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

over a field F, where $n \ge 2$.

The main motivation for this paper is, presented by Havlicek and Saniga in [6], classification of the vectors of the free left module ${}^{2}T_{2}$ over a ring of ternions T_{2} up to the natural action of $GL_{2}(T_{2})$. It is shown that such classification yields exactly two (one unimodular and one non-unimodular) orbits of free cyclic submodules of ${}^{2}T_{2}$ under the action of $GL_{2}(T_{2})$. In case of any associative ring R with unity the unimodular $GL_{2}(R)$ -orbit is the projective line $\mathbb{P}(R)$ which is described and used in many articles (see e.g. [4, 2, 3]).

In [1] are described all orbits of free cyclic submodules in the ring T_3 under the action of $GL_2(T_3)$. Throughout the paper we aim at extending and generalisation these findings of [1] to an arbitrary ring T_n and we present the full classification of orbits of free cyclic submodules of 2T_n .

The starting point of our investigation is giving the complete characterisation of free cyclic submodules, unimodular pairs and outliers generating free cyclic submodules in case of considered class of rings, i.e. T_n . Next, we introduce representatives of all $GL_2(T_n)$ -orbits (Theorems 1 and 2). Finally, we prove that the total number of $GL_2(T_n)$ -orbits is equal to the Bell number B_n (Theorem 3).

2 Free cyclic submodules $T_n(A, B)$

Consider the free left module ${}^{2}R$ over a ring R with unity. Let $(a, b) \subset {}^{2}R$, then the set $R(a, b) = \{r(a, b); r \in R\}$ is a left cyclic submodule of ${}^{2}R$. If the equation (ra, rb) = (0, 0) implies that r = 0, then R(a, b) is free. We give now the condition for a cyclic submodule $M_n(A, B) \subset {}^{2}M_n$, where M_n is the ring of $n \times n$ matrices over F, to be free. To this aim we use the expression

$$[AB] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} & b_{11} & b_{12} & \dots & b_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} & b_{21} & b_{22} & \dots & b_{2n} \\ \vdots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} & b_{n1} & b_{n2} & \dots & b_{nn} \end{bmatrix}$$
for any $A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}, B = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ b_{n1} & b_{n2} & \dots & b_{nn} \end{bmatrix} \in M_n.$

Lemma 1. A cyclic submodule $M_n(A, B)$ is free if, and only if, the rank of [AB] is n.

Proof. Let

$$k_{j} = \begin{cases} \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \\ b_{1j} \\ b_{2j} \\ \vdots \\ b_{nj} \end{bmatrix}, j = 1, 2, \dots, n$$

and let $w_i = [x_{i1}x_{i2}\dots x_{in}], i = 1, 2, \dots, n.$

Then
$$[AB] = [k_1k_2...k_{2n}]$$
 and $X = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1n} \\ x_{21} & x_{22} & \dots & x_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ x_{n1} & x_{n2} & \dots & x_{nn} \end{bmatrix} = \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix}.$

" \Leftarrow " Suppose that the rank of [AB] is n. Equivalently exactly n column k_i of [AB] is a set of linearly independent vectors. Then the equation

m]

$$\begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix} [k_1 k_2 \dots k_{2n}] = 0,$$

(where 0 denotes the zero matrix of size $n \ge 2n$) implies that linear functionals $w_1, w_2, \ldots, w_n : F^n \to F$ are zeros on n linearly independent vectors of the space F^n . So they are zeros and hence $M_n(A, B)$ is free.

" \Rightarrow "Assume that a submodule $M_n(A, B)$ is free. Then the fact that any set of linear functionals $w_1, w_2, \ldots, w_n : F^n \to F$ are zeros on vectors k_1, k_2, \ldots, k_{2n} implies $w_1 = w_2 = \ldots = w_n = 0$. Consequently there exists a set of n linearly independent vectors k_i and so the rank of [AB] is n.

Remark 1. The statement of Lemma 1 remains true if one substitutes the ring M_n by any its subring, in particular by the ring T_n .

It is well known that any unimodular pair generates a free cyclic submodule, that is why we also call such submodules unimodular. Now we give the description of the unimodularity in case of T_n .

Remark 2. A pair $(A, B) \subset {}^{2}T_{n}$ is unimodular if, and only if, $a_{ii} \neq 0 \lor b_{ii} \neq 0$ for any i = 1, 2, ..., n.

Proof. According to the definition of the unimodularity (see [7, 5]) a pair $(A, B) \subset {}^{2}T_{n}$ is unimodular, if there exist matrices $X, Y \in T_{n}$ such that $AX + BY = U \in T_{n}^{*}$, where T_{n}^{*} denotes the group of invertible elements of the ring T_{n} .

$$AX + BY = \begin{bmatrix} a_{11}x_{11} + b_{11}y_{11} & 0 \\ & \ddots & \\ & * & a_{nn}x_{22} + b_{nn}y_{nn} \end{bmatrix}$$

for any $X, Y \in T_n(q)$. Thus $AX + BY = U \Leftrightarrow a_{ii}x_{ii} + b_{ii}y_{ii} \neq 0$ for any i = 1, 2, ..., n. This is equivalent to the following condition: $a_{ii}x_{ii} \neq -b_{ii}y_{ii}$ for any i = 1, 2, ..., n. Hence $(A, B) \subset {}^2T_n$ is unimodular if, and only if, $a_{ii} \neq 0 \lor b_{ii} \neq 0$ for any i = 1, 2, ..., n.

Another type of free cyclic submodules is the one represented by pairs not contained in any cyclic submodule generated by an unimodular pair, so called, *outliers* (as first introduced in [8]). As it was shown in [6, 1] the class of non-unimodular free cyclic submodules can be wide, although not all outliers generate free cyclic submodules.

Lemma 2. A pair $(A, B) \subset {}^{2}T_{n}$ is an outlier generating free cyclic submodule if, and only if, (A, B) is non-unimodular and rank[AB] = n.

Proof. According to [1, Theorem 1(2)] if a non-unimodular pair $(A, B) \subset {}^{2}T_{n}$ generates a free cyclic submodule, then (A, B) is an outlier. Lemma 1 and Remark 1 now yield to desired claim.

It follows then that there is no other free cyclic submodule $T_n(A, B)$ except generated by an unimodular pair or an outlier. Furthermore, a cyclic submodule $T_n(A, B)$ generated by an unimodular pair cannot have non-unimodular generators (see [1, Corollary 1]).

3 Representatives of $GL_2(T_n)$ -orbits

The general linear group $GL_2(T_n)$ acts in natural way (from the right) on the free left T_n -module. Orbits of the set of all free cyclic submodules $T_n(q)(A, B) \subset {}^2T_n(q)$ under the action of $GL_2(T_n)$ are called $GL_2(T_n)$ -orbits. Before we proceed to show the form of representatives of such orbits, we prove the following lemma.

Lemma 3. Let $X, Y, W, Z \in T_n$. The matrix $\begin{bmatrix} X & Y \\ W & Z \end{bmatrix}$ is invertible, i.e. is an element of the group $GL_2(T_n)$, if, and only if, $x_{ii}z_{ii} - y_{ii}w_{ii} \neq 0$ for all i = 1, 2, ..., n.

Proof. If X, Y, W, Z are elements of the ring T_n over F, then the matrix $\begin{bmatrix} X & Y \\ W & Z \end{bmatrix}$ can be identified with

$$V_n = \begin{bmatrix} x_{11} & 0 & \dots & 0 & y_{11} & 0 & \dots & 0 \\ x_{21} & x_{22} & \dots & 0 & y_{21} & y_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \dots & x_{nn} & y_{n1} & y_{n2} & \dots & y_{nn} \\ w_{11} & 0 & \dots & 0 & z_{11} & 0 & \dots & 0 \\ w_{21} & w_{22} & \dots & 0 & z_{21} & z_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ w_{n1} & w_{n2} & \dots & w_{nn} & z_{n1} & z_{n2} & \dots & z_{nn} \end{bmatrix},$$

where $x_{ij}, y_{ij}, w_{ij}, z_{ij} \in F$ for any i = 1, 2, ..., n; j = 1, 2, ..., i. Hence the matrix $\begin{bmatrix} X & Y \\ W & Z \end{bmatrix}$ is invertible if, and only if, $|V_n| \neq 0$. So, an equivalent formulation of the above lemma is now: $|V_n| \neq 0 \Leftrightarrow x_{ii}z_{ii} - y_{ii}w_{ii} \neq 0$ for all i = 1, 2, ..., n. It is easy to check that this equivalence is true for n = 2. Suppose now that it holds for any natural k > 2 and let k + 1 = l. Then

$$|V_l| = \begin{vmatrix} x_{11} & 0 & \dots & 0 & 0 & y_{11} & 0 & \dots & 0 & 0 \\ x_{21} & x_{22} & \dots & 0 & 0 & y_{21} & y_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ x_{k1} & x_{k2} & \dots & x_{kk} & 0 & y_{k1} & y_{k2} & \dots & y_{kk} & 0 \\ x_{l1} & x_{l2} & \dots & x_{lk} & x_{ll} & y_{l1} & y_{l2} & \dots & y_{lk} & y_{ll} \\ w_{11} & 0 & 0 & \dots & 0 & z_{11} & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ w_{k1} & w_{k2} & \dots & w_{kk} & 0 & z_{k1} & z_{k2} & \dots & z_{kk} & 0 \\ w_{l1} & w_{l2} & \dots & w_{lk} & w_{ll} & z_{l1} & z_{l2} & \dots & z_{lk} & z_{ll} \end{vmatrix} =$$

$$+w_{ll}(-1)^{3l} \begin{vmatrix} x_{11} & 0 & \dots & 0 & y_{11} & 0 & \dots & 0 & 0 \\ x_{21} & x_{22} & \dots & 0 & y_{21} & y_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ x_{k1} & x_{k2} & \dots & x_{kk} & y_{k1} & y_{k2} & \dots & y_{kk} & 0 \\ w_{11} & 0 & \dots & 0 & z_{11} & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ w_{k1} & w_{k2} & \dots & w_{kk} & z_{k1} & z_{k2} & \dots & z_{kk} & 0 \\ w_{l1} & w_{l2} & \dots & w_{lk} & z_{l1} & z_{l2} & \dots & z_{lk} & z_{ll} \end{vmatrix} + \\ +w_{ll}(-1)^{3l} \begin{vmatrix} x_{11} & 0 & \dots & 0 & y_{11} & 0 & \dots & 0 & 0 \\ x_{21} & x_{22} & \dots & 0 & y_{21} & y_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ x_{k1} & x_{k2} & \dots & x_{kk} & y_{k1} & y_{k2} & \dots & y_{lk} & 0 \\ x_{l1} & x_{l2} & \dots & 0 & z_{l1} & 0 & \dots & 0 & 0 \\ w_{21} & w_{22} & \dots & 0 & z_{21} & z_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ w_{k1} & w_{k2} & \dots & w_{kk} & z_{k1} & z_{k2} & \dots & z_{kk} & 0 \\ w_{l1} & w_{l2} & \dots & 0 & z_{l1} & 0 & \dots & 0 & 0 \\ w_{21} & w_{22} & \dots & 0 & z_{21} & z_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ w_{k1} & w_{k2} & \dots & w_{kk} & z_{k1} & z_{k2} & \dots & z_{kk} & 0 \end{vmatrix} = \\ = x_{ll} z_{ll}(-1)^{4l+2k} |V_{k}| + y_{ll} w_{ll}(-1)^{5l+k} |V_{k}| = (x_{ll} z_{ll} - y_{ll} w_{ll}) |V_{k}|.$$

By the assumption we get:

$$|V_{k+1}| \neq 0 \Leftrightarrow x_{(k+1)(k+1)}z_{(k+1)(k+1)} \neq y_{(k+1)(k+1)}w_{(k+1)(k+1)} \wedge x_{ii}z_{ii} \neq y_{ii}w_{ii}$$

for all $i = 1, 2, \dots, k$. Equivalently

$$|V_{k+1}| \neq 0 \Leftrightarrow x_{ii} z_{ii} \neq y_{ii} w_{ii}$$
 for all $i = 1, 2, \dots, k+1$.

Thus, the lemma holds by induction.

Let (A, B) be a pair of ${}^{2}T_{n}$ such that for all i = 1, 2, ..., n, j = 1, 2, ..., i - 1:

 \Box

- $a_{ii}, b_{ij} \in \{0, 1\},$
- $a_{ij} = b_{ii} = 0$,
- the number of non-zero entries of [AB] is equal to its rank and it is n.

We will use the grotesk type face for such pairs. According to Lemma 1 and Remark 1 any pair $(\mathtt{A},\mathtt{B})\in{}^{2}T_{n}$ generates free cyclic submodule. **Theorem 1.** Let $(\mathbf{A}, \mathbf{B}), (\mathbf{C}, \mathbf{D}) \in {}^{2}T_{n}$. $T_{n}(\mathbf{A}, \mathbf{B}), T_{n}(\mathbf{C}, \mathbf{D})$ are representatives of distinct $GL_{2}(T_{n})$ -orbits if, and only if, $(\mathbf{A}, \mathbf{B}) \neq (\mathbf{C}, \mathbf{D})$.

Proof. " \Rightarrow " This is obvious.

" \Leftarrow " We give the proof by contradiction. Assume that $T_n(\mathbf{A}, \mathbf{B}), T_n(\mathbf{C}, \mathbf{D})$ are in the same $GL_2(T_n)$ -orbit, i.e. there exists $\begin{bmatrix} X & Y \\ W & Z \end{bmatrix} \in GL_2(T_n)$ such that

$$(\mathbf{A}, \mathbf{B}) \begin{bmatrix} X & Y \\ W & Z \end{bmatrix} = (\mathbf{C}, \mathbf{D}).$$
 Hence

$$\begin{cases} a_{ii}x_{ii} = c_{ii} \\ a_{ii}y_{ii} = 0 \end{cases}$$

for any $i \in \{1, 2, ..., n\}$.

 $T_n(\mathbf{A}, \mathbf{B}), T_n(\mathbf{C}, \mathbf{D})$ are free, thus $a_{11}, c_{11} \in F^* = F \setminus \{0\}$, and so $a_{11} = c_{11} = 1$. Let $i \in \{2, 3, \dots, n\}$. Consider two cases:

- 1. $a_{ii} = 1$; Then by the above system of equations and according to Lemma 3 we get $y_{ii} = 0$, $x_{ii} \neq 0$ and $c_{ii} \neq 0$. Consequently $c_{ii} = a_{ii} = 1$, and so all remaining entries in the *i*-th row of matrices A, B, C, D are equal zero.
- 2. $a_{ii} = 0$; Then we get immediately $c_{ii} = a_{ii} = 0$. The form of $(\mathbf{A}, \mathbf{B}) \in {}^{2}T_{n}$ implies that there exists exactly one non-zero entry b_{ij} in the *i*-th row of the matrix **B**. Suppose then that $b_{ij} = 1$ for some $j \in \{1, 2, ..., i-1\}$. We obtain the system of equations:

$$\begin{cases} c_{ij} = 0 = a_{ii}x_{ij} + b_{ij}w_{jj} + b_{i(j+1)}w_{(j+1)j} + \dots + b_{i(i-1)}w_{(i-1)j} = b_{ij}w_{jj} \\ d_{ij} = a_{ii}y_{ij} + b_{ij}z_{jj} + b_{i(j+1)}z_{(j+1)j} + \dots + b_{i(i-1)}z_{(i-1)j} = b_{ij}z_{jj} \end{cases}$$

Therefore $w_{jj} = 0$. Lemma 3 leads to $z_{jj} \neq 0$, and so $d_{ij} \neq 0$. Thus $d_{ij} = b_{ij} = 1$ and all remaining entries in the *i*-th row of matrices A, B, C, D are equal zero.

It means that (A, B) = (C, D), which completes the proof. \Box

Theorem 2. Any $GL_2(T_n)$ -orbit has a unique representative $T_n(A, B) \subset {}^2T_n$.

Proof. Taking into account Theorem 1 it suffices to show that any $GL_2(T_n)$ -orbit has a representative of the form $T_n(\mathbf{A}, \mathbf{B}) \subset {}^2T_n$. This is equivalent to saying that for any free cyclic submodule $T_n(A, B) \subset {}^2T_n$ there exist matrices $Q \in GL_2(T_n)$ and $U \in T_n^*$ such that $U(A, B)Q = (\mathbf{A}, \mathbf{B})$. We show now that this holds.

Let $T_n(A, B)$ be any free cyclic submodule of 2T_n . We multiply $(A, B) \in {}^2T_n$ by $\begin{bmatrix} X & Y \\ W & Z \end{bmatrix} \in GL_2(T_n)$, with entries $y_{ii} = -a_{ii}^{-1}b_{ii}z_{ii}$, if $a_{ii} \neq 0$, and $z_{ii} = 0$, if $a_{ii} = 0, i = 1, 2, ..., n$. It gives

$$(A,B)\left[\begin{array}{cc} X & Y \\ W & Z \end{array}\right] = (C,D),$$

where $d_{ii} = 0$ for any i = 1, 2, ..., n. Next step is multiplying

$$P^{-1}(C,D) \begin{bmatrix} P & 0\\ 0 & I \end{bmatrix} = \left(P^{-1}CP, P^{-1}D\right) = (J,E),$$

where $P \in T_n$, J is a Jordan normal form of C with block matrices J_i , which are just elements of F. Of course, $e_{ii} = 0$ for any i = 1, 2, ..., n.

Let J' be the matrix obtained from J by replacement any element $J_i \neq 0$ by J_i^{-1} , and $J_i = 0$ by 1. The result of multiplication J'(J, E) = (J'J, J'E) is a pair (A, F), where, for any i = 1, 2, ..., n, $a_{ii} = \begin{cases} 1, J_i \neq 0 \\ 0, J_i = 0 \end{cases}$ and $f_{ii} = 0$.

We multiply again last pair (\mathbf{A}, F) by $\begin{bmatrix} I & -F \\ 0 & I \end{bmatrix}$ and we obtain a pair $(\mathbf{A}, -\mathbf{A}F + F) = (\mathbf{A}, G)$, where $g_{ij} = 0$ for any $i, j = 1, 2, \ldots, n$ such that $a_{ii} = 1$.

Suppose that $i_t, t \in T$ are the numbers of non-zero rows of matrix G, where $i_t < i_s$ if t < s. Let now

- 1. j_1 be the number of column of G such that $g_{i_1j_1} \neq 0$ and $g_{i_1k} = 0$ for any $k > j_1$;
- 2. j_t be the number of column of G such that $j_t \neq j_l$ for any $l = 1, 2, \ldots, j_t 1$, $g_{i_t j_t} \neq 0$ and $g_{i_t k} = 0$ for any k satisfying the two conditions: $k > j_t$ and $k \neq j_l$ for any $l = 1, 2, \ldots, j_t 1$.

Consider the multiplication $(\mathbf{A}, G) \begin{bmatrix} I & 0 \\ 0 & V \end{bmatrix}$, where V is an invertible matrix of T_n such that $v_{j_t j_t} = g_{i_t j_t}^{-1}$ if $j_t = i_t - 1$, in other case

 $v_{j_t j_t} = g_{i_t j_t}^{-1} \left(1 - g_{i_t (j_t+1)} v_{(j_t+1) j_t} - g_{i_t (j_t+2)} v_{(j_t+2) j_t} - \dots - g_{i_t (i_t-1)} v_{(i_t-1) j_t} \right),$ and $v_{j_t l} = -g_{i_t j_t}^{-1} \left(g_{i_t l} v_{ll} + g_{i_t (l+1)} v_{(l+1) l} + \dots + g_{i_t (j_t-1)} v_{(j_t-1) l} \right)$ if $j_t = i_t - 1$, in other case

$$v_{jtl} = -g_{i_t j_t}^{-1} \left[\left(g_{i_t l} v_{ll} + g_{i_t (l+1)} v_{(l+1)l} + \dots + g_{i_t (j_t - 1)} v_{(j_t - 1)l} \right) + \right]$$

+
$$(g_{i_t(j_t+1)}v_{(j_t+1)l} + g_{i_t(j_t+2)}v_{(j_t+2)l} + \dots + g_{i_t(i_t-1)}v_{(i_t-1)l})]$$

for any $l = 1, 2, \ldots, j_t - 1$.

It gives a pair (\mathbf{A}, H) , where for any pair of indexes $i_t, j_t, t \in T$ and for any $l = 1, 2, \ldots, j_t - 1$ we have: $h_{i_t j_t} = g_{i_t j_t} v_{j_t j_t} = g_{i_t j_t} g_{i_t j_t}^{-1} = 1$ if $j_t = i_t - 1$, in other case

$$h_{i_t j_t} = g_{i_t j_t} v_{j_t j_t} + g_{i_t (j_t+1)} v_{(j_t+1)j_t} + \dots + g_{i_t (i_t-1)} v_{(i_t-1)j_t} =$$

= $g_{i_t j_t} g_{i_t j_t}^{-1} \left(1 - g_{i_t (j_t+1)} v_{(j_t+1)j_t} - g_{i_t (j_t+2)} v_{(j_t+2)j_t} - \dots - g_{i_t (i_t-1)} v_{(i_t-1)j_t} \right) +$
 $+ g_{i_t (j_t+1)} v_{(j_t+1)j_t} + \dots + g_{i_t (i_t-1)} v_{(i_t-1)j_t} = 1,$

and

$$h_{itl} = g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(it-1)}v_{(it-1)l} =$$

$$= g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(jt-1)}v_{(jt-1)l} + g_{itjt}v_{jtl} =$$

$$= g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(jt-1)}v_{(jt-1)l} +$$

$$+ g_{itjt} \left[-g_{itjt}^{-1} \left(g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(jt-1)}v_{(jt-1)l} \right) \right] = 0$$

if $j_t = i_t - 1$, in other case

$$\begin{aligned} h_{itl} &= g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(it-1)}v_{(it-1)l} = \\ &= g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(jt-1)}v_{(jt-1)l} + \\ &+ g_{itjt}v_{jtl} + g_{it(jt+1)}v_{(jt+1)l} + \dots + g_{it(it-1)}v_{(it-1)l} = \\ &= g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(jt-1)}v_{(jt-1)l} + \\ &+ g_{itjt}\left\{ -g_{itjt}^{-1}\left[\left(g_{itl}v_{ll} + g_{it(l+1)}v_{(l+1)l} + \dots + g_{it(jt-1)}v_{(jt-1)l} \right) + \\ &+ \left(g_{it(jt+1)}v_{(jt+1)l} + g_{it(jt+2)}v_{(jt+2)l} + \dots + g_{it(it-1)}v_{(it-1)l} \right) \right] \right\} + \\ &+ g_{it(jt+1)}v_{(jt+1)l} + g_{it(jt+2)}v_{(jt+2)l} + \dots + g_{it(it-1)}v_{(it-1)l} = 0. \end{aligned}$$

Clearly, for any i = 1, 2, ..., n, j = 1, 2, ..., i - 1 we have: $h_{ii} = g_{ii}v_{ii} = 0$, and $h_{ij} = g_{ij}v_{jj} + g_{i(j+1)}v_{(j+1)j} + \cdots + g_{i(i-1)}v_{(i-1)j} = 0$ if $g_{ij} = 0$ for all j = 1, 2, ..., i - 1.

Let us illustrate this part of the proof by the example of a pair

where $g_{32}, g_{54}, g_{63}, g_{71} \in F^*$ First, we find indexes $i_t, j_t, t \in T$ and determine respective entries of $V \in T_7^*$:

- $i_1 = 3, j_1 = 2,$ $v_{22} = g_{32}^{-1},$ $v_{21} = -g_{32}^{-1}g_{31}v_{11};$
- $i_2 = 5, j_2 = 4,$ $v_{44} = g_{54}^{-1},$ $v_{41} = -g_{54}^{-1} (g_{51}v_{11} + g_{52}v_{21} + g_{53}v_{31}),$ $v_{42} = -g_{54}^{-1} (g_{52}v_{22} + g_{53}v_{32}),$ $v_{43} = -g_{54}^{-1} g_{53}v_{33};$
- $i_3 = 6, j_3 = 3,$ $v_{33} = g_{63}^{-1} (1 - g_{64} v_{43}),$ $v_{31} = -g_{63}^{-1} (g_{61} v_{11} + g_{62} v_{21} + g_{64} v_{41}),$ $v_{32} = -g_{63}^{-1} (g_{62} v_{22} + g_{64} v_{42});$
- $i_4 = 7, j_4 = 1,$ $v_{11} = g_{71}^{-1} \left(1 - g_{72} v_{21} - g_{73} v_{31} - g_{74} v_{41} \right).$

The result of multiplication $(\mathbf{A}, G) \begin{bmatrix} I & 0 \\ 0 & V \end{bmatrix}$ is a pair $(\mathbf{A}, H) \in {}^{2}T_{7}$, where: $h_{32} = g_{32}v_{22} = g_{32}g_{32}^{-1} = 1$, $h_{31} = g_{31}v_{11} + g_{32}v_{21} = g_{31}v_{11} + g_{32}\left(-g_{32}^{-1}g_{31}v_{11}\right) = 0$, $h_{54} = g_{54}v_{44} = g_{54}g_{54}^{-1} = 1$, $h_{51} = g_{51}v_{11} + g_{52}v_{21} + g_{53}v_{31} + g_{54}v_{41} = g_{51}v_{11} + g_{52}v_{21} + g_{53}v_{31} + g_{54}v_{42} = g_{52}v_{22} + g_{53}v_{32} + g_{54}v_{42} = g_{52}v_{22} + g_{53}v_{32} + g_{54}v_{42} = g_{52}v_{22} + g_{53}v_{32} + g_{54}\left[-g_{54}^{-1}\left(g_{51}v_{11} + g_{52}v_{21} + g_{53}v_{31}\right)\right] = 0$, $h_{52} = g_{52}v_{22} + g_{53}v_{32} + g_{54}v_{42} = g_{52}v_{22} + g_{53}v_{33} + g_{54}\left(-g_{54}^{-1}\left(g_{52}v_{22} + g_{53}v_{32}\right)\right)\right] = 0$, $h_{53} = g_{53}v_{33} + g_{54}v_{43} = g_{53}v_{33} + g_{54}\left(-g_{54}^{-1}g_{53}v_{33}\right) = 0$, $h_{63} = g_{63}v_{33} + g_{64}v_{43} = g_{63}g_{63}^{-1}\left(1 - g_{64}v_{43}\right) + g_{64}v_{43} = 1$, $h_{61} = g_{61}v_{11} + g_{62}v_{21} + g_{63}v_{31} + g_{64}v_{41} = g_{61}v_{11} + g_{62}v_{21} + g_{64}v_{42} = g_{62}v_{22} + g_{63}\left[-g_{63}^{-1}\left(g_{62}v_{22} + g_{64}v_{42}\right)\right] + g_{64}v_{42} = 0$, $h_{71} = g_{71}v_{11} + g_{72}v_{21} + g_{73}v_{31} + g_{74}v_{41} = g_{71}g_{71}^{-1}\left(1 - g_{72}v_{21} - g_{73}v_{31} - g_{74}v_{41}\right) + g_{72}v_{21} + g_{73}v_{31} + g_{74}v_{41} = 1$. As it was mentioned above $h_{ii} = g_{ii}v_{ii} = 0$ for any i = 1, 2, ..., 7 and $h_{21} = g_{21}v_{11} = 0$, $h_{4j} = g_{4j}v_{jj} + g_{4(j+1)}v_{(j+1)j} + \dots + g_{43}v_{3j} = 0$ for j = 1, 2, 3. Moreover, $h_{65} = g_{65}v_{55} = 0$, $h_{75} = g_{75}v_{55} + g_{76}v_{65} = 0$, $h_{76} = g_{76}v_{66} = 0$. Thus

•

Now we present the last part of the proof. Let K be a matrix of T_n^* such that $k_{ii} = 1$ for all i = 1, 2, ..., n and

$$k_{ii'} = -k_{i(i'+1)}h_{(i'+1)j} - k_{i(i'+2)}h_{(i'+2)j} - \dots - k_{ii}h_{ij}$$

if j-th column of H is of the form: $h_{i'j} = 1, h_{mj} = 0$ for all m < i' and $h_{ij} \neq 0$ for some i > i'. Then $K(\mathbf{A}, H) = (\mathbf{A}, L)$, where

$$l_{ij} = k_{i(j+1)}h_{(j+1)j} + k_{i(j+2)}h_{(j+2)j} + \dots + k_{ii}h_{ij} =$$

= $k_{ii'} + k_{i(i'+1)}h_{(i'+1)j} + k_{i(i'+2)}h_{(i'+2)j} + \dots + k_{ii}h_{ij} = 0.$

$$= \kappa_{ii'} + \kappa_{i(i'+1)}n_{(i'+1)j} + \kappa_{i(i'+2)}n_{(i'+2)j} + \dots + \kappa_{ii}n_{ij} =$$

Therefore $(\mathbf{A}, L) = (\mathbf{A}, \mathbf{B})$ what finishes the proof. Turning to the example we get

Corollary 1. The total number of $GL_2(T_n)$ -orbits is equal to the number of pairs $(\mathbf{A}, \mathbf{B}) \in {}^2T_n$.

Theorem 3. The total number of $GL_2(T_n)$ -orbits is equal to the Bell number B_n .

Proof. Recall that the Bell number B_n is the number of partitions Π of the set $\{1, 2, \ldots n\}$. Therefore, in the light of Corollary 1 it is enough to show that the set of pairs $(\mathbf{A}, \mathbf{B}) \in {}^2T_n$ and the family of partitions Π are equipotent. Let t_k be the rank of a matrix A_k obtained from $\mathbf{A} = (a_{kl})$ by removing its rows and columns number $k+1, k+2, \ldots, n$ and let $T = \{t_k; k = 1, 2, \ldots, n\}$. There exists a bijection that associates to every pair $(\mathbf{A}, \mathbf{B}) \in {}^2T_n$ the partition $\Pi = \{U_{t_k}, t_k \in T\}$ of the form

$$\Pi = \{\{k\} \cup \{i; b_{ij} = 1, t_k = j - rankB_{ij}\}; k \in \{1, 2, \dots, n\}, a_{kk} = 1\},\$$

where B_{ij} is the matrix obtained from $B = (b_{ij})$ by changing its rows number $i, i+1, \ldots, n$ and columns number $j, j+1, \ldots, n$ into zero rows and columns. For example, consider the pair

 $a_{kk} = 1$ for k = 1, 2, 4, so $\Pi = \{U_{t_1}, U_{t_2}, U_{t_4}\}$, where $t_1 = 1, t_2 = 2, t_4 = 3$. Of course, $1 \in U_1, 2 \in U_2, 4 \in U_3$. Moreover,

- $b_{32} = 1$, thus $j rankB_{32} = 2 0 = 2$ and hence $3 \in U_2$;
- $b_{54} = 1$, thus $j rankB_{54} = 4 1 = 3$ and hence $5 \in U_3$;
- $b_{63} = 1$, thus $j rankB_{63} = 3 1 = 2$ and hence $6 \in U_2$.

Therefore, the partition $\Pi = \{\{1\}, \{2, 3, 6\}, \{4, 5\}\}$ corresponds to such pair (A, B).

Conversely, to a partition $\Pi = \{U_s, s \in S\}$ of the set $\{1, 2, \ldots n\}$ corresponds exactly one pair $(\mathbf{A}, \mathbf{B}) \in {}^2T_n$. To show this correspondence suppose that u_s is the least element of the set U_s , where $s \in S$. An entry a_{ij} of $\mathbf{A} = (\mathbf{a_{ij}})$ is equal 1 if $i = j = u_s$ for some $s \in S$, and is equal 0 in other cases. Elements of the set

$$\{1, 2, \dots, n\} \setminus \{u_s, s \in S\} = \{x_m; m = 1, 2, \dots, n - rankA\},\$$

where $x_1 < x_2 < \cdots < x_{n-rankA}$, are numbers of rows with the entry 1 of the matrix B. Any other rows of B are zero rows. We have to indicate now the column number y_m containing the entry 1 from a row x_m for any $m = 1, 2, \ldots, n - rankA$. We do it for each row one by one, from the least x_m , i.e. x_1 , to the greatest, i.e. $x_{n-rankA}$. This order is crucial. For any $m = 1, 2, \ldots, n - rankA$ the column y_m is sth column without the entry 1 (from the left). Thereby, the entry 1 in x_1 th row is in sth column, where $x_1 \in U_s$.

Consider now the partition $\Pi = \{\{1\}, \{2, 3, 6\}, \{4, 5\}\}$ from the above example. $\{u_s, s \in S\} = \{1, 2, 4\}$, so $a_{11} = a_{22} = a_{44} = 1$ and all remaining entries of **A** are zero. $x_1 = 3, x_2 = 5, x_3 = 6$, hence rows 3th, 5th and 6th of **B** contain the entry 1. $3 \in U_2$, thus $b_{32} = 1$. $5 \in U_3$ and 3th column without the entry 1 (from the left) is now the column number 4, therefore $b_{54} = 1$. By the same way we get that $b_{63} = 1$ and all remaining entries of **B** are zero. Finally, the pair $(\mathbf{A}, \mathbf{B}) \in {}^2T_6$ corresponding to Π is of the form

Corollary 2. An orbit of unimodular submodules of ${}^{2}T_{n}$ represented by pair (A, B) = (1, 0) of ${}^{2}T_{n}$ corresponds to the partition $\Pi = \{\{1\}, \{2\}, \ldots, \{n\}\}$ of the set $\{1, 2, \ldots, n\}$.

Example 1. Consider the ring T_4 . By Theorem 3 there are exactly $B_4 = 15$ $GL_2(T_4)$ -orbits with the following representatives and corresponding partitions of the set $\{1, 2, 3, 4\}$:

$$\begin{split} T_4\left(\left[\begin{array}{cccc}1&0&0&0\\0&1&0&0\\0&0&0&0\end{array}\right], \left[\begin{array}{cccc}0&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right], \Pi = \{\{1\},\{2,3,4\}\}; \\ T_4\left(\left[\begin{array}{cccc}1&0&0&0\\0&0&0&0\\0&0&1&0\\0&0&0&0\end{array}\right], \left[\begin{array}{cccc}0&0&0&0\\1&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right], \left[\begin{array}{cccc}0&0&0&0\\1&0&0&0\\0&1&0&0\end{array}\right]\right), \Pi = \{\{1,2,4\},\{3\}\}; \\ T_4\left(\left[\begin{array}{cccc}1&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right], \left[\begin{array}{cccc}0&0&0&0\\1&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right], \left[\begin{array}{cccc}0&0&0&0\\1&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right]\right), \Pi = \{\{1,2\},\{3,4\}\}; \\ T_4\left(\left[\begin{array}{cccc}1&0&0&0\\0&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right], \left[\begin{array}{cccc}0&0&0&0\\1&0&0&0\\0&1&0&0\\0&0&0&0\end{array}\right]\right), \Pi = \{\{1,2,3\},\{4\}\}; \\ T_4\left(\left[\begin{array}{cccc}1&0&0&0\\0&0&0&0\\0&0&0&0\\0&0&0&0\end{array}\right], \left[\begin{array}{ccccc}0&0&0&0\\1&0&0&0\\0&0&0&0\end{array}\right]\right), \Pi = \{\{1,2,3\},\{4\}\}; \\ \end{array}\right]$$

References

- Bartnicka, E., Matraś, A.: 2016, Free Cyclic Submodules in the Context of the Projective Line, Results. Math., Vol. 70, pp. 567–580.
- [2] Bartnicka, E., Matraś, A.: 2017, The distant graph of the projective line over a finite ring with unity, Results. Math., Vol. 72, pp. 1943–1958.
- [3] Bartnicka, E., Saniga M.: 2020, Affine and projective planes linked with projective lines over certain rings of lower triangular matrices, Linear Algebra and its Applications, Vol. 586, pp. 377–409.
- [4] Blunck A., Havlicek H., Projective Representations I. Projective lines over rings, Abh. Math. Sem. Univ. Hamburg 70, 287-299 (2000)
- [5] Havlicek, H.: 2012, Divisible Designs, Laguerre Geometry, and Beyond, J. Math. Sci., New York, 186, 882-926.

- [6] Havlicek, H., and Saniga, M.: 2009, Vectors, Cyclic Submodules and Projective Spaces Linked with Ternions, Journal of Geometry, Vol. 92, Nos. 1-2, pp. 79–90.
- [7] Herzer, A.: 1995, Chain Geometries, In Buekenhout, F., editor, Handbook of Incidence Geometry, pp. 781–842, Elsevier, Amsterdam.
- [8] Saniga, M., Havlicek, H., Planat, M., and Pracna, P.: 2008, Twin 'Fano-Snowflakes' over the Smallest Ring of Ternions, Symmetry, Integrability and Geometry: Methods and Applications, Vol. 4, Paper 050, 7 pages.

Edyta Bartnicka University of Warmia and Mazury Faculty of Mathematics and Computer Science Słoneczna 54 Street, P-10710 Olsztyn Poland E-mail: edytabartnicka@wp.pl