

GEVREY SERIES SOLUTIONS IN ANALYTIC FUNCTIONS OF FIRST ORDER HOLOMORPHIC PDES

SERGIO A. CARRILLO AND CARLOS A. HURTADO

ABSTRACT. The goal of this work is to exhibit a Gevrey type, in an analytic function P , of formal power series solutions of some families of first order holomorphic PDEs. The approach is based on the classical majorant series technique by applying Nagumo norms joint with a division algorithm. Our main result recovers systematically many situations studied in the literature on the Gevrey type of formal solutions of these equations. We also provide a relation between Gevrey series in P and Gevrey series in several variables.

1. INTRODUCTION

In the study of ordinary and partial differential equations at irregular singular points or in the case of singular perturbation problems, a technique to obtain holomorphic solutions from formal ones is by applying certain summability methods such as Borel–summability and multisummability. These solutions represent asymptotically the formal power series solution as the variables approach the singular locus in adequate domains. In general, the first step to follow this method is to determine the existence, uniqueness and divergence rate (Gevrey order) of these series. The study of their summability is determined by the nature of the equation and it is a much harder problem. We refer to [12, 13, 20, 27, 31] for some examples of PDEs, including Navier-Stokes equation in \mathbb{R}^3 , which are susceptible to this type of analysis.

The goal of this paper is to provide a proof on the Gevrey type of formal power series solutions $\hat{\mathbf{y}}$ of holomorphic ordinary and partial differential equations of first order at a singular locus S . We will show that under a suitable geometric condition, the germ of analytic function P that generates S is the generic source of divergence: $\hat{\mathbf{y}}$ is 1–Gevrey in the germ P (P –1–Gevrey for short). Roughly speaking, this means that we can write $\hat{\mathbf{y}} = \sum_{n=0}^{\infty} y_n P^n$ as a power series in P , where the coefficients y_n are holomorphic in a common polydisc D at the origin and $\sup_{\mathbf{x} \in D} |y_n(\mathbf{x})| \leq CA^n n!$, for some constants $C, A > 0$. This notion was introduced recently by J. Mozo and R. Schäfke [23] in the framework of asymptotic expansions and summability with respect to a germ of an analytic function, and it generalizes the notion of Gevrey series in one variable.

More specifically, if $\mathbf{x} = (x_1, \dots, x_d) \in (\mathbb{C}^d, \mathbf{0})$ and $\mathbf{y} = (y_1, \dots, y_N) \in \mathbb{C}^N$, we consider a germ P of a non-zero holomorphic function on $(\mathbb{C}^d, \mathbf{0})$ such that $P(\mathbf{0}) = 0$, and the system of partial differential equations

$$(1) \quad P(\mathbf{x})L(\mathbf{y})(\mathbf{x}) = F(\mathbf{x}, \mathbf{y}), \quad \text{where } L := a_1(\mathbf{x})\partial_{x_1} + \dots + a_d(\mathbf{x})\partial_{x_d},$$

is a first order differential operator with holomorphic coefficients a_j near the origin –not all identically zero–, and F is a \mathbb{C}^N -valued holomorphic map defined in some neighborhood of $(\mathbf{0}, \mathbf{0}) \in \mathbb{C}^d \times \mathbb{C}^N$. The *singular locus* of (1) is the germ at the origin of the analytic set

$$S := \{\mathbf{x} \in (\mathbb{C}^d, \mathbf{0}) : P(\mathbf{x})a_j(\mathbf{x}) = 0, j = 1, \dots, d\},$$

Date: February 13, 2020.

2010 *Mathematics Subject Classification.* Primary 35F35, Secondary 35F05, 35C10, 34M25, 34M60.

The first author was supported by the projects “Análisis complejo, ecuaciones diferenciales y sumabilidad” (IN.BG.086.20.002 Univ. Sergio Arboleda) and “Álgebra y Geometría en Dinámica Real y Compleja III” (MTM2013-46337-C2-1-P Ministerio de Economía y Competitividad, Spain).

where the nature of equation (1) changes from differential to implicit one. Note that S contains the zero set of P , and they coincide if $a_j(\mathbf{0}) \neq 0$, for at least some index j . Furthermore, if $\frac{\partial F}{\partial \mathbf{y}}(\mathbf{0}, \mathbf{0})$ is an invertible matrix, P cannot be canceled from (1), so its zero set is a non-removable singular part of the equation. Under these conditions our main result can be stated as follows.

Theorem 1.1. *Consider the partial differential equation (1) where $F(\mathbf{0}, \mathbf{0}) = \mathbf{0}$, and $\mu := \frac{\partial F}{\partial \mathbf{y}}(\mathbf{0}, \mathbf{0})$ is an invertible matrix. If P divides $L(P)$, equation (1) has a unique formal power series solution $\hat{\mathbf{y}} \in \mathbb{C}[[\mathbf{x}]]^N$. Moreover, $\hat{\mathbf{y}}$ is a P -1-Gevrey series.*

Equation (1) falls into the category of singular first order holomorphic PDEs of the form

$$(2) \quad L_1(\mathbf{y})(\mathbf{x}) = F(\mathbf{x}, \mathbf{y}),$$

where $L_1 = \sum_{j=1}^d X_j(\mathbf{x})\partial_{x_j}$ is a germ of a holomorphic vector field, singular at $\mathbf{0} \in \mathbb{C}^d$, i.e., $X_j(\mathbf{0}) = 0$, for all $j = 1, \dots, d$. The convergence vs. rate of divergence of formal power series solutions of (2) has been studied extensively by several authors, see e.g., [25, 18, 14, 15, 32, 26]. These growth properties depend on conditions on $S = \{\mathbf{x} \in \mathbb{C}^d : X_j(\mathbf{x}) = 0, j = 1, \dots, d\}$ or on its associated ideal $(X_1, \dots, X_d) \subseteq \mathbb{C}\{\mathbf{x}\}$, and on non-resonance conditions on μ and the Jacobian matrix $\Lambda := (\partial_{x_i} X_j(\mathbf{0}))_{i,j}$ that we will explain below. Then, if S is an analytic submanifold, c.f. [18, 32], by choosing a suitable analytic coordinate system $\boldsymbol{\xi} = (\xi_1, \dots, \xi_d)$ of $(\mathbb{C}^d, \mathbf{0})$ where S is the zero set of some of these coordinates, and Λ is in canonical Jordan form, the convergence or a Gevrey type of solutions can be obtained. For instance, by means of a Newton polyhedron associated to L_1 that generalizes the Newton–Malgrange polygon which is familiar in the study of ODEs at irregular singular points.

Let us set $\text{Spec}(\mu) = \{\mu_1, \dots, \mu_d\}$ and $\text{Spec}(\Lambda) = \{\lambda_1, \dots, \lambda_m, 0, \dots, 0\}$, where $\mu_k \neq 0$, $\lambda_j \neq 0$, and all eigenvalues are repeated according multiplicity. If $m \geq 1$, the classical non-resonance *Poincaré condition* requests that

$$(3) \quad |\lambda_1\beta_1 + \dots + \lambda_m\beta_m - \mu_k| \geq \nu|\boldsymbol{\beta}|, \quad \text{for all } \boldsymbol{\beta} \in \mathbb{N}^m, k = 1, \dots, d,$$

for some constant $\nu > 0$. Then, if (3) is valid and $m = d$, i.e., Λ is invertible, the solution of (2) is convergent [25, 14, 15]. Otherwise, the solution is generically divergent, but of some Gevrey order in the variable $\boldsymbol{\xi}$, depending on the sizes of the blocks of the canonical Jordan form of Λ associated to the zero eigenvalue [14, 15].

It is worth remarking that Poincaré condition (3) is better known in the theory of normal forms [4, 26] or in the problem of existence of analytic invariant manifolds [9], both for holomorphic vector fields defined near a singular point. In particular, in the problem of their local analytic linearization where much more complicated non-resonance rules appear such as Siegel’s or Bruno’s ones.

Returning to our main problem, the linear part of $L_1 = P \cdot L$ in equation (1) can be highly degenerated and Λ is generically the zero matrix. In fact, $\Lambda = (a_i(\mathbf{0})p_j)_{i,j}$, where $p_j = \partial_{x_j} P(\mathbf{0})$. This is a very special type of matrix and its canonical Jordan form is the diagonal matrix $\text{diag}(\text{tr}(\Lambda), 0, \dots, 0)$, where

$$\text{tr}(\Lambda) = a_1(\mathbf{0})p_1 + \dots + a_d(\mathbf{0})p_d = L(P)(\mathbf{0}).$$

Thus, the only case in which $m \geq 1$, in fact, $m = 1$, is when $L(P)(\mathbf{0}) \neq 0$. Furthermore, Poincaré condition (3) is satisfied if and only if

$$\mu_k - nL(P)(\mathbf{0}) \neq 0, \quad \text{for all } n \in \mathbb{N}, k = 1, \dots, d.$$

In the aforementioned papers, our situation ($m = 0$ or 1) is covered in [14, Thm. 1.1], [15, Thm. 1.2] claiming the solution is $(1, \dots, 1)$ -Gevrey while working in the variable $\boldsymbol{\xi}$, see Section 4 for definitions. For the case $m = 0$, Theorem 1.1 improves the divergence rate of the formal solution by showing it is $(1/k, \dots, 1/k)$ -Gevrey, where $k = o(P)$ is the order of P , see Proposition 4.4. But more importantly, it identifies a possible variable to study summability phenomena. Finally, in

the case $m = 1$, $L(P)(\mathbf{0}) \neq 0$, the formal solution is actually convergent. In fact, by reordering the coordinates we can assume $a_1(\mathbf{0})p_1 \neq 0$, thus, $\xi_1 = P(\mathbf{x}), \xi_2 = x_2, \dots, \xi_d = x_d$ is a local change of variables in which our differential operator takes the form

$$P \cdot L = \xi_1 \cdot (U(\boldsymbol{\xi})\partial_{\xi_1} + \bar{a}_2(\boldsymbol{\xi})\partial_{\xi_2} + \dots + \bar{a}_d(\boldsymbol{\xi})\partial_{\xi_d}),$$

where $\bar{a}_j(\boldsymbol{\xi}) = a_j(\mathbf{x})$, and $U(\boldsymbol{\xi}) = \bar{a}_1\partial_{x_1}P + \dots + \bar{a}_d\partial_{x_d}P$ is a unit since $U(\mathbf{0}) = L(P)(\mathbf{0})$. Then, a standard majorant argument by working in the variable ξ_1 proves the convergence of the solution. We can also prove this by a slight modification of the proof of Theorem 1.1. In this way we find our second result.

Theorem 1.2. *Suppose the hypotheses of Theorem 1.1, but now assume $L(P)(\mathbf{0}) \neq 0$. Then, if $\mu - nL(P)(\mathbf{0})I_N$ is invertible, for all $n \in \mathbb{N}$, equation (1) has a unique analytic solution at the origin $\hat{\mathbf{y}} \in \mathbb{C}\{\mathbf{x}\}^N$.*

Theorem 1.1 has a general nature and recovers many examples of Gevrey type formal power solutions of ODEs and PDEs that have been treated in the literature. We would like to mention the following situations where it can be applied (see the beginning of Section 2 for notations):

- (a) Equation (1) includes the case of singularly perturbed and doubly singular ODEs by taking all a_j but one identically zero. Relabeling the variables, and under the previous hypotheses on F , we can consider systems of type

$$(4) \quad Q(\varepsilon)x^{k+1}\frac{\partial \mathbf{y}}{\partial x}(x, \varepsilon) = F(x, \varepsilon, \mathbf{y}),$$

where $x \in (\mathbb{C}, 0)$, $\varepsilon = (\varepsilon_1, \dots, \varepsilon_m) \in (\mathbb{C}^m, \mathbf{0})$, Q is analytic at the origin and $k \geq -1$ is an integer. In the regular case, i.e., $k = -1$ or 0 and $Q(\mathbf{0}) \neq 0$, if there exists a formal solution, it is convergent.

In the irregular case, if $Q(\mathbf{0}) \neq 0$, we can interpret ε as regular parameters and the classical theory establishes that the formal solution of (4) is $1/k$ -Gevrey in x , uniformly in ε [29]. In our setting, this means precisely that the solution is a x^{k-1} -Gevrey series. Equation (4) was studied by W. Balser and V. Kostov [3] for $m = 1$, $k = 0$, and by W. Balser and J. Mozo [2] for $m = 1$, $k \geq 1$, both when $Q(\varepsilon) = \varepsilon$ and in the linear case $F(x, \varepsilon, \mathbf{y}) = A(x, \varepsilon)\mathbf{y} - f(x, \varepsilon)$, proving the summability of the formal solution in the perturbation parameter ε , in adequate domains of x . On the other hand, M. Canalis-Durand, J.P. Ramis, R. Schäfke and Y. Sibuya [6] studied this equation when $m = 1$, $k = -1$ and $Q(\varepsilon) = \varepsilon^\sigma$, $\sigma \geq 1$ a positive integer. In particular, they showed that the solution is $1/\sigma$ -Gevrey in ε , uniformly in x . Later on, M. Canalis-Durand, J. Mozo and R. Schäfke [5] considered the case $m = 1$, $Q(\varepsilon) = \varepsilon^q$, and $k, q \geq 1$, and they proved the $\varepsilon^q x^{k-1}$ -summability of the formal power series solution and the singular directions are determined by the solutions of $\det(k\eta^q \xi^k I_N - \mu) = 0$, in the two-dimensional (ξ, η) -Borel space.

We can recover all these Gevrey type properties by applying Theorem 1.1 as follows:

- (i) If $k = -1$ and $Q(\mathbf{0}) = 0$, by choosing $P(x, \varepsilon) = Q(\varepsilon)$ and $L = \partial_x$, we have $L(P) = 0$. Thus, the solution is $Q(\varepsilon)$ -1-Gevrey.
- (ii) If $k \geq 0$, we take $P(x, \varepsilon) = x^k Q(\varepsilon)$ and $L = x\partial_x$, since $L(P) = kx^k Q(\varepsilon) = kP$. Thus, the solution is $x^k Q(\varepsilon)$ -1-Gevrey. If $Q(\mathbf{0}) \neq 0$, this means the solution is a x^{k-1} -Gevrey series.
- (b) Let ε and Q be as before, and assume $P(\mathbf{0}) = 0$. If $L = a_1(\mathbf{x}, \varepsilon)\partial_{x_1} + \dots + a_d(\mathbf{x}, \varepsilon)\partial_{x_d}$, the system of PDEs

$$(5) \quad Q(\varepsilon)P(\mathbf{x})L(\mathbf{y})(\mathbf{x}, \varepsilon) = F(\mathbf{x}, \varepsilon, \mathbf{y}),$$

can be seen as a singularly perturbed problem where the perturbation is given by Q if $Q(\mathbf{0}) = 0$. If P divides $L(P)$, then $L(QP) = QL(P)$ is divisible by QP and we can apply

Theorem 1.1 to conclude the system has a unique formal power series solution which is $Q(\varepsilon)P(\mathbf{x})$ -1-Gevrey. Particular situations are:

- (i) When $P \in \mathbb{C}[\mathbf{x}]$ is a quasi-homogeneous polynomial in \mathbf{x} , i.e., $P(t^{\lambda_1}x_1, \dots, t^{\lambda_d}x_d) = t^\lambda P(\mathbf{x})$, for some rational numbers $\lambda, \lambda_1, \dots, \lambda_d > 0$. Then, the operator

$$L_\lambda := \lambda_1 x_1 \partial_{x_1} + \dots + \lambda_d x_d \partial_{x_d},$$

satisfies $L_\lambda(P) = \lambda P$, and the solution of (5) is $Q(\varepsilon)P(\mathbf{x})$ -1-Gevrey.

- (ii) When $P(\mathbf{x}) = \mathbf{x}^\alpha$ is a monomial, $\alpha = (\alpha_1, \dots, \alpha_d)$ a tuple of non-negative integers, and $L = \sum_{j=1}^d b_j(\mathbf{x})x_j \partial_{x_j}$, with b_j holomorphic near the origin. Then $L(\mathbf{x}^\alpha) = \mathbf{x}^\alpha \sum_{j=1}^d \alpha_j b_j(\mathbf{x})$, thus the solution of (5) is $Q(\varepsilon)\mathbf{x}^\alpha$ -1-Gevrey. Another simple situation is $L = \frac{\partial P}{\partial x_i} \partial_{x_j} - \frac{\partial P}{\partial x_j} \partial_{x_i}$, $i \neq j$, since $L(P) = 0$.
- (iii) Families of PDEs with normal crossings given by

$$\varepsilon^{\alpha'} \mathbf{x}^\alpha L_\lambda(\mathbf{y})(\mathbf{x}, \varepsilon) = F(\mathbf{x}, \varepsilon, \mathbf{y}),$$

where L_λ and α are as before, and $\alpha' = (\alpha'_1, \dots, \alpha'_m)$ is another tuple of non-negative integers, but $\lambda = (\lambda_1, \dots, \lambda_d) \in (\mathbb{C} \setminus \{0\})^d$. Since $L_\lambda(\varepsilon^{\alpha'} \mathbf{x}^\alpha) = \langle \lambda, \alpha \rangle \varepsilon^{\alpha'} \mathbf{x}^\alpha$, where $\langle \lambda, \alpha \rangle := \lambda_1 \alpha_1 + \dots + \lambda_d \alpha_d$, we obtain a $\varepsilon^{\alpha'} \mathbf{x}^\alpha$ -1-Gevrey solution.

These equations have been studied by H. Yamazawa and M. Yoshino [33] in the case $m = 1$, $\alpha = \mathbf{0}$, $\mu = \text{diag}(\mu_1, \dots, \mu_d)$ a diagonal matrix and $\lambda_j, \text{Re}(\mu_k) > 0$, for all $j, k = 1, \dots, d$. In fact, the authors proved the 1-summability in $\varepsilon = \eta$ of the formal solution, uniformly in \mathbf{x} . In this trend, and assuming that λ has, up to a non-zero constant, positive entries, J. Mozo and the first author [7] studied these equations for the case $d = 2$ and $m = 0$ proving the solution is actually $x_1^{\alpha_1} x_2^{\alpha_2}$ -1-summable. Later on, this was generalized by the first author [10] for any $d \geq 2$ and m by using an adapted Borel-Laplace method: the formal solution is $\varepsilon^{\alpha'} \mathbf{x}^\alpha$ -1-summable and the singular directions are determined by the solutions of $\det(\langle \lambda, \alpha \rangle \xi^\alpha \eta^{\alpha'} I_N - \mu) = 0$, in the $(d + m)$ -dimensional (ξ, η) -Borel space.

- (iv) The family of scalar singular first-order linear PDEs of nilpotent type given by

$$(\alpha(x) + \beta(x, y))y \partial_x u + (a + b(x, y))y^2 \partial_y u + (1 + a(x, y))y u = f(x, y),$$

where $\alpha(0) \neq 0$ and $\beta(x, 0) \equiv b(x, 0) \equiv 0$. We obtain a unique y -1-Gevrey series solution by taking $P(x, y) = y$ and $L = (\alpha + \beta)\partial_x + (a + b)y\partial_y$. These equations were studied by M. Hibino [16, 17] proving the 1-summability in y , uniformly in x , under conditions on α , and on the analytic continuation and exponential growth of β, b, a and f .

We will give two more examples at the end of the paper, where after punctual blow-ups and ramifications we can apply Theorem 1.1. One in the setting of singular PDEs [21], Example 6.4, and the other in the framework of confluence of singularities of nonlinear ODEs [19], Example 6.5.

The technique to prove Theorems 1.1 and 1.2 is based on modified Nagumo norms for several variables, as introduced in [6], joint with a generalized Weierstrass division theorem that allows to write a power series as a series in the germ P , although the decomposition depends on the monomial order employed. Due to the compatibility of these tools, we can use the typical majorant series argument to establish the results.

The structure of the paper is as follows: Section 2 and 3 contain the technical parts of the work where we explain and develop the properties we will need on modified Nagumo norms, the Weierstrass division theorem and their compatibility. In Section 4 we recall the notions of (s, \dots, s) - and P - s -Gevrey series, $s \geq 0$, and we develop some properties relating them. Section 5 contains the proofs of Theorems 1.1 and 1.2, and also Corollary 5.1 which explains a simple extension for higher order systems. Finally, we include in Section 6 examples, including one where we show the hypotheses of Theorem 1.1 are necessary to conclude the desired Gevrey type.

2. NAGUMO NORMS

Let us start by fixing some notation: \mathbb{N} is the set of natural numbers including 0, $\mathbb{N}^+ = \mathbb{N} \setminus \{0\}$, and \mathbb{R}^+ is the set of positive real numbers. For a coordinate t , we will write $\frac{\partial}{\partial t} = \partial_t$ for the corresponding derivative. If $\boldsymbol{\beta} = (\beta_1, \dots, \beta_d) \in \mathbb{N}^d$, we use the multi-index notation $|\boldsymbol{\beta}| = \beta_1 + \dots + \beta_d$, $\boldsymbol{\beta}! = \beta_1! \cdots \beta_d!$, $\mathbf{x}^\beta = x_1^{\beta_1} \cdots x_d^{\beta_d}$ and $\frac{\partial^\beta}{\partial \mathbf{x}^\beta} = \frac{\partial^{|\beta|}}{\partial x_1^{\beta_1} \cdots \partial x_d^{\beta_d}}$.

Let $d \geq 1$ be an integer. We will work with $(\mathbb{C}^d, \mathbf{0})$ and local coordinates $\mathbf{x} = (x_1, \dots, x_d)$. We also write $\mathbf{x}' = (x_2, \dots, x_d)$ by removing the first coordinate. $\widehat{\mathcal{O}} = \mathbb{C}[[\mathbf{x}]]$ and $\mathcal{O} = \mathbb{C}\{\mathbf{x}\}$ denote the rings of formal and convergent power series in \mathbf{x} with complex coefficients, respectively. $\mathcal{O}^* = \{U \in \mathcal{O} : U(\mathbf{0}) \neq 0\}$ will denote the corresponding groups of units. Given $\hat{f} = \sum a_\beta \mathbf{x}^\beta \in \widehat{\mathcal{O}}$, $o(\hat{f})$ will denote its order: if $\hat{f} = \sum_{n=0}^{\infty} f_n$, $f_n = \sum_{|\beta|=n} a_\beta \mathbf{x}^\beta$, is written as sum of its homogeneous components, $o(\hat{f})$ is the least integer k such that $f_k \neq 0$.

For any $\mathbf{r} = (r_1, \dots, r_d) \in (\mathbb{R}^+)^d$, $D_{\mathbf{r}} = \{\mathbf{x} \in \mathbb{C}^d : |x_j| < r_j, j = 1, \dots, d\}$ is the polydisc centered at the origin with polyradius \mathbf{r} . If $r_j = r$, for all j , we write $D_{\mathbf{r}} = D_r^d$ as a Cartesian product instead. By using the norm $|\mathbf{x}| := \max_{1 \leq j \leq d} |x_j|$, we can write

$$D_r^d = \{\mathbf{x} \in \mathbb{C}^d : |\mathbf{x}| < r\}.$$

Also, $\mathcal{O}(D_{\mathbf{r}})$ and $\mathcal{O}_b(D_{\mathbf{r}})$ will be denote the sets of holomorphic and bounded holomorphic C -valued functions on the given polydisc. We denote by $J : \mathcal{O}(D_{\mathbf{r}})^N \rightarrow \mathcal{O}^N$ the Taylor map sending a vector function to its Taylor series at the origin.

Nagumo norms were introduced originally by M. Nagumo in [24] in his study of analytic partial differential equations. We will use a variant as it appears in [6] for the case of one complex variable. Let us fix two numbers $0 < \rho < r$ and consider the function

$$d_r(x) = \begin{cases} r - |x| & \text{if } |x| \geq \rho, \\ r - \rho & \text{if } |x| < \rho, \end{cases}$$

which satisfies

$$(6) \quad |d_r(x) - d_r(y)| \leq |x - y|, \quad x, y \in D_r.$$

The number ρ can be chosen arbitrarily but for our purposes we will choose always $\rho = r/2$.

Fix a polyradius $\mathbf{r} = (r_1, \dots, r_d)$. If $f \in \mathcal{O}(D_{\mathbf{r}})$ and $m \in \mathbb{N}$, we consider the family of *Nagumo norms*

$$(7) \quad \|f\|_m := \sup_{\mathbf{x} \in D_{\mathbf{r}}} |f(\mathbf{x})| d_{r_1}(x_1)^m \cdots d_{r_d}(x_d)^m.$$

These norms depend on \mathbf{r} , but to simplify notation we omit this dependence. There is no reason for these values to be finite, for instance, if $m = 0$ this norm reduced to the maximum norm. Note that if $\|f\|_k$ is finite and $m > k$, then

$$(8) \quad \|f\|_m \leq (r_1/2)^{m-k} \cdots (r_d/2)^{m-k} \|f\|_k.$$

In particular, if $k = 0$, i.e., if $f \in \mathcal{O}_b(D_{\mathbf{r}})$, all its Nagumo norms are finite.

We collect in the next proposition the main properties of these norms we will use in the proof of Theorems 1.1 and 1.2, including their behavior under the shift operators

$$(9) \quad S_j(f)(\mathbf{x}) = \begin{cases} (f(\mathbf{x}) - f(x_1, \dots, x_{j-1}, 0, x_{j+1}, \dots, x_d)) / x_j & \text{if } x_j \neq 0, \\ \frac{\partial f}{\partial x_j}(\mathbf{x}) & \text{if } x_j = 0. \end{cases}$$

Proposition 2.1. *Consider $m, k \in \mathbb{N}$ and $j = 1, \dots, d$. If $f, g \in \mathcal{O}(D_{\mathbf{r}})$, then*

- (1) $\|f + g\|_m \leq \|f\|_m + \|g\|_m$ and $\|fg\|_{m+k} \leq \|f\|_m \|g\|_k$.
- (2) $\left\| \frac{\partial f}{\partial x_j} \right\|_{m+1} \leq e(m+1) \prod_{i \neq j} (r_i/2) \|f\|_m$.
- (3) $\|S_j(f)\|_m \leq \frac{4}{r_j} \|f\|_m$.

Proof. The inequalities in (1) are clear from the definition. We prove (2) and (3) for the variable x_1 . If $\mathbf{x} = (x_1, \mathbf{x}') \in D_{\mathbf{r}}$, we have

$$(10) \quad |f(\mathbf{x})| d_{r_1}(x_1)^m \cdots d_{r_d}(x_d)^m \leq \|f\|_m.$$

To establish (2), we use Cauchy's formula

$$\left| \frac{\partial f}{\partial x_1}(\mathbf{x}) \right| = \left| \frac{1}{2\pi} \int_{|\xi - x_1| = R} \frac{f(\xi, \mathbf{x}')}{(\xi - x_1)^2} d\xi \right| \leq \frac{1}{R} \sup_{|\xi - x_1| = R} |f(\xi, \mathbf{x}')|,$$

valid for any $0 < R < r - |x_1|$. Note that if $|\xi - x_1| = R$, then $d_{r_1}(x_1) - R \leq d_{r_1}(\xi)$ by applying inequality (6). In particular, if $0 < R < d_{r_1}(x_1)$ it holds that

$$|f(\xi, \mathbf{x}')| \leq \|f\|_m d_{r_2}(x_2)^{-m} \cdots d_{r_d}(x_d)^{-m} (d_{r_1}(x_1) - R)^{-m}.$$

If we choose $R = \frac{d_{r_1}(x_1)}{m+1}$, we find that

$$\left| \frac{\partial f}{\partial x_1}(\mathbf{x}) \right| \leq \frac{(m+1)\|f\|_m}{d_{r_1}(x_1)^{m+1} d_{r_2}(x_2)^m \cdots d_{r_d}(x_d)^m} \left(1 + \frac{1}{m}\right)^m.$$

Therefore, by using the well-known inequality $(1 + 1/m)^m < e$ we conclude that

$$\left| \frac{\partial f}{\partial x_1}(\mathbf{x}) d_{r_1}(x_1)^{m+1} \cdots d_{r_d}(x_d)^{m+1} \right| \leq e(m+1) d_{r_2}(x_2) \cdots d_{r_d}(x_d) \|f\|_m \leq e(m+1) \left(\frac{r_2}{2} \cdots \frac{r_d}{2}\right) \|f\|_m,$$

as we wanted to show.

Finally, to prove (3), note that by inequality (10) we have

$$|f(0, \mathbf{x}')| \leq \|f\|_m (r_1/2)^{-m} d_{r_2}(x_2)^{-m} \cdots d_{r_d}(x_d)^{-m} \leq \|f\|_m d_{r_1}(x_1)^{-m} d_{r_2}(x_2)^{-m} \cdots d_{r_d}(x_d)^{-m},$$

for all $\mathbf{x} \in D_{\mathbf{r}}$. Hence, if $|x_1| \geq r_1/2$,

$$\left| \frac{f(x_1, \mathbf{x}') - f(0, \mathbf{x}')}{x_1} \right| \leq \frac{4}{r_1} \|f\|_m d_{r_1}(x_1)^{-m} \cdots d_{r_d}(x_d)^{-m}.$$

For $|x_1| < r_1/2$ we can use the maximum modulus principle and the above estimate to see that

$$|S_1(f)(\mathbf{x})| \leq \max_{|\xi| = r_1/2} |S_1(f)(\xi, \mathbf{x}')| \leq \frac{4}{r_1} \|f\|_m (r_1/2)^{-m} d_{r_2}(x_2)^{-m} \cdots d_{r_d}(x_d)^{-m}.$$

Since $d_{r_1}(x_1) = r_1/2$ if $|x_1| < r_1/2$, we find that in all cases that $|S_1(f)(\mathbf{x}) d_{r_1}(x_1)^m \cdots d_{r_d}(x_d)^m| \leq (4/r_1) \|f\|_m$ as required. \square

For vector-valued $\mathbf{y} = (y_1, \dots, y_N) \in \mathcal{O}(D_{\mathbf{r}})^N$, and matrix-valued $A = (A_{i,j}) \in \mathcal{O}(D_{\mathbf{r}})^{N \times N}$ maps we extend Nagumo norms by the rules

$$(11) \quad \|\mathbf{y}\|_m := \max_{1 \leq i \leq N} \|y_i\|_m, \quad \|A\|_m := \max_{1 \leq i \leq N} \sum_{j=1}^N \|A_{i,j}\|_m.$$

Then, it is immediate to check that

$$\|f \cdot \mathbf{y}\|_{m+k} \leq \|f\|_m \|\mathbf{y}\|_k, \quad \|A \cdot \mathbf{y}\|_{m+k} \leq \|A\|_m \|\mathbf{y}\|_k, \quad \|A \cdot B\|_{m+k} \leq \|A\|_m \|B\|_k,$$

for all $f \in \mathcal{O}(D_{\mathbf{r}})$, $\mathbf{y} \in \mathcal{O}(D_{\mathbf{r}})^N$, and $A, B \in \mathcal{O}(D_{\mathbf{r}})^{N \times N}$.

3. THE DIVISION ALGORITHM

We recall here a generalized Weierstrass division theorem by following closely [23], and whose original version is due to J.M. Aroca, H. Hironaka, and J. L. Vicente [1]. For the sake of completeness we include the proof for convergent series including the compatibility of the division algorithm with the Nagumo norms introduced in the previous section.

We will use the partial order \leq on \mathbb{N}^d defined by $\alpha \leq \beta$ if $\alpha_j \leq \beta_j$, for all $j = 1, \dots, d$. Thus $\alpha \not\leq \beta$ means there is an index j such that $\beta_j < \alpha_j$. We also use the notation

$$\Delta_\alpha := \left\{ \sum g_\beta \mathbf{x}^\beta \in \widehat{\mathcal{O}} : g_\beta = 0 \text{ if } \alpha \leq \beta \right\}.$$

Given $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d \setminus \{\mathbf{0}\}$, any power series $\hat{f} = \sum_{\beta \in \mathbb{N}^d} f_\beta \mathbf{x}^\beta \in \widehat{\mathcal{O}}$ can be written uniquely as a series in the monomial \mathbf{x}^α as

$$(12) \quad \hat{f} = \sum_{n=0}^{\infty} \hat{f}_{\alpha, n}(\mathbf{x}) \mathbf{x}^{n\alpha}, \quad \hat{f}_{\alpha, n}(\mathbf{x}) = \sum_{\alpha \not\leq \beta} f_{n\alpha + \beta} \mathbf{x}^\beta \in \Delta_\alpha.$$

This decomposition can be obtained by a repeated use of the canonical division algorithm by \mathbf{x}^α : given $\hat{f} \in \widehat{\mathcal{O}}$, there are unique $q \in \widehat{\mathcal{O}}$, $r \in \Delta_\alpha$ such that

$$\hat{f} = q\mathbf{x}^\alpha + r, \quad \text{where} \quad q = \sum_{\alpha \leq \beta} f_\beta \mathbf{x}^{\beta - \alpha}, \quad r = \sum_{\alpha \not\leq \beta} f_\beta \mathbf{x}^\beta.$$

Moreover, if $f \in \mathcal{O}(D_r)$, then $q, r \in \mathcal{O}(D_r)$. We can actually use the shift operators (9) introduced in the previous section to write

$$q = Q_\alpha(f) := S_1^{\alpha_1} \circ \dots \circ S_d^{\alpha_d}(f), \quad r = R_\alpha(f) := f - Q_\alpha(f) \cdot \mathbf{x}^\alpha.$$

In particular, Proposition 2.1 (3) shows that

$$(13) \quad \|Q_\alpha(f)\|_m \leq \frac{4^{|\alpha|}}{r^\alpha} \|f\|_m, \quad \|R_\alpha(f)\|_m \leq \|f\|_m + 4^{|\alpha|} \|f\|_m = (1 + 4^{|\alpha|}) \|f\|_m, \quad m \in \mathbb{N}.$$

By taking $m = 0$ we conclude that $Q_\alpha, R_\alpha : \mathcal{O}_b(D_r) \rightarrow \mathcal{O}_b(D_r)$ are linear continuous maps.

The generalized Weierstrass division allows to extend the previous considerations by dividing by a non-zero element of $\widehat{\mathcal{O}} \setminus \{0\}$ with zero constant term, but not in a canonical way. We will focus on division by an analytic germ $P \in \mathcal{O} \setminus \{0\}$, $P(\mathbf{0}) = 0$. The division is determined by P and an injective linear form $\ell : \mathbb{N}^d \rightarrow \mathbb{R}^+$, $\ell(\alpha) = \ell_1 \alpha_1 + \dots + \ell_d \alpha_d$ used to order the monomials by the rule

$$\mathbf{x}^\alpha <_\ell \mathbf{x}^\beta \quad \text{if} \quad \ell(\alpha) < \ell(\beta).$$

Then, any $\hat{f} = \sum f_\beta \mathbf{x}^\beta \in \widehat{\mathcal{O}} \setminus \{0\}$, has a *minimal exponent* $\nu_\ell(\hat{f})$ with respect to ℓ , i.e., $\nu_\ell(\hat{f}) = \alpha$ where $\mathbf{x}^\alpha = \min_\ell \{\mathbf{x}^\beta : f_\beta \neq 0\}$, and the minimum is taken according to $<_\ell$. The division process, for formal and convergent series, can be stated as follows, c.f. [23, Lemmas 2.4, 2.6].

Proposition 3.1 (Generalized Weierstrass Division). *Let P and ℓ as above. For every $\hat{g} \in \widehat{\mathcal{O}}$, there are unique $\hat{q} \in \widehat{\mathcal{O}}$, $\hat{r} \in \Delta_{\nu_\ell(P)}$ such that*

$$\hat{g} = \hat{q}P + \hat{r}.$$

Moreover, if $\rho > 0$ is sufficiently small, then for every $g \in \mathcal{O}_b(D_{\rho(\ell)})$, $\rho(\ell) = (\rho^{\ell_1}, \dots, \rho^{\ell_d})$, there are unique $q \in \mathcal{O}_b(D_{\rho(\ell)})$, $r \in \mathcal{O}_b(D_{\rho(\ell)})$ with $J(r) \in \Delta_{\nu_\ell(P)}$ such that

$$g = qP + r, \quad Q_{P, \ell}(g) := q, \quad R_{P, \ell}(g) := r.$$

The corresponding operators $Q_{P, \ell}, R_{P, \ell} : \mathcal{O}_b(D_{\rho(\ell)}) \rightarrow \mathcal{O}_b(D_{\rho(\ell)})$ are linear and continuous. In fact, if ρ is sufficiently small, then

$$\|Q_{P, \ell}(g)\|_m \leq \frac{2 \cdot 4^{|\nu_\ell(P)|}}{\rho^{\ell(\nu_\ell(P))}} \|g\|_m, \quad \|R_{P, \ell}(g)\|_m \leq 2(1 + 4^{|\nu_\ell(P)|}) \|g\|_m \quad \text{for all } m \in \mathbb{N}.$$

Proof. By the choice of the polyradius $\rho(\ell)$, we have $|\mathbf{x}^\beta| \leq \rho^{\ell(\beta)}$ if $\mathbf{x} \in D_{\rho(\ell)}$. Let us write $\boldsymbol{\alpha} = \nu_\ell(P)$. Without loss of generality we can assume $P = \mathbf{x}^\alpha + \tilde{P}$, where $\tilde{P} \in \mathcal{O} \setminus \{0\}$ and $\nu_\ell(\tilde{P}) >_\ell \mathbf{x}^\alpha$. Then, solving $g = qP + r$ or $q\mathbf{x}^\alpha + r = g - q\tilde{P}$ for q and r is equivalent to find a fixed point for the equation

$$(14) \quad q = Q_\alpha(g - q\tilde{P}),$$

joint with $r = R_\alpha(g - q\tilde{P})$. If ρ is sufficiently small, we can choose a constant $K > 0$ such that $\|\tilde{P}\|_0 \leq K\rho^{\ell(\nu_\ell(\tilde{P}))}$. Consider the map $\phi_g : \mathcal{O}_b(D_{\rho(\ell)}) \rightarrow \mathcal{O}_b(D_{\rho(\ell)})$ given by $\phi_g(h) = Q_\alpha(g - h\tilde{P})$. By using the first inequality in (13) for $m = 0$ we see that

$$\|\phi_g(h_1) - \phi_g(h_2)\|_0 = \left\| Q_\alpha((h_1 - h_2)\tilde{P}) \right\|_0 \leq 4^{|\alpha|} K \rho^{\ell(\nu_\ell(\tilde{P})) - \ell(\alpha)} \|h_1 - h_2\|_0.$$

Thus, ϕ_g defines a contraction if ρ is small enough, i.e., if $4^{|\alpha|} K \rho^{\ell(\nu_\ell(\tilde{P})) - \ell(\alpha)} < 1$, and it has a unique fixed point q . This determines the existence and uniqueness of q and r .

Finally, from equation (14) we find that

$$\|Q_{P,\ell}(g)\|_m \leq \frac{4^{|\alpha|} / \rho^{\ell(\alpha)}}{1 - 4^{|\alpha|} K \rho^{\ell(\nu_\ell(\tilde{P})) - \ell(\alpha)}} \|g\|_m,$$

and from $r = R_\alpha(g - q\tilde{P})$, that

$$\|R_{P,\ell}(g)\|_m \leq (1 + 4^{|\alpha|}) \left(1 + \frac{\|\tilde{P}\|_0 4^{|\alpha|} / \rho^{\ell(\alpha)}}{1 - 4^{|\alpha|} K \rho^{\ell(\nu_\ell(\tilde{P})) - \ell(\alpha)}} \right) \|g\|_m = \frac{1 + 4^{|\alpha|}}{1 - 4^{|\alpha|} K \rho^{\ell(\nu_\ell(\tilde{P})) - \ell(\alpha)}} \|g\|_m.$$

Therefore the result follows by taking additionally $\rho > 0$ such that $4^{|\alpha|} K \rho^{\ell(\nu_\ell(\tilde{P})) - \ell(\alpha)} < 1/2$. \square

By a repeated application of the previous proposition [23, Coro. 2.5], any $\hat{f} \in \hat{\mathcal{O}}$ can be written uniquely as

$$(15) \quad \hat{f} = \sum_{n=0}^{\infty} \hat{f}_{P,\ell,n} P^n, \quad \hat{f}_{P,\ell,n} \in \Delta_{\nu_\ell(P)}.$$

For the convergent case, we have a similar result that we state in the following corollary, c.f. [23, Coro. 2.7].

Corollary 3.2. *If $s > 0$ is such that the operators $Q_{P,\ell}$ and $R_{P,\ell}$ are defined over $\mathcal{O}_b(D_{s(\ell)})$, there is $r = r(s) > 0$, depending only on s , such that for any $f \in \mathcal{O}_b(D_{s(\ell)})$ we can find a unique sequence $(f_n)_{n \in \mathbb{N}} \subset \mathcal{O}_b(D_r^d)$ with $J(f_n) \in \Delta_{\nu_\ell(P)}$, such that*

$$(16) \quad f = \sum_{n=0}^{\infty} f_n P^n, \quad f_n = R_{P,\ell} \circ Q_{P,\ell}^n(f),$$

both series being convergent for $|\mathbf{x}| < r$.

Proof. By applying the preceding lemma we obtain

$$f = \sum_{n=0}^{N-1} R_{P,\ell}(Q_{P,\ell}^n(f)) P^n + Q^N(f) P^N, \quad \text{for all } N \in \mathbb{N}.$$

If we choose $0 < r \leq s$ such that $M = \sup_{|\mathbf{x}| < r} |P(\mathbf{x})| < s^{\ell(\nu_\ell(P))} / 2 \cdot 4^{|\nu_\ell(P)|} = 1/b$, then we can estimate

$$\sup_{|\mathbf{x}| < r} \left| f(\mathbf{x}) - \sum_{n=0}^{N-1} R_{P,\ell}(Q_{P,\ell}^n(f))(\mathbf{x}) P(\mathbf{x})^n \right| \leq (bM)^N \sup_{\mathbf{y} \in D_{s(\ell)}} |f(\mathbf{y})|.$$

The result follows by taking $N \rightarrow +\infty$. \square

To finish this section we would like to remark that if $\hat{f} = \sum f_n P^n$ and $\hat{g} = \sum g_n P^n$, where $f_n, g_n \in \mathcal{O}_b(D_r^d)$ for a common r , decomposition (15) for their product is given by

$$(17) \quad \begin{aligned} \hat{f} \cdot \hat{g} &= \sum_{k=0}^{\infty} \sum_{j=0}^k f_j g_{k-j} P^k = \sum_{k,m=0}^{\infty} \sum_{j=0}^k R_{P,\ell}(Q_{P,\ell}^m(f_j g_{k-j})) P^{k+m} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \sum_{j=0}^k R_{P,\ell}(Q_{P,\ell}^{n-k}(f_j g_{k-j})) \right) P^n. \end{aligned}$$

Similar formulas hold for the product of more than two series.

4. GEVREY SERIES

If $\mathbf{s} = (s_1, \dots, s_d) \in \mathbb{R}_{\geq 0}^d$ and $\hat{f} = \sum_{\beta \in \mathbb{N}^d} a_{\beta} \mathbf{x}^{\beta} \in \widehat{\mathcal{O}}$, we say that \hat{f} is a \mathbf{s} -Gevrey series if we can find constants $C, A > 0$ such that $|a_{\beta}| \leq CA^{|\beta|} |\beta|!^{\mathbf{s}}$, for all $\beta \in \mathbb{N}^d$. Note that $\mathbf{s} = \mathbf{0}$ means convergence. We will be interested in the case $s_1 = \dots = s_d = s > 0$. Thanks to the inequalities

$$\beta! \leq |\beta|! \leq d^{|\beta|} \beta!,$$

a series \hat{f} is (s, \dots, s) -Gevrey if and only if there are constants $C, A > 0$ such that

$$|a_{\beta}| \leq CA^{|\beta|} |\beta|!^s, \quad \beta \in \mathbb{N}^d.$$

We denote by $\widehat{\mathcal{O}}_s$ the set of (s, \dots, s) -Gevrey series.

Remark 4.1. For any $s \geq 0$, $\widehat{\mathcal{O}}_s$ is closed under sums, products, partial derivatives, composition, and it contains \mathcal{O} . This can be seen as a consequence of a more general situation in the setting of ultradifferentiable functions. In that framework the Gevrey sequence $(n!^s)_{n \in \mathbb{N}}$ is generalized by a sequence of positive numbers $(M_n)_{n \in \mathbb{N}}$ satisfying log-convexity ($M_n^2 \leq M_{n-1} M_{n+1}$), stability under derivatives ($M_{n+1} \leq K^n M_n$ for some $K > 0$) and the condition $M_n^{1/n} \rightarrow +\infty$ as $n \rightarrow \infty$, see e.g. [30, 28] including other stability properties in more general contexts.

According to the previous remark, if $\hat{f} \in \widehat{\mathcal{O}}_s$, the same is true for $\hat{f}(A\mathbf{x})$, for all matrices $A \in \mathbb{C}^{d \times d}$, c.f. [14, Lemma 2.1]. In particular, we highlight the following simple statement we will need later.

Lemma 4.2. *Let $s \geq 0$. Then, $\hat{f} \in \widehat{\mathcal{O}}_s$ if and only if $\hat{f}(A\mathbf{x}) \in \widehat{\mathcal{O}}_s$, for all $A \in GL_n(\mathbb{C})$.*

Consider a germ $P \in \mathcal{O} \setminus \{0\}$ such that $P(\mathbf{0}) = 0$, and $s \geq 0$. There are equivalent definitions for Gevrey series with respect to the germ P [23, Def./Prop. 7.5]. For simplicity, we will use the characterization given in [8, Lemma 4.1].

Definition 4.3. A series $\hat{f} \in \widehat{\mathcal{O}}$ is P - s -Gevrey series if there is a polyradius \mathbf{r} , constants $C, A > 0$ and a sequence $\{f_n\}_{n \in \mathbb{N}} \in \mathcal{O}_b(D_{\mathbf{r}})$ such that

$$(18) \quad \hat{f} = \sum_{n=0}^{\infty} f_n P^n, \quad \text{where } \sup_{\mathbf{x} \in D_{\mathbf{r}}} |f_n(\mathbf{x})| \leq CA^n n!^s.$$

We will use the notation $\widehat{\mathcal{O}}^{P,s}$ for the set of P - s -Gevrey series.

This clearly generalizes the notion of s -Gevrey series in x_j , uniformly in the other variables (x_j - s -Gevrey series in our notation). In fact, setting $j = 1$ to fix ideas, the classical notions required that when we write $\hat{f} = \sum_{n=0}^{\infty} f_n x_1^n$ as a power series in x_1 , there is a polyradius $\mathbf{r}' \in \mathbb{R}^{d-1}$ such that $f_n \in \mathcal{O}_b(D_{\mathbf{r}'})$ and $\sup_{\mathbf{x}' \in D_{\mathbf{r}'}} |f_n(\mathbf{x}')| \leq CA^n n!^s$, for adequate constants C, A .

By using the generalized Weierstrass division we can show the notion of P - s -Gevrey series is well-defined, in the sense that it is independent of the decomposition (18). Note it is enough to check the definition for the decomposition (15) induced by a given injective linear form $\ell : \mathbb{N}^d \rightarrow \mathbb{R}^+$. In fact, if (18) holds, and since all f_n are defined in a common polydisc, we can use decomposition (16) to find $\rho > 0$ and sequences $\{f_{n,j}\}_{n \in \mathbb{N}} \subset \mathcal{O}_b(D_\rho^d)$ with $J(f_{n,j}) \in \Delta_\ell(P)$, such that $f_n = \sum_{j=0}^{\infty} f_{n,j} P^j$, valid for $|\mathbf{x}| < \rho$, where $f_{n,j} = R_{P,\ell} \circ Q_{P,\ell}^j(f_n)$. Therefore, the decomposition (15) of \hat{f} is given by

$$\hat{f} = \sum_{n=0}^{\infty} g_n P^n, \quad g_n = \sum_{j=0}^n f_{j,n-j} \in \mathcal{O}_b(D_\rho^d), \quad J(g_n) \in \Delta_\ell(P),$$

and the sequence $(g_n)_{n \in \mathbb{N}}$ exhibits s -Gevrey bounds since

$$|g_n(\mathbf{x})| \leq \sum_{j=0}^n \|R_{P,\ell}\| \|Q_{P,\ell}\|^{n-j} \sup_{|\mathbf{y}| \leq \rho} |f_j(\mathbf{y})| \leq \sum_{j=0}^n \|R_{P,\ell}\| \|Q_{P,\ell}\|^{n-j} C A^j j!^s, \quad |\mathbf{x}| < \rho,$$

as we wanted to show.

From the previous definition it is easy to deduce many properties on this type of series. We recall the following, valid for $P, Q \in \mathcal{O} \setminus \{0\}$ such that $P(\mathbf{0}) = Q(\mathbf{0}) = 0$, c.f. [8, Coro. 4.2, Lemma 4.3]:

- (a) $\widehat{\mathcal{O}}^{P,s}$ is stable under sums, products, partial derivatives, and it contains \mathcal{O} .
- (b) For any $k \in \mathbb{N}^+$, $\widehat{\mathcal{O}}^{P^k, ks} = \widehat{\mathcal{O}}^{P,s}$.
- (c) If Q divides P , then $\widehat{\mathcal{O}}^{P,s} \subseteq \widehat{\mathcal{O}}^{Q,s}$. In particular, if $Q = U \cdot P$, $U \in \mathcal{O}^*$, then $\widehat{\mathcal{O}}^{P,s} = \widehat{\mathcal{O}}^{Q,s}$.
- (d) Let $\phi : (\mathbb{C}^d, \mathbf{0}) \rightarrow (\mathbb{C}^d, \mathbf{0})$ be analytic, $\phi(\mathbf{0}) = \mathbf{0}$, and assume $P \circ \phi$ is not identically zero. If $\hat{f} \in \widehat{\mathcal{O}}^{P,s}$, then $\hat{f} \circ \phi \in \widehat{\mathcal{O}}^{P \circ \phi, s}$.
- (e) If $P(\mathbf{x}) = \mathbf{x}^\alpha$, $\alpha \in \mathbb{N}^d \setminus \{\mathbf{0}\}$, then $\hat{f} = \sum f_\beta \mathbf{x}^\beta \in \widehat{\mathcal{O}}^{\mathbf{x}^\alpha, s}$ if and only if there are constants $C, A > 0$ satisfying

$$(19) \quad |f_\beta| \leq C A^{|\beta|} \min\{\beta_j!^{s/\alpha_j} : j = 1, \dots, d, \alpha_j \neq 0\}, \quad \beta \in \mathbb{N}^d.$$

It follows from (19) that if $\hat{f} \in \widehat{\mathcal{O}}^{\mathbf{x}^\alpha, s}$, then $\hat{f} \in \widehat{\mathcal{O}}_{s/|\alpha|}$. Indeed, this is a consequence of the inequality $\min\{a_1, \dots, a_d\} \leq a_1^{\tau_1} \cdots a_d^{\tau_d}$, valid for all $a_j > 0$ and $\tau_j \geq 0$ such that $\tau_1 + \dots + \tau_d = 1$, by applying it to $\tau_j = \alpha_j/|\alpha|$. This property can be generalized to an arbitrary germ and we have the following new inclusion of rings of Gevrey series.

Proposition 4.4. *Consider $P \in \mathcal{O}$ with $o(P) = k \geq 1$. Then, a P - s -Gevrey series is a $(s/k, \dots, s/k)$ -Gevrey series. In symbols,*

$$\widehat{\mathcal{O}}^{P,s} \subseteq \widehat{\mathcal{O}}_{s/k}.$$

Proof. Write $P = \sum_{j=k}^{\infty} P_j$ as sum of homogeneous polynomials. Since $P_k \neq 0$, we can find $\mathbf{a} \neq \mathbf{0}$ such that $P_k(\mathbf{a}) \neq 0$. Choose $A \in \text{GL}_n(\mathbb{C})$ having \mathbf{a} as first column. If we set $Q(\mathbf{x}) = P(A\mathbf{x})$ and we write it as sum of its homogeneous components $Q = \sum Q_j$, then $Q_j(\mathbf{x}) = P_j(A\mathbf{x})$, and $Q_k(\mathbf{x}) = P_k(\mathbf{a})x_1^k + \dots$, i.e., $o(Q) = k$ and $Q_k(1, 0, \dots, 0) \neq 0$.

Consider a P - s -Gevrey series \hat{f} . Then $\hat{f}_0(\mathbf{x}) = \hat{f}(A\mathbf{x}) = \sum a_\beta \mathbf{x}^\beta$ is a Q - s -Gevrey series. We now consider the change of variables

$$(20) \quad x_1 = z_1, \quad x_2 = z_1 z_2, \quad \dots, \quad x_d = z_1 z_d,$$

that geometrically corresponds to a local expression for the blow-up of the origin in \mathbb{C}^d [22]. If $R(\mathbf{z}) = Q(\mathbf{x})$ and $\hat{f}_1(\mathbf{z}) = \hat{f}_0(\mathbf{x})$, we see \hat{f}_1 is a R - s -Gevrey series.

On the one hand,

$$R(\mathbf{z}) = Q(z_1, z_1 z_2, \dots, z_1 z_d) = \sum_{j=k}^{\infty} z_1^j Q_j(1, z_2, \dots, z_d) = z_1^k U(\mathbf{z}),$$

where U is a unit, since $U(\mathbf{0}) = Q_k(1, 0, \dots, 0) \neq 0$. Thus, we conclude \hat{f}_1 is z_1^k - s -Gevrey, or equivalently, a z_1 - s/k -Gevrey series. On the other hand, since \hat{f}_1 is written as

$$\hat{f}_1(\mathbf{z}) = \hat{f}_0(z_1, z_1 z_2, \dots, z_1 z_d) = \sum_{\boldsymbol{\beta} \in \mathbb{N}^d} a_{\boldsymbol{\beta}} z_1^{|\boldsymbol{\beta}|} z_2^{\beta_2} \dots z_d^{\beta_d} = \sum_{\substack{(n, \boldsymbol{\gamma}) \in \mathbb{N} \times \mathbb{N}^{d-1} \\ n \geq |\boldsymbol{\gamma}|}} a_{n-|\boldsymbol{\gamma}|, \boldsymbol{\gamma}} z_1^n \mathbf{z}'^{\boldsymbol{\gamma}},$$

we can find constants $C, A > 0$ such that $|a_{n-|\boldsymbol{\gamma}|, \boldsymbol{\gamma}}| \leq CA^{n+|\boldsymbol{\gamma}|} n!^{s/k}$. Therefore, in the index $\boldsymbol{\beta} = (n, \boldsymbol{\gamma})$, we find the bound

$$|a_{\boldsymbol{\beta}}| \leq CA^{\beta_1 + 2\beta_2 + \dots + 2\beta_d} |\boldsymbol{\beta}|!^{s/k}, \quad \text{for all } \boldsymbol{\beta} \in \mathbb{N}^d.$$

This means \hat{f}_0 and \hat{f} are $(s/k, \dots, s/k)$ -Gevrey series, thanks to Lemma 4.2. \square

Remark 4.5. Proposition 4.4 and Lemma 4.2 shows that if $\hat{f} \in \widehat{\mathcal{O}}^{\boldsymbol{\alpha}, s}$, then $\hat{f}(A\mathbf{x}) \in \widehat{\mathcal{O}}_{s/|\boldsymbol{\alpha}|}$, for all $A \in \mathbb{C}^{d \times d}$. However, being $\boldsymbol{\alpha}$ - s -Gevrey is not stable under linear changes of variable. We illustrate this by a simple example: the series $\hat{f}(x_1, x_2) = \sum_{n=0}^{\infty} n! (x_1 x_2)^n$ is $x_1 x_2$ -1-Gevrey, but

$$\hat{f}_0(\xi_1, \xi_2) = \hat{f}(\xi_1 + \xi_2, \xi_1 - \xi_2) = \sum_{j, k \geq 0} \binom{j+k}{j} (j+k)! (-1)^j \xi_1^{2j} \xi_2^{2k},$$

is $(1/2, 1/2)$ -Gevrey in ξ_1, ξ_2 , but not $\xi_1 \xi_2$ -1-Gevrey.

5. PROOF OF THE MAIN RESULTS

Proof of Theorem 1.1. The idea of the proof is to write

$$(21) \quad \widehat{\mathbf{y}} = \sum_{n=0}^{\infty} y_n P^n,$$

as a power series in the germ P according to decomposition (15) for a given injective linear form $\ell : \mathbb{N}^d \rightarrow \mathbb{R}_+$, and then to find recursively the coefficients y_n , $J(y_n) \in \Delta_{\nu_\ell(P)}^N$. Then, with the aid of Nagumo norms and the majorant series technique we will establish the required Gevrey property.

Before we start we note that if $F(\mathbf{x}, \mathbf{0})$ is identically zero, the unique formal power series solution is the zero series. Thus, we can assume $c(\mathbf{x}) := F(\mathbf{x}, \mathbf{0}) \neq 0$.

We divide the proof in several steps.

Step 0 (Preliminaries) Let us fix an injective linear form $\ell : \mathbb{N}^d \rightarrow \mathbb{R}_+$ and $\rho' > 0$ sufficiently small such that $Q_{P, \ell}, R_{P, \ell} : \mathcal{O}_b(D_{\rho'(\ell)}) \rightarrow \mathcal{O}_b(D_{\rho'(\ell)})$ from Proposition 3.1 are defined. We consider the linear operators $Q, R : \mathcal{O}_b(D_{\rho'(\ell)})^N \rightarrow \mathcal{O}_b(D_{\rho'(\ell)})^N$ given by $Q(f_1, \dots, f_N) = (Q_{P, \ell}(f_1), \dots, Q_{P, \ell}(f_N))$, and $R(f_1, \dots, f_N) = (R_{P, \ell}(f_1), \dots, R_{P, \ell}(f_N))$. Then, by using the norms (11) we see that

$$(22) \quad \|Q(\mathbf{f})\|_m \leq \|Q\| \cdot \|\mathbf{f}\|_m, \quad \|R(\mathbf{f})\|_m \leq \|R\| \cdot \|\mathbf{f}\|_m,$$

for all $m \in \mathbb{N}$ and $\mathbf{f} \in \mathcal{O}_b(D_{\rho'(\ell)})^N$, where to simplify notation we write $\|Q\| = 2 \cdot 4^{|\nu_\ell(P)|} / \rho'^{\ell(\nu_\ell(P))}$ and $\|R\| = 2(1 + 4^{|\nu_\ell(P)|})$. The same considerations and inequalities are valid for matrix-valued maps. It will be important for later to note that $\|R\|$ is independent of the radius, since we will shrink ρ' during this proof.

Since F is analytic we can write it as a convergent power series in \mathbf{y} , say

$$F(\mathbf{x}, \mathbf{y}) = c(\mathbf{x}) + (\mu + A(\mathbf{x}))\mathbf{y} + \sum_{|I| \geq 2} A_I(\mathbf{x})\mathbf{y}^I,$$

where $c, A_I \in \mathcal{O}_b(D_{r'}^d)^N$, $A \in \mathcal{O}_b(D_{r'}^d)^{N \times N}$, $A(\mathbf{0}) = \mathbf{0}$, and the summation is taken over all $I = (i_1, \dots, i_N) \in \mathbb{N}^N$ such that $|I| = i_1 + \dots + i_N \geq 2$. Furthermore, we can find $K, \delta > 0$ such that

$$(23) \quad \|A_I\|_0 \leq K\delta^{|I|}, \quad \text{for all } I \in \mathbb{N}^N.$$

Note that all the previous coefficients are defined in the common polydisc of polyradius (r', \dots, r') . By reducing r' if necessary, we can assume $a_1, \dots, a_d \in \mathcal{O}_b(D_{r'}^d)$ which are not all identically zero.

Now, choose $0 < s < \rho'$ such that $s^{l_j} \leq r'$, for all j , in order to be able to apply Corollary 3.2 to the previous functions. Then, we conclude there is $r > 0$ such that we can write

$$a_j = \sum_{n=0}^{\infty} a_{j,n} P^n, \quad c = \sum_{n=0}^{\infty} c_n P^n, \quad A = \sum_{n=0}^{\infty} A_n P^n, \quad A_I = \sum_{n=0}^{\infty} A_{I,n} P^n,$$

where $A_n \in (\mathcal{O}_b(D_r^d) \cap \Delta_{\nu_\ell(P)})^{N \times N}$, $c_n, A_{I,n} \in (\mathcal{O}_b(D_r^d) \cap \Delta_{\nu_\ell(P)})^N$, $a_{j,n} \in \mathcal{O}_b(D_r^d) \cap \Delta_{\nu_\ell(P)}$, for all $j = 1, \dots, d$, $I \in \mathbb{N}^N$ and $n \in \mathbb{N}$.

Step 1 (The coefficient y_0) Our first step is to determine the first term in (21), namely, y_0 . Note that $\hat{\mathbf{y}}(\mathbf{0}) = y_0(\mathbf{0}) = \mathbf{0}$ since $F(\mathbf{0}, \mathbf{0}) = \mathbf{0}$ and $P(\mathbf{0}) = 0$.

When we plug $\hat{\mathbf{y}}$ into equation (1) and equate in common powers of P , we find y_0 must be an analytic solution of

$$(24) \quad 0 = R(c + (\mu + A)y + \sum_{|I| \geq 2} A_I y^I) = c_0 + \mu y_0 + R(A_0 y) + \sum_{|I| \geq 2} R(A_{I,0} y^I),$$

such that $J(y_0) \in \Delta_{\nu_\ell(P)}^N$. We will prove this problem has a unique solution in $\mathcal{O}_b(D_r^d)$ if $r > 0$ is taken small enough. In order to proceed, we write (24) as the fixed point equation

$$y = G(y), \quad G(y) := -\mu^{-1} R \left(c_0 + A_0 y + \sum_{|I| \geq 2} A_{I,0} y^I \right).$$

We will show we can reduce $r > 0$ and find $\epsilon > 0$ sufficiently small such that $G : \overline{B}_\epsilon \rightarrow \overline{B}_\epsilon$ is well-defined and a contraction, where $\overline{B}_\epsilon := \{y \in \mathcal{O}_b(D_r^d)^N : \|y\|_0 \leq \epsilon, y(\mathbf{0}) = \mathbf{0}\}$ which is closed. Then, by Banach fixed point theorem G has a unique fixed point.

Let us check first that G maps $\overline{B}_{1/2\delta}$ to $\mathcal{O}_b(D_r^d)^N$, where δ is as in (23): by using (22) for R and (23) we see that

$$\|G(y)\|_0 \leq \|\mu^{-1}\|_0 \|R\|^2 \left(\|c\|_0 + \|A\|_0 + \sum_{|I| \geq 2} K\delta^{|I|} \|y\|_0^{|I|-1} \right) \|y\|_0.$$

But the identity $\sum_{|I| \geq 1} \alpha^{|I|} = (1 - \alpha)^{-N} - 1$, $|\alpha| < 1$, shows that

$$\|G(y)\|_0 \leq \|\mu^{-1}\|_0 \|R\|^2 \left(\|c\|_0 + \|A\|_0 + K\delta (2^N - 1) \right) \|y\|_0,$$

thus, $\|G(y)\|_0$ is finite as we wanted to show.

On the other hand, if $y + h, y \in \overline{B}_{1/2\delta}$ we also have

$$\|G(y + h) - G(y)\|_0 \leq \|\mu^{-1}\|_0 \|R\|^2 \left(\|A\|_0 \|h\|_0 + \sum_{|I| \geq 2} K\delta^{|I|} \|(y + h)^I - y^I\|_0 \right).$$

Taking into account the inequality $\|(y+h)^I - y^I\|_0 \leq |I|(\|y\|_0 + \|h\|_0)^{|I|-1}\|h\|_0$ that follows readily by induction on $|I|$, we obtain

$$\begin{aligned} \|G(y+h) - G(y)\|_0 &\leq \|\mu^{-1}\|_0 \|R\|^2 \left(\|A\|_0 + K \sum_{|I| \geq 2} |I| \delta^{|I|} (\|y\|_0 + \|h\|_0)^{|I|-1} \right) \|h\|_0 \\ &= \|\mu^{-1}\|_0 \|R\|^2 \left(\|A\|_0 + K \delta g(\delta(\|y\|_0 + \|h\|_0)) \right) \|h\|_0, \end{aligned}$$

where $g(\alpha) = \sum_{|I| \geq 2} |I| \alpha^{|I|-1} = \frac{d}{d\alpha} \left(\sum_{|I| \geq 2} \alpha^{|I|} \right) = N((1-\alpha)^{-N-1} - 1)$, for $|\alpha| < 1$. Since $g(0) = 0$, by its continuity, we can choose $0 < \epsilon < \min\{1, 1/2\delta\}$ such that $\|\mu^{-1}\|_0 \|R\|^2 K \delta \cdot g(\epsilon) < 1/4$. Also, since $A(\mathbf{0}) = \mathbf{0}$, we can reduce r to have $\|\mu^{-1}\|_0 \|R\|^2 \|A\|_0 < 1/4$. Therefore, if $\|y\|_0 + \|h\|_0 \leq \epsilon$, we conclude

$$\|G(y+h) - G(y)\|_0 \leq \frac{1}{2} \|h\|_0.$$

But $c(\mathbf{0}) = F(\mathbf{0}, \mathbf{0}) = \mathbf{0}$, and since ϵ has been fixed, we can reduce r again to have $\|\mu^{-1}\|_0 \|R\|^2 \|c\|_0 < \epsilon/2$. By applying the previous inequality to $y = 0$ we find $\|G(h)\|_0 \leq \|G(h) - G(0)\|_0 + \|G(0)\|_0 \leq \frac{1}{2} \|h\|_0 + \epsilon/2$. Thus, $G : \overline{B}_\epsilon \rightarrow \overline{B}_\epsilon$ has the desired properties.

Several remarks are at hand. First, if y_0 is the solution of equation (24), $J(y_0)$ will be the unique formal solution of (24) and it is convergent. But $R(y_0)$ is another analytic solution of (24), thus $J(y_0) = J(R(y_0)) \in \Delta_{\nu_\ell(P)}^N$. Second, there is a direct way to find a solution of (24) as follows: we find first a solution $Y_0(\mathbf{x})$ of $F(\mathbf{x}, \mathbf{y}(\mathbf{x})) = \mathbf{0}$, with the aid of the holomorphic implicit function theorem - it can be applied since $F(\mathbf{0}, \mathbf{0}) = \mathbf{0}$ and $\frac{\partial F}{\partial \mathbf{y}}(\mathbf{0}, \mathbf{0}) = \mu$ is invertible-. Then, it follows by applying R to the previous equation that $y_0 = R(Y_0)$ is the solution of (24), since we already know it is unique.

Step 2 (Recurrence equations for y_n) We can now assume $y_0 = 0$ by making the change of variables $y \mapsto y - y_0$ in the initial equation (1). In fact, after doing so, we obtain a similar PDE such that P divides c and we search for a formal solution $\hat{\mathbf{y}} = \sum_{n=1}^{\infty} y_n P^n$ which is divisible by P .

To find the recurrence equations satisfied by the y_n we start with the right side of (1). By using the identity (17) for the product of series we find

$$A \cdot \hat{\mathbf{y}} = \sum_{n=1}^{\infty} \left(\sum_{k=1}^n \sum_{j=1}^k RQ^{n-k} (A_{k-j} y_j) \right) P^n.$$

For the non-linear term we have the decomposition

$$\begin{aligned} \sum_{|I| \geq 2} A_I \hat{\mathbf{y}}^I &= \sum_{k=2}^{\infty} \sum_{*k} A_{I,m} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} y_{l,n_{l,j}} P^k = \sum_{k=2}^{\infty} \sum_{p=0}^{\infty} \sum_{*k} RQ^p \left(A_{I,m} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} y_{l,n_{l,j}} \right) P^{k+p} \\ &= \sum_{n=2}^{\infty} \left(\sum_{k=2}^n \sum_{*k} RQ^{n-k} \left(A_{I,m} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} y_{l,n_{l,j}} \right) \right) P^n. \end{aligned}$$

where the sum \sum_{*k} is taken over all $I \in N^N$ such that $2 \leq |I| \leq k$, m satisfying $0 \leq m \leq k - |I|$, and $n_{l,j} \geq 1$ such that $k = m + n_{1,1} + \dots + n_{1,i_1} + \dots + n_{N,1} + \dots + n_{N,i_N}$. Note in particular that $n_{l,j} < k \leq n$ and thus no component of $y_n = (y_{n,1}, \dots, y_{n,d})$ appears in the coefficient corresponding to P^n .

For the left-side of (1) we can write

$$(25) \quad P \cdot L(\hat{\mathbf{y}}) = \sum_{n=1}^{\infty} (L(y_{n-1}) + n y_n L(P)) P^n.$$

At this point we use the hypothesis $L(P) = P \cdot h$, for some $h \in \mathcal{O}_b(D_r^d)$, so the previous equations becomes

$$P \cdot L(\hat{\mathbf{y}}) = \sum_{n=2}^{\infty} (L(y_{n-1}) + (n-1)hy_{n-1}) P^n.$$

Now, we can equate both sides of equation (1) in common power series of P , to obtain the recurrence

$$\begin{aligned} \sum_{k=2}^n RQ^{n-k}(L(y_{k-1})) + \sum_{k=2}^n (k-1)RQ^{n-k}(hy_{k-1}) = c_n + \mu y_n + \sum_{k=1}^n \sum_{j=1}^k RQ^{n-k}(A_{k-j}y_j) \\ + \sum_{k=2}^n \sum_{*k} RQ^{n-k} \left(A_{I,m} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} y_{l,n_{l,j}} \right), \end{aligned}$$

or equivalently

$$(26) \quad \mu y_n + R(A_0 y_n) = b_n := e_n + \sum_{k=2}^n (k-1)RQ^{n-k}(hy_{k-1}), \quad \text{for all } n \geq 1,$$

where

$$\begin{aligned} e_n = -c_n + \sum_{k=2}^n RQ^{n-k}(L(y_{k-1})) - \sum_{k=1}^{n-1} \sum_{j=1}^k RQ^{n-k}(A_{k-j}y_j) - \sum_{j=1}^{n-1} R(A_{n-j}y_j) \\ - \sum_{k=2}^n \sum_{*k} RQ^{n-k} \left(A_{I,m} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} y_{l,n_{l,j}} \right) \end{aligned}$$

Note in particular that $b_1 = -c_1$.

Equation (26) can be solved as follows: consider $Y_n = (\mu + A_0)^{-1}(b_n)$, where we have if necessary, reduced r to ensure $\mu + A_0(\mathbf{x})$ is invertible for all $|\mathbf{x}| \leq r$. Then $R(Y_n)$ solves (26), as we see by applying R to $(\mu + A_0)Y_n = b_n$ and recalling that $R(b_n) = b_n$. To check uniqueness, note that if y_n and w_n are solutions, then $R((\mu + A_0)(y_n - w_n)) = 0$, so $(\mu + A_0)(y_n - w_n) = h_1 P$, for some $h_1 \in \mathcal{O}$. Thus, $y_n - w_n = R(y_n - w_n) = R((\mu + A_0)^{-1}h_1 P) = 0$.

In conclusion, we can find recursively the coefficients y_n by means of the formulas

$$(27) \quad y_n = R((\mu + A_0)^{-1}(b_n)),$$

and equation (1) has a unique formal power series solution.

Step 3 (Majorant series) We use the majorant series technique to show that $\hat{\mathbf{y}}$ is P -1-Gevrey by proving that $\sum_{n=1}^{\infty} \|y_n\|_n \tau^n$ is 1-Gevrey in τ .

We have chosen $r > 0$ satisfying all previous requirements in order to find y_n recursively. Now, we take $0 < \rho < \min\{r, 1\}$ satisfying $\rho^{l_j} < r$, $j = 1, \dots, d$, in order to apply the bounds (22) for functions in $\mathcal{O}_b(D_{\rho(\ell)})^N$.

Let $M = \|(\mu + A_0)^{-1}\|_0 > 0$. By applying the Nagumo norm $\|\cdot\|_n$ to equation (27) and taking into account the properties developed in Propositions 2.1 and 3.1 we find that

$$\begin{aligned} \frac{\|y_n\|_n}{M\|R\|} \leq \|c_n\|_n + \sum_{k=2}^n \|R\| \|Q\|^{n-k} (\|L(y_{k-1})\|_n + (k-1)\|hy_{k-1}\|_n) \\ + \sum_{k=1}^{n-1} \|R\| \|Q\|^{n-k} \sum_{j=1}^k \|A_{k-j}\|_{k-j} \|y_j\|_j + \sum_{j=1}^{n-1} \|R\| \|A_{n-j}\|_{n-j} \|y_j\|_j \\ + \sum_{k=2}^n \|R\| \|Q\|^{n-k} \sum_{*k} \|A_{I,m}\|_m \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} \|y_{l,n_{l,j}}\|_{n_{l,j}}. \end{aligned}$$

To bound effectively $\|L(y_{k-1})\|_n$, note that since $0 < \rho < 1$, by Proposition 2.1 (2) we have

$$\left\| a_j \frac{\partial y_{k-1}}{\partial x_j} \right\|_n \leq \|a_j\|_{n-k} \left\| \frac{\partial y_{k-1}}{\partial x_j} \right\|_k \leq ek \prod_{i \neq j} (\rho^{\ell_i} / 2) \|a_j\|_{n-k} \|y_{k-1}\|_{k-1} \leq ek \|a_j\|_{n-k} \|y_{k-1}\|_{k-1}.$$

But inequality (8) implies that $\|a_j\|_{n-k} \leq \|a_j\|_0$. Therefore, if $a = \|a_1\|_0 + \dots + \|a_d\|_0$, by hypothesis $a > 0$, and

$$\|L(y_{k-1})\|_n \leq eak \|y_{k-1}\|_{k-1}.$$

On the other hand,

$$\|hy_{k-1}\|_n \leq \|h\|_{n-k+1} \|y_{k-1}\|_{k-1} \leq \|h\|_0 \|y_{k-1}\|_{k-1}.$$

Thus, we find that

$$(28) \quad \begin{aligned} \frac{\|y_n\|_n}{M\|R\|} &\leq \|c_n\|_n + \|R\|(ea + \|h\|_0) \sum_{k=2}^n k \|Q\|^{n-k} \|y_{k-1}\|_{k-1} + \|R\| \sum_{k=1}^{n-1} \|Q\|^{n-k} \sum_{j=1}^k \|A_{k-j}\|_{k-j} \|y_j\|_j \\ &+ \|R\| \sum_{j=1}^{n-1} \|A_{n-j}\|_{n-j} \|y_j\|_j + \|R\| \sum_{k=2}^n \|Q\|^{n-k} \sum_{*_k} \|A_{I,m}\|_m \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} \|y_{n_l,j}\|_{n_l,j}. \end{aligned}$$

If we divide by $n!$ and using that $m!k! \leq (m+k)!$ we conclude that

$$\begin{aligned} \frac{\|y_n\|_n}{M\|R\|n!} &\leq \frac{\|c_n\|_n}{n!} + \|R\|(ea + \|h\|_0) \sum_{k=2}^n \frac{\|Q\|^{n-k}}{(n-k)!} \frac{\|y_{k-1}\|_{k-1}}{(k-1)!} + \|R\| \sum_{k=1}^{n-1} \frac{\|Q\|^{n-k}}{(n-k)!} \sum_{j=1}^k \frac{\|A_{k-j}\|_{k-j}}{(k-j)!} \frac{\|y_j\|_j}{j!} \\ &+ \|R\| \sum_{j=1}^{n-1} \frac{\|A_{n-j}\|_{n-j}}{(n-j)!} \frac{\|y_j\|_j}{j!} + \|R\| \sum_{k=2}^n \frac{\|Q\|^{n-k}}{(n-k)!} \sum_{*_k} \frac{\|A_{I,m}\|_m}{m!} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} \frac{\|y_{n_l,j}\|_{n_l,j}}{n_l,j!}. \end{aligned}$$

Let us define the sequence z_n recursively by

$$(29) \quad \begin{aligned} \frac{z_n}{M\|R\|} &= \frac{\|c_n\|_n}{n!} + \|R\|(ea + \|h\|_0) \sum_{k=2}^n \frac{\|Q\|^{n-k}}{(n-k)!} z_{k-1} + \|R\| \sum_{k=1}^{n-1} \frac{\|Q\|^{n-k}}{(n-k)!} \sum_{j=1}^k \frac{\|A_{k-j}\|_{k-j}}{(k-j)!} z_j \\ &+ \|R\| \sum_{j=1}^{n-1} \frac{\|A_{n-j}\|_{n-j}}{(n-j)!} z_j + \|R\| \sum_{k=2}^n \frac{\|Q\|^{n-k}}{(n-k)!} \sum_{*_k} \frac{\|A_{I,m}\|_m}{m!} \prod_{\substack{1 \leq l \leq N \\ 1 \leq j \leq i_l}} z_{n_l,j}, \end{aligned}$$

where $z_1 = M\|R\|\|c_1\|_1$. Since the terms of the previous equation are all non-negative real numbers, we find inductively that

$$(30) \quad \frac{\|y_n\|_n}{n!} \leq z_n.$$

On the other hand, we consider the generating power series

$$\bar{c}(\tau) = \sum_{n=1}^{\infty} \frac{\|c_n\|_n}{n!} \tau^n, \quad \bar{A}(\tau) = \sum_{n=0}^{\infty} \frac{\|A_n\|_n}{n!} \tau^n, \quad \bar{F}(\tau, Y) = \sum_{m \geq 0, |I| \geq 2} \frac{\|A_{I,m}\|_m}{m!} \tau^m Y^{|I|}.$$

which are in fact convergent. For instance, the coefficient of $\tau^m Y^j$ is given and bounded by $\sum_{|I|=j} \frac{\|A_{I,m}\|_m}{m!} \leq \sum_{|I|=j} K \|R\| \frac{\|Q\|_m^m}{m!} \delta^{|I|} = K \|R\| \frac{\|Q\|_m^m}{m!} \delta^j \binom{j+N-1}{N-1} \leq K \|R\| \frac{\|Q\|_m^m}{m!} \delta^j 2^{j+N-1}$.

By using these series and equation (29), we find $Z(\tau) = \sum_{n=1}^{\infty} z_n \tau^n$ is a formal solution of the analytic equation

$$\frac{Z(\tau)}{M\|R\|} = \bar{c}(\tau) + \|R\|(ea + \|h\|_0) e^{\|Q\|\tau} \tau Z(\tau) + \|R\|(e^{\|Q\|\tau} \bar{A}(\tau) - \|A_0\|_0) Z(\tau) + \|R\| e^{\|Q\|\tau} \bar{F}(\tau, Z(\tau)),$$

but the holomorphic implicit function theorem implies this equation has a unique convergent power series solution at the origin, thus it must be $Z(\tau)$, so it is convergent. By (30) the series $\sum_{n=1}^{\infty} \|y_n\|_n \tau^n$ is 1-Gevrey in τ as we wanted to show. \square

Proof of Theorem 1.2. Regarding the previous proof, only some minor changes are required so establish the result. While Step 0 and Step 1 remain the same, in Step 2 the recurrence for y_n takes the form

$$(31) \quad \mu y_n - nR(L(P)y_n) + R(A_0 y_n) = d_n := e_n + \sum_{k=1}^{n-1} kRQ^{n-k}(L(P)y_k), \quad \text{for all } n \geq 1,$$

and e_n as before. Then d_n and b_n differ only in the previous sum, that we will bound by shifting one index, as follows

$$(32) \quad \left\| \sum_{k=2}^n (k-1)RQ^{n-k+1}(L(P)y_{k-1}) \right\|_n \leq \|R\| \|Q\| \|L(P)\|_0 \sum_{k=2}^n k \|Q\|^{n-k} \|y_{k-1}\|_{k-1}.$$

In the current case, the solution of (31) is given by

$$y_n = R((\mu - nL(P)I_N + A_0)^{-1}(d_n)),$$

where I_N is the identity matrix of size N . To make this formula meaningful, it is enough to prove $\mu - nL(P)(\mathbf{x})I_N + A_0(\mathbf{x})$ is invertible for all $n \geq 1$ and $|\mathbf{x}| \leq r$, if r is sufficiently small. To proceed let us recall that if $B \in \mathbb{C}^{N \times N}$ is such that $|B| < 1$ for a matrix norm $|\cdot|$, then $I_N - B$ is invertible, $(I_N - B)^{-1} = \sum_{n=0}^{\infty} B^n$, and $|(I_N - B)^{-1}| \leq (1 - |B|)^{-1}$. Here as before we use $|B| = \max_{1 \leq i \leq N} \sum_{j=1}^N |B_{i,j}|$. Now, since $L(P)(\mathbf{0}) \neq 0$, we can choose a small $r > 0$ such that $\alpha = \inf_{|\mathbf{x}| \leq r} |L(P)(\mathbf{x})| > 0$. Thus, if $n > \|\mu + A_0\|_0 / \alpha$, we see that $|\mu + A_0(\mathbf{x})| / |nL(P)(\mathbf{x})| \leq \|\mu + A_0\|_0 / n\alpha < 1$, for all $|\mathbf{x}| \leq r$, so $\mu - nL(P) + A_0$ is invertible with inverse bounded by

$$\begin{aligned} |(\mu - nL(P)(\mathbf{x}) + A_0(\mathbf{x}))^{-1}| &= \frac{1}{n|L(P)(\mathbf{x})|} \left| \left(I_N - \frac{1}{nL(P)(\mathbf{x})} (\mu + A_0(\mathbf{x})) \right)^{-1} \right| \\ &\leq \frac{1}{\alpha n} \frac{1}{1 - \frac{\|\mu + A_0\|_0}{\alpha n}} = \frac{1}{\alpha n - \|\mu + A_0\|_0}. \end{aligned}$$

For $n \leq \|\mu + A_0\|_0 / \alpha$, by hypothesis the remaining finite number of matrices $\mu - nL(P)(\mathbf{x}) + A_0(\mathbf{x})$ are invertible at the origin. Thus, we can shrink r and assume they are invertible for all $|\mathbf{x}| \leq r$. In conclusion, all these matrices are invertible, and we can find $M > 0$ such that

$$(33) \quad \|(\mu - nL(P) + A_0)^{-1}\|_0 \leq M/n, \quad \text{for all } n \geq 1.$$

At this stage we can proceed with Step 3 by using the Nagumo norms and taking into account (32). However, the factor M/n in (33) improves our bounds and shows that

$$\frac{\|y_n\|_n}{M\|R\|} \leq \|c_n\|_n + \|R\|(ea + \|Q\|\|L(P)\|_0) \sum_{k=2}^n \|Q\|^{n-k} \|y_{k-1}\|_{k-1} + \dots,$$

where the dots indicate the remaining terms are the same as in (28). In this case it is not necessary to divide by $n!$, just by defining z_n accordingly we find $\|y_n\|_n \leq z_n$, for all $n \geq 1$, and $Z(\tau)$ satisfies the analytic equation

$$\frac{Z(\tau)}{M\|R\|} = \bar{c}(\tau) + \|R\|(ea + \|Q\|\|L(P)\|_0) \frac{\tau Z(\tau)}{1 - \|Q\|\tau} + \|R\| \left(\frac{\bar{A}(\tau)}{1 - \|Q\|\tau} - \|A_0\|_0 \right) Z(\tau) + \|R\| \frac{\bar{F}(\tau, Z(\tau))}{1 - \|Q\|\tau},$$

with analytic coefficients $\bar{c}(\tau) = \sum_{n=1}^{\infty} \|c_n\|_n \tau^n$, $\bar{A}(\tau) = \sum_{n=0}^{\infty} \|A_n\|_n \tau^n$, and $\bar{F}(\tau, Y) = \sum_{m \geq 0, |I| \geq 2} \|A_{I,m}\|_m \tau^m Y^{|I|}$. Therefore, by the holomorphic implicit function theorem $Z(\tau)$ and also $\hat{\mathbf{y}}$ are convergent as required. \square

There is a straightforward way to extend the theorems for some systems of PDEs of higher order by augmenting the size of the given equation, as we explain in the following result.

Corollary 5.1. *Let P , L and F be as in Theorem 1.1, fix $u_1, \dots, u_{k-1} \in \mathcal{O}$, and consider the system of PDEs*

$$(34) \quad (P \cdot L)^k(\mathbf{y})(\mathbf{x}) + u_{k-1}(\mathbf{x})(P \cdot L)^{k-1}(\mathbf{y})(\mathbf{x}) + \dots + u_1(\mathbf{x})(P \cdot L)(\mathbf{y})(\mathbf{x}) = F(\mathbf{x}, \mathbf{y}).$$

Then, the following statements hold:

- (1) If P divides $L(P)$, (34) has a unique formal power series solution which is P -1-Gevrey.
- (2) If $L(P)(\mathbf{0}) \neq 0$, and $\sigma - nL(P)(\mathbf{0}) \neq 0$, for all $n \in \mathbb{N}$ and all solutions σ of the polynomial equation

$$(35) \quad p_\mu(\sigma^k + u_{k-1}(\mathbf{0})\sigma^{k-1} + \dots + u_2(\mathbf{0})\sigma^2 + u_1(\mathbf{0})\sigma) = 0,$$

where p_μ is the characteristic polynomial of μ , then (34) has a unique convergent power series solution.

Proof. In the variable $\mathbf{w} = (\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_{k-1}) \in \mathbb{C}^{Nk}$, where $\mathbf{w}_0 = \mathbf{y}$, $\mathbf{w}_1 = (P \cdot L)(\mathbf{w}_0)$, $\mathbf{w}_2 = (P \cdot L)(\mathbf{w}_1)$, \dots , $\mathbf{w}_{k-1} = (P \cdot L)(\mathbf{w}_{k-2})$, (34) can be written as

$$P \cdot L(\mathbf{w}) = G(\mathbf{x}, \mathbf{w}) := (\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_{k-1}, F(\mathbf{x}, \mathbf{w}_0) - u_1(\mathbf{x})\mathbf{w}_1 - \dots - u_{k-1}(\mathbf{x})\mathbf{w}_{k-1}),$$

which has the form of equation (1). Then, the results follow from Theorem 1.1 and 1.2 by noticing that

$$\frac{\partial G}{\partial \mathbf{w}}(\mathbf{0}, \mathbf{0}) = \begin{pmatrix} 0 & I_N & 0 & \dots & 0 \\ 0 & 0 & I_N & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & I_N \\ \mu & -u_1(\mathbf{0})I_N & -u_2(\mathbf{0})I_N & \dots & -u_{k-1}(\mathbf{0})I_N \end{pmatrix} \in \mathbb{C}^{Nk} \times \mathbb{C}^{Nk}$$

is an invertible matrix with eigenvalues given by the solutions of (35). \square

6. EXAMPLES

Example 6.1. The solution of (1) is generically divergent, but there are cases where it can be convergent. This is evidenced already in the case of one variable: while Euler's equation $x^2 y' + y = x$ has the x -1-Gevrey solution $\hat{y}(x) = \sum_{n=0}^{\infty} (-1)^n n! x^{n+1}$, the equation $x^2 y' + y = x + x^2$ has $\hat{y}(x) = x$ as analytic solution. More examples can be obtained by taking $f \in \mathbb{C}\{z\}$ and $P \in \mathcal{O}$, $P(\mathbf{0}) = 0$. If $L(P) = 0$, then the solution of

$$P(\mathbf{x})L(y) = y - f(P(\mathbf{x})), \quad \text{is} \quad \hat{y}(\mathbf{x}) = f(P(\mathbf{x})),$$

which is convergent.

Example 6.2. We consider the equation $x_1 x_2 \frac{\partial y}{\partial x_1} = \mu y - \frac{x_1}{1-x_1}$, where $\mu \neq 0$ is constant. A way to find its unique formal power series solution is to plug $\hat{y} = \sum_{n=0}^{\infty} y_n(x_2) x_1^n$ into the equation and then equate common powers of x_1 . Thus, we find $y_0(x_2) = 0$, $y_n(x_2) = (\mu - n x_2)^{-1}$, $n \geq 1$, and the formal solution is equal to

$$\hat{y}(x_1, x_2) = \sum_{n \geq 1, m \geq 0} \frac{n^m}{\mu^{m+1}} x_1^n x_2^m.$$

We see \hat{y} is x_2 -1-Gevrey by direct inspection or by applying Theorem 1.1 to $P = x_2$ and $L = x_1 \frac{\partial}{\partial x_1}$ since $L(P) = 0$. However, \hat{y} is not x_1 -1-Gevrey, i.e., $P = x_1$, $L = x_2 \frac{\partial}{\partial x_1}$ is not a valid choice: as a power series in x_1 , $y_n(x_2)$ is analytic on the disc $\{x_2 \in \mathbb{C} : |x_2| < |\mu|/n\}$, so there is no common neighborhood of the origin where all $y_n(x_2)$ are defined.

Example 6.3. Consider $\boldsymbol{\alpha} \in \mathbb{N}^d \setminus \{\mathbf{0}\}$, $\mu \in \mathbb{C}^*$, $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_d) \in (\mathbb{C} \setminus \{0\})^d$, and $c(\mathbf{x}) = \sum_{\boldsymbol{\beta} \in \mathbb{N}^d} a_{\boldsymbol{\beta}} \mathbf{x}^{\boldsymbol{\beta}} \in \mathbb{C}\{\mathbf{x}\}$. Then the PDE

$$\mathbf{x}^{\boldsymbol{\alpha}} L_{\boldsymbol{\lambda}}(\mathbf{y}) = \mathbf{x}^{\boldsymbol{\alpha}} (\lambda_1 x_1 \partial_{x_1} \mathbf{y} + \dots + \lambda_d x_d \partial_{x_d} \mathbf{y}) = \mu \mathbf{y} - c(\mathbf{x}),$$

has generically a $\mathbf{x}^{\boldsymbol{\alpha}}$ -1-Gevrey formal solution \widehat{y} . To find it we can reduce the problem to solve a family of ODEs as follows: write

$$(36) \quad \widehat{y}(\mathbf{x}) = \sum_{\boldsymbol{\alpha} \preceq \boldsymbol{\beta}} \mathbf{x}^{\boldsymbol{\beta}} \widehat{y}_{\boldsymbol{\beta}}(\mathbf{x}^{\boldsymbol{\alpha}}), \quad c(\mathbf{x}) = \sum_{\boldsymbol{\alpha} \preceq \boldsymbol{\beta}} \mathbf{x}^{\boldsymbol{\beta}} c_{\boldsymbol{\beta}}(\mathbf{x}^{\boldsymbol{\alpha}}),$$

as power series with coefficients series in $\mathbf{x}^{\boldsymbol{\alpha}}$, according to the decomposition $c_{\boldsymbol{\beta}}(t) = \sum_{n=0}^{\infty} a_{n\boldsymbol{\alpha}+\boldsymbol{\beta}} t^n$. Then, plug \widehat{y} into the equation and equate the common terms in $\mathbf{x}^{\boldsymbol{\beta}}$. It follows the initial problem is equivalent to solve the family of independent ODEs

$$\langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle t^2 \widehat{y}'_{\boldsymbol{\beta}}(t) = (\mu - \langle \boldsymbol{\lambda}, \boldsymbol{\beta} \rangle t) \widehat{y}_{\boldsymbol{\beta}}(t) - c_{\boldsymbol{\beta}}(t), \quad \boldsymbol{\alpha} \preceq \boldsymbol{\beta}.$$

Thus each $\widehat{y}_{\boldsymbol{\beta}}$ is uniquely determined and generically t -1-Gevrey.

Let us consider two explicit examples. First, take $c(\mathbf{x}) = \mathbf{x}^{\boldsymbol{\beta}}$. If $\langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle = 0$, the solution is $\widehat{y}(\mathbf{x}) = \frac{\mathbf{x}^{\boldsymbol{\beta}}}{1 - \langle \boldsymbol{\lambda}, \boldsymbol{\beta} \rangle \mathbf{x}^{\boldsymbol{\alpha}}}$ which is convergent. Otherwise, after some calculations we find the solution

$$\widehat{y}(\mathbf{x}) = \sum_{n=0}^{\infty} \binom{-\langle \boldsymbol{\lambda}, \boldsymbol{\beta} \rangle / \langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle}{n} (-1)^n n! \frac{\langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle^n}{\mu^{n+1}} \mathbf{x}^{n\boldsymbol{\alpha}+\boldsymbol{\beta}},$$

which is $\mathbf{x}^{\boldsymbol{\alpha}}$ -1-Gevrey. In fact, \widehat{y} is $\mathbf{x}^{\boldsymbol{\alpha}}$ -1-summable in direction θ , see [5], with sum given by

$$y_{\theta}(\mathbf{x}) = \mathbf{x}^{\boldsymbol{\beta}-\boldsymbol{\alpha}} \int_0^{e^{i\theta}\infty} (1 + \langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle \xi / \mu)^{-\langle \boldsymbol{\lambda}, \boldsymbol{\beta} \rangle / \langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle} e^{-\xi / \mathbf{x}^{\boldsymbol{\alpha}}} d\xi, \quad \text{for } \theta \neq \arg(-\mu / \langle \boldsymbol{\lambda}, \boldsymbol{\alpha} \rangle).$$

Note \widehat{y} reduces to a polynomial when $\langle \boldsymbol{\lambda}, m\boldsymbol{\alpha} + \boldsymbol{\beta} \rangle = 0$, for some $m \geq 0$.

As second example consider $x_1 x_2 (x_1 \partial_{x_1} y - x_2 \partial_{x_2} y) = \mu y - (1 - x_1)^{-1} (1 - x_2)^{-1}$. Then, decomposition (36) takes the form

$$\widehat{y}(x_1, x_2) = \sum_{n=0}^{\infty} y_{(n,0)}(x_1 x_2) x_1^n + \sum_{n=1}^{\infty} y_{(0,n)}(x_1 x_2) x_2^n, \quad \sum_{n,m \geq 0} x_1^n x_2^m = \frac{1}{1 - x_1 x_2} + \sum_{n=1}^{\infty} \frac{x_1^n + x_2^n}{1 - x_1 x_2},$$

and we find the coefficients are equal to

$$y_{(n,0)}(t) = \frac{1}{(1-t)(\mu - nt)}, \quad y_{(0,n)}(t) = \frac{1}{(1-t)(\mu + nt)}, \quad \text{valid for } |t| < \frac{|\mu|}{n}.$$

By using the Taylor series at the origin of the previous functions we can determine \widehat{y} . The relation between \widehat{y} and the solution

$$y_0(x_1, x_2) = \frac{1}{1 - x_1 x_2} \sum_{n=0}^{\infty} \frac{x_1^n}{\mu - n x_1 x_2} + \frac{1}{1 - x_1 x_2} \sum_{n=1}^{\infty} \frac{x_2^n}{\mu + n x_1 x_2},$$

which is analytic on $\{(x_1, x_2) \in \mathbb{C}^2 : |x_1|, |x_2| < 1, x_1 x_2 \neq \mu/n, n \geq 1\}$, is that y_0 is the $x_1 x_2$ -1-sum of \widehat{y} , c.f. [11, Example 2.1] for more details on similar calculations with these series.

Theorem 1.1 can also be applied in other situations after suitable changes of variables of blow-up type. We illustrate this fact in the following example.

Example 6.4. Consider the system of PDEs given by

$$(37) \quad x_1^{p_1+1} c_1(\mathbf{x}) \partial_{x_1} \mathbf{y} + \dots + x_d^{p_d+1} c_d(\mathbf{x}) \partial_{x_d} \mathbf{y} = \mathbf{F}(\mathbf{x}, \mathbf{y}),$$

where $\mu = \frac{\partial \mathbf{F}}{\partial \mathbf{y}}(\mathbf{0}, \mathbf{0})$ is invertible, and we assume $1 \leq p_1 \leq p_j$, for all j . The system has a unique formal power series solution $\widehat{\mathbf{y}}$, and we show it is a $(1/p_1, \dots, 1/p_1)$ -Gevrey series: by using the punctual blow-up (20), we find

$$z_1 \partial_{z_1} = x_1 \partial_{x_1} + \dots + x_d \partial_{x_d}, \quad z_j \partial_{z_j} = x_j \partial_{x_j}, \quad j = 2, \dots, d,$$

thus (37) takes the form

$$z_1^{p_1} L'(\mathbf{u}) = z_1^{p_1+1} c'_1(\mathbf{z}) \partial_{z_1} \mathbf{u} + z_1^{p_1} \sum_{j=2}^d (z_1^{p_j-p_1} z_j^{p_j} c'_j(\mathbf{z}) - c'_1(\mathbf{z})) z_j \partial_{z_j} \mathbf{u} = \mathbf{F}'(\mathbf{z}, \mathbf{u}),$$

where $c'_j(\mathbf{z}) = c_j(\mathbf{x})$, $\mathbf{F}'(\mathbf{z}, \mathbf{u}) = \mathbf{F}(\mathbf{x}, \mathbf{y})$, and $\mathbf{u}(\mathbf{z}) = \mathbf{y}(\mathbf{x})$. Since $L'(z_1^{p_1}) = p_1 z_1^{p_1-1} c'_1(\mathbf{z})$ and $\frac{\partial \mathbf{F}'}{\partial \mathbf{z}}(\mathbf{0}, \mathbf{0}) = \mu$ is invertible, Theorem 1.1 implies that $\widehat{\mathbf{u}}(\mathbf{z}) = \widehat{\mathbf{y}}(\mathbf{x})$ is a $z_1^{p_1}$ -1-Gevrey series, and the proof of Proposition 4.4 shows $\widehat{\mathbf{y}}$ is a $(1/p_1, \dots, 1/p_1)$ -Gevrey series as desired.

This is the generic situation, as we can exemplify with the scalar PDE

$$x_1^2 \partial_{x_1} y + \dots + x_d^2 \partial_{x_d} y + y = \mathbf{x}^1, \quad \widehat{\mathbf{y}}(\mathbf{x}) = \sum_{\beta \in \mathbb{N}^d} (-1)^{|\beta|} |\beta|! \mathbf{x}^{\beta+1},$$

where $\mathbf{1} = (1, \dots, 1)$, that can be seen as a multidimensional Euler's equation. On the other hand, it is worth remarking (37) has been recently studied by Z. Luo, H. Chen and C. Zhang [21] for the case

$$x_1^2 c_1(\mathbf{x}) \partial_{x_1} u + x_2^2 c_2(\mathbf{x}) \partial_{x_2} u = b(\mathbf{x}) u - a(\mathbf{x}),$$

where a, b, c_1, c_2 are analytic near $\mathbf{0} \in \mathbb{C}^2$ and $b(\mathbf{0})c_1(\mathbf{0})c_2(\mathbf{0}) \neq 0$. This scalar equation has a unique formal power series solution \hat{u} which is Borel-summable in the variables (x_1, x_2) . In particular, \hat{u} is $(1, 1)$ -Gevrey as we showed before.

Example 6.5. Consider parametric families of ODEs unfolding $k+1$ singularities

$$(38) \quad (x^{k+1} - \varepsilon) \frac{d\mathbf{y}}{dx} = \mu \mathbf{y} - f(x, \varepsilon, \mathbf{y}),$$

where k is a positive integer, μ is an invertible matrix, f is analytic near the origin in $\mathbb{C} \times \mathbb{C} \times \mathbb{C}^d$, $\frac{\partial f}{\partial \mathbf{y}}(0, 0, \mathbf{0}) = \mathbf{0}$, and $\varepsilon \in (\mathbb{C}, 0)$ is a small parameter. These systems have been studied by M. Klimeš [19] for the case $k=1$ by using an adapted (unfolded) Borel-Laplace method in order to obtain parametric solutions defined and bounded on certain ramified domains attached to both singularities $x = \pm \sqrt{\varepsilon}$, at which they possess a limit in a spiraling manner.

As it is remarked in Section 2.4 of [19], the system above has a unique formal power series solution $\widehat{\mathbf{y}}$ which is $(\frac{1}{k}, \frac{k+1}{k})$ -Gevrey in (x, ε) . We can prove this readily as follows: consider the ramification $\varepsilon = \eta^{k+1}$, and afterwards the punctual blow-up $x = z, \eta = z\zeta$. In these coordinates equation (38) takes the form

$$z^k \left(z \frac{\partial \mathbf{u}}{\partial z} - \zeta \frac{\partial \mathbf{u}}{\partial \zeta} \right) = (1 - \eta^{k+1})^{-1} (\mu \mathbf{u} - f(z, z^{k+1} \zeta^{k+1}, \mathbf{u})),$$

where $\mathbf{u}(z, \zeta) = \mathbf{y}(z, z^{k+1} \zeta^{k+1}) = \mathbf{y}(x, \varepsilon)$. By applying Theorem 1.1 to $P = z^k$ and $L = z \partial_z - \zeta \partial_\zeta$ we find $\widehat{\mathbf{u}}(z, \zeta) = \widehat{\mathbf{y}}(x, \varepsilon)$ is z^k -1-Gevrey, since $L(P) = k z^k$. Thus, $\widehat{\mathbf{y}}(x, \eta^{k+1})$ is $(\frac{1}{k}, \frac{1}{k})$ -Gevrey in (x, η) , and therefore $\widehat{\mathbf{y}}(x, \varepsilon)$ is $(\frac{1}{k}, \frac{k+1}{k})$ -Gevrey in (x, ε) as we claimed.

REFERENCES

- [1] J. M. Aroca, H. Hironaka and J. L. Vicente, *The Theory of the Maximal Contact*. Memorias de Matemática del Instituto "Jorge Juan" 29, Instituto "Jorge Juan" de Matemáticas, Consejo Superior de Investigaciones Científicas, Madrid, 1975.
- [2] W. Balser and J. Mozo-Fernández, *Multisummability of formal solutions of singular perturbation problems*. J. Differential Equations, vol. 183, (2002) 526–545.

- [3] W. Balsler and V. Kostov, *Singular perturbation of linear systems with a regular singularity*. J. Dynam. Control. Syst. 8, 3 (2002) 313–322.
- [4] B. L. J. Braaksma and L. Stolovitch, *Small Divisors and Large Multipliers*. Ann. Inst. Fourier 57, no. 2 (2007) 603–628.
- [5] M. Canalis-Durand, J. Mozo-Fernández and R. Schäfke, *Monomial summability and doubly singular differential equations*. J. Differential Equations, vol. 233, (2007) 485–511.
- [6] M. Canalis-Durand, J.P. Ramis, R. Schäfke and Y. Sibuya, *Gevrey solutions of singularly perturbed differential equations*. J. Reine Angew. Math, vol. 518, (2000) 95–129.
- [7] S.A. Carrillo and J. Mozo-Fernández, *An extension of Borel-Laplace methods and monomial summability*. J. Math. Anal. Appl, vol. 457, Issue 1, (2018) 461–477.
- [8] S.A. Carrillo, J. Mozo-Fernández and R. Schäfke, *Tauberian theorems for summability in analytic functions*. Submitted to publication. Available at <https://arxiv.org/abs/1903.08898>.
- [9] S.A. Carrillo and F. Sanz, *Briot-Bouquet’s theorem in high dimension*. Publ. Mat. 58 suppl. (2014) 135–152.
- [10] S.A. Carrillo, *Summability in a monomial for some classes of singularly perturbed partial differential equations*. Submitted to publication. Available at <https://arxiv.org/abs/1803.06719>.
- [11] S.A. Carrillo and J. Mozo-Fernández, *Tauberian properties for monomial summability with applications to Pfaffian systems*. J. Differential Equations vol. 261 (2016) 7237–7255.
- [12] O. Costin and S. Tanveer, *Nonlinear evolution PDEs in $\mathbb{R}^+ \times \mathbb{C}^d$: Existence and uniqueness of solutions, asymptotic and Borel summability properties*. Annales De L’Institut Henri Poincaré (C) Analyse Non Linéaire 25 (2007) 795–823.
- [13] O. Costin and S. Tanveer, *Short time existence and Borel summability in the Navier–Stokes equation in \mathbb{R}^3* . Comm. Partial Differential Equations 34(7–9) (2009) 785–817.
- [14] M. Hibino, *Divergence property of formal solutions for singular first order linear partial differential equations*. Publ. Res. Inst. Math. Sci. 35 (1999), no. 6, 893–919.
- [15] M. Hibino, *Formal Gevrey theory for singular first order semi-linear partial differential equations*. Osaka J. Math. 41, no. 1 (2004) 159–191.
- [16] M. Hibino, *Borel Summability of Divergent Solutions for Singular First-order Partial Differential Equations with Variable Coefficients. Part I*. J. Differential Equations 227, no. 2 (2006) 499–533.
- [17] M. Hibino, *Borel Summability of Divergent Solutions for Singular First-order Partial Differential Equations with Variable Coefficients. Part II*. J. Differential Equations 227, no. 2 (2006) 534–563.
- [18] S. Kaplan, *Formal and convergent power series solutions of singular partial differential equations*. Trans. Amer. Math. Soc. 256 (1979) 163–183.
- [19] M. Klimeš, *Confluence of Singularities of Nonlinear Differential Equations via Borel-Laplace Transformations*. J. Dyn. Control Syst. 22 (2016) 285–324.
- [20] A. Lastra and S. Malek, *On parametric Gevrey asymptotics for some nonlinear initial value Cauchy problems*. J. Differential Equations 259, no. 10 (2015) 5220–5270.
- [21] Z. Luo, H. Chen, and C. Zhang, *On the summability of divergent power series satisfying singular PDEs*. C. R. Math. Acad. Sci. Paris 357, no. 3 (2019) 258–262.
- [22] J.-F. Mattei and R. Moussu, *Holonomie et intégrales premières*. Ann. Sci. École Norm. Sup. 4, no. 13 (1980) no. 4, 469–523.
- [23] J. Mozo-Fernández and R. Schäfke, *Asymptotic expansions and summability with respect to an analytic germ*. Publ. Mat. 63 (2019) 3–79.
- [24] M. Nagumo, *Über das anfangswertproblem partieller differentialgleichunge*. Jap. J. Math. 18 (1942) 41–47.
- [25] T. Oshima, *On the theorem of Cauchy-Kowalevsky for first order linear differential equations with degenerate principal symbols*. Proc. Japan Acad. Ser. A Math. Sci. 49 (1973) 83–87.
- [26] S. Ōuchi, *Borel summability of formal solutions of some first order singular partial differential equations and normal forms of vector fields*. J. Math. Soc. Japan 57, no. 2 (2005) 415–460.
- [27] S. Ōuchi, *Multisummability of formal solutions of some linear partial differential equations*. J. Differential Equations 185, no. 2 (2002) 513–549.
- [28] A. Rainer and G. Schindl, *Equivalence of stability properties for ultradifferentiable function classes*. Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM 110, no. 1 (2016) 17–32.
- [29] Y. Sibuya, *Convergence of formal solutions of meromorphic differential equations containing parameters*. Funkcial. Ekvac. 37 (1994) 395–400.
- [30] V. Thilliez, *On Quasianalytic Local Rings*. Expo. Math. 26, no. 1 (2008) 1–23.
- [31] H. Tahara and H. Yamazawa, *Multisummability of formal solutions to the Cauchy Problem for some linear partial differential equations*. J. Differential Equations 255, no. 10 (2013) 3592–3637.
- [32] H. Yamazawa, *Formal Gevrey Class of formal power series solution for singular first order linear partial differential operators*. Tokyo J. Math. 23, no. 2 (2000) 537–561.
- [33] H. Yamazawa and M. Yoshino, *Parametric Borel summability for some semilinear system of partial differential equations*. Opuscula Math. 35, no. 5 (2015) 825–845.

ESCUELA DE CIENCIAS EXACTAS E INGENIERÍA, UNIVERSIDAD SERGIO ARBOLEDA, CALLE 74, # 14-14, BOGOTÁ, COLOMBIA.

E-mail address: `sergio.carrillo@usa.edu.co`

UNIVERSIDAD PRIVADA DEL NORTE, CAMPUS BREÑA, AVENIDA TINGO MARIA 1122, LIMA, PERÚ

E-mail address: `carlos.hurtado@upn.edu.pe`