

THRESHOLD FOR BLOWUP AND STABILITY FOR NONLINEAR SCHRÖDINGER EQUATION WITH ROTATION

NYLA BASHARAT, HICHEM HAJAIEJ, YI HU AND SHIJUN ZHENG

ABSTRACT. We consider the focusing NLS with an angular momentum and a harmonic potential, which models Bose-Einstein condensate under a rotating magnetic trap. We give a sharp condition on the global existence and blowup in the mass-critical case. We further consider the stability of such systems via variational method. We determine that at the critical exponent $p = 1 + 4/n$, the mass of Q , the ground state for the NLS with zero potential, is the threshold for both finite time blowup and orbital instability. Moreover, we prove similar results for the rotational NLS with an inhomogeneous nonlinearity. The analysis relies on the existence of ground state as well as a virial identity for the associated kinetic-magnetic operator.

CONTENTS

1. Introduction	2
2. Preliminaries	7
3. Virial identity for NLS with rotation	10
4. Blowup criterion for $\Omega \neq 0$	14
5. Proof of Theorem 1.1	16
6. Virial identity for inhomogeneous NLS	20
7. Ground state solutions	23
8. Orbital stability of standing waves	29
9. Concluding remarks	32
References	34

Date: January 30, 2020.

2010 Mathematics Subject Classification. 35Q55, 37K45, 35P25.

Key words and phrases. NLS, angular momentum, ground states, blowup, orbital stability.

1. INTRODUCTION

The Cauchy problem for the nonlinear Schrödinger equation (NLS), or Gross-Pitaevskii equation with rotation in $(t, x) \in \mathbb{R}^{1+n}$, $n \geq 2$ reads

$$(1) \quad \begin{aligned} iu_t &= -\frac{1}{2}\Delta u + V(x)u - \lambda|u|^{p-1}u + L_\Omega u \\ u(0, x) &= u_0 \in \mathcal{H}^1, \end{aligned}$$

where $p \in [1, 1 + 4/(n-2))$, $V(x) := \frac{1}{2}\gamma^2|x|^2$ is the harmonic potential with frequency $\gamma > 0$ that models the magnetic trap and λ is a positive constant. Adding a spin symmetry, the angular momentum operator is denoted by $L_\Omega := -\Omega L_z = iA \cdot \nabla$, where $\Omega \in \mathbb{R}$ is the rotation frequency, $L_z = i(x_2\partial_{x_1} - x_1\partial_{x_2})$ and $A = \Omega\langle -x_2, x_1, 0, \dots, 0 \rangle$. The weighted Sobolev space is given by $\mathcal{H}^s = \mathcal{H}^{s,2}$, where

$$\mathcal{H}^{s,r} := \{f \in L^r(\mathbb{R}^n) : (-\Delta)^{s/2}f \in L^r, \langle x \rangle^s f \in L^r\},$$

with $\langle x \rangle = (1 + |x|^2)^{1/2}$. The weighted Sobolev space arises naturally as a suitable space in this context.

When $n = 2, 3$, the NLS above models Bose-Einstein condensation with rotation [1, 4, 5, 9, 35, 44], which is a remarkable system arising in optics, plasma, superfluids, spinor particles, quantized vortices and surface waves. Extensions to higher dimensions can be found in [6, 21]. Mathematically, it can be viewed as the mean field limit of rotating many-body bosons in a confining trap [31, 32]. Equation (1) can be formally derived from

$$i\frac{\partial u}{\partial t} = \frac{\delta H}{\delta \bar{u}},$$

where H is the associated Hamiltonian

$$H[u] = \int \left(\frac{1}{2}|\nabla u|^2 + V|u|^2 - \frac{2\lambda}{p+1}|u|^{p+1} - \Omega \bar{u}L_z u \right).$$

It is desirable to provide rigorous mathematical descriptions for the rotating BEC model. As is known, the well-posedness and blowup for the standard NLS have been extensively studied for a few decades in the energy sub-critical regime. For the rotational NLS (RNLS) in (1), the local wellposedness (l.w.p.) has been considered in e.g. [2, 13, 27] when $n = 2, 3$, and [16, 20, 39, 50] for general magnetic NLS (mNLS), to list a few. RNLS (1) can be written in the magnetic form (5), thus the wellposedness follows from that of mNLS if $p \in (1, 1 + 4/(n-2))$ in view of Proposition 2.1.

In the mass-subcritical case $p < 1 + 4/n$ and $\lambda > 0$, it is known that all \mathcal{H}^1 -solutions of equation (1) exist globally in time. However, if

$p \in [1 + \frac{4}{n}, 1 + \frac{4}{n-2})$, there exist finite time blowup solutions for (1), see e.g. [2, 8, 10, 23]. Such situation becomes more subtle and complex, where the occurrence of global existence or wave collapse depends on both the size and the profile of the initial data. Thus, it is an open question to find a threshold condition that distinguishes the g.w.p. and blowup for the RNLS.

In this paper, we are mainly concerned with the mass-critical case $p = 1 + 4/n$. Inspired by the work in [45] and [49], we address this so-called “minimal mass blowup” problem for (1). To our knowledge, previous techniques and ideas do not directly apply to solving this threshold problem. One delicate issue is that the threshold profile for (1) turns out *not* to be the minimizer for the associated energy! This is one essential reason that makes the problem much more challenging. Technically, in the presence of V and L_Ω , some of the key symmetries for the standard NLS (including scaling and translation invariance) are broken, whose geometry of the trajectories of motion of particles has the effect on relevant physical phenomenon of wave collapse as well as stability of solitons.

In our first main theorem, we give a sharp threshold in terms of the unique positive radial ground state $Q = Q_{\lambda,1}$ in the usual Sobolev space $H^1(\mathbb{R}^n)$ if $p = 1 + 4/n$:

$$(2) \quad -\frac{1}{2}\Delta Q - \lambda|Q|^{p-1}Q = -Q.$$

The existence of Q is well-known [29, 37, 45].

Theorem 1.1 (threshold for g.w.p. and blowup). *Let $p = 1 + \frac{4}{n}$ and $\lambda > 0$. Let (Ω, γ) be any given pair in $\mathbb{R} \times \mathbb{R}_+$. Suppose $u_0 \in \mathcal{H}^1$.*

- (a) *If $\|u_0\|_2 < \|Q\|_2$, then there exists a unique global in time solution u of (1) in $C(\mathbb{R}, \mathcal{H}^1) \cap L_{loc}^{2+\frac{4}{n}}(\mathbb{R}, L^{2+\frac{4}{n}})$.*
- (b) *The condition in (a) is sharp in the sense that for all $c \geq \|Q\|_2$, there exists u_0 in \mathcal{H}^1 satisfying $\|u_0\|_2 = c$ such that (1) has a finite time blowup solution.*

When $n = 2$, the threshold value in the theorem can be evaluated at $\lambda^{-\frac{1}{2}}\|Q_{1,1}\|_2 = \lambda^{-\frac{1}{2}}(\pi \cdot 1.86225 \cdots)^{\frac{1}{2}}$, see [25, 45]. For blowup solutions, in general, the wave collapse depends on the delicate balance between kinetic and potential energies (linear vs. nonlinear) as well as angular momentum for the profile of the solution, under the constriction of mass and energy conservation.

Theorem 1.1 shows that (1) has the same minimal mass $\|Q\|_2^2$ for blowup as in the free case $\Omega = V = 0$. The heuristic reason is that in (1) neither V nor $-\Omega L_z$ sees the scaling. Note that, however, at the

threshold, $u_0 = Q$ leads to a soliton solution in the absence of potential while the same initial data leads to blowup solution for (1).

The proof of Theorem 1.1 relies on a sharp criterion in part (a) of Lemma 4.1, where we prove the blowup for (1) for all Ω, γ if $p = 1 + 4/n$, namely, u blows up in finite time provided $E_{0,0}(u_0) = \frac{1}{2} \int |\nabla u_0|^2 - \frac{\lambda n}{n+2} \int |u_0|^{2+\frac{4}{n}} \leq 0$. This lemma is sharp in the sense that it dictates the blowup for initial data $u_0 = Q$ with $E_{0,0}(Q) = 0$ such that the Q -blowup profile is attainable, which is not covered by existing results in the literature as far as the authors know. In Lemma 4.1 (c)-(d) we also obtain some blowup conditions if $p > 1 + 4/n$, which allows us to show in Proposition 5.3 a blowup result above the ground state level. Our approach is mainly motivated by the treatment in [45] and [49], where blowup results were proven with the same minimal mass in the cases $\Omega = V = 0$ and $\Omega = 0, V = |x|^2$ respectively. See also related discussions in [10, 11, 16] for NLS with a sub-quadratic potential basically under the condition $|\Omega| < \gamma$.

In Section 6, we further consider the threshold problem for the inhomogeneous equation (41). In doing so, we derive a virial type identity for (41), then we obtain Theorem 6.6, an analogue of Theorem 1.1 by following the same line of proof of the theorem presented in Section 5. As is commented in Remark 6.7, concerning the exact threshold, there remains a gap $\|Q_{\lambda_{max}}\|_2 < \|Q_{\lambda_{min}}\|_2$ in Theorem 6.6. This suggests that an improvement on the blowup criterion lemma, an inhomogeneous version of Lemma 4.1 would be needed.

Observe that the threshold in Theorem 1.1 is valid if V is an isotropic harmonic potential. When V is anisotropic, it remains an open question in the “fast rotating” regime, where $|\Omega| > \underline{\gamma} := \min_{1 \leq j \leq n} (\gamma_j)$, concerning the threshold on finite time blowup for either $p = 1 + 4/n$ or $p > 1 + 4/n$, see some numerical results in [4, 5]. In physics, this question concerns the scenario where the rotation frequency is stronger than the trapping frequency, in which case it would be worthwhile to study the behavior of minimal mass wave collapse quantitatively.

Here we would like to briefly review the previous work on blowup for RNLS. The initial result on the blowup for (1) was obtained in [23] in the case $|\Omega| = \gamma, n = 3$ if $p > 1 + 4/n$. Recent blowup results were proven in [2] in the case for all Ω, γ if $p \geq 1 + 4/n, n = 2, 3$ and in [8, 21] for general electromagnetic potentials. However, these results only give a general sufficient condition, not a *sharp criterion* to address the threshold, or minimal mass blowup problem. The pre-existing results mainly assume either $E_{\Omega,V}(u_0)$ or $E_{0,V}(u_0)$ is negative, from which follows the blowup of the solution based on a virial identity, a convexity

argument for the variance $J(t) = \int |x|^2 |u|^2$. Such assumption seems too strong in the mass-critical case. Notice that some generic data does not satisfy this condition: e.g. $E_{\Omega, V}(Q) = E_{0, V}(Q) > 0$. This example indicates that we need a sharper blowup condition. Lemma 4.1 thus provides such criteria with weaker conditions. The proof of the lemma is based on the magnetic virial identity (Lemma 3.1) and an explicit solution of the o.d.e. (22) for $J(t)$. Then, the blowup time can be located by examining the first zero of $J(t)$ in a more accurate way. Consequently, the lemma enable us to apply the sharp Gagliardo-Nirenberg inequality involving the ground state given by (2) to prove Theorem 1.1 for all (Ω, γ) in all dimensions. Note that if $p = 1 + 4/n$, by evaluating the derivative of $J(t)$ at its zero $t = T_*$, we show in passing a lower bound of the blowup rate $\|\nabla u\|_2 \gtrsim \|u_0\|_2 (T_* - t)^{-1/2}$ in (26).

The approach along with the techniques elaborated above allows one to prove similar result in Section 6 for the inhomogeneous equation (41). We first prove a local virial identity in Lemma 6.4. An alternative proof of this lemma can be performed through the method from a hydrodynamical system [1, 2]. Then, we prove Lemma 6.5, the analogue of Lemma 4.1 (a) for equation (41), in which case we encounter the difficulty where the solution of the o.d.e. (45) for $J(t)$ is not solvable. We overcome this obstacle by writing the unknown $J(t)$ in an implicit integral equation so that we are able to obtain some good estimate for $J(t)$ as shown in (47), which leads to establishing Lemma 6.5 for the inhomogeneous RNLS.

Concerning the blowup rate (26) shown in the proof for RNLS (1), we add that when $p = 1 + 4/n$ and $\|u_0\|_2 = \|Q\|_2$, like in the cases where $\Omega = V = 0$ and $\Omega = 0, V = |x|^2$ in [36] and [11] respectively, via the \mathcal{R} -transform, a pseudo-conformal type transform, we were able to determine the profile for all blowup solutions with minimal mass at the ground state level. Hence all such blowup solutions in \mathcal{H}^1 have blowup rate $(T_* - t)^{-1}$, which is however unstable, see [6, Proposition 4.5]. When $p = 1 + 4/n$ and $\|u_0\|_2$ is slightly greater than $\|Q\|_2$, N.B., Y.H. and S.Z. proved in [6] the log-log law for RNLS (1), i.e., there exists a universal constant $\alpha^* := \alpha_n^* > 0$ such that if $\int |Q|^2 < \int |u_0|^2 < \int |Q|^2 + \alpha^*$ with negative energy $E_{0,0}(u_0) < 0$, then $u \in C([0, T_*); \mathcal{H}^1)$ is a blowup solution to (1) on its lifespan $[0, T_*)$ satisfying

$$(3) \quad \|\nabla u(t, \cdot)\|_2 = \frac{\|\nabla Q\|_2}{\sqrt{2\pi}} \sqrt{\frac{\log |\log(T_* - t)|}{T_* - t}} + o(1), \quad \text{as } t \rightarrow T_*$$

where Q is the unique solution of (2). The analogue for the standard NLS was initially proven in Merle and Raphaël [38] under the hypothesis of certain spectral property conjecture. We can now state the log-log law (3) for RNLS (1) when $n \leq 12$, in light of the recent numerical verification of the spectral property conjecture in [48]. Such a blowup rate is also known to be stable in \mathcal{H}^1 . An example of the initial data is given in the form $u_0 = \eta Q$ with $1 < \eta < \sqrt{1 + \frac{\alpha^*}{\|Q\|_2^2}}$, which can be easily verified to satisfy the condition above by the Pohozaev identity (29).

In the second component of this paper we consider the stability of *ground state solutions* for (1). A *ground state solution* (g.s.s.) at mass level $c > 0$ is a minimizer of the energy functional with constant mass constraint that is defined by Definition 7.1. For the usual focusing NLS

$$(4) \quad iu_t = -\frac{1}{2}\Delta u - \lambda|u|^{p-1}u, \quad u(0, x) = u_0 \in H^1,$$

the ground state Q is unique and orbitally stable if $p < 1 + 4/n$, and unstable if $p \geq 1 + 4/n$. In Section 7, in the slow rotating setting $|\Omega| < \gamma$, we show the existence of g.s.s. for the RNLS (48) with more general class of nonlinearities. Our construction of the g.s.s relies on the concentration compactness method in [14] and [26]. Then in Section 8, we prove the orbital stability for (48) via standard argument. Theorem 8.3 shows that Z_c , the set of g.s.s. for (48) is orbitally stable if either $p < 1 + 4/n$, or $p = 1 + 4/n$ and the mass level is below that of $Q_{\lambda,1}$ for some optimal constant λ in the focusing case.

Theorem 1.2. *Let $p = 1 + 4/n$ and $|\Omega| < \gamma$. Let Q be the unique ground state of (2). Suppose $c < \|Q\|_2$. Then the set of minimizers Z_c is orbitally stable for (1).*

This result is part (b) in Theorem 8.3, which suggests that the “minimal mass” for the blowup is the same threshold for orbital stability problem under the condition $|\Omega| < \gamma$. When $p = 1 + 4/n$, the energy functional $E_{\Omega,\gamma}(u)$ for (1) is unbounded from below on the mass level set S_{c_0} with $c_0 = \|Q\|_2$, thus, an absolute minimum of the energy does not exist on S_{c_0} , see [24, 25]. This also indicates the existence of the threshold we have obtained in Theorem 1.1 and Theorem 1.2. Concerning mass-subcritical case $p < 1 + 4/n$, the orbital stability was proven in [13, 17] if $|\Omega| = \gamma$, and recently in [3] if $|\Omega| < \gamma$, $n = 2, 3$. The frequency condition $|\Omega| \leq \gamma$ is critical to guarantee stability. Physically, if the angular velocity of rotation exceeds the trapping frequency, that is, $|\Omega| > \gamma$, then V cannot provide the necessary centripetal force (that counteract the centrifugal force caused by the rotation), and the gas

may fly apart. The counterexample we show in Section 8 agrees with physics observation. For attractive particle interactions, it may lead to centrifugal forces destabilizing all rotating states.

2. PRELIMINARIES

One can write (1) in the form of mNLS

$$(5) \quad iu_t = -\frac{1}{2}(\nabla - iA)^2 u + V_e u - \lambda|u|^{p-1}u,$$

where V_e denotes the effective electric potential for the Hamiltonian operator $H_{A,V} = -\frac{1}{2}(\nabla - iA)^2 + V_e$. In particular, if $V = \frac{\gamma^2}{2}|x|^2$,

$$(6) \quad A = \Omega \langle -x_2, x_1, 0, \dots, 0 \rangle,$$

then $V_e = \frac{1}{2}(\gamma^2 - \Omega^2)(x_1^2 + x_2^2) + \frac{\gamma^2}{2}x_3^2 + \dots + \frac{\gamma^2}{2}x_n^2$ and $iA \cdot \nabla = -\Omega L_z = L_\Omega$. Define

$$(7) \quad H_{\Omega,V} := -\frac{1}{2}\Delta + V - \Omega L_z.$$

Then $H_{\Omega,V} = H_{A,V}$. In \mathbb{R}^3 , the operator L_Ω generates a rotation in the sense that: If (r, θ, z) is the cylindrical coordinate, then $e^{-itL_\Omega} f(r, \theta, z) = f(r, \theta + t\Omega, z)$.

Now let us review some results on l.w.p and g.w.p. for the Cauchy theory for (1) in the energy subcritical regime. For $1 \leq p < 1 + 4/(n-2)$, i.e., the \mathcal{H}^1 -subcritical case, the local existence and uniqueness for (1) follow from those of the magnetic NLS (5), based on the fundamental solution constructed in [47], see [16, 39]. The \mathcal{H}^s -subcritical result for (5) was considered in [50] for $1 \leq p < 1 + 4/(n-2s)$ if $V_e(x)$ is subquadratic and bounded from below. The local wellposedness for $V_e(x) = -\sum_j \tilde{\gamma}_j^2 x_j^2$ follow from a Strichartz estimate in Lemma 2.4, see Proposition 2.1. In the mass-subcritical case $p < 1 + 4/n$ with any data in L^2 and in the mass-critical case $p = 1 + 4/n$ with small data in L^2 , the global in time solution exists and is unique [20, 50] in \mathbb{R}^n .

Proposition 2.1. *Let $1 \leq p < 1 + \frac{4}{n-2}$. Let $u_0 \in \mathcal{H}^1$, $r = p + 1$ and $q = \frac{2(p+1)}{p-1}$.*

- (a) *(local existence) Then there exist a maximal time interval $I = (-T_{min}, T_{max})$, $T_{max}, T_{min} > 0$ such that (1) has a unique solution $u \in C(I, \mathcal{H}^1) \cap L^q(I, \mathcal{H}^{1,r})$.*
- (b) *(global existence) There exists a unique, \mathcal{H}^1 -bounded global solution in $C(\mathbb{R}, \mathcal{H}^1) \cap L^q_{loc}(\mathbb{R}, \mathcal{H}^{1,r})$ if one of the following conditions is satisfied:*
 - (i) $1 < p < 1 + \frac{4}{n-2}$, $\lambda < 0$ (defocusing),

- (ii) $1 \leq p < 1 + \frac{4}{n}$, $\lambda > 0$ (*focusing*),
- (iii) $1 + \frac{4}{n} \leq p < 1 + \frac{4}{n-2}$, $\lambda > 0$ (*focusing*), $\|u_0\|_2 < \varepsilon$ for some $\varepsilon = \varepsilon(\lambda, n)$ sufficiently small.
- (c) If T_{max} (respectively T_{min}) is finite, then $\|\nabla u\|_2 \rightarrow \infty$ as $t \rightarrow T_{max}$ (respectively $-T_{min}$).
- (d) On the lifespan interval $(-T_{min}, T_{max})$, the following quantities are conserved in time:

(8)

$$(mass) \quad M(u) = \int |u|^2$$

(9)

$$(energy) \quad E_{\Omega, V}(u) = \int \left(\frac{1}{2} |\nabla u|^2 + V|u|^2 - \frac{2\lambda}{p+1} |u|^{p+1} \right) + \langle L_{\Omega} u, u \rangle$$

- (e) The angular momentum $\ell_{\Omega}(u) := \langle L_{\Omega} u, u \rangle$ is real-valued and

$$(10) \quad \langle L_{\Omega} u, u \rangle = -\Omega \int \bar{u} L_z u,$$

where the inner product is defined as $\langle f, g \rangle := \int f \bar{g}$.

The continuity and conservation laws for the solution map $u_0 \mapsto u$ in \mathcal{H}^1 follow from a standard argument using the Duhamel formula [16]. The above results extend to l.w.p. and g.w.p. in \mathcal{H}^k for (1) through a similar proof in view of Lemma 2.4.

The local in time result for (5) requires the dispersive and Strichartz estimates for the time-dependent propagator $U(t) = e^{-itH_{A,V}}$. Yajima [47] combines the oscillatory integral operators, bicharacteristics and integral equation method developed by Fujiwara and Kitada et al's work to obtain

$$U(t)f(x) = (2\pi it)^{-n/2} \int e^{iS(t,x,y)} a(t,x,y) f(y) dy$$

$$(11) \quad \Rightarrow |U(t,x,y)| \leq \frac{C_n}{|t|^{n/2}}, \quad |t| < \delta.$$

Then it is well known that the dispersive estimate for $|t| < \delta$,

$$(12) \quad \|U(t)f\|_{L^\infty} \lesssim \frac{1}{|t|^{n/2}} \|f\|_{L^1},$$

leads to Strichartz estimates (13) and (14), and hence the local existence on $(-\delta, \delta)$, see [16, 28, 39, 50].

2.2. Dispersive and Strichartz estimates for $e^{-itH_{\Omega,V}}$.

Definition 2.3. We call $(q, r) = (q, r, n)$ an admissible pair if $q, r \in [2, \infty]$ satisfy $(q, r, n) \neq (2, \infty, 2)$ and

$$\frac{2}{q} + \frac{n}{r} = \frac{n}{2}.$$

By Duhamel formula, u is a weak solution of (1) is equivalent to

$$u = U(t)u_0 + i \int_0^t U(t-s)\lambda|u|^{p-1}uds.$$

From (12) and [47, Lemma 3.1] we have from above the following Strichartz estimates in weighted Sobolev spaces.

Lemma 2.4. *Let $I = [-T, T]$, $T < \delta$ small. Let (q, r) and (\tilde{q}, \tilde{r}) be any admissible pairs. Then*

$$\begin{aligned} & \|U(t)f\|_{L^q(I, L^r)} \leq C\|f\|_{L^2(\mathbb{R}^n)}, \\ & \left\| \int_0^t U(t-s)Fds \right\|_{L^q(I, L^r)} \leq C_{n,q,\tilde{q}}\|F\|_{L^{\tilde{q}'}(I, L^{\tilde{r}'})}, \\ (13) \quad & \|U(t)f\|_{L^q(I, \mathcal{H}^{1,r})} \leq C\|f\|_{\mathcal{H}^1}, \end{aligned}$$

$$(14) \quad \left\| \int_0^t U(t-s)Fds \right\|_{L^q(I, \mathcal{H}^{1,r})} \leq C_{n,q,\tilde{q}}\|F\|_{L^{\tilde{q}'}(I, \mathcal{H}^{1,\tilde{r}'})},$$

where

$$\begin{aligned} \|u\|_{L^q(I, L^r)} &= \left(\int_I \left(\int |u(t, x)|^r dx \right)^{q/r} dt \right)^{1/q}, \\ \|u\|_{L^q(I, \mathcal{H}^{1,r})} &= \left(\int_I \|u(t, \cdot)\|_{\mathcal{H}^{1,r}}^q dt \right)^{1/q}, \\ \|u\|_{\mathcal{H}^{1,r}} &:= \|\nabla u\|_{L^r} + \|xu\|_{L^r} + \|u\|_{L^r}. \end{aligned}$$

The lemma here applies to the case where V_e is subquadratic, e.g. $V_e(x) = -\sum_j \tilde{\gamma}_j^2 x_i^2$ for $\tilde{\gamma}_j \geq 0$. This is a generalized version for the Strichartz estimates given in [16, 50] where V_e is required being bounded from below. In the proof of the lemma, we directly study the action of $U(t-s)$ on the space $\mathcal{H}^{1,r}$ based on [47, Lemma 3.1], an oscillatory integral operator result of Yajima. This provides a treatment for more general A and V in the time-dependent case that covers those in Proposition 2.1. Such treatment are also different from the commutator method used in [2, 12].

2.5. **Minimal mass of blowup is $\|Q\|_2^2$.** Let $Q_{\Omega,\gamma,c} \in \Sigma$ be a ground state solution (g.s.s.) for the minimization problem:

$$(15) \quad E(Q_{\Omega,\gamma,c}) = \inf\{E_{\Omega,V}(u) : u \in \Sigma, \|u\|_2^2 = c^2 > 0\},$$

where $E_{\Omega,V}$ is given as in (9), also see (52) in Sections 7 for the definition of g.s.s. in the setting of general nonlinearity. Let $p = 1+4/n$. Theorem 1.1 shows that $\|Q\|_2^2$ is the threshold of minimal mass blowup if Q is the ground state of (2). This is consistent with the profile description for blowup solutions [30], where it is essentially shown that if $p = 3$, $n = 2$, then as $c^2 \nearrow \|Q\|_2^2$, the ground state $Q_{\Omega,\gamma,c}$ blowup occurs at the lowest point of the trap V , that is, if $c = \|Q\|_2$, there is no minimizer for (15). See also the characterization of blowup profiles in [6, 11, 24, 25].

3. VIRIAL IDENTITY FOR NLS WITH ROTATION

In this section we derive the virial identity associated with equation (1). Let A be given as in (6) and $L_\Omega = iA \cdot \nabla$. Proof of Theorem 1.1 is based on the following lemma for the variance

$$(16) \quad J(t) := \int |x|^2 |u|^2.$$

Lemma 3.1 (Virial identity). *Let u be solution of (1) with initial data $u_0 \in \mathcal{H}^1$. Then*

(17)

$$J'(t) = 2\Im \int x \bar{u} \cdot \nabla u$$

$$J''(t) = 2 \int |\nabla u|^2 - 2\gamma^2 \int |x|^2 |u|^2 - 2n\lambda \frac{p-1}{p+1} \int |u|^{p+1}$$

(18)

$$= 4E_{\Omega,V}(u) - 4\gamma^2 \int |x|^2 |u|^2 + \frac{2\lambda}{p+1} (4 - n(p-1)) \int |u|^{p+1} - 4\langle L_\Omega(u), u \rangle.$$

Note that $\langle L_\Omega u, u \rangle$ is real since L_Ω is selfadjoint. The virial inequality will be used to analyze wave collapse in finite time. Identity (18) can also be derived from the magnetic analog [18, 21, 8]. In Section 6, we will give a version of the virial identity in the case of inhomogeneous nonlinearity.

Proof of Lemma 3.1. We may assume $u \in C^1(I, C_0^2 \cap \mathcal{H}^1)$ and $I = [0, T_{max})$. For general data the identities (17) and (18) follow from a standard approximation argument. First we obtain the identity (17) from (16) and (1) by integration by parts, where note that

- (i) V and λ are real-valued;

- (ii) $x \cdot A = 0$;
- (iii) $L_z = i(x_2 \partial_{x_1} - x_1 \partial_{x_2})$ is self-adjoint and $L_z(|x|^2) = 0$.

Then, it follows by differentiating in t on (17) that

$$\begin{aligned} J''(t) &= 2\Im \left(\int x \bar{u} \cdot \nabla u_t + \int x \bar{u}_t \cdot \nabla u \right) = 2\Im \left(- \int \nabla \cdot (x \bar{u}) u_t + \int x \bar{u}_t \cdot \nabla u \right) \\ &= -2n\Im \left(\int \bar{u} u_t \right) - 4\Im \left(\int x \cdot \nabla \bar{u} u_t \right) \\ &:= -2nS - 4T. \end{aligned}$$

Noting that $\int \bar{u} L_z u$ is real, we have by a simple calculation

$$S = -\frac{1}{2} \int |\nabla u|^2 - \frac{\gamma^2}{2} \int |x|^2 |u|^2 + \lambda \int |u|^{p+1} + \Omega \int \bar{u} L_z u.$$

To compute T , one has

$$\begin{aligned} T &= \Im \left(\frac{i}{2} \int x \cdot \nabla \bar{u} \Delta u \right) + \Im \left(-i \frac{\gamma^2}{2} \int x \cdot \nabla \bar{u} (|x|^2 u) \right) \\ &\quad + \Im \left(i \lambda \int x \cdot \nabla \bar{u} |u|^{p-1} u \right) + \Im \left(i \Omega \int x \cdot \nabla \bar{u} L_z u \right) \\ &:= T_1 + T_2 + T_3 + T_4. \end{aligned}$$

For T_1 , integration by parts gives

$$\begin{aligned} T_1 &= \Im \left(-i \frac{1}{2} \sum_{j,k=1}^n \int (\delta_{j,k} \bar{u}_{x_j} + x_j \bar{u}_{x_j x_k}) u_{x_k} \right) \\ &= \Im \left(-i \frac{1}{2} \int |\nabla u|^2 \right) + \Im \left(-i \frac{1}{2} \sum_{j,k=1}^n \int x_j \bar{u}_{x_j x_k} u_{x_k} \right) := -\frac{1}{2} \int |\nabla u|^2 + T_{1,1}, \end{aligned}$$

where $\delta_{j,k}$ denote the Kronecker delta. To compute $T_{1,1}$, one has

$$\begin{aligned} T_{1,1} &= \Im \left(\frac{i}{2} \sum_{j,k=1}^n \int (x_j u_{x_k})_{x_j} \bar{u}_{x_k} \right) = \Im \left(\frac{i}{2} \sum_{j,k=1}^n \int (u_{x_k} + x_j u_{x_k x_j}) \bar{u}_{x_k} \right) \\ &= \frac{n}{2} \int |\nabla u|^2 + \Im \left(\frac{i}{2} \sum_{j,k=1}^n \int x_j u_{x_k x_j} \bar{u}_{x_k} \right) = \frac{n}{2} \int |\nabla u|^2 - T_{1,1}. \end{aligned}$$

This shows that $T_{1,1} = \frac{n}{4} \int |\nabla u|^2$, and so $T_1 = \frac{n-2}{4} \int |\nabla u|^2$. For T_2 and T_3 , applying divergence theorem shows that

$$\begin{aligned} T_2 &= \frac{n+2}{4} \gamma^2 \int |x|^2 |u|^2. \\ T_3 &= \frac{-\lambda n}{p+1} \int |u|^{p+1}. \end{aligned}$$

For T_4 , integration by parts gives

$$\begin{aligned} T_4 &= \Im \left(-\Omega \int \sum_{k=1}^n x_k \bar{u}_{x_k} (x_2 u_{x_1} - x_1 u_{x_2}) \right) \\ &= \Im \left(\Omega \int \left(\sum_{k=1}^n x_k \bar{u}_{x_k} x_2 \right)_{x_1} u \right) - \Im \left(\Omega \int \left(\sum_{k=1}^n x_k \bar{u}_{x_k} x_1 \right)_{x_2} u \right) \\ &= \Im \left(i\Omega \int u \overline{L_z u} \right) + \sum_{k=1}^n \Im \left(-\Omega \int (x_k x_2 u)_{x_k} \bar{u}_{x_1} \right) - \sum_{k=1}^n \Im \left(-\Omega \int (x_k x_1 u)_{x_k} \bar{u}_{x_2} \right) \\ &= \Omega \int u \overline{L_z u} + \Im \left(-n\Omega \int u (x_2 \bar{u}_{x_1} - x_1 \bar{u}_{x_2}) \right) + \Im \left(-\Omega \int u (x_2 \bar{u}_{x_1} - x_1 \bar{u}_{x_2}) \right) \\ &\quad + \Im \left(-\Omega \int x \cdot \nabla u (x_2 \bar{u}_{x_1} - x_1 \bar{u}_{x_2}) \right) \\ &= -n\Omega \int u \overline{L_z u} - T_4. \end{aligned}$$

This shows that $T_4 = -\frac{n}{2} \Omega \int u \overline{L_z u}$. Collecting all T_i terms yields

$$T = \frac{n-2}{4} \int |\nabla u|^2 + \frac{(n+2)\gamma^2}{4} \int |x|^2 |u|^2 - \frac{n\lambda}{p+1} \int |u|^{p+1} - \frac{n}{2} \Omega \int u \overline{L_z u}.$$

Finally, we obtain the virial identity (18)

$$\begin{aligned} J''(t) &= -2n \left(-\frac{1}{2} \int |\nabla u|^2 - \frac{\gamma^2}{2} \int |x|^2 |u|^2 + \lambda \int |u|^{p+1} + \Omega \int \bar{u} L_z u \right) \\ &\quad - 4 \left(\frac{n-2}{4} \int |\nabla u|^2 + \frac{(n+2)\gamma^2}{4} \int |x|^2 |u|^2 - \frac{n\lambda}{p+1} \int |u|^{p+1} - \frac{n}{2} \Omega \int u \overline{L_z u} \right) \\ &= 2 \int |\nabla u|^2 - 2\gamma^2 \int |x|^2 |u|^2 - 2n\lambda \left(\frac{p-1}{p+1} \right) \int |u|^{p+1}. \end{aligned}$$

□

3.2. The conservation of angular momentum. For A given in (6) we show (10) in Proposition 2.1 that is, the angular momentum $\ell_\Omega(u)$ is conserved in t .

Proof. Recall that $\ell_\Omega(u) = -\Omega \int \bar{u} L_z u$. It suffices to show $\frac{d}{dt} \int \bar{u} L_z u = 0$. Differentiating $\langle L_z u, u \rangle$ with respect to t and substituting u_t from equation (1), we have

$$\begin{aligned} \frac{d}{dt} (\langle L_z u, u \rangle) &= 2\Re \int \bar{u}_t L_z u \\ &= \Re \int \Delta \bar{u} (x_2 \partial_{x_1} u - x_1 \partial_{x_2} u) - 2\Re \int V \bar{u} (x_2 \partial_{x_1} - x_1 \partial_{x_2}) u \\ &\quad + 2\Re \int \lambda \bar{u} |u|^{p-1} (x_2 \partial_{x_1} - x_1 \partial_{x_2}) u \\ &=: 2\Re \left(\frac{1}{2} I_1 - I_2 + I_3 \right). \end{aligned}$$

Noting $[\Delta, L_z] = 0$, we obtain $I_1 = -\overline{I_1} \Rightarrow \Re I_1 = 0$. For I_2 , we have

$$\begin{aligned} I_2 &= \int V \bar{u} (x_2 \partial_{x_1} - x_1 \partial_{x_2}) u \\ &= - \int (\tilde{L}_z V) |u|^2 - \int V u \tilde{L}_z \bar{u} \\ &= 0 - \int V u \tilde{L}_z \bar{u} \Rightarrow \Re I_2 = 0, \end{aligned}$$

where we have defined $\tilde{L}_z = x_2 \partial_{x_1} - x_1 \partial_{x_2}$ and noted $\tilde{L}_z V(x) = 0$.

For I_3 , since \tilde{L}_z is skew-symmetric, we have

$$\begin{aligned} I_3 &= \lambda \int \bar{u} |u|^{p-1} \tilde{L}_z u \\ &= -\lambda \int u |u|^{p-1} (x_2 \partial_{x_1} - x_1 \partial_{x_2}) \bar{u} - \lambda \int |u|^2 (x_2 \partial_{x_1} - x_1 \partial_{x_2}) (|u|^{p-1}) \\ &= -\lambda \int u |u|^{p-1} \overline{\tilde{L}_z u} - \lambda(p-1) \Re \int (|u|^{p-1} u) (x_2 \partial_{x_1} - x_1 \partial_{x_2}) \bar{u}. \end{aligned}$$

Taking the real part gives

$$\Re(I_3) = -p\Re(I_3) \Rightarrow \Re(I_3) = 0.$$

Combining the results of I_1, I_2, I_3 above, we obtain

$$\frac{d}{dt} (\ell_\Omega(u)) = 0.$$

□

4. BLOWUP CRITERION FOR $\Omega \neq 0$

In this section, we prove some general criteria on finite time blowup for the solutions of RNLS (1) based on the virial identity and conservation of mass, energy and angular momentum.

Lemma 4.1. *Let $p \in [1 + \frac{4}{n}, 1 + \frac{4}{n-2})$. Suppose $0 \neq u_0 \in \mathcal{H}^1$ satisfies one of the following conditions:*

- (a) $E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0) \leq \frac{\gamma^2}{2}J(0)$, if $p = 1 + 4/n$.
- (b) $\frac{\gamma^2}{2}J(0) < E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0) \leq \frac{\gamma}{2}|J'(0)|$, if $p = 1 + 4/n$.
- (c) $E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0) < 0$, if $p > 1 + 4/n$.
- (d) $E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0) = 0$ and $J'(0) < 0$, if $p > 1 + 4/n$.

Then there exists $0 < T_* < \infty$ such that the corresponding solution u of equation (1) blows up on $[0, T_*)$ satisfying

$$(19) \quad \|\nabla u\|_2 \rightarrow \infty \quad \text{as } t \rightarrow T_*.$$

Remark 4.2. The condition in part (a) of the lemma is equivalent to

$$(20) \quad E_{0,0}(u_0) = \frac{1}{2} \int |\nabla u_0|^2 - \frac{\lambda n}{n+2} \int |u_0|^{2+\frac{4}{n}} \leq 0.$$

We will also need the uncertainty principle given in [45, 49].

Lemma 4.3.

$$(21) \quad \int_{\mathbb{R}^n} |u|^2 \leq \frac{2}{n} \int |\nabla u|^2 \int |x|^2 |u|^2,$$

where the equality is achieved with $u = e^{-|x|^2/2}$.

Now we prove Lemma 4.1.

Proof of Lemma 4.1. We divided the proof into three cases.

Case 1. Let $p = 1 + 4/n$. From (18) in Lemma 3.1, along with (10) and (9) we have if $p = 1 + 4/n$,

$$(22) \quad J''(t) + 4\gamma^2 J(t) = 4E_{\Omega,V}(u_0) - 4\ell_{\Omega}(u_0).$$

Solving this o.d.e. gives

$$(23) \quad \begin{aligned} J(t) &= (J(0) - \frac{1}{\gamma^2}(E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0))) \cos 2\gamma t + \frac{J'(0)}{2\gamma} \sin 2\gamma t + \frac{1}{\gamma^2}(E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0)) \\ &= C \sin(2\gamma t + \beta) + \frac{1}{\gamma^2}(E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0)), \end{aligned}$$

where $C > 0$ and $\beta \in (0, \pi)$ are constants given by

$$\begin{aligned} C^2 &= \left(J(0) - \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0)) \right)^2 + \frac{1}{4\gamma^2} (J'(0))^2, \\ \sin \beta &= \frac{J(0) - \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0))}{C} \geq 0, \quad (\text{by condition (a)}) \\ \cos \beta &= \frac{J'(0)}{2\gamma C}. \end{aligned}$$

Claim.

$$(24) \quad C \geq \frac{1}{\gamma^2} |E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0)|.$$

Indeed, the condition (a) always implies (24) if $E(u_0) - \ell_{\Omega}(u_0)$ is either positive or negative. Now, from the equation

$$(25) \quad 0 \leq J(t) = C \sin(2\gamma t + \beta) + \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0))$$

we see that there must exist $T_* \in [\frac{\pi}{4\gamma}, \frac{3\pi}{4\gamma}]$ such that

$$\lim_{t \rightarrow T_*} J(t) = 0.$$

Furthermore, since $\lim_{t \rightarrow T_*} J'(t) = 2\gamma C \cos(2\gamma T_* + \beta)$ is finite, we deduce from (21) that there exists some constant c_0 such that

$$(26) \quad \|\nabla u\|_2 \geq \frac{c_0 \|u_0\|_2}{\sqrt{T_* - t}}.$$

Case 2. Let $p = 1 + 4/n$. If $\frac{\gamma^2}{2} J(0) < E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0) \leq \frac{\gamma}{2} |J'(0)|$, then (24) still holds. It follows from the same line of proof in Case 1 that there exists some T_* such that (19) holds as $t \rightarrow T_*$.

Case 3. Let u_0 satisfy either condition (c) or (d). From (18) we have if $p \geq 1 + 4/n$,

$$J''(t) \leq 4(E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0)).$$

Then it follows from Taylor formula that

$$J(t) \leq J(0) + J'(0)t + 2(E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0))t^2.$$

We see either condition (c) or (d) suggests that there exists $T_* < \infty$ such that

$$J(t) \rightarrow 0 \quad \text{as } t \rightarrow T_*$$

which implies (19) by (21). This concludes the proof of Lemma 4.1. \square

Remark 4.4. Condition (a) in the lemma suggests that if the initial energy $E_{\Omega,V}(u_0) \leq \int V|u_0|^2 + \ell_{\Omega}(u_0) \iff E_{0,0}(u_0) \leq 0$, then finite time blowup occurs. However, when there is no potential, where $\gamma = 0$, the condition $E_{0,0}(u_0)$ alone may not imply blowup solutions in the mass-critical case. In the presence of potential, noting $T_* \leq \frac{3\pi}{4\gamma}$, we infer that V may make the blowup time come sooner. If in addition $J'(0) = 0$, then a closer look over the proof shows $T_* \in [\frac{\pi}{4\gamma}, \frac{\pi}{2\gamma})$.

5. PROOF OF THEOREM 1.1

The proof of part (a) of Theorem 1.1 relies on the sharp Gagliardo-Nirenberg inequality, where the optimal constant is evaluated in terms of the ground state $Q = Q_{\lambda,1}$ as given in (2). Part (b) of Theorem 1.1 follows as an application of Lemma 4.1. We begin with the following lemma on the Pohozaev identity.

Lemma 5.1. *Let $p \in (1, 1 + 4/(n - 2))$. Suppose $u \in H^1$ is a positive solution of the elliptic equation*

$$(27) \quad a\Delta u + bu^p = cu.$$

Then

$$(28) \quad \begin{aligned} 2a \int |\nabla u|^2 &= bn \frac{p-1}{p+1} \int |u|^{p+1} \\ 2c \int |u|^2 &= b \left(2 - \frac{n(p-1)}{p+1} \right) \int |u|^{p+1} \\ a \left(\frac{2(p+1)}{n(p-1)} - 1 \right) \int |\nabla u|^2 &= c \int |u|^2. \end{aligned}$$

Proof. Multiplying $x \cdot \nabla u$ and then integrating on both sides of the equation (27), we obtain

$$a \left(\frac{2}{n} - 1 \right) \int |\nabla u|^2 = c \int |u|^2 - \frac{2b}{p+1} \int |u|^{p+1}.$$

This and the obvious equation

$$a \int |\nabla u|^2 - b \int |u|^{p+1} = -c \int |u|^2$$

combine to prove the the three identities in the lemma. \square

For $a = \frac{1}{2}$, $\lambda > 0$, $c = 1$ the equation (2) has a unique positive radially symmetric solution $Q = Q_{\lambda}$ in $H^1 \cap C^2$ which is decreasing exponentially, see [29, 45]. If $p = 1 + 4/n$, we have $\|Q_{\lambda,1}\|_2^{4/n} = \lambda^{-1} \|Q_{1,1}\|_2^{4/n}$,

and by Lemma 5.1, $Q = Q_{\lambda,1}(x)$ satisfies

$$(29) \quad \int |\nabla Q_{\lambda,1}|^2 = \frac{2\lambda n}{n+2} \int |Q_{\lambda,1}|^{2+4/n}.$$

This is equivalent to

$$(30) \quad E_{0,0}(Q) = 0,$$

where

$$E_{0,0}(u) = \frac{1}{2} \int |\nabla u|^2 - \frac{2\lambda}{p+1} \int |u|^{p+1}.$$

Next we turn to the sharp Gagliardo-Nirenberg inequality, which states that there exists an optimal constant $c_{GN} = c(p, n)$ such that for all u in H^1

$$\int |u|^{p+1} \leq c_{GN} \|u\|_2^{2-\frac{(p-1)(n-2)}{2}} \|\nabla u\|_2^{n(p-1)/2}.$$

If $p = 1 + 2\sigma$, the constant c_{GN}^{-1} is given by the minimal of the J -functional:

$$(31) \quad c_{GN}^{-1} = \min_{0 \neq u \in H^1} J_{\sigma,n}(u),$$

where

$$(32) \quad J_{\sigma,n}(u) = \frac{\|u\|_2^{2+2\sigma-n\sigma} \|\nabla u\|_2^{n\sigma}}{\|u\|_{p+1}^{p+1}}.$$

Note that $J_{\sigma,n}$ has the scaling invariance $J_{\sigma,n}(\beta u(\alpha x)) = J_{\sigma,n}(u)$ for all $\alpha, \beta > 0$. Thus $Q = Q_{\lambda,1}$ is a minimizer of $J_{\sigma,n}$. From Lemma 5.1 it follows that the sharp constant in (31) is given by

$$(33) \quad \begin{aligned} c_{GN}^{-1} &= J_{\sigma,n}(Q) \\ &= \frac{\lambda \|Q_{\lambda,1}\|_2^{2\sigma}}{2^{1-\sigma n/2}} \left(2 - \frac{\sigma n}{\sigma + 1}\right) \left(\frac{\sigma n}{2\sigma + 2 - \sigma n}\right)^{\sigma n/2}, \end{aligned}$$

where $u = Q_{\lambda,\omega} > 0$ solves the elliptic problem (27) if taking $a = \frac{1}{2}$, $b = \lambda$, $c = \omega$. In particular, if $p = 1 + 4/n$, then we have

$$(34) \quad c_{GN}^{-1} = \frac{2\lambda n}{n+2} \|Q_{\lambda,1}\|_2^{4/n} = \frac{2n}{n+2} \|Q_{1,1}\|_2^{4/n},$$

where note that $u = Q_{\lambda,1}$ is the solution of (2) and equation (27) enjoys the following scaling invariance:

- (i) ω -scaling: $u_{\lambda,\omega}(x) = \omega^{1/(p-1)} u_{\lambda,1}(\omega^{1/2}x)$
- (ii) λ -scaling: $u_{\lambda,\omega}(x) = \lambda^{-1/p} u_{1,\lambda^{-1/p}\omega}(\lambda^{1/2}x)$.

Summarizing the above gives the following lemma.

Lemma 5.2. *Let $p \in (1, 1 + \frac{4}{n-2})$, $\sigma = (p-1)/2$ and $\theta = n(\frac{1}{2} - \frac{1}{p+1})$. Then*

(a)

$$(35) \quad \|u\|_{p+1}^{p+1} \leq c_{GN} \|u\|_2^{(1-\theta)(p+1)} \|\nabla u\|_2^{\theta(p+1)},$$

where c_{GN} is defined as in (33):

$$(36) \quad \begin{aligned} c_{GN}^{-1} &= \frac{\lambda \|Q_{\lambda,1}\|_2^{2\sigma}}{2^{1-\sigma n/2}} \left(2 - \frac{\sigma n}{\sigma+1}\right) \left(\frac{\sigma n}{2\sigma+2-\sigma n}\right)^{\sigma n/2} \\ &= \frac{\|Q_{1,1}\|_2^{2\sigma}}{2^{1-\sigma n/2}} \left(2 - \frac{\sigma n}{\sigma+1}\right) \left(\frac{\sigma n}{2\sigma+2-\sigma n}\right)^{\sigma n/2}. \end{aligned}$$

(b) *In particular, if $p = 1 + 4/n$,*

$$(37) \quad \|u\|_{2+4/n}^{2+4/n} \leq c_{GN} \|u\|_2^{4/n} \|\nabla u\|_2^2,$$

where c_{GN} is given by (34).

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. (1) Let $u \in C([0, T_{max}), \mathcal{H}^1)$ be the solution of equation (1). Let $Q = Q_{\lambda,1}$ be the g.s.s of (2). From (9) and Lemma 5.2 we have

$$\frac{1}{2} \int |\nabla u|^2 + \frac{\gamma^2}{2} \int |x|^2 |u|^2 - E_{\Omega,V}(u_0) + \ell_{\Omega}(u_0) \leq \frac{1}{2} \|Q\|_2^{-4/n} \|\nabla u\|_2^2 \|u\|_2^{4/n}.$$

Then with $\gamma > 0, \lambda > 0$ in (1), we obtain

$$\begin{aligned} &\frac{1}{2} \left(1 - \left(\frac{\|u_0\|_2}{\|Q\|_2}\right)^{4/n}\right) \int |\nabla u|^2 + \frac{\gamma^2}{2} \int |x|^2 |u|^2 \\ &\leq E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0). \end{aligned}$$

Since $\|u_0\|_2 < \|Q\|_2$, it follows that $\|\nabla u\|_2$ and $\|xu\|_2$ are bounded for any $t \in [0, T_{max})$. Therefore $T_{max} = \infty$, in view of Proposition 2.1 (c), which proves part (a).

(2) To prove the second part we consider the following initial data

$$u_0(x) = ce^{i\theta} \alpha^{n/2} Q(\alpha x)$$

with $c \geq 1, \alpha > 0, \theta \in \mathbb{R}$. Then $\|u_0\|_2 = c \|Q\|_2$. It is easy to calculate with $p = 1 + 4/n$

$$E_{\Omega,V}(u_0) - \ell_{\Omega}(u_0) \begin{cases} = \frac{\gamma^2}{2} J(0) & \text{if } c = 1, \\ < \frac{\gamma^2}{2} J(0) & \text{if } c > 1. \end{cases}$$

Indeed, we have from (30)

$$\begin{aligned} E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0) &= \frac{1}{2} \int |\nabla u_0|^2 + \frac{\gamma^2}{2} \int |x|^2 |u_0|^2 - \frac{2\lambda}{p+1} \int |u_0|^{p+1} \\ &= \frac{c^2 \alpha^2}{2} \int |\nabla Q(x)|^2 + \frac{c^2 \gamma^2}{2\alpha^2} \int |x|^2 |Q(x)|^2 - \frac{2\lambda}{p+1} c^{p+1} \alpha^{\frac{n}{2}(p-1)} \int |Q(x)|^{p+1} \\ &\leq c^2 \alpha^2 E_{0,0}(Q) + \frac{c^2 \gamma^2}{2\alpha^2} \int |x|^2 |Q(x)|^2 = \frac{\gamma^2}{2} J(0), \end{aligned}$$

where we note $J(0) = \int |x|^2 |u_0|^2 = \frac{c^2}{\alpha^2} \int |x|^2 |Q|^2$. Observe in addition from (17) that $J'(0) = 2\Im \int x \bar{u}_0 \nabla u_0 = 2\frac{c^2}{\alpha} \Im \int x \bar{Q} \nabla Q dx = \Im (n \int |Q|^2 dx) = 0$. Therefore, according to Lemma 4.1 (a) and Remark 4.4, we conclude that the solution of (1) corresponding to the given u_0 blows up on $[0, T_*)$ for $T_* \in [\frac{\pi}{4\gamma}, \frac{\pi}{2\gamma})$. \square

The following proposition provides a blowup result as an application of Lemma 4.1 in the mass supercritical regime.

Proposition 5.3. *Let $p \in (1 + 4/n, 1 + 4/(n - 2))$. Let Q be the g.s.s in (2). Then there exists u_0 such that the corresponding solution u of (1) blows up in finite time.*

Proof. Take

$$(38) \quad u_0(x) = c e^{i\nu|x|^2} Q(x), \quad \nu \in \mathbb{R}.$$

A straightforward calculation shows for $p > 1 + 4/n$ and c sufficiently large, by (28)

$$(39) \quad E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0) < 0,$$

which suggests finite time blowup solution according to Lemma 4.1 (c). \square

Remark 5.4. If $p = 1 + 4/n$, $\|u_0\|_2 = \|Q\|_2$, then the so-called \mathcal{R} -transform, yields all blowup solutions modular symmetries (scaling, translation and phase invariance) with initial data given by either Q or for $\nu > 0$

$$(40) \quad u_0(x) = \nu^{n/2} e^{i\nu} e^{-i\frac{\nu}{2}|x|^2} Q(\nu x),$$

see [6, Subsection 4.1]. Note that Q is the profile for blowup, while the standing waves $u = e^{it} Q_{\Omega, \gamma}$ apparently are global solutions.

Remark 5.5. In the case $\Omega = 0, \gamma \neq 0$, Carles [11] also tells that Q is the critical minimal mass blowup. However, Carles's result is obtained by lens transform and Merle's characterization for the profile, while our proof relies on a virial identity for (1), which seems more direct and

simpler. In particular, the blowup condition in Lemma 4.1 is sharper than those in [11, Corollary 4.3 (ii)] for that case and [8] for mNLS.

5.6. Focusing NLS with angular momentum: Profile for blowup solution. In view of [6, 11, 36], we are able to apply similar argument and techniques to describe the behavior of minimal mass blowup solutions of (1) at the mass level $\|Q\|_2$ in the mass-critical case by showing that all blowup solutions result from concentration of energy as $t \rightarrow T$. Let $u_0 \in \mathcal{H}^1$ and $\|u_0\|_2 = \|Q\|_2$. If $u \in C([0, T], \mathcal{H}^1)$ is the blowup solution of (1) with $A = \Omega\langle -x_2, x_1, 0 \dots, 0 \rangle$, $V = \frac{\gamma^2}{2}|x|^2$, then there exist functions $x(t) \in \mathbb{R}^n$ and $\theta(t) \in \mathbb{R}$ so that the following holds in \mathcal{H}^1 :

$$u(t, x) = \frac{1}{\lambda(t)^{n/2}} Q\left(\frac{x - x(t)}{\lambda(t)}\right) e^{i\theta(t)} + o(1) \quad \text{as } t \rightarrow T.$$

Here $Q \in H^1(\mathbb{R}^n)$ is given by (2) and $\lambda(t) = \|\nabla Q\|_2 / \|\nabla u\|_2$. However, such behavior of collapse is unstable at the ground state level, see [6, 38]. This is in consistence with the stability result at $p = 1 + 4/n$ where Q is the threshold, as is shown in Theorem 1.1 and Theorem 1.2. We shall prove Theorem 1.2 in Sections 7 and 8 that when the initial data is below the ground state level, namely, $c < \|Q_{\lambda,1}\|_2$, then (1) is orbital stable. The analogous result is valid for NLS (4) with zero potential. Moreover, Theorem 1.1 shows the solution is strongly unstable when $M(u_0) = M(Q)$, where $\|\nabla u\|_2 \approx \frac{\|\nabla Q\|_2}{T-t}$ as $t \rightarrow T$. The question concerning blowup rate above the ground state level has been briefly discussed in (3) in Section 1.

6. VIRIAL IDENTITY FOR INHOMOGENEOUS NLS

In this section, we mainly prove a virial identity for (41), which will allow us to show a similar blowup result in the case of an inhomogeneous nonlinearity. Consider the inhomogeneous RNLS in \mathbb{R}^{1+n}

$$(41) \quad iu_t = -\frac{1}{2}\Delta u + \frac{\gamma^2}{2}|x|^2 u - \lambda(x)|u|^{p-1}u - \Omega L_z u, \quad u_0 \in \Sigma = \mathcal{H}^1$$

where $p \in [1, 1 + 4/(n-2))$ and $\lambda = \lambda(x)$ satisfies the conditions in Hypothesis 1.

Hypothesis 1. Assume $\lambda \in C^1(\mathbb{R}^n)$ satisfies the following:

- (i) λ is radially symmetric
- (ii) $\lambda_{min} \leq \lambda(x) \leq \lambda_{max}$
- (iii) $x \cdot \nabla \lambda(x) \leq 0$,

where we write $\lambda_{min} = \inf_x \lambda(x) > 0$ and $\lambda_{max} = \sup_x \lambda(x) > 0$.

Examples of λ 's satisfying the hypothesis include, e.g. $\lambda = \lambda_0 + \langle x \rangle^{-m}$, where $\lambda_0, m > 0$.

6.1. Conservation laws for equation (41). Let $p \in (1, 1+4/(n-2))$ and $\lambda = \lambda(x)$ as above. Let $V = \frac{\gamma^2}{2}|x|^2$ and $L_\Omega = -\Omega L_z$ be the same notation as for (1). Suppose $u_0 \in \Sigma$ and u is the corresponding solution of (41). Following the same line of proof as in the case λ being a constant, one can show the local wellposedness and blowup alternative theorem like Proposition 2.1. Moreover, the mass, energy and angular momentum are conserved in the lifespan $I = (-T_{min}, T_{max})$:

$$\begin{aligned}
 M(u) &= \int |u|^2 = M(u_0) \\
 (42) \quad E_{\Omega, V}(u) &= \frac{1}{2} \int |\nabla u|^2 + \int V|u|^2 - \frac{2}{p+1} \int \lambda(x)|u|^{p+1} + \ell_\Omega(u) \\
 &= E_{\Omega, V}(u_0) \\
 \ell_\Omega(u) &= -\Omega \int \bar{u} L_z u = \ell_\Omega(u_0).
 \end{aligned}$$

6.2. Localized virial identity for (41). Let $\rho(x) = \rho(|x|)$ be radial. Define $J(t) := J_\rho(t) = \int \rho|u|^2$. By similar argument as in the proof of Lemma 3.1, we are able to show the localized virial identity for equation (41), whose detailed proof will be omitted. The magnetic versions were obtained in [18, 21] and earlier [22] for homogeneous equations.

Lemma 6.3. *Let $p \in [1, 1 + 4/(n - 2))$ and $\rho \in C_0^\infty(\mathbb{R}^n)$ be radially symmetric. If u is a solution of (41) in $C^1(I, \Sigma)$, then we have*

$$J'_\rho(t) = \Im \int \nabla \rho \cdot \nabla u \bar{u}$$

and

$$\begin{aligned}
 (43) \quad J''_\rho(t) &= -\frac{1}{4} \int \Delta^2 \rho |u|^2 - \frac{p-1}{p+1} \int \lambda \Delta \rho |u|^{p+1} + \int \nabla \bar{u} \cdot \text{Hessian}(\rho) \nabla u \\
 &\quad - \gamma^2 \int x \cdot \nabla \rho |u|^2 + \frac{2}{p+1} \int \nabla \lambda \cdot \nabla \rho |u|^{p+1}.
 \end{aligned}$$

Ansatz $\rho = |x|^2$ gives us the virial identity in the following lemma.

Lemma 6.4. *Let $p \in [1, 1 + 4/(n - 2))$ and $u_0 \in \Sigma$. Let u be the corresponding solution of (41) in $C^1(I, \Sigma)$. Then it holds*

$$\begin{aligned} J''(t) &= 2 \int |\nabla u|^2 - 2\gamma^2 \int |x|^2 |u|^2 \\ &\quad - 2n \frac{p-1}{p+1} \int \lambda |u|^{p+1} + \frac{4}{p+1} \int x \cdot \nabla \lambda |u|^{p+1} \\ &= 4E_{\Omega, V}(u_0) + 4\Omega \int (L_z u_0) \bar{u}_0 - 4\gamma^2 \int |x|^2 |u|^2 \\ &\quad + \frac{2}{p+1} (4 - n(p-1)) \int \lambda(x) |u|^{p+1} + \frac{4}{p+1} \int x \cdot \nabla \lambda |u|^{p+1}. \end{aligned}$$

An application of Lemma 6.4 yields all the counterpart parts (a)-(d) of Lemma 4.1 for equation (41). For our purpose we state the blowup criterion in the mass-critical case only.

Lemma 6.5. *Let $p = 1 + \frac{4}{n}$. Suppose $0 \neq u_0 \in \Sigma$ satisfies $E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0) \leq \frac{\gamma^2}{2} J(0)$, that is $E_{0,0}(u_0) \leq 0$. Then, there exists $0 < T_* < \infty$ such that the corresponding solution u of equation (41) satisfies*

$$(44) \quad \|\nabla u\|_2 \rightarrow \infty \quad \text{as } t \rightarrow T_*.$$

The proof of the lemma is similar to that for part (a) of Lemma 4.1. We only mention a technical point concerning dealing with the inhomogeneous term in the virial identity. Following the proof of the lemma in the homogeneous case, instead of (22) we arrive at

$$(45) \quad J''(t) + 4\gamma^2 J(t) = 4E_{\Omega, V}(u_0) - 4\ell_{\Omega}(u_0) + \frac{4}{p+1} \int x \cdot \nabla \lambda |u|^{p+1}$$

by applying Lemma 6.4. An integral representation for $J(t)$ is given by

$$\begin{aligned} J(t) &= (J(0) - \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0))) \cos 2\gamma t + \frac{J'(0)}{2\gamma} \sin 2\gamma t \\ &\quad + \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0)) + \int_0^t \frac{\sin(2\gamma(t-s))}{2\gamma} f(s) ds \end{aligned} \tag{46}$$

$$= C \sin(2\gamma t + \beta) + \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0)) + \int_0^t \frac{\sin(2\gamma(t-s))}{2\gamma} f(s),$$

where the constant C is given as in (23) and $f(t) := \frac{4}{p+1} \int x \cdot \nabla \lambda |u(t, x)|^{p+1}$. Since $f(t) \leq 0$ by condition (iii) in Hypothesis 1, it follows that

$$(47) \quad J(t) \leq C \sin(2\gamma t + \beta) + \frac{1}{\gamma^2} (E_{\Omega, V}(u_0) - \ell_{\Omega}(u_0)).$$

Then, this inequality implies Lemma 6.5 for RNLS (41).

Let $Q_\lambda := Q_{\lambda,1} \in H^1 \cap \Sigma$ be the unique positive radial ground state solution of equation (2). Then $Q_{\lambda,1}(x) = \lambda^{-n/4} Q_{1,1}(x)$. Proceeding as in the proof of Theorem 1.1, applying (37) and Lemma 6.5 we can prove the following theorem for RNLS (41).

Theorem 6.6. *Let $p = 1 + \frac{4}{n}$ and (Ω, γ) be any pair in $\mathbb{R} \times \mathbb{R}_+$. Suppose the initial data $u_0 \in \Sigma$. The following holds true for (41).*

- (a) *If $\|u_0\|_{L^2} < \|Q_{\lambda_{max}}\|_2$, then the solution $u \in C(\mathbb{R}, \Sigma)$ exists globally in time.*
- (b) *There exists u_0 in Σ satisfying $\|u_0\|_2 \geq \|Q_{\lambda_{min}}\|_2$ such that the solution $u \in C(I, \Sigma)$ blows up near both ends of $I = (-T_{min}, T_{max})$ in the sense that $\|\nabla u\|_2 \rightarrow \infty$ as $t \rightarrow T_{max}$ and as $t \rightarrow -T_{min}$.*

Remark 6.7. From the proof of part (b) of this theorem, in view of Lemma 6.5, we can construct a blowup solution by taking $u_0(x) = c\alpha^{n/2} Q_{\lambda_{min}}(\alpha x)$ with $c \geq 1$ and $\alpha > 0$. However, we do not know the exact minimal mass for blowup due to the gap $\|Q_{\lambda_{max}}\|_2 < \|Q_{\lambda_{min}}\|_2$. We conjecture that the exact minimal mass level is given by $\|\tilde{Q}\|_2$, where $\tilde{Q} \in H^1$ obeys

$$-\frac{1}{2}\Delta\tilde{Q} - \lambda(x)|\tilde{Q}|^{\frac{4}{n}}\tilde{Q} = -\tilde{Q},$$

see [37] for related discussions. The analogue of Theorem 6.6 was studied in [33] when $\Omega = 0$, $V = |x|^2$. For further investigation we refer to [41] for the treatment on the minimal mass problem in the case of the inhomogeneous NLS.

7. GROUND STATE SOLUTIONS

In this section, we will prove the existence for the set of ground states for (1) in the case $|\Omega| < \gamma$. In fact, we will elaborate the result for more general nonlinearity as given in the following RNLS on \mathbb{R}^{1+n}

$$(48) \quad iu_t = -\frac{1}{2}\Delta u + \frac{\gamma^2}{2}|x|^2 u - \kappa N(x, u) - \Omega L_z u, \quad u(0, x) = u_0 \in \Sigma.$$

The associated energy functional $E_{\Omega, \gamma}$ is given by

$$(49) \quad E_{\Omega, \gamma}(u) = \int \left(\frac{1}{2}|\nabla u|^2 + \frac{\gamma^2}{2}|x|^2|u|^2 - \kappa G(|u|^2) - \Omega \bar{u} L_z u \right) dx,$$

where $\kappa = \pm 1$ and $G(x, s^2) = 2 \int_0^s N(x, \tau) d\tau$ for $s \geq 0$. Throughout this section and next, we assume conditions $(\Omega, \gamma)_0$ and $(G)_0$ given in the following hypothesis.

Hypothesis 2. *Let Ω , γ and G satisfy the following conditions.*

$(\Omega, \gamma)_0: |\Omega| < \gamma$.

(G_0) : Let $G : \mathbb{R}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be continuous and differentiable such that for some constant $C > 0$ and $p > 1$,

$$(50) \quad 0 \leq G(x, v) \leq C(v + v^{\frac{p+1}{2}}).$$

Note that the nonlinearity $N(u) = G'(|u|^2)u$ includes $\mu(x)|u|^{p-1}u$, and the combined inhomogeneous terms $\mu_1(x)|u|^{p_1-1}u + \mu_2(x)|u|^{p_2-1}u$, μ, μ_1, μ_2 being positive and bounded functions. In what follows we will use the abbreviation $G(s) = G(x, s)$ unless otherwise specified.

Recall $\Sigma = \{u \in H^1 : \int |x|^2 |u|^2 < \infty\}$ is a Hilbert space endowed with the inner product

$$\langle u, v \rangle_\Sigma = \langle H_{\Omega, \gamma} u, v \rangle_{L^2},$$

where the norm is given by $\|u\|_\Sigma^2 = \langle H_{\Omega, \gamma} u, u \rangle \approx \|\nabla u\|_2^2 + \|xu\|_2^2 + \|u\|_2^2$ and $H_{\Omega, \gamma}$ is defined as in (7). Then we have Σ is compactly embedded in L^r for all $r \in [2, \infty)$ in view of [49, Lemma 5.1].

Definition 7.1. Let

$$(51) \quad I_c = \inf\{E_{\Omega, \gamma}(u) : u \in S_c\}, \quad S_c = \{u \in \Sigma : \int |u|^2 = c^2\}.$$

Then $u \in S_c$ is called a ground state solution (g.s.s.) provided u is a solution to the following minimization problem

$$(52) \quad E_{\Omega, \gamma}(u) = I_c.$$

Note that any solution of (52) solves the Euler-Lagrange equation

$$(53) \quad -\omega u = -\frac{1}{2}\Delta u + \frac{\gamma^2}{2}|x|^2 u - \kappa N(x, u) - \Omega L_z u,$$

where $-\omega \in \mathbb{R}$ is a Lagrange multiplier. Once we construct a g.s.s. $u = Q_{\Omega, \gamma} := Q_{\Omega, \gamma, c}$, then $e^{i\omega t} Q_{\Omega, \gamma}$ is a solitary wave solution to (48). Write

$$(54) \quad E_{\Omega, \gamma}(u) = E_{\Omega, \gamma}^1(u) - \kappa \int G(|u|^2).$$

Then $E_{\Omega, \gamma}^1$ denotes the linear energy

$$E_{\Omega, \gamma}^1(u) = \frac{1}{2} \|\nabla_A u\|_2^2 + \int V_e |u|^2,$$

where A is given in (6) and $V_e = \frac{1}{2}(\gamma^2 - \Omega^2)(x_1^2 + x_2^2) + \frac{\gamma^2}{2}x_3^2 + \dots + \frac{\gamma^2}{2}x_n^2$ as in Section 2. It follows by the assumption $(\Omega, \gamma)_0$ and [50, Proposition 4.4] that for all $u \in \Sigma$

$$(55) \quad E_{\Omega, \gamma}^1(u) = \langle H_{\Omega, \gamma} u, u \rangle = \|u\|_\Sigma^2.$$

Now we state the first result in this section.

Theorem 7.2. *Assume Hypothesis 2. Let $c > 0$.*

- (a) *If $1 \leq p < 1 + \frac{4}{n}$ and $\kappa = 1$, then the problem (52) has a minimizer, i.e., for any $c > 0$, there exists $u := Q_{\Omega, \gamma} \in S_c$ such that $E_{\Omega, \gamma}(u) = I_c$.*
- (b) *If $1 \leq p < 1 + \frac{4}{n-2}$ and $\kappa = -1$, then (52) always has a minimizer solution for any $c > 0$.*

Proof. (a) Let $p \in (1, 1 + \frac{4}{n})$ and G satisfy (G_0) . Then $0 < \sigma n < 2$. First we show that for all u in S_c , $E_{\Omega, \gamma}^1(u) = \|u\|_{\Sigma}^2$ are bounded from below, which implies

$$(56) \quad I_c \neq -\infty.$$

Indeed, we obtain from (50) and (35) and the Young's inequality

$$(57) \quad \begin{aligned} \int G(|u|^2) &\leq C \int |u|^2 + C \int |u|^{p+1} \\ &\leq C \|u\|_2^2 + C \frac{\|u\|_2^{(2\sigma+2-\sigma n)q}}{\varepsilon^q} + C \varepsilon^{q'} \|\nabla u\|_2^2, \quad \forall \varepsilon > 0, \end{aligned}$$

where we set $a = \varepsilon^{-1} \|u\|_2^{2\sigma+2-\sigma n}$, $b = \varepsilon \|\nabla u\|_2^{\sigma n}$, $q' = \frac{2}{\sigma n}$, $q = \frac{2}{2-\sigma n}$ in the inequality $ab \leq \frac{a^q}{q} + \frac{b^{q'}}{q'}$. Then, by taking ε sufficiently small, it is easy to see from (54) and (57) that there exists some $C_0 > 0$ such that for all u in S_c

$$(58) \quad E_{\Omega, \gamma}(u) \geq -C_0.$$

Secondly, we show that any given minimizing sequence $\{u_n\} \subset S_c$ of (52) is relatively compact in Σ . To do this we prove the following properties:

- (i) $\|u_n\|_{\Sigma}$ are uniformly bounded.
- (ii) There exists a subsequence $\{u_{n_j}\}$ weakly converging to some $w \in \Sigma$ such that $\lim_{j \rightarrow \infty} \int G(|u_{n_j}|^2) = \int G(|w|^2)$.
- (iii) The limit w is a minimizer solution to (52).
- (iv) The sequence $\{u_{n_j}\}$ converges to w in the Σ -norm.

Since $\{u_n\}$ is a minimizing sequence in S_c , $E_{\Omega, \gamma}(u_n)$ are bounded. It follows from (54) and (57) that

$$\begin{aligned} E_{\Omega, \gamma}^1(u_n) &= E_{\Omega, \gamma}(u_n) + \int G(|u_n|^2) \\ &\leq C + \frac{C}{\varepsilon^q} + C \varepsilon^{q'} \|\nabla u_n\|_2^2. \end{aligned}$$

By taking ε sufficiently small, we obtain, in view of (55),

$$\|u_n\|_{\Sigma}^2 = E_{\Omega, \gamma}^1(u_n) \leq C.$$

This proves (i).

The property (i) suggests that $\{u_n\}$ is weakly precompact in Σ , namely, there exists $w \in \Sigma$ such that, up to a subsequence, $\{u_n\} \rightharpoonup w$ weakly in Σ . Since Σ is compactly embedded in L^r for any $r \in [2, \frac{2n}{n-2})$, we see that there exists a subsequence, which is still denoted by $\{u_n\}$, such that

$$(59) \quad u_n \rightarrow w \text{ in } L^r, \quad \forall 2 \leq r < \infty.$$

Claim.

$$(60) \quad \int G(|u_n|^2) dx \rightarrow \int G(|w|^2).$$

Indeed, since $u_n \rightarrow w$ in L^2 and L^{p+1} , by the continuity of G we have, up to a subsequence, for almost all x

$$(61) \quad \lim_n G(|u_n(x)|^2) = G(|w(x)|^2).$$

Moreover, by the inverse dominated convergence theorem, we can find a subsequence $\{u_{n_j}\}$ of $\{u_n\}$ and two functions h_1 and h_2 such that:

- (1) $h_1 \in L^2$ and $h_2 \in L^{p+1}$
- (2) $|u_{n_j}(x)| \leq h_1(x)$ and $|u_{n_j}(x)| \leq h_2(x)$ a.e.

Thus $G(|u_{n_j}(x)|^2) \leq C(h_1(x)^2 + h_2(x)^{p+1})$. Since $h_1^2 + h_2^{p+1} \in L^1$, applying the dominated convergence theorem, we obtain

$$(62) \quad \lim_{j \rightarrow \infty} \int G(|u_{n_j}(x)|^2) = \int G(|w(x)|^2).$$

This actually implies that the whole sequence in the Claim verifies (60), because otherwise one can easily find a contradiction by a subsequence argument. Thus we have proven property (ii).

To show (iii), we note that, on the other hand, by the lower semi-continuity of $\|\cdot\|_\Sigma$ for weakly convergence sequence,

$$(63) \quad \|w\|_\Sigma^2 \leq \liminf \|u_n\|_\Sigma^2.$$

Now combining (59), (60) and (63) we obtain

$$(64) \quad \begin{cases} \|w\|_2^2 = c^2, \\ E_{\Omega, \gamma}(w) \leq \liminf E_{\Omega, \gamma}(u_n) = I_c. \end{cases}$$

Consequently w is a minimizer of (51). In turn, since w is a weak limit of u_n in Σ , plus $\lim_n \|u_n\|_\Sigma = \|w\|_\Sigma$, we conclude that $\{u_n\}$ converges to w in Σ . This proves (iv) and the existence of a ground state in the focusing case.

(b) Let $p \in (1, 1 + \frac{4}{n-2})$ and $\kappa = -1$ (defocusing). We have

$$(65) \quad E_{\Omega, \gamma}(u) = \|u\|_{\Sigma}^2 + \int G(|u|^2).$$

Then the problem (51) always has $0 < I_c < \infty$. Let $\{u_n\} \subset S_c$ be a minimizing sequence of (52): $\|u_n\|_2^2 = c^2$ and $\lim_n E_{\Omega, \gamma}(u_n) = I_c$.

It follows from (65) that $\|u_n\|_{\Sigma}^2 \leq E_{\Omega, \gamma}(u_n)$. Therefore $\{u_n\}$ is bounded in Σ . The remaining of the proof proceeds the same as in the focusing case. \square

From the proof of Theorem 7.2, we have the following corollary.

Corollary 7.3. *Under the assumptions in either (a) or (b) of Theorem 7.2, any minimizing sequence of (52) is relatively compact in Σ .*

Remark 7.4. If $N = \mu(x)|u|^{p-1}u$ and $u = Q_{\Omega, \gamma}$ is a g.s.s. of (52), then

$$-\omega c^2 = I_c - \frac{p-1}{p+1} \int \mu(x)|Q_{\Omega, \gamma}|^{p+1}.$$

This identity can provide information on the signs of the ω and I_c .

Remark 7.5. The convergence for the integrals in (60) essentially relies on the so called inverse dominated convergence theorem. We can also prove (60) by the fact that any given $f_n \rightarrow f$ a.e. satisfying $\{f_n\}$ being uniformly absolute continuous in the sense of integration, then $\lim_n \int f_n \rightarrow \int f$ holds (without recourse to the Lebesgue dominated convergence theorem).

7.6. Mass-critical case. When $p = 1 + 4/n$, the minimization problem (52) still makes sense for small values of $c > 0$. We show in Theorem 7.8 that there exists a threshold mass level concerning the existence of g.s.s. A priori, we need the following lemma by the Gagliardo-Nirenberg inequality (37) and the diamagnetic inequality

$$\|\nabla|u|\|_2 \leq \|(\nabla - iA)u\|_2,$$

which can be found in e.g. [17] or [8].

Lemma 7.7. *There is a sharp constant c_{GN} such that for all $u \in \Sigma$*

$$(66) \quad \|u\|_{2+4/n}^{2+4/n} \leq c_{GN} \|\nabla_A u\|_2^2 \|u\|_2^{4/n},$$

where $c_{GN} = \frac{n+2}{2n\|Q_{1,1}\|_2^{4/n}} = \frac{n+2}{2\lambda n\|Q_{\lambda,1}\|_2^{4/n}}$ for any $\lambda > 0$.

Theorem 7.8. *Let $p = 1 + 4/n$ and $\kappa = 1$. Let Ω, γ and G verify Hypothesis 2. Let $Q_{\lambda,1}$ be the positive radial solution of (2), where $\lambda = \frac{n+2}{n}C$ and C is the constant in (50). We have:*

- (a) If $c < \|Q_{\lambda,1}\|_2$, then there exists a ground state solution of the problem (52). Moreover, any minimizing sequence for (52) is relatively compact in Σ .
- (b) In particular, let $G(v) = \frac{\lambda n}{n+2} v^{1+2/n}$ corresponding to (1). If $c < \|Q_{\lambda,1}\|_2$, then there exists a g.s.s. $u = Q_{\Omega,\gamma}$ of (52) satisfying for some $\omega = \omega_c \in \mathbb{R}$

$$-\frac{1}{2}\Delta u + \frac{\gamma^2}{2}|x|^2 u - \lambda|u|^{\frac{4}{n}}u - \Omega L_z u = -\omega u, \quad \|u\|_2 = c.$$

Proof. From (50) and Lemma 7.7 we have for all u in S_c

$$\begin{aligned} E_{\Omega,\gamma}(u) &= \frac{1}{2} \int |\nabla_A u|^2 + \int V_e |u|^2 - \int G(|u|^2) \\ &\geq \frac{1}{2} \int |\nabla_A u|^2 + \int V_e |u|^2 - \frac{\lambda n}{n+2} \int |u|^2 - \frac{\lambda n}{n+2} c_{GN} \|u\|_2^{\frac{4}{n}} \|\nabla_A u\|_2^2 \\ (67) \quad &= \frac{1}{2} \left(1 - \frac{\|u\|_2^{4/n}}{\|Q_{\lambda,1}\|_2^{4/n}} \right) \int |\nabla_A u|^2 + \int V_e |u|^2 - C_0. \end{aligned}$$

Like in the proof of Theorem 7.2, we have, given $c < \|Q_{\lambda,1}\|_2$, then the above estimate shows:

- (i) For all $u \in S_c$, $E_{\Omega,\gamma}^1(u)$ is bounded from below by a constant. Hence I_c exists.
- (ii) Let $\{u_n\} \subset S_c$ be a minimizing sequence for (52). Then it is bounded in Σ .
- (iii) Since Σ compactly embedded in L^r , $r \in [2, \infty)$, there exists $w \in \Sigma$ and a subsequence, still denoted by $\{u_n\}$, such that $\int G(|u_n|^2) \rightarrow \int G(|w|^2)$.
- (iv) This implies that up to a subsequence, $E_{\Omega,\gamma}(w) = \lim_n E_{\Omega,\gamma}(u_n)$ and $u_n \rightarrow w$ in Σ . In particular, w is a minimizer.

This proves part (a) of the theorem. Part (b) is a special case of part (a). \square

Remark 7.9. The theorem suggests that in the focusing, mass-critical case, there exists ground state solutions on the mass level set S_c that are below the free ground state $Q_{\lambda,1}$. However, such solutions are not unique, consult e.g. [17]. If $c > \|Q\|_2$, then there may not exist a solution for (52), see [25] for the case where V is a quadratic potential and $\Omega = 0$. Compare also [3] where the case of anisotropic V is considered for $p < 1 + 4/n$, $n = 2, 3$.

8. ORBITAL STABILITY OF STANDING WAVES

In this section, we prove the orbital stability of the standing waves of (48). Throughout this section we will assume G satisfies:

$(\Omega, \gamma)_0$: $|\Omega| < \gamma$.

(G'_0) : Let $G : \mathbb{R}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be continuous and differentiable such that for some constant $C > 0$ and $1 \leq p < 1 + \frac{4}{n-2}$,

$$(68) \quad |\partial_v G(x, v)| \leq C(1 + v^{\frac{p-1}{2}}).$$

Note that (G'_0) is slightly stronger than (G_0) . It is easy to see from the condition (G'_0) and the embedding $H^1 \hookrightarrow L^r$, $r \in [2, \frac{2n}{n-2})$

$$(69) \quad E_{\Omega, \gamma} : \Sigma \rightarrow \mathbb{R} \quad \text{is continuous.}$$

Proposition 8.1. *Let G satisfy the conditions $(\Omega, \gamma)_0$ and (G'_0) for $p > 1$ and $\kappa = \pm 1$. Let $r = p + 1$ and $q = \frac{2(p+1)}{(p-1)}$.*

- (a) *Given any u_0 in Σ , there exists a unique maximal solution u of (48) in $C(I, \Sigma) \cap L^q(I, H^{1,r}) \cap C^1(I, \Sigma^{-1})$ on $I = (-T_{min}, T_{max})$ such that if T_{max} (or T_{min}) is finite, then $\|u(t)\|_{\Sigma} \rightarrow \infty$ as $t \rightarrow T_{max}$, (respectively $t \rightarrow -T_{min}$). Moreover, the following conservation laws hold on I :*

$$\begin{aligned} \|u\|_2 &= \|u_0\|_2 \\ E_{\Omega, \gamma}(u) &= E_{\Omega, \gamma}(u_0). \end{aligned}$$

- (b) *The solution u of (48) is global in time if one of the following holds:*
- (i) $\kappa = 1$, $p = 1 + \frac{4}{n}$ and $\|u_0\|_2 < \|Q_{\lambda, 1}\|_2$, where λ is given as in Theorem 7.8.
 - (ii) $\kappa = -1$ and $1 < p < 1 + \frac{4}{n-2}$.

The proof of the local theory above is similar to that of Proposition 2.1 and Theorem 1.1 (a), hence omitted. Also, see [13, Theorem 1] for the case $|\Omega| = \gamma$. The global existence follows from the estimate $\|u(t, \cdot)\|_{\Sigma} \leq C \|u_0\|_{\Sigma}$ in view of (67) in the case $\kappa = 1$ and $\|u_0\|_2 < \|Q_{\lambda, 1}\|_2$. As we will see, the stability requires the local existence and uniqueness for $u_0 \in \mathcal{H}^1$ along with the mass and energy conservation laws in the proposition above.

For $c > 0$ let

$$(70) \quad Z_c = \{u \in \Sigma : E_{\Omega, \gamma}(u) = I_c, \|u\|_2 = c\},$$

where I_c is the minimum given by (51). Then Z_c is the set of all ground state solutions for (52).

Definition 8.2. The set Z_c is called orbitally stable if given any $\varphi \in Z_c$, the following property holds: $\forall \varepsilon > 0, \exists \delta > 0$ such that

$$(71) \quad \|u_0 - \varphi\|_\Sigma < \delta \Rightarrow \sup_t \inf_{\phi \in Z_c} \|u(t, \cdot) - \phi\|_\Sigma < \varepsilon,$$

where $u_0 \mapsto u(t, \cdot)$ is the unique solution flow map of (48).

Theorem 8.3. *Assume the condition $(\Omega, \gamma)_0$ and (G'_0) . Let $c > 0$. Suppose one of the following is satisfied:*

- (a) $\kappa = 1, 1 < p < 1 + \frac{4}{n}$;
- (b) $\kappa = 1, p = 1 + \frac{4}{n}$ and $c < \|Q_{\lambda,1}\|_2$;
- (c) $\kappa = -1, 1 < p < 1 + \frac{4}{n-2}$.

Then Z_c is orbitally stable for (48). In particular, if $G(v) = \frac{2\lambda}{p+1}v^{(p+1)/2}$ for $v \geq 0$ with $\kappa = 1$, we obtain Theorem 1.2.

Proof. From Theorem 7.2 and Theorem 7.8 we know $Z_c \neq \emptyset$. According to Proposition 8.1, we have

- (i) RNLS (48) has global in time solution, that is, the lifespan is $(-\infty, \infty)$;
- (ii) RNLS (48) enjoys the conservation laws for mass and energy.

In order to prove the orbital stability, we argue by contradiction. Suppose Z_c is not stable, then there exists $w_0 \in Z_c, \varepsilon_0 > 0$ and a sequence $\{\psi_{0,n}\} \subset \Sigma$ such that

$$(72) \quad \|\psi_{0,n} - w_0\|_\Sigma \rightarrow 0 \text{ as } n \rightarrow \infty \text{ but } \inf_{\phi \in Z_c} \|\psi_n(t_n, \cdot) - \phi\|_\Sigma \geq \varepsilon_0$$

for some sequence $\{t_n\} \subset \mathbb{R}$, where ψ_n is the solution of (48) corresponding to the initial data $\psi_{0,n}$.

Now let $z_n = \psi_n(t_n, \cdot)$. Since $w_0 \in S_c$ and $E_{\Omega,\gamma}(w_0) = I_c$, it follows from the continuity of $\|\cdot\|_2$ and the continuity of $E_{\Omega,\gamma}$ on Σ , that $\|\psi_{0,n}\|_2^2 \rightarrow c^2$ and $E_{\Omega,\gamma}(\psi_{0,n}) \rightarrow I_c$. By the conservation of the mass $\|\cdot\|_2$ and the energy $E_{\Omega,\gamma}$, we certainly have $\|z_n\|_2^2 \rightarrow c^2$ and $E_{\Omega,\gamma}(z_n) \rightarrow I_c$. Therefore, $\{z_n\}$ is a minimizing sequence. According to Corollary 7.3 and Theorem 7.8, under any one of the hypotheses in (a), (b) or (c), $\{z_n\}$ contains a subsequence $\{z_{n_j}\}$ converging to an element w in Σ as $j \rightarrow \infty$, such that $\|w\|_2^2 = c^2$ and $E_{\Omega,\gamma}(w) = I_c$. Thus $w \in Z_c$, and as a consequence

$$\inf_{\phi \in Z_c} \|\phi_{n_j}(t_{n_j}, \cdot) - \phi\|_\Sigma \leq \|z_{n_j} - w\|_\Sigma \rightarrow 0,$$

as $j \rightarrow \infty$. This contradicts (72). Therefore, we have established the orbital stability. \square

Remark 8.4. Concerning the usual nonlinearity $N(x, u) = |u|^{p-1}u$ with $\kappa = 1$, if $p = 1 + \frac{4}{n}$, $c \geq \|Q\|_2$, or $p > 1 + \frac{4}{n}$, then strong instability, namely, blowup can occur for (1), see Theorem 1.1, Lemma 4.1 and [2, 8].

Remark 8.5. The stability of ground state solution for mNLS in \mathbb{R}^3 was initially considered in [13] and certain instability was studied in [22] and [19] for the mass-supercritical case. Our treatment is motivated by [14] and [26] on the concentration compactness method and the extension to NLS with a general nonlinearity given by (50) by showing any minimizing sequence is relatively compact in Σ . Comparing [13] the major improvement of our results are the following:

- (i) We treat the case $V_e = V - \frac{|A|^2}{2} \neq 0$ in all dimensions, which extends the results in [13, 17] where $V_e = 0$. Note that this extension requires a norm characterization $\|u\|_\Sigma \approx \|\nabla_A u\|_2 + \|xu\|_2$, which was obtained in [50] and entails a non-trivial proof using harmonic analysis.
- (ii) On a remarkable level, we prove a sharp threshold result on the stability in the mass-critical case for (48), which can be viewed as a complement on the mass-subcritical result in [13] or [3] if $n = 2, 3$. To prove this we have used a sharp diamagnetic Gagliardo-Nirenberg type inequality in Lemma 7.7.

8.6. Instability when $|\Omega| > \gamma$: A counterexample. In this subsection, we provide an example in \mathbb{R}^2 to show non-existence of minimizers for (70) if $|\Omega| > \gamma$ (fast rotating). Our consideration is motivated by [9], [44], and [5]. This counterexample shows there does not exist a ground state solution to (52) in either focusing or defocusing case. Note that in physics the state of the system is stable only when it attains its lowest energy. So physically, there exhibits instability in the fast-rotating scenario, in consistence with the numerical simulations.

Let $\Omega > \gamma > 0$ and let $\kappa = 1$, $G(v) = K(v + v^{a/2})$, $a \in (2, \infty)$. Define a sequence of vortex (test) functions $\{\psi_m\} \subset \Sigma$ for $m \in \mathbb{Z}$ using $x = (r, \theta)$, the polar coordinate,

$$\psi_m(x) = \frac{\gamma^{\frac{|m|+1}{2}}}{\sqrt{\pi(|m|)!}} |x|^{|m|} e^{-\gamma|x|^2/2} e^{im\theta}.$$

Then $\|\psi_m\|_2 = 1$ for all m . We have $E_{\Omega, \gamma}(\psi_m) \rightarrow -\infty$ as $m \rightarrow +\infty$.

Indeed, we compute by (49)

$$\begin{aligned} E_{\Omega,\gamma}(\psi_m) &= \frac{1}{2} \int |\nabla \psi_m|^2 + \int V |\psi_m|^2 - \Omega \int \overline{\psi_m} L_z(\psi_m) - \int G(|\psi_m|^2) \\ &= \frac{|m|+1}{2} \gamma + \frac{|m|+1}{2} \gamma - \Omega m - K - 2\pi K \int_0^\infty |\psi_m|^a r dr \\ &= (|m|+1)\gamma - \Omega m - K + o(1), \end{aligned}$$

where it is easy to calculate for $L_z = -i\partial_\theta$, $V(x) = \frac{\gamma^2}{2}|x|^2$ and $a > 2$

$$\begin{aligned} \int |\nabla \psi_m|^2 &= (|m|+1)\gamma \\ \int V |\psi_m|^2 &= \frac{|m|+1}{2} \gamma \\ \int \overline{\psi_m} L_z(\psi_m) &= m \\ \int |\psi_m|^a &= \gamma^{\frac{a}{2}-1} \pi^{\frac{3}{2}(1-\frac{a}{2})} \frac{1}{2^{\frac{a}{4}-1} \sqrt{a} (\sqrt{|m|})^{\frac{a}{2}-1}} + o(1) \quad \text{as } |m| \rightarrow \infty. \end{aligned}$$

Remark 8.7. From the calculations we see that V may augment the kinetic energy, while $L_\Omega = -\Omega L_z$ may add to

$$\begin{cases} \text{negative angular momentum energy} & \text{as } m \rightarrow +\infty \\ \text{positive angular momentum energy} & \text{as } m \rightarrow -\infty \end{cases}$$

and have larger effect over the nonlinearity in the fast rotating regime.

9. CONCLUDING REMARKS

We conclude our paper with some remarks regarding the applicability and extension of our study to other related problems. This paper mainly addresses the threshold for the blowup of mass critical focusing RNLS (1). We are able to solve a hard problem using new ideas and techniques that we think would open the way to the study on the threshold dynamics theory that generally includes the blowup rate, blowup profile and location of the blowup points of the solutions as well as the stability *near the threshold*. These would provide precise descriptions of the blowup solutions near certain ground state level, in consistence with the experimental evidence in physics lab owing to the principle that the state of the particle maintains stable when it attains the lowest energy.

1. Equations (1) and (41) have extensions to general rotational NLS by reformulating the term $L_\Omega = iA \cdot \nabla$, where $A = Mx$ with $x \in \mathbb{R}^n$ and M being any skew-symmetric matrix [4, 6, 21]. The results in

this paper, e.g. Theorem 1.1 and Theorem 6.6 extend to such RNLS following the same proofs. We wish to emphasize that although some (non-bordering) results can be obtained from those of (4) by means of the \mathcal{R} -transform, a pseudo-conformal or lens type transform introduced in [6], cf. also [12], certain crucial tools and properties, e.g., Lemma 4.1 and the blowup of the endpoint Q -profile, cannot be obtained this way. Indeed, the solution of the variance $J(t)$ in (46) reveals that the standard NLS is the limiting case of the RNLS as $\gamma \rightarrow 0$, but that limiting case does not directly provide an ultimate threshold information for the blowup profile.

2. In the mass supercritical case $p \in (1 + \frac{4}{n}, 1 + \frac{4}{n-2})$, the blowup dynamics appears very subtle for $\|u_0\|_2 = \|Q\|_2$, where Q is the unique ground state of (2). Within an arbitrarily small neighborhood of Q , there always exist φ_0 and ψ_0 such that the flow $\varphi_0 \mapsto \varphi$ blowups in finite time, and, $\psi_0 \mapsto \psi$ exists globally in time in \mathcal{H}^1 . We have seen that the criteria in Lemma 4.1 give sufficient blowup conditions if $p > 1 + 4/n$. It would be of interest to further investigate the *sharp* threshold problem in the mass-supercritical regime. Moreover, one should be able to adapt the method and techniques in this paper to prove the analogue for some nonlocal problems such as RNLS with a Hartree nonlinearity. In the energy-critical regime, one can consider (1) e.g. in the case where the power nonlinearity becomes exponential type in two dimensions [15, 42].

3. Technically, in order to give a deeper sharp description of the singularity formation and the local asymptotic stability of the “self-similar” blow-up profile for $p \geq 1 + 4/n$, it would require computing the trajectory of the solution on the soliton manifold [34, 46]. In our setting, it is natural and easier to consider the blowup and stability on a submanifold \mathcal{O} of \mathcal{H}^1 . In \mathbb{R}^3 , \mathcal{O} can be taken as the set of all the m -vortex solitons of the form $Q^{(m)}(r, \theta, z) = e^{im\theta} \phi(r, z)$ using cylindrical coordinates [13, 17, 43]. We need to analyze the spectral properties of the associated linearization operators $L_+^{(m)} = H_{\Omega, V} + 1 - p|Q^{(m)}|^{p-1}$ and $L_-^{(m)} = H_{\Omega, V} + 1 - |Q^{(m)}|^{p-1}$, which may lead to solving the profile equation of (1).

4. The method and results on the orbital stability and the construction of the ground state solutions of (1) and (48) apply to focusing and defocusing cases. Such treatment might enable one to further study the vortex soliton stability for the Pauli-Schrödinger system with sharper insight.

5. Finally, we raise the question on the relativistic spin- $\frac{1}{2}$ model NLD [7, 40]: Determine the threshold for the blowup and stability for the nonlinear Dirac equation in higher dimensions?

Acknowledgments This work was initiated when S.Z. visited the NYU institute in Shanghai during summer 2015. Further, S. Z. would like to thank W. A. Strauss and J. Holmer for helpful discussions during his visit at Brown University fall 2015. He also thanks the hospitality and support of the Department of Mathematics at Brown for an enjoyable environment.

REFERENCES

- [1] A. Aftalion, Vortices in Bose-Einstein condensates. Progress in Nonlinear Differential Equations and Their Applications **67**, Birkhäuser, 2006.
- [2] P. Antonelli, D. Marahrens, C. Sparber, On the Cauchy problem for nonlinear Schrödinger equations with rotation. Discrete Contin. Dyn. Syst. **32** (2012), no. 3, 703–715.
- [3] J. Arbutich, I. Nenciu, C. Sparber, Stability and instability properties of rotating Bose-Einstein condensates. Lett Math Phys **109** (2019), 1415–1432.
- [4] W. Bao, Y. Cai, Mathematical theory and numerical methods for Bose-Einstein condensation. Kinetic and Related Models **6** (2013), 1–135.
- [5] W. Bao, H. Wang, P. Markowich, Ground, symmetric and central vortex states in rotating Bose-Einstein condensates. Comm. Math. Sci. **3** (2005), no. 1, 57–88.
- [6] N. Basharat, Y. Hu, S. Zheng, Blowup rate for mass critical rotational nonlinear Schrödinger equations. Contemporary Mathematics **725** (2019), 1–12.
- [7] N. Boussaïd, A. Comech, Nonlinear Dirac Equation: Spectral Stability of Solitary Waves. Mathematical Surveys and Monographs **244** AMS 2019.
- [8] N. Boussaïd, H. Hajaiej, S. Ibrahim, L. Michel, On the global Cauchy problem for non-linear Schrödinger equation with magnetic potential. Preprint.
- [9] D. Butts, D. Rokhsar, Predicted signatures of rotating Bose-Einstein condensates. Nature **397** (1999), 327–329.
- [10] R. Carles, Remarks on nonlinear Schrödinger equations with harmonic potential. Annales Henri Poincaré **3** (2002), no. 4, 757–772.
- [11] R. Carles, Critical nonlinear Schrödinger equations with and without harmonic potential. Math. Mod. Meth. Appl. Sci. **12** (2002), no.10, 1513–1523.
- [12] R. Carles, Nonlinear Schrödinger equation with time dependent potential. Commun. Math. Sci. **9** (2011), no. 4, 937–964.
- [13] T. Cazenave, M. Esteban, On the stability of stationary states for nonlinear Schrödinger equations with an external magnetic field. Matemática Aplicada Comput. **7** (1988), no. 3, 155–168.
- [14] T. Cazenave, P.-L. Lions, Orbital stability of standing waves for some nonlinear Schrödinger equations. Comm. Math. Phys. **85** (1982), 549–561.
- [15] J. Colliander, S. Ibrahim, M. Majdoub, N. Masmoudi, Energy critical NLS in two space dimensions. J. Hyper. Diff. Eqns. **6** (2009), no.3, 549–575.

- [16] A. De Bouard, Nonlinear Schrödinger equations with magnetic fields. *Differential Integral Equations* **4** (1991), no. 1, 73–88.
- [17] M. Esteban, P.-L. Lions, Stationary solutions of nonlinear Schrödinger equations with an external magnetic field. In: *PDEs and Calculus of Variations. Progress in Nonlinear Differential Equations and Their Applications* **1**, 401–449, Birkhäuser, 1989.
- [18] L. Fanelli, L. Vega, Magnetic virial identities, weak dispersion and Strichartz inequalities. *Math. Ann.* **344** (2009), 249–278.
- [19] R. Fukuizumi, M. Ohta, Instability of standing waves for nonlinear Schrödinger equations with potentials. *Differential Integral Equations* **16** (2003), no. 6, 691–706.
- [20] L. Galati, S. Zheng, Nonlinear Schrödinger equations for Bose-Einstein condensates. *Nonlinear and Modern Mathematical Physics. Amer. Institute Phys. Conference Proceedings* **50** (2013), 1562.
- [21] A. Garcia, Magnetic virial identities and applications to blow-up for Schrödinger and wave equations. *Journal of Physics A: Mathematical and Theoretical* **45** (2012), no. 1, 15202–15217.
- [22] J. Gonçalves Ribeiro, Instability of symmetric stationary states for some nonlinear Schrödinger equations with an external magnetic field. *Annales de l' IHP Physique théorique. sec. A* **54** (1991), no. 4, 403–433.
- [23] J. Gonçalves Ribeiro, Finite time blow-up for some nonlinear Schrödinger equations with an external magnetic field. *Nonl. Anal., Theory Methods & Applications* **16** (1991) no. 11, 941–948.
- [24] Y. Guo, Y. Luo, W. Yang, Refined mass concentration of rotating Bose-Einstein condensates with attractive interactions. *ArXiv:1901.09619*, 2019.
- [25] Y. Guo, R. Seiringer, On the mass concentration for Bose-Einstein condensates with attractive interactions. *Lett. Math. Phys.* **104** (2014), 141–156.
- [26] H. Hajaiej, C. Stuart, On the variational approach to the stability of standing waves for the nonlinear Schrödinger equation. *Adv. Nonlinear Studies* **4** (2004), 469–501.
- [27] C. Hao, L. Hsiao, H. Li, Global well posedness for the Gross-Pitaevskii equation with an angular momentum rotational term in three dimensions. *J. Math. Phys.* **48** (2007), no. 10, 102105.
- [28] M. Keel, T. Tao, Endpoint Strichartz estimates. *Amer. J. Math.* **120** (1998), no. 5, 955–980.
- [29] M.-K. Kwong, Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbb{R}^n . *Arch. Rational Mech. Anal.* **105** (1989), no. 3, 243–266.
- [30] M. Lewin, P. Nam, N. Rougerie, Blow-up profile of rotating 2D focusing Bose gases. In: *Macroscopic Limits of Quantum Systems. Springer Proceedings in Mathematics & Statistics* **270** (2018), 145–170.
- [31] M. Lewin, P. Nam, N. Rougerie, The mean-field approximation and the nonlinear Schrödinger functional for trapped Bose gases. *Trans. Amer. Math. Soc.* **368** (2016), 6131–6157.
- [32] E. Lieb, R. Seiringer, Derivation of the Gross-Pitaevskii equation for rotating Bose gases. *Comm. Math. Physics* **264** (2006), no. 2, 505–537.
- [33] Q. Liu, Y. Zhou, J. Zhang, W. Zhang, Sharp condition of global existence for nonlinear Schrödinger equation with a harmonic potential. *Appl. Math. Comput.* **177** (2006), no. 2, 482–487.

- [34] Y. Martel, F. Merle, P. Raphaël, J. Szeftel, Near soliton dynamics and singularity formation for L^2 critical problems. *Uspekhi Mat. Nauk* **69** (2014), no.2, 77–106.
- [35] M. Matthews, B. Anderson, P. Haljan, D. Hall, C. Wieman, E. Cornell, Vortices in a Bose-Einstein Condensate. *Phys. Rev. Lett.* **83** (1999), 2498.
- [36] F. Merle, Determination of blow-up solutions with minimal mass for nonlinear Schrödinger equations with critical power. *Duke Math. J.* **69** (1993), no. 2, 427–454.
- [37] F. Merle, Nonexistence of minimal blow-up solutions of equations $iu_t = -\Delta u - k(x)|u|^{4/N}u$ in \mathbb{R}^N . *Annales de l’institut Henri Poincaré (A) Physique théorique* **64** (1996), no. 1, 33–85.
- [38] F. Merle, P. Raphaël, Blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation. *Ann. Math.* **16** (2005), 157–222.
- [39] L. Michel, Remarks on non-linear Schrödinger equation with magnetic fields. *Comm. Partial Differential Equations* **33** (2008), 1198–1215.
- [40] D. Pelinovsky, Y. Shimabukuro, Orbital stability of Dirac soliton. *Lett Math Phys* **104** (2014), 21–41.
- [41] P. Raphaël, J. Szeftel, Existence and uniqueness of minimal blow-up solutions to an inhomogeneous mass critical NLS. *J. Amer. Math. Soc.* **24** (2011), 471–546.
- [42] T. Saanouni, Global well-posedness and instability of a nonlinear Schrödinger equation with harmonic potential. *J. Austral. Math. Soc.* **98** (2015), no. 1, 78–10.
- [43] G. Simpson, I. Zwiars, Vortex collapse for the L^2 -critical nonlinear Schrödinger equation. *J. Math. Physics* **52** (2011), no.8, 083503.
- [44] R. Seiringer, Gross-Pitaevskii theory of the rotating Bose gas. *Comm. Math. Phys.* **229** (2002), 491–509.
- [45] M. Weinstein, Nonlinear Schrödinger equations and sharp interpolation estimates. *Comm. Math. Phys.* **87** (1983), no. 4, 567–576.
- [46] M. Weinstein, Modulational stability of ground states of nonlinear Schrödinger equations, *SIAM J. Math. Anal.* **16** (1985), 472–491.
- [47] K. Yajima, Schrödinger evolution equations with magnetic fields. *Journal d’Analyse Mathématique* **56** (1991), 29–76.
- [48] K. Yang, S. Roudenko, Y. Zhao, Blow-up dynamics and spectral property in the L^2 -critical nonlinear Schrödinger equation in high dimensions. *Nonlinearity* **31** (2018), 4354–4392.
- [49] J. Zhang, Stability of attractive Bose-Einstein condensates. *J. Statist. Phys.* **101** (2000), 731–746.
- [50] S. Zheng, Fractional regularity for nonlinear Schrödinger equations with magnetic fields. *Contemporary Mathematics* **581** (2012), 271–285.

(Nyla Basharat) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, SASKATOON, SK S7N 5E6, CANADA

E-mail address: nylabasharat12@gmail.com

(Hichem Hajaiej) DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, LOS ANGELES, CA 90032

E-mail address: hhajaie@calstatela.edu

(Yi Hu) DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGIA SOUTHERN UNIVERSITY, STATESBORO, GA 30460

E-mail address: `yihu@GeorgiaSouthern.edu`

(Shijun Zheng) DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGIA SOUTHERN UNIVERSITY, STATESBORO, GA 30460

E-mail address: `szheng@GeorgiaSouthern.edu`