

The finite dimensional subalgebra classification of infinite dimensional symmetry algebra of two dimensional coupled nonlinear Schrödinger equations*

YueXing Bai¹, Temuer Chaolu^{2†}, Yan Li²

(1. College of Arts and Sciences, Shanghai Maritime University, Shanghai 200135, China)

(2. Inner Mongolia University of Technology, Huhhot, 010051, China)

Abstract: The symmetry group structures of two dimensional coupled nonlinear Schrödinger equations are considered. We first show that the equations admit infinite dimensional symmetry algebra as well as the corresponding symmetry group depending on four arbitrary functions of one variable. Then we show some physical symmetries and an affine loop algebra contained in the symmetry algebra of the equations. Third, we give the complete classifications of finite dimension (less than four) subalgebras of the symmetry algebra under the adjoint group of the symmetry group. These results provide the theoretical and computational basis for the further study of the equations with symmetry methods.

Keywords: Infinite dimensional symmetry group (algebra); 2D-coupled nonlinear Schrödinger Equation; Subalgebra classification

1 Introduction

The nonlinear Schrödinger typed equations are of considerable importance in both theory and applications. Particularly, the various coupled nonlinear Schrödinger systems are used in different fields of physics and mathematics [1]. One of them, a system of (2+1)-dimensional coupled nonlinear Schrödinger (2D-CNLS) system

$$\Omega : \begin{cases} iu_t - pu_{xx} + qu_{yy} + r|u|^2u - 2uv = 0, \\ pv_{xx} + qv_{yy} - pr(|u|^2)_{xx} = 0, \end{cases} \quad (1)$$

arising in the describing the evolution of the wave packet on a two-dimensional water surface under gravity [2, 3], has been investigated in series of papers [4–12]. The evolutions of packets of various waves were discussed in [4–7]; the multi-soliton solutions were obtained in [8] in which the two-soliton resonant interaction and a triple soliton structure were found; different types of lump solutions were found in [9, 10]; particular symmetry reductions and corresponding similarity solutions are obtained in [11]. The one dimensional optimal system of an eight dimensional symmetry algebra and corresponding reductions of system (1) are given in [12].

*Supported by National Natural Science Foundation of China (11571008).

†Corresponding author E-mail: tmchaolu@shmtu.edu.cn

Actually, as we showed in [12], the system (1) admits infinite dimensional symmetry (Lie) algebra of the system (we call it as 2D-CNLS algebra) depending on four arbitrary functions of single variable t with corresponding infinite dimensional symmetry group (we call it as 2D-CNLS group). As usual, to solve system (1) by the symmetry method, we need to identify the subalgebras of the 2D-CNLS algebra. More particularly, we need subalgebras that correspond to Lie groups having orbits of codimension less than four [13]. In general sense, what structures of finite dimensional subalgebras does an infinite dimensional algebra have and how to identify them are nontrivial. This problem is equivalent to the subalgebra classification problem under an algebraic equivalent criterion. For finite dimensional Lie algebra, the classification methods have been developed in [13–16]. In [17, 18], the authors applying the methods gave the subalgebra classification of infinite dimensional symmetry algebras of the Kadomtsev- Petviashvili equation and the selfdual Einstein equations. These are the pioneering works of investigating the structure of infinite dimensional symmetry group (algebra) admitted by a partial differential equations (PDEs). It has been turned out that many of the important nonlinear PDEs of modern physics admit infinite dimensional symmetry algebra depending on arbitrary functions. Such are the cases studied in [17, 18] and 2D-CNLS treated in this article, but also of other PDEs in higher dimensions. For instance, in Earth science, the PDEs of dynamic convection in a sea [19] admit infinite dimensional symmetry algebra with five arbitrary functions; In incompressible fluids, the two (three) dimensional Navier- Stokes equations admit symmetry algebra with three (four) arbitrary functions [20]; in nonlinear optics and acoustics, the transparent atmosphere equations admit symmetry algebra with five arbitrary functions [21]; in geophysical fluid dynamics the two dimensional nonlinear Boussinesq equations admit symmetry algebra with three arbitrary functions [22]. The presence of arbitrary functions in the symmetry algebra is a characteristic property of such science fields [20,23]. However the structure of the finite dimensional subalgebras of the such infinite dimensional symmetry algebra of a PDEs and the corresponding complete reductions of the PDEs have yet not been fully studied.

In this article, using the methods in [17], we give the classification of the subalgebras of dimension less than four of the 2D-CNLS algebra under the adjoint group (action) of the 2D-CNLS group (the group of inner automorphisms of the 2D-CNLS algebra). It will derive a better understanding of the structure of the 2D-CNLS algebra and clarify the invariant solutions of different classes of the system (1) under the adjoint group. The classification also further show the applicability of the methods developed for classifying optimal subalgebras of finite-dimensional Lie algebras.

The present paper is organized as follows: In Sec. II, we show the 2D-CNLS algebra and the corresponding 2D-CNLS group. We establish a Levi decomposition of the algebra. For Laurent polynomials form of the arbitrary functions in the algebra, we identify the Lie algebra as a subalgebra of the loop algebra $A_4^{(1)}$, a Kac-Moody type of algebra. We also present the physical meaning of the finite-dimensional algebras obtained by restricting the arbitrary functions involved in 2D-CNLS algebra to first and second degree polynomials. In Sec. III, the properties of the action of the adjoint group corresponding to the basis elements of the 2D-CNLS algebra are presented. These properties are frequently used in the computations for obtaining the classification of the 2D-CNLS algebra. In Sec. IV, as the heart of present article, the classification of the less then four dimension subalgebras under the adjoint action is presented. Finally, in Sec.V, we give some brief conclusion remarks on our results.

Here a remark about some notations frequently used in this article.

Remark 1:

Use \mathbb{R} denote the complex number and real number fields respectively.

Notation $C^\infty(\mathbb{R})$ represents the set of infinite order differential functions on \mathbb{R} .

Use notations $f, g, h, k, \phi, \psi, \Phi, \Psi, F, G, K, H$ to denote the arbitrary functions of single variable t in $C^\infty(\mathbb{R})$ and use $\alpha, \beta, \lambda, \delta, \mu, \nu, \sigma, \gamma, \rho, \kappa, a, b, c, d$ to denote arbitrary constants in \mathbb{R} .

Denote $\epsilon = q/p \in \mathbb{R}$ for physical constants p and q in 2D-CNLS system (1).

2 The 2D-CNLS algebra and group

2.1 The 2D-CNLS algebra

Let the 2D-CNLS group, i.e. an one-parameter Lie's symmetry group of infinitesimal transformation of system (1) mapping (x, y, t, u, v) into (x', y', t', u', v') , be given as [16, 24]

$$\begin{cases} x' = x + \epsilon\xi(x, y, t, u, v) + O(\epsilon^2), \\ y' = y + \epsilon\eta(x, y, t, u, v) + O(\epsilon^2), \\ t' = t + \epsilon\tau(x, y, t, u, v) + O(\epsilon^2), \\ u' = u + \epsilon\zeta(x, y, t, u, v) + O(\epsilon^2), \\ v' = v + \epsilon\varsigma(x, y, t, u, v) + O(\epsilon^2). \end{cases} \quad (2)$$

Thus the symmetry group is generated by the infinitesimal generator

$$\mathcal{X} = \xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \tau \frac{\partial}{\partial t} + \zeta \frac{\partial}{\partial u} + \varsigma \frac{\partial}{\partial v}, \quad (3)$$

where $\xi = \xi(x, y, t, u, v)$, $\eta = \eta(x, y, t, u, v)$, $\tau = \tau(x, y, t, u, v)$, $\zeta = \zeta(x, y, t, u, v)$ and $\varsigma = \varsigma(x, y, t, u, v)$ are the infinitesimal functions of \mathcal{X} belonging to $C^\infty(\mathbb{R}^5)$. The invariant condition of equation (1) under (2) is given by $\text{Pr}^{(2)}\mathcal{X}(\Omega)|_{\Omega=0} = 0$, where $\text{Pr}^{(2)}\mathcal{X}$ is the second prolongation of generator \mathcal{X} on the jet space of u and v . Algorithms for determining the generator (5) of symmetry groups of a PDEs are well known [16, 24] and also see recent new algorithm in [25, 26]. For the 2D-CNLS system (1), a straightforward application of the algorithm has yielded an infinite dimensional Lie algebra of the symmetry group with the infinitesimal functions given by

$$\begin{aligned} \xi &= f(t) + \frac{1}{2}x\dot{h}(t), \quad \eta = g(t) + \frac{1}{2}y\dot{h}(t), \quad \tau = h(t), \\ \zeta &= \left(ik(t) - \frac{1}{2}\dot{h}(t) - \frac{i}{8pq}(4(qx\dot{f}(t) - py\dot{g}(t)) + (qx^2 - py^2)\ddot{h}(t)) \right) u, \\ \varsigma &= -\dot{h}(t)v - \frac{1}{2}\dot{k}(t) - \frac{1}{16pq} \left(8(qx\ddot{f}(t) - py\ddot{g}(t)) + (qx^2 - py^2)\ddot{\dot{h}}(t) \right). \end{aligned}$$

They depend on four arbitrary functions f, g, h and k of variable t in $C^\infty(\mathbb{R})$. Corresponding to the four arbitrary functions, we have operators

$$\begin{aligned} Z_h &= h(t)\partial_t + \frac{1}{2}\dot{h}(t)(x\partial_x + y\partial_y - u\partial_u - 2v\partial_v) - \frac{qx^2 - py^2}{16pq} \left(2i\ddot{h}(t)u\partial_u - \ddot{\dot{h}}(t)\partial_v \right), \\ X_f &= f(t)\partial_x - \frac{1}{4p}x \left(2ui\dot{f}(t)\partial_u - \ddot{f}(t)\partial_v \right), \\ Y_g &= g(t)\partial_y + \frac{1}{4q}y(2ui\dot{g}(t)\partial_u - \ddot{g}(t)\partial_v), \\ W_k &= ik(t)u\partial_u - \frac{1}{2}\dot{k}(t)\partial_v. \end{aligned} \quad (4)$$

These operators span an infinite dimensional Lie algebra, i.e., 2D-CNLS algebra, denoted it by $\mathfrak{L} = \langle X_f, Y_g, Z_h, W_k \rangle$, for arbitrary differential real functions $f = f(t), g = g(t), h = h(t)$ and $k = k(t)$ of class $C^\infty(\mathbb{R})$. Thus, the general element of the 2D-CNLS algebra \mathfrak{L} can be written as

$$V = X_f + Y_g + Z_h + W_k, \quad (5)$$

and the commutation relations for $V \in \{X_F, Y_G, Z_H, W_K\}$ and $v \in \{X_f, Y_g, Z_h, W_k\}$ of 2D-CNLS algebra \mathfrak{L} is given by

Table 1: Commutator of 2D-CNLS algebra \mathfrak{L}

$[V, v]$	Z_h	X_f	Y_g	W_k
Z_H	$Z_{Hh-\dot{H}h}$	$X_{Hf-\frac{1}{2}\dot{H}f}$	$Y_{H\dot{g}-\frac{1}{2}\dot{H}g}$	W_{Hk}
X_F	$X_{\frac{1}{2}Fh-\dot{F}h}$	$-\frac{1}{2p}W_{Ff-\dot{F}f}$	0	0
Y_G	$Y_{\frac{1}{2}Gh-\dot{G}h}$	0	$\frac{1}{2q}W_{G\dot{g}-\dot{G}g}$	0
W_K	$-W_{h\dot{K}}$	0	0	0

From the commutator, we notice that the 2D-CNLS algebra has a Levi decomposition

$$\mathfrak{L} = \mathfrak{S} \oplus \mathfrak{N}, \quad (6)$$

where $\mathfrak{N} = \{X_f, Y_g, W_k\}$ is a solvable ideal in \mathfrak{L} and $\mathfrak{S} = \{Z_h\}$ is a simple Lie algebra which is isomorphic to the simple algebra [27]

$$\mathfrak{J}(\mathbb{R}) = \{h(\xi)\partial_\xi | h(\xi) \in C^\infty(\mathbb{R})\},$$

of real vector fields on \mathbb{R} defined by the mapping

$$\psi : \mathfrak{J}(\mathbb{R}) \rightarrow S \text{ with } h(\xi)\partial_\xi \mapsto Z_{h(t)}.$$

Particularly, it is noticed that 2D-CNLS algebra \mathfrak{L} can be written

$$\mathfrak{L} = \mathfrak{L}_X + \mathfrak{L}_Y, \quad (7)$$

as linear space, where $\mathfrak{L}_X = \langle Z_h, X_f, W_k \rangle$ and $\mathfrak{L}_Y = \langle Z_h, Y_g, W_k \rangle$ are subalgebras of algebra \mathfrak{L} .

The following properties of the four operators in (4) are obvious and frequently used as a basic arithmetic operations in the computations of classifying the 2D-CNLS algebra.

Lemma 1 *Operators X_f, Y_g, Z_h and W_k are linear in their labeling functions f, g, h and k respectively and they are linear independent for not all zero functions f, g, h and k .*

2.2 The 2D-CNLS group

It is well known that the symmetry groups of a differential equations are the solutions to the initial value problem of Lie equations of operator (??).

For our case, the 2D-CNLS groups are determined in the following two cases.

First, the function $h(t) \neq 0$ and $f(t), g(t)$ and $k(t)$ in (5) are arbitrary. Solving the Lie equations

$$\begin{aligned}
\frac{dt'}{d\varepsilon} &= h(t'), \\
\frac{dx'}{d\varepsilon} &= f(t') + \frac{1}{2}x'\dot{h}(t'), \\
\frac{dy'}{d\varepsilon} &= g(t') + \frac{1}{2}y'\dot{h}(t'), \\
\frac{du'}{d\varepsilon} &= \frac{1}{2}u' \left(2ik(t') - \frac{i}{p}x'\dot{f}(t') + \frac{i}{q}y'\dot{g}(t') - \dot{h}(t') - \frac{i}{4pq}(qx'^2 - py'^2)\ddot{h}(t') \right), \\
\frac{dv'}{d\varepsilon} &= -\frac{1}{2} \left(\dot{k}(t') - \frac{1}{2p}x'\ddot{f}(t') + \frac{1}{2q}y'\ddot{g}(t') + 2v'\dot{h}(t') - \frac{i}{8pq}(qx'^2 - py'^2)\ddot{h}(t') \right), \tag{8}
\end{aligned}$$

with initial values $t'(0) = t, x'(0) = x, y'(0) = y, u'(0) = u, v'(0) = v$, we obtain the 2D-CNLS groups corresponding to 2D-CNLS algebra as the following

$$\begin{aligned}
t' &= \Phi^{-1}(\varepsilon + \Phi(t)), \\
x' &= [x + F(t', t)] \left[\frac{h(t')}{h(t)} \right]^{\frac{1}{2}}, \\
y' &= [y + G(t', t)] \left[\frac{h(t')}{h(t)} \right]^{\frac{1}{2}}, \\
u' &= u(t, x, y) \exp \left(\int^{t'} \left(ik(t') - \frac{ix'\dot{f}(t')}{2p} + \frac{iy'\dot{g}(t')}{2q} - \frac{\dot{h}(t')}{2} - i\ddot{h}(t') \left(\frac{x'^2}{8p} - \frac{y'^2}{8q} \right) \right) \frac{1}{h(t')} dt' \right), \\
v' &= \left(\frac{h(t')}{h(t)} \right)^{-1} \left\{ v(t, x, y) - H(t', t) - \frac{1}{4h(t)} \left(\frac{x'}{p} \left(\frac{\dot{h}(t')f(t')}{2h(t')} - \dot{f}(t') \right) - \frac{y'}{q} \left(\frac{\dot{h}(t')g(t')}{2h(t')} - \dot{g}(t') \right) \right) \right. \\
&\quad + \frac{1}{8h(t)} \left(\frac{h(t')^{-1}g(t')^2 - h(t)^{-1}g(t)^2}{q} + \frac{h(t)^{-1}f(t)^2 - h(t')^{-1}f(t')^2}{p} \right) \\
&\quad \left. + \frac{1}{16h(t)h(t')} \left(\frac{x'^2}{p} - \frac{y'^2}{q} \right) \left(h(t')\ddot{h}(t') - \frac{1}{2}\dot{h}(t')^2 \right) \right\}, \tag{9}
\end{aligned}$$

where

$$\begin{aligned}
\Phi(t) &= \int_{t_0}^t \frac{1}{h(s)} ds, \quad F(t', t) = h(t)^{1/2} \int_t^{t'} f(s)h(s)^{-3/2} ds, \\
G(t', t) &= h(t)^{1/2} \int_t^{t'} g(s)h(s)^{-3/2} ds, \quad H(t', t) = h(t)^{-1} \int_t^{t'} \frac{1}{2}\dot{k}(s) ds,
\end{aligned}$$

and Φ^{-1} denotes the inverse of Φ and ε is the group parameter.

Next, we consider the case $h(t) \equiv 0$ and $f(t), k(t)$ and $g(t)$ are arbitrary. Solving the corresponding initial value problem of the Lie equations, we obtain the 2D-CNLS groups as following

$$\begin{cases} t' = t, \\ x' = x + \varepsilon f(t), \\ y' = y + \varepsilon g(t), \\ u' = u \exp \left(\frac{i}{4pq} \varepsilon (4pqk(t) - q(2x + \varepsilon f(t))\dot{f}(t) + p(2y + \varepsilon g(t))\dot{g}(t)) \right), \\ v' = v - \frac{\varepsilon}{8pq} \left(4pq\dot{k}(t) - q(2x + \varepsilon f(t))\ddot{f}(t) + p(2y + \varepsilon g(t))\ddot{g}(t) \right). \end{cases} \tag{10}$$

The immediate application of the symmetry groups (9) and (10) is to give us new solutions (u', v') from known ones (u, v) . For instance, if we have a initial constant solution $u = a, v = b$, then (9) and (10)

yield a family of solutions to 2D-CNLS system (1) depending on arbitrary functions $f(t), g(t), h(t), k(t)$ and constants a and b .

It is obviously here that 2D-CNLS system (1) is also invariant under the reflections R_x, R_y, R_{xy} and R_u (discrete symmetries),

$$\begin{aligned}
R_x &: t \mapsto t, x \mapsto -x, y \mapsto y, u \mapsto u, v \mapsto v, \\
R_y &: t \mapsto t, x \mapsto x, y \mapsto -y, u \mapsto u, v \mapsto v, \\
R_{xy} &: t \mapsto t, x \mapsto -x, y \mapsto -y, u \mapsto u, v \mapsto v, \\
R_u &: t \mapsto t, x \mapsto x, y \mapsto y, u \mapsto -u, v \mapsto v,
\end{aligned} \tag{11}$$

which are not obtained from (5) by solving its Lie equations.

2.3 Some finite-dimensional subalgebras of physical transformations

In the section, we show that the infinite dimensional 2D-CNLS algebra \mathfrak{L} contains many specific subalgebras. They are interested in physical applications.

1. First order subalgebras

Taking the arbitrary functions $f(t), g(t), h(t)$ and $k(t)$ as the first-order polynomials in t , then we have

$$\begin{aligned}
X_1 &= \partial_x, \quad Y_1 = \partial_y, \quad Z_1 = \partial_t, \quad W_1 = iu\partial_u, \\
X_t &= t\partial_x - \frac{i}{2p}xu\partial_u, \quad Y_t = t\partial_y + \frac{i}{2q}yu\partial_u, \\
Z_t &= \frac{1}{2}x\partial_x + \frac{1}{2}y\partial_y + t\partial_t - \frac{1}{2}u\partial_u - v\partial_v, \quad W_t = itu\partial_u - \frac{1}{2}\partial_v,
\end{aligned}$$

as a basis of an eight-dimensional solvable Lie algebra $\mathfrak{L}_8 = \langle X_t, Y_t, Z_t, W_t, X_1, Y_1, Z_1, W_1 \rangle$. It has a seven-dimensional nilpotent ideal $\mathfrak{N} = \langle X_t, Y_t, W_t, X_1, Y_1, Z_1, W_1 \rangle$.

2. Quadratic subalgebras

a. A three-dimensional subalgebra is obtained from $Z(h)$ by restricting $h = h(t)$ into quadratic polynomials in t . This yields

$$\begin{aligned}
Z_1 &= \partial_t, \quad Z_t = \frac{1}{2}x\partial_x + \frac{1}{2}y\partial_y + t\partial_t - \frac{1}{2}u\partial_u - v\partial_v, \\
Z_{t^2} &= xt\partial_x + yt\partial_y + t^2\partial_t - (t + \frac{qx^2 - py^2}{4pq}i)u\partial_u - 2tv\partial_v.
\end{aligned} \tag{12}$$

Their commutation relations are $[Z_1, Z_{t^2}] = 2Z_t, [Z_t, Z_{t^2}] = Z_{t^2}, [Z_1, Z_t] = Z_1$, so we obtain the algebra $\mathfrak{L}_s = \langle Z_1, Z_t, Z_{t^2} \rangle$ which is isomorphism to $\mathfrak{sl}(2, \mathbb{R})$. The Z_{t^2} generates the projective symmetry group

$$\begin{aligned}
x' &= x/(1 - \varepsilon t), \quad y' = y/(1 - \varepsilon t), \quad t' = t/(1 - \varepsilon t), \\
u' &= (1 - \varepsilon t)u \exp(i(py^2 - qx^2)\varepsilon/4pq(1 - \varepsilon t)), \quad v' = (1 - \varepsilon t)^2v.
\end{aligned}$$

b. Another three-dimensional algebra is obtained by restricting $k = k(t)$ in W_k to quadratic polynomials in t and obtain the basis $W_1 = iu\partial_u, W_t = itu\partial_u - \frac{1}{2}\partial_v, W_{t^2} = it^2u\partial_u - t\partial_v$ with $[W_1, W_t] = [W_1, W_{t^2}] = [W_t, W_{t^2}] = 0$. This is a three-dimensional Abelian Lie algebra. In fact, for any an integer n , we can obtain n -dimensional Abelian algebra from W_k by taking the $k = k(t)$ as an $n - 1$ -degree polynomials in t since the Abelian property of operator W_k for arbitrary functions k .

3. An affine loop algebra

One interesting aspect of the 2D-CNLS algebra is that the two subalgebras \mathfrak{L}_X and \mathfrak{L}_Y in (7) can be simultaneously embedded into an affine loop algebra if the arbitrary functions contained in the algebras are taken as Laurent polynomials. Indeed, let us consider the subalgebra \mathfrak{L}^n of the 2D-CNLS algebra obtained by restricting the functions f, g, h and k in (4) to be Laurent polynomials in t . A basis for this subalgebra is given by

$$\begin{aligned} X_{t^n} &= t^n \partial_x - \frac{i}{2p} n t^{n-1} x u \partial_u + \frac{1}{4p} n(n-1) t^{n-2} x \partial_v, \\ Y_{t^n} &= t^n \partial_y + \frac{i}{2q} n t^{n-1} y u \partial_u - \frac{1}{4q} n(n-1) t^{n-2} y \partial_v, \\ Z_{t^n} &= t^n \partial_t + \frac{1}{2} n t^{n-1} (x \partial_x + y \partial_y - u \partial_u - 2v \partial_v) - \frac{q x^2 - p y^2}{16 p q} n(n-1) (2 u i t^{n-2} \partial_u - (n-2) t^{n-3} \partial_v), \\ W_{t^n} &= i t^n u \partial_u - \frac{1}{2} n t^{n-1} \partial_v, \end{aligned}$$

where $n \in \mathbb{N}$ (Integer set).

The Table 2 in the following is the commutation relations between $\{X_{t^n}, Y_{t^n}, Z_{t^n}, W_{t^n}\}$ and $\{X_{t^m}, Y_{t^m}, Z_{t^m}, W_{t^m}\}$ derived from Table 1.

Table 2: Commutators of $\{X_{t^i}, Y_{t^i}, Z_{t^i}, W_{t^i}\}$ for $i = n, m$

$[\cdot, \cdot]$	Z_{t^m}	X_{t^m}	Y_{t^m}	W_{t^m}
Z_{t^n}	$(m-n)Z_{t^{m+n-1}}$	$(m-\frac{1}{2}n)X_{t^{m+n-1}}$	$(m-\frac{1}{2}n)Y_{t^{m+n-1}}$	$mW_{t^{m+n-1}}$
X_{t^n}	$(\frac{1}{2}m-n)X_{t^{m+n-1}}$	$-\frac{m-n}{2p}W_{t^{m+n-1}}$	0	0
Y_{t^n}	$(\frac{1}{2}m-n)Y_{t^{m+n-1}}$	0	$\frac{m-n}{2q}W_{t^{m+n-1}}$	0
W_{t^n}	$-nW_{t^{m+n-1}}$	0	0	0

Let $\mathfrak{L}_X^n = \langle X_{t^n}, Z_{t^n}, W_{t^n} \rangle$ and $\mathfrak{L}_Y^n = \langle Y_{t^n}, Z_{t^n}, W_{t^n} \rangle$, then $\mathfrak{L}^n = \mathfrak{L}_X^n + \mathfrak{L}_Y^n$ as linear space.

Let us now consider the eleven-dimensional Lie algebra \mathfrak{L}_{11} generated by the following vector fields:

$$\begin{aligned} \Delta &= (1/2)(x \partial_x + y \partial_y - u \partial_u - 2v \partial_v); \quad U = i u \partial_u; \quad V = -(1/2) \partial_v; \\ A &= ((p y^2 - q x^2)/8 p q) U; \quad P = ((p y^2 - q x^2)/8 p q) V; \quad U_x = (-x/2 p) U; \\ V_x &= (-x/2 p) V; \quad U_y = (y/2 q) U; \quad V_y = (y/2 q) V, \quad X = \partial_x; \quad Y = \partial_y. \end{aligned}$$

This is a solvable algebra and its nilradical is spanned by $\{A, P, X, Y, U, V, U_x, U_y, V_x, V_y\}$ and it contains an eight dimensional Abelian ideal spanned by $\{A, P, U, V, U_x, U_y, V_x, V_y\}$. Obviously, algebra \mathfrak{L}_{11} is not a subalgebra of 2D-CNLS algebra. We can write

$$\mathfrak{L}_{11} = \mathfrak{L}_x + \mathfrak{L}_y, \tag{13}$$

as a linear space, where $\mathfrak{L}_x = \langle \Delta, X, U, V, A, P, U_x, V_x \rangle$ and $\mathfrak{L}_y = \langle \Delta, Y, U, V, A, P, U_y, V_y \rangle$. Both \mathfrak{L}_x and \mathfrak{L}_y is solvable subalgebras containing five- dimensional Abelian ideals spanned by $\langle V, A, P, U_x, V_x \rangle$ and $\langle V, A, P, U_y, V_y \rangle$ respectively, with common center $\langle U \rangle$. The two subalgebras should be embedded into simple Lie algebras. It is known that algebra A_4 or in our case $\mathfrak{sl}(5, \mathbb{R})$ is a simple Lie algebra of lowest dimension that contains a five dimensional Abelian subalgebra. Indeed, the following traceless matrices with

eight parameters $\delta, \rho, \gamma, \nu, \alpha, \mu, \sigma$ and κ

$$\mathcal{A} = \begin{pmatrix} \frac{11}{10}\delta & 0 & -\rho & 2\nu & v \\ 0 & \frac{1}{10}\delta & -\alpha & 2\mu & \sigma \\ 0 & 0 & -\frac{9}{10}\delta & \kappa & 0 \\ 0 & 0 & 0 & -\frac{2}{5}\delta & \frac{1}{4p}\kappa \\ 0 & 0 & 0 & 0 & \frac{1}{10}\delta \end{pmatrix} \quad (14)$$

provide a representation of the Lie algebra \mathfrak{L}_x . Each entry in the basis of \mathfrak{L}_x corresponds to a matrix obtained from \mathcal{A} in which one parameter is taken 1 and all other are zero. For instance, the Δ corresponds to the matrix by setting $\delta = 1$ and all other parameters equal to zero in \mathcal{A} . Similarly for X, U, V, A, P, U_x, V_x , we have correspondences: $X \rightarrow \kappa, U \rightarrow \sigma, V \rightarrow v, A \rightarrow \alpha, P \rightarrow \rho, U_x \rightarrow \mu, V_x \rightarrow \nu$. By the same manner, for \mathfrak{L}_y , we have the same form matrix representatives obtained by replacing the p in \mathcal{A} by the q .

Let us now define a natural grading on \mathfrak{L}_x (\mathfrak{L}_y) by attributing the degree “ μ ” to entries of the basis of \mathfrak{L}_x (\mathfrak{L}_y). The degree is equal to the distance between, in the same column, the corresponding parameter term and the diagonal term of \mathcal{A} . Thus Δ has degree 0, A and X (Y) degree 1, P and U_x (U_y) degree 2, U and V_x (V_y) degree 3, V degree 4. Consequently, $\mathfrak{L}_{11} = \mathfrak{L}_x + \mathfrak{L}_y$ embedded in to $\mathfrak{sl}(5, \mathbb{R})$. Meantime, each element in the basis of \mathfrak{L}_X^n (\mathfrak{L}_Y^n) has a well-defined degrees (sum of degrees in t and entry of X, U, V, A, P, U_x (U_y) and V_x (V_y) just defined above), namely $n - 1, n + 1$, and $n + 3$.

From the embedding constructed above for \mathfrak{L}_{11} through \mathfrak{L}_x and \mathfrak{L}_y into $\mathfrak{sl}(5, \mathbb{R})$ and from the representations of \mathfrak{L}^n given below

$$\begin{aligned} Z_{t^n} &= nt^{n-1}\Delta + n(n-1)t^{n-2}A + n(n-1)(n-2)t^{n-3}P + t^n\partial_t, \\ Y_{t^n} &= t^nY + nt^{n-1}U_y + n(n-1)t^{n-2}V_y, \\ X_{t^n} &= t^nX + nt^{n-1}U_x + n(n-1)t^{n-2}V_x, \\ W_{t^n} &= t^nU + nt^{n-1}V, \end{aligned}$$

we see that \mathfrak{L}^n is a subalgebra of the affine loop algebra $A_4^{(1)}$ defined by

$$A_4^{(1)} := \{R[t, t^{-1}] \otimes \mathfrak{sl}(5, R)\} \oplus R[t, t^{-1}] \frac{d}{dt}.$$

The Levi decomposition (6) also holds for \mathfrak{L}^n . Indeed, from the commutation relation in Table 1 we see that $\mathfrak{N} = \{X_{t^n}, Y_{t^n}, W_{t^n}\}$ forms a nilpotent ideal. The element Z_{t^n} form a Lie algebra \mathfrak{S} isomorphic to the simple \mathbb{N} -graded algebra $\mathfrak{L}_t = \mathbb{R}[t, t^{-1}] \frac{d}{dt}$. A basis for \mathfrak{L}_t is given by the set of derivations $(d_n)_{n \in \mathbb{N}}$ given by $d_n = t^n \frac{d}{dt}$ with commutation relations

$$[d_n, d_m] = (m - n)d_{m+n-1}.$$

From the above discussion, we see that the 2D-CNLS system (1) admits rich finite and infinite algebras as well as physical symmetries. In section four, we will give lower dimensional subalgebra classification under the adjoint action of the 2D-CNLS group, which leads to a list of distinguishable equivalent classess of finite dimensional subalgebras of the 2D-CNLS algebra. Consequently, it will clarify the how many equivalent classes of invariant solutions exist for the 2D-CNLS system (1).

3 Adjoint action of 2D-CNLS group

3.1 Adjoint action of 2D-CNLS group

In convenience, let $z = (t, x, y, u, v)$ and $\partial_z = (\partial_t, \partial_x, \partial_y, \partial_u, \partial_v)$.

Let $\Gamma : z \rightarrow \hat{z}$ be a symmetry of system (1) and $\mathcal{X} = z \cdot \partial_z \in \mathfrak{L}$ be the generator of an one-parameter symmetry group of system (1) mapping z to z' . The dot represents the scalar production of z and ∂_z as being seen vectors. Thus, under transformation Γ , we conclude that the transformed operator $\hat{\mathcal{X}}$ of \mathcal{X} by

$$\hat{\mathcal{X}} = \Gamma \mathcal{X} \Gamma^{-1} = \hat{z}' \cdot \partial_{\hat{z}}, \quad (15)$$

is the generator for the conjugate group of the group generated by \mathcal{X} with respect to Γ with $\hat{z}' = \mathcal{X}(\hat{z}) = (\mathcal{X}(\hat{t}), \mathcal{X}(\hat{x}), \mathcal{X}(\hat{y}), \mathcal{X}(\hat{u}), \mathcal{X}(\hat{v}))$.

In particular, if the symmetry Γ is taken as an element Γ_δ of 2D-CNLS group with a parameter δ generated by $\mathcal{Z} \in \mathfrak{L}$, then (15) induces the inner automorphism group $\exp(\delta \cdot ad(\mathcal{Z}))$ of 2D-CNLS algebra \mathfrak{L} (adjoint representation of 2D-CNLS group). The action induced by the inner automorphism group on the Lie algebra \mathfrak{L} is called the adjoint action of the 2D-CNLS group. The conjugating between \mathcal{X} and $\hat{\mathcal{X}}$ is implemented by formula given in [16] as

$$\hat{\mathcal{X}} = \exp(\delta \cdot ad(\mathcal{Z}))\mathcal{X} = \sum_{k=0}^{\infty} \frac{\delta^k}{k!} ad(\mathcal{Z})^k \mathcal{X}, \quad (16)$$

where $ad(\mathcal{Z})$ is the adjoint operator of \mathcal{Z} defined by $ad(\mathcal{Z})\mathcal{Y} = [\mathcal{Y}, \mathcal{Z}]$ with Lie product $[\cdot, \cdot]$ of \mathfrak{L} for all $\mathcal{Y} \in \mathfrak{L}$. We call that two operators \mathcal{X} and $\hat{\mathcal{X}}$ are equivalent if (16) is satisfied for some $\mathcal{Z} \in \mathfrak{L}$ and denote it as $\mathcal{X} \sim \hat{\mathcal{X}}$. This is an equivalent relationship on the algebra \mathfrak{L} . Hence it yields equivalent classification of 2D-CNLS algebra \mathfrak{L} .

3.2 Properties of adjoint action of 2D-CNLS group

Based on the commutator Table 1 and formulae (15) and (16), the adjoint actions $\exp(\delta \cdot ad(V))v$ of 2D-CNLS group with $V \in \{X_F, Y_G, Z_H, W_K\}$ on $v \in \{X_f, Y_g, Z_h, W_k\}$ are given in Table 3 in which $\tilde{f} = f(t) \left(\frac{H(t')}{H(t)} \right)^{\frac{1}{2}}$, $\tilde{g} = g(t) \left(\frac{H(t')}{H(t)} \right)^{\frac{1}{2}}$, $\tilde{h} = h(t) \frac{H(t')}{H(t)}$, $\tilde{F} = \frac{1}{2p}(F\dot{f} - \dot{F}f)$, $\tilde{G} = \frac{1}{2q}(G\dot{g} - \dot{G}g)$, $\tilde{F} = \frac{1}{4p}(\frac{1}{2}F\dot{h} - \dot{F}h)$, $\tilde{G} = \frac{1}{4q}(\frac{1}{2}G\dot{h} - \dot{G}h)$ and $\tilde{k} = \tilde{k}(t') = k(t)$ with $t = \Phi(-\delta + \Phi(t'))$.

Table 3: Adjoint actions of the 2D-CNLS group on \mathfrak{L}

$\exp(\delta \cdot ad(V))v$	Z_h	X_f	Y_g	W_k
$\exp(\delta \cdot ad(Z_H))$	$Z_{\tilde{h}}$	$X_{\tilde{f}}$	$Y_{\tilde{g}}$	$W_{\tilde{k}}$
$\exp(\delta \cdot ad(X_F))$	$Z_h - 4p\delta X_{\tilde{F}} - \delta^2 W_{F\tilde{F} - \dot{F}\tilde{F}}$	$X_f + \delta W_{\tilde{F}}$	Y_g	W_k
$\exp(\delta \cdot ad(Y_G))$	$Z_h - 4q\delta Y_{\tilde{G}} + \delta^2 W_{G\tilde{G} - \dot{G}\tilde{G}}$	X_f	$Y_g - \delta W_{\tilde{G}}$	W_k
$\exp(\delta \cdot ad(W_K))$	$Z_h + \delta W_{h\tilde{K}}$	X_f	Y_g	W_k

We frequently use the following three Theorems as basic operations to normalize the elements of Lie algebra \mathfrak{L} .

Theorem 1 *The symmetry group generated by generator Z_H in 2D-CNLS algebra \mathfrak{L} for a differential nonzero function $H = H(t)$ is given by*

$$\begin{aligned} t' &= \phi^{-1}(\delta + \phi(t)), \\ x' &= x \left(\frac{H(t')}{H(t)} \right)^{\frac{1}{2}}, \\ y' &= y \left(\frac{H(t')}{H(t)} \right)^{\frac{1}{2}}, \\ u' &= u(t, x, y) \exp \left(-\frac{1}{2} \int^{t'} \left(\frac{\dot{H}(t') + \frac{i}{4pq}(qx'^2 - py'^2)\ddot{H}(t')}{H(t')} \right) dt' \right), \\ v' &= \left(\frac{H(t')}{H(t)} \right)^{-1} \left\{ v(t, x, y) + \frac{1}{16pqH(t)H(t')} (qx'^2 - py'^2) \left(H(t')\ddot{H}(t') - \frac{1}{2}\dot{H}(t')^2 \right) \right\}, \end{aligned}$$

where $\Phi(t) = \int^t \frac{1}{H(s)} ds$ and δ is group parameter.

Proof: This is obtained from (9) by setting $h = H$ and $f = g = k = 0$.

Theorem 2 *If $h \neq 0$ in $V = Z_h + X_f + Y_g + W_k \in \mathfrak{L}$, then V is equivalent to Z_1 for arbitrary functions h, f, g and k of variable t , i.e., $V \sim Z_1$.*

Proof: Successively using adjoint actions $\exp(\delta \cdot ad(X_F))$, $\exp(\lambda \cdot ad(Y_G))$ and $\exp(\nu \cdot ad(W_K))$ on V and choosing suitable $F, G, K, \delta, \lambda, \nu$ for given f, g, k , we can normalize V into Z_1 by the following steps.

Step 1. According to Table 3, we first can choose G so as to transform g away from V by acting on it with $\exp(\delta \cdot ad(Y_G))$. Indeed, it can be verified that if we take G satisfying $g - \delta(G\dot{h}/2 - \dot{G}h) = 0$, i.e.,

$$G(t) = \sqrt{h(t)} \left(a \int^t g(s)h(s)^{-3/2} ds + c \right),$$

where $a \neq 0$ and c are arbitrary constants, as the function labeling the element Y_G of the 2D-CNLS algebra and $\delta = -a^{-1}$ as the value of group parameter δ , then we have $\exp(\delta \cdot ad(Y_G))V = Z_h + X_f + W_k$, where actually k is a new function depends on the chosen G and we denote it again k .

Step 2. The next step is to choose the function F so as to transform f away from $Z_h + X_f + W_k$ by performing on it with $\exp(\delta \cdot ad(X_F))$. Again if we take F satisfying $f - \delta(F\dot{h}/2 - \dot{F}h) = 0$, i.e.,

$$F(t) = \sqrt{h(t)} \left(a \int^t f(s)h(s)^{-3/2} ds + c \right),$$

where $a \neq 0$ and c are arbitrary constants, as the function labeling the element X_F of the 2D-CNLS algebra and $\delta = -a^{-1}$ as the value of group parameter δ , then we have $\exp(\delta \cdot ad(X_F))(Z_h + X_f + W_k) = Z_h + W_k$.

Step 3. The third step is to choose the function K so as to drop k away from $Z_h + W_k$ by acting on the $Z_h + W_k$ with $\exp(\delta \cdot ad(W_K))$. Again if we take K satisfying $k + \delta h\dot{K} = 0$, i.e., $K(t) = a \int^t k(s)h(s)^{-1} ds + c$ with $a = -\delta^{-1}$, then we have

$$\exp(\delta \cdot ad(W_K))(Z_h + W_k) = Z_h.$$

Step 4. The last step is to choose the function H so as to normalize h to 1 in Z_h by acting on it with adjoint transformation $\exp(\delta \cdot ad(Z_H))$. Indeed, $\exp(\delta \cdot ad(Z_H))Z_h = Z_{\tilde{h}}$ by formulas (15) and $\tilde{h} = h(t) \frac{H(t')}{H(t)}$

with $t = \Phi^{-1}(-\delta + \Phi(t'))$. It is shown in [17] and references there that there exists a function H satisfying the relation

$$h(t) \frac{H(t')}{H(t)} = 1. \quad (17)$$

Let H be a solution to (17). Then we normalize h to 1. Dropping primes, we have

$$\exp(\delta \cdot ad(Z_H))Z_h = Z_1.$$

This has proved $V \sim Z_1$.

Theorem 3 *For arbitrary functions f, g and k of t , we have*

- a. *if $f \neq 0$, then $X_f + Y_g + W_k \sim X_1 + Y_{\tilde{g}}$ for an arbitrary function \tilde{g} of t ;*
- b. *if $g \neq 0$, then $X_f + Y_g + W_k \sim Y_1 + X_{\tilde{f}}$ for an arbitrary function \tilde{f} of t ;*
- c. *if $fg \neq 0$, then there exist functions \tilde{f}, \tilde{g} of t such that $X_f + Y_g + W_k \sim X_{\tilde{f}} + Y_1 \sim X_1 + Y_{\tilde{g}}$.*

Proof: The proof of a) and b) is included in the proof of c). Hence we give the proof of c) in the following.

As suggested by the commutation relations in Table 1 for the 2D-CNLS algebra, we can first choose F so as to eliminate k from V by acting on it with $\exp(\delta \cdot ad(X_F))$. Since

$$\exp(\delta \cdot ad(X_F))V = X_f + Y_g + W_{k + \frac{\delta}{2p}(F\dot{f} - \dot{F}f)},$$

taking $k + \frac{\delta}{2p}(F\dot{f} - \dot{F}f) = 0$, i.e.,

$$F(t) = f(t) \left(2ap \int^t k(s)f(s)^{-2} ds + c \right),$$

with $a = \delta^{-1}$ and arbitrary constant c , we transform V into $V' = X_f + Y_g$. Then, applying the adjoint map $\exp(\delta \cdot ad(Z_H))$ and the symmetry generated by Z_H (see Theorem 1) on V' , we have

$$\exp(\delta \cdot ad(Z_H))V' = X_{\tilde{f}} + Y_{\tilde{g}},$$

where $\tilde{f} = f(t) \left(\frac{H(t')}{H(t)} \right)^{\frac{1}{2}}$, $\tilde{g} = g(t) \left(\frac{H(t')}{H(t)} \right)^{\frac{1}{2}}$ with $t = \Phi^{-1}(-\delta + \Phi(t'))$.

Like discussion about the existence of function H in (17), we can set either $\tilde{f} = 1$ or $\tilde{g} = 1$ for a selection of H . Consequently, we have the conclusions of the theorem.

3.3 General structures of two-, three- dimensional Lie subalgebras

The following two lemmas are theoretical bases for algebra classification.

Lemma 2 *Any two- dimensional Lie algebra with a basis $\{A_1, A_2\}$ over \mathbb{R} is isomorphic to one of the following ones:*

- (1) *Abelian:* $[A_1, A_2] = 0$.
- (2) *Solvable, non- Abelian:* $[A_1, A_2] = A_1$.

Lemma 3 [17, 18] *Any three- dimensional Lie algebra with a basis $\{A_1, A_2, A_3\}$ over \mathbb{R} can be either simple or solvable. In simple case, the algebra is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$ with a two dimensional non Abelian suabalgebra $\langle A_1, A_2 \rangle$ with the commutations*

$$[A_1, A_2] = A_1, [A_1, A_3] = 2A_2, [A_2, A_3] = A_3. \quad (18)$$

In the solvable case, it always will have a two- dimensional Abelian ideal, i.e., there a basis $\{A_1, A_2, A_3\}$ satisfying

$$[A_1, A_2] = 0, [A_1, A_3] = aA_1 + bA_2, [A_2, A_3] = cA_1 + dA_2, \quad (19)$$

In the isomorphism classification of the three dimensional Lie algebra over \mathbb{R} , the matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ from (19) have six standard forms, i.e.,

$$M_1 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, M_2 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, M_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, M_4 = \begin{bmatrix} 1 & 0 \\ 0 & \alpha \end{bmatrix}, M_5 = \begin{bmatrix} \alpha & 1 \\ -1 & \alpha \end{bmatrix}, M_6 = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}. \quad (20)$$

The details of the classification of three dimensional Lie algebra are seen in [28].

4 Classifications of Low Dimensional Lie Subalgebras

In this section we give the classification of all one-, two-, and three- dimensional subalgebras of the 2D-CNLS algebra under the adjoint action of the 2D-CNLS group.

4.1 One-dimensional subalgebras

We will show that there are four conjugacy classes of one-dimensional subalgebras of the 2D-CNLS algebra under the adjoint action of the 2D-CNLS group, with representatives $Z_1, X_1, X_\phi + Y_1$ and W_ψ , respectively, for arbitrary differential functions ϕ and ψ of t in $C^\infty(\mathbb{R})$. The approach we take is similar to the one used in [17, 18]. The focus here is that one obtains differential conditions on the arbitrary functions labeling the group elements, as seen in proof of Theorem 2. The adjoint action is used to turn the generators of the subalgebras into normal forms.

There are three cases to be considered in the classification of the one- dimensional subalgebras of the 2D-CNLS algebra, generated by typical elements of the form (5) into conjugacy classes under the adjoint group of the 2D-CNLS group.

Case A: $h \neq 0$. By Theorems 1-2, we claim that general element $V = Z_h + X_f + Y_g + W_k$ always can be normalized to Z_1 by the adjoint group of the 2D-CNLS group. Hence, we have a subalgebra $\mathfrak{L}_{1,1} = \langle Z_1 \rangle$.

Case B: $h = 0$ and $f^2 + g^2 \neq 0$. Using Theorem 3, we claim that $V = X_f + Y_g + W_k$ can be transformed into X_1 or $X_f + Y_1$ or Y_1 by the adjoint group of the 2D-CNLS group. That is,

For the case $f \neq 0, g = 0$, we have an essential subalgebra $\mathfrak{L}_{1,2} = \langle X_1 \rangle$.

If $fg \neq 0$, we have subalgebra $\mathfrak{L}_{1,3} = \langle X_\phi + Y_1 \rangle$ for an arbitrary function $\phi \neq 0$.

For the case $f = 0$ and $g \neq 0$, we can prove $V \sim Y_1$. This can be merged into the $\mathfrak{L}_{1,3}$ by allowing $\phi = 0$.

Case C: $h = f = g = 0$ and $k \neq 0$. Because of the invariance of W_k under adjoint actions of 2D-CNLS group, we have subalgebra $\mathfrak{L}_{1,4} = \langle W_\psi \rangle$ for the arbitrary nonzero function ψ .

To summarize, we have proved the following theorem.

Theorem 4 *an arbitrary one - dimensional subalgebra of 2D-CNLS algebra \mathfrak{L} is conjugate, under adjoint group of 2D-CNLS group, to precisely one of the following.*

$$\mathfrak{L}_{1,1} = \langle Z_1 \rangle, \mathfrak{L}_{1,2} = \langle X_1 \rangle, \mathfrak{L}_{1,3} = \langle X_\phi + Y_1 \rangle, \mathfrak{L}_{1,4} = \langle W_\psi \rangle, \quad (21)$$

for arbitrary differential functions ϕ and $\psi \neq 0$.

4.2 Two-dimensional subalgebras

By Lemma 2, there exist two types of two-dimensional Lie algebras both over \mathbb{R} , namely Abelian and solvable non-Abelian algebras spanned by a basis $\{A_1, A_2\}$, satisfying $[A_1, A_2] = A_1$.

We will take A_1 in one of the four possible forms obtained in Theorem 4, and let $A_2 = Z_h + X_f + Y_g + W_k$ be a general element of the 2D-CNLS algebra. We first impose the commutation relations, then simplify A_2 using the stabilizer (isotropy) group of A_1 in the adjoint group of 2D-CNLS algebra.

4.2.1 Abelian algebras

(1.1) $A_1 = Z_1$. Requiring $[A_1, A_2] = 0$ and using Table 1 we find $\dot{h} = \dot{f} = \dot{g} = \dot{k} = 0$. Hence $A_2 = aZ_1 + bX_1 + cY_1 + dW_1$. Replacing A_2 by $A'_2 = A_2 - aA_1$ we effectively set $a = 0$ in A_2 . Subsequently using $\exp(\lambda \cdot ad(W_t))$, $\exp(\delta \cdot ad(X_t))$ and $\exp(\sigma \cdot ad(Y_t))$ on $A_1 = Z_1$ and $A_2 = bX_1 + cY_1 + dW_1$, we normalize A_2 and keep the invariance of A_1 .

After applying these groups, we have

$$\begin{aligned} A_1 &\sim Z_1 + \delta X_1 + \sigma Y_1 + \left(\lambda - \frac{1}{4q}(\delta^2\epsilon - \sigma^2)\right)W_1, \\ A_2 &\sim bX_1 + cY_1 + \left(d - \frac{1}{2q}(b\delta\epsilon - c\sigma)\right)W_1. \end{aligned} \quad (22)$$

Let

$$\lambda - \frac{1}{4q}(\delta^2\epsilon - \sigma^2) = 0, \quad d - \frac{1}{2q}(b\delta\epsilon - c\sigma) = 0, \quad \delta = Ab, \quad \sigma = Ac, \quad (23)$$

for a nonzero constant A .

Suppose $d \neq 0$, then it implies $c^2 \neq \epsilon b^2$. Hence system (23) has solution $\lambda = \frac{1}{2}dA, \delta = Ab, \sigma = Ac$ with $A = 2dq/(\epsilon b^2 - c^2)$. Further we suppose $b \neq 0$, then we get equivalent algebras $\langle A_1, A_2 \rangle \sim \langle Z_1 + A(bX_1 + cY_1), bX_1 + cY_1 \rangle$ from (22). By scaling, we can take $A_1 = Z_1$ and $A_2 = X_1 + \alpha Y_1$. Therefore, we have $\mathfrak{L}_{2,1} = \langle Z_1, X_1 + \alpha Y_1 \rangle$ for arbitrary constant α . Similarly, if $b = 0, c \neq 0$, then we get $\mathfrak{L}_{2,2} = \langle Z_1, Y_1 \rangle$. If $b = c = 0$, we get $\mathfrak{L}_{2,3} = \langle Z_1, W_1 \rangle$.

In the case where $d = 0$, the two-dimensional subalgebra belongs to $\mathfrak{L}_{2,1}$ or $\mathfrak{L}_{2,2}$.

$$(1.2) A_1 = X_1.$$

We again take $A_2 = Z_h + X_f + Y_g + W_k$. The condition $[A_1, A_2] = 0$ implies $\dot{f} = \dot{h} = 0$ and $A_2 = aZ_1 + bX_1 + Y_g + W_k$. We eliminate b by linear combination with A_1 . If $a \neq 0$, conjugating by $\exp(\lambda ad(Y_G) + \nu ad(W_k))$ and appropriately choosing the F, G, K, δ, ν and λ like doing in Theorem 2, we can arrive at $A_2 = Z_1$. This can be merged into the subalgebra $\mathfrak{L}_{2,1}$ by setting $\alpha = 0$. If $a = 0$ and $g \neq 0$, conjugating A_2 by $\exp(\delta \cdot ad(Y_G))$ and selecting G satisfying $(G\dot{g} - \dot{G}g)/(2q) = k$, we can transform $k \rightarrow 0$. Hence we have $A_2 = Y_g$ for arbitrary function $g = g(t) \neq 0$. If $g = 0$, we have $A_2 = W_k$ for arbitrary function k of t . Hence we get essential subalgebras $\mathfrak{L}_{2,4} = \langle X_1, Y_\psi \rangle$ and $\mathfrak{L}_{2,5} = \langle X_1, W_\omega \rangle$ for arbitrary functions $\psi \neq 0$ and $\omega \neq 0$.

$$(1.3) A_1 = X_\phi + Y_1 \text{ for an arbitrary differential function } \phi = \phi(t).$$

Let $A_2 = Z_h + X_f + Y_g + W_k$. The condition $[A_1, A_2] = 0$ implies two kinds of conditions on f, h, g and ϕ :

Condition 1: $h = 0, \dot{g} = \epsilon(\phi\dot{f} - \dot{\phi}f).$

Condition 2: $\dot{\phi} = 0, h = b \neq 0, \dot{g} = \epsilon\phi\dot{f}.$

For the first case, conjugating A_1 and A_2 by $\exp(\lambda \cdot ad(X_F) + \delta \cdot ad(Y_G))$, one obtains

$$\begin{aligned} A_1 &\sim A_1 + W_{\lambda\bar{F}_0 - \delta\bar{G}_0}, \text{ with } \bar{G}_0 = -\frac{1}{2q}\dot{G}, \bar{F}_0 = \frac{1}{2p}(F\dot{\phi} - \dot{F}\phi), \\ A_2 &\sim X_f + Y_g + W_{k + \lambda\bar{F} - \delta\bar{G}}, \text{ with } \bar{G} = \frac{1}{2q}(G\dot{g} - \dot{G}g), \bar{F} = \frac{1}{2p}(F\dot{f} - \dot{F}f). \end{aligned} \quad (24)$$

Setting

$$\lambda\bar{F}_0 - \delta\bar{G}_0 = 0, \quad k + \lambda\bar{F} - \delta\bar{G} = 0. \quad (25)$$

and after tedious calculation, we have the following conclusions about the solutions to system (25).

Case 1: $\dot{g} \neq 0$.

The system (25) has infinite number of solutions F and G satisfying

$$\begin{aligned} G &= -\frac{\lambda}{\delta}\epsilon \left((f - \phi g)\dot{F} - (\dot{f} - \dot{\phi}g)F \right) / \dot{g}, \\ \dot{g}\ddot{F} - \ddot{g}\dot{F} + \epsilon \left((\dot{\phi}\ddot{f} - \dot{f}\ddot{\phi}) \right) F &= 0, \end{aligned}$$

for $f \neq \phi g$ and

$$G = \frac{\alpha\lambda}{\delta}\epsilon F(t),$$

for $f = \phi g$ with $\phi = \alpha$ being a nonzero constant. Taking a solution F and G of the equations as parameters in the above conjugating action (24), we can choose $A_2 = X_f + Y_g$ for ϕ, f and g satisfying Condition 1. Thus, we have algebra $\mathfrak{L}_{2,6} = \langle X_\phi + Y_1, X_\psi + Y_\Phi \rangle$. For linear independence of $X_\phi + Y_1$ and $X_\psi + Y_\Phi$, we need additional condition: Φ is nonconstant or ϕ and ψ are linear independent.

Case 2: $\dot{g} = 0$.

By the same procedure given above, we can prove that $\langle A_1, A_2 \rangle \sim \langle X_\phi + Y_1, \alpha X_\phi + \beta Y_1 \rangle$ for arbitrary functions ϕ and arbitrary constants α and β with $\alpha \neq \beta$. It is obvious that $\langle X_\phi + Y_1, \alpha X_\phi + \beta Y_1 \rangle = \langle X_\phi, Y_1 \rangle$. While, the last one is equivalent to algebra $\mathfrak{L}_{2,4}$ under adjoint action of $\exp(\delta Z_H)$ for properly selection of parameter H (also see b in Theorem 3). Hence, in this case no new algebra is arisen.

Turning to Condition 2, we can choose $A_2 = bZ_1 + X_f + Y_{af+c}$ for $\phi = \epsilon a, h = b, g = af + c$ where a, b and c are arbitrary constants. Now $A_1 = Y_1 + \epsilon a X_1$.

By Theorem 2, we can take $A_2 = Z_1$. This does not lead to new algebra since the algebra belongs to $\mathfrak{L}_{2,1}$ if $a \neq 0$ or $\mathfrak{L}_{2,2}$ if $a = 0$.

(1.4) $A_1 = W_\phi$ for an arbitrary nonzero differential function $\phi = \phi(t)$.

In this case, for $A_2 = Z_h + X_f + Y_g + W_k$, the Lie production $[A_1, A_2] = 0$ implies $h\dot{\phi} = 0$. Then two cases are arisen.

Case 1: $\phi = a$ is a nonzero constant and $h \neq 0$.

Case 2: $h = 0$ and ϕ is nonzero arbitrary function of t .

For the first case, we let $A_1 = W_1$. The invariance of W_k under adjoint actions of 2D-CNLS group and adjoint connection between $Z_h + X_f + Y_g + W_k$ and Z_1 proved in Theorem 2, here we can take $A_2 = Z_1$. This is the same as case $\mathfrak{L}_{2,3}$.

For the second case, The equivalences among $X_f + Y_g + W_k, X_1 + Y_{\bar{g}}$ and $X_{\bar{f}} + Y_1$ when $fg \neq 0$ proved in Theorem 2 yields algebra $\mathfrak{L}_{2,7} = \langle W_\phi, X_\psi + Y_1 \rangle$ for arbitray functions ϕ and ψ . For $g = 0, f \neq 0$, the

algebra $\langle W_\phi, X_1 \rangle$ is the same as $\mathfrak{L}_{2,5}$. For the case of $f = 0, g \neq 0$, we obtain an algebra $\langle W_\phi, Y_1 \rangle$ which can be merged into $\mathfrak{L}_{2,7}$ allowing $\psi = 0$.

Obviously, there is an additional algebra $\mathfrak{L}_{2,8} = \langle W_\phi, W_\psi \rangle$ for $h = f = g = 0$ and arbitrary function $\phi\psi \neq 0$ and $\phi/\psi \neq \text{constant}$.

4.2.2 Non- Abelian algebras

$$(2.1)A_1 = Z_1.$$

We take $A_2 = Z_h + X_f + Y_g + W_k$. Requiring $[A_1, A_2] = A_1$ and using in Table 1 we find $\dot{h} = 1, \dot{f} = \dot{g} = \dot{k} = 0$. Hence $A_2 = Z_t + bX_1 + cY_1 + dW_1$. Conjugating by $\exp(2c \cdot ad(Y_1)), \exp(2b \cdot ad(X_1))$ and $\exp(ad(W_{-d \ln t}))$, we can arrive at $A_2 = Z_t$ and algebra $\mathfrak{L}_{2,9} = \langle Z_1, Z_t \rangle$.

$$(2.2)A_1 = X_1.$$

We take $A_2 = Z_h + X_f + Y_g + W_k$. Requiring $[A_1, A_2] = A_1$ and using in Table 1 we find $\dot{h} = 2, \dot{f} = 0$. Hence $A_2 = Z_{2t+a} + Y_g + W_k$. Conjugating by $\exp(ad(Y_G))$ and $\exp(ad(W_k))$ and appropriately choosing the G, K like doing in Theorem 2, we can arrive at $A_2 = Z_{2t+a}$. Then using Theorem 1 with $H = 1/2$, one equivalently map Z_{2t+a} to $Z_{t'}$. Hence, we have algebra $\mathfrak{L}_{2,10} = \langle X_1, Z_t \rangle$ after deleting the prime.

$$(2.3)A_1 = Y_1 + X_\phi \text{ for an arbitrary differential function } \phi = \phi(t).$$

Let $A_2 = Z_h + X_f + Y_g + W_k$. Then, the condition $[A_1, A_2] = A_1$ implies conditions $h = 2t + a, \phi = b, g = \epsilon bf + c$ for arbitrary constants a, b and c .

Choosing a nontrivial solutions F, G and K to the following equations

$$\begin{aligned} 4p\lambda\hat{F} - f &= 0, \quad 4q\delta\hat{G} - g = 0, \\ \delta^2(G\dot{\hat{G}} - \dot{\hat{G}}\hat{G}) - \lambda^2(F\dot{\hat{F}} - \dot{\hat{F}}\hat{F}) + \lambda\bar{F} - \delta\bar{G} + \nu(2t+a)\dot{K} &= 0, \end{aligned}$$

with $\hat{F} = \frac{1}{4p}(F - (2t+a)\dot{F}), \hat{G} = \frac{1}{4q}(G - (2t+a)\dot{G}), \bar{F} = \frac{1}{2p}(F\dot{f} - \dot{F}f), \bar{G} = \frac{1}{2q}(G\dot{g} - \dot{G}g)$, and conjugating A_2 by $\exp(\delta \cdot ad(Y_G)), \exp(\lambda \cdot ad(X_F))$ and $\exp(\nu \cdot ad(W_K))$ successively we obtain $A_2 = Z_{2t+a}$. It can be proved that the solutions F and G to above equations satisfy $\delta\dot{\hat{G}} = \epsilon b\lambda\dot{\hat{F}}$ which guarantees the invariance of A_1 . Then using the symmetry transformation in Theorem 1 with $H = 1/2$, we equivalently transform Z_{2t+a} into Z_t . Hence we have subalgebra $\mathfrak{L}_{2,11} = \langle Y_1 + \alpha X_1, Z_t \rangle$ for arbitrary constant α .

(2.4) $A_1 = W_\phi$ for an arbitrary differential function $\phi = \phi(t) \neq 0$. we take $A_2 = Z_h + X_f + Y_g + W_k$. Requiring $[A_1, A_2] = A_1$ yields $\phi + \dot{\phi}h = 0$ with $h \neq 0$. By theorem 2, we transform Z_h into Z_1 and $\phi + \dot{\phi} = 0$ in coordinates with prime. Hence we have new algebra $\mathfrak{L}_{2,12} = \langle W_{e^{-t}}, Z_1 \rangle$ after delating the prime.

Summarizing above results, we have the following theorem.

Theorem 5 *Every two- dimensional subalgebra of 2D-CNLS algebra is conjugate under the adjoint group of the 2D-CNLS group to precisely one of the following algebras.*

1. *Abelian algebras:*

$$\mathfrak{L}_{2,1} = \langle Z_1, X_1 + \alpha Y_1 \rangle, \quad \forall \alpha \in \mathbb{R}.$$

$$\mathfrak{L}_{2,2} = \langle Z_1, Y_1 \rangle.$$

$$\mathfrak{L}_{2,3} = \langle Z_1, W_1 \rangle.$$

$$\mathfrak{L}_{2,4} = \langle X_1, Y_\psi \rangle, \quad \forall \psi \in C^\infty(R) \setminus \{0\}.$$

$$\mathfrak{L}_{2,5} = \langle X_1, W_\omega \rangle, \quad \forall \omega \in C^\infty(R) \setminus \{0\}.$$

$$\mathfrak{L}_{2,6} = \langle Y_1 + X_\phi, X_\psi + Y_\Phi \rangle, \text{ either } \Phi \text{ or } \phi/\psi \neq \text{constant with } \psi^2 + \Phi^2 \neq 0 \text{ and } \dot{\Phi} = \epsilon(\phi\dot{\psi} - \dot{\phi}\psi).$$

$$\mathfrak{L}_{2,7} = \langle W_\phi, X_\psi + Y_1 \rangle, \quad \forall \phi, \psi \in C^\infty(R), \phi^2 + \psi^2 \neq 0.$$

$\mathfrak{L}_{2,8} = \langle W_\phi, W_\psi \rangle, \forall \phi, \psi \in C^\infty(\mathbb{R})$ with $\phi/\psi \neq \text{constant}$.

2. *Non-Abelian algebras satisfying $[A_1, A_2] = A_1$:*

$\mathfrak{L}_{2,9} = \langle Z_1, Z_t \rangle$.

$\mathfrak{L}_{2,10} = \langle X_1, Z_t \rangle$.

$\mathfrak{L}_{2,11} = \langle Y_1 + \alpha X_1, Z_t \rangle, \forall \alpha \in \mathbb{R}$.

$\mathfrak{L}_{2,12} = \langle W_{e^{-t}}, Z_1 \rangle$.

4.3 Three-dimensional subalgebras

A real three-dimensional Lie algebra can be either simple or solvable. We will consider these two cases separately.

1. Simple Lie algebras

By Lemma 3, this algebra has a two-dimensional non-Abelian subalgebra with commutation relations (18). We will identify $\{A_1, A_2\}$ with one of the algebras in $\mathfrak{L}_{2,9} \sim \mathfrak{L}_{2,12}$, i.e., consider it to be already in standard form.

Let us start with

$\mathfrak{L}_{2,9} : A_1 = Z_1, A_2 = Z_t, A_3 = Z_h + X_f + Y_g + W_k$.

Imposing (18) we find $A_3 = Z_{t^2}$.

Then consider

$\mathfrak{L}_{2,10} : A_1 = X_1, A_2 = Z_t, A_3 = Z_h + X_f + Y_g + W_k$;

$\mathfrak{L}_{2,11} : A_1 = Y_1 + \alpha X_1, A_2 = Z_t, A_3 = Z_h + X_f + Y_g + W_k$;

$\mathfrak{L}_{2,12} : A_1 = W_{e^{-t}}, A_2 = Z_1, A_3 = Z_h + X_f + Y_g + W_k$.

It is easy to see that $[A_1, A_3] = 2A_2$ cannot be satisfied for all the three cases.

We thus have reobtained algebras \mathfrak{L}_s spanned by $\{Z_1, Z_t, Z_{t^2}\}$ given in (12). This is unique three dimensional simple algebra admitted by 2D-CNLS system (1).

2. Solvable Lie algebras

A solvable three dimensional Lie algebra always have a two-dimensional Abelian ideal. We assume that the ideal spanned by $\{A_1, A_2\}$ is already in standard form $\mathfrak{L}_{2,1} \sim \mathfrak{L}_{2,8}$ and look for a third element $A_3 = Z_h + X_f + Y_g + W_k$ so that A_1, A_2 and A_3 span the representative of the class of subalgebras.

We first use the commutation relations, then simplify A_3 using the stabilizer (isotropy) group of $\{A_1, A_2\}$ in the invariance group of the 2D-CNLS equations.

Let us now realize this procedure for each two-dimensional Abelian subalgebras in $\mathfrak{L}_{2,1} \sim \mathfrak{L}_{2,8}$.

a. $\mathfrak{L}_{2,1} = \langle Z_1, X_1 + \alpha Y_1 \rangle$ for arbitrary constant α .

Imposing (19) and using the commutator in Table 1, we obtain

$$\dot{h} = a = 2d, \dot{f} = b, \dot{g} = \alpha b, \dot{k} = 0, c = 0, (\alpha^2 - \epsilon)b = 0. \quad (26)$$

Naturally, two cases are arisen.

Case $a \neq 0$. In this case, according to the conditions (26), we can set $A_3 = aZ_t + b(X_t + \alpha Y_t) + c_1 Y_1 + c_2 W_1$. Conjugating the A_3 by $\exp(\delta \cdot ad(Y_1))$ and $\exp(\mu \cdot ad(W_1))$ with selections of δ and μ satisfying $\frac{\delta}{2} - c_2 = 0$ and $c_2 - \frac{\alpha b \delta}{2q} + \mu = 0$, we can arrive $A_3 \sim Z_t + b(X_t + \alpha Y_t)$. For $b \neq 0$, then, conjugating further by $\exp(-2\alpha b \cdot ad(Y_t))$ and $\exp(2\alpha b \cdot ad(X_t))$, we have $A_3 \sim Z_{2t}$. Hence we have algebra $\mathfrak{L}_{3,1} = \langle Z_1, X_1 + \alpha Y_1, Z_{2t} \rangle$. Here

$$M = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ for arbitrary constant } \alpha.$$

Case $a = 0$. If $b = 0$, then this algebra is Abelian $\mathfrak{L}_{3,1}^0 = \langle Z_1, X_1 + \alpha Y_1, \beta X_1 + \gamma Y_1 + \mu W_1 \rangle$ for arbitrary constants α, β, γ , and μ . If $b \neq 0$ and after scaling, we set $b = 1$ and $A_3 = X_t + \alpha Y_t + c_1 Y_1 + c_2 W_1$ for arbitrary constants c_1, c_2 and $\alpha^2 = \epsilon$. Conjugating the A_3 through $\exp(\delta \cdot ad(Y_1))$ by selecting $\delta = 2qc_2/\alpha$, we can remove the term of W_1 from A_3 . Hence we have Lie algebra $\mathfrak{L}_{3,2} = \langle Z_1, X_1 + \alpha Y_1, X_t + \alpha Y_t + \beta Y_1 \rangle$ for α satisfying $\alpha^2 = \epsilon$ and arbitrary constant β . Here $M = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.

b. $\mathfrak{L}_{2,2} = \langle Z_1, Y_1 \rangle$.

Imposing (19) and using the commutators in Table 1, we obtain

$$b = c = 0, \dot{h} = a = 2d, \dot{f} = \dot{g} = \dot{k} = 0. \quad (27)$$

Hence we can write $A_3 = aZ_t + c_1X_1 + c_2W_1$ for arbitrary constants c_1 and c_2 . If $a = 0$, then this is an Abelian algebra $\mathfrak{L}_{3,2}^0 = \langle Z_1, Y_1, \alpha X_1 + \beta W_1 \rangle$ for arbitrary not all zero constants α and β . If $a \neq 0$, scaling and conjugating through $\exp(\lambda \cdot ad(X_1))$ with $\lambda = 2c_1/a$, we have $A_3 \sim Z_t + c_2W_1$ for arbitrary constant c_2 . Hence we have Lie algebra $\mathfrak{L}_{3,3} = \langle Z_1, Y_1, Z_{2t} + \alpha W_1 \rangle$ for arbitrary constant α and corresponding matrix

$$M = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

c. $\mathfrak{L}_{2,3} = \langle Z_1, W_1 \rangle$.

Imposing (19) and using the commutators in Table 1, we obtain

$$d = c = 0, \dot{h} = a, \dot{f} = \dot{g} = 0, \dot{k} = b. \quad (28)$$

Hence we can write $A_3 = aZ_t + c_1X_1 + c_2Y_1 + bW_t$ for arbitrary constants c_1 and c_2 . Similar to the above cases, if $a = 0$, we have an Abelian algebra $\mathfrak{L}_{3,3}^0 = \langle Z_1, W_1, \alpha X_1 + \beta Y_1 \rangle$ for arbitrary not all zero constants α and β . If $a \neq 0$, applying $\exp(\delta \cdot ad(Y_1))$ and $\exp(\lambda \cdot ad(X_1))$ on the A_3 for suitable selections of δ and λ , we can remove the X_1 and Y_1 terms from the A_3 and have $A_3 \sim Z_t + \alpha W_t$ for arbitrary constant α . Further, conjugating by $\exp(-\alpha W_t)$, we make $A_3 \sim Z_t$; If $a = 0$, then either $c_1 \neq 0$ or $c_2 \neq 0$ including $b = 0$, we always make $A_3 \sim c_1X_1 + c_2Y_1$ by applying $\exp(\lambda \cdot ad(Y_{t^2}))$ or $\exp(\lambda \cdot ad(X_{t^2}))$. This leads to Abelian algebras $\mathfrak{L}_{3,3}^0$. Hence we only left the case $c_2 = c_3 = 0$ and $b \neq 0$. In this case $A_3 \sim W_t$.

Consequently, we have algebras $\mathfrak{L}_{3,4} = \langle Z_1, W_1, Z_t \rangle$ and $\mathfrak{L}_{3,5} = \langle Z_1, W_1, W_t \rangle$ with corresponding matrix $M = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $M = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, respectively.

d. $\mathfrak{L}_{2,4} = \langle X_1, Y_\psi \rangle$ for arbitrary function ψ of t .

Imposing (19) and using the commutators in Table 1, we obtain

$$b = c = 0, \dot{h} = 2a, \dot{f} = 0, (a - d)\psi = h\dot{\psi}, \psi\dot{g} - \dot{\psi}g = 0. \quad (29)$$

The last equation implies $g = \tilde{c}\psi$ for arbitrary constant \tilde{c} . Hence we can write $A_3 = Z_h + W_k$.

If $h \neq 0$, conjugating through $\exp(ad(W_K))$, we can remove W_k term from A_3 by selecting K satisfying $h\dot{K} + k = 0$. Hence $A_3 \sim Z_h$. From (29), we have $\psi = \tilde{c}h^{\frac{a-d}{2a}}$ for $a \neq 0$ and $\psi = \tilde{c}\exp(-\frac{d}{h})t$ for $a = 0$. Here \tilde{c} denotes an arbitrary constant. For the both cases of ψ , we use Theorem 3 with translating and scaling variable t , we get algebras $\mathfrak{L}_{3,6} = \langle X_1, Y_{(2t)^\alpha}, Z_{2t} \rangle$ for arbitrary constant α and $\mathfrak{L}_{3,7} = \langle X_1, Y_{e^{-t}}, Z_1 \rangle$.

They have matrices $M = \begin{bmatrix} 1 & 0 \\ 0 & 1 - \alpha \end{bmatrix}$ and $M = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$, respectively.

If $h = 0$, we get Abelian algebra $\mathfrak{L}_{3,4}^0 = \langle X_1, Y_\psi, W_\omega \rangle$ for arbitrary nonzero functions ψ and ω .

e. $\mathfrak{L}_{2,5} = \langle X_1, W_\omega \rangle$ for arbitrary function ω of t .

Imposing (19) and using the commutators in Table 1, we obtain

$$c = 0, \dot{h} = 2a, \dot{f} = -2pb\omega, h\dot{\omega} = -d\omega. \quad (30)$$

For simplification of A_3 , three cases are considered.

Case $\dot{h} \neq 0$.

Applying $\exp(\delta \cdot ad(Y_G))$ and letting G be any solution of equation $g - \delta(\frac{1}{2}G\dot{h} - \dot{G}h) = 0$, we remove the Y_g term from A_3 , i.e., $A_3 \sim A'_3 = Z_h + X_f + W_{\tilde{k}}$ for some function \tilde{k} depending on G and k . Now applying $\exp(\lambda \cdot ad(X_F))$ on A'_3 and letting F satisfy

$$f - \lambda(\frac{1}{2}F\dot{h} - \dot{F}h) = 0. \quad (31)$$

Meantime, in order to keep the invariance of algebra $\mathfrak{L}_{2,5}$, we also let

$$\dot{F} = 2p\rho\omega, \quad (32)$$

for some constant ρ .

Under the condition $a \neq d$, there exist solutions F and $\rho = b/(a - d)$ to equations (31) and (32). Therefore, under $a \neq d$, we reach $A_3 \sim Z_h + W_{\hat{k}}$ for another function \hat{k} depending on G, F and ρ . Finally applying $\exp(\mu \cdot ad(W_K))$, we can remove the W term from A_3 by requiring $h\dot{K} + \hat{k} = 0$, i.e., $A_3 \sim Z_h$.

Solving equations (30), we have $\omega = \tilde{c}h^{-\frac{d}{2a}}$ for arbitrary constant \tilde{c} .

Consequently, after using the scaling and translation for the variable t , we obtain equivalent algebra $\mathfrak{L}_{3,8} = \langle X_1, W_{(2t)^\nu}, Z_{2t} \rangle$ with $\nu \neq -\frac{1}{2}$. It corresponds to matrix $M = \begin{bmatrix} 1 & 0 \\ 0 & -2\nu \end{bmatrix}$.

Similarly, when $a = d$, we obtain equivalence algebra $\mathfrak{L}_{3,9} = \langle X_1, W_{1/\sqrt{2t}}, Z_{2t} + 4p\alpha X_{\sqrt{2t}} \rangle$ with matrix $M = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ for arbitrary constant α .

Case $\dot{h} = 0, h \neq 0$. We obtain algebra $\mathfrak{L}_{3,10} = \langle X_1, W_{e^{-t}}, Z_1 \rangle$ which corresponds to matrix $M = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$.

Case $h = 0$. Conjugating A_3 through $\exp(\delta \cdot ad(Y_G))$, we can remove the term W_k from it. Hence we obtain algebra $\mathfrak{L}_{3,11} = \langle X_1, W_\omega, X_f + Y_\psi \rangle$ with $\dot{f} = -2p\alpha^2\omega$ and matrix $M = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ for arbitrary functions ω, ψ and constant $\alpha \neq 0$.

If $\alpha = 0$, we have algebra $\mathfrak{L}_{3,4}^0$ again.

f. $\mathfrak{L}_{2,6} = \langle Y_1 + X_\phi, X_\psi + Y_\Phi \rangle$ for arbitrary functions ϕ, ψ and Φ satisfying $\dot{\Phi} = \epsilon(\phi\dot{\psi} - \dot{\phi}\psi)$.

Let $A_1 = X_\phi + Y_1, A_2 = X_\psi + Y_\Phi$ and $A_3 = Z_h + X_f + Y_g + W_k$. Imposing (19) and noticing Abelian property of A_1 and A_2 , we obtain

$$(I) \begin{cases} \frac{1}{2}\phi\dot{h} - \dot{\phi}h = a\phi + b\psi \\ \frac{1}{2}\dot{h} = a + b\Phi \\ \dot{g} = \epsilon(\phi\dot{f} - \dot{\phi}f) \end{cases} \quad (II) \begin{cases} \frac{1}{2}\psi\dot{h} - \dot{\psi}h = c\phi + d\psi \\ \frac{1}{2}\Phi\dot{h} - h\dot{\Phi} = c + d\Phi \\ \Phi\dot{g} - \dot{\Phi}g = \epsilon(\psi\dot{f} - \dot{\psi}f) \end{cases} \quad (III) \quad \dot{\Phi} = \epsilon(\phi\dot{\psi} - \dot{\phi}\psi)$$

Particularly, for the linear combinations of A_1 and A_2 : $B_1 = \alpha A_1 + \beta A_2$ and $B_2 = \gamma A_1 + \sigma A_2$ with $\alpha\gamma - \beta\sigma \neq 0$, we can transform B_1 into A_1 and B_2 into A_2 in different functions ϕ, ψ and Φ by $\exp(\mu \cdot ad(Z_H))$. Thus we assume that M is already in standard form (see Lemma 3).

$$(f.1) \quad M = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Corresponding to the cases $h \neq 0$ and $h = 0$, we obtain two algebras, namely $\langle X_1, Y_1, Z_1 \rangle$ which belongs to $\mathfrak{L}_{3,2}^0$ when $\beta = 0$ and $\mathfrak{L}_{3,5}^0 = \langle X_\phi + Y_1, X_\psi + Y_\Phi, X_f + Y_g + W_k \rangle$ for arbitrary solutions ϕ, ψ, f, g to (I)₃, (II)₃ and (III) with either $\{\phi, \psi\}$ is linear independent or Φ is not constant, where k is an arbitrary function of t .

$$(f.2) \quad M = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

Solving (I)-(II) with this matrix M , we obtain two algebras $\mathfrak{L}_{3,12} = \langle \beta X_1 + Y_1, X_{\sqrt{2t}} + \epsilon\beta Y_{\sqrt{2t}}, Z_{2t} \rangle$ for $\epsilon\beta^2 \neq 1$ and $\mathfrak{L}_{3,13} = \langle \beta X_1 + Y_1, \beta X_{\sqrt{2t}} + Y_{\sqrt{2t}}, Z_{2t} + \beta X_g + Y_g \rangle$ for $\epsilon\beta^2 = 1$ and arbitrary function g of t .

$$(f.3) \quad M = \begin{bmatrix} 1 & 0 \\ 0 & \alpha \end{bmatrix}.$$

Solving (I)-(II) with this matrix M , we obtain three algebras: $\mathfrak{L}_{3,14} = \langle \beta X_1 + Y_1, X_{(2t)^{\frac{1-\alpha}{2}}} + \epsilon\beta Y_{(2t)^{\frac{1-\alpha}{2}}}, Z_{2t} \rangle$ for both $\alpha \neq 1$ and $\epsilon\beta^2 \neq 1$; $\mathfrak{L}_{3,15} = \langle \beta X_1 + Y_1, X_{(2t)^{\frac{1-\alpha}{2}}} + \epsilon\beta Y_{(2t)^{\frac{1-\alpha}{2}}}, Z_{2t} + \beta X_g + Y_g \rangle$ for $\alpha \neq 1$ and $\epsilon\beta^2 = 1$ and arbitrary function g of t ; $\mathfrak{L}_{3,16} = \langle X_1, Y_1, Z_{2t} \rangle$ for $\alpha = 1$.

$$(f.4) \quad M = \begin{bmatrix} \alpha & 1 \\ -1 & \alpha \end{bmatrix}.$$

In this case, from (I)-(II) we see that we have to $h \neq 0$. In order to solve system (I)-(III), we use a way differ from above procedure. At first, we perform a conjugating by $\exp(\mu \cdot ad(Z_H))$ changing the basis A_1, A_2 and A_3 into $B_1 = X_\phi + Y_\Psi, B_2 = X_\psi + Y_\Phi$ and $B_3 = Z_1 + X_f + Y_g + W_k$, respectively (without loss of generality, we take same notations on these group labeled functions). The conditions (19) now implies

$$(I)' \begin{cases} -\dot{\phi} = \alpha\phi + \psi, \\ -\dot{\Psi} = \alpha\Psi + \Phi, \\ \Psi\dot{g} - \dot{\Psi}g = \epsilon(\phi\dot{f} - \dot{\phi}f). \end{cases} \quad (II)' \begin{cases} -\dot{\psi} = \alpha\psi - \phi, \\ -\dot{\Phi} = \alpha\Phi - \Psi, \\ \Phi\dot{g} - \dot{\Phi}g = \epsilon(\psi\dot{f} - \dot{\psi}f). \end{cases} \quad (III)' \quad \Psi\dot{\Phi} - \dot{\Psi}\Phi = \epsilon(\phi\dot{\psi} - \dot{\phi}\psi).$$

Solving first two equations in (I)'₁₋₂ and (II)'₁₋₂ respectively, we obtain

$$\begin{aligned} \phi &= e^{-\alpha t}(c_1 \cos t + c_2 \sin t); & \psi &= e^{-\alpha t}(-c_2 \cos t + c_1 \sin t); \\ \Phi &= e^{-\alpha t}(c_3 \cos t + c_4 \sin t); & \Psi &= e^{-\alpha t}(c_4 \cos t - c_3 \sin t), \end{aligned}$$

for arbitrary constants $c_1 \sim c_4$ satisfying $\epsilon(c_1^2 + c_2^2) = (c_3^2 + c_4^2)$ from (III)', which implies that $\phi = \psi = 0$ is equivalent to $\Psi = \Phi = 0$.

Hence, we have $c_1^2 + c_2^2 \neq 0$. Under this condition, we always choice a basis of algebra $\mathfrak{L}_{2,6} = \langle A_1, A_2 \rangle$ through linear combinations of A_1 and A_2 such that $c_1 = 1$ and $c_2 = 0$, which leads to $\phi = e^{-\alpha t} \cos t, \psi = e^{-\alpha t} \sin t$ and $\Phi = \rho\phi + \beta\psi, \Psi = \beta\phi - \rho\psi$ for arbitrary constants ρ and β satisfying $\epsilon = \rho^2 + \beta^2$.

Solving (I)'₃ and (II)'₃, we obtain

$$f(t) = \delta(\rho\phi - \beta\psi) + \sigma\psi, \quad g(t) = \sigma(\rho\phi + \beta\psi) - \delta\epsilon\psi. \quad (33)$$

if $\rho \neq 0$. Here δ, σ, ρ and β are arbitrary constants with satisfying $\epsilon = \rho^2 + \beta^2$; and

$$g(t) = \beta f(t) \quad (34)$$

if $\rho = 0$. Here f is arbitrary function of t and $\epsilon = \beta^2$.

Hence we obtain algebra $\mathfrak{L}_{3,17}$ and $\mathfrak{L}_{3,18}$ spanned by $\{A_1, A_2, A_3\}$ with $A_1 = X_\phi + Y_\Psi, A_2 = X_\psi + Y_\Phi$ and $A_3 = Z_1 + X_f + X_g$ in which f and g are given by (33) and (34), respectively.

$$(f.5) \quad M = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$

In this case, we obtain two algebras $\mathfrak{L}_{3,19}$ spanned by $A_1 = \beta X_1 + Y_1, A_2 = X_{\beta t + \nu} + Y_t$ and $A_3 = -Z_1$ for arbitrary constant ν and $\mathfrak{L}_{3,20}$ spanned by $A_1 = \beta X_1 + Y_1, A_2 = \beta X_t + Y_t$ and $A_3 = -Z_1 + \beta X_g + Y_g$ for arbitrary function g of t . In both cases, β satisfies $\epsilon\beta^2 = 1$.

$$(f.6) \quad M = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

We obtain two algebras: $\mathfrak{L}_{3,21}$ spanned by $A_1 = \beta X_1 + Y_1, A_2 = X_{-\frac{\beta}{2} \ln t + \nu} + Y_{-\frac{1}{2} \ln t + \sigma}$ and $A_3 = Z_{2t}$ for arbitrary constants ν and σ ; and $\mathfrak{L}_{3,22}$ spanned by $A_1 = \beta X_1 + Y_1, A_2 = X_{-\frac{\beta}{2} \ln t} + Y_{-\frac{1}{2} \ln t}$ and $A_3 = Z_{2t} + \beta X_g + Y_g$ for arbitrary function g of t . In both cases, β satisfies $\epsilon\beta^2 = 1$.

g. $\mathfrak{L}_{2,7} = \langle W_\phi, X_\psi + Y_1 \rangle$ for arbitrary functions ϕ and ψ .

Corresponding to each case of standard matrices M , we obtain seven algebras.

(g.1) $M = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. In this case we have two algebras. Algebra $\mathfrak{L}_{3,6}^0$ is spanned by $A_1 = W_1, A_2 = \beta X_1 + Y_1$ and $A_3 = Z_1$ for arbitrary constant β ; Algebra $\mathfrak{L}_{3,7}^0$ is spanned by $A_1 = W_\phi, A_2 = X_\psi + Y_1$ and $A_3 = X_f + Y_g$ for arbitrary functions f, g, ϕ with $f^2 + g^2 \neq 0$ and satisfying $\dot{g} = \epsilon(\psi \dot{f} - \dot{\psi} f)$. If $f = g = 0$, we have an Abelian algebra $\mathfrak{L}'_{3,7} = \langle W_\phi, X_\psi + Y_1, W_k \rangle$ where arbitrary functions ϕ and k are linear independent.

(g.2) $M = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. In this case we have algebra $\mathfrak{L}_{3,23}$ spanned by $A_1 = W_{e^{-t}}, A_2 = \beta X_1 + Y_1 + \alpha W_1$ and $A_3 = Z_1$ for arbitrary constants α, β ;

(g.3) $M = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$. In this case we have two algebras. Algebra $\mathfrak{L}_{3,24}$ spanned by $A_1 = W_1, A_2 = \beta X_1 + Y_1 + W_t$ and $A_3 = Z_1$ for arbitrary constant β ; Algebra $\mathfrak{L}_{3,25}$ spanned by $A_1 = W_\phi, A_2 = X_\psi + Y_1$ and $A_3 = X_f + Y_g$ for arbitrary functions f, g, ϕ and ψ satisfying $\dot{g} = 2q\phi + \epsilon(\psi \dot{f} - f \dot{\psi})$; $\mathfrak{L}_{3,26} = \langle W_\phi, Y_1, Y_g + W_k \rangle$ for arbitrary functions g, k and $\phi \neq 0$ satisfying $\dot{g} = 2q\phi$;

(g.4) $M = \begin{bmatrix} 1 & 0 \\ 0 & d \end{bmatrix}$. In this case we have algebra $\mathfrak{L}_{3,27} = \langle W_{t^{-\nu}}, \beta X_1 + Y_1, Z_{\nu^{-1}t} \rangle$ for arbitrary constants β and $d = (2\nu)^{-1} \neq 0$.

(g.5) $M = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$. In this case we have no new algebra;

k. $\mathfrak{L}_{2,8} = \langle W_\phi, W_\psi \rangle$ for arbitrary nonzero functions ϕ and ψ with $\phi/\psi \neq \text{constant}$.

We consider two cases $\neq 0$ and $h = 0$ separately.

Case: $h \neq 0$. According to the Theorem 1 and adjoint action Table 4, we can take $A_3 = Z_h$ without changing $\langle A_1 = W_\phi, A_2 = W_\psi \rangle$. Conjugating by $\exp(\mu \cdot \text{ad}(Z_H))$ and changing the basis A_1, A_2 and A_3 into $B_1 = X_\phi + Y_\psi, B_2 = X_\psi + Y_\phi$ and $B_3 = Z_1 + X_f + Y_g + W_k$, respectively (without loss of generality, we take same notations on these group labeled functions). Thus the conditions (19) implies

$$-\dot{\phi} = a\phi + b\psi, \quad -\dot{\psi} = c\psi + d\phi. \quad (35)$$

Corresponding to each case of standard matrices M , we solve equations (35) and obtain following algebras.

(k.1) $M = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. Algebra $\mathfrak{L}_{3,28} = \langle W_{e^{-t}}, W_1, Z_1 \rangle$.

(k.2) $M = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$. Algebra $\mathfrak{L}_{3,29} = \langle W_1, W_{-t}, Z_1 \rangle$.

$$(k.3) \quad M = \begin{bmatrix} 1 & 0 \\ 0 & d \end{bmatrix}. \text{ Algebra } \mathfrak{L}_{3,30} = \langle W_{e^{-t}}, W_{e^{-dt}}, Z_1 \rangle, \forall d \in \mathbb{R} \setminus \{0\}.$$

$$(k.4) \quad M = \begin{bmatrix} a & 1 \\ -1 & a \end{bmatrix}. \text{ Algebra } \mathfrak{L}_{3,31} = \langle W_{e^{-at \cos t}}, W_{e^{-at \sin t}}, Z_1 \rangle, \forall a \in \mathbb{R}.$$

$$(k.5) \quad M = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}. \text{ Algebra } \mathfrak{L}_{3,32} = \langle W_{e^t}, W_{te^t}, Z_{-1} \rangle.$$

Case: $h = 0$. In this case, we have only Abelian algebra $\mathfrak{L}_{3,8}^0 = \langle W_\phi, W_\psi, X_f + Y_g \rangle$ for arbitrary functions ϕ, ψ, f and g with satisfying $\phi/\psi \neq \text{constant}$ and $f^2 + g^2 \neq 0$. Here we notice that algebra $\mathfrak{L}_{3,7}'$ is contained in algebra $\mathfrak{L}_{3,8}^0$.

Remark 2: The algebras $\mathfrak{L}_{3,2}, \mathfrak{L}_{3,13} - \mathfrak{L}_{3,22}$ only exist for $\epsilon > 0$. This shows the effect of the physical constants p and q on the extending of symmetries of system (1) when they have the same sign.

In summarizing this section, we present a list of representatives of all equivalent classes of three dimensional subalgebras of 2D-CNLS algebra by specifying the matrix M in (20). The solvable algebras are given in order $\{A_1, A_2, A_3\}$ and $\{A_1, A_2\}$ is Abelian ideal and the action of A_3 is given in (19).

Theorem 6 *Every three- dimensional subalgebra of 2D-CNLS algebra is conjugate under the adjoint group of the 2D-CNLS group to precisely one of the following algebras of lists 1-7.*

1. Abelian algebras with $M = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$:

$$\mathfrak{L}_{3,1}^0 = \langle Z_1, X_1 + \alpha Y_1, \beta X_1 + \gamma Y_1 + \mu W_1 \rangle, \forall \alpha, \beta, \gamma, \mu \in \mathbb{R}.$$

$$\mathfrak{L}_{3,2}^0 = \langle Z_1, Y_1, \alpha X_1 + \beta W_1 \rangle, \forall \alpha, \beta \in \mathbb{R}, \alpha^2 + \beta^2 \neq 0.$$

$$\mathfrak{L}_{3,3}^0 = \langle Z_1, W_1, \alpha X_1 + \beta Y_1 \rangle, \forall \alpha, \beta \in \mathbb{R}, \alpha^2 + \beta^2 \neq 0.$$

$$\mathfrak{L}_{3,4}^0 = \langle X_1, Y_\psi, W_\omega \rangle, \forall \psi, \omega \in C^\infty(\mathbb{R}).$$

$$\mathfrak{L}_{3,5}^0 = \langle X_\phi + Y_1, X_\psi + Y_\Phi, X_f + Y_g + W_k \rangle, \forall \text{ solutions } \phi, \psi, \Phi, f, g \in C^\infty(\mathbb{R}) \text{ to (I)}_3, \text{ (II)}_3 \text{ and (III)}$$

with either $\{\phi, \psi\}$ is linear independent or Φ is not constant.

$$\mathfrak{L}_{3,6}^0 = \langle W_1, \beta X_1 + Y_1, Z_1 \rangle, \forall \beta \in \mathbb{R}.$$

$$\mathfrak{L}_{3,7}^0 = \langle W_\phi, X_\psi + Y_1, X_f + Y_g \rangle, \forall \phi, \psi, f, g \in C^\infty(\mathbb{R}) \text{ satisfying } f^2 + g^2 \neq 0 \text{ and } \dot{g} = \epsilon(\psi \dot{f} - \dot{\psi} f).$$

$$\mathfrak{L}_{3,8}^0 = \langle W_\phi, W_\psi, X_f + Y_g \rangle, \forall \phi, \psi, f, g \in C^\infty(\mathbb{R}) \text{ with } \phi/\psi \neq \text{constant and } f^2 + g^2 \neq 0.$$

2. Decomposable, non-Abelian algebras, with $M = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$:

$$\mathfrak{L}_{3,4} = \langle Z_1, W_1, Z_t \rangle.$$

$$\mathfrak{L}_{3,7} = \langle Y_{e^{-t}}, X_1, Z_1 \rangle.$$

$$\mathfrak{L}_{3,10} = \langle W_{e^{-t}}, X_1, Z_1 \rangle.$$

$$\mathfrak{L}_{3,12} = \langle \beta X_1 + Y_1, X_{\sqrt{2t}} + \epsilon \beta Y_{\sqrt{2t}}, Z_{2t} \rangle, \text{ for } \epsilon \beta^2 \neq 1.$$

$$\mathfrak{L}_{3,13} = \langle \beta X_1 + Y_1, \beta X_{\sqrt{2t}} + Y_{\sqrt{2t}}, Z_{2t} + \beta X_g + Y_g \rangle, \text{ for } \epsilon \beta^2 = 1, \forall g \in C^\infty(\mathbb{R}).$$

$$\mathfrak{L}_{3,23} = \langle W_{e^{-t}}, \beta X_1 + Y_1 + \alpha W_1, Z_1 \rangle, \forall \alpha, \beta \in \mathbb{R}.$$

$$\mathfrak{L}_{3,28} = \langle W_{e^{-t}}, W_1, Z_1 \rangle.$$

3. Nilpotent algebras with $M = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$:

$$\mathfrak{L}_{3,2} = \langle X_1 + \alpha Y_1, Z_1, X_t + \alpha Y_t + \beta Y_1 \rangle, \alpha^2 = \epsilon, \forall \beta \in \mathbb{R}.$$

$$\mathfrak{L}_{3,5} = \langle W_1, Z_1, W_t \rangle.$$

$$\mathfrak{L}_{3,11} = \langle W_\omega, X_1, X_f + Y_\psi \rangle, \dot{f} = -2p\alpha^2\omega, \forall \omega, \psi \in C^\infty(\mathbb{R}), \alpha \in \mathbb{R} \setminus \{0\}.$$

$$\mathfrak{L}_{3,19} = \langle \beta X_1 + Y_1, X_{\beta t + \nu} + Y_t, A_3 = -Z_1 \rangle, \forall \nu \in \mathbb{R} \text{ and } \epsilon \beta^2 = 1.$$

$$\mathfrak{L}_{3,20} = \langle \beta X_1 + Y_1, \beta X_t + Y_t, -Z_1 + \beta X_g + Y_g \rangle, \forall g \in C^\infty(\mathbb{R}) \text{ and } \epsilon \beta^2 = 1.$$

$$\mathfrak{L}_{3,24} = \langle W_1, A_2 = \beta X_1 + Y_1 + W_t, Z_1 \rangle, \forall \beta \in \mathbb{R}.$$

$$\mathfrak{L}_{3,25} = \langle W_\phi, X_\psi + Y_1, X_f + Y_g \rangle, \forall f, g, \phi, \psi \in C^\infty(\mathbb{R}) \text{ satisfying } \dot{g} = 2\alpha q\phi + \epsilon(\psi \dot{f} - f \dot{\psi}), \alpha \in \mathbb{R} \setminus \{0\}.$$

$$\mathfrak{L}_{3,26} = \langle W_\phi, Y_1, Y_g + W_k \rangle, \forall g, k, \phi \in C^\infty(\mathbb{R}) \text{ satisfying } \dot{g} = 2q\phi.$$

$$\mathfrak{L}_{3,29} = \langle W_1, W_{-t}, Z_1 \rangle.$$

4. Diagonal action on ideal with $M = \begin{bmatrix} 1 & 0 \\ 0 & a \end{bmatrix}, a \neq 0$:

$$\mathfrak{L}_{3,1} = \langle Z_1, X_1 + \alpha Y_1, Z_{2t} \rangle, a = 1, \forall \alpha \in \mathbb{R}.$$

$$\mathfrak{L}_{3,3} = \langle Z_1, Y_1, Z_{2t} + \alpha W_1 \rangle, \forall \alpha \in \mathbb{R}, a = 1.$$

$$\mathfrak{L}_{3,6} = \langle X_1, Y_{(2t)^\alpha}, Z_{2t} \rangle, \forall \alpha \in \mathbb{R}, a = 1 - \alpha.$$

$$\mathfrak{L}_{3,8} = \langle X_1, W_{(2t)^\nu}, Z_{2t} \rangle, \nu \neq -\frac{1}{2}, a = -2\nu,$$

$$\mathfrak{L}_{3,9} = \langle X_1, W_{1/\sqrt{2t}}, Z_{2t} + 4p\alpha X_{\sqrt{2t}} \rangle, \forall \alpha \in \mathbb{R}, a = 1.$$

$$\mathfrak{L}_{3,14} = \langle \beta X_1 + Y_1, X_{(2t)^{\frac{1-a}{2}}} + \epsilon \beta Y_{(2t)^{\frac{1-a}{2}}}, Z_{2t} \rangle, \forall a \neq 1, \epsilon \beta^2 \neq 1.$$

$$\mathfrak{L}_{3,15} = \langle \beta X_1 + Y_1, X_{(2t)^{\frac{1-a}{2}}} + \epsilon \beta Y_{(2t)^{\frac{1-a}{2}}}, Z_{2t} + \beta X_g + Y_g \rangle, \forall a \neq 1, \epsilon \beta^2 = 1, \forall g \in C^\infty(\mathbb{R}).$$

$$\mathfrak{L}_{3,16} = \langle X_1, Y_1, Z_{2t} \rangle, a = 1.$$

$$\mathfrak{L}_{3,27} = \langle W_{t^{-\nu}}, \beta X_1 + Y_1, Z_{\nu^{-1}t} \rangle, \forall \beta, \nu \in \mathbb{R}, a = (2\nu)^{-1} \neq 0.$$

$$\mathfrak{L}_{3,30} = \langle W_{e^{-t}}, W_{e^{-at}}, Z_1 \rangle, \forall a = d \in \mathbb{R}.$$

5. Complex action on ideal with $M = \begin{bmatrix} a & 1 \\ -1 & a \end{bmatrix}, \forall a \in \mathbb{R}$:

$\mathfrak{L}_{3,17} = \{A_1, A_2, A_3\}$, where $A_1 = X_\phi + Y_\Psi$, $A_2 = X_\psi + Y_\Phi$ and $A_3 = Z_1 + X_f + X_g$; $\phi = e^{-at} \cos t$, $\psi = e^{-at} \sin t$, $\Phi = \rho\phi + \beta\psi$, $\Psi = \beta\phi - \rho\psi$ and $\{f, g\}$ are given by (33) and $\epsilon = \rho^2 + \beta^2, \forall \delta, \sigma, \beta \in \mathbb{R}$ and $\rho \in \mathbb{R} \setminus \{0\}$.

$\mathfrak{L}_{3,18} = \{A_1, A_2, A_3\}$ where $A_1 = X_\phi + Y_\Psi$, $A_2 = X_\psi + Y_\Phi$ and $A_3 = Z_1 + X_f + X_g$; $\phi = e^{-at} \cos t$, $\psi = e^{-at} \sin t$, $\Phi = \beta\psi$, $\Psi = \beta\phi$ and f and g are given by (34) $\forall f \in C^\infty(\mathbb{R})$ and $\epsilon = \beta^2, \forall \beta \in \mathbb{R}$.

$$\mathfrak{L}_{3,31} = \langle W_{e^{-at} \cos t}, W_{e^{-at} \sin t}, Z_1 \rangle, \forall a \in \mathbb{R}.$$

6. Jordan action on idea with $M = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$:

$$\mathfrak{L}_{3,21} = \langle \beta X_1 + Y_1, X_{-\frac{\beta}{2} \ln t + \nu} + Y_{-\frac{1}{2} \ln t + \sigma}, Z_{2t} \rangle, \forall \nu, \sigma \in \mathbb{R} \text{ and } \epsilon \beta^2 = 1.$$

$$\mathfrak{L}_{3,22} = \langle \beta X_1 + Y_1, X_{-\frac{\beta}{2} \ln t} + Y_{-\frac{1}{2} \ln t}, Z_{2t} + \beta X_g + Y_g \rangle, \forall g \in C^\infty(\mathbb{R}) \text{ and } \epsilon \beta^2 = 1.$$

$$\mathfrak{L}_{3,32} = \langle W_{e^t}, W_{te^t}, Z_{-1} \rangle.$$

7. The simple Lie algebra $\mathfrak{sl}(2, \mathbb{R})$:

$$\mathfrak{L}_s = \langle Z_1, Z_t, Z_{t^2} \rangle [\text{see (12)}].$$

5 Conclusions

The main results of this paper are summed up in Theorems 4-6 providing representatives of the conjugacy classes of one-, two-, and three- dimensional real subalgebras of the 2D-CNLS algebra. The classification is performed under the action of 2D-CNLS group on 2D-CNLS algebra. We obtain several inequivalent classes for subalgebras with different dimensions: four classes of one-dimensional subalgebras, twelve classes of two- dimensional subalgebras and thirty three classes of three- dimensional subalgebras. The classification shows the finite dimensional algebra structure of infinite dimensional Lie algebra of the 2D-CNLS system (1). It will provide the way to know the infinite number of the invariant solutions of the 2D-CNLS equations

under the 2D-CNLS group with finite solutions of the same equations. To construct invariant solutions to system (1), two equivalent approaches can be adopted. One is to perform the symmetry reduction using a general element V expressed by (5). Another procedure is to make use of the classification of subalgebras of the 2D-CNLS algebra, shown in previous sections. In both cases, we have standard procedure to get the invariant solutions [16,24]. In the second method, we first go through the procedure using the representatives of each conjugacy classes and then use the transformations (9) and (10) to produce the solutions belong to the equivalent classes of the representatives. The number of the representatives shows the finite number of inequivalent classes of the invariant solutions of the 2D-CNLS system (1). Moreover, we have noted that there are four arbitrary functions in the obtained solutions. This may be useful to solve initial or initial-boundary problems of the system (1) by the renormal group methods [31]. Hence, the state is now set for performing the actual symmetry reduction of the 2D-CNLS system (1) and obtaining *all* solutions of the 2D-CNLS equations that are invariant under the action of 2D-CNLS subgroups. For this rich topic, due to the lack of space and different focus, we will discuss it in detail in another article.

Acknowledgments This work is supported by National Natural Science Foundation of China (No. 11571008).

References

- [1] Wu-Ming Liu, Schrödinger Equations in Nonlinear Systems, Spring-Verlag: Singapore, 2019.
- [2] D.J. Benney and G.J. Roskes, Wave Instabilities, Stud. Appl. Math. **48** (1969) 377-385.
- [3] A. Davey and K. Stewartson, On Three-Dimensional Packets of Surface Waves, Proc. Roy. Soc. London. A338 (1974) 101-109.
- [4] V. D. Djordjevic and L. G. Redekopp, On two-dimensional packets of capillary-gravity waves, J. Fluid Mech. Great Britain. **79** (1977) 703-714.
- [5] C. N. Freeman and A. Davey, On the evolution of packets of long surface waves, Proc. R. Soc. London. A344 (1975) 427-433.
- [6] M. J. Ablowitz and H. Segur, On the evolution of packets of water waves, J. Fluid Mech. Great Britain. **92** (1979) 691-715.
- [7] M. J. Ablowitz and R. Haberman, Nonlinear Evolution Equations- Two and Three Dimensions, Phys. Rev. Lett. **35** (1975) 1185-1188.
- [8] D. Anker and N. C. Freeman, On the soliton solutions of the Davey- Stewartson equations for long waves, Proc. R. Soc. London. A360 (1978) 529-540.
- [9] J. Satsuma and M. J. Ablowitz, Two-dimensional lumps in nonlinear dispersive systems, J. Math. Phys. **20** (1979) 1496-1503.
- [10] A. Nakamura, Explode-decay mode lump solutions of a two-dimensional nonlinear Schrödinger equation, Phys. Lett. **88A** (1982) 55-56.

- [11] M. Tajiri and M. Hagiwar, Similarity solutions of the Two-Dimensional Coupled Nonlinear Schrödinger Equation, *J. Phys. Soci. Jap.* Vol.52. No.11(1983) 3727-3734.
- [12] A. Li, Chaolu Temuer, Lie Symmetries, One-Dimensional Optimal System and Optimal Reduction of (2 + 1)-Coupled nonlinear Schrödinger Equations, *Journal of Applied Mathematics and Physics*, Vol.2 No.7, (2014) 667-6690.
- [13] L. V. Ovsyannikov, Lectures on the Theory of Group Properties of Differential Equations, Higher Education Press: Beijing, 2013.
- [14] J. Patera, R. T. Sharp, and P. Winternitz and H. Zassenhaus, Invariants of real low dimension Lie algebras, *J. Math. Phys.* 17, (1976)986-994.
- [15] J. Patera and P. Winternitz and H. Zassenhaus, Quantum numbers for particles in de Sitter space, *J. Math. Phys.* 17, 717 (1976),717-728.
- [16] Olver P. J, Applications of Lie groups to differential equations, 2nd ed. New York, Springer-Verlag, 1993.
- [17] D. David, N. Kamran, D. Levi, and P. Winternitz, Symmetry reduction for the Kadomtsev- Petviashvili equation using a loop algebra, *J. Math. Phys.* 27(1986) 1225-1237.
- [18] C. P. Boyer and P. Winternitz, Symmetries of selfdual Einstein equations. I: The infinite-dimensional symmetry group and its low dimensional subgroups, *J. Math. Phys.* 30(1989)1081-1094.
- [19] N. H. Ibragimov, *CRC Handbook of Lie Group Analysis of Differential Equations. Vol 2: Applications in Engineering and Physical sciences*, CRC press, Boca Raton, Ann Arbor, London, Tokyo, 1994.
- [20] Buchnev, A. A, Lie group admitted by the equations of motion of an ideal incompressible fluid, *Continuum Dynamics.* 7 (1971),212-214.
- [21] Kudryavtsev, A. G, Group analysis of equations of laser radiation thermal blooming in a moving medium, *Differentsial'niye Uravneniya.* 29(10)1993.
- [22] N.H. Ibragimov and R. N. Ibragimov, Applications of Lie group analysis in geophysical fluid dynamics, Higher Education Press, Beijing, 2011.
- [23] Andreev, V. K., Kaptsov, O.V., Pukhnachev, V.V. and Rodionov, Applications of group theoretic methods in hydrodynamics, Nauk, Novosibirsk(Russian. English translation by Kluwer Academic Publishers), 1998.
- [24] Bluman G W, Kumei S, Symmetries and Differential Equations, Springer-Verlag, New York, Berlin, 1989.
- [25] Temuer Chaolu and Bluman G, An algorithmic method for showing existence of nontrivial non-classical symmetries of partial differential equations without solving determining equations, *J. Math. Anal. Appl.* 411 (2014) 281-296.
- [26] Temuer Chaolu, Bai Yushan, A new algorithmic theory for determining and classifying classical and nonclassical symmetries of partial differential equations, *Science in China: mathematics (Chinese)*, 4(2010) 331-348.

- [27] Nathan Jacobson. Lie algebras Dover Publications, Inc. New York., 1962.
- [28] Allegra Fowler- Wright, The classification of three-dimensional Lie algebras (Thesis), mathematics Institute the univesity of Warwick, 2014.
- [29] Hamidođlu Ali. On General Form of Tanh Method and Its Application to Medical Problems. Springer Singapore, 2017.
- [30] Sudaο Bilige, Temuer Chaolu, An extended simplest equation method and its application to several forms of the fifth-order KdV equation, Appl. Math. Comput. 216(2010) 3146-3153.
- [31] N. H. Ibragimov and V. F. Kovlev, Approxiamate and Renormgroup Symmmtries, Higher Education Press, Beijing, 2009.