

MATCHING HOM-SETTING OF ROTA-BAXTER ALGEBRAS, DENDRIFORM ALGEBRAS AND PRE-LIE ALGEBRAS

DAN CHEN, XIAO-SONG PENG, CHIA ZARGEH, AND YI ZHANG

ABSTRACT. In this paper, we introduce the Hom-algebra setting of the notions of matching Rota-Baxter algebras, matching (tri)dendriform algebras and matching (pre)Lie algebras. Moreover, we study the properties and relationships between categories of these matching Hom-algebraic structures.

CONTENTS

1. Introduction	1
1.1. Hom-algebraic structures	1
1.2. Motivations for matching Hom-algebraic structures	2
1.3. Outline of the paper and summary of results	3
2. Matching Hom-associative, matching Hom-preLie and matching Hom-Lie algebras	4
3. Matching Rota-Baxter algebras and Hom-associative algebras	7
4. Matching Hom-dendriform algebras and matching Hom-tridendriform algebras	10
5. Matching Rota-Baxter operators and Hom-nonassociative algebras	15
References	18

1. INTRODUCTION

1.1. **Hom-algebraic structures.** The origin of Hom-structures may be found in the study of Hom-Lie algebras which were first introduced by Hartwig, Larsson and Silvestrov [24]. Hom-Lie algebras, as a generalization of Lie algebras, are introduced to describe the structures on deformations of the Witt algebra and the Virasoro algebra. More precisely, a Hom-Lie algebra is a triple $(L, [-, -], \alpha)$ consisting of a \mathbf{k} -module L , a bilinear skew-symmetric bracket $[-, -] : L \otimes L \rightarrow L$ and an algebra endomorphism $\alpha : L \rightarrow L$ satisfying the following Hom-Jacobi identity:

$$[\alpha(x), [y, z]] + [\alpha(y), [z, x]] + [\alpha(x), [x, y]] = 0 \text{ for all } x, y, z \in L.$$

Recently, there have been several interesting developments of Hom-Lie algebras in mathematics and mathematical physics, including Hom-Lie bialgebras [10, 13], quadratic Hom-Lie algebras [7], involutive Hom-semigroups [55], deformed vector fields and differential calculus [26], representations [40, 49], cohomology and homology theory [3, 46], Yetter-Drinfeld categories [44], Hom-Yang-Baxter equations [11, 12, 28, 42, 48], Hom-Lie 2-algebras [41, 43], (m, n) -Hom-Lie algebras [33], Hom-left-symmetric algebras [35] and enveloping algebras [20].

Date: June 3, 2020.

2010 Mathematics Subject Classification. 16W99, 17B38, 17B61.

Key words and phrases. Matching Hom-Rota-Baxter algebras; Matching Hom-dendriform algebras; Matching Hom-pre-Lie algebras; Matching Hom-Lie algebras.

In particular, the Hom-Lie algebra on semisimple Lie algebras was studied in [25] and the Hom-Lie structure on affine Kac-Moody was constructed in [38].

In 2008, Makhlouf and Silvestrov [35] introduced the notation of Hom-associative algebras whose associativity law is twisted by a linear map. Usual functors between the categories of Lie algebras and associative algebras have been extended to the Hom-setting. It is shown that a Hom-associative algebra gives rise to a Hom-Lie algebra using the commutator. Since then, various Hom-analogues of some classical algebraic structures have been introduced and studied intensively, such as Hom-coalgebras, Hom-bialgebras and Hom-Hopf algebras [36, 37], Hom-groups [22, 23], Hom-Hopf modules [21], Hom-Lie superalgebras [4, 31], generalize Hom-Lie algebras [32], Hom-Poisson algebras [5].

Dendriform algebras were introduced by Loday [29] with motivation from algebraic K -theory. Latter, tridendriform algebras were proposed by Loday and Ronco [30] in their study of polytopes and Koszul duality. The classical links between Rota-Baxter algebras and (tri)dendriform algebras were given in [1, 14], resembling the structure of Lie algebras on an associative algebra. In 2012, Makhlouf [34] generalized the concepts of dendriform algebras and Rota-Baxter algebras by twisting the identities by mean of a linear map, which were called Hom-dendriform algebras and Rota-Baxter Hom-algebras, respectively. The connections between all these categories of Hom algebras were also investigated in [34]. Due to the fundamental work of Makhlouf [34], we have the following commutative diagram of categories (the arrows will go in the opposite direction for the corresponding operads)

$$\begin{array}{ccc}
 \begin{array}{c} \text{Rota-Baxter} \\ \text{Hom-associative algebra} \end{array} & \xrightarrow{\text{commutator}} & \begin{array}{c} \text{Rota-Baxter} \\ \text{Hom-Lie algebra} \end{array} \\
 \downarrow & & \downarrow \\
 \text{Hom-dendriform algebra} & \xrightarrow{\text{commutator}} & \text{Hom-pre-Lie algebra}
 \end{array}$$

1.2. Motivations for matching Hom-algebraic structures. The recent concept of a matching or multiple Rota-Baxter [52] came from the study of multiple pre-Lie algebras [15] originated from the pioneering work of Bruned, Hairer and Zambotti [8] on algebraic renormalization of regularity structures. It is shown that the matching Rota-Baxter algebra was motivated by the studies of associative Yang-Baxter equations, Volterra integral equations and linear structure of Rota-Baxter operators [52]. More precisely, for exploring the relationship between associative Yang-Baxter equations and classical Yang-Baxter equations, Aguiar [2] proposed a polarized form of the expression on the left hand side of the associative Yang-Baxter equation:

$$\{r, s\} := r_{13}s_{12} - r_{12}s_{23} + r_{23}s_{13},$$


where $r, s \in A \otimes A$ and A is a unitary associative algebra. The corresponding equation

$$r_{13}s_{12} - r_{12}s_{23} + r_{23}s_{13} = 0$$


was called polarized associative Yang-Baxter equation (PAYBE) by Guo and etc [52]. Paralleled to the fact that solutions of the associative Yang-Baxter equation naturally give Rota-Baxter operators, the matching Rota-Baxter operators are determined by solutions of a PAYBE [52].

The basic theory of matching Rota-Baxter algebras was originally established in [52, 17], has proven useful not only in (compatible) multiple operations [27, 50, 51, 53, 54], but also in other areas of mathematics as well, such as polarized associative Yang-Baxter equation [52], algebraic

combinatorics [52, 18], matching shuffle product [17], algebraic integral equation [19], Gröbner-Shirshov bases and Hopf algebras [18]. Based on the close relationships between matching Rota-Baxter algebras, matching dendriform algebras and matching pre-Lie algebras, Guo et al. [52] previously showed the following commutative diagram of categories.



The main purpose of this paper is to extend these matching algebraic structures to the Hom-algebra setting and study the connections between these categories of Hom-algebras. These results give rise to the following commutative diagram of categories.



We would like to emphasize that the notation of matching Hom-Lie Rota-Baxter algebras will play a curial role in mathematical physics. The Rota-Baxter equation on a Lie algebra is the operator form of the classical Yang-Baxter equation [39]. Similarly, there should be a close relationship between the matching Hom Rota-Baxter equation in (18) with weight zero and the polarized classical Yang-Baxter equation, as a Hom-Lie algebra variation of the Hom version of polarized associative Yang-Baxter equation.

1.3. Outline of the paper and summary of results. In section 2, we provide definitions concerning generalization of matching associative algebras, matching pre-Lie algebras to Hom-algebras setting and describe some specific cases of matching Hom-algebraic structures. Also, the close relationship between matching Hom-Lie algebras and Hom-Lie algebras will be shown.

In section 3, we extend the notion of matching Rota-Baxter algebras to the Hom-associative algebra setting. It is also shown that matching Hom-associative Rota-Baxter algebras can be reduced from a matching Rota-Baxter algebra. At the end of this section, the construction of Hom-algebras using elements of the centroid is generalized to the matching Rota-Baxter algebras.

Section 4 is devoted to the definition of matching Hom-(tri)dendriform algebras and the approach of construction of a matching Hom-(tri)dendriform algebra from a matching (tri)dendriform algebra. Some results related to the connections between matching Hom-(tri)dendriform algebras and compatible Hom-associative algebras as well as between matching Hom-dendriform algebras and matching Hom-preLie algebras will be established.

In section 5, the concepts of matching Hom-Lie Rota-Baxter algebras and matching Rota-Baxter algebras involving elements of the centroid of matching Lie Rota-Baxter algebras will be established. Also, some results related to the connection between matching Hom-Lie Rota-Baxter algebra of weight zero and matching Hom-preLie algebra will be obtained.

Notation. Throughout this paper, let \mathbf{k} be a unitary commutative ring unless the contrary is specified, which will be the base ring of all modules, algebras, tensor products, operations as well as linear maps. We always suppose that Ω is a nonempty set. We denote by $P_\Omega := (P_\omega)_{\omega \in \Omega}$ the collection of operations P_ω , $\omega \in \Omega$, where Ω is a set indexing the linear operators.

2. MATCHING HOM-ASSOCIATIVE, MATCHING HOM-PRELIE AND MATCHING HOM-LIE ALGEBRAS

In this section, we give the definitions of matching Hom-associative algebras, compatible Hom-associative algebras, compatible Hom-preLie algebras and compatible Hom-Lie algebras, which generalize the corresponding matching algebraic structures introduced in [52]. Then we explore the relationships between these categories from the point of view of Hom-algebras.

Definition 2.1. A **matching Hom-associative algebra** is a \mathbf{k} -module A together with a collection of binary operations $\cdot_\omega : A \otimes A \rightarrow A$, $\omega \in \Omega$ and a linear map $p : A \rightarrow A$ such that

$$(x \cdot_\alpha y) \cdot_\beta p(z) = p(x) \cdot_\alpha (y \cdot_\beta z) \text{ for all } x, y, z \in A \text{ and } \alpha, \beta \in \Omega.$$

A matching Hom-associative algebra is called **totally compatible** if it satisfies

$$(x \cdot_\alpha y) \cdot_\beta p(z) = p(x) \cdot_\beta (y \cdot_\alpha z) \text{ for all } x, y, z \in A \text{ and } \alpha, \beta \in \Omega.$$

More generally,

Definition 2.2. A **compatible Hom-associative algebra** is a \mathbf{k} -module A together with a collection of binary operations $\cdot_\omega : A \otimes A \rightarrow A$, $\omega \in \Omega$ and a linear map $p : A \rightarrow A$ such that

$$(1) \quad (x \cdot_\alpha y) \cdot_\beta p(z) + (x \cdot_\beta y) \cdot_\alpha p(z) = p(x) \cdot_\alpha (y \cdot_\beta z) + p(x) \cdot_\beta (y \cdot_\alpha z)$$

for all $x, y, z \in A$ and $\alpha, \beta \in \Omega$. For simplicity, we denote it by (A, \cdot_Ω, p) .

Remark 2.3. (a) Any matching Hom-associative algebra or totally compatible Hom-associative algebra is a compatible Hom-associative algebra.

(b) By taking $p = \text{id}$, we recover to the definition of matching associative algebras, totally compatible associative algebra and compatible associative algebra given in [52].

(c) If Ω is a singleton and the characteristic of \mathbf{k} is not 2, then the notation of matching Hom-associative algebras and the notation of compatible Hom-associative algebras are equivalent and recover to the Hom-associative algebras introduced in [35].

Definition 2.4. A **matching Hom-Lie algebra** is a \mathbf{k} -module \mathfrak{g} equipped with a collection of binary operations $[\cdot, \cdot]_\omega : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$, $\omega \in \Omega$ and a linear map $p : \mathfrak{g} \rightarrow \mathfrak{g}$ such that

$$[x, x]_\omega = 0$$

and

$$(2) \quad [p(x), [y, z]_\beta]_\alpha + [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\alpha]_\beta = 0$$

for all $x, y, z \in \mathfrak{g}$ and $\alpha, \beta, \omega \in \Omega$.

Remark 2.5. A totally compatible Hom-associative algebra (A, \cdot_Ω, p) has a natural matching Hom-Lie algebra structure with the Lie bracket defined by

$$[x, y]_\omega := x \cdot_\omega y - y \cdot_\omega x, \text{ for } x, y \in A \text{ and } \omega \in \Omega.$$

The matching Hom-Lie algebra has a close relationship with Hom-Lie algebras. We first record a lemma for a preparation.

Lemma 2.6. *Let $(\mathfrak{g}, [\cdot, \cdot]_\Omega, p)$ be a matching Hom-Lie algebra. Consider linear combinations*

$$(3) \quad [\cdot, \cdot]_A := \sum_{\alpha \in \Omega} a_\alpha [\cdot, \cdot]_\alpha \text{ and } [\cdot, \cdot]_B := \sum_{\beta \in \Omega} b_\beta [\cdot, \cdot]_\beta,$$

where $a_\alpha, b_\beta \in \mathbf{k}$ for $\alpha, \beta \in \Omega$ with finite supports. Then

$$[p(x), [y, z]_B]_A + [p(y), [z, x]_A]_B + [p(z), [x, y]_A]_B = 0 \text{ for } x, y, z \in \mathfrak{g}.$$

Proof. By Eq. (3), for $x, y, z \in \mathfrak{g}$, we have

$$\begin{aligned} [p(x), [y, z]_B]_A &= [p(x), \sum_{\beta \in \Omega} b_\beta [y, z]_\beta]_A = \sum_{\alpha \in \Omega} a_\alpha [p(x), \sum_{\beta \in \Omega} b_\beta [y, z]_\beta]_\alpha \\ &= \sum_{\alpha \in \Omega} \sum_{\beta \in \Omega} a_\alpha b_\beta [p(x), [y, z]_\beta]_\alpha. \end{aligned}$$

Similarly, we also have

$$[p(y), [z, x]_A]_B = \sum_{\alpha \in \Omega} \sum_{\beta \in \Omega} b_\beta a_\alpha [p(y), [z, x]_\alpha]_\beta \quad \text{and} \quad [p(z), [x, y]_A]_B = \sum_{\alpha \in \Omega} \sum_{\beta \in \Omega} b_\beta a_\alpha [p(z), [x, y]_\alpha]_\beta.$$

Since $(\mathfrak{g}, [,]_\Omega, p)$ is a matching Hom-Lie algebra, then

$$[p(x), [y, z]_\beta]_\alpha + [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\alpha]_\beta = 0 \quad \text{for all } x, y, z \in \mathfrak{g} \text{ and } \alpha, \beta \in \Omega.$$

Thus

$$[p(x), [y, z]_B]_A + [p(y), [z, x]_A]_B + [p(z), [x, y]_A]_B = 0,$$

as desired. \square

Proposition 2.7. *Let $(\mathfrak{g}, [,]_\Omega, p)$ be a matching Hom-Lie algebra. Consider linear combinations*

$$(4) \quad [,]_A := \sum_{\omega \in \Omega} a_\omega [,]_\omega, \quad a_\omega \in \mathbf{k},$$

with a finite support. Then $(\mathfrak{g}, [,]_A)$ is a Hom-Lie algebra.

Proof. It follows from Lemma 2.6 by taking $(a_\omega)_{\omega \in \Omega} = (b_\omega)_{\omega \in \Omega}$. \square

More generally, we propose

Definition 2.8. A **compatible Hom-Lie algebra** is a \mathbf{k} -module \mathfrak{g} together with a set of binary operations $[,]_\omega : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$, $\omega \in \Omega$ and a linear map $p : \mathfrak{g} \rightarrow \mathfrak{g}$ such that

$$[x, x]_\omega = 0$$

and

$$(5) \quad \begin{aligned} &[p(x), [y, z]_\alpha]_\beta + [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\alpha]_\beta \\ &+ [p(x), [y, z]_\beta]_\alpha + [p(y), [z, x]_\beta]_\alpha + [p(z), [x, y]_\beta]_\alpha = 0 \end{aligned}$$

for all $x, y, z \in \mathfrak{g}$ and $\omega, \alpha, \beta \in \Omega$.

Remark 2.9. (a) Every matching Hom-Lie algebra is a compatible Hom-Lie algebra.

(b) Given two Hom-Lie algebras $(\mathfrak{g}, [,]_\alpha, p)$ and $(\mathfrak{g}, [,]_\beta, p)$. Define a new bracket $[,] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ as follows:

$$[x, y] := a_\alpha [x, y]_\alpha + b_\beta [x, y]_\beta \quad \text{for some } a_\alpha, b_\beta \in \mathbf{k}.$$

Clearly, this new bracket is both skew symmetric and bilinear. Then $(\mathfrak{g}, [,], p)$ is further a Hom-Lie algebra if $[,]$ satisfies the Hom-Jacobi identity

$$[p(x), [y, z]] + [p(y), [z, x]] + [p(z), [x, y]] = 0.$$

By a direct calculation, we get that this condition is equivalent to Eq. (5).

Proposition 2.10. *Let $(\mathfrak{g}, [,]_\Omega, p)$ be a matching Hom-Lie algebra. Then for $x, y, z \in \mathfrak{g}$ and $\alpha, \beta \in \Omega$, we have*

$$\begin{aligned} [p(x), [y, z]_\alpha]_\beta &= [p(x), [y, z]_\beta]_\alpha, \\ [p(x), [y, z]_\alpha]_\beta + [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\alpha]_\beta &= 0. \end{aligned}$$

Proof. Since Eq. (2) holds for any $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we get

$$(6) \quad [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\beta]_\alpha + [p(x), [y, z]_\beta]_\alpha = 0.$$

Eqs. (2) and (6) result in

$$[p(z), [x, y]_\alpha]_\beta - [p(z), [x, y]_\beta]_\alpha = 0.$$

By the arbitrariness of x, y, z , we have

$$[p(x), [y, z]_\alpha]_\beta = [p(x), [y, z]_\beta]_\alpha$$

and so

$$[p(x), [y, z]_\alpha]_\beta + [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\alpha]_\beta = 0.$$

□

Generalizing the well known result that an associative algebra has a Lie algebra structure via the commutator bracket, we show that a compatible Hom-associative algebra has a compatible Hom-Lie algebra structure.

Proposition 2.11. *Let (A, \cdot_Ω, p) be a compatible Hom-associative algebra. Then $(A, [,]_\Omega, p)$ is a compatible Hom-Lie algebra, where*

$$(7) \quad [,]_\omega : A \otimes A \rightarrow A, [x, y]_\omega := x \cdot_\omega y - y \cdot_\omega x \text{ for } x, y \in A \text{ and } \omega \in \Omega.$$

Proof. For $x, y, z \in A$ and $\alpha, \beta, \omega \in \Omega$, by Eq. (7), we get $[x, x]_\omega = 0$ and

$$\begin{aligned} [p(x), [y, z]_\alpha]_\beta &= [p(x), y \cdot_\alpha z - z \cdot_\alpha y]_\beta \\ &= p(x) \cdot_\beta (y \cdot_\alpha z - z \cdot_\alpha y) - (y \cdot_\alpha z - z \cdot_\alpha y) \cdot_\beta p(x) \\ &= p(x) \cdot_\beta (y \cdot_\alpha z) - p(x) \cdot_\beta (z \cdot_\alpha y) - (y \cdot_\alpha z) \cdot_\beta p(x) + (z \cdot_\alpha y) \cdot_\beta p(x). \end{aligned}$$

Similarly, we have

$$\begin{aligned} [p(y), [z, x]_\alpha]_\beta &= p(y) \cdot_\beta (z \cdot_\alpha x) - p(y) \cdot_\beta (x \cdot_\alpha z) - (z \cdot_\alpha x) \cdot_\beta p(y) + (x \cdot_\alpha z) \cdot_\beta p(y), \\ [p(z), [x, y]_\alpha]_\beta &= p(z) \cdot_\beta (x \cdot_\alpha y) - p(z) \cdot_\beta (y \cdot_\alpha x) - (x \cdot_\alpha y) \cdot_\beta p(z) + (y \cdot_\alpha x) \cdot_\beta p(z), \\ [p(x), [y, z]_\beta]_\alpha &= p(x) \cdot_\alpha (y \cdot_\beta z) - p(x) \cdot_\alpha (z \cdot_\beta y) - (y \cdot_\beta z) \cdot_\alpha p(x) + (z \cdot_\beta y) \cdot_\alpha p(x), \\ [p(y), [z, x]_\beta]_\alpha &= p(y) \cdot_\alpha (z \cdot_\beta x) - p(y) \cdot_\alpha (x \cdot_\beta z) - (z \cdot_\beta x) \cdot_\alpha p(y) + (x \cdot_\beta z) \cdot_\alpha p(y), \\ [p(z), [x, y]_\beta]_\alpha &= p(z) \cdot_\alpha (x \cdot_\beta y) - p(z) \cdot_\alpha (y \cdot_\beta x) - (x \cdot_\beta y) \cdot_\alpha p(z) + (y \cdot_\beta x) \cdot_\alpha p(z). \end{aligned}$$

By Eq. (1), we get

$$\begin{aligned} [p(x), [y, z]_\alpha]_\beta + [p(y), [z, x]_\alpha]_\beta + [p(z), [x, y]_\alpha]_\beta \\ + [p(x), [y, z]_\beta]_\alpha + [p(y), [z, x]_\beta]_\alpha + [p(z), [x, y]_\beta]_\alpha = 0. \end{aligned}$$

Hence $(A, [,]_\Omega, p)$ is a compatible Hom-Lie algebra. □

Now we give the definition of matching Hom-preLie algebras.

Definition 2.12. A **matching Hom-preLie algebra** is a \mathbf{k} -module A together with a family of binary operations $*_{\omega} : A \otimes A \rightarrow A$, $\omega \in \Omega$ and a linear map $p : A \rightarrow A$ such that

$$(8) \quad p(x) *_{\alpha} (y *_{\beta} z) - (x *_{\alpha} y) *_{\beta} p(z) = p(y) *_{\beta} (x *_{\alpha} z) - (y *_{\beta} x) *_{\alpha} p(z)$$

for all $x, y, z \in A$ and $\alpha, \beta \in \Omega$.

Now we give the relationship between matching Hom-preLie algebras and compatible Hom-Lie algebras.

Proposition 2.13. *Let $(A, *_{\Omega}, p)$ be a matching Hom-preLie algebra. Then $(A, [,]_{\Omega}, p)$ is a compatible Hom-Lie algebra, where*

$$(9) \quad [,]_{\omega} : A \otimes A \rightarrow A, [x, y]_{\omega} := x *_{\omega} y - y *_{\omega} x, \text{ for all } x, y \in A \text{ and } \omega \in \Omega.$$

Proof. For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, by Eq. (9) we have $[x, x]_{\omega} = 0$ and

$$\begin{aligned} [p(x), [y, z]_{\alpha}]_{\beta} &= [p(x), y *_{\alpha} z - z *_{\alpha} y]_{\beta} \\ &= p(x) *_{\beta} (y *_{\alpha} z - z *_{\alpha} y) - (y *_{\alpha} z - z *_{\alpha} y) *_{\beta} p(x) \\ &= p(x) *_{\beta} (y *_{\alpha} z) - p(x) *_{\beta} (z *_{\alpha} y) - (y *_{\alpha} z) *_{\beta} p(x) + (z *_{\alpha} y) *_{\beta} p(x). \end{aligned}$$

Similarly, we have

$$\begin{aligned} [p(y), [z, x]_{\alpha}]_{\beta} &= p(y) *_{\beta} (z *_{\alpha} x) - p(y) *_{\beta} (x *_{\alpha} z) - (z *_{\alpha} x) *_{\beta} p(y) + (x *_{\alpha} z) *_{\beta} p(y), \\ [p(z), [x, y]_{\alpha}]_{\beta} &= p(z) *_{\beta} (x *_{\alpha} y) - p(z) *_{\beta} (y *_{\alpha} x) - (x *_{\alpha} y) *_{\beta} p(z) + (y *_{\alpha} x) *_{\beta} p(z), \\ [p(x), [y, z]_{\beta}]_{\alpha} &= p(x) *_{\alpha} (y *_{\beta} z) - p(x) *_{\alpha} (z *_{\beta} y) - (y *_{\beta} z) *_{\alpha} p(x) + (z *_{\beta} y) *_{\alpha} p(x), \\ [p(y), [z, x]_{\beta}]_{\alpha} &= p(y) *_{\alpha} (z *_{\beta} x) - p(y) *_{\alpha} (x *_{\beta} z) - (z *_{\beta} x) *_{\alpha} p(y) + (x *_{\beta} z) *_{\alpha} p(y), \\ [p(z), [x, y]_{\beta}]_{\alpha} &= p(z) *_{\alpha} (x *_{\beta} y) - p(z) *_{\alpha} (y *_{\beta} x) - (x *_{\beta} y) *_{\alpha} p(z) + (y *_{\beta} x) *_{\alpha} p(z). \end{aligned}$$

Then by Eq. (8), we get

$$\begin{aligned} [p(x), [y, z]_{\alpha}]_{\beta} + [p(y), [z, x]_{\alpha}]_{\beta} + [p(z), [x, y]_{\alpha}]_{\beta} \\ + [p(x), [y, z]_{\beta}]_{\alpha} + [p(y), [z, x]_{\beta}]_{\alpha} + [p(z), [x, y]_{\beta}]_{\alpha} = 0. \end{aligned}$$

Hence $(A, [,]_{\Omega}, p)$ is a compatible Hom-Lie algebra. \square

3. MATCHING ROTA-BAXTER ALGEBRAS AND HOM-ASSOCIATIVE ALGEBRAS

In this section, we extend the notion of matching Rota-Baxter algebras to the Hom-associative algebra setting.

Definition 3.1. [52] Let $\lambda_{\Omega} := (\lambda_{\omega})_{\omega \in \Omega} \subseteq \mathbf{k}$ be a set of scalars indexed by Ω . A **matching Rota-Baxter algebra** of weight λ_{Ω} is an associative algebra A equipped with a family $P_{\Omega} := (P_{\omega})_{\omega \in \Omega}$ of linear operators $P_{\omega} : R \rightarrow R$, $\omega \in \Omega$, that satisfy the matching Rota-Baxter equation

$$(10) \quad P_{\alpha}(x) \cdot P_{\beta}(y) = P_{\alpha}(x \cdot P_{\beta}(y)) + P_{\beta}(P_{\alpha}(x) \cdot y) + \lambda_{\beta} P_{\alpha}(x \cdot y), \text{ for all } x, y \in A \text{ and } \alpha, \beta \in \Omega.$$

Definition 3.2. A **matching Hom-associative Rota-Baxter algebra** is a quadruples $(A, \cdot, P_{\Omega}, p)$, where (A, P_{Ω}) is a matching Rota-Baxter algebra and (A, \cdot, p) is a Hom-associative algebra.

Taking $p = \text{id}$, we recover to matching Rota-Baxter associative algebras and denote it by (A, \cdot, P_{Ω}) . If Ω is a singleton, a matching Hom-associative Rota-Baxter algebra becomes a Hom-associative Rota-Baxter algebra given in [34].

A Hom-associative Rota-Baxter algebra can be induced from an associative Rota-Baxter algebra with a particular algebra endomorphism [34]. The following result generalize it to the matching Rota-Baxter case.

Theorem 3.3. *Let (A, \cdot, P_Ω) be a matching Rota-Baxter algebra and $p : A \rightarrow A$ be an algebra endomorphism which commutes with P_ω for all $\omega \in \Omega$. Then $(A, \cdot_p, P_\Omega, p)$, where $x \cdot_p y := p(x \cdot y)$, is a matching Hom-associative Rota-Baxter algebra.*

Proof. The Hom-associative structure of the algebra follows from Yau's Theorem in [45]. We only need to show that the matching Rota-Baxter equation holds. For $x, y \in A$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned}
& P_\alpha(x) \cdot_p P_\beta(y) = p(P_\alpha(x) \cdot P_\beta(y)) \text{ (by the definition of } \cdot_p) \\
& = p(P_\alpha(x \cdot P_\beta(y)) + P_\beta(P_\alpha(x) \cdot y) + \lambda_\beta P_\alpha(x \cdot y)) \text{ (by Eq. (10))} \\
& = p(P_\alpha(x \cdot P_\beta(y))) + p(P_\beta(P_\alpha(x) \cdot y)) + \lambda_\beta p(P_\alpha(x \cdot y)) \\
& = P_\alpha(p(x \cdot P_\beta(y))) + P_\beta(p(P_\alpha(x) \cdot y)) + \lambda_\beta P_\alpha(p(x \cdot y)) \text{ (by } p \circ P_\omega = P_\omega \circ p) \\
& = P_\alpha(x \cdot_p P_\beta(y)) + P_\beta(P_\alpha(x) \cdot_p y) + \lambda_\beta P_\alpha(x \cdot_p y),
\end{aligned}$$

as required. \square

Given a matching Hom-associative Rota-Baxter algebra (A, \cdot, P_Ω, p) , it is natural to wonder that whether this matching Hom-associative Rota-Baxter algebra is induced by an ordinary associative matching Rota-Baxter algebra (A, \cdot', P_Ω) , i.e. p is an algebra endomorphism with respect to \cdot' and $\cdot = p \circ \cdot'$.

Let (A, \cdot, p) be a multiplicative Hom-associative algebra, i.e. $p(a \cdot b) = p(a) \cdot p(b)$ for all $a, b \in A$. It was proved in [16] that in case p is invertible, $(A, p^{-1} \circ \cdot)$ an associative algebra. It is generalized to the multiplicative Hom-associative Rota-Baxter algebras in [34] and the following result generalize it to the multiplicative matching Hom-associative Rota-Baxter algebras.

Proposition 3.4. *Let (A, \cdot, P_Ω, p) be a multiplicative matching Hom-associative Rota-Baxter algebra, where p is invertible and $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Then $(A, \cdot' := p^{-1} \circ \cdot, P_\Omega)$ is an associative matching Rota-Baxter algebra.*

Proof. For $x, y, z \in A$, we have

$$\begin{aligned}
& (x \cdot' y) \cdot' z - x \cdot' (y \cdot' z) \\
& = p^{-1}(p^{-1}(x \cdot y) \cdot z) - p^{-1}(x \cdot p^{-1}(y \cdot z)) \text{ (by } \cdot' = p^{-1} \circ \cdot) \\
& = p^{-2}((x \cdot y) \cdot p(z) - p(x) \cdot (y \cdot z)) \text{ (by } p(x) \cdot p(y) = p(x \cdot y)) \\
& = 0.
\end{aligned}$$

Hence the associativity condition holds. For $\alpha, \beta \in \Omega$, we have

$$\begin{aligned}
& P_\alpha(x) \cdot' P_\beta(y) \\
& = p^{-1}(P_\alpha(x) \cdot P_\beta(y)) \\
& = p^{-1}(P_\alpha(x \cdot P_\beta(y)) + P_\beta(P_\alpha(x) \cdot y) + \lambda_\beta P_\alpha(x \cdot y)) \\
& = P_\alpha(p^{-1}(x \cdot P_\beta(y))) + P_\beta(p^{-1}(P_\alpha(x) \cdot y)) + \lambda_\beta P_\alpha(p^{-1}(x \cdot y)) \\
& = P_\alpha(x \cdot' P_\beta(y)) + P_\beta(P_\alpha(x) \cdot' y) + \lambda_\beta P_\alpha(x \cdot' y).
\end{aligned}$$

Hence the matching Rota-Baxter equation holds for the new multiplication and (A, \cdot', P_Ω) is an associative matching Rota-Baxter algebra. \square

There are two new ways of constructing Hom-associative algebras from a given multiplicative Hom-associative algebra [34, 47].

Definition 3.5. ([34, 47]) Let (A, \cdot, p) be a multiplicative Hom-algebra and $n \geq 0$. Then the following two algebras are also Hom-associative algebras:

- (a) the n -th derived Hom-algebra of type 1 of A defined by

$$A^n = (A, \cdot^{(n)} = p^n \circ \cdot, p^{n+1}),$$

- (b) the n -th derived Hom-algebra of type 2 of A defined by

$$A^n = (A, \cdot^{(n)} = p^{2^n-1} \circ \cdot, p^{2^n}).$$

Now we show that the n -th derived Hom-algebra of type 1 and 2 of a multiplicative matching Hom-associative Rota-Baxter algebra is also a matching Hom-associative Rota-Baxter algebra generalizing the Rota-Baxter case in [34].

Theorem 3.6. Let (A, \cdot, P_Ω, p) be a multiplicative matching Hom-associative Rota-Baxter algebra such that $p \circ P_\omega = P_\omega \circ p$ for all $\omega \in \Omega$. Then

- (a) the n -th derived Hom-algebra of type 1 $(A, \cdot^{(n)} = p^n \circ \cdot, p^{n+1})$ is a matching Hom-associative Rota-Baxter algebra.
(b) the n -th derived Hom-algebra of type 2 $(A, \cdot^{(n)} = p^{2^n-1} \circ \cdot, p^{2^n})$ is a matching Hom-associative Rota-Baxter algebra.

Proof. (a) By [47], (A, \cdot^n, p^{n+1}) is a Hom-associative algebra. Now we show the matching Rota-Baxter equation holds. For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} & P_\alpha(x) \cdot^n P_\beta(y) \\ &= p^n(P_\alpha(x) \cdot P_\beta(y)) \\ &= p^n(P_\alpha(x \cdot P_\beta(y)) + P_\beta(P_\alpha(x) \cdot y) + \lambda_\beta P_\alpha(x \cdot y)) \\ &= P_\alpha(p^n(x \cdot P_\beta(y))) + P_\beta(p^n(P_\alpha(x) \cdot y)) + \lambda_\beta P_\alpha(p^n(x \cdot y)) \\ &= P_\alpha(x \cdot^n P_\beta(y)) + P_\beta(P_\alpha(x) \cdot^n y) + \lambda_\beta P_\alpha(x \cdot^n y). \end{aligned}$$

Thus the matching Rota-Baxter equation holds for the new multiplication.

(b) By [47], $(A, \cdot^{(n)} = p^{2^n-1} \circ \cdot, p^{2^n})$ is also a Hom-associative algebra. For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} & P_\alpha(x) \cdot^n P_\beta(y) \\ &= p^{2^n-1}(P_\alpha(x) \cdot P_\beta(y)) \\ &= p^{2^n-1}(P_\alpha(x \cdot P_\beta(y)) + P_\beta(P_\alpha(x) \cdot y) + \lambda_\beta P_\alpha(x \cdot y)) \\ &= P_\alpha(p^{2^n-1}(x \cdot P_\beta(y))) + P_\beta(p^{2^n-1}(P_\alpha(x) \cdot y)) + \lambda_\beta P_\alpha(p^{2^n-1}(x \cdot y)) \\ &= P_\alpha(x \cdot^n P_\beta(y)) + P_\beta(P_\alpha(x) \cdot^n y) + \lambda_\beta P_\alpha(x \cdot^n y). \end{aligned}$$

This completes the proof. \square

Let (A, \cdot) be an associative algebra. The centroid of A is defined by

$$\text{Cent}(A) := \left\{ p \in \text{End}(A) \mid p(x \cdot y) = p(x) \cdot y = x \cdot p(y) \text{ for all } x, y \in A \right\}.$$

The same definition of the centroid is assumed for Hom-associative algebras.

In [7], Benayadi and Makhoulf gave the construction of Hom-algebras using elements of the centroid for Lie algebras. In [34], the construction was extended to Rota-Baxter algebras. Now we generalize it to the matching Rota-Baxter case.

Proposition 3.7. *Let (A, \cdot, P_Ω) be an associative matching Rota-Baxter algebra. For $p \in \text{Cent}(A)$ and $x, y \in A$, define*

$$x \cdot_p^1 y := p(x) \cdot y \text{ and } x \cdot_p^2 y := p(x) \cdot p(y).$$

If $p \circ P_\omega = P_\omega \circ p$ for all $\omega \in \Omega$, then $(A, \cdot_p^1, P_\Omega, p)$ and $(A, \cdot_p^2, P_\Omega, p)$ are matching Hom-associative Rota-Baxter algebras.

Proof. By [34], (A, \cdot_p^1, p) and (A, \cdot_p^2, p) are Hom-associative algebras. Now we show that they are also matching Rota-Baxter algebras. For $x, y \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} P_\alpha(x) \cdot_p^1 P_\beta(y) &= p(P_\alpha(x)) \cdot P_\beta(y) = P_\alpha(p(x)) \cdot P_\beta(y) \\ &= P_\alpha(p(x) \cdot P_\beta(y)) + P_\beta(P_\alpha(p(x)) \cdot y) + \lambda_\beta P_\alpha(p(x) \cdot y) \\ &= P_\alpha(x \cdot_p^1 P_\beta(y)) + P_\beta(P_\alpha(x) \cdot_p^1 y) + \lambda_\beta P_\alpha(x \cdot_p^1 y) \end{aligned}$$

and

$$\begin{aligned} P_\alpha(x) \cdot_p^2 P_\beta(y) &= p(P_\alpha(x)) \cdot p(P_\beta(y)) = P_\alpha(p(x)) \cdot P_\beta(p(y)) \\ &= P_\alpha(p(x) \cdot P_\beta(p(y))) + P_\beta(P_\alpha(p(x)) \cdot p(y)) + \lambda_\beta P_\alpha(p(x) \cdot p(y)) \\ &= P_\alpha(p(x) \cdot p(P_\beta(y))) + P_\beta(p(P_\alpha(x)) \cdot p(y)) + \lambda_\beta P_\alpha(p(x) \cdot p(y)) \\ &= P_\alpha(x \cdot_p^2 P_\beta(y)) + P_\beta(P_\alpha(x) \cdot_p^2 y) + \lambda_\beta P_\alpha(x \cdot_p^2 y). \end{aligned}$$

This completes the proof. \square

4. MATCHING HOM-DENDRIFORM ALGEBRAS AND MATCHING HOM-TRIDENDRIFORM ALGEBRAS

In this section, we introduce the notions of matching Hom-dendriform algebras and matching Hom-tridendriform algebras generalizing the definitions of matching dendriform algebras and matching tridendriform algebras given in [52].

Definition 4.1. A **matching Hom-dendriform algebra** is a \mathbf{k} -module D together with a family of binary operations $\odot_\omega : D \otimes D \rightarrow D$, where $\odot \in \{<, >\}$ and $\omega \in \Omega$, and a linear map $p : D \rightarrow D$ such that for all $x, y, z \in D$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned} (x <_\alpha y) <_\beta p(z) &= p(x) <_\alpha (y <_\beta z) + p(x) <_\beta (y >_\alpha z), \\ (x >_\alpha y) <_\beta p(z) &= p(x) >_\alpha (y <_\beta z), \\ (x <_\beta y) >_\alpha p(z) + (x >_\alpha y) >_\beta p(z) &= p(x) >_\alpha (y >_\beta z). \end{aligned}$$

For simplicity, we denote it by $(D, <_\Omega, >_\Omega, p)$.

Definition 4.2. A **matching Hom-tridendriform algebra** is a \mathbf{k} -module D together with a family of binary operations $\odot_\omega : D \otimes D \rightarrow D$, where $\odot \in \{<, \bullet, >\}$ and $\omega \in \Omega$, and a linear map $p : D \rightarrow D$ such that for all $x, y, z \in D$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned} (11) \quad (x <_\alpha y) <_\beta p(z) &= p(x) <_\alpha (y <_\beta z) + p(x) <_\beta (y >_\alpha z) + p(x) <_\alpha (y \bullet_\beta z), \\ (12) \quad (x >_\alpha y) <_\beta p(z) &= p(x) >_\alpha (y <_\beta z), \\ (13) \quad p(x) >_\alpha (y >_\beta z) &= (x <_\beta y) >_\alpha p(z) + (x >_\alpha y) >_\beta p(z) + (x \bullet_\beta y) >_\alpha p(z), \\ (14) \quad (x >_\alpha y) \bullet_\beta p(z) &= p(x) >_\alpha (y \bullet_\beta z), \\ (15) \quad (x <_\alpha y) \bullet_\beta p(z) &= p(x) \bullet_\beta (y >_\alpha z), \\ (16) \quad (x \bullet_\alpha y) <_\beta p(z) &= p(x) \bullet_\alpha (y <_\beta z), \end{aligned}$$

$$(17) \quad (x \bullet_{\alpha} y) \bullet_{\beta} p(z) = p(x) \bullet_{\alpha} (y \bullet_{\beta} z).$$

Definition 4.3. (a) Let $(D, \langle_{\Omega}, \rangle_{\Omega}, p)$ and $(D', \langle'_{\Omega}, \rangle'_{\Omega}, p')$ be two matching Hom-dendriform algebras. A linear map $f : D \rightarrow D'$ is called a **matching Hom-dendriform algebra morphism** if for all $\omega \in \Omega$,

$$\langle'_{\omega} \circ (f \otimes f) = f \circ \langle_{\omega}, \rangle_{\omega} \circ (f \otimes f) = f \circ \rangle_{\omega} \text{ and } p' \circ f = f \circ p.$$

(b) Let $(D, \langle_{\Omega}, \bullet_{\Omega}, \rangle_{\Omega}, p)$ and $(D', \langle'_{\Omega}, \bullet'_{\Omega}, \rangle'_{\Omega}, p')$ be two matching Hom-tridendriform algebras. A linear map $f : D \rightarrow D'$ is called a **matching Hom-tridendriform algebra morphism** if for all $\omega \in \Omega$,

$$\langle'_{\omega} \circ (f \otimes f) = f \circ \langle_{\omega}, \bullet'_{\omega} \circ (f \otimes f) = f \circ \bullet'_{\omega}, \rangle_{\omega} \circ (f \otimes f) = f \circ \rangle_{\omega} \text{ and } p' \circ f = f \circ p.$$

The following results show that we can construct a matching Hom-(tri)dendriform algebra from a matching (tri)dendriform algebra, generalizing the (tri)dendriform case in [34].

Theorem 4.4. (a) Let $(D, \langle_{\Omega}, \rangle_{\Omega})$ be a matching dendriform algebra and $p : D \rightarrow D$ be a matching dendriform algebra endomorphism. Then $A_p = (A, \langle_{p,\Omega}, \rangle_{p,\Omega}, p)$, where $\langle_{p,\omega} := p \circ \langle_{\omega}$ and $\rangle_{p,\omega} := p \circ \rangle_{\omega}$ for each $\omega \in \Omega$, is a matching Hom-dendriform algebra. Moreover, suppose that $(A', \langle'_{\Omega}, \rangle'_{\Omega})$ is another matching dendriform algebra and $p' : A' \rightarrow A'$ is a matching dendriform algebra endomorphism. If $f : A \rightarrow A'$ is a matching dendriform algebra morphism that satisfies $f \circ p = p' \circ f$, then

$$f : (D, \langle_{p,\Omega}, \rangle_{p,\Omega}, p) \rightarrow (D', \langle'_{p,\Omega}, \rangle'_{p,\Omega}, p')$$

is a morphism of matching Hom-dendriform algebras.

(b) Let $(D, \langle_{\Omega}, \bullet_{\Omega}, \rangle_{\Omega})$ be a matching tridendriform algebra and $p : D \rightarrow D$ be a matching tridendriform algebra endomorphism. Then $A_p = (A, \langle_{p,\Omega}, \bullet_{p,\Omega}, \rangle_{p,\Omega}, p)$, where $\langle_{p,\omega} := p \circ \langle_{\omega}$, $\bullet_{p,\omega} := p \circ \bullet_{\omega}$ and $\rangle_{p,\omega} := p \circ \rangle_{\omega}$ for each $\omega \in \Omega$, is a matching Hom-tridendriform algebra. Moreover, suppose that $(A', \langle'_{\Omega}, \bullet'_{\Omega}, \rangle'_{\Omega})$ is another matching tridendriform algebra and $p' : A' \rightarrow A'$ is a matching tridendriform algebra endomorphism. If $f : A \rightarrow A'$ is a matching tridendriform algebra morphism that satisfies $f \circ p = p' \circ f$, then

$$f : (D, \langle_{p,\Omega}, \bullet_{p,\Omega}, \rangle_{p,\Omega}, p) \rightarrow (D', \langle'_{p,\Omega}, \bullet'_{p,\Omega}, \rangle'_{p,\Omega}, p')$$

is a morphism of matching Hom-tridendriform algebras.

Proof. We just prove Item (b) and Item (a) can be proved similarly. For any $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} (x \langle_{p,\alpha} y) \langle_{p,\beta} p(z) &= p(p(x \langle_{\alpha} y) \langle_{\beta} p(z)) = p^2((x \langle_{\alpha} y) \langle_{\beta} z); \\ p(x) \langle_{p,\alpha} (y \langle_{p,\beta} z) &= p(p(x) \langle_{\alpha} p(y \langle_{\beta} z)) = p^2(x \langle_{\alpha} (y \langle_{\beta} z)); \\ p(x) \langle_{p,\beta} (y \rangle_{p,\alpha} z) &= p(p(x) \langle_{\beta} p(y \rangle_{\alpha} z)) = p^2(x \langle_{\beta} (y \rangle_{\alpha} z)); \\ p(x) \langle_{p,\alpha} (y \bullet_{p,\beta} z) &= p(p(x) \langle_{\alpha} p(y \bullet_{\beta} z)) = p^2(x \langle_{\alpha} (y \bullet_{\beta} z)). \end{aligned}$$

Hence

$$(x \langle_{p,\alpha} y) \langle_{p,\beta} p(z) = p(x) \langle_{p,\alpha} (y \langle_{p,\beta} z) + p(x) \langle_{p,\beta} (y \rangle_{p,\alpha} z) + p(x) \langle_{p,\alpha} (y \bullet_{p,\beta} z),$$

that is, Eq. (11) holds for $(A, \langle_{p,\Omega}, \bullet_{p,\Omega}, \rangle_{p,\Omega}, p)$. Similarly, Eqs. (12)-(17) holds. Hence $(A, \langle_{p,\Omega}, \bullet_{p,\Omega}, \rangle_{p,\Omega}, p)$ is a matching Hom-tridendriform algebra. And

$$\begin{aligned} f(x) \langle_{p',\alpha} f(y) &= p'(f(x) \langle_{\alpha} f(y)) = p' \circ f(x \langle_{\alpha} y) = f \circ p(x \langle_{\alpha} y) = f(x \langle_{p,\alpha} y); \\ f(x) \rangle_{p',\alpha} f(y) &= p'(f(x) \rangle_{\alpha} f(y)) = p' \circ f(x \rangle_{\alpha} y) = f \circ p(x \rangle_{\alpha} y) = f(x \rangle_{p,\alpha} y); \end{aligned}$$

$$f(x) \bullet_{p',\alpha} f(y) = p'(f(x) \bullet_{\alpha} f(y)) = p' \circ f(x \bullet_{\alpha} (y)) = f \circ p(x \bullet y) = f(x \bullet_{p,\alpha} y).$$

Hence $f : (D, \langle_{p,\Omega}, \bullet_{p,\Omega}, \rangle_{p,\Omega}, p) \rightarrow (D', \langle'_{p,\Omega}, \bullet'_{p,\Omega}, \rangle'_{p,\Omega}, p')$ is a morphism of matching Hom-tridendriform algebras. \square

Now we show that any linear combinations of the operations of a matching Hom-dendriform algebra still result in a matching Hom-dendriform algebra, generalizing the matching dendriform case in [52].

Proposition 4.5. *Let I be an nonempty set. For each $i \in I$, let $A_i : \Omega \rightarrow k$ be a map with finite supports, identified with finite set $A_i = (a_{i,\omega})_{\omega \in \Omega}$, $a_{i,\omega} \in \mathbf{k}$.*

(a) *Let $(D, \langle_{\Omega}, \rangle_{\Omega}, p)$ be a matching Hom-dendriform algebra. Define the following binary operations:*

$$\odot_i := \sum_{\omega \in \Omega} a_{i,\omega} \odot_{\omega}, \text{ where } \odot \in \{\langle, \rangle\} \text{ and } i \in I.$$

Then $(D, \langle_I, \rangle_I, p)$ is also a matching Hom-dendriform algebra.

(b) *Let $(T, \langle_{\Omega}, \bullet_{\Omega}, \rangle_{\Omega}, p)$ be a matching Hom-tridendriform algebra. Define the following binary operations:*

$$\odot_i := \sum_{\omega \in \Omega} a_{i,\omega} \odot_{\omega}, \text{ where } \odot \in \{\langle, \bullet, \rangle\} \text{ and } i \in I.$$

Then $(T, \langle_I, \bullet_I, \rangle_I, p)$ is also a matching Hom-tridendriform algebra.

Proof. We just prove Item (b) and Item (a) can be proved similarly. For $x, y, z \in D$ and $i, j \in I$, we have

$$\begin{aligned} (x \langle_i y) \langle_j p(z) &= \sum_{\beta \in \Omega} b_{j,\beta} \left(\sum_{\alpha \in \Omega} a_{i,\alpha} x \langle_{\alpha} y \right) \langle_{\beta} p(z) = \sum_{\alpha \in \Omega} \sum_{\beta \in \Omega} a_{i,\alpha} b_{j,\beta} (x \langle_{\alpha} y) \langle_{\beta} p(z) \\ &= \sum_{\alpha \in \Omega} \sum_{\beta \in \Omega} a_{i,\alpha} b_{j,\beta} (p(x) \langle_{\alpha} (y \langle_{\beta} z) + p(x) \langle_{\beta} (y \rangle_{\alpha} z) + p(x) \langle_{\alpha} (y \bullet_{\beta} z)) \\ &= \sum_{\alpha \in \Omega} a_{i,\alpha} p(x) \langle_{\alpha} \left(\sum_{\beta \in \Omega} b_{j,\beta} y \langle_{\beta} z \right) + \sum_{\beta \in \Omega} b_{j,\beta} p(x) \langle_{\beta} \left(\sum_{\alpha \in \Omega} a_{i,\alpha} y \rangle_{\alpha} z \right) + \sum_{\alpha \in \Omega} a_{i,\alpha} p(x) \langle_{\alpha} \left(\sum_{\beta \in \Omega} b_{j,\beta} y \bullet_{\beta} z \right) \\ &= \sum_{\alpha \in \Omega} a_{i,\alpha} p(x) \langle_{\alpha} (y \langle_j z) + \sum_{\beta \in \Omega} b_{j,\beta} p(x) \langle_{\beta} (y \rangle_i z) + \sum_{\alpha \in \Omega} a_{i,\alpha} p(x) \langle_{\alpha} (y \bullet_j z) \\ &= p(x) \langle_i (y \langle_j z) + p(x) \langle_j (y \rangle_i z) + p(x) \langle_i (y \bullet_j z). \end{aligned}$$

Hence, Eq. (11) holds. Similarly, Eqs. (12)-(17) hold. Hence $(T, \langle_I, \bullet_I, \rangle_I, p)$ is a matching Hom-tridendriform algebra. \square

The following results establish the connections between matching Hom-(tri)dendriform algebras and compatible Hom-associative algebras, generalizing the well known result that a (tri)dendriform algebra has an associative algebraic structure.

Theorem 4.6. (a) *Let $(A, \langle_{\Omega}, \rangle_{\Omega}, p)$ be a matching Hom-dendriform algebra. Then (A, \cdot_{Ω}, p) is a compatible Hom-associative algebra, where*

$$\cdot_{\omega} : A \otimes A \rightarrow A, \quad x \cdot_{\omega} y := x \langle_{\omega} y + x \rangle_{\omega} y \text{ for } x, y \in A \text{ and } \omega \in \Omega.$$

(b) Let $(A, \prec_\Omega, \bullet_\Omega, \succ_\Omega, p)$ be a matching Hom-tridendriform algebra. Then (A, \cdot_Ω, p) is a compatible Hom-associative algebra, where

$$\cdot_\omega : A \otimes A \rightarrow A, \quad x \cdot_\omega y := x \prec_\omega y + x \bullet_\omega y + x \succ_\omega y \text{ for } x, y \in A \text{ and } \omega \in \Omega.$$

Proof. We only prove Item (b) and Item (a) can be proved similarly. For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} & (x \cdot_\alpha y) \cdot_\beta p(z) + (x \cdot_\beta y) \cdot_\alpha p(z) \\ &= (x \prec_\alpha y + x \bullet_\alpha y + x \succ_\alpha y) \cdot_\beta p(z) + (x \prec_\beta y + x \bullet_\beta y + x \succ_\beta y) \cdot_\alpha p(z) \\ &= (x \prec_\alpha y) \prec_\beta p(z) + (x \bullet_\alpha y) \prec_\beta p(z) + (x \succ_\alpha y) \prec_\beta p(z) + (x \prec_\alpha y) \bullet_\beta p(z) + (x \bullet_\alpha y) \bullet_\beta p(z) \\ & \quad + (x \succ_\alpha y) \bullet_\beta p(z) + (x \prec_\alpha y) \succ_\beta p(z) + (x \bullet_\alpha y) \succ_\beta p(z) + (x \succ_\alpha y) \succ_\beta p(z) + (x \prec_\beta y) \prec_\alpha p(z) \\ & \quad + (x \succ_\beta y) \prec_\alpha p(z) + (x \bullet_\beta y) \prec_\alpha p(z) + (x \prec_\beta y) \bullet_\alpha p(z) + (x \succ_\beta y) \bullet_\alpha p(z) + (x \bullet_\beta y) \bullet_\alpha p(z) \\ & \quad + (x \prec_\beta y) \succ_\alpha p(z) + (x \bullet_\beta y) \succ_\alpha p(z) + (x \succ_\beta y) \succ_\alpha p(z), \end{aligned}$$

and

$$\begin{aligned} & p(x) \cdot_\alpha (y \cdot_\beta z) + p(x) \cdot_\beta (y \cdot_\alpha z) \\ &= p(x) \cdot_\alpha (y \prec_\beta z + y \bullet_\beta z + y \succ_\beta z) + p(x) \cdot_\beta (y \prec_\alpha z + y \bullet_\alpha z + y \succ_\alpha z) \\ &= p(x) \prec_\alpha (y \prec_\beta z) + p(x) \prec_\alpha (y \bullet_\beta z) + p(x) \prec_\alpha (y \succ_\beta z) + p(x) \bullet_\alpha (y \prec_\beta z) + p(x) \bullet_\alpha (y \bullet_\beta z) \\ & \quad + p(x) \bullet_\alpha (y \succ_\beta z) + p(x) \succ_\alpha (y \prec_\beta z) + p(x) \succ_\alpha (y \bullet_\beta z) + p(x) \succ_\alpha (y \succ_\beta z) + p(x) \prec_\beta (y \prec_\alpha z) \\ & \quad + p(x) \prec_\beta (y \bullet_\alpha z) + p(x) \prec_\beta (y \succ_\alpha z) + p(x) \bullet_\beta (y \prec_\alpha z) + p(x) \bullet_\beta (y \bullet_\alpha z) + p(x) \bullet_\beta (y \succ_\alpha z) \\ & \quad + p(x) \succ_\beta (y \prec_\alpha z) + p(x) \succ_\beta (y \bullet_\alpha z) + p(x) \succ_\beta (y \succ_\alpha z). \end{aligned}$$

By Eqs (11-17), we get

$$(x \cdot_\alpha y) \cdot_\beta p(z) + (x \cdot_\beta y) \cdot_\alpha p(z) = p(x) \cdot_\alpha (y \cdot_\beta z) + p(x) \cdot_\beta (y \cdot_\alpha z).$$

Hence (A, \cdot_Ω, p) is a compatible Hom-associative algebra. \square

Now we explore the relationship between matching Hom-dendriform algebras and matching Hom-preLie algebras.

Theorem 4.7. Let $(A, \prec_\Omega, \succ_\Omega, p)$ be a matching Hom-dendriform algebra. Then $(A, *_\Omega, p)$ is a matching Hom-preLie algebra, where

$$*_\omega : A \otimes A \rightarrow A, \quad x *_\omega y := x \succ_\omega y - y \prec_\omega x \text{ for } x, y \in A \text{ and } \omega \in \Omega.$$

Proof. For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} & p(x) *_\alpha (y *_\beta z) - (x *_\alpha y) *_\beta p(z) \\ &= p(x) *_\alpha (y \succ_\beta z - z \prec_\beta y) - (x \succ_\alpha y - y \prec_\alpha x) *_\beta p(z) \\ &= p(x) \succ_\alpha (y \succ_\beta z) - p(x) \succ_\alpha (z \prec_\beta y) - (y \succ_\beta z) \prec_\alpha p(x) + (z \prec_\beta y) \prec_\alpha p(x) - (x \succ_\alpha y) \succ_\beta p(z) \\ & \quad + (y \prec_\alpha x) \succ_\beta p(z) + p(z) \prec_\beta (x \succ_\alpha y) - p(z) \prec_\beta (y \prec_\alpha x) \end{aligned}$$

and

$$\begin{aligned} & p(y) *_\beta (x *_\alpha z) - (y *_\beta x) *_\alpha p(z) \\ &= p(y) *_\beta (x \succ_\alpha z - z \prec_\alpha x) - (y \succ_\beta x - x \prec_\beta y) *_\alpha p(z) \\ &= p(y) \succ_\beta (x \succ_\alpha z) - p(y) \succ_\beta (z \prec_\alpha x) - (x \succ_\alpha z) \prec_\beta p(y) + (z \prec_\alpha x) \prec_\beta p(y) - (y \succ_\beta x) \succ_\alpha p(z) \\ & \quad + (x \prec_\beta y) \succ_\alpha p(z) + p(z) \prec_\alpha (y \succ_\beta x) - p(z) \prec_\alpha (x \prec_\beta y). \end{aligned}$$

By Eqs (11)-(17), we get

$$p(x) *_\alpha (y *_\beta z) - (x *_\alpha y) *_\beta p(z) = p(y) *_\beta (x *_\alpha z) - (y *_\beta x) *_\alpha p(z).$$

Hence $(A, *_\Omega, p)$ is a matching Hom-preLie algebra. \square

A matching Rota-Baxter algebra (A, \cdot, P_Ω) is of weight 0 if the set $\lambda_\Omega = \{0\}$. The connections between Rota-Baxter algebras and (tri)dendriform algebras are given in [2, 14] and extended to matching Rota-Baxter algebras. Now we generalize it to matching Hom-associative Rota-Baxter algebra.

Proposition 4.8. (a) *Let (A, \cdot, P_Ω, p) be a matching Hom-associative Rota-Baxter algebra of weight 0. Assume that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Define the operations $<_\omega$ and $>_\omega$ for $\omega \in \Omega$ by*

$$x <_\omega y := x \cdot P_\omega(y) \text{ and } x >_\omega y = P_\omega(x) \cdot y, \text{ for } x, y \in A.$$

Then $(A, <_\Omega, >_\Omega, p)$ is a matching Hom-dendriform algebra.

(b) *Let (A, \cdot, P_Ω, p) be a matching Hom-associative Rota-Baxter algebra. Assume that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Define the operations $<_\omega, >_\omega, \omega \in \Omega$ by*

$$x <_\omega y := x \cdot P_\omega(y) + \lambda_\omega x \cdot y \text{ and } x >_\omega y = P_\omega(x) \cdot y, \text{ for } x, y \in A.$$

Then $(A, <_\Omega, >_\Omega, p)$ is a matching Hom-dendriform algebra.

Proof. Since Item (a) can be seen as a special case of Item (b) by taking $\lambda_\Omega = \{0\}$, we only prove Item (b). For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} & p(x) <_\alpha (y <_\beta z) + p(x) <_\beta (y >_\alpha z) \\ &= p(x) <_\alpha (y \cdot P_\beta(z) + \lambda_\beta y \cdot z) + p(x) <_\beta (P_\alpha(y) \cdot z) \\ &= p(x) \cdot P_\alpha(y \cdot P_\beta(z) + \lambda_\beta y \cdot z) + \lambda_\alpha p(x) \cdot (y \cdot P_\beta(z) + \lambda_\beta y \cdot z) + p(x) \cdot P_\beta(P_\alpha(y) \cdot z) \\ &\quad + \lambda_\beta p(x) \cdot (P_\alpha(y) \cdot z) \\ &= p(x)(P_\alpha(y) \cdot P_\beta(z)) + \lambda_\alpha p(x) \cdot (y \cdot P_\beta(z)) + \lambda_\alpha \lambda_\beta p(x) \cdot (y \cdot z) + \lambda_\beta p(x) \cdot (P_\alpha(y) \cdot z) \\ &= (x \cdot P_\alpha(y) + \lambda_\alpha x \cdot y) \cdot P_\beta(p(z)) + \lambda_\beta (x \cdot P_\alpha(y) + \lambda_\alpha x \cdot y) \cdot p(z) \\ &= (x \cdot P_\alpha(y) + \lambda_\alpha x \cdot y) <_\beta p(z) = (x <_\alpha y) <_\beta p(z). \end{aligned}$$

Also,

$$\begin{aligned} (x >_\alpha y) <_\beta p(z) &= (P_\alpha(x) \cdot y) <_\beta p(z) = (P_\alpha(x) \cdot y) \cdot P_\beta(p(z)) + \lambda_\beta (P_\alpha(x) \cdot y) \cdot p(z) \\ &= P_\alpha(p(x)) \cdot (y \cdot P_\beta(z)) + \lambda_\beta P_\alpha(p(x)) \cdot (y \cdot z) = P_\alpha(p(x)) \cdot (y \cdot P_\beta(z) + \lambda_\beta y \cdot z) \\ &= P_\alpha(p(x)) \cdot (y <_\beta z) = p(x) >_\alpha (y <_\beta z) \end{aligned}$$

and

$$\begin{aligned} & (x <_\beta y) >_\alpha p(z) + (x >_\alpha y) >_\beta p(z) \\ &= (x \cdot P_\beta(y) + \lambda_\beta x \cdot y) >_\alpha p(z) + (P_\alpha(x) \cdot y) >_\beta p(z) \\ &= P_\alpha(x \cdot P_\beta(y) + \lambda_\beta x \cdot y) \cdot p(z) + P_\beta(P_\alpha(x) \cdot y) \cdot p(z) \\ &= (P_\alpha(x \cdot P_\beta(y)) + P_\beta(P_\alpha(x) \cdot y) + \lambda_\beta P_\alpha(x \cdot y)) \cdot p(z) \\ &= (P_\alpha(x) \cdot P_\beta(y)) \cdot p(z) \\ &= P_\alpha(p(x)) \cdot (P_\beta(y) \cdot z) \end{aligned}$$

$$= p(x) \succ_{\alpha} (y \succ_{\beta} z).$$

Hence $(A, \prec_{\Omega}, \succ_{\Omega}, p)$ is a matching Hom-dendriform algebra. \square

Proposition 4.9. *Let $(A, \cdot, P_{\Omega}, p)$ be a matching Hom-associative Rota-Baxter algebra. Assume that $p \circ P_{\omega} = P_{\omega} \circ p$ for each $\omega \in \Omega$. Define the operations \prec_{ω} , \succ_{ω} and \bullet_{ω} for $\omega \in \Omega$ by*

$$x \prec_{\omega} y := x \cdot P_{\omega}(y), \quad x \succ_{\omega} y = P_{\omega}(x) \cdot y \quad \text{and} \quad x \bullet_{\omega} y = \lambda_{\omega} x \cdot y, \quad \text{for } x, y \in A.$$

Then $(A, \prec_{\Omega}, \bullet_{\Omega}, \succ_{\Omega}, p)$ is a matching Hom-tridendriform algebra.

Proof. For $x, y, z \in A$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} (x \prec_{\alpha} y) \prec_{\beta} (p(z)) &= (x \cdot P_{\alpha}(y)) \cdot P_{\beta}(p(z)) = p(x) \cdot (P_{\alpha}(y) \cdot P_{\beta}(z)) \\ &= p(x) \cdot (P_{\alpha}(y \cdot P_{\beta}(z)) + P_{\beta}(P_{\alpha}(y) \cdot z) + \lambda_{\beta} P_{\alpha}(y \cdot z)), \\ &= p(x) \prec_{\alpha} (y \prec_{\beta} (z)) + p(x) \prec_{\beta} (y \succ_{\alpha} z) + x \prec_{\alpha} (y \bullet_{\beta} z), \\ (x \succ_{\alpha} y) \prec_{\beta} p(z) &= (P_{\alpha}(x) \cdot y) \cdot P_{\beta}(p(z)) = P_{\alpha}(p(x)) \cdot (y \cdot P_{\beta}(z)) = p(x) \succ_{\alpha} (y \prec_{\beta} z), \\ p(x) \succ_{\alpha} (y \succ_{\beta} z) &= P_{\alpha}(p(x)) \cdot (P_{\beta}(y) \cdot z) = (P_{\alpha}(x) \cdot P_{\beta}(y)) \cdot p(z) \\ &= (P_{\alpha}(x \cdot P_{\beta}(y)) + P_{\beta}(P_{\alpha}(x) \cdot y) + \lambda_{\beta} P_{\alpha}(x \cdot y)) \cdot p(z) \\ &= (x \prec_{\beta} y) \succ_{\alpha} p(z) + (x \succ_{\alpha} y) \succ_{\beta} p(z) + (x \bullet_{\beta} y) \succ_{\alpha} p(z), \\ (x \succ_{\alpha} y) \bullet_{\beta} p(z) &= \lambda_{\beta} (P_{\alpha}(x) \cdot y) \cdot p(z) = \lambda_{\beta} P_{\alpha}(p(x)) \cdot (y \cdot z) = p(x) \succ_{\alpha} (y \bullet_{\beta} z), \\ (x \prec_{\alpha} y) \bullet_{\beta} p(z) &= \lambda_{\beta} (x \cdot P_{\alpha}(y)) \cdot p(z) = \lambda_{\beta} p(x) \cdot (P_{\alpha}(y) \cdot z) = p(x) \bullet_{\beta} (y \succ_{\alpha} z), \\ (x \bullet_{\alpha} y) \prec_{\beta} p(z) &= \lambda_{\alpha} (x \cdot y) \cdot P_{\beta}(p(z)) = \lambda_{\alpha} p(x) \cdot (y \cdot P_{\beta}(z)) = p(x) \bullet_{\alpha} (y \prec_{\beta} z), \\ (x \bullet_{\alpha} y) \bullet_{\beta} p(z) &= \lambda_{\alpha} \lambda_{\beta} (x \cdot y) \cdot p(z) = \lambda_{\alpha} \lambda_{\beta} p(x) \cdot (y \cdot z) = p(x) \bullet_{\alpha} (y \bullet_{\beta} z), \end{aligned}$$

as required. \square

Corollary 4.10. (a) *Let $(A, \cdot, P_{\Omega}, p)$ be a matching Hom-associative Rota-Baxter algebra of weight 0. Then $(A, *__{\Omega})$ is a matching Hom-preLie algebra, where*

$$x *__{\omega} y := P_{\omega}(x) \cdot y - y \cdot P_{\omega}(x) \quad \text{for } x, y \in A \text{ and } \omega \in \Omega.$$

(b) *Let $(A, \cdot, P_{\Omega}, p)$ be a matching Hom-associative Rota-Baxter algebra. Then $(A, *__{\Omega})$ is a matching Hom-preLie algebra, where*

$$x *__{\omega} y := P_{\omega}(x) \cdot y - y \cdot P_{\omega}(x) - \lambda_{\omega} y \cdot x \quad \text{for } x, y \in A \text{ and } \omega \in \Omega.$$

Proof. (a) It follows from Theorem 4.7 and Proposition 4.8 (a).

(b) It follows from Theorem 4.7 and Proposition 4.8 (b). \square

5. MATCHING ROTA-BAXTER OPERATORS AND HOM-NONASSOCIATIVE ALGEBRAS

Rota-Baxter Lie algebras were introduced independently by Belavin and Drinfeld and Semenov-Tian-Shansky in [6, 39] and were related to solutions of the (modified) Yang-Baxter equation. Makhlof extended Rota-Baxter operators to the context of Hom-Lie algebras. Now we generalize it to the matching Rota-Baxter case.

Definition 5.1. Let $\lambda_{\Omega} := (\lambda_{\omega})_{\omega \in \Omega} \subseteq \mathbf{k}$ be a family indexed by Ω . A matching Hom-Lie Rota-Baxter algebra is a Hom-Lie algebra $(\mathfrak{g}, [,], p)$ endowed with a set of linear maps $P_{\omega} : \mathfrak{g} \rightarrow \mathfrak{g}$, where $\omega \in \Omega$, subject to the relation

$$(18) \quad [P_{\alpha}(x), P_{\beta}(y)] = P_{\alpha}([x, P_{\beta}(y)]) + P_{\beta}([P_{\alpha}(x), y]) + \lambda_{\beta} P_{\alpha}([x, y]),$$

for all $x, y \in \mathfrak{g}$ and $\alpha, \beta \in \Omega$. For simplicity, we denote it by $(\mathfrak{g}, [,], P_{\Omega}, p)$.

Theorem 5.2. *Let $(\mathfrak{g}, [,], P_\Omega)$ be a matching Lie Rota-Baxter algebra and $p : \mathfrak{g} \rightarrow \mathfrak{g}$ be a Lie algebra endomorphism such that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Then $(\mathfrak{g}, [,]_p, P_\Omega, p)$, where $[,]_p := p \circ [,]$, is a matching Hom-Lie Rota-Baxter algebra.*

Proof. Since $[p(x), [y, z]]_p = p[p(x), p[y, z]] = p^2[x, [y, z]]$, the Hom-Jacobi identity for $(\mathfrak{g}, [,]_p, p)$ follows from the Jacobi identity of $(\mathfrak{g}, [,])$. The skew-symmetry of $(\mathfrak{g}, [,]_p, p)$ holds from the skew-symmetry of $(\mathfrak{g}, [,])$, hence $(\mathfrak{g}, [,]_p, p)$ is a Hom-Lie algebra.

For $x, y \in \mathfrak{g}$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} [P_\alpha(x), P_\beta(y)]_p &= p[P_\alpha(x), P_\beta(y)] \\ &= p(P_\alpha([x, P_\beta(y)]) + P_\beta([P_\alpha(x), y]) + \lambda_\beta P_\alpha([x, y])) \\ &= P_\alpha(p[x, P_\beta(y)]) + P_\beta(p[P_\alpha(x), y]) + \lambda_\beta P_\alpha(p[x, y]) \\ &= P_\alpha([x, P_\beta(y)]_p) + P_\beta([P_\alpha(x), y]_p) + \lambda_\beta P_\alpha([x, y]_p), \end{aligned}$$

as required. \square

Proposition 5.3. *Let $(\mathfrak{g}, [,], P_\Omega, p)$ be a matching Hom-Lie Rota-Baxter algebra such that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Then $(\mathfrak{g}, [,]_{p^{-1}}, P_\Omega)$ is a matching Lie Rota-Baxter algebra.*

Proof. Since $[x, [y, z]]_{p^{-1}} = p^{-1}[x, p^{-1}[y, z]]$, the Jacobi identity of $(\mathfrak{g}, [,]_{p^{-1}})$ holds from the Hom-Jacobi identity of $(\mathfrak{g}, [,], p)$. The skew-symmetry of $(\mathfrak{g}, [,]_{p^{-1}})$ holds from skew symmetry of $(\mathfrak{g}, [,], p)$, hence $(\mathfrak{g}, [,]_{p^{-1}})$ is a Lie algebra.

Since $p \circ P_\omega = P_\omega \circ p$, $p^{-1} \circ P_\omega = P_\omega \circ p^{-1}$. Then

$$\begin{aligned} [P_\alpha(x), P_\beta(y)]_{p^{-1}} &= p^{-1}([P_\alpha(x), P_\beta(y)]) \\ &= p^{-1}(P_\alpha([x, P_\beta(y)]) + P_\beta([P_\alpha(x), y]) + \lambda_\beta P_\alpha([x, y])) \\ &= P_\alpha(p^{-1}([x, P_\beta(y)])) + P_\beta(p^{-1}([P_\alpha(x), y])) + \lambda_\beta P_\alpha(p^{-1}([x, y])) \\ &= P_\alpha([x, P_\beta(y)]_{p^{-1}}) + P_\beta([P_\alpha(x), y]_{p^{-1}}) + \lambda_\beta P_\alpha([x, y]_{p^{-1}}), \end{aligned}$$

as required. \square

Definition 5.4. Let $(\mathfrak{g}, [,], p)$ be a multiplicative Hom-Lie algebra and $n \geq 0$. The n th derived Hom-algebra of \mathfrak{g} is defined by

$$\mathfrak{g}^{(n)} = (\mathfrak{g}, [,]^{(n)} = p^n \circ [,], p^{n+1}).$$

Theorem 5.5. *Let $(\mathfrak{g}, [,], P_\Omega, p)$ be a multiplicative matching Hom-Lie Rota-Baxter algebra and assume that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Then its n th derived Hom-algebra is a matching Hom-Lie Rota-Baxter algebra.*

Proof. Following [47], the n -th derived Hom-algebra is a Hom-Lie algebra. For $x, y \in \mathfrak{g}$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned} [P_\alpha(x), P_\beta(y)]^{(n)} &= p^n([P_\alpha(x), P_\beta(y)]) \\ &= p^n(P_\alpha([x, P_\beta(y)]) + P_\beta([P_\alpha(x), y]) + \lambda_\beta P_\alpha([x, y])) \\ &= P_\alpha(p^n([x, P_\beta(y)])) + P_\beta(p^n([P_\alpha(x), y])) + \lambda_\beta P_\alpha(p^n([x, y])) \\ &= P_\alpha([x, P_\beta(y)]^{(n)}) + P_\beta([P_\alpha(x), y]^{(n)}) + \lambda_\beta P_\alpha([x, y]^{(n)}), \end{aligned}$$

as required. \square

In the following we construct matching Hom-Lie Rota-Baxter algebras involving elements of the centroid of matching Lie Rota-Baxter algebras. Let $(\mathfrak{g}, [,], \Omega, R)$ be a matching Lie Rota-Baxter algebra. The centroid is defined by

$$\text{Cent}(\mathfrak{g}) := \{p \in \text{End}(\mathfrak{g}) : p[x, y] = [p(x), y], \forall x, y \in \mathfrak{g}\}.$$

Proposition 5.6. *Let $(\mathfrak{g}, [,], P_\Omega)$ be a matching Lie Rota-Baxter algebra. Let $p \in \text{Cent}(\mathfrak{g})$ and set for $x, y \in \mathfrak{g}$*

$$[x, y]_p^1 := [p(x), y] \quad \text{and} \quad [x, y]_p^2 := [p(x), p(y)].$$

Assume that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Then $(\mathfrak{g}, [,],_p^1, P_\Omega, p)$ and $(\mathfrak{g}, [,],_p^2, P_\Omega, p)$ are matching Hom-Lie Rota-Baxter algebras.

Proof. Following Proposition 1.12 of [7], $(\mathfrak{g}, [,],_p^1, p)$ and $(\mathfrak{g}, [,],_p^2, p)$ are Hom-Lie algebras. Also,

$$\begin{aligned} [P_\alpha(x), P_\beta(y)]_p^1 &= [p(P_\alpha(x)), P_\beta(y)] = p([P_\alpha(x), P_\beta(y)]) \\ &= p(P_\alpha([x, P_\beta(y)]) + P_\beta([P_\alpha(x), y]) + \lambda_\beta P_\alpha([x, y])) \\ &= P_\alpha([p(x), P_\beta(y)]) + P_\beta([p(P_\alpha(x)), y]) + \lambda_\beta P_\alpha([p(x), y]) \\ &= P_\alpha([x, P_\beta(y)]_p^1) + P_\beta([P_\alpha(x), y]_p^1) + \lambda_\beta P_\alpha([x, y]_p^1) \end{aligned}$$

and

$$\begin{aligned} [P_\alpha(x), P_\beta(y)]_p^2 &= [p(P_\alpha(x)), p(P_\beta(y))] = p([P_\alpha(x), p(P_\beta(y))]) \\ &= -p^2([P_\beta(y), P_\alpha(x)]) = p^2([P_\alpha(x), P_\beta(y)]) \\ &= p^2(P_\alpha([x, P_\beta(y)]) + P_\beta([P_\alpha(x), y]) + \lambda_\beta P_\alpha([x, y])) \\ &= P_\alpha([p(x), p(P_\beta(y))]) + P_\beta([p(P_\alpha(x)), p(y)]) + \lambda_\beta P_\alpha([p(x), p(y)]) \\ &= P_\alpha([x, P_\beta(y)]_p^2) + P_\beta([P_\alpha(x), y]_p^2) + \lambda_\beta P_\alpha([x, y]_p^2). \end{aligned}$$

This completes the proof. \square

Proposition 5.7. *Let $(A, [,], P_\Omega, p)$ be a matching Hom-Lie Rota-Baxter algebra of weight zero (i.e. $\lambda_\omega = 0$ for all $\omega \in \Omega$). Assume that $p \circ P_\omega = P_\omega \circ p$ for each $\omega \in \Omega$. Then $(A, \{*_\omega \mid \omega \in \Omega\}, p)$ is a matching Hom-pre-Lie algebra, where*

$$(19) \quad x *_\omega y = [P_\omega(x), y] \quad \text{for } x, y \in A \text{ and } \omega \in \Omega.$$

Proof. For $x, y, z \in \mathfrak{g}$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} &p(x) *_\alpha (y *_\beta z) - (x *_\alpha y) *_\beta z \\ &= [P_\alpha(p(x)), [P_\beta(y), z]] - [P_\beta([P_\alpha(x), y]), p(z)] \quad (\text{by Eq. (19)}) \\ &= [P_\alpha(p(x)), [P_\beta(y), z]] - [[P_\alpha(x), P_\beta(y)], p(z)] + [P_\alpha([x, P_\beta(y)]), p(z)] \quad (\text{by Eq. (18)}) \\ &= [p(P_\alpha(x)), [P_\beta(y), z]] + [p(z), [P_\alpha(x), P_\beta(y)]] - [P_\alpha([P_\beta(y), x]), p(z)] \quad (\text{by } p \circ P_\alpha = P_\alpha \circ p) \\ &= -[p(P_\beta(y)), [z, P_\alpha(x)]] - [P_\alpha([P_\beta(y), x]), p(z)] \quad (\text{by Hom-Jacobi identity}) \\ &= [P_\beta(p(y)), [P_\alpha(x), z]] - [P_\alpha([P_\beta(y), x]), p(z)] \\ &= p(y) *_\beta (x *_\alpha z) - (y *_\beta x) *_\alpha p(z). \end{aligned}$$

This completes the proof. \square

Data Availability: No data were used to support this study.

Disclosure statement: No potential conflict of interest was reported by the authors.

Acknowledgments: The authors would like to thank the anonymous reviewers for their valuable comments which improved the paper. This work was supported by the National Natural Science Foundation of China (Grant No. 11771191). The work of Chia Zargeh was partially supported by CNPq grant.

REFERENCES

- [1] M. Aguiar, Pre-Poisson algebras, *Lett. Math. Phys.*, 54, (2000), 263-277. [2](#)
- [2] M. Aguiar, On the associative analog of Lie bialgebras, *J. Algebra* **54** (2000), 263-277. [2](#), [14](#)
- [3] F. Ammar, Z. Ejbehi and A. Makhlouf, Cohomology and deformations of Hom-algebras, *J. Lie Theory* **21** (2011), 813-836. [1](#)
- [4] H. Albuquerque, E. Barreiro, A. J. Calderón and J. M. Sánchez, On split regular Hom-Lie superalgebras, *J. Geom. Phys.* **128** (2018), 1-11. [2](#)
- [5] M. J. Argón Perrián and A. J. Calderó Martín, The structure of split regular Hom-Poisson algebras, *Colloq. Math.* **145** (2016), 1-13. [2](#)
- [6] A. A. Belavin and V.G. Drinfeld, Solutions of the classical Yang-Baxter equation for simple Lie algebras, *Funct. Anal. Appl.* **16** (1982), 159-180. [15](#)
- [7] S. Benayadi and A. Makhlouf, Hom-Lie algebras with symmetric invariant nondegenerate bilinear form, *J. Geom. Phys.* **76** (2014), 38-60. [1](#), [9](#), [17](#)
- [8] Y. Bruned, M. Hairer and L. Zambotti, Algebraic renormalisation of regularity structures, *Invent. math.* **215** (2019), 1039-1156. [2](#)
- [9] D. Chen, Y. F. Luo, Y. Zhang and Y. Y. Zhang, Free Ω -Rota-Baxter algebras and Gröbner-Shirshov bases, accepted by *International Journal of Algebra and Computation* (2020).
- [10] Y. Chen, Z. Wang and L. Zhang, Quasitriangular Hom-Lie bialgebras, *J. Lie Theory* **22** (2012), 1075-1089. [1](#)
- [11] Y. Chen and L. Zhang, Hom- \mathcal{O} -operators and Hom-Yang-Baxter equations, *Adv. Math. Phys.* (2015), Art. ID 823756, 11 pp. [1](#)
- [12] Y. Chen and L. Zhang, Hom-Yang-Baxter equations and Frobenius monoidal Hom-algebras, *Adv. Math. Phys.* (2018), Art. ID 2912578, 10 pp. [1](#)
- [13] Y. Chen, H. Zheng and L. Zhang, Double Hom-Associative Algebra and Double Hom-Lie Bialgebra, *Adv. Appl. Clifford Algebra* **30** (2020), Paper No. 8. [1](#)
- [14] K. Ebrahimi-Fard, Loday-type algebras and the Rota-Baxter relation, *Letters in Mathematical Physics* **61** (2002), 130-147. [2](#), [14](#)
- [15] L. Foissy, Algebraic structures on typed decorated rooted trees, arXiv:1811.07572. [2](#)
- [16] A. Gohr, On Hom-algebras with surjective twisting, arXiv:0906.3270v3. [8](#)
- [17] X. Gao, L. Guo and Y. Zhang, Commutative matching Rota-Baxter operators, decorated shuffle products and matching Zinbiel algebras, preprint (2020). [2](#), [3](#)
- [18] X. Gao, L. Guo and Y. Zhang, Hopf algebra of multi-decorated rooted forests, free matching Rota-Baxter algebras and Gröbner-Shirshov bases, arXiv:2002.02864. [3](#)
- [19] L. Guo, R. Gustavson and Y. Li, Volterra integral operators and generalized Reynolds algebras, in preparation. [3](#)
- [20] L. Guo, B. Zhang and S. Zheng, Universal enveloping algebras and Poincaré-Birkhoff-Witt theorem for involutive Hom-Lie algebras, *J. Lie Theory* **26** (2018), 739-759. [1](#)
- [21] S. Guo, X. Zhang and S. Wang, Relative Hom-Hopf modules and total integrals, *J. Math. Phys.* **56** (2015), 021701, 16 pp. [2](#)
- [22] M. Hassanzadeh, Hom-groups, representations and homological algebra, *Colloq. Math.* **158** (2019), 21-38. [2](#)
- [23] M. Hassanzadeh, Lagrange's theorem for Hom-groups, *Rocky Mountain J. Math.* **49** (2019), 773-787. [2](#)
- [24] J. T. Hartwig, D. Larsson and S. D. Silvestrov, Deformations of Lie algebras using σ -derivations, *J. Algebra* **295** (2006), 3140-361. [1](#)

- [25] Q. Jin and X. Li, Hom-Lie algebra structures on semi-simple Lie algebras, *J. Algebra* **319** (2008), 1398-1408. [2](#)
- [26] D. Larsson and S. D. Silvestrov, Quasi-hom-Lie algebras, central extensions and 2-cocycle-like identities, *J. Algebra* **288** (2005), 321-344. [1](#)
- [27] J. Liu, C. Bai and Y. Sheng, Compatible \mathcal{O} -operators on bimodules over associative algebras, *J. Algebra* **532** (2019), 80-118. [2](#)
- [28] L. Liu, A. Makhlouf, C. Menini and F. Panaite, (σ, τ) -Rota-Baxter operators, infinitesimal Hom-bialgebras and the associative (Bi)Hom-Yang-Baxter equation, *Canad. Math. Bull.* **62** (2019), 355-372. [1](#)
- [29] J. L. Loday, Dialgebras, in *Lecture Notes in Math.*, vol. 1763, Springer, Berlin, 2001, 7-66. [2](#)
- [30] J. L. Loday and M. Ronco, Algèbre de Hopf colibres, *C.R. Acad. Sci. Paris*, 337, (2003), 153-158. [2](#)
- [31] L. Ma, L. Chen and J. Zhao, δ -hom-Jordan Lie superalgebras, *Comm. ALgebra* **46** (2018), 1668-1697. [2](#)
- [32] T. Ma, L. Dong and H. Li, General Hom-Lie algebra, *J. Algebra Appl.* **15** (2016), 1650081, 13 pp. [2](#)
- [33] T. Ma and H. Zheng, (m, n) -Hom-Lie algebras, *Publ. Math. Debrecen* **92** (2018), 59-78. [1](#)
- [34] A. Makhlouf, Hom-dendriform algebras and Rota-Baxter Hom-algebras, (English summary) *Operads and universal algebra*, 147-171, *Nankai Ser. Pure Appl. Math. Theoret. Phys.* **9**, World Sci. Publ., Hackensack, NJ, 2012. [2](#), [7](#), [8](#), [9](#), [10](#), [11](#)
- [35] A. Makhlouf and S. Silvestrov, Hom-algebra structures. *J. Gen. Lie Theory Appl.* **2** (2008), 51-64. [1](#), [2](#), [4](#)
- [36] A. Makhlouf and S. Silvestrov, Hom-Lie admissible Hom-coalgebras and Hom-Hopf algebras, (English summary) *Generalized Lie theory in mathematics, physics and beyond*, 189-206, Springer, Berlin, 2009. [2](#)
- [37] A. Makhlouf and S. Silvestrov, Hom-algebras and Hom-coalgebras, *J. Algebra Appl.* **9** (2010), 553-589. [2](#)
- [38] A. Makhlouf and P. Zusmanovich, Hom-Lie structures on Kac-Moody algebras, *J. Algebra* **515** (2018), 278-297. [2](#)
- [39] M. A. Semenov-Tian-Shansky, What is a classical r -matrix?, *Funct. Ana. Appl.* **17** (1983), 259-272. [3](#), [15](#)
- [40] Y. Sheng, Representations of hom-Lie algebras, *Algebr. Represent. Theory* **15** (2012), 1081-1098. [1](#)
- [41] Y. Sheng and D. Chen, Hom-Lie 2-algebras, *J. Algebra* **376** (2013), 174-195. [1](#)
- [42] Y. Sheng and C. Bai, A new approach to hom-Lie bialgebras, *J. Algebra* **399** (2014), 232-250. [1](#)
- [43] L. Song and R. Tang, Derivation Hom-Lie 2-algebras and non-abelian extensions of regular Hom-Lie algebras, *J. Algebra Appl.* **17** (2018), 1850081. [1](#)
- [44] S. Wang and S. Wang, Hom-Lie algebras in Yetter-Drinfeld categories, *Comm. Algebra* **42** (2014), 4526-4547. [1](#)
- [45] D. Yau, Hom-algebras as deformations and homology, *J. Lie Theory* **19** (2009), 409-421. [8](#)
- [46] D. Yau, Hom-algebras and homology, *J. Lie Theory* **19** (2009), 409-421. [1](#)
- [47] D. Yau, Hom-Maltsev, Hom-alternative, and Hom-Jordan algebras, arXiv:1002.3944 [8](#), [9](#), [16](#)
- [48] D. Yau, The Hom-Yang-Baxter equation and Hom-Lie algebras, *J. Math. Phys.* **52** (2011), 053502. [1](#)
- [49] J. Zhao, L. Chen and L. Ma, Representations and T^* -extensions of hom-Jordan-Lie algebras, *Comm. Algebra* **44** (2016), 2786-2812. [1](#)
- [50] Y. Zhang, C. Bai and L. Guo, The category and opera of matching dialgebras, *Appl. Categ. Structures* **21** (2013), 851-865. [2](#)
- [51] Y. Zhang, C. Bai and L. Guo, Totally compatible associative and Lie dialgebras, tridendriform algebras and PostLie algebras, *Sci. China Math.* **57** (2014), 259-273. [2](#)
- [52] Y. Zhang, X. Gao and L. Guo, Matching Rota-Baxter algebras, matching dendriform algebras and matching pre-Lie algebras, *J. Algebra* **552** (2020), 134-170. [2](#), [3](#), [4](#), [7](#), [10](#), [12](#)
- [53] Y. Y. Zhang and X. Gao, Free Rota-Baxter family algebras and (tri)dendriform family algebras, *Pacific J. Math.* **301** (2019), 741-766. [2](#)
- [54] Y. Y. Zhang, X. Gao and D. Manchon, Free (tri)dendriform family algebras, *J. Algebra* **547** (2020), 456-493. [2](#)
- [55] S. Zheng and L. Guo, Free involutive Hom-semigroups and Hom-associative algebras, *Front. Math. China* **11** (2016), 497-508. [1](#)

SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, P. R. CHINA
E-mail address: gaoxing@lzu.edu.cn

SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, P. R. CHINA
E-mail address: pengxiaosong3@163.com

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, SÃO PAULO, SP, BRASIL
E-mail address: ch.zarge@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, P. R. CHINA
E-mail address: zhangy2016@lzu.edu.cn