

Mild assumptions for the derivation of Einstein's effective viscosity formula

David Gérard-Varet, Richard M. Höfer

February 13, 2020

Abstract

We provide a rigorous derivation of Einstein's formula for the effective viscosity of dilute suspensions of n rigid balls, $n \gg 1$, set in a volume of size 1. So far, most justifications were carried under a strong assumption on the minimal distance between the balls: $d_{min} \geq cn^{-\frac{1}{3}}$, $c > 0$. We relax this assumption into a set of two much weaker conditions: one expresses essentially that the balls do not overlap, while the other one gives a control of the number of balls that are close to one another. In particular, our analysis covers the case of suspensions modelled by standard Poisson processes with almost minimal hardcore condition.

1 Introduction

Mixtures of particles and fluids, called *suspensions*, are involved in many natural phenomena and industrial processes. The understanding of their rheology, notably the so-called *effective viscosity* μ_{eff} induced by the particles, is therefore crucial. Many experiments or simulations have been carried out to determine μ_{eff} [9]. For λ large enough, they seem to exhibit some generic behaviour, in terms of the ratio between the solid volume fraction λ and the maximal flowable solid volume fraction λ_c , cf. [9]. Still, a theoretical derivation of the relation $\mu_{eff} = \mu_{eff}(\lambda/\lambda_c)$ observed experimentally is missing, due to the complex interactions involved: hydrodynamic interactions, direct contacts, . . . Mathematical works related to the analysis of suspensions are mostly limited to the *dilute regime*, that is when λ is small.

In these mathematical works, the typical model under consideration is as follows. One considers n rigid balls $B_i = \overline{B(x_i, r_n)}$, $1 \leq i \leq n$, in a fixed compact subset of \mathbb{R}^3 , surrounded by a viscous fluid.

The inertia of the fluid is neglected, leading to the Stokes equations

$$\begin{cases} -\mu\Delta u_n + \nabla p_n = f_n, & x \in \Omega_n = \mathbb{R}^3 \setminus \cup B_i, \\ \operatorname{div} u_n = 0, & x \in \Omega_n, \\ u_n|_{B_i} = u_{n,i} + \omega_{n,i} \times (x - x_i). \end{cases} \quad (1)$$

The last condition expresses a no-slip condition at the rigid spheres, where the velocity is given by some translation velocities $u_{n,i}$ and some rotation vectors $\omega_{n,i}$, $1 \leq i \leq n$. We neglect the inertia of the balls: the $2n$ vectors $u_{n,i}, \omega_{n,i}$ can then be seen as Lagrange multipliers for the $2n$ conditions

$$\int_{\partial B_i} \sigma_\mu(u, p)\nu = - \int_{B_i} f_n, \quad \int_{\partial B_i} \sigma_\mu(u, p)\nu \times (x - x_i) = - \int_{B_i} (x - x_i) \times f_n \quad (2)$$

where $\sigma_\mu = 2\mu D(u)\nu - p\nu$ is the usual Newtonian tensor, and ν the normal vector pointing outward B_i .

The general belief is that one should be able to replace (1)-(2) by an effective Stokes model, with a modified viscosity taking into account the average effect of the particles:

$$\begin{cases} -\operatorname{div}(2\mu_{eff}Du_{eff}) + \nabla p_{eff} = f, & x \in \mathbb{R}^3, \\ \operatorname{div} u_{eff} = 0, & x \in \mathbb{R}^3, \end{cases} \quad (3)$$

with $D = \frac{1}{2}(\nabla + \nabla^t)$ the symmetric gradient. Of course, such average model can only be obtained asymptotically, namely when the number of particles n gets very large. Moreover, for averaging to hold, it is very natural to impose some averaging on the distribution of the balls itself. Our basic hypothesis will therefore be the existence of a limit density, through

$$\frac{1}{n} \sum_i \delta_{x_i} \xrightarrow{n \rightarrow +\infty} \rho(x) dx \quad \text{weakly in the sense of measures} \quad (A0)$$

where $\rho \in L^\infty(\mathbb{R}^3)$ is assumed to be zero outside a smooth open bounded set \mathcal{O} . After playing on the length scale, we can always assume that $|\mathcal{O}| = 1$. Of course, we expect μ_{eff} to be different from μ only in this region \mathcal{O} where the particles are present.

The volume fraction of the balls is then given by $\lambda = \frac{4\pi}{3}nr_n^3$. We shall consider the case where λ is small (dilute suspension), but independent of n so as to derive a non-trivial effect as $n \rightarrow +\infty$. The mathematical questions that follow are:

- Q1 : Can we approximate system (1)-(2) by a system of the form (3) for large n ?
- Q2 : If so, can we provide a formula for μ_{eff} inside \mathcal{O} ? In particular, for small λ , can we derive an expansion

$$\mu_{eff} = \mu + \lambda\mu_1 + \dots \quad ?$$

Regarding Q1, the only work we are aware of is the recent paper [5]. It shows that u_n converges to the solution u_{eff} of an effective model of the type (2), under two natural conditions:

- i) the balls satisfy the separation condition $\inf_{i \neq j} |x_i - x_j| \geq M r_n$, $M > 2$. Note that this is a slight reinforcement of the natural constraint that the balls do not overlap.
- ii) the centers of the balls are obtained from a stationary ergodic point process.

We refer to [5] for all details. Note that in the scalar case, with the Laplacian instead of the Stokes operator, similar results were known since the work of Kozlov, see [13, paragraph 8.6].

Q2, and more broadly quantitative aspects of dilute suspensions, have been studied for long. The pioneering work is due to Einstein [6]. By *neglecting the interaction between the particles*, he computed a first order approximation of the effective viscosity of homogeneous suspensions:

$$\mu_{eff} = (1 + \frac{5}{2}\lambda)\mu \quad \text{in } \mathcal{O}.$$

This celebrated formula was confirmed experimentally afterwards. It was later extended to the inhomogenous case, with formula

$$\mu_{eff} = (1 + \frac{5}{2}\lambda\rho)\mu, \quad (4)$$

see [1, page 16]. Further works investigated the $o(\lambda^2)$ approximation of the effective viscosity, cf. [2] and the recent analysis [8].

Our concern in the present paper is the justification of Einstein's formula. To our knowledge, the first rigorous studies on this topic are [16] and [14]: they rely on homogenization

techniques, and are restricted to suspensions that are periodically distributed in a bounded domain. A more complete justification, still in the periodic setting but based on variational principles, can be found in [10]. Recently, the periodicity assumption was relaxed in [11], [15], and replaced by an assumption on the minimal distance:

$$\text{There exists an absolute constant } c, \text{ such that } \quad \forall n, \forall 1 \leq i \neq j \leq n, \quad |x_i - x_j| \geq cn^{-\frac{1}{3}}. \quad (\text{A1})$$

For instance, introducing the solution u_E of the Einstein's approximate model

$$-\operatorname{div}(2\mu_E Du_E) + \nabla p_E = f, \quad \operatorname{div} u = 0 \quad \text{in } \mathbb{R}^3 \quad (5)$$

with $\mu_E = (1 + \frac{5}{2}\lambda\mu)\mu$, it is shown in [11] that for all $1 \leq p < \frac{3}{2}$,

$$\limsup_{n \rightarrow \infty} \|u_n - u_E\|_{L^p_{loc}(\mathbb{R}^3)} = O(\lambda^{1+\theta}), \quad \theta = \frac{1}{p} - \frac{2}{3}.$$

We refer to [11] for refined statements, including quantitative convergence in n and treatment of polydisperse suspensions.

Although it is a substantial gain over the periodicity assumption, hypothesis (A1) on the minimal distance is still strong. In particular, it is much more stringent than the condition that the rigid balls can not overlap. Indeed, this latter condition reads: $\forall i \neq j, |x_i - x_j| \geq 2r_n$, or equivalently $|x_i - x_j| \geq c\lambda^{1/3}n^{-\frac{1}{3}}$, with $c = 2(\frac{3\pi}{4})^{1/3}$. It follows from (A1) at small λ . On the other hand, one could argue that a simple non-overlapping condition is not enough to ensure the validity of Einstein's formula. Indeed, it is based on neglecting interaction between particles, which is incompatible with too much clustering in the suspension. Still, one can hope that if the balls are not too close from one another *on average*, the formula still holds.

This is the kind of result that we prove here. Namely, we shall replace (A1) by a set of two relaxed conditions:

$$\text{There exists } M > 2, \text{ such that } \quad \forall n, \forall 1 \leq i \neq j \leq n, \quad |x_i - x_j| \geq Mr_n. \quad (\text{B1})$$

$$\text{There exists } C > 0, \text{ such that } \quad \forall \eta > 0, \quad \#\{i, \exists j, |x_i - x_j| \leq \eta n^{-\frac{1}{3}}\} \leq C\eta^3 n \quad (\text{B2})$$

Note that (B1) is slightly stronger than the non-overlapping condition, and was already present in the work [5] to ensure the existence of an effective model. As regards (B2), one can show that it is satisfied almost surely in the case when the particle positions are generated by a stationary ergodic point process if the process does not favor too much close pairs of points. In particular, it is satisfied by a (hard-core) Poisson point process. We postpone further discussion to Section 5.

Under these general assumptions, we obtain:

Theorem 1. *Let $\lambda > 0$, $f \in L^1(\mathbb{R}^3) \cap L^\infty(\mathbb{R}^3)$. For all n , let r_n such that $\lambda = \frac{4\pi}{3}nr_n^3$, let $f_n \in L^{\frac{6}{5}}(\mathbb{R}^3)$, and u_n in $\dot{H}^1(\mathbb{R}^3)$ the solution of (1)-(2). Assume (A0)-(B1)-(B2), and that $f_n \rightarrow f$ in $L^{\frac{6}{5}}(\mathbb{R}^3)$. Then, there exists $p_{min} > 1$ such that for any $p < p_{min}$, any $q < \frac{3p_{min}}{3-p_{min}}$, one can find $\delta > 0$ with the estimate*

$$\|\nabla(u - u_E)\|_{L^p} + \limsup_{n \rightarrow +\infty} \|u_n - u_E\|_{L^q_{loc}} = O(\lambda^{1+\delta}) \quad \text{as } \lambda \rightarrow 0,$$

where u is any weak accumulation point of u_n in $\dot{H}^1(\mathbb{R}^3)$ and u_E satisfies Einstein's approximate model (5).

Here, we use the notation $\dot{H}^1(\mathbb{R}^3)$ for the homogeneous Sobolev space $\dot{H}^1(\mathbb{R}^3) = \{w \in L^6(\mathbb{R}^3) : \nabla w \in L^2(\mathbb{R}^3)\}$ equipped with the L^2 norm of the gradient.

The rest of the paper is dedicated to the proof of this theorem.

2 Main steps of proof

To prove Theorem 1, we shall rely on an enhancement of the general strategy explained in [7], to justify various effective models for conducting and fluid media. Let us point out that one of the examples considered in [7] is the scalar version of (1)-(2). It leads to a proof of a scalar analogue of Einstein's formula, under assumptions (A0), (B1), plus an abstract assumption intermediate between (A1) and (B2). We refer to the discussion at the end of [7] for more details. Nevertheless, to justify the effective fluid model (5) under the mild assumption (B2) will require several new steps. The main difficulty will be to handle particles that are close to one another, and will involve sharp $W^{1,q}$ estimates obtained in [12].

Concretely, let φ be a smooth and compactly supported divergence-free vector field. For each n , we introduce the solution $\phi_n \in \dot{H}^1(\mathbb{R}^3)$ of

$$\begin{aligned} -\operatorname{div}(2\mu D\phi_n) + \nabla q_n &= \operatorname{div}(5\lambda\mu\rho D\varphi) \quad \text{in } \Omega_n, \\ \operatorname{div}\phi_n &= 0 \quad \text{in } \Omega_n, \\ \phi_n &= \varphi + \phi_{n,i} + w_{n,i} \times (x - x_i) \quad \text{in } B_i, \quad 1 \leq i \leq n \end{aligned} \quad (6)$$

where the constant vectors $\phi_{n,i}$, $w_{n,i}$ are associated to the constraints

$$\begin{aligned} \int_{\partial B_i} \sigma_\mu(\phi_n, q_n)\nu &= - \int_{\partial B_i} 5\lambda\mu\rho D\varphi\nu, \\ \int_{\partial B_i} (x - x_i) \times \sigma_\mu(\phi_n, q_n)\nu &= - \int_{\partial B_i} (x - x_i) \times 5\lambda\mu\rho D\varphi\nu. \end{aligned} \quad (7)$$

Testing (1) with $\varphi - \phi_n$, we find after a few integration by parts that

$$\int_{\mathbb{R}^3} 2\mu_E Du_n : D\varphi = \int_{\mathbb{R}^3} f_n \cdot \varphi - \int_{\mathbb{R}^3} f_n \cdot \phi_n.$$

Testing (5) with φ , we find

$$\int_{\mathbb{R}^3} 2\mu_E Du_E : D\varphi = \int_{\mathbb{R}^3} f \cdot \varphi.$$

Combining both, we end up with

$$\int_{\mathbb{R}^3} 2\mu_E D(u_n - u_E) : D\varphi = \int_{\mathbb{R}^3} (f_n - f) \cdot \varphi - \int_{\mathbb{R}^3} f_n \cdot \phi_n. \quad (8)$$

We remind that vector fields u_n, u_E, ϕ_n depend implicitly on λ .

The main point will be to show

Proposition 2. *There exists $p_{\min} > 1$ such that for all $p < p_{\min}$, there exists $\delta > 0$ and $C > 0$, independent of φ , such that*

$$\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f_n \cdot \phi_n \right| \leq C\lambda^{1+\delta} \|\nabla\varphi\|_{L^{p'}}, \quad p' = \frac{p}{p-1}. \quad (9)$$

Let us show how the theorem follows from the proposition. First, by standard energy estimates, we find that u_n is bounded in $\dot{H}^1(\mathbb{R}^3)$ uniformly in n . Let $u = \lim u_{n_k}$ be a weak accumulation point of u_n in this space. Taking the limit in (8), we get

$$\int_{\mathbb{R}^3} 2\mu_E D(u - u_E) : D\varphi = \langle R, \varphi \rangle$$

where $\langle R, \varphi \rangle = \lim_{k \rightarrow +\infty} \int_{\mathbb{R}^3} f_{n_k} \cdot \phi_{n_k}$. Recall that φ is an arbitrary smooth and compactly supported divergence-free vector field and that such functions are dense in the homogeneous Sobolev space of divergence-free functions $\dot{W}_\sigma^{1,p}$. Thus, Proposition 2 implies that R is an element of $\dot{W}_\sigma^{-1,p}$ with $\|R\|_{\dot{W}_\sigma^{-1,p}} = O(\lambda^{1+\delta})$. Moreover, the previous identity is the weak formulation of

$$-\operatorname{div}(2\mu_E D(u - u_E)) + \nabla q = R, \quad \operatorname{div}(u - u_E) = 0 \quad \text{in } \mathbb{R}^3.$$

Writing these Stokes equations with non-constant viscosity as

$$-\mu \Delta(u - u_E) + \nabla q = R + \operatorname{div}(5\lambda\mu\rho D(u - u_E)), \quad \operatorname{div}(u - u_E) = 0 \quad \text{in } \mathbb{R}^3.$$

and using standard estimates for this system, we get

$$\|\nabla(u - u_E)\|_{L^p} \leq C \left(\|R\|_{\dot{W}_\sigma^{-1,p}} + \lambda \|\nabla(u - u_E)\|_{L^p} \right).$$

For λ small enough, the last term is absorbed by the left-hand side, and finally

$$\|\nabla(u - u_E)\|_{L^p(\mathbb{R}^3)} \leq C\lambda^{1+\delta}$$

which implies the first estimate of the theorem. Then, by Sobolev imbedding, for any $q \leq \frac{3p}{3-p}$, and any compact K ,

$$\|u - u_E\|_{L^q(K)} \leq C_{K,q} \lambda^{1+\delta}. \quad (10)$$

We claim that $\limsup_{n \rightarrow \infty} \|u_n - u_E\|_{L^q(K)} \leq C_{K,q} \lambda^{1+\delta}$. Otherwise, there exists a subsequence u_{n_k} and $\varepsilon > 0$ such that $\|u_{n_k} - u_E\|_{L^q(K)} \geq C_{K,q} \lambda^{1+\delta} + \varepsilon$ for all k . Denoting by u a (weak) accumulation point of u_{n_k} in \dot{H}^1 , Rellich's theorem implies that, for a subsequence still denoted u_{n_k} , $\|u_{n_k} - u\|_{L^q(K)} \rightarrow 0$, because $q < 6$ (for p_{\min} taken small enough). Combining this with (10), we reach a contradiction. As p is arbitrary in $(1, p_{\min})$, $q \leq \frac{3p}{3-p}$ is arbitrary in $(1, \frac{3p_{\min}}{3-p_{\min}})$. The last estimate of the theorem is proved.

It remains to prove Proposition 2. Therefore, we need a better understanding of the solution ϕ_n of (6)-(7). Neglecting any interaction between the balls, a natural attempt is to approximate ϕ_n by

$$\phi_n \approx \phi_{\mathbb{R}^3} + \sum_i \phi_{i,n} \quad (11)$$

where $\phi_{\mathbb{R}^3}$ is the solution of

$$-\mu \Delta \phi_{\mathbb{R}^3} + \nabla p_{\mathbb{R}^3} = \operatorname{div}(5\lambda\mu\rho D\varphi), \quad \operatorname{div} \phi_{\mathbb{R}^3} = 0 \quad \text{in } \mathbb{R}^3 \quad (12)$$

and $\phi_{i,n}$ solves

$$-\mu \Delta \phi_{i,n} + \nabla p_{i,n} = 0, \quad \operatorname{div} \phi_{i,n} = 0 \quad \text{outside } B_i, \quad \phi_{i,n}|_{B_i}(x) = D\varphi(x_i)(x - x_i) \quad (13)$$

Roughly, the idea of approximation (11) is that $\phi_{\mathbb{R}^3}$ adjusts to the source term in (6), while for all i , $\phi_{i,n}$ adjusts to the boundary condition at the ball B_i . Indeed, using a Taylor expansion of φ at x_i , and splitting $\nabla\varphi(x_i)$ between its symmetric and skew-symmetric part, we find

$$\phi_{i,n}|_{B_i}(x) \approx D\varphi(x_i)(x - x_i) + \text{rigid vector field} = \phi_{i,n}|_{B_i}(x) + \text{rigid vector field}.$$

Moreover, $\phi_{i,n}$ can be shown to generate no force and torque, so that the extra rigid vector fields (whose role is to ensure the no-force and no-torque conditions), should be small.

Still, approximation (11) may be too crude : the vector fields $\phi_{j,n}$, $j \neq i$, have a non-trivial contribution at B_i , and for the balls B_j close to B_i , which are not excluded by our relaxed

assumption (B1), these contributions may be relatively big. We shall therefore modify the approximation, restricting the sum in (11) to balls far enough from the others.

Therefore, for $\eta > 0$, we introduce a *good* and a *bad* set of indices:

$$\mathcal{G}_\eta = \{1 \leq i \leq n, \forall j \neq i, |x_i - x_j| \geq \eta n^{-\frac{1}{3}}\}, \quad \mathcal{B}_\eta = \{1, \dots, n\} \setminus \mathcal{G}_\eta. \quad (14)$$

The good set \mathcal{G}_η corresponds to balls that are at least $\eta n^{-\frac{1}{3}}$ away from all the others. The parameter $\eta > 0$ will be specified later: we shall consider $\eta = \lambda^\theta$ for some appropriate power $0 < \theta < 1/3$. We set

$$\phi_{app,n} = \phi_{\mathbb{R}^3} + \sum_{i \in \mathcal{G}_\eta} \phi_{i,n} \quad (15)$$

Note that $\phi_{\mathbb{R}^3}$ and $\phi_{i,n}$ are explicit:

$$\phi_{\mathbb{R}^3} = \mathcal{U} \star \operatorname{div} (5\lambda\rho D\varphi), \quad \mathcal{U}(x) = \frac{1}{8\pi} \left(\frac{I}{|x|} + \frac{x \otimes x}{|x|^3} \right)$$

and

$$\phi_{i,n} = r_n V[D\varphi(x_i)] \left(\frac{x - x_i}{r_n} \right) \quad (16)$$

where for all trace-free symmetric matrix S , $V[S]$ solves

$$-\Delta V[S] + \nabla P[S] = 0, \quad \operatorname{div} V[S] = 0 \quad \text{outside } B(0,1), \quad V[S](x) = Sx, \quad x \in B(0,1).$$

with expressions

$$V[S] = \frac{5}{2} S : (x \otimes x) \frac{x}{|x|^5} + Sx \frac{1}{|x|^5} - \frac{5}{2} (S : x \otimes x) \frac{x}{|x|^7}, \quad P[S] = 5 \frac{S : x \otimes x}{|x|^5}.$$

Eventually, we denote

$$\psi_n = \phi_n - \phi_{app,n}.$$

Tedious but straightforward calculations show that

$$-\operatorname{div} (\sigma_\mu(V[S], P[S])) = 5\mu Sx s^1 = -\operatorname{div} (5\mu S 1_{B(0,1)}) \quad \text{in } \mathbb{R}^3$$

where s^1 denotes the surface measure at the unit sphere. It follows that

$$-\mu \Delta \phi_{app,n} + \nabla p_{app,n} = \operatorname{div} \left(5\lambda\mu\rho D\varphi - \sum_{i \in \mathcal{G}_\eta} 5\mu D\varphi(x_i) 1_{B_i} \right), \quad \operatorname{div} \phi_{app,n} = 0 \quad \text{in } \mathbb{R}^3, \quad (17)$$

Moreover, for all $1 \leq i \leq n$,

$$\begin{aligned} \int_{\partial B_i} \sigma_\mu(\phi_{app,n}, p_{app,n}) \nu &= - \int_{\partial B_i} 5\lambda\mu\rho D\varphi, \\ \int_{\partial B_i} (x - x_i) \times \sigma_\mu(\phi_{app,n}, p_{app,n}) \nu &= - \int_{\partial B_i} (x - x_i) \times 5\lambda\mu\rho D\varphi. \end{aligned}$$

Hence, the remainder ψ_n satisfies

$$\begin{aligned} -\mu \Delta \psi_n + \nabla q_n &= 0 \quad \text{in } \Omega_n, \\ \operatorname{div} \psi_n &= 0 \quad \text{in } \Omega_n, \\ \psi_n &= \varphi - \phi_{app,n} + \psi_{n,i} + w_{n,i} \times (x - x_i) \quad \text{in } B_i, \quad 1 \leq i \leq n \end{aligned} \quad (18)$$

where the constant vectors $\psi_{n,i}$, $w_{n,i}$ are associated to the constraints

$$\begin{aligned} \int_{\partial B_i} \sigma_\mu(\psi_n, q_n) \nu &= 0, \\ \int_{\partial B_i} (x - x_i) \times \sigma_\mu(\psi_n, q_n) \nu &= 0. \end{aligned} \tag{19}$$

Estimates on $\phi_{app,n}$ and ψ_n will be postponed to sections 3 and 4 respectively. Regarding $\phi_{app,n}$, we shall prove

Proposition 3. *For all $p \geq 1$,*

$$\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f_n \cdot \phi_{app,n} \right| \leq C_{p,f} (\lambda \eta^3)^{\frac{1}{p}} \|\nabla \varphi\|_{L^{p'}}. \tag{20}$$

Regarding the remainder ψ_n , we shall prove

Proposition 4. *One can find $p_0 \in (1, 2)$, $\theta_0 > 0$, $p_{min} > 1$ satisfying: for all $1 < p < p_{min}$, for all $1 < q < p'$, there exists $c, C > 0$ such that for all $\eta \geq c\lambda^{1/3}$,*

$$\limsup_{n \rightarrow \infty} \|\nabla \psi_n\|_{L^{p_0}(\mathbb{R}^3) + L^2(\mathbb{R}^3) + L^q(\mathbb{R}^3)} \leq C \left(\lambda^{1+\theta_0} + \lambda^{1+\frac{2-p}{2p}} \eta^{-\frac{3}{p}} + (\eta^3 \lambda)^{\frac{1}{r}} \right) \|\nabla \varphi\|_{L^{p'}}.$$

with $\frac{1}{r} + \frac{1}{p'} = \frac{1}{q}$.

Let us explain how to deduce Proposition 2 from these two propositions. Let $1 < p < p_{min}$, with p_{min} as in Proposition 4. By standard estimates, we see that ϕ_n and $\phi_{app,n}$ are bounded uniformly in n in \dot{H}^1 , and so the same holds for ψ_n . It follows that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f_n \cdot \phi_n \right| &= \limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f \cdot \phi_n \right| \leq \limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f \cdot \phi_{app,n} \right| + \limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f \cdot \psi_n \right| \\ &\leq C_{p,f} (\lambda \eta^3)^{\frac{1}{p}} \|\nabla \varphi\|_{L^{p'}} + \left| \int_{\mathbb{R}^3} f \cdot \psi \right| \end{aligned}$$

with ψ a weak accumulation point of ψ_n . Note that we used Proposition 3 to go from the first to the second inequality. By Proposition 4, for all $1 < q < p'$, for all $\eta \geq c\lambda^{1/3}$,

$$\|\nabla \psi\|_{L^{p_0}(\mathbb{R}^3) + L^2(\mathbb{R}^3) + L^q(\mathbb{R}^3)} \leq C \left(\lambda^{1+\theta_0} + \lambda^{1+\frac{2-p}{2p}} \eta^{-\frac{3}{p}} + (\eta^3 \lambda)^{\frac{1}{r}} \right) \|\nabla \varphi\|_{L^{p'}}$$

with $\frac{1}{r} + \frac{1}{p'} = \frac{1}{q}$. As $f \in L^1 \cap L^\infty$, it belongs to the dual of $\dot{W}^{1,p_0}(\mathbb{R}^3) + \dot{W}^{1,2}(\mathbb{R}^3) + \dot{W}^{1,q}(\mathbb{R}^3)$ so that eventually

$$\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f_n \cdot \phi_n \right| \leq C \left(\lambda^{1+\theta_0} + \lambda^{1+\frac{2-p}{2p}} \eta^{-\frac{3}{p}} + (\eta^3 \lambda)^{\frac{1}{r}} \right) \|\nabla \varphi\|_{L^{p'}}$$

To conclude, we adjust properly the various parameters. We look for η in the form $\eta = \lambda^\theta$, with $\theta = \theta(p) < \frac{1}{3}$, so that the lower bound on η needed in Proposition 4 will be satisfied for small enough λ . First, we notice that for any $p < 2$, if θ is such that $2 - p - 6\theta > 0$, then $\lambda^{1+\frac{2-p}{2p}} \eta^{-\frac{3}{p}} = \lambda^{1+\delta_1}$ for some $\delta_1 > 0$. Such a choice of θ being made, we can choose $r = r(p)$ close enough to 1 in a way that $(\eta^3 \lambda)^{\frac{1}{r}} = \lambda^{1+\delta_2}$ for some $\delta_2 > 0$. Upon taking smaller p_{min} , we can further ensure that $\frac{1}{r} + \frac{1}{p'} < 1$ so that we can define q by $\frac{1}{r} + \frac{1}{p'} = \frac{1}{q}$, and use the previous estimates. Eventually, for any $1 < p < p_{min}$, for λ small enough,

$$\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} f_n \cdot \phi_n \right| \leq C \lambda^{1+\delta} \|\nabla \varphi\|_{L^{p'}(\mathbb{R}^3)}$$

where $\delta = \min(\theta_0, \delta_1, \delta_2)$.

3 Bound on the approximation

This section is devoted to the proof of Proposition 3. We decompose

$$\phi_{app,n} = \phi_{app,n}^1 + \phi_{app,n}^2 + \phi_{app,n}^3$$

where

$$\begin{aligned} -\mu\Delta\phi_{app,n}^1 + \nabla p_{app,n}^1 &= \operatorname{div} \left(5\lambda\mu\rho D\varphi - \sum_{1 \leq i \leq n} 5\mu D\varphi(x_i) \mathbf{1}_{B_i} \right), \quad \operatorname{div} \phi_{app,n}^1 = 0 \quad \text{in } \mathbb{R}^3, \\ -\mu\Delta\phi_{app,n}^2 + \nabla p_{app,n}^2 &= \operatorname{div} \left(\sum_{i \in \mathcal{B}_\eta} 5\mu D\varphi(x) \mathbf{1}_{B_i} \right), \quad \operatorname{div} \phi_{app,n}^2 = 0 \quad \text{in } \mathbb{R}^3, \\ -\mu\Delta\phi_{app,n}^3 + \nabla p_{app,n}^3 &= \operatorname{div} \left(\sum_{i \in \mathcal{B}_\eta} 5\mu (D\varphi(x_i) - D\varphi(x)) \mathbf{1}_{B_i} \right), \quad \operatorname{div} \phi_{app,n}^3 = 0 \quad \text{in } \mathbb{R}^3. \end{aligned}$$

By standard energy estimates, $\phi_{app,n}^k$ is seen to be bounded in n in \dot{H}^1 , for all $1 \leq k \leq 3$. We shall prove next that $\phi_{app,n}^1$ and $\phi_{app,n}^3$ converge in the sense of distributions to zero, while for any f with $D(\Delta)^{-1}\mathbb{P}f \in L^\infty$ (\mathbb{P} denoting the standard Helmholtz projection), for any $p \geq 1$,

$$\left| \int_{\mathbb{R}^3} f \cdot \phi_{app,n}^2 \right| \leq C_{f,p} (\lambda\eta^3)^{\frac{1}{p}} \|\nabla\varphi\|_{L^{p'}}, \quad p' = \frac{p}{p-1}. \quad (21)$$

Proposition 3 follows easily from those properties.

We start with

Lemma 5. *Under assumption (A0), $\sum_{1 \leq i \leq n} D\varphi(x_i) \mathbf{1}_{B_i} \rightharpoonup \lambda\rho D\varphi$ weakly* in L^∞ .*

Proof. As the balls are disjoint, $|\sum_{1 \leq i \leq n} D\varphi(x_i) \mathbf{1}_{B_i}| \leq \|D\varphi\|_{L^\infty}$. Let $g \in C_c(\mathbb{R}^3)$, and denote $\delta_n = \frac{1}{n} \sum_i \delta_{x_i}$ the empirical measure. We write

$$\begin{aligned} \int_{\mathbb{R}^3} \sum_{1 \leq i \leq n} D\varphi(x_i) \mathbf{1}_{B_i}(y) g(y) dy &= \sum_{1 \leq i \leq n} D\varphi(x_i) \int_{B(0,r_n)} g(x_i + y) dy \\ &= n \int_{\mathbb{R}^3} D\varphi(x) \int_{B(0,r_n)} g(x + y) dy d\delta_n(x) \\ &= nr_n^3 \int_{\mathbb{R}^3} \int_{B(0,1)} g(x + r_n z) dz d\delta_n(x). \end{aligned}$$

The sequence of bounded continuous functions $x \rightarrow \int_{B(0,1)} g(x + r_n z) dz$ converges uniformly to the function $x \rightarrow \frac{4\pi}{3} g(x)$ as $n \rightarrow +\infty$. We deduce:

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^3} \sum_{1 \leq i \leq n} D\varphi(x_i) \mathbf{1}_{B_i}(y) g(y) dy = \lim_{n \rightarrow \infty} \lambda \int_{\mathbb{R}^3} D\varphi(x) g(x) d\delta_n(x) = \lambda \int_{\mathbb{R}^3} D\varphi(x) g(x) \rho(x) dx$$

where the last equality comes from (A0). The lemma follows by density of C_c in L^1 .

Let now $h \in C_c^\infty(\mathbb{R}^3)$ and $v = (\Delta)^{-1}\mathbb{P}h$. We find

$$\begin{aligned} \langle \phi_{app,n}^1, h \rangle &= \langle \phi_{app,n}^1, \Delta v \rangle = \langle \Delta \phi_{app,n}^1, v \rangle \\ &= \int_{\mathbb{R}^3} \left(5\lambda\mu\rho D\varphi - \sum_{1 \leq i \leq n} 5\mu D\varphi(x_i) \mathbf{1}_{B_i} \right) \cdot Dv \rightarrow 0 \quad \text{as } n \rightarrow +\infty \end{aligned}$$

where we used the previous lemma and the fact that Dv belongs to L^1_{loc} and φ has compact support. Hence, $\phi^1_{app,n}$ converges to zero in the sense of distributions. As regards $\phi^3_{app,n}$, we notice that

$$\begin{aligned} \left\| \sum_{i \in \mathcal{B}_\eta} 5\mu(D\varphi(x) - D\varphi(x_i))1_{B_i} \right\|_{L^1} &\leq \|\nabla^2\varphi\|_{L^\infty} \sum_{1 \leq i \leq n} \int_{B_i} |x - x_i| dx \\ &\leq \|\nabla^2\varphi\|_{L^\infty} \lambda r_n \rightarrow 0 \quad \text{as } n \rightarrow +\infty \end{aligned}$$

Using the same duality argument as for $\phi^1_{app,n}$ (see also below), we get that $\phi^3_{app,n}$ converges to zero in the sense of distributions.

It remains to show (21). We use a simple Hölder estimate, and write for all $p \geq 1$:

$$\begin{aligned} \left\| \sum_{i \in \mathcal{B}_\eta} 5\mu D\varphi 1_{B_i} \right\|_{L^1} &\leq 5\mu \left\| \sum_{i \in \mathcal{B}_\eta} 1_{B_i} \right\|_{L^p} \|D\varphi\|_{L^{p'}} = 5\mu (\text{card } \mathcal{B}_\eta)^{\frac{1}{p}} \left(\frac{4\pi}{3} r_n^3 \right)^{\frac{1}{p}} \|D\varphi\|_{L^{p'}} \\ &\leq C(\eta^3 \lambda)^{\frac{1}{p}} \|D\varphi\|_{L^{p'}} \end{aligned}$$

where the last inequality follows from (B2). Denoting $v = (\Delta)^{-1}\mathbb{P}f$, we have this time

$$\int_{\mathbb{R}^3} f \cdot \phi^2_{app,n} = \int_{\mathbb{R}^3} Dv \cdot \sum_{i \in \mathcal{B}_\eta} 5\mu D\varphi 1_{B_i} \leq C \|Dv\|_{L^\infty} (\eta^3 \lambda)^{\frac{1}{p}} \|D\varphi\|_{L^{p'}}$$

which implies (21).

4 Bound on the remainder

We focus here on estimates for the remainder $\psi_n = \phi_n - \phi_{app,n}$, which satisfies (18)-(19). We will use as a black box sharp $W^{1,q}$ estimates derived recently in [12], which notably apply to systems of the form

$$-\mu\Delta\psi + \nabla p = 0, \quad \text{div } \psi = 0 \quad \text{in } \Omega_n, \quad D\psi = D\tilde{\psi} \quad \text{in } B_i, \quad 1 \leq i \leq n \quad (22)$$

together with the constraints

$$\int_{\partial B_i} \sigma_\mu(\psi, p)\nu = \int_{\partial B_i} (x - x_i) \times \sigma_\mu(\psi, p)\nu = 0, \quad 1 \leq i \leq n. \quad (23)$$

We first recall that, *under assumption* (B1), one has the somehow standard estimate

$$\|\nabla\psi\|_{L^2(\mathbb{R}^3)} \leq C \|D\tilde{\psi}\|_{L^2(\cup_i B_i)} \quad (24)$$

for a constant C independent of n . A sketch of proof is as follows. By a classical variational characterization of ψ , we have

$$\|\nabla\psi\|_{L^2(\mathbb{R}^3)}^2 = 2\|D\psi\|_{L^2(\mathbb{R}^3)}^2 = \inf \{ 2\|DU\|_{L^2(\mathbb{R}^3)}^2, \quad DU = D\tilde{\psi} \text{ on } \cup_i B_i \}.$$

Hence it is enough to build one vector field U satisfying the same condition on $\cup_i B_i$ and fulfilling the estimate. By adding an appropriate rigid vector field to $\tilde{\psi}$ on each B_i , so that $D\tilde{\psi}$ is unchanged, one can always assume a Poincaré and a Korn inequality: $r_n^{-2} \|\tilde{\psi}\|_{L^2(B_i)}^2 \leq C \|\nabla\tilde{\psi}\|_{L^2(B_i)}^2 \leq C' \|D\tilde{\psi}\|_{L^2(B_i)}^2$, where the factor r_n^{-2} comes from scaling considerations. Hence, after these simplifications, it is enough to find a U satisfying $U = \tilde{\psi}$ on each B_i ,

and such that $\|\nabla U\|_{L^2(\mathbb{R}^3)}^2 \leq C \left(r_n^{-2} \|\psi\|_{L^2(\cup B_i)}^2 + \|\nabla \tilde{\psi}\|_{L^2(\cup B_i)}^2 \right)$. This can then be done under assumption (B1), using standard considerations on the Bogovskii operator: one can even choose U to be supported in a vicinity of $\cup B_i$. We refer to [7, Lemma 5] for details.

However, the derivation of an L^p analogue of (24) is much more unclear, as no variational characterization is available. This problem was tackled recently by the second author in [12], through a careful analysis of the so-called *method of reflections*, introduced by Smoluchowski. We shall use the following

Theorem 6. [12, Proposition 5.2 and Remark 5.6]

Assume that $\inf_{i \neq j} |x_i - x_j| \geq M r_n$. Then,

- i) for all $M > 2$, there exists $\varepsilon_M > 0$ such that for all $q \in [2 - \varepsilon_M, 2 + \varepsilon_M]$, and $\tilde{\psi} \in L^q(\cup_i B_i)$, there exists a unique solution $\psi \in \dot{W}^{1,q}(\mathbb{R}^3)$ of (22)-(23) such that

$$\|\nabla \psi\|_{L^q(\mathbb{R}^3)} \leq C_M \|D\tilde{\psi}\|_{L^q(\cup B_i)}$$

- ii) for all $q > 1$, there exists M_q such that for all $M \geq M_q$, and $\tilde{\psi} \in L^q(\cup_i B_i)$, there exists a unique solutions $\psi \in \dot{W}^{1,q}(\mathbb{R}^3)$ of (22)-(23) such that

$$\|\nabla \psi\|_{L^q(\mathbb{R}^3)} \leq C_q \|D\tilde{\psi}\|_{L^q(\cup B_i)}$$

We now come back to the analysis of ψ_n . We decompose the field $\varphi - \phi_{app,n}$ that appears in the third line of (18): we write

$$\varphi - \phi_{app,n} = \tilde{\psi}_n^1 + \tilde{\psi}_n^2 + \tilde{\psi}_n^3$$

where

$$\forall 1 \leq i \leq n, \forall x \in B_i, \quad \tilde{\psi}_n^1(x) = -\phi_{\mathbb{R}^3}(x) - \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_n}} \phi_{j,n}(x)$$

and

$$\begin{aligned} \forall i \in \mathcal{G}_\eta, \forall x \in B_i, \quad \tilde{\psi}_n^2(x) &= \varphi(x) - \varphi(x_i) - \nabla \varphi(x_i)(x - x_i) + \left(\varphi(x_i) + \frac{1}{2} \operatorname{curl} \varphi(x_i) \times (x - x_i) \right), \\ \forall i \in \mathcal{B}_\eta, \forall x \in B_i, \quad \tilde{\psi}_n^2(x) &= 0, \\ \forall i \in \mathcal{G}_\eta, \forall x \in B_i, \quad \tilde{\psi}_n^3(x) &= 0, \\ \forall i \in \mathcal{B}_\eta, \forall x \in B_i, \quad \tilde{\psi}_n^3(x) &= \varphi(x). \end{aligned}$$

We remind that the sum in (15) is restricted to indices $i \in \mathcal{G}_\eta$ and that $\phi_{i,n}(x) = D\varphi(x_i)(x - x_i)$ for x in B_i . This explains the distinction between $\tilde{\psi}_n^2$ and $\tilde{\psi}_n^3$.

Clearly, $\psi_n = \sum_{k=1}^3 \psi_n^k$, where ψ_n^k is the solution of (22)-(23) associated to data $\tilde{\psi}_n^k$. The control of ψ_n^2 is the simplest to obtain: we apply estimate (24), valid under the sole assumption (B1). We find

$$\|\nabla \psi_n^2\|_{L^2(\mathbb{R}^3)} \leq C \|D\tilde{\psi}_n^2\|_{L^2(\cup B_i)} \leq C \|D^2 \varphi\|_{L^\infty} \left(\sum_{i \in \mathcal{G}_\eta} \int_{B_i} |x - x_i|^2 dx \right)^{1/2} \leq C' \lambda^{1/2} r_n.$$

Hence,

$$\lim_{n \rightarrow +\infty} \|\nabla \psi_n^2\|_{L^2(\mathbb{R}^3)} = 0. \quad (25)$$

As regards ψ_n^1 , we apply again (24) and find

$$\|\nabla\psi_n^1\|_{L^2(\mathbb{R}^3)} \leq C \left(\|D\phi_{\mathbb{R}^3}\|_{L^2(\cup B_i)} + \left(\sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} D\phi_{j,n} \right|^2 \right)^{1/2} \right) \quad (26)$$

For any $r, s < +\infty$ with $\frac{1}{r} + \frac{1}{s} = \frac{1}{2}$, we obtain

$$\|D\phi_{\mathbb{R}^3}\|_{L^2(\cup B_i)} \leq \|1_{\cup B_i}\|_{L^r(\mathbb{R}^3)} \|D\phi_{\mathbb{R}^3}\|_{L^s(\mathbb{R}^3)} \leq C \|1_{\cup B_i}\|_{L^r(\mathbb{R}^3)} \|\lambda\rho D\varphi\|_{L^s(\mathbb{R}^3)} \quad (27)$$

using standard L^s estimate for system (12). Hence,

$$\|D\phi_{\mathbb{R}^3}\|_{L^2(\cup B_i)} \leq C' \lambda^{1+\frac{1}{r}} \|D\varphi\|_{L^s(\mathbb{R}^3)}.$$

Note that we can choose any $s > 2$, this lower bound coming from the requirement $\frac{1}{r} + \frac{1}{s} = \frac{1}{2}$. Introducing p such that $s = p'$, we find that for any $p < 2$,

$$\|D\phi_{\mathbb{R}^3}\|_{L^2(\cup B_i)} \leq C' \lambda^{\frac{1}{2}+\frac{1}{p}} \|D\varphi\|_{L^{p'}(\mathbb{R}^3)}. \quad (28)$$

The treatment of the second term at the r.h.s. of (26) is more delicate. We write, see (16):

$$D\phi_{j,n}(x) = DV[D\varphi(x_j)] \left(\frac{x-x_j}{r_n} \right) = \mathcal{V}[D\varphi(x_j)] \left(\frac{x-x_j}{r_n} \right) + \mathcal{W}[D\varphi(x_j)] \left(\frac{x-x_j}{r_n} \right) \quad (29)$$

where $\mathcal{V}[S] = D\left(\frac{5}{2}S : (x \otimes x) \frac{x}{|x|^5}\right)$, $\mathcal{W}[S] = D\left(\frac{Sx}{|x|^5} - \frac{5}{2}(S : x \otimes x) \frac{x}{|x|^7}\right)$.

We have:

$$\sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \mathcal{W}[D\varphi(x_j)] \left(\frac{x-x_j}{r_n} \right) \right|^2 dx \leq C r_n^{10} \sum_i \int_{B_i} \left(\sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} |D\varphi(x_j)| |x-x_j|^{-5} \right)^2 dx$$

For all i , for all $j \in \mathcal{G}_\eta$ with $j \neq i$, and all $(x, y) \in B_i \times B(x_j, \frac{\eta}{4}n^{-\frac{1}{3}})$, we have for some absolute constants $c, c' > 0$:

$$|x-x_j| \geq c|x-y| \geq c'\eta n^{-\frac{1}{3}}.$$

Denoting $B_j^* = B(x_j, \frac{\eta}{4}n^{-\frac{1}{3}})$ We deduce

$$\begin{aligned} & \sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \mathcal{W}[D\varphi(x_j)] \left(\frac{x-x_j}{r_n} \right) \right|^2 dx \\ & \leq C r_n^{10} \sum_i \int_{B_i} \left(\sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \frac{1}{|B_j^*|} \int_{B_j^*} |x-y|^{-5} 1_{\{|x-y|>c'\eta n^{-\frac{1}{3}}\}} (x-y) |D\varphi(x_j)| dy \right)^2 dx \\ & \leq C' n^2 \frac{r_n^{10}}{\eta^6} \int_{\mathbb{R}^3} 1_{\cup B_i}(x) \left(\int_{\mathbb{R}^3} |x-y|^{-5} 1_{\{|x-y|>c'\eta n^{-\frac{1}{3}}\}} (x-y) \sum_{1 \leq j \leq n} |D\varphi(x_j)| 1_{B_j^*}(y) dy \right)^2 dx \end{aligned}$$

Using Hölder and Young's convolution inequalities, we find that for all r, s with $\frac{1}{r} + \frac{1}{s} = 1$,

$$\begin{aligned} & \int_{\mathbb{R}^3} 1_{\cup B_i}(x) \left(\int_{\mathbb{R}^3} |x-y|^{-5} 1_{\{|x-y|>c'\eta n^{-\frac{1}{3}}\}} (x-y) \sum_{1 \leq j \leq n} |D\varphi(x_j)| 1_{B_j^*}(y) dy \right)^2 dx \\ & \leq \|1_{\cup B_i}\|_{L^r} \|(|x|^{-5} 1_{\{|x|>c'\eta n^{-\frac{1}{3}}\}}) \star \sum_{1 \leq j \leq n} |D\varphi(x_j)| 1_{B_j^*}\|_{L^{2s}}^2 \\ & \leq \|1_{\cup B_i}\|_{L^r} \| |x|^{-5} 1_{\{|x|>c'\eta n^{-\frac{1}{3}}\}} \|_{L^1}^2 \left\| \sum_{1 \leq j \leq n} |D\varphi(x_j)| 1_{B_j^*} \right\|_{L^{2s}}^2 \\ & \leq C \lambda^{\frac{1}{r}} (\eta n^{-\frac{1}{3}})^{-4} \left(\sum_j |D\varphi(x_j)|^{2s} \eta^3 n^{-1} \right)^{\frac{1}{s}} \end{aligned}$$

Note that, by (A0), $\frac{1}{n} \sum_j |D\varphi(x_j)|^t \rightarrow \int_{\mathbb{R}^3} |D\varphi|^t \rho$ as $n \rightarrow +\infty$. We end up with

$$\limsup_{n \rightarrow \infty} \sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \mathcal{W}[D\varphi(x_j)] \left(\frac{x - x_j}{r_n} \right) \right|^2 dx \leq C \lambda^{\frac{10}{3} + \frac{1}{r}} \eta^{-10 + \frac{3}{s}} \|D\varphi\|_{L^{2s}(\mathcal{O})}^2.$$

We can take any $s > 1$, which yields by setting p such that $p' = 2s$: for any $p < 2$

$$\limsup_{n \rightarrow \infty} \sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \mathcal{W}[D\varphi(x_j)] \left(\frac{x - x_j}{r_n} \right) \right|^2 dx \leq C \lambda^{\frac{10}{3} + \frac{2-p}{p}} \eta^{-4 - \frac{6}{p}} \|D\varphi\|_{L^{2s}(\mathcal{O})}^2. \quad (30)$$

To treat the first term in the decomposition (29), we write

$$\mathcal{V}[D\varphi(x_j)] \left(\frac{x - x_j}{r_n} \right) = r_n^3 \mathcal{M}(x - x_j) D\varphi(x_j)$$

for \mathcal{M} a matrix-valued Calderon-Zygmund operator.

We use that for all i and all $j \neq i, j \in \mathcal{G}_\eta$ we have for all $(x, y) \in B_i \times B_j^*$

$$|\mathcal{M}(x - x_j) - \mathcal{M}(x - y)| \leq C \eta n^{-1/3} |x - y|^{-4}$$

Thus, by similar manipulations as before

$$\begin{aligned} & \sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \mathcal{V}[D\varphi(x_j)] \left(\frac{x - x_j}{r_n} \right) \right|^2 dx \\ & \leq C r_n^6 \sum_i \int_{B_i} \left(\sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \frac{1}{|B_j^*|} \int_{B_j^*} \mathcal{M}(x - y) 1_{\{|x-y| > c\eta n^{-\frac{1}{3}}\}}(x - y) |D\varphi(x_j)| dy \right)^2 dx \\ & + C \frac{\eta^2}{n^{2/3}} r_n^6 \sum_i \int_{B_i} \left(\sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \frac{1}{|B_j^*|} \int_{B_j^*} |x - y|^{-4} 1_{\{|x-y| > c\eta n^{-\frac{1}{3}}\}}(x - y) |D\varphi(x_j)| dy \right)^2 dx \\ & \leq C n^2 \frac{r_n^6}{\eta^6} \|1_{\cup B_i}\|_{L^r} \|(\mathcal{M}(x) 1_{\{|x| > \eta n^{-\frac{1}{3}}\}}) \star \sum_{1 \leq j \leq n} |D\varphi(x_j)| 1_{B_j^*}\|_{L^{2s}}^2 \\ & + C n^2 \frac{\eta^2}{n^{2/3}} \frac{r_n^6}{\eta^6} \|1_{\cup B_i}\|_{L^r} \| |x|^{-4} 1_{\{|x| > c\eta n^{-\frac{1}{3}}\}} \|_{L^1}^2 \left\| \sum_{1 \leq j \leq n} |D\varphi(x_j)| 1_{B_j^*} \right\|_{L^{2s}}^2 \end{aligned}$$

As seen in [8, Lemma 2.4], the kernel $\mathcal{M}(x) 1_{\{|x| > c\eta n^{-\frac{1}{3}}\}}$ defines a singular integral that is continuous over L^t for any $1 < t < \infty$, with operator norm bounded independently of the value $\eta n^{-\frac{1}{3}}$ (by scaling considerations). Applying this continuity property with $t = 2s$, writing as before $p' = 2s$, we get for all $p < 2$,

$$\limsup_{n \rightarrow \infty} \sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} \mathcal{V}[D\varphi(x_j)] \left(\frac{x - x_j}{r_n} \right) \right|^2 dx \leq C \lambda^{2 + \frac{2-p}{p}} \eta^{-\frac{6}{p}} \|D\varphi\|_{L^{2s}(\mathcal{O})}^2$$

Combining this last inequality with (29) and (30), we finally get: for all $p < 2$,

$$\limsup_{n \rightarrow \infty} \left(\sum_i \int_{B_i} \left| \sum_{\substack{j \neq i, \\ j \in \mathcal{G}_\eta}} D\phi_{j,n} \right|^2 \right)^{1/2} \leq C' \lambda^{1 + \frac{2-p}{2p}} \eta^{-\frac{3}{p}} \|D\varphi\|_{L^{p'}(\mathcal{O})} \quad (31)$$

Finally, if we inject (28) and (31) in (26), we obtain that for any $p < 2$,

$$\limsup_{n \rightarrow \infty} \|\nabla \psi_n^1\|_{L^2(\mathbb{R}^3)} \leq C \left(\lambda^{\frac{1}{2} + \frac{1}{p}} + \lambda^{1 + \frac{2-p}{2p}} \eta^{-\frac{3}{p}} \right) \|D\varphi\|_{L^{p'}(\mathbb{R}^3)} \quad (32)$$

The final step in the proof of Proposition 4 is to establish bounds on ψ_n^3 . This term expresses the effect of the balls that are close to one another, and to control it will require the sharp estimates of Theorem 6. Let M the constant in assumption (B1), and $\varepsilon_M > 0$ given in point i) of Theorem 6. We fix once for all $p_0 \in (2 - \varepsilon_M, 2)$ and $t_0 \in (p_0, 2)$. Set q_0 such that $\frac{1}{t_0} + \frac{1}{q_0} = \frac{1}{p_0}$. Let $1 < p < (q_0)'$, $q < p'$, and M_q as given in point ii) of Theorem 6. We set $\eta_q = \max(M_{q_0}, M_q) \left(\frac{3}{4\pi} \lambda\right)^{\frac{1}{3}}$, so that $\eta_q n^{-\frac{1}{3}} = \max(M_{q_0}, M_q) r_n$. We restrict here to $\eta \geq \eta_q$, a condition that will be guaranteed by our choice of η . We further divide

$$\mathcal{B}_\eta = \left(\mathcal{B}_\eta \cap \mathcal{G}_{\eta_q} \right) \cup \left(\mathcal{B}_\eta \cap \mathcal{B}_{\eta_q} \right) = \mathcal{B}' \cup \mathcal{B}''.$$

and correspondingly: $\psi_n^3 = \psi' + \psi''$, where

$$-\mu \Delta \psi' + \nabla p' = 0, \quad \operatorname{div} \psi' = 0, \quad x \in \mathbb{R}^3 \setminus \cup_{i \in \mathcal{G}_\eta \cup \mathcal{B}'} B_i$$

and

$$D\psi'|_{B_i} = 0, \quad \forall i \in \mathcal{G}_\eta, \quad D\psi'|_{B_i} = D\varphi|_{B_i}, \quad i \in \mathcal{B}'$$

plus no torque and no force at balls B_i , $i \in \mathcal{G}_\eta \cup \mathcal{B}'$. We insist that balls B_i with $i \in \mathcal{B}''$ are completely left aside in the definition of ψ' . The point is that for all $\eta \geq \eta_q$, for all $i \neq j \in \mathcal{G}_\eta \cup \mathcal{B}'$, $|x_i - x_j| \geq M_q r_n$. We can apply point ii) of Theorem 6 to deduce

$$\begin{aligned} \|\nabla \psi'\|_{L^q(\mathbb{R}^3)} &\leq C \|D\varphi\|_{L^q(\cup_{i \in \mathcal{B}'} B_i)} \leq C \|1_{\cup_{i \in \mathcal{B}'} B_i}\|_{L^r(\mathbb{R}^3)} \|D\varphi\|_{L^s(\cup_{i \in \mathcal{B}'} B_i)} \\ &\leq C' \left(\operatorname{card} \mathcal{B}' r_n^3 \right)^{\frac{1}{r}} \|D\varphi\|_{L^s(\cup_{i \in \mathcal{B}'} B_i)} \end{aligned}$$

for all r, s with $\frac{1}{r} + \frac{1}{s} = \frac{1}{q}$. Moreover, by assumption (B2), $\operatorname{card} \mathcal{B}' \leq \operatorname{card} \mathcal{B}_\eta \leq C \eta^3 n$. Finally, taking $s = p'$, we find that for all $p > 1$, for all $q < p'$, and all $\eta > \eta_q$:

$$\|\nabla \psi'\|_{L^q(\mathbb{R}^3)} \leq C (\eta^3 \lambda)^{\frac{1}{r}} \|D\varphi\|_{L^{p'}(\mathbb{R}^3)}, \quad \frac{1}{r} + \frac{1}{p'} = \frac{1}{q}. \quad (33)$$

A similar reasoning holds with q_0 instead of q . As $p < (q_0)'$, we find that for r_0 such that $\frac{1}{r_0} + \frac{1}{p'} = \frac{1}{q_0}$,

$$\|\nabla \psi'\|_{L^{q_0}(\mathbb{R}^3)} \leq C (\eta^3 \lambda)^{\frac{1}{r_0}} \|D\varphi\|_{L^{p'}(\mathbb{R}^3)}. \quad (34)$$

Eventually, the last term ψ'' satisfies

$$-\mu \Delta \psi'' + \nabla p'' = 0, \quad \operatorname{div} \psi'' = 0, \quad x \in \Omega_n$$

and

$$D\psi''|_{B_i} = 0, \quad \forall i \in \mathcal{G}_\eta \cup \mathcal{B}', \quad D\psi''|_{B_i} = D\varphi|_{B_i} - D\psi'|_{B_i}, \quad i \in \mathcal{B}''$$

plus no force and no torque at all balls. Here, we can only rely on assumption (B1) and point i) of Theorem 6: we find

$$\begin{aligned} \|\nabla\psi''\|_{L^{p_0}(\mathbb{R}^3)} &\leq C\left(\|D\varphi\|_{L^{p_0}(\cup_{i\in\mathcal{B}''}B_i)} + \|D\psi'\|_{L^{p_0}(\cup_{i\in\mathcal{B}''}B_i)}\right) \\ &\leq C\|1_{\cup_{i\in\mathcal{B}''}B_i}\|_{L^{t_0}(\mathbb{R}^3)}\left(\|D\varphi\|_{L^{q_0}(\cup_{i\in\mathcal{B}'}B_i)} + \|D\psi'\|_{L^{q_0}(\mathbb{R}^3)}\right) \end{aligned}$$

as $\frac{1}{t_0} + \frac{1}{q_0} = \frac{1}{p_0}$. By assumption (B2), $\text{card}\mathcal{B}'' \leq C\eta_q^3 n \leq C'_q \lambda n$. Hence,

$$\|\nabla\psi''\|_{L^{p_0}(\mathbb{R}^3)} \leq C(\lambda^2)^{\frac{1}{t_0}} \left(\|D\varphi\|_{L^{q_0}(\cup_{i\in\mathcal{B}'}B_i)} + \|D\psi'\|_{L^{q_0}(\mathbb{R}^3)} \right)$$

Remember that $t_0 < 2$. Combining with (34), we see that there exists $\theta_0 > 0$ such that for $p < (q_0)'$,

$$\|\nabla\psi''\|_{L^{p_0}(\mathbb{R}^3)} \leq \lambda^{1+\theta_0} \|D\varphi\|_{L^{p'}(\mathbb{R}^3)}$$

Combining with (33), we arrive at: for all $1 < p < (q_0)'$, for all $q < p'$, and r such that $\frac{1}{r} + \frac{1}{p'} = \frac{1}{q}$,

$$\|\nabla\psi_n^3\|_{L^{p_0}(\mathbb{R}^3)+L^q(\mathbb{R}^3)} \leq C\left((\eta^3\lambda)^{\frac{1}{r}} + \lambda^{1+\theta_0}\right) \|D\varphi\|_{L^{p'}(\mathbb{R}^3)} \quad (35)$$

Proposition 4 follows from collecting (25), (32) and (35).

5 Discussion of assumption (B2)

Let $\Phi^\delta = \{y_i\}_i \subset \mathbb{R}^3$ be a stationary ergodic point process on \mathbb{R}^3 with intensity δ and hard-core radius R , i.e., $|y_i - y_j| \geq R$ for all $i \neq j$. An example of such a process is a hard-core Poisson point process, which is obtained from a Poisson point process upon deleting all points with a neighboring point closer than R . We refer to [3, 4] for the construction and properties of such processes.

Assume that \mathcal{O} is convex and contains the origin. For $\varepsilon > 0$, we consider the set

$$\varepsilon\Phi^\delta \cap \mathcal{O} =: \{x_i^\varepsilon, i = 1, \dots, n_\varepsilon\}.$$

Let $r < R/2$ and denote $r_\varepsilon = \varepsilon r$ and consider $B_i = \overline{B(x_i, r_\varepsilon)}$. The volume fraction of the particles depends on ε in this case. However, it is not difficult to generalize our result to the case when the volume fraction converges to λ and this holds in the setting under consideration since

$$\frac{4\pi}{3} n_\varepsilon r_\varepsilon^3 \rightarrow \frac{4\pi}{3} \delta r^3 =: \lambda(r, \delta) \quad \text{almost surely as } \varepsilon \rightarrow 0.$$

Clearly, $\lambda(r, \delta) \rightarrow 0$, both if $r \rightarrow 0$ and if $\delta \rightarrow 0$. However, the process behaves fundamentally different in those cases. Indeed, if we take $r \rightarrow 0$ (for δ and R fixed), we find that condition (A1), which implies (B2), is satisfied almost surely for ε sufficiently small as

$$n_\varepsilon^{1/3} |x_i^\varepsilon - x_j^\varepsilon| \geq n_\varepsilon^{1/3} \varepsilon R \rightarrow \delta^{1/3} R.$$

In the case when we fix r and consider $\delta \rightarrow 0$ (e.g. by randomly deleting points from a process), (A1) is in general not satisfied. We want to characterize processes for which (B2) is still fulfilled almost surely as $\varepsilon \rightarrow 0$. Indeed, using again the relation between ε and n_ε , it suffices to show

$$\forall \eta > 0, \quad \#\{i, \exists j, |x_i - x_j| \leq \eta\varepsilon\} \leq C\eta^3 \delta^2 \varepsilon^{-3}. \quad (36)$$

Let Φ_η^δ be the process obtained from Φ^δ by deleting those points y with $B(y, \eta) \cap \Phi^\delta = \{y\}$. Then, the process Φ_η^δ is again stationary ergodic (since deleting those points commutes with translations¹), so that almost surely as $\varepsilon \rightarrow 0$

$$\varepsilon^3 \#\{i, \exists j, |x_i - x_j| \leq \eta\varepsilon\} \rightarrow \mathbb{E}[\#\Phi_\eta^\delta \cap Q],$$

where $Q = [0, 1]^3$. Clearly,

$$\mathbb{E}[\#\Phi_\eta^\delta \cap Q] \leq \mathbb{E} \sum_{y \in \Phi^\delta \cap Q} \sum_{y' \neq y \in \Phi^\delta} 1_{B(0, \eta)}(y' - y).$$

We can express this expectation in terms of the 2-point correlation function $\rho_2^\delta(y, y')$ of Φ^δ yielding

$$\mathbb{E}[\#\Phi_\eta^\delta \cap Q] \leq \int_{\mathbb{R}^6} 1_Q(y) 1_{B(0, \eta)}(y' - y) \rho_2^\delta(y, y') \, dy \, dy'.$$

Hence, (36) and therefore also (B2) is in particular satisfied if $\rho_2^\delta \leq C\delta^2$ which is the case for a (hard-core) Poisson point process.

Acknowledgement

The first author acknowledges the support of the Institut Universitaire de France, and of the SingFlows project, grant ANR-18-CE40-0027 of the French National Research Agency (ANR)

The second author has been supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the collaborative research center “The Mathematics of Emerging Effects” (CRC 1060, Projekt-ID 211504053) and the Hausdorff Center for Mathematics (GZ 2047/1, Projekt-ID 390685813).

References

- [1] Y. Almog and H. Brenner. Global homogenization of a dilute suspension of spheres.
- [2] G. Batchelor and J. Green. The determination of the bulk stress in a suspension of spherical particles at order c^2 . *J. Fluid Mech.*, 56:401–427, 1972.
- [3] D. J. Daley and D. Vere-Jones. *An introduction to the theory of point processes. Vol. I.* Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2003. Elementary theory and methods.
- [4] D. J. Daley and D. Vere-Jones. *An introduction to the theory of point processes. Vol. II.* Probability and its Applications (New York). Springer, New York, second edition, 2008. General theory and structure.
- [5] M. Duerinckx and A. Gloria. Corrector equations in fluid mechanics: Effective viscosity of colloidal suspensions. arXiv:1909.09625.
- [6] A. Einstein. Eine neue bestimmung der moleküldimensionen. *Ann. Physik.*, 19:289–306, 1906.

¹In detail: let \mathcal{E}_η be the operator that erases all points without a neighboring point closer than η , and let T_x denote a translation by x . Now, let μ be the measure for the original process Φ^δ . Then the measure for Φ_η^δ is given by $\mu_\eta = \mu \circ \mathcal{E}_\eta^{-1}$. Since $\mathcal{E}_\eta T_x = T_x \mathcal{E}_\eta$ (for all x , in particular for $T_{-x} = T_x^{-1}$), we have for any measurable set A that $T_x \mathcal{E}_\eta^{-1} A = \mathcal{E}_\eta^{-1} T_x A$. This immediately implies that the new process adopts stationarity and ergodicity.

- [7] D. Gérard-Varet. A simple justification of effective models for conducting or fluid media with dilute spherical inclusions
- [8] D. Gérard-Varet and M. Hillairet. Analysis of the viscosity of dilute suspensions beyond Einstein’s formula. arXiv:1905.08208, May 2019.
- [9] E. Guazelli and O. Pouliquen. Rheology of dense granular suspensions. *Journal of Fluid Mechanics*, Cambridge University Press (CUP), 2018, 852
- [10] B. M. Haines and A. L. Mazzucato. A proof of Einstein’s effective viscosity for a dilute suspension of spheres. *SIAM J. Math. Anal.*, 44(3):2120–2145, 2012.
- [11] M. Hillairet and D. Wu. Effective viscosity of a polydispersed suspension. Preprint arXiv:1905.12306, 2019.
- [12] R. M. Höfer. Convergence of the method of reflections for particle suspensions in Stokes flows. Preprint arXiv:1912.04388, 2019.
- [13] V. V. Jikov, S. M. Kozlov, and O. A. Oleĭnik. *Homogenization of differential operators and integral functionals*. Springer-Verlag, Berlin, 1994. Translated from the Russian by G. A. Yosifian.
- [14] T. Lévy and E. Sánchez-Palencia. Einstein-like approximation for homogenization with small concentration. II. Navier-Stokes equation. *Nonlinear Anal.*, 9(11):1255–1268, 1985.
- [15] B. Niethammer and R. Schubert. A local version of Einstein’s formula for the effective viscosity of suspensions. arXiv:1903.08554.
- [16] E. Sánchez-Palencia. Einstein-like approximation for homogenization with small concentration. I. Elliptic problems. *Nonlinear Anal.*, 9(11):1243–1254, 1985.