

Gravitational Waves in Axion Dark Matter

Chong-Sun Chu^{*,**}, Jiro Soda[‡], and Daiske Yoshida[‡]

**Physics Division, National Center for Theoretical Sciences,
National Tsing-Hua University, Hsinchu, 30013, Taiwan*

***Department of Physics, National Tsing-Hua University, Hsinchu 30013, Taiwan*

‡Department of Physics, Kobe University, Kobe 657-8501, Japan

(Dated: February 13, 2020)

Abstract

Axion dark matter is interesting as it allows a natural coupling to the gravitational Chern-Simons term. In the presence of an axion background, the gravitational Chern-Simons term produces parity violating effects in the gravitational sector, in particular on the propagation of gravitational waves. Previously, it has been shown that the coherent oscillation of the axion field leads to a parametric amplification of gravitational waves with a specific frequency. In this paper we focus on the parity violating effects of the Chern-Simon coupling and show the occurrence of gravitational birefringence. We also find deviation from the speed of light of the velocity of the gravitational waves. We give constraints on the axion-Chern-Simons coupling constant and the abundance of the axion dark matter from the observation of GW170817 and GRB170817A.

PACS numbers: 04.50.Kd

I. INTRODUCTION

The direct detection of gravitational waves in 2015 has widened the frontiers in research in fundamental physics [1]. Indeed, we are now in the era of gravitational wave astrophysics and multi-messenger astronomy and it is now possible to use gravitational wave signals to discriminate the various models beyond the standard model of particle physics as well as of cosmology. Moreover, the discovery of gravitational waves GW170817 [2] from a neutron star binary has sparked new interests in the study of nuclear physics. Remarkably, the observation of the optical counterpart of GW170817, GRB170817A [3], has given a constraint on the velocity of gravitational waves, which killed many modified theories of gravity [4–6].

In gravitational physics, there are three processes to be studied, namely, the production, propagation and detection of gravitational waves. The production process and detection process have been well studied. On the other hand, the propagation process has been mostly regarded as a trivial problem. In fact, in a Minkowski background, the gravitational wave equation is merely a conventional scalar wave equation. Even in the presence of the conventional matter, it is easy to solve the propagation problem of gravitational waves in the curved background. However, things can get more interesting with axions. As a pseudoscalar, coupling to a gravitational Chern-Simons term is allowed [7–9]. As a result, parity symmetry is broken in the presence of an axion background. A peculiar feature of the axion dark matter is coherent oscillations of the axion field, which may affect the propagation of electromagnetic waves [10] and gravitational waves [11, 12].

According to string theory, axions are ubiquitous in the universe [13]. Remarkably, the mass of string axions can take values in the broad range from 10^{-33} eV to 10^{18} GeV. In fact, it has been known that the axion is a natural candidate of an inflaton and induces circularly polarized gravitational waves [14, 15]. Recently, an axion has been intensively studied as a candidate for the dark matter. As the dark matter, we can consider the axion with mass from 10^{-23} eV to 10^3 eV. The lower bound comes from observations of cosmic background radiations and the upper bound comes from observations of X-ray backgrounds [16].

In this paper, we comprehensively investigate gravitational waves propagating in an axion background. First of all, we review the results of the previous work on the parametric resonance of gravitational waves [11, 12]. Then, we focus on the gravitational birefringence and the velocity modulation of gravitational waves. In particular, we discuss constraints on

the Chern-Simons coupling and the abundance of the axion dark matter from observations of the velocity of gravitational waves.

The organization of the paper is as follows. In section II, we introduce basic equations for gravitational waves in axion-Chern-Simons gravity. We also discuss a potential ghost mode and a cutoff scale that is needed in order to avoid the occurrence of this un-physical feature. In section III, we analysis the propagation of gravitational waves in a background of coherently oscillating axions. In section IV, we review the parametric amplification of gravitational waves and present consistency checks. In section V, we consider the gravitational birefringence, which can be regarded as a gravitational Faraday rotation. In section VI, we derive the velocity of gravitational waves. We discuss constraints on the Chern-Simons coupling constant and the abundance of the axion dark matter using the observation of the velocity of gravitational waves. The final section is devoted to the conclusion.

II. GRAVITATIONAL WAVES IN DYNAMICAL CHERN-SIMONS GRAVITY

Let us consider the action of dynamical Chern-Simons gravity

$$S = \kappa \int d^4x \sqrt{-g} R - \int d^4x \sqrt{-g} \left[\frac{1}{2} \nabla^\mu \Phi \nabla_\mu \Phi + V(\Phi) \right] + S_{\text{dCS}}, \quad (1)$$

where the first term of action is the Einstein-Hilbert term, g is the determinant of the metric $g_{\mu\nu}$ and $\kappa = 1/(16\pi G)$. The second term describes an action of an axion field Φ . In this paper we will take Φ as a dark matter candidate and exploit existing observational constraints on its mass. The last term in (1)

$$S_{\text{dCS}} = \frac{1}{4} \int d^4x \sqrt{-g} F(\Phi) R \tilde{R}, \quad R \tilde{R} := \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} R_{\alpha\beta\rho\sigma} R_{\gamma\delta}{}^{\rho\sigma} \quad (2)$$

is the dynamical Chern-Simons action with $\epsilon^{\alpha\beta\gamma\delta}$ being the Levi-Civita tensor density. We have allowed a nontrivial coupling $F(\Phi)$ of the axion field to the $R \tilde{R}$ Chern-Simons term. Otherwise, the dynamical Chern-Simons term is topological and won't affect the equation of motion.

We are interested in the effect of the dark matter Chern-Simons coupling on the propagation of gravitational wave. Before we start, we need to fix the background. Let us consider a background spacetime with spatial isotropy and homogeneity

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j. \quad (3)$$

Due to its structure, the Chern-Simons term does not contribute to the equation of motion of the isotropic and homogeneous universe. As a result, we have the equations of motion,

$$3H^2 = \frac{1}{2\kappa} \left(\frac{1}{2} \dot{\Phi}^2 + V(\Phi) \right) \quad (4)$$

$$\dot{H} + 3H^2 = \frac{1}{2\kappa} V(\Phi) \quad (5)$$

$$\ddot{\Phi} + 3H\dot{\Phi} + V'(\Phi) = 0, \quad (6)$$

where a dot denotes a time derivative and $H = \dot{a}/a$ is the Hubble parameter. Generally the dark matter has a mass

$$V(\Phi) = \frac{1}{2} m^2 \Phi^2. \quad (7)$$

We will ignore self interaction as it is not relevant for our analysis.

A. Action for gravitational waves

Let us now derive the quadratic action for the gravitational waves from the action (1). The tensor perturbation reads

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -dt^2 + a^2(t) (\delta_{ij} + h_{ij}) dx^i dx^j, \quad (8)$$

where h_{ij} satisfies the transverse-traceless conditions $h_{ij,j} = h_{ii} = 0$. Substituting the metric into (1), we obtain the quadratic action

$$S = \frac{\kappa}{4} \int dt d^3x a^3 \left[\dot{h}^{ij} \dot{h}_{ij} - \frac{1}{a^2} h^{ij|k} h_{ij|k} + \frac{\dot{F}}{\kappa a} \epsilon^{ijk} \left(\dot{h}_{ai} \dot{h}_{k|j}^a - \frac{1}{a^2} h_{ai|b} h^a{}_{k|j}{}^b \right) \right], \quad (9)$$

where the stroke $|$ denotes a covariant derivative with respect to the spatial coordinates. Here, we have used the convention $\epsilon^{0ijk} = -\epsilon^{ijk}$ with $\epsilon^{123} \equiv 1$. It is convenient to expand h_{ij} in terms of circular polarization basis

$$h_{ij}(\eta, \mathbf{x}) = \sum_{A=R,L} \int \frac{d^3k}{(2\pi)^3} h_A(t, \mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} p_{ij}^A, \quad (10)$$

where p_{ij}^A are the circular polarization tensor defined by

$$p_{ij}^R \equiv \frac{1}{\sqrt{2}} (p_{ij}^+ + ip_{ij}^\times), \quad p_{ij}^L \equiv \frac{1}{\sqrt{2}} (p_{ij}^+ - ip_{ij}^\times) \quad (11)$$

and the polarization tensors p_{ij}^+ and p_{ij}^\times are the plus and cross modes respectively. The polarization tensors are normalized as

$$p_{ij}^{*A} p_{ij}^B = 2\delta^{AB}, \quad A, B = \text{R, L} \quad (12)$$

and satisfies the helicity condition

$$i\epsilon_i^{sj} \hat{k}_s p_{jk}^A = \rho_A p_{ik}^A \quad \text{with} \quad \rho_{\text{R}} = +1, \quad \rho_{\text{L}} = -1, \quad (13)$$

where $\hat{k}^i := k^i/k$ is the unit vector in the direction of propagation of the gravitational wave. The circular polarization modes are special as they diagonalize the dynamical Chern-Simons action term and we obtain the quadratic action (9)

$$S = \frac{\kappa}{2} \sum_{A=\text{R,L}} \int dt \int \frac{d^3k}{(2\pi)^3} a^3 B_A \left(\dot{h}_A^2 - \frac{k^2}{a^2} h_A^2 \right). \quad (14)$$

where the function B_A is defined by

$$B_A(t, k) := 1 - \frac{k\rho_A \dot{F}}{\kappa a}. \quad (15)$$

In order for the perturbation to be stable, it is required that

$$B_A > 0 \quad (16)$$

for both polarizations A . One can derive from this a natural upper bound to the energy scale of the gravitational wave in the theory [17]. This is in contrast to the parity violating gravitational waves in Lorentz violating gravity [18].

III. GRAVITATIONAL WAVES PROPAGATING IN AXION DARK MATTER

In this section, we suppose that a source in our cosmological horizon emits gravitational waves propagating in the axion dark matter background. In this situation, we can neglect the effect of cosmic expansion of the Universe in the dynamical equation (6) of the axion. To see this, let us note that the Friedmann equation (4) set the Hubble parameter to be of the order of $H \sim \sqrt{\rho/\kappa} \sim \sqrt{\rho}/M_{\text{pl}}$, where $M_{\text{pl}} := \sqrt{2\kappa}$ is the reduced Planck mass. On the other hand, the scalar field changes at a rate determined by the mass scale: $\dot{\Phi} \sim m$. The cosmological expansion in (6) is negligible if

$$H/m \sim \sqrt{\rho}/(mM_{\text{pl}}) \ll 1. \quad (17)$$

Since an upper bound of the dark matter density is given by

$$\rho = 0.3\text{GeV}/\text{cm}^3 = 2.3 \times 10^{-6}\text{eV}^4, \quad (18)$$

the condition (17) is always satisfied for the axion dark matter with a mass $m > 10^{-30}$ eV.

For simplicity, let us consider a linear dynamical Chern-Simon coupling

$$F(\Phi) = \alpha\Phi \quad (19)$$

Following [19], we express the Chern-Simon coupling constant α in terms of a length ℓ as

$$\alpha = \frac{M_{\text{pl}}}{2}\ell^2, \quad (20)$$

where $M_{\text{pl}} = \sqrt{2\kappa}$ is the reduced Planck mass. The coupling constant ℓ is experimentally constrained by the Gravity Probe B as [21]

$$\ell \leq 10^8\text{km}. \quad (21)$$

Now, we can write down the condition (16) in the present context. The axion satisfies the equation of motion

$$\ddot{\Phi} + m^2\Phi = 0. \quad (22)$$

This can be solved as

$$\Phi = \Phi_0 \cos(mt), \quad (23)$$

where, without loss of generality, we have made a choice of time so that the phase in (23) is zero. In this case, the no-ghost condition gives

$$k < k_g := \frac{1}{\alpha m \Phi_0 / \kappa} = \frac{\kappa}{\alpha \sqrt{2\rho}} \quad (24)$$

where

$$\rho := \frac{1}{2}m^2\Phi_0^2 \quad (25)$$

is the energy density of the dark matter field. Once we used the observed energy density, the amplitude can be determined as

$$\Phi_0 \simeq 2.1 \times 10^7\text{eV} \left(\frac{10^{-10}\text{eV}}{m} \right) \sqrt{\frac{\rho}{0.3\text{GeV}/\text{cm}^3}}. \quad (26)$$

Finally, we can deduce the cutoff frequency f_g as

$$f_g \equiv \frac{k_g}{2\pi} = 1.1 \times 10^9\text{Hz} \left(\frac{10^8\text{km}}{\ell} \right)^2 \sqrt{\frac{0.3\text{GeV}/\text{cm}^3}{\rho}}. \quad (27)$$

Above this scale, we cannot use the Chern-Simons gravity to describe the propagation of gravitational waves in the axion dark matter.

On energy scales below this, the equation of motion of gravitational waves in the Chern-Simons gravity is given by

$$\ddot{h}_A + D_A \dot{h}_A + \frac{k^2}{a^2} h_A = 0 \quad (28)$$

where the function D_A is defined by

$$D_A(t, k) := 3H + \frac{\dot{B}_A}{B_A}. \quad (29)$$

The first term is due to cosmological expansion, which we can ignore. The second term is due to the dark matter background (23). It is convenient to introduce the dimensionless parameter

$$\delta := m^2 \alpha \Phi_0 / \kappa \quad (30)$$

in terms of which we have

$$B_A = 1 + \frac{\rho_A \delta}{m} k \sin(mt). \quad (31)$$

The parameter can be estimated as

$$\delta \simeq 2.3 \times 10^{-5} \left(\frac{m}{10^{-10} \text{eV}} \right) \left(\frac{\ell}{10^8 \text{km}} \right)^2 \sqrt{\frac{\rho}{0.3 \text{GeV/cm}^3}}. \quad (32)$$

In the following sections, we analyze the features of gravitational waves in the axion dark matter background.

IV. PARAMETRIC RESONANCES

Apparently, the dynamical Chern-Simons coupling induces the parity violation in the presence of the axion dark matter. In fact, the equations for each circular polarization modes are different. Generically, the polarization dependent effect in gravitational waves characterized by δ is small for typical values of model parameters. However, the effect can be exponentially enhanced due to the resonance.

To see this, let us introduce a new variable $\Psi_A(t, k)$ defined by

$$h_A(t, k) = \exp\left(-\frac{1}{2} \int^t D_A(t', k) dt'\right) \Psi_A(t, k). \quad (33)$$

Then the equation of motion for the gravitational wave becomes

$$\ddot{\Psi}_A + \omega_A^2 \Psi_A = 0, \quad (34)$$

where

$$\omega_A^2 := \frac{k^2}{a^2} - \frac{1}{2}\dot{D}_A - \frac{1}{4}D_A^2. \quad (35)$$

To the leading order of δ , the angular frequency is given by

$$\omega_A^2 = k^2 (1 + \rho_A f_0 \sin(mt)) \quad (36)$$

and eq.(34) takes the form of the Mathieu equation

$$\ddot{\Psi}_A + k^2 (1 + \rho_A f_0 \sin(mt)) \Psi_A = 0, \quad \text{where} \quad f_0 := \frac{1}{2} \frac{m}{k} \delta. \quad (37)$$

This describes an oscillator with a frequency k pumped by the polarization dependent periodic force with a magnitude f_0 and a frequency m . As is well known, the resonance occurs when

$$k \sim m/2. \quad (38)$$

In this case, we obtain

$$f_0 \sim \delta. \quad (39)$$

The amplitude of gravitational waves h_A grows exponentially $|h_A| \sim e^{\Gamma t}$ with the growth rate given by

$$\begin{aligned} \Gamma &= \frac{m\delta}{8} \\ &= 2.8 \times 10^{-16} \text{eV} \left(\frac{m}{10^{-10} \text{eV}} \right)^2 \left(\frac{\ell}{10^8 \text{km}} \right)^2 \sqrt{\frac{\rho}{0.3 \text{GeV/cm}^3}}. \end{aligned} \quad (40)$$

We can estimate the length $R_{\times 10}$ which the gravitational wave grows ten times bigger from the growth rate as follows,

$$\begin{aligned} R_{\times 10} &= \frac{8}{m\delta} \\ &= 5.2 \times 10^{-8} \text{pc} \left(\frac{10^{-10} \text{eV}}{m} \right)^2 \left(\frac{10^8 \text{km}}{\ell} \right)^2 \sqrt{\frac{0.3 \text{GeV/cm}^3}{\rho}}. \end{aligned}$$

The range of the wave number for the resonance is given by

$$\frac{m}{2} - \frac{m}{8}\delta \lesssim k \lesssim \frac{m}{2} + \frac{m}{8}\delta.$$

This corresponds to a resonance width Δk_{res} :

$$\Delta k_{\text{res}} = \frac{1}{4}m\delta. \quad (41)$$

The phenomenological consequence of this result has been discussed in a previous paper [11]. Recently, more serious comparison with gravitational observation are made in [20]

A. Coherence length

In the above analysis, we have ignored the coherence issue of the dark matter background. In principle, if the length scale of the gravitational perturbation becomes comparable to the Jeans length scale, the dark matter cloud can no longer remain homogeneous and gravitational collapse will occur. In other words, the coherence can be sustained only within the Jeans scale. As is known, the Jeans length r_J of the axion dark matter can be deduced as

$$\begin{aligned} r_J &= 6.7 \times 10^{20} \text{eV}^{-1} \left(\frac{m}{10^{-10} \text{eV}} \right)^{-\frac{1}{2}} \left(\frac{\rho}{0.3 \text{ GeV/cm}^3} \right)^{-\frac{1}{4}} \\ &= 4.3 \times 10^{-3} \text{pc} \left(\frac{m}{10^{-10} \text{eV}} \right)^{-\frac{1}{2}} \left(\frac{\rho}{0.3 \text{ GeV/cm}^3} \right)^{-\frac{1}{4}}. \end{aligned} \quad (42)$$

Thus, the condition $R_{\times 10} \leq r_J$ is necessary for the resonance to occur. This is satisfied for $m \geq m_c$ where the critical mass m_c is given by

$$m_c = 5.3 \times 10^{-14} \text{eV} \left(\frac{10^8 \text{km}}{\ell} \right)^{\frac{4}{3}} \left(\frac{0.3 \text{ GeV/cm}^3}{\rho} \right)^{\frac{1}{6}}. \quad (43)$$

B. One more consistency check

For having the resonance, we need one more condition. Axion dark matter within the dark matter halo has a virial velocity. From the simple dimensional analysis, one can estimate the virial velocity as

$$v_{\text{vir}} = 1.5 \times 10^{-11} \left(\frac{m}{10^{-10} \text{eV}} \right)^{-\frac{1}{2}} \left(\frac{\rho}{0.3 \text{ GeV/cm}^3} \right)^{\frac{1}{4}}.$$

This velocity induces fluctuations in the frequency of the axion dark matter given by

$$\Delta k_{\text{vir}} = \frac{1}{2}mv_{\text{vir}}^2 \quad (44)$$

If the fluctuations are larger than the band width Δk_{res} of the parametric resonance, the amplitude cannot grow efficiently. This is characterized by the ratio

$$\gamma := \frac{\Delta k_{\text{vir}}}{\Delta k_{\text{res}}}. \quad (45)$$

If γ satisfies

$$\gamma \ll 1, \quad (46)$$

the amplitude of gravitational waves grows. On the other hand, if γ satisfies

$$\gamma \geq 1, \quad (47)$$

then the frequency of the axion dark matter easily escape from the resonance band and the gravitational wave never grows. Since it is calculated as

$$\gamma = 2.0 \times 10^{-17} \left(\frac{m}{10^{-10}\text{eV}} \right)^{-2} \left(\frac{l}{10^8\text{km}} \right)^{-2}, \quad (48)$$

we see that the amplitude of gravitational waves grows.

V. GRAVITATIONAL FARADAY ROTATION

The fact that the angular frequency (36) is different for different circular polarization modes implies that the phase velocity

$$v_A^{(p)} := \frac{\omega_A}{k} = \sqrt{1 + \rho_A f_0 \sin(mt)}. \quad (49)$$

is different for different circular polarization modes. It also implies a phase shift between the R and L polarization arises as the wave propagates:

$$\Delta\phi = \int_0^t dt' (\omega_R - \omega_L) = \frac{\delta}{2} \cos(mt) + \mathcal{O}(\delta^2). \quad (50)$$

This is characterized by a period of phase oscillation

$$T \sim \left(\frac{m}{10^{-10}\text{eV}} \right)^{-1} \sim 10^{-6} \text{ s}. \quad (51)$$

The amplitude of the gravitational Faraday rotation is given by

$$(\Delta\phi)_{\text{max}} = \frac{\delta}{2} = 1.1 \times 10^{-5} \left(\frac{m}{10^{-10}\text{eV}} \right) \left(\frac{\ell}{10^8\text{km}} \right)^2 \sqrt{\frac{\rho}{0.3\text{GeV}/\text{cm}^3}}. \quad (52)$$

Hence, the gravitational Faraday rotation is sizable for axion with $m \gtrsim 10^{-5}$ eV.

VI. VELOCITY OF GRAVITATIONAL WAVES

The propagation of gravitational waves is characterized by the group velocity $v_A^{(g)} := \partial\omega_A/\partial k$. To the leading order of δ , we obtain

$$v_A^{(g)} = 1 + \frac{1}{32} \frac{m^2}{k^2} \delta^2 \sin^2(mt) + \mathcal{O}(\delta^3), \quad (53)$$

where we assumed $\delta \ll 1$ and $m\delta/k \ll 1$. Note that it is independent of polarization up to the second order of δ . The group velocity is always greater than the speed of light and has a maximal deviation of

$$\begin{aligned} \Delta c &:= \frac{1}{32} \frac{m^2}{k^2} \delta^2 \\ &= 9.4 \times 10^{-7} \left(\frac{100\text{Hz}}{f_{\text{gw}}} \right)^2 \left(\frac{\ell}{10^8\text{km}} \right)^4 \left(\frac{m}{10^{-10}\text{eV}} \right)^4 \left(\frac{\rho}{0.3\text{GeV/cm}^3} \right). \end{aligned} \quad (54)$$

The current upper bound of Δc coming from the observation of GW170817 and GRB170817A is

$$\Delta c \leq 5 \times 10^{-16}. \quad (55)$$

If we observe gravitational waves oscillating in the low frequency and the axion dark matter which have the heavier mass, we may give the stronger constraint on the coupling constant ℓ . Especially, if the gravitational waves through the core of Galaxy, the velocity of the gravitational waves will be modified strongly. For example, if we use gravitational waves which have the frequency about 1Hz and we assume the density of the axion dark matter is about 0.3GeV/cm^3 and the mass of the axion dark matter is about 10^{-10}eV , the constraint on the Chern-Simons coupling constant reads

$$\ell \leq 4.8 \times 10^4\text{km}. \quad (56)$$

Once we obtain this constraint, by observing at a lower frequency, say 10^{-4}Hz , we can further constrain the Chern-Simons coupling constant as

$$\ell \leq 4.8 \times 10^2\text{km}. \quad (57)$$

For the extreme case of $f_{\text{gw}} = 10^{-9}\text{Hz}$ and $m = 10^3\text{eV}$, we obtain the stringent constraint on the coupling constant

$$\ell \leq 1.5 \times 10^{-13}\text{km}. \quad (58)$$

On the other hand, assuming the coupling constant $\ell = 10^2$ km and $f_{\text{gw}} = 10^{-9}$ Hz, we obtain the constraint on the abundance of the axion with $m = 10^{-10}$ eV as

$$\Omega_{\text{axion}} < 3.4 \times 10^{-3} . \quad (59)$$

Thus, we see the gravitational waves can provide useful constraints to the Chern-Simons coupling constant and the abundance of the axion dark matter.

VII. CONCLUSION

We studied gravitational waves propagating in the axion dark matter. In the presence of the axion, it is natural to consider the coupling of the axion to the gravitational Chern-Simons term. Since the axion condensation violates the parity symmetry, there is a chance to observe parity violation effects in the gravity sector using gravitational waves. We found that the coherent oscillation of the axion field leads to the parametric amplification of gravitational waves with a specific frequency. We investigated the gravitational birefringence induced by the difference in the phase velocity of the different polarization modes. We also derived the group velocity of gravitational waves which is independent on the polarization at the leading order. In particular, we have given a constraint on the Chern-Simons coupling constant and the abundance of the axion dark matter from the observation of GW170817 and GRB170817A.

There are various ways to proceed. It is important to perform a comparison of our result with real data. It is interesting to study gravitational wave propagating in the ultralight vector dark matter [22, 23]. It is also possible to extend the analysis to other higher spin dark matter.

ACKNOWLEDGMENTS

J. S. would like to thank Sunghoon Jung for useful comments and TaeHun Kim and Yuko Urakawa for fruitful discussions. C.S.C. was supported in part by NCTS and the grant MOST 107-2119-M-007-014-MY3 of the Ministry of Science and Technology of Taiwan. J. S. was supported in part by JSPS KAKENHI Grant Numbers JP17H02894, JP17K18778, JP15H05895, JP17H06359, JP18H04589. J. S. was also supported by JSPS Bilateral Joint

Research Projects (JSPS-NRF collaboration) “String Axion Cosmology.” Discussions during the YITP workshop YITP-T-19-02 on “Resonant instabilities in cosmology” were useful for this work. D. Y. was supported by Grant-in-Aid for JSPS Research Fellow and JSPS KAKENHI Grant Numbers 17J00490.

- [1] B. P. Abbott *et al.* [LIGO Scientific and Virgo Collaborations], *Phys. Rev. Lett.* **116**, no. 6, 061102 (2016) doi:10.1103/PhysRevLett.116.061102 [arXiv:1602.03837 [gr-qc]].
- [2] B. P. Abbott *et al.* [LIGO Scientific and Virgo Collaborations], *Phys. Rev. Lett.* **119**, no. 16, 161101 (2017) doi:10.1103/PhysRevLett.119.161101 [arXiv:1710.05832 [gr-qc]].
- [3] B. P. Abbott *et al.* [LIGO Scientific and Virgo and Fermi-GBM and INTEGRAL Collaborations], *Astrophys. J.* **848**, no. 2, L13 (2017) doi:10.3847/2041-8213/aa920c [arXiv:1710.05834 [astro-ph.HE]].
- [4] A. Nishizawa, *Phys. Rev. D* **97**, no. 10, 104037 (2018) doi:10.1103/PhysRevD.97.104037 [arXiv:1710.04825 [gr-qc]].
- [5] S. Arai and A. Nishizawa, *Phys. Rev. D* **97**, no. 10, 104038 (2018) doi:10.1103/PhysRevD.97.104038 [arXiv:1711.03776 [gr-qc]].
- [6] A. Nishizawa and T. Kobayashi, *Phys. Rev. D* **98**, no. 12, 124018 (2018) doi:10.1103/PhysRevD.98.124018 [arXiv:1809.00815 [gr-qc]].
- [7] B. A. Campbell, M. J. Duncan, N. Kaloper and K. A. Olive, *Nucl. Phys. B* **351**, 778 (1991). doi:10.1016/S0550-3213(05)80045-8
- [8] A. Lue, L. M. Wang and M. Kamionkowski, *Phys. Rev. Lett.* **83**, 1506 (1999) [arXiv:astro-ph/9812088].
- [9] R. Jackiw and S. Y. Pi, *Phys. Rev. D* **68**, 104012 (2003) [arXiv:gr-qc/0308071].
- [10] D. Yoshida and J. Soda, *PTEP* **2018**, no. 4, 041E01 (2018) doi:10.1093/ptep/pty029 [arXiv:1710.09198 [hep-th]].
- [11] D. Yoshida and J. Soda, *Int. J. Mod. Phys. D* **27**, no. 09, 1850096 (2018) doi:10.1142/S0218271818500967 [arXiv:1708.09592 [gr-qc]].
- [12] J. Soda and D. Yoshida, *Galaxies* **5**, no. 4, 96 (2017). doi:10.3390/galaxies5040096
- [13] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper and J. March-Russell, *Phys. Rev. D* **81**, 123530 (2010) doi:10.1103/PhysRevD.81.123530 [arXiv:0905.4720 [hep-th]].

- [14] M. Satoh, S. Kanno and J. Soda, Phys. Rev. D **77**, 023526 (2008) doi:10.1103/PhysRevD.77.023526 [arXiv:0706.3585 [astro-ph]].
- [15] M. Satoh and J. Soda, JCAP **0809**, 019 (2008) doi:10.1088/1475-7516/2008/09/019 [arXiv:0806.4594 [astro-ph]].
- [16] D. J. E. Marsh, Phys. Rept. **643**, 1 (2016) doi:10.1016/j.physrep.2016.06.005 [arXiv:1510.07633 [astro-ph.CO]].
- [17] S. Dyda, E. E. Flanagan and M. Kamionkowski, Phys. Rev. D **86**, 124031 (2012) doi:10.1103/PhysRevD.86.124031 [arXiv:1208.4871 [gr-qc]].
- [18] T. Takahashi and J. Soda, Phys. Rev. Lett. **102**, 231301 (2009) doi:10.1103/PhysRevLett.102.231301 [arXiv:0904.0554 [hep-th]].
- [19] K. Yagi, N. Yunes and T. Tanaka, Phys. Rev. Lett. **109**, 251105 (2012) Erratum: [Phys. Rev. Lett. **116**, no. 16, 169902 (2016)] doi:10.1103/PhysRevLett.116.169902, 10.1103/PhysRevLett.109.251105 [arXiv:1208.5102 [gr-qc]].
- [20] S. Jung, T. Kim, J. Soda and Y. Urakawa, “Constraining axion dark matter with stimulated decays into chirping GW and light” .
- [21] Y. Ali-Haïmoud and Y. Chen, Phys. Rev. D **84**, 124033 (2011) doi:10.1103/PhysRevD.84.124033 [arXiv:1110.5329 [astro-ph.HE]].
- [22] K. Nakayama, JCAP **1910**, no. 10, 019 (2019) doi:10.1088/1475-7516/2019/10/019 [arXiv:1907.06243 [hep-ph]].
- [23] K. Nomura, A. Ito and J. Soda, arXiv:1912.10210 [gr-qc].