


Complexity of Combinatorial Matrix Completion With Diameter Constraints

Tomohiro Koana 

Technische Universität Berlin, Faculty IV, Algorithmics and Computational Complexity
koana@campus.tu-berlin.de

Vincent Froese 

Technische Universität Berlin, Faculty IV, Algorithmics and Computational Complexity
vincent.froese@tu-berlin.de

Rolf Niedermeier 

Technische Universität Berlin, Faculty IV, Algorithmics and Computational Complexity
rolf.niedermeier@tu-berlin.de

Abstract

We thoroughly study a novel and still basic combinatorial matrix completion problem: Given a binary incomplete matrix, fill in the missing entries so that the resulting matrix has a specified maximum diameter (that is, upper-bounding the maximum Hamming distance between any two rows of the completed matrix) as well as a specified minimum Hamming distance between any two of the matrix rows. This scenario is closely related to consensus string problems as well as to recently studied clustering problems on incomplete data.

We obtain an almost complete complexity dichotomy between polynomial-time solvable and NP-hard cases in terms of the minimum distance lower bound and the number of missing entries per row of the incomplete matrix. Further, we develop polynomial-time algorithms for maximum diameter three, which are based on Deza's theorem from extremal set theory. On the negative side we prove NP-hardness for diameter at least four. For the parameter number of missing entries per row, we show polynomial-time solvability when there is only one missing entry and NP-hardness when there can be at least two missing entries. In general, our algorithms heavily rely on Deza's theorem and the correspondingly identified sunflower structures pave the way towards solutions based on computing graph factors and solving 2-SAT instances.

2012 ACM Subject Classification Theory of computation → Parameterized complexity and exact algorithms; Theory of computation → Pattern matching

Keywords and phrases sunflowers, binary matrices, Hamming distance, stringology, consensus problems, complexity dichotomy, fine-grained complexity, combinatorial algorithms, graph factors, 2-Sat, Hamming radius

Funding *Tomohiro Koana*: Partially supported by the DFG project MATE (NI 369/17).

Acknowledgements We are grateful to Christian Komusiewicz for helpful feedback on an earlier version of this work and to Stefan Szeider for pointing us to the work on clustering incomplete data [7].

	1	1	0	1
	1	0	1	0
1	0	0	1	0
0		1	0	1

1	1	1	0	1
0	1	0	1	0
1	0	0	1	0
0	0	1	0	1

0	1	1	0	1
0	1	0	1	0
1	0	0	1	0
0	0	1	0	1

0	0	0	0	0
---	---	---	---	---

■ **Figure 1** An illustration of matrix completion problems with the input matrix (left). Missing entries (and their completions) are framed by thick lines. The middle matrix is a completion of diameter four and the right matrix is a completion of radius three with the center vector below. Note that missing entries in the same column might be filled with different values to meet the diameter constraint, whereas this is never necessary for the radius constraint.

1 Introduction

In combinatorial matrix completion problems one is given an incomplete matrix over a fixed alphabet with some missing entries, and the goal is to fill in the missing entries such that the resulting “completed matrix” (over the same alphabet) fulfills a desired property. Performing a parameterized complexity analysis, Ganian et al. [11] and Eiben et al. [7] recently contributed to this growing field by studying various desirable properties. More specifically, Ganian et al. [11] studied the two properties of minimizing the rank or of minimizing the number of distinct rows of the completed matrix. Eiben et al. [7] investigated clustering problems where one wants to partition the rows of the completed matrix into a given number of clusters of small radius or of small diameter. In addition, in companion work [16] we studied two cases of completing the matrix into one which has small (local) radius. The latter two papers rely on Hamming distance as a measure; in general, all considered matrix completion problems are NP-hard and thus the above three papers [7, 11, 16] mostly focused on parameterized complexity studies. In this work, we research on a desirable property closely related to achieving small radius, namely achieving diameter bounds. Doing so, we focus on the case of binary alphabet. For a matrix $\mathbf{S} \in \{0, 1\}^{n \times \ell}$, let $\gamma(\mathbf{S}) = \min_{i \neq i' \in [n]} d(\mathbf{S}[i], \mathbf{S}[i'])$ and $\delta(\mathbf{S}) = \max_{i \neq i' \in [n]} d(\mathbf{S}[i], \mathbf{S}'[i])$, where d denotes the Hamming distance. Specifically, we study the following core problem.

DIAMETER MATRIX COMPLETION (DMC)

Input: An incomplete matrix $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ and $\alpha \leq \beta \in \mathbb{N}$.

Question: Is there a completion $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ of \mathbf{S} with $\alpha \leq \gamma(\mathbf{T})$ and $\delta(\mathbf{T}) \leq \beta$?

Before further motivating the study of DMC, let us first consider an example (see Figure 1) that also illustrates significant differences between radius minimization [16] and diameter minimization (the latter referring to $\delta(\mathbf{T}) \leq \beta$ above).

Compare DMC with **CONSTRAINT RADIUS MATRIX COMPLETION** as studied in our companion work [16]:

CONSTRAINT RADIUS MATRIX COMPLETION (CONRMC)

Input: An incomplete matrix $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ and $r \in \mathbb{N}^n$.

Question: Is there a row vector $v \in \{0, 1\}^\ell$ with $d(v, \mathbf{S}[i]) \leq r[i]$ for all $i \in [n]$?

An important difference between DMC and CONRMC is that in DMC we basically have to compare all rows against each other, but in CONRMC we have to compare one “center row”

against all others. Indeed, this makes these two similarly defined problems quite different in many computational complexity aspects.

Now let us consider potential application scenarios where DMC may appear. It is a natural combinatorial matrix problems which may appear in the following contexts:

- In coding theory, one may want to “design” (by filling in the missing entries) codewords that are pairwise neither too close (parameter α in DMC) nor too far (parameter β in DMC) from each other.
- In computational biology, one may want to minimize the maximum distance of sequences in order to determine their degree of relatedness (thus minimizing β); missing entries refer to missing data points.¹
- In data science, each row may represent an entity with its attributes, and solving the DMC problem may fulfill some constraints with respect to the pairwise (dis)similarity of the completed entities.
- In stringology, DMC seems to constitute a new and natural problem, closely related to many intensively studied consensus problems [4, 14].

Somewhat surprisingly, although simple to define and well-motivated, in the literature there seems to be no systematic study of DMC and its computational complexity. The two closest studies are the work of Eiben et al. [7] and our companion work [16]. On the problem level, Eiben et al. [7] focus on clustering while we focus on only finding one cluster (that is, the whole resulting matrix with small diameter). However, other than Eiben et al. [7] with the lower-bound parameter α we also model the aspect of achieving a minimum pairwise distance (not only a maximum diameter); actually, one may say that we essentially combine their “dispersion” and diameter clustering problems (for the special case of a single cluster). Indeed, in this sense the problems are incomparable. We perform a more fine-grained study of special cases, identifying polynomial-time cases as well as already NP-hard cases; actually, we make significant steps towards a computational complexity dichotomy (polynomial-time solvable versus NP-hard), leaving fairly few cases open. While the focus of the previous works [7, 16] is on parameterized complexity studies, in this work we settle very basic questions on the DMC problem, relying on several combinatorial insights, including results from (extremal) combinatorics (most prominently, Deza’s theorem [6]). Indeed, we believe that exploiting sunflowers based on Deza’s theorem in combination with corresponding use of algorithms for graph factors and for 2-SAT is perhaps our contribution of highest technical interest. In this context, we also observe the phenomenon that the running time bounds that we can prove for odd values of α (the “lower bound for dissimilarity”) are significantly better than the ones for even values of α —indeed, for even values of α the running time exponentially depends on α while it is independent of α for odd values of α .


We survey our results in Figure 2, also spotting few remaining open cases in terms of polynomial-time versus NP-hard classification.

2 Preliminaries

For $m \leq n \in \mathbb{N}$, let $[m, n] := \{m, \dots, n\}$ and let $[n] := [1, n]$.

For a matrix $\mathbf{T} \in \{0, 1\}^{n \times \ell}$, we denote by $\mathbf{T}[i, j]$ the entry in the i -th row and j -th column ($i \in [n]$ and $j \in [\ell]$) of \mathbf{T} . We use $\mathbf{T}[i, :]$ (or $\mathbf{T}[i]$ in short) to denote the *row*

¹ Here, it would be particularly natural to also study the case of non-binary alphabets; however, most of our positive results probably only hold for binary alphabet.



(a) Complexity of DMC with respect to combinations of constant parameter values α and β .

(b) Complexity of DMC with respect to combinations of parameters k and $\beta - \alpha$.

■ **Figure 2** Overview of our results. Green denotes polynomial-time solvability and Red denotes NP-hardness. White cells indicate open cases.

vector $(\mathbf{T}[i, 1], \dots, \mathbf{T}[i, \ell])$ and $\mathbf{T}[:, j]$ to denote the *column vector* $(\mathbf{T}[1, j], \dots, \mathbf{T}[n, j])^T$. For subsets $I \subseteq [n]$ and $J \subseteq [\ell]$, we write $\mathbf{T}[I, J]$ to denote the submatrix containing only the rows in I and the columns in J . We abbreviate $\mathbf{T}[I, [\ell]]$ and $\mathbf{T}[[n], J]$ as $\mathbf{T}[I, :]$ (or $\mathbf{T}[I]$ for short) and $\mathbf{T}[:, J]$, respectively. We use the special character \square for a *missing* entry. A matrix $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ called *incomplete* if it contains a missing entry and *complete* otherwise. We say that $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ is a *completion* of $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ if either $\mathbf{S}[i, j] = \square$ or $\mathbf{S}[i, j] = \mathbf{T}[i, j]$ holds for all $i \in [n]$ and $j \in [\ell]$.

Let $u, w \in \{0, 1, \square\}^\ell$ be row vectors. Let $D(u, w) := \{j \in [\ell] \mid u[j] \neq w[j] \wedge u[j] \neq \square \wedge w[j] \neq \square\}$ be the set of column indices where u and w disagree (not considering positions with missing entries). The *Hamming distance* between u and w is $d(u, w) := |D(u, w)|$. Note that the Hamming distance obeys the triangle inequality $d(u, w) \leq d(u, v) + d(v, w)$ for a complete vector $v \in \{0, 1\}^\ell$. For a subset $J \subseteq [\ell]$, we also define $d_J(u, w) := d(u, w)$. Let $u', v', w' \in \{0, 1\}^\ell$ be complete row vectors. Then, it holds that $d(u', w') = |D(u', v') \Delta D(v', w')| = |D(u', v')| + |D(v', w')| - 2|D(u', v') \cap D(v', w')|$.

The binary operation $u \oplus v$ replaces the missing entries of u with the corresponding entries in v for $v \in \{0, 1\}^\ell$. We sometimes use string notation to represent row vectors, such as 001 for $(0, 0, 1)$.

3 Constant Diameter Bounds α and β

In this section we consider the special case (α, β) -DMC of DMC, where $\alpha \leq \beta$ are some fixed constants. We prove the results depicted in Figure 2a. To start with, we show the following simple linear-time special case which will subsequently be used several times.

► **Lemma 1.** DMC can be solved in linear time for matrices with a constant number ℓ of columns.

Proof. Consider a set $\mathcal{V} \subseteq \{0, 1\}^\ell$ in which the pairwise Hamming distances are all between α and β . We simply check whether each row vector in the input matrix can be completed to some row vector in \mathcal{V} in $O(n \cdot 2^\ell) = O(n)$ time. Since there are at most 2^{2^ℓ} (that is, constantly many) choices for \mathcal{V} , this procedure can be done in linear time. ◀

3.1 Polynomial time for $\alpha = 0$ and $\beta \leq 3$

As an entry point, we show that $(0, 1)$ -DMC is easily solvable. To this end, we call a column vector *dirty* if it contains both 0 and 1. Clearly, for $\alpha = 0$, we can ignore columns that are not dirty since they can always be completed without increasing the Hamming distances between rows. Hence, throughout this subsection, we assume that the input matrix contains only dirty columns. Now, any $(0, 1)$ -DMC instance is a **Yes**-instance if and only if there is at most one dirty column in the input matrix:

► **Lemma 2.** *A matrix $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ admits a completion $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ with $\delta(\mathbf{T}) \leq 1$ if and only if \mathbf{S} contains at most one dirty column.*

Proof. Suppose that \mathbf{S} contains two dirty columns $\mathbf{S}[j_0]$ and $\mathbf{S}[j_1]$ for $j_0 \neq j_1 \in [\ell]$. We claim that $\delta(\mathbf{T}) \geq 2$ holds for any completion \mathbf{T} of \mathbf{S} . Let $i \in [n]$. Then, there exist $i_0, i_1 \in [n]$ with $\mathbf{T}[i, j_0] \neq \mathbf{T}[i_0, j_0]$ and $\mathbf{T}[i, j_1] \neq \mathbf{T}[i_1, j_1]$. If $\delta(\mathbf{T}) \leq 1$, then we obtain $\mathbf{T}[i_0, j_1] = \mathbf{T}[i, j_1]$ and $\mathbf{T}[i_1, j_0] = \mathbf{T}[i, j_0]$. Now we have $d(\mathbf{T}[i_0], \mathbf{T}[i_1]) \geq 2$ because $\mathbf{T}[i_0, j_0] \neq \mathbf{T}[i_1, j_0]$ and $\mathbf{T}[i_0, j_1] \neq \mathbf{T}[i_1, j_1]$. The reverse direction follows easily. ◀

Lemma 2 implies that one can solve $(0, 1)$ -DMC in linear time. In the following, we extend this to a linear-time algorithm for $(0, 2)$ -DMC (Theorem 12) and a polynomial-time algorithm for $(0, 3)$ -DMC (Theorem 18).

For these algorithms, we make use of a concept from extremal set theory, known as Δ -systems [15]. We therefore consider matrices as certain set systems.

► **Definition 3.** *For a matrix $\mathbf{T} \in \{0, 1\}^{n \times \ell}$, let \mathcal{T} denote the set system $\{D(\mathbf{T}[i], \mathbf{T}[n]) \mid i \in [n-1]\}$. Moreover, for $x \in \mathbb{N}$, let \mathcal{T}_x denote the set system $\{D(\mathbf{T}[i], \mathbf{T}[n]) \mid i \in [n-1], d(\mathbf{T}[i], \mathbf{T}[n]) = x\}$.*

The set system \mathcal{T} contains the subsets (without duplicates) of column indices corresponding to the columns where the row vectors $\mathbf{T}[1], \dots, \mathbf{T}[n-1]$ differ from $\mathbf{T}[n]$. For given $\mathbf{T}[n]$, since we have binary alphabet this clearly determines the full matrix \mathbf{T} .

The concept of Δ -systems has previously been used to obtain efficient algorithms [8, 9, 7]. They are defined as follows:

► **Definition 4 (Weak Δ -system).** *A set family $\mathcal{F} = \{S_1, \dots, S_m\}$ is a weak Δ -system if there exists an integer $\lambda \in \mathbb{N}$ such that $|S_i \cap S_j| = \lambda$ for any pair of distinct sets $S_i, S_j \in \mathcal{F}$. The integer λ is called the intersection size of \mathcal{F} .*

► **Definition 5 (Strong Δ -system, Sunflower).** *A set family $\mathcal{F} = \{S_1, \dots, S_m\}$ is a strong Δ -system (or sunflower) if there exists a subset $C \subseteq S_1 \cup \dots \cup S_m$ such that $S_i \cap S_j = C$ for any pair of distinct sets $S_i, S_j \in \mathcal{F}$. We call the set C the core and the sets $P_i = S_i \setminus C$ the petals of \mathcal{F} .*

Clearly, every strong Δ -system is a weak Δ -system.

Our algorithms are based on the combinatorial property that under certain conditions the set system \mathcal{T} of a matrix \mathbf{T} with bounded diameter forms a strong Δ -system (which can be algorithmically exploited). We say that a family \mathcal{F} of sets is h -uniform if $|S| = h$ holds for each $S \in \mathcal{F}$. Deza [6] showed that an h -uniform weak Δ -system is a strong Δ -system if its cardinality is sufficiently large (more precisely, if $|\mathcal{F}| \geq h^2 - h + 2$). Moreover, Deza [5] also proved a stronger lower bound for uniform weak Δ -systems in which the intersection size is exactly half of the cardinality of each set.

► **Lemma 6 ([5, Théorème 1.1]).** *Let \mathcal{F} be a (2μ) -uniform weak Δ -system with intersection size μ . If $|\mathcal{F}| \geq \mu^2 + \mu + 2$, then \mathcal{F} is a strong Δ -system.*

We extend this result to the case in which the set size is odd.

► **Lemma 7.** *Let \mathcal{F} be a $(2\mu + 1)$ -uniform weak Δ -system.*

- (i) *If the intersection size of \mathcal{F} is $\mu + 1$ and $|\mathcal{F}| \geq \mu^2 + \mu + 3$, then \mathcal{F} is a strong Δ -system.*
- (ii) *If the intersection size of \mathcal{F} is μ and $|\mathcal{F}| \geq (\mu + 1)^2 + \mu + 3$, then \mathcal{F} is a strong Δ -system.*

Proof. (i) Let $S \in \mathcal{F}$ and let $\mathcal{F}' = \{T \Delta S \mid T \in \mathcal{F} \setminus \{S\}\}$. Here $T \Delta S$ denotes the symmetric difference $(T \setminus S) \cup (S \setminus T)$. Note that \mathcal{F}' is a 2μ -uniform weak Δ -system with intersection size μ :

- For each $T \in \mathcal{F} \setminus \{S\}$, we have $|T \Delta S| = |S \setminus T| + |T \setminus S| = 2\mu$.
- We show that $|(T \Delta S) \cap (U \Delta S)| = \mu$ for each distinct $T, U \in \mathcal{F} \setminus \{S\}$. We rewrite

$$\begin{aligned} & |(T \Delta S) \cap (U \Delta S)| \\ &= |((T \setminus S) \cup (S \setminus T)) \cap ((U \setminus S) \cup (S \setminus U))| \\ &= |((T \setminus S) \cap (U \setminus S)) \cup ((T \setminus S) \cap (S \setminus U)) \\ &\quad \cup ((S \setminus T) \cap (U \setminus S)) \cup ((S \setminus T) \cap (S \setminus U))| \\ &= |((T \cap U) \setminus S) \cup (S \setminus (T \cup U))| \\ &= |((T \cap U) \setminus S)| + |S \setminus (T \cup U)|. \end{aligned}$$

Here the third equality follows from $(T \setminus S) \cap (S \cap U) = (S \setminus T) \cap (U \setminus S) = \emptyset$. Let $\kappa = |S \cap T \cap U|$. Since $|S \cap T| = |S \cap U| = \mu + 1$, it follows that $|(S \cap T) \setminus U| = |(S \cap U) \setminus T| = \mu - \kappa + 1$. Thus, we obtain

$$\begin{aligned} |S \setminus (T \cup U)| &= |S| - |(S \cap T) \setminus U| - |(S \cap U) \setminus T| - |S \cap T \cap U| \\ &= (2\mu + 1) - (\mu - \kappa + 1) - (\mu - \kappa + 1) - \kappa = \kappa - 1. \end{aligned}$$

Moreover, we obtain

$$|((T \cap U) \setminus S)| = |T \cap U| - |S \cap T \cap U| = \mu - \kappa + 1.$$


Now we have $|(T \Delta S) \cap (U \Delta S)| = |S \setminus (T \cup U)| + |((T \cap U) \setminus S)| = \mu$.

Now, Lemma 6 tells us that \mathcal{F}' is a strong Δ -system. Let C' be the core of \mathcal{F}' . Note that $|(T \Delta S) \cap S| = |S \setminus T| = \mu$ for each $T \in \mathcal{F} \setminus \{S\}$ or equivalently $|T' \cap S| = \mu$ for each $T' \in \mathcal{F}'$. We claim that $T' \cap S = C'$ for each $T' \in \mathcal{F}'$. Suppose not. Then, we have $C' \setminus S \neq \emptyset$ because $|T' \cap S| = \mu$. It follows that there exists an element $x \in (T' \setminus C') \cap S$ for each $T' \in \mathcal{F}'$. Since the set family $\{T' \setminus C' \mid T' \in \mathcal{F}'\}$ is pairwise disjoint, it gives us $|S| \geq \mu^2 + \mu + 2 > 2\mu + 1$, a contradiction. Thus, \mathcal{F} is a sunflower with its core being $S \setminus C'$.

(ii) Let x be an element which is not included in any set of \mathcal{F} . Consider the set family $\mathcal{F}' = \{S \cup \{x\} \mid S \in \mathcal{F}\}$. It is easy to see that \mathcal{F}' is a $(2\mu + 2)$ -uniform weak Δ -system with intersection size $\mu + 1$. Since \mathcal{F}' is a sunflower by Lemma 6, so is \mathcal{F} . ◀

In order to obtain a linear-time algorithm for DMC with $\alpha = 0$ and $\beta = 2$, we will prove that for $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ with $\delta(\mathbf{T}) \leq 2$ the set system \mathcal{T} is a sunflower if ℓ is sufficiently large. This yields a linear-time algorithm via a reduction to a linear-time solvable special case of CONRMC. We start with a simple observation on matrices of diameter two, which will be helpful in the subsequent proofs.

► **Observation 8.** *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix with $\delta(\mathbf{T}) \leq 2$. For each $T_1 \in \mathcal{T}_1$ and $T_2, T_2' \in \mathcal{T}_2$, it holds that $T_1 \subseteq T_2$ and that $|T_2 \cap T_2'| \geq 1$ (otherwise there exists a pair of rows with Hamming distance three).*

(a) The case $\mathcal{T}_2 = \emptyset$.(b) The case $|\mathcal{T}_2| \geq 4$.

■ **Figure 3** Illustration of Lemma 11 with $n = 6$. A black cell denotes a value different from row vector $\mathbf{T}[6]$. In (b) the set system \mathcal{T}_2 forms a sunflower with core $\{2\}$. In both cases the radius is one.

The next lemma states that $|\mathcal{T}_2|$ restricts the number of columns.

► **Lemma 9.** *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix consisting of dirty columns with $\delta(\mathbf{T}) \leq 2$. If $\mathcal{T}_2 \neq \emptyset$, then $\ell \leq |\mathcal{T}_2| + 1$.*

Proof. First, observe that $\ell = |\bigcup_{T_1 \in \mathcal{T}_1} T_1 \cup \bigcup_{T_2 \in \mathcal{T}_2} T_2|$ because each column of \mathbf{T} is dirty. Thus, it follows from Observation 8 that $\ell = |\bigcup_{T_2 \in \mathcal{T}_2} T_2|$. We prove the lemma by induction on $|\mathcal{T}_2|$. Clearly, we have at most two columns if $|\mathcal{T}_2| = 1$. Suppose that $|\mathcal{T}_2| \geq 2$. For $T_2 \in \mathcal{T}_2$, we claim that

$$\ell = \left| \bigcup_{T'_2 \in \mathcal{T}_2} T'_2 \right| = \left| \bigcup_{T'_2 \in \mathcal{T}_2 \setminus \{T_2\}} T'_2 \right| + \left| T_2 \setminus \bigcup_{T'_2 \in \mathcal{T}_2 \setminus \{T_2\}} T'_2 \right| \leq |\mathcal{T}_2| + 1.$$

The induction hypothesis gives us that $|\bigcup_{T'_2 \in \mathcal{T}_2 \setminus \{T_2\}} T'_2| \leq |\mathcal{T}_2|$. For the other term, observe that $|T_2 \setminus \bigcup_{T'_2 \in \mathcal{T}_2 \setminus \{T_2\}} T'_2| \leq |T_2 \setminus T'_2| = |T_2| - |T_2 \cap T'_2|$ for $T'_2 \in \mathcal{T}_2 \setminus \{T_2\}$. Hence, it follows from Observation 8 that the second term is at most 1. ◀

Next, we show that a matrix with diameter at most two has radius at most one as long as it has at least five columns. Thus, we can solve DMC by solving CONRMC with radius one, which can be done in linear time [16]. We use the following lemma concerning certain intersections of a set with elements of a sunflower.

► **Lemma 10** ([9, Lemma 8]). *Let $\lambda \in \mathbb{N}$, let \mathcal{F} be a sunflower with core C , and let X be a set such that $|X \cap S| \geq \lambda$ for all $S \in \mathcal{F}$. If $|\mathcal{F}| > |X|$, then $\lambda \leq |C|$ and $|X \cap C| \geq \lambda$.*

► **Lemma 11.** *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix with $\delta(\mathbf{T}) \leq 2$. If $\ell \geq 5$, then there exists a vector $v \in \{0, 1\}^\ell$ such that $d(v, \mathbf{T}[i]) \leq 1$ for all $i \in [n]$.*

Proof. If $\mathcal{T}_2 = \emptyset$, then we are immediately done by definition, because $d(\mathbf{T}[n], \mathbf{T}[i]) \leq 1$ for all $i \in [n]$ (see Figure 3a for an illustration). Since $\ell \geq 5$, Lemma 9 implies $|\mathcal{T}_2| \geq 4$.

It follows from Observation 8 that \mathcal{T}_2 is a 2-uniform weak Δ -system with intersection size one (see Figure 3b). Thus, \mathcal{T}_2 is a sunflower by Lemma 6. Let $\{j_{\text{core}}\}$ denote the core of \mathcal{T}_2 . Note that $|T_1 \cap T_2| \geq 1$ holds for each $T_1 \in \mathcal{T}_1$ and $T_2 \in \mathcal{T}_2$ by Observation 8. Now we can infer from Lemma 10 that $T_1 = \{j_{\text{core}}\}$ for each $T_1 \in \mathcal{T}_1$.

Hence, it holds that $d(v, \mathbf{T}[i]) \leq 1$ for all $i \in [n]$, where $v \in \{0, 1\}^\ell$ is a row vector such that $v[j_{\text{core}}] = 1 - \mathbf{T}[n, j_{\text{core}}]$ and $v[j] = \mathbf{T}[n, j]$ for each $j \in [\ell] \setminus \{j_{\text{core}}\}$. ◀

We arrive at the following theorem.

► **Theorem 12.** *(0, 2)-DMC can be solved in $O(n\ell)$ time.*

Proof. Let $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ be an input matrix of $(0, 2)$ -DMC. If $\ell \leq 4$, then we use the linear-time algorithm of Lemma 1. Henceforth, we assume that $\ell \geq 5$.

We claim that \mathbf{S} is a **Yes**-instance if and only if the CONRMC instance $I' = (\mathbf{S}, 1^n)$ is a **Yes**-instance.

(\Rightarrow) Let \mathbf{T} be a completion of \mathbf{S} with $\delta(\mathbf{T}) \leq 2$. Since $\ell \geq 5$, there exists a vector v such that $d(v, \mathbf{T}[i]) \leq 1$ for all $i \in [n]$ by Lemma 11. It follows that I' is a **Yes**-instance.

(\Leftarrow) Let v be a solution of I' . Let \mathbf{T} be the matrix such that $\mathbf{T}[i] = \mathbf{S}[i] \oplus v$ for each $i \in [n]$. Then, we have $d(v, \mathbf{T}[i]) \leq 1$ for each $i \in [n]$. By the triangle inequality, we obtain $d(\mathbf{T}[i], \mathbf{T}[i']) \leq d(v, \mathbf{T}[i]) + d(v, \mathbf{T}[i']) \leq 2$ for each $i, i' \in [n]$.

Since CONRMC can be solved in linear time when $\max_{i \in [n]} r[i] = 1$ [16, Theorem 1], it follows that $(0, 2)$ -DMC can be solved in linear time. \blacktriangleleft

Next, we show polynomial-time solvability of $(0, 3)$ -DMC. The overall idea is, albeit technically more involved, similar to $(0, 2)$ -DMC. We first show that the set family \mathcal{T} of a matrix \mathbf{T} with $\delta(\mathbf{T}) = 3$ contains a sunflower by Lemma 7. We then show that such a matrix has a certain structure (Lemma 16) which again allows us to reduce the problem to the linear-time solvable special case of CONRMC with radius one (Theorem 18). We start with an observation on a matrix whose diameter is at most three.

► Observation 13. *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix with $\delta(\mathbf{T}) \leq 3$. It holds for each $T_1 \in \mathcal{T}_1$, $T_2 \in \mathcal{T}_2$, and $T_3, T'_3 \in \mathcal{T}_3$ that $T_1 \subseteq T_3$, $T_2 \cap T_3 \neq \emptyset$, and $|T_3 \cap T'_3| \geq 2$ (otherwise there exists a pair of rows with Hamming distance four).*

From Observation 13, we obtain (by induction) the following lemma analogously to Lemma 9.

► Lemma 14. *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix consisting of dirty columns with $\delta(\mathbf{T}) \leq 3$. If $\mathcal{T}_3 \neq \emptyset$, then $\ell \leq |\mathcal{T}_2| + |\mathcal{T}_3| + 2$.*

Our goal is to use Lemma 7 to derive that \mathcal{T}_3 forms a sunflower, that is, we need that $|\mathcal{T}_3| \geq 5$. The next lemma shows that this holds when \mathbf{T} has at least 14 dirty columns.

► Lemma 15. *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix consisting of $\ell \geq 14$ dirty columns with $\delta(\mathbf{T}) = 3$. Then, $|\mathcal{T}_3| \geq 5$ (for some permutation of rows).*

Proof. Assume that the rows are permuted such that $|\mathcal{T}_3|$ is maximized. If $|\mathcal{T}_3| \leq 4$, then we have $|\mathcal{T}_2| \geq 8$ by Lemma 14. Let $T_3 \in \mathcal{T}_3$. By Observation 13, $T_2 \cap T_3 \neq \emptyset$ holds for each $T_2 \in \mathcal{T}_2$. There are at most three sets $T_2 \in \mathcal{T}_2$ with $T_2 \subseteq T_3$. Thus, there are at least five sets $T_2 \in \mathcal{T}_2$ such that $|T_2 \cap T_3| = 1$. For each of these five sets, it holds that $|T_2 \triangle T_3| = |T_2| + |T_3| - 2|T_2 \cap T_3| = 3$. This contradicts the choice of the row permutation. \blacktriangleleft


With Lemma 15 at hand, we are ready to reveal the structure of a diameter-three matrix (see Figure 4 for an illustration).

► Lemma 16. *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix with $\delta(\mathbf{T}) \leq 3$ and $|\mathcal{T}_3| \geq 5$. Then, there exist $j_1 \neq j_2 \in [\ell]$ such that the following hold:*

- $T_1 \subseteq \{j_1, j_2\}$ for each $T_1 \in \mathcal{T}_1$.
- $T_2 \cap \{j_1, j_2\} \neq \emptyset$ for each $T_2 \in \mathcal{T}_2$.
- $T_3 \supseteq \{j_1, j_2\}$ for each $T_3 \in \mathcal{T}_3$.

Moreover, exactly one of the following holds for $\mathcal{T}'_2 = \{T_2 \in \mathcal{T}_2 \mid j_1 \in T_2 \wedge j_2 \notin T_2\}$ and $\mathcal{T}''_2 = \{T_2 \in \mathcal{T}_2 \mid j_1 \notin T_2 \wedge j_2 \in T_2\}$:

- (a) $\mathcal{T}'_2 = \emptyset$ or $\mathcal{T}''_2 = \emptyset$.



■ **Figure 4** Illustration (for smaller ℓ) of Lemma 16 (a) (left) and (b) (right). A black cell indicates that the entry differs from the last row vector in the corresponding column.

(b) $\mathcal{T}'_2 = \{\{j_1, j_3\}\}$ and $\mathcal{T}'_3 = \{\{j_2, j_3\}\}$ for $j_3 \in [\ell]$.

Proof. Note that \mathcal{T}_3 is 3-uniform by definition and note also that it is a weak Δ -system with intersection size two by Observation 13. Hence, Lemma 7 gives us column indices $j_1, j_2 \in [\ell]$ such that $\{j_1, j_2\} \subseteq T_3$ for each $T_3 \in \mathcal{T}_3$. It follows from Observation 13 and Lemma 10 that $T_1 \subseteq \{j_1, j_2\}$ for each $T_1 \in \mathcal{T}_1$ and $T_2 \cap \{j_1, j_2\} \neq \emptyset$ for each $T_2 \in \mathcal{T}_2$.

Now we show that either (a) or (b) holds. Suppose that $|\mathcal{T}'_1| \geq 2$ and $|\mathcal{T}'_2| \geq 1$, and let $T_2 \neq T'_2 \in \mathcal{T}'_2$ and $T''_2 \in \mathcal{T}'_2$. Then, either $T_2 \cap T''_2 = \emptyset$ or $T'_2 \cap T''_2 = \emptyset$ must hold. However, this is a contradiction because the corresponding row vectors have Hamming distance four. Thus, we have that $|\mathcal{T}'_2| \leq 1$ or $\mathcal{T}'_2 = \emptyset$. Analogously, we obtain $\mathcal{T}'_2 = \emptyset$ or $|\mathcal{T}'_2| \leq 1$. If $\mathcal{T}'_2 = \emptyset$ or $\mathcal{T}'_2 = \emptyset$, then (a) is satisfied. Otherwise, we have $|\mathcal{T}'_2| = |\mathcal{T}'_2| = 1$. Since $|T'_2 \Delta T''_2| = |T'_2| + |T''_2| - 2|T'_2 \cap T''_2| \leq \delta(\mathbf{T}) \leq 3$, we obtain $T'_2 \cap T''_2 \neq \emptyset$ for each $T'_2 \in \mathcal{T}'_2$ and $T''_2 \in \mathcal{T}'_2$. Hence, (b) holds. ◀

The following lemma establishes a connection to CONRMC. For $v, v' \in \{0, 1\}^\ell$ and $J \subseteq [\ell]$, we write $d_J(v, v')$ to denote $d(v[J], v'[J])$.

► **Lemma 17.** Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ be a matrix consisting of dirty columns with $\delta(\mathbf{T}) = 3$. If $\ell \geq 14$, then there exists a $v \in \{0, 1\}^\ell$ such that at least one of the following holds:

- (a) There exists $j \in [\ell]$ such that $d_{[\ell] \setminus \{j\}}(v, \mathbf{T}[i]) \leq 1$ for all $i \in [n]$.
- (b) There exist three column indices $J = \{j_1, j_2, j_3\} \subseteq [\ell]$ such that all of the following hold for each $i \in [n]$:

- $d(v_J, t_{i,J}) \leq 2$.
- If $d(v_J, t_{i,J}) \geq 1$, then $d_{[\ell] \setminus J}(v, \mathbf{T}[i]) = 0$.
- If $d(v_J, t_{i,J}) = 0$, then $d_{[\ell] \setminus J}(v, \mathbf{T}[i]) \leq 1$.

Here $v_J = (v[j_1], v[j_2], v[j_3])$ and $t_{i,J} = (\mathbf{T}[i, j_1], \mathbf{T}[i, j_2], \mathbf{T}[i, j_3])$ for each $i \in [n]$.

Proof. From Lemma 15, we can assume that $|\mathcal{T}_3| \geq 5$. Hence, Lemma 16 applies. Let j_1 and j_2 be the according column indices. Let $v \in \{0, 1\}^\ell$ be the row vector with

$$v[j] = \begin{cases} 1 - \mathbf{T}[n, j] & \text{if } j \in \{j_1, j_2\}, \\ \mathbf{T}[n, j] & \text{otherwise,} \end{cases}$$

for each $j \in [\ell]$. We claim that (a) corresponds to Lemma 16 (a), and (b) corresponds to Lemma 16 (b).

Suppose that Lemma 16 (a) holds with $\mathcal{T}'_2 = \emptyset$ (the case $\mathcal{T}'_2 = \emptyset$ is completely analogous). We prove that $d_{[\ell] \setminus \{j_2\}}(v, \mathbf{T}[i]) \leq 1$ for all $i \in [n]$. Since $d_{[\ell] \setminus \{j_2\}}(v, \mathbf{T}[i]) =$

$|\{j_1\} \Delta (D(\mathbf{T}[i], \mathbf{T}[n]) \setminus \{j_2\})|$, it suffices to show that $|\{j_1\} \Delta (T \setminus \{j_2\})| \leq 1$ holds for all $T \in \mathcal{T}$. Due to Lemma 16, we have

- $\{j_1\} \Delta (T \setminus \{j_2\}) \subseteq \{j_1\}$ for each $T \in \mathcal{T}_1$ and
- $|\{j_1\} \Delta (T \setminus \{j_2\})| = |T \setminus \{j_1, j_2\}| \leq 1$ for each $T \in \mathcal{T}_2 \cup \mathcal{T}_3$ since $j_1 \in T$.

Hence, (a) is true.

Now, assume that Lemma 16 (b) holds and let $J = \{j_1, j_2, j_3\}$. If there exists $i \in [n]$ with $d(v_J, t_{i,J}) = 3$, then this implies $\{j_3\} \in \mathcal{T}_1$ which yields the contradiction $\{j_3\} \not\subseteq \{j_1, j_2\}$.

Further, for each $T \in \mathcal{T}_1 \cup \mathcal{T}_2$, we have $T \setminus J = \emptyset$. Hence, for the row vector $\mathbf{T}[i]$ corresponding to T , we have $d_{[\ell] \setminus J}(v, \mathbf{T}[i]) = 0$. Now, let $T_3 \in \mathcal{T}_3$ with corresponding row vector $\mathbf{T}[i]$. If $T_3 = \{j_1, j_2, j_3\}$, then $d(v_J, t_{i,J}) = 1$ and $d_{[\ell] \setminus J}(v, \mathbf{T}[i]) = 0$. Otherwise, we have $T_3 = \{j_1, j_2, j\}$ for some $j \in [\ell] \setminus J$. Hence, $d(v_J, t_{i,J}) = 0$ and $d_{[\ell] \setminus J}(v, \mathbf{T}[i]) = |T_3 \setminus \{j_1, j_2\}| = 1$. Hence, (b) is true. \blacktriangleleft

Based on the connection to CONRMC, we obtain a polynomial-time algorithm.

► **Theorem 18.** *(0, 3)-DMC can be solved in $O(n^{\ell^4})$ time.*

Proof. We first apply Theorem 12 to determine whether there exists a completion $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ of $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ such that $\delta(\mathbf{T}) \leq 2$. If not, then it remains to determine whether there exists a completion \mathbf{T} with $\delta(\mathbf{T}) = 3$. We can assume that $\ell \geq 14$ by Lemma 1. We solve the problem by solving several instances of CONRMC based on Lemma 17.

For $j \in [\ell]$, let $I_j = (\mathbf{S}[:, [\ell] \setminus \{j\}], 1^n)$ be a CONRMC instance and let $\mathcal{I}_1 = \{I_j \mid j \in [\ell]\}$. These instances correspond to Lemma 17 (a).

Now, we describe the instances corresponding to Lemma 17 (b). Let $j_1, j_2, j_3 \in [\ell]$ be three distinct column indices and let $v_1, v_2, v_3 \in \{0, 1\}$. We define an instance $I_{j_1, j_2, j_3}^{v_1, v_2, v_3} = (\mathbf{S}_{j_1, j_2, j_3}, r)$ of CONRMC as follows:

- $\mathbf{S}_{j_1, j_2, j_3} = \mathbf{S}[:, [\ell] \setminus \{j_1, j_2, j_3\}]$.
- For each $i \in [n]$, let

$$r[i] = \begin{cases} 0 & \text{if } (\mathbf{S}[i, j_1] = 1 - v_1) \vee (\mathbf{S}[i, j_2] = 1 - v_2) \vee (\mathbf{S}[i, j_3] = 1 - v_3). \\ 1 & \text{otherwise.} \end{cases}$$

Let \mathcal{I}_2 contain those instances $I_{j_1, j_2, j_3}^{v_1, v_2, v_3}$ in which for each $i \in [n]$ at least one of $\mathbf{S}[i, j_1] \neq 1 - v_1$, $\mathbf{S}[i, j_2] \neq 1 - v_2$, or $\mathbf{S}[i, j_3] \neq 1 - v_3$ holds. We claim that \mathbf{S} is a **Yes**-instance if and only if at least one instance in \mathcal{I}_1 or \mathcal{I}_2 is a **Yes**-instance.

- If $I_j \in \mathcal{I}_1$ is a **Yes**-instance, then there exists $v \in \{0, 1\}^{\ell-1}$ such that $d(v', \mathbf{S}[i, [\ell] \setminus \{j\}]) \leq 1$ for each $i \in [n]$. Let \mathbf{T} be the completion of \mathbf{S} in which $\mathbf{T}[i, [\ell] \setminus \{j\}] = \mathbf{S}[i, [\ell] \setminus \{j\}] \oplus v$ for each $i \in [n]$. Then, we have

$$\begin{aligned} d(\mathbf{T}[i], \mathbf{T}[i']) &\leq d(\mathbf{T}[i, [\ell] \setminus \{j\}], \mathbf{T}[i', [\ell] \setminus \{j\}]) + 1 \\ &\leq d(v, \mathbf{T}[i, [\ell] \setminus \{j\}]) + d(v, \mathbf{T}[i', [\ell] \setminus \{j\}]) + 1 \leq 3 \end{aligned}$$

for each $i, i' \in [n]$.

- If $I_{j_1, j_2, j_3}^{v_1, v_2, v_3} = (\mathbf{S}_{j_1, j_2, j_3}, r) \in \mathcal{I}_2$ is a **Yes**-instance with solution $v' \in \{0, 1\}^{\ell-3}$, then let $v \in \{0, 1\}^\ell$ be the row vector obtained from v' by inserting v_1, v_2 , and v_3 in the j_1 -th, j_2 -th, and j_3 -th column, respectively, and let \mathbf{T} be the completion of \mathbf{S} in which $\mathbf{T}[i] = \mathbf{S}[i] \oplus v$ or each $i \in [n]$. We prove that $\delta(\mathbf{T}) \leq 3$. Let $R_x \subseteq [n]$ be the set of row indices i with $r[i] = x$ for $x \in \{0, 1\}$. Then, we have that

- $d(v, \mathbf{T}[i]) = d_{\{j_1, j_2, j_3\}}(v, \mathbf{T}[i]) + d_{[\ell] \setminus \{j_1, j_2, j_3\}}(v, \mathbf{T}[i]) \leq 2 + 0 = 2$ for each $i \in R_0$ and

– $d(v, \mathbf{T}[i]) = d_{\{j_1, j_2, j_3\}}(v, \mathbf{T}[i]) + d_{[\ell] \setminus \{j_1, j_2, j_3\}}(v, \mathbf{T}[i]) \leq 0 + 1 = 1$ for each $i \in R_1$.
 By the triangle inequality, we obtain $d(\mathbf{T}[i], \mathbf{T}[i']) \leq 3$ for each $i, i' \in [n]$ with $i \in R_1$ or $i' \in R_1$. Thus, it suffices to show $\delta(\mathbf{T}[R_0]) \leq 3$. Since $\mathbf{T}[i, [\ell] \setminus \{j_1, j_2, j_3\}] = \mathbf{T}[i', [\ell] \setminus \{j_1, j_2, j_3\}] = v'$ for each $i, i' \in R_0$, this clearly holds.

The reverse direction is easily verified using Lemma 17.

Overall, we construct $O(\ell^3)$ CONRMC instances, each of which can be solved in $O(n\ell)$ time [16, Theorem 1]. Hence, $(0, 3)$ -DMC can be solved in $O(n\ell^4)$ time. ◀

Our algorithms presented here work via reductions to CONRMC. Although CONRMC imposes an upper bound on the diameter implicitly by the triangle inequality, it is seemingly difficult to enforce any lower bounds (that is, $\alpha > 0$). In the next section, we will see polynomial-time algorithms for $\alpha > 0$, based on reductions to the graph factorization problem.

3.2 Polynomial time for $\beta \leq \alpha + 1$

We now give polynomial-time algorithms for (α, β) -DMC with constant $\alpha > 0$ given that $\beta \leq \alpha + 1$. As in Section 3.1, our algorithms exploit combinatorial structures revealed by Deza's theorem (Lemmas 6 and 7). Recall that \mathcal{T} denotes a set system obtained from a complete matrix \mathbf{T} (Definition 3). We show that \mathcal{T} essentially is a sunflower when $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \alpha + 1$. For the completion into such a sunflower, it suffices to solve the following matrix completion problem, which we call SUNFLOWER MATRIX COMPLETION.

SUNFLOWER MATRIX COMPLETION (SMC)

Input: An incomplete matrix $\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}$ and $s, m \in \mathbb{N}$.

Question: Is there a completion $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ of \mathbf{S} such that $D(\mathbf{T}[1], \mathbf{T}[n]), \dots, D(\mathbf{T}[n-1], \mathbf{T}[n])$ are pairwise disjoint sets of each size at most s and $\sum_{i \in [n-1]} |D(\mathbf{T}[i], \mathbf{T}[n])| \geq m$.

Intuitively speaking, the problem asks for a completion into a sunflower with empty core and bounded petal sizes. All algorithms presented in this subsection are via reductions to SMC. First, we show that SMC is indeed polynomial-time solvable. We prove this using a well-known polynomial-time algorithm for the graph problem (g, f) -FACTOR [10].

(g, f) -FACTOR


Input: A graph $G = (V, E)$, functions $f, g: V \rightarrow \mathbb{N}$, and $m' \in \mathbb{N}$.

Question: Does G contain a subgraph $G' = (V, E')$ such that $|E'| \geq m'$ and $g(v) \leq \deg_{G'}(v) \leq f(v)$ for all $v \in V$?

► **Lemma 19.** *For constant $s > 0$, SMC can be solved in $O(n\ell\sqrt{n+\ell})$ time.*

Proof. Let (\mathbf{S}, s, m) be an SMC instance. Let a_j^x be the number of occurrences of $x \in \{0, 1\}$ in $\mathbf{S}[:, j]$ for each $j \in [\ell]$. We can assume that $a_j^0 \geq a_j^1$ for each $j \in [\ell]$ (otherwise swap the occurrences of 0's and 1's in the column). If $a_j^0 \geq 2$ and $\mathbf{S}[n, j] = 1$ for some $j \in [\ell]$, then we can return **No** since there will be two intersecting sets. Also, if $a_j^1 \geq 2$, then we can return **No**.

We construct an instance of (g, f) -FACTOR as follows. We introduce a vertex u_i for each $i \in [n-1]$ and a vertex v_j for each $j \in [\ell]$. The resulting graph G will be a bipartite graph with one vertex subset $\{u_1, \dots, u_{n-1}\}$ representing rows and the other $\{v_1, \dots, v_\ell\}$ representing columns. Essentially, we add an edge between u_i and v_j if the column $\mathbf{S}[:, j]$ can



■ **Figure 5** A completion of a 5×5 incomplete matrix (left). The known entries are highlighted in gray. A bipartite graph as constructed in the reduction (right). Note that the entries framed by thick lines (which differ from all others in the same column) correspond to the subgraph represented by the thick lines.

be completed such that the i -th entry differs from all others (see Figure 5 for an illustration). Intuitively, such an edge encodes the information that column index j can be contained in a petal of the sought sunflower. Formally, there is an edge $\{u_i, v_j\}$ if and only if there is a completion $t_j \in \{0, 1\}^n$ of $\mathbf{S}[:, j]$ in which $t_j[i] = 1 - t_j[n]$ and $t_j[h] = t_j[n]$ for all $h \in [n-1] \setminus \{i\}$. We set $g(u_i) := 0$ and $f(u_i) := s$ for each $i \in [n-1]$, $g(v_j) := a_j^1$ and $f(v_j) := 1$ for each $j \in [\ell]$, and $m' := m$. This construction can be done in $O(n\ell)$ time. To see this, note that the existence of an edge $\{u_i, v_j\}$ only depends on a_j^0 , a_j^1 , and $\mathbf{S}[i, j]$.

- If $a_j^0 \leq 1$ and $a_j^1 = 0$, then add the edge $\{u_i, v_j\}$. The corresponding completion t_j can be seen as follows:
 - If $\mathbf{S}[h, j] = \square$ for all $h \in [n-1]$, then let $t_j[i] := 1$ and let $t_j[h] := 0$ for all $h \in [n] \setminus \{i\}$.
 - If $\mathbf{S}'[h, j] = 0$ for some $h \in [n-1]$, then $\mathbf{S}'[h', j] = \square$ for all $h' \in [n] \setminus \{h\}$. If $h \neq i$, then let $t_j[i] := 1$ and let $t_j[h] := 0$ for all $h \in [n] \setminus \{i\}$. Otherwise, let $t_j[h] := 1$ for all $h \in [n] \setminus \{i\}$.
- If $a_j^0 = 1$ and $a_j^1 = 1$, then add the edge $\{u_i, v_j\}$ if $\mathbf{S}[i, j] \neq \square$.
- If $a_j^0 \geq 2$ and $a_j^1 = 0$, then add the edge $\{u_i, v_j\}$ if $\mathbf{S}[i, j] = \square$.
- If $a_j^0 \geq 2$ and $a_j^1 = 1$, then add the edge $\{u_i, v_j\}$ if $\mathbf{S}[i, j] = 1$ (because $\mathbf{S}[n, j]$ must be completed with 0).

The correctness of the reduction easily follows from the definition of an edge: If \mathbf{T} is a solution for (\mathbf{S}, s, m) , then the corresponding subgraph of G contains the edge $\{u_i, v_j\}$ for each $i \in [n-1]$ and each $j \in D(\mathbf{T}[i], \mathbf{T}[n])$. Conversely, a completion of \mathbf{S} is obtained from a subgraph G' by taking for each edge $\{u_i, v_j\}$ the corresponding completion t_j as the j -th column. Note that no vertex v_j can have two incident edges since $f(v_j) = 1$. Moreover, if v_j has no incident edges, then this implies $g(v_j) = a_j^1 = 0$. Hence, we can complete all missing entries in column j by 0.

As regards the running time, note that the constructed graph G has at most $n\ell$ edges and $\sum_{i \in [n-1]} f(u_i) \in O(n)$ and $\sum_{j \in [\ell]} f(v_j) \in O(\ell)$. Since (g, f) -FACTOR can be solved in $O(|E|\sqrt{f(V)})$ time [10] for $f(V) = \sum_{v \in V} f(v)$, SMC can be solved in time $O(n\ell\sqrt{n+\ell})$. ◀

As a first application of Lemma 19, we show that (α, α) -DMC can be solved in polynomial time.

► **Theorem 20.** (α, α) -DMC can be solved in $O(n\ell\sqrt{n+\ell})$ time.

Proof. We first show that (α, α) -DMC can easily be solved if α is odd. Consider row vectors $u, v, w \in \{0, 1\}^\ell$ and let $U := D(u, v)$ and $W := D(v, w)$. Then, $d(u, v) + d(v, w) + d(w, u) = |U| + |W| + (|U| + |W| - 2|U \cap W|) = 2(|U| + |W| - |U \cap W|)$ and hence $d(u, v) + d(v, w) + d(w, u)$

is even. Thus, we can immediately answer **No** if $n \geq 3$. It is also easy to see that DMC can be solved in linear time if $n \leq 2$.

We henceforth assume that α is even. Eiben et al. [7, Theorem 34] provided a linear-time algorithm for $(0, \alpha)$ -DMC with constant n (and arbitrary α) using reductions to ILP. It is straightforward to adapt their ILP formulation to show that (α, α) -DMC can also be solved in linear time for constant n (basically, we just need the additional constraint that each pairwise distance is at least α). So we can assume that $n \geq (\alpha/2)^2 + (\alpha/2) + 3$. We claim that there is a completion \mathbf{T} of \mathbf{S} with $\gamma(\mathbf{T}) = \delta(\mathbf{T}) = \alpha$ if and only if the SMC instance $(\mathbf{S}', \alpha/2, \alpha n/2)$ is a **Yes**-instance for the matrix $\mathbf{S}' \in \{0, 1, \square\}^{(n+1) \times \ell}$ obtained from \mathbf{S} with an additional row vector \square^ℓ .

(\Rightarrow) Let \mathbf{T} be a completion of \mathbf{S} with $\gamma(\mathbf{T}) = \delta(\mathbf{T}) = \alpha$. Then, \mathcal{T} is a weak Δ -system with intersection size $\alpha/2$. Since $|\mathcal{T}| \geq (\alpha/2)^2 + (\alpha/2) + 2$, Lemma 6 tells us that \mathcal{T} is a sunflower. Let C be the core of \mathcal{T} . Consider the completion \mathbf{T}' of \mathbf{S}' such that

- $\mathbf{T}'[[n], :] = \mathbf{T}$,
- $\mathbf{T}'[n+1, j] = 1 - \mathbf{T}[n, j]$ for each $j \in C$, and
- $\mathbf{T}'[n+1, j] = \mathbf{T}[n, j]$ for each $j \in [\ell] \setminus C$.

Note that $D(\mathbf{T}'[i], \mathbf{T}'[n+1]) = D(\mathbf{T}'[i], \mathbf{T}'[n]) \setminus C$ for each $i \in [n-1]$. Note also that $D(\mathbf{T}'[n], \mathbf{T}'[n+1]) = C$. Hence, $D(\mathbf{T}'[1], \mathbf{T}'[n+1]), \dots, D(\mathbf{T}'[n], \mathbf{T}'[n+1])$ are pairwise disjoint sets of size $\alpha/2$.

(\Leftarrow) Let \mathbf{T}' be a completion of \mathbf{S}' such that $D(\mathbf{T}'[1], \mathbf{T}'[n+1]), \dots, D(\mathbf{T}'[n], \mathbf{T}'[n+1])$ are pairwise disjoint sets of size $\alpha/2$. For the completion $\mathbf{T} = \mathbf{T}'[[n], :]$ of \mathbf{S} , it holds that $d(\mathbf{T}[i], \mathbf{T}[i']) = |D(\mathbf{T}'[i], \mathbf{T}'[n+1]) \Delta D(\mathbf{T}'[i'], \mathbf{T}'[n+1])| = |D(\mathbf{T}'[i], \mathbf{T}'[n+1])| + |D(\mathbf{T}'[i'], \mathbf{T}'[n+1])| = \alpha$ for each $i, i' \in [n]$. \blacktriangleleft

Now we proceed to develop polynomial-time algorithms for the case $\alpha + 1 = \beta$. We will make use of the following observation made by Froese et al. [9, Proof of Theorem 9].

► **Observation 21.** *Let $\mathbf{T} \in \{0, 1\}^{n \times \ell}$ with $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \beta = \alpha + 1$. For $T_\alpha \neq T'_\alpha \in \mathcal{T}_\alpha$ and $T_\beta \neq T'_\beta \in \mathcal{T}_\beta$, it holds that $|T_\alpha \cap T'_\alpha| = \lfloor \alpha/2 \rfloor$, $|T_\alpha \cap T_\beta| = \lceil \alpha/2 \rceil = \lfloor \beta/2 \rfloor$, and $|T_\beta \cap T'_\beta| = \lceil \beta/2 \rceil$.*

Surprisingly, odd α seems to allow for significantly more efficiently algorithms than even α .

► **Theorem 22.** *(α, β) -DMC with $\beta = \alpha + 1$ can be solved in*

- (i) $O(n\ell\sqrt{n+\ell})$ time for odd α , and
- (ii) $(n\ell)^{O(\alpha^3)}$ time for even α .

Proof. (i) We can assume that $n \geq \beta^2/2 + \beta + 7$ holds since otherwise the problem is linear-time solvable (as in the proof of Theorem 20). Suppose that \mathbf{S} admits a completion \mathbf{T} with $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \beta$. Since $\mathcal{T} = \mathcal{T}_\alpha \cup \mathcal{T}_\beta$ and $|\mathcal{T}| \geq \beta^2/2 + \beta + 6$, it follows that $\max\{|\mathcal{T}_\alpha|, |\mathcal{T}_\beta|\} \geq c := (\beta/2)^2 + (\beta/2) + 3$. We consider two cases depending on the size of \mathcal{T}_α and \mathcal{T}_β .

- Suppose that $|\mathcal{T}_\alpha| \geq c$. Since \mathcal{T}_α is a weak Δ -system with intersection size $(\alpha-1)/2$, \mathcal{T}_α is a sunflower with a core of size $(\alpha-1)/2$ and petals of size $(\alpha+1)/2$ by Lemma 7 (ii). We claim that $\mathcal{T}_\beta = \emptyset$. Suppose not and let $T_\beta \in \mathcal{T}_\beta$. Consequently, we obtain $|T_\alpha \cap T_\beta| = (\alpha+1)/2$ for all $T_\alpha \in \mathcal{T}_\alpha$ by Observation 21, which contradicts Lemma 10.
- Suppose that $|\mathcal{T}_\beta| \geq c$. Again, \mathcal{T}_β is a sunflower whose core C has size $\beta/2$ by Lemma 6. By Observation 21 and Lemma 10, $T_\alpha \supseteq C$ holds for each $T_\alpha \in \mathcal{T}_\alpha$. Now suppose that there exist $T_\alpha \neq T'_\alpha \in \mathcal{T}_\alpha$. Since $C \subseteq T_\alpha$ and $C \subseteq T'_\alpha$, it follows that $|T_\alpha \cap T'_\alpha| \geq \beta/2$, thereby contradicting Observation 21. Hence, we have $|\mathcal{T}_\alpha| \leq 1$.

We construct an instance I of SMC covering both cases above, as in Theorem 20. We use the matrix \mathbf{S}' obtained from \mathbf{S} by appending a row vector \square^ℓ , and we set $s := \beta/2$ and $m := ns - 1$. Basically, we allow at most one “petal” to have size $s - 1$. We return **Yes** if and only if I is a **Yes**-instance. The correctness can be shown analogously to the proof of Theorem 20.

(ii) Suppose that there is a completion \mathbf{T} of \mathbf{S} with $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \beta$. Again, we can assume that $n > 2c$ for $c := (\beta/2)^2 + (\beta/2) + 4$, and consider a case distinction regarding the size of \mathcal{T}_α and \mathcal{T}_β .

- Suppose that $|\mathcal{T}_\alpha| \geq c$ and $|\mathcal{T}_\beta| \geq c$. It follows from Observation 21 and Lemmas 6 and 7 that \mathcal{T}_α and \mathcal{T}_β are sunflowers. Let C_α and C_β be the cores of \mathcal{T}_α and \mathcal{T}_β , respectively. Note that $|C_\alpha| = \alpha/2$ and $|C_\beta| = \alpha/2 + 1$, and hence $C_\alpha \subsetneq C_\beta$ holds by Observation 21 and Lemma 10. Let $j \in [\ell]$ be such that $C_\alpha \cup \{j\} = C_\beta$ and let $\mathbf{T}' := \mathbf{T}[:, [\ell] \setminus \{j\}]$. Then, the set family \mathcal{T}' is a sunflower with a core of size $\alpha/2$ and petals of size $\alpha/2$. Hence, there exists $j \in [\ell]$ such that the (α, α) -DMC instance $\mathbf{S}[:, [\ell] \setminus \{j\}]$ is a **Yes**-instance. On the other hand, if there is a completion \mathbf{T}' of $\mathbf{S}[:, [\ell] \setminus \{j\}]$ with $\gamma(\mathbf{T}') = \delta(\mathbf{T}') = \alpha$, then $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \alpha + 1$ hold for any completion \mathbf{T} of \mathbf{S} with $\mathbf{T}[:, [\ell] \setminus \{j\}] = \mathbf{T}'$.
- Suppose that $|\mathcal{T}_\alpha| \geq c$ and $|\mathcal{T}_\beta| < c$. The same argument as above shows that $T_\alpha \cap T_\beta = C$ holds for each $T_\alpha \in \mathcal{T}_\alpha$ and $T_\beta \in \mathcal{T}_\beta$, where C is the size- $\alpha/2$ core of sunflower \mathcal{T}_α . Let $I_\beta = \{i \in [n-1] \mid d(\mathbf{T}[i], \mathbf{T}[n]) = \beta\}$ be the row indices that induce the sets in \mathcal{T}_β and let $J_\beta = \bigcup_{T_\beta \in \mathcal{T}_\beta} T_\beta$. Consider $\mathbf{T}' = \mathbf{S}[[n] \setminus I_\beta, [\ell] \setminus (C \cup J_\beta)]$ and note that the family \mathcal{T}' consists of pairwise disjoint sets, each of size $\alpha/2$. We use this observation to obtain a reduction to SMC. The idea is to test all possible choices for \mathbf{T}' , that is, we simply try out all possibilities to choose the following sets:
 - $C \subseteq [\ell]$ of size exactly $\alpha/2$.
 - $I_\beta \subseteq [n-1]$ of size at most c .
 - $J_\beta \subseteq [\ell] \setminus C$ of size at most $\beta \cdot c$ such that $d_{[\ell] \setminus (C \cup J_\beta)}(\mathbf{S}[i_\beta], \mathbf{S}[n]) = 0$ for all $i_\beta \in I_\beta$.
For each possible choice, we check whether it allows for a valid completion. Formally, it is necessary that the following exist:
 - A completion t_C of $\mathbf{S}[n, C]$ such that $\mathbf{S}[i, j] \neq t_C[j]$ for all $i \in [n-1]$ and $j \in C$.
 - A completion t_{J_β} of $\mathbf{S}[n, J_\beta]$ such that $d(t_{J_\beta}, \mathbf{S}[i, J_\beta]) = 0$ for all $i \in [n-1] \setminus I_\beta$.
 - A completion t_{i_β} of $\mathbf{S}[i_\beta, J_\beta]$ for each $i_\beta \in I_\beta$ such that $d(t_{i_\beta}, t_{J_\beta}) = \alpha/2 + 1$ for each $i_\beta \in I_\beta$ and $d(t_{i_\beta}, t_{i'_\beta}) = \alpha$ for each $i_\beta \neq i'_\beta \in I_\beta$.
The existence of the above completions can be checked in $O(n)$ time. We then construct an SMC instance $(\mathbf{S}', \alpha/2, (n - |I_\beta| - 1) \cdot \alpha/2)$, where \mathbf{S}' is an incomplete matrix with $n' = n - |I_\beta|$ rows and $\ell - |C| - |J_\beta|$ columns defined as follows:
 - $\mathbf{S}'[[n'] - 1] = \mathbf{S}[[n-1] \setminus I_\beta, [\ell] \setminus (C \cup J_\beta)]$.
 - $\mathbf{S}'[n', j] = \square$ for each $j \in [\ell] \setminus (C \cup J_\beta)$ such that $\mathbf{S}[i_\beta, j] = \square$ for all $i_\beta \in I_\beta \cup \{n\}$.
 - $\mathbf{S}'[n', j] = \mathbf{S}[i_\beta, j]$ for each $j \in [\ell] \setminus (C \cup J_\beta)$ such that $\mathbf{S}[i_\beta, j] \neq \square$ for some $i_\beta \in I_\beta \cup \{n\}$.
Overall, we solve at most $(n\ell)^{O(\alpha^3)}$ many SMC instances. Hence, this case is solvable in $(n\ell)^{O(\alpha^3)}$ time.
- Suppose that $|\mathcal{T}_\alpha| < c$ and $|\mathcal{T}_\beta| \geq c$. Let $i \in [n-1]$ be such that $d(\mathbf{T}[i], \mathbf{T}[n]) = \beta$. Then, $d(\mathbf{T}[i], \mathbf{T}[i']) = \alpha$ holds for each $i' \in [n-1] \setminus \{i\}$. Since there are at least $n-2 > 2c-2 > c$ such row indices, it follows that this case is essentially equivalent to the previous case (by considering row i as the last row).

◀

A natural question is whether one can extend our approach above to the case $\beta = \alpha + 2$ (especially $\alpha = 1$ and $\beta = 3$). The problem is that the petals of the sunflowers \mathcal{T}_2 and \mathcal{T}_3

may have nonempty intersections. Therefore, reducing to SMC to obtain a polynomial-time algorithm is probably impossible.

3.3 NP-hardness

Hermelin and Rozenberg [14, Theorem 5] proved that CONRMC (under the name CLOSEST STRING WITH WILDCARDS) is NP-hard even if $r[i] = 2$ for all $i \in [n]$. We use this result to prove the following.

► **Theorem 23.** (α, β) -DMC is NP-hard if $\beta \geq 2\lceil\alpha/2\rceil + 4$.

Proof. We show a polynomial-time reduction from CONRMC. Let $(\mathbf{S} \in \{0, 1, \square\}^{n \times \ell}, r)$ be a CONRMC instance with $r[i] = 2$ for all $i \in [n]$.

Let $\mathbf{C} \in \{0, 1\}^{(n+1) \times m}$ be the binary matrix with $m = (n-1) \cdot \lceil\alpha/2\rceil + \beta - 2$ columns obtained by horizontally stacking

- the $(n+1) \times (n+1)$ identity matrix $\lceil\alpha/2\rceil$ times and
- the column vector $(0^n 1)^T$ $\beta - 2\lceil\alpha/2\rceil - 2$ times.

Since the pairwise row Hamming distances in the identity matrix are all two, we have that:

- $d(\mathbf{C}[i], \mathbf{C}[i']) = 2\lceil\alpha/2\rceil$ for each $i \neq i' \in [n]$ and
- $d(\mathbf{C}[i], \mathbf{C}[n+1]) = \beta - 2$ for each $i \in [n]$.

Consider the matrix $\mathbf{S}' \in \{0, 1, \square\}^{(n+1) \times (\ell+m)}$ obtained from \mathbf{S} by adding the row \square^ℓ and then horizontally appending \mathbf{C} . We show that there exists a vector $v \in \{0, 1\}^\ell$ with $d(v, \mathbf{S}[i]) \leq 2$ for all $i \in [n]$ if and only if \mathbf{S}' admits a completion \mathbf{T}' with $\gamma(\mathbf{T}') \geq \alpha$ and $\delta(\mathbf{T}') \leq \beta$.

(\Rightarrow) Let \mathbf{T}' be the completion of \mathbf{S}' such that for each $i \in [n+1]$, $\mathbf{T}'[i, \ell] = \mathbf{S}'[i] \oplus v$.

Then, we have the following:

- $\gamma(\mathbf{T}') \geq \gamma(\mathbf{C}) = 2\lceil\alpha/2\rceil \geq \alpha$.
- $d(\mathbf{T}'[i], \mathbf{T}'[n+1]) = d(\mathbf{S}[i], v) + \beta - 2 \leq \beta$ for each $i \in [n]$.
- By the triangle inequality, $d(\mathbf{T}'[i], \mathbf{T}'[i']) \leq d(v, \mathbf{S}[i]) + d(v, \mathbf{S}[i']) + d(\mathbf{C}[i], \mathbf{C}[i']) \leq 2 + 2 + 2\lceil\alpha/2\rceil \leq \beta$ holds for each $i, i' \in [n]$

(\Leftarrow) Let $v = \mathbf{T}'[n+1, \ell]$. It is easy to see that $d(\mathbf{S}[i], v) = d(\mathbf{T}'[i], \mathbf{T}'[n+1]) - d(\mathbf{C}[i], \mathbf{C}[n+1]) \leq \beta - (\beta - 2) \leq 2$ holds for each $i \in [n]$. ◀

It remains open whether NP-hardness also holds for $(\alpha, \alpha + 3)$ -DMC with $\alpha \geq 1$ (recall that $(0, 3)$ -DMC is polynomial-time solvable). In Section 4.3, however, we show NP-hardness for $\beta = \alpha + 3$ when α and β are part of the input.

4 Bounded number k of missing entries per row

In this section, we consider DMC with α and β being part of the input, hence not necessarily being constants. We consider the maximum number k of missing entries in any row as a parameter. Clearly, DMC is trivial for $k = 0$. We will show that DMC is polynomial-time solvable for $k = 1$, and also for $k = 2$ if $\alpha = \beta$. On the negative side, we show that DMC is NP-hard for $k \geq 2$ with $\beta - \alpha \geq 3$ and for $k \geq 3$ with $\beta - \alpha = 0$.

4.1 Polynomial time for $k = 1$

We show that DMC can be solved in polynomial time when $k = 1$, via a reduction to 2-SAT. For a Boolean variable x , we use $(x = 1)$ and $(x \neq 0)$ to denote the positive literal x . Similarly, we use $(x = 0)$ and $(x \neq 1)$ for the negative literal $\neg x$.

► **Theorem 24.** DMC can be solved in $O(n^2\ell)$ time when $k = 1$.

Proof. We construct a 2-CNF formula ϕ of polynomial size such that ϕ is satisfiable if and only if the input matrix \mathbf{S} admits a completion \mathbf{T} with $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \beta$.

First, we compute the distances $d(\mathbf{S}[i], \mathbf{S}[i'])$ for each $i, i' \in [n]$ in $O(n^2\ell)$ time. Clearly, if there exists a pair with distance less than $\alpha - 2$ or larger than β , then we have a **No**-instance. Let $I \subseteq [n]$ be the set of row indices corresponding to row vectors with a missing entry and let $j_i \in [\ell]$ be such that $\mathbf{S}[i, j_i] = \square$.

We introduce a variable x_i for each $i \in I$, where x_i is set to true if $\mathbf{S}[i, j_i]$ is completed with a 1. We construct the formula ϕ as follows:

- For each $i < i' \in [n]$ with $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 2$:
 - If $i \notin I$ or $i' \notin I$, or $j_i = j_{i'}$, then return **No**. Otherwise, add the clauses $(x_i = 1 - \mathbf{S}[i', j_i])$ and $(x_{i'} = 1 - \mathbf{S}[i, j_{i'}])$.
- For each $i < i' \in [n]$ with $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 1$:
 - If $i \notin I$ and $i' \notin I$, then return **No**.
 - If $i \in I$ and $i' \notin I$, then add the clause $(x_i \neq \mathbf{S}[i', j_i])$.
 - If $i \notin I$ and $i' \in I$, then add the clause $(x_{i'} \neq \mathbf{S}[i, j_i])$.
 - If $i \in I$ and $i' \in I$ and $j_i = j_{i'}$, then add the clauses $(x_i \vee x_{i'})$ and $(\neg x_i \vee \neg x_{i'})$.
 - If $i \in I$ and $i' \in I$ and $j_i \neq j_{i'}$, then add the clause $(x_i \neq \mathbf{S}[i', j_i] \vee x_{i'} \neq \mathbf{S}[i, j_{i'}])$.

It is easy to see that these clauses ensure that $\gamma(\mathbf{T}) \geq \alpha$. Similarly, to ensure that $\delta(\mathbf{T}) \leq \beta$, we add the following clauses:

- For each $i < i' \in [n]$ with $d(\mathbf{S}[i], \mathbf{S}[i']) = \beta$:
 - If $i \in I$ and $i' \notin I$, then add the clause $(x_i = \mathbf{S}[i', j_i])$.
 - If $i \notin I$ and $i' \in I$, then add the clause $(x_{i'} = \mathbf{S}[i, j_{i'}])$.
 - If $i \in I$ and $i' \in I$ and $j_i = j_{i'}$, then add the clauses $(x_i \vee \neg x_{i'})$ and $(\neg x_i \vee x_{i'})$.
 - If $i \in I$ and $i' \in I$ and $j_i \neq j_{i'}$, then add the clauses $(x_i = \mathbf{S}[i', j_i])$ and $(x_{i'} = \mathbf{S}[i, j_{i'}])$.
- For each $i < i' \in [n]$ with $d(\mathbf{S}[i], \mathbf{S}[i']) = \beta - 1$:
 - If $i \in I$ and $i' \in I$ and $j_i \neq j_{i'}$, then add the clause $(x_i = \mathbf{S}[i', j_i] \vee x_{i'} = \mathbf{S}[i, j_{i'}])$.

Thus, ϕ is of size $O(n^2)$ and can be solved in $O(n^2)$ time [1]. The correctness follows directly from the construction. ◀

We remark that the quadratic dependence on n in the running time of Theorem 24 is presumably inevitable. To prove this, we will use the **ORTHOGONAL VECTORS** conjecture, which states that **ORTHOGONAL VECTORS** cannot be solved in time $O(n^{2-\epsilon} \cdot \ell^c)$ for any $\epsilon, c > 0$ (assuming the Strong Exponential Time Hypothesis) [12].

ORTHOGONAL VECTORS

Input: Sets \mathcal{U}, \mathcal{V} of row vectors in $\{0, 1\}^\ell$ with $|\mathcal{U}| = |\mathcal{V}| = n$.

Question: Are there row vectors $u \in \mathcal{U}$ and $v \in \mathcal{V}$ such that $u[j] \cdot v[j] = 0$ holds for all $j \in [\ell]$?

► **Theorem 25.** DMC cannot be solved in time $O(n^{2-\epsilon} \cdot \ell^c)$ for any $c, \epsilon > 0$, unless the **ORTHOGONAL VECTORS** conjecture breaks.

Proof. We reduce from **ORTHOGONAL VECTORS**. Let $u_1, \dots, u_n, v_1, \dots, v_n \in \{0, 1\}^\ell$ be row

$$\mathbf{T} = \left[\begin{array}{c|c|c|c} 001 & 111 & 001 & 00000000 \\ 111 & 001 & 001 & 00000000 \\ \hline 111 & 111 & 010 & 11111111 \\ 111 & 010 & 111 & 11111111 \end{array} \right]$$

■ **Figure 6** An illustration of the reduction from ORTHOGONAL VECTORS, where $\mathcal{U} = \{010, 110\}$ and $\mathcal{V} = \{110, 101\}$.

vectors. Consider the matrix $\mathbf{T} \in \{0, 1\}^{2n \times 6\ell}$ where

$$\mathbf{T}[i, [3j-2, 3j]] = \begin{cases} 001 & \text{if } j \leq \ell \text{ and } u_i[j] = 0 \\ 111 & \text{if } j \leq \ell \text{ and } u_i[j] = 1 \\ 000 & \text{otherwise} \end{cases}$$

$$\mathbf{T}[n+i, [3j-2, 3j]] = \begin{cases} 010 & \text{if } j \leq \ell \text{ and } v_i[j] = 0 \\ 111 & \text{if } j \leq \ell \text{ and } v_i[j] = 1 \\ 111 & \text{otherwise} \end{cases}$$

for each $i \in [n]$ and $j \in [2\ell]$. It is easy to see that there are $i, i' \in [n]$ such that u_i and $v_{i'}$ are orthogonal if and only if $\delta(\mathbf{T}) = 5\ell$ (see Figure 6 for an illustration). ◀

4.2 Polynomial time for $k = 2$ with $\alpha = \beta$

We extend the polynomial-time solvability for $k = 1$ to the case $k = 2$ when $\alpha = \beta$. Again, we show a polynomial-time reduction to 2-SAT based on a more extensive case distinction.

► **Theorem 26.** DMC can be solved in $O(n^2\ell)$ time for $k = 2$ and $\alpha = \beta$.

Proof. Since $k = 2$, there are at most four possible ways to complete the row vector $\mathbf{S}[n]$. Assuming that $\mathbf{S}[n]$ is complete (without loss of generality, $\mathbf{S}[n] = 0^\ell$) we show that DMC can be solved in $O(n^2\ell)$ time.

First, we check whether $\alpha - 2 \leq d(\mathbf{S}[i], 0^\ell) \leq \alpha$ holds for each $i \in [n-1]$ (otherwise, we return **No**). We do the following for each $i \in [n-1]$:

- $\mathbf{S}[i]$ contains exactly one missing entry: If $d(\mathbf{S}[i], 0^\ell) = \alpha - 1$, then fill the missing entry by 1. If $d(\mathbf{S}[i], 0^\ell) = \alpha$, then fill the missing entry by 0. If $d(\mathbf{S}[i], 0^\ell) = \alpha - 2$, then return **No**.
- $\mathbf{S}[i]$ contains exactly two missing entries: If $d(\mathbf{S}[i], 0^\ell) = \alpha - 2$, then fill both missing entries by 1. If $d(\mathbf{S}[i], 0^\ell) = \alpha$, then fill both missing entries by 0.

Now, each row vector either contains no missing entry or exactly two missing entries. Let $I_0 \subseteq [n]$ be the set of row indices corresponding to row vectors without any missing entry and let $I_2 = [n] \setminus I_0$. We check in $O(n^2\ell)$ time whether all pairwise Hamming distances in $\mathbf{S}[I_0]$ are α . If not, then we return **No**. Note that we have $d(\mathbf{S}[i], 0^\ell) = \alpha - 1$ for each $i \in I_2$, and thus there are exactly two ways to complete $\mathbf{S}[i]$: One missing entry filled by 1 and the other by 0, or vice versa. For each $i \in I_2$, let $j_i^1 < j_i^2 \in [\ell]$ be such that $\mathbf{S}[i, j_i^1] = \mathbf{S}[i, j_i^2] = \square$. We verify whether the following necessary conditions hold:

- $d(\mathbf{S}[i_0], \mathbf{S}[i]) = \alpha - 1$ for each $i_0 \in I_0$ and $i \in I_2$ with $\mathbf{S}[i_0, j_i^1] = \mathbf{S}[i_0, j_i^2]$.
- $d(\mathbf{S}[i_0], \mathbf{S}[i]) \in \{\alpha - 2, \alpha\}$ for each $i_0 \in I_0$ and $i \in I_2$ with $\mathbf{S}[i_0, j_i^1] \neq \mathbf{S}[i_0, j_i^2]$.
- Let $i \neq i' \in I_2$ be such that $\{j_i^1, j_i^2\} \cap \{j_{i'}^1, j_{i'}^2\} = \emptyset$. Observe that if $\mathbf{S}[i', j_i^1] = \mathbf{S}[i', j_i^2]$, then the completion of $\mathbf{S}[i]$ increases the distance between $\mathbf{S}[i]$ and $\mathbf{S}[i']$ by exactly one.

Otherwise, the distance either stays the same or increases by exactly two. It is analogous for the completion of $\mathbf{S}[i']$. Thus, the following must hold for each $i \neq i' \in I_2$ with $\{j_i^1, j_i^2\} \cap \{j_{i'}^1, j_{i'}^2\} = \emptyset$:

- $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 2$ if $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i, j_{i'}^2]$ and $\mathbf{S}[i', j_i^1] = \mathbf{S}[i', j_i^2]$.
- $d(\mathbf{S}[i], \mathbf{S}[i']) = \{\alpha - 3, \alpha - 1\}$ if either $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i, j_{i'}^2]$ and $\mathbf{S}[i', j_i^1] \neq \mathbf{S}[i', j_i^2]$, or $\mathbf{S}[i, j_{i'}^1] \neq \mathbf{S}[i, j_{i'}^2]$ and $\mathbf{S}[i', j_i^1] = \mathbf{S}[i', j_i^2]$.
- $d(\mathbf{S}[i], \mathbf{S}[i']) = \{\alpha - 4, \alpha - 2, \alpha\}$ if $\mathbf{S}[i, j_{i'}^1] \neq \mathbf{S}[i, j_{i'}^2]$ and $\mathbf{S}[i', j_i^1] \neq \mathbf{S}[i', j_i^2]$.
- Let $i \neq i' \in I_2$ be such that $j_i^2 = j_{i'}^1$. Note that $\mathbf{S}[i, j_i^1]$ and $\mathbf{S}[i', j_{i'}^1]$ are completed by the same value, if and only if $\mathbf{S}[i, j_i^1]$ and $\mathbf{S}[i', j_{i'}^1]$ are completed by the same value. Hence, the following must hold for each $i \neq i' \in I_2$ with $j_i^2 = j_{i'}^1$:
 - $d(\mathbf{S}[i], \mathbf{S}[i']) \in \{\alpha - 2, \alpha\}$ if $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i', j_i^2]$.
 - $d(\mathbf{S}[i], \mathbf{S}[i']) \in \{\alpha - 3, \alpha - 1\}$ if $\mathbf{S}[i, j_{i'}^1] \neq \mathbf{S}[i', j_i^2]$.
- $d(\mathbf{S}[i], \mathbf{S}[i']) \in \{\alpha - 2, \alpha\}$ for each $i \neq i' \in I_2$ with $j_i^1 = j_{i'}^1$ and $j_i^2 = j_{i'}^2$.

We return **No** if at least one of the above fails. Clearly this requires $O(n^2\ell)$ time.

Now, we construct a 2-CNF formula which is satisfiable if and only if our DMC instance is a **Yes**-instance. We introduce a variable x_i for each $i \in I_2$, which basically encodes the completion of $\mathbf{S}[i]$. Intuitively speaking, $(\mathbf{S}[i, j_i^1], \mathbf{S}[i, j_i^2])$ are filled by $(1, 0)$ if x_i is true and by $(0, 1)$ if x_i is false. We add clauses as follows:

- For each $i_0 \in I_0$ and $i \in I_2$ with $d(\mathbf{S}[i_0], \mathbf{S}[i]) = \alpha - 2$, add a singleton clause (x_i) if $(\mathbf{S}[i_0, j_i^1], \mathbf{S}[i_0, j_i^2]) = (0, 1)$ and $(\neg x_i)$ otherwise.
- For each $i \neq i' \in I_2$ with $\{j_i^1, j_i^2\} \cap \{j_{i'}^1, j_{i'}^2\} = \emptyset$, $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i, j_{i'}^2]$, and $\mathbf{S}[i, j_{i'}^1] \neq \mathbf{S}[i, j_{i'}^2]$, add a clause (x_i) if either $(\mathbf{S}[i, j_i^1], \mathbf{S}[i, j_i^2]) = (0, 1)$ and $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 3$, or $(\mathbf{S}[i, j_i^1], \mathbf{S}[i, j_i^2]) = (1, 0)$ and $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 1$. Analogously, we add a clause $(x_{i'})$ or $(\neg x_{i'})$ for the case $\{j_{i'}^1, j_{i'}^2\} \cap \{j_i^1, j_i^2\} = \emptyset$, $\mathbf{S}[i', j_i^1] = \mathbf{S}[i', j_i^2]$, and $\mathbf{S}[i', j_i^1] \neq \mathbf{S}[i', j_i^2]$.
- We do as follows for each $i \neq i' \in I_2$ with $\{j_i^1, j_i^2\} \cap \{j_{i'}^1, j_{i'}^2\} = \emptyset$, $\mathbf{S}[i, j_{i'}^1] \neq \mathbf{S}[i, j_{i'}^2]$, and $\mathbf{S}[i, j_{i'}^1] \neq \mathbf{S}[i, j_{i'}^2]$.
 - Suppose that $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 4$. Add a clause (x_i) if $(\mathbf{S}[i', j_i^1], \mathbf{S}[i', j_i^2]) = (0, 1)$ and $(\neg x_i)$ otherwise. Moreover, add a clause $(x_{i'})$ if $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i, j_{i'}^2]) = (0, 1)$ and $(\neg x_{i'})$ otherwise.
 - Suppose that $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 2$. If $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i, j_{i'}^2]) = (\mathbf{S}[i', j_i^1], \mathbf{S}[i', j_i^2])$, then add clauses $(x_i \vee x_{i'})$ and $(\neg x_i \vee \neg x_{i'})$. Note that these clauses are satisfied if and only if $x_i \neq x_{i'}$. Otherwise, we add clauses $(x_i \vee \neg x_{i'})$ and $(\neg x_i \vee x_{i'})$, which are satisfied if and only if $x_i = x_{i'}$.
 - Suppose that $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha$. Add a clause (x_i) if $(\mathbf{S}[i', j_i^1], \mathbf{S}[i', j_i^2]) = (1, 0)$ and $(\neg x_i)$ otherwise. Moreover, add a clause $(x_{i'})$ if $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i, j_{i'}^2]) = (1, 0)$ and $(\neg x_{i'})$ otherwise.
- We do as follows for each $i \neq i' \in I_2$ with $j_i^2 = j_{i'}^1$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 2$ and $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i', j_i^2] = 0$, then add a clause $(x_i \vee \neg x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha$ and $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i', j_i^2] = 0$, then add clauses $(\neg x_i)$ and $(x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 2$ and $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i', j_i^2] = 1$, then add a clause $(\neg x_i \vee x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha$ and $\mathbf{S}[i, j_{i'}^1] = \mathbf{S}[i', j_i^2] = 1$, then add clauses (x_i) and $(\neg x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 3$ and $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i', j_i^2]) = (1, 0)$, then add clauses (x_i) and $(x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 1$ and $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i', j_i^2]) = (1, 0)$, then add a clause $(\neg x_i \vee \neg x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 3$ and $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i', j_i^2]) = (0, 1)$, then add clauses $(\neg x_i)$ and $(\neg x_{i'})$.
 - If $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 1$ and $(\mathbf{S}[i, j_{i'}^1], \mathbf{S}[i', j_i^2]) = (0, 1)$, then add a clause $(x_i \vee x_{i'})$.
- For each $i \neq i' \in I_2$ with $j_i^1 = j_{i'}^1$ and $j_i^2 = j_{i'}^2$, add clauses $(x_i \vee \neg x_{i'})$ and $(\neg x_i \vee x_{i'})$ if $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha - 2$ and add clauses $(x_i \vee x_{i'})$ and $(\neg x_i \vee \neg x_{i'})$ if $d(\mathbf{S}[i], \mathbf{S}[i']) = \alpha$.

It is easy to check that the constructed formula is correct. The formula contains $O(n^2)$ clauses and can thus be solved in $O(n^2)$ time [1]. ◀

4.3 NP-hardness

In this subsection we prove two NP-hardness results. Namely, we prove that DMC is NP-hard for $k = 2$ and $\beta \geq \alpha + 3$ (Theorem 28) and for $k = 3$ and $\beta = \alpha$ (Theorem 33). To start with, we introduce a tool to increase the distance of one specific pair of row vectors relative to all other pairs. This will be useful for the subsequent reductions.

▶ **Lemma 27.** *For each $n \geq 3$ and each $i < i' \in [n]$, one can construct in $n^{O(1)}$ time, a matrix $\mathbf{B}_{i,i'}^n \in \{0, 1\}^{n \times \ell}$ with n rows and $\ell = \binom{n}{2} - 1$ columns such that*

$$d(\mathbf{B}_{i,i'}^n[h], \mathbf{B}_{i,i'}^n[h']) = \begin{cases} \gamma(\mathbf{B}_{i,i'}^n) + 2 & \text{if } (h, h') = (i, i'), \\ \gamma(\mathbf{B}_{i,i'}^n) & \text{otherwise.} \end{cases}$$

for all $h \neq h' \in [n]$.

Proof. First, we define a binary matrix $\mathbf{A}^n \in \{0, 1\}^{n \times (2n-1)}$ as follows:

$$\mathbf{A}^n := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 1 & & & & & & & & \\ 1 & 1 & 1 & & \mathbf{I} & & & \mathbf{I} & & & \\ & & \vdots & & & & & & & & \\ 1 & 1 & 1 & & & & & & & & \end{bmatrix},$$

where \mathbf{I} is the $(n-2) \times (n-2)$ identity matrix. Note that $d(\mathbf{A}^n[1], \mathbf{A}^n[2]) = 2$ and $d(\mathbf{A}^n[h], \mathbf{A}^n[h']) = 4$ for all $h < h' \in [n]$ with $(h, h') \neq (1, 2)$. We also define the matrix $\mathbf{A}_{h,h'}^n$ obtained from \mathbf{A}^n by swapping the row vectors $\mathbf{A}^n[1]$ (and $\mathbf{A}^n[2]$) with $\mathbf{A}^n[h]$ (and $\mathbf{A}^n[h']$, respectively) for each $h < h' \in [n]$. The matrix $\mathbf{A}_{h,h'}^n$ is a matrix in which the distance between the h -th and h' -th row vectors are exactly two smaller than all other pairs. Now we use the matrix $\mathbf{A}_{h,h'}^n$ to obtain a binary matrix in which the distance of a certain pair of row vectors is exactly two greater than all others. We define $\mathbf{B}_{i,i'}^n \in \{0, 1\}^{n \times \ell}$ as the matrix obtained by horizontally stacking $\binom{n}{2} - 1$ matrices $\mathbf{A}_{h,h'}^n$ for all $h < h' \in [n]$ with $(h, h') \neq (i, i')$:

$$\mathbf{B}_{i,i'}^n := [\mathbf{A}_{1,1}^n \cdots \mathbf{A}_{1,n}^n \cdots \mathbf{A}_{i,i+1}^n \cdots \mathbf{A}_{i,i'-1}^n \mathbf{A}_{i,i'+1}^n \cdots \mathbf{A}_{i,n}^n \cdots \mathbf{A}_{n-1,n}^n].$$

Observe that $d(\mathbf{B}_{i,i'}^n[i], \mathbf{B}_{i,i'}^n[i']) = 4 \cdot (\binom{n}{2} - 1) = 2n(n-1) - 4$, since $d(\mathbf{A}_{i,i'}^n[h], \mathbf{A}_{i,i'}^n[h']) = 4$ for all $h < h' \in [n]$ with $(h, h') \neq (i, i')$. Note also that for each $h < h' \in [n]$ with $(h, h') \neq (i, i')$, we have $d(\mathbf{B}_{i,i'}^n[h], \mathbf{B}_{i,i'}^n[h']) = 2n(n-1) - 6$ because the distance between $\mathbf{A}_{i,i'}^n[\tilde{h}]$ and $\mathbf{A}_{i,i'}^n[\tilde{h}']$ is four for every $\tilde{h} < \tilde{h}' \in [n]$ except that it is smaller by two for the pair $\mathbf{A}_{i,i'}^n[i]$ and $\mathbf{A}_{i,i'}^n[i']$. It is easy to see that the matrix $\mathbf{B}_{i,i'}^n$ can be constructed in polynomial time. ◀

We now prove that DMC is NP-hard for $k = 2$ and $\alpha + 3 \leq \beta$. Our proof is based on a polynomial-time reduction from the following NP-hard variant of 3-SAT [2].

(3, B2)-SAT

Input: A Boolean formula in conjunctive normal form, in which each literal occurs exactly twice and each clause contains exactly three literals of distinct variables.

Question: Is there a satisfying truth assignment?

The most challenging technical aspect of the reduction is to ensure the upper and lower bounds on the Hamming distances. We achieve this by making heavy use of Lemma 27.

► **Theorem 28.** *DMC is NP-hard for $k = 2$ and $\alpha + 3 \leq \beta$.*

Proof. We reduce from (3, B2)-SAT. We divide our proof into two parts as follows. We first provide a set \mathcal{C} of incomplete matrices and describe rules under which the matrices of \mathcal{C} are completed. We prove that the given formula of (3, B2)-SAT is satisfiable if and only if the matrices \mathcal{C} can be completed under those rules. We then show that one can construct in polynomial time a single incomplete matrix \mathbf{S} containing each matrix in \mathcal{C} as a submatrix, such that \mathbf{S} admits a solution if and only if the completions to \mathcal{C} according to the rules are feasible. We are going to exploit the matrix $\mathbf{B}_{i,i'}^n$ of Lemma 27 for this construction.

Part I.

Let ϕ be an instance of (3, B2)-SAT with clauses C_0, \dots, C_{m-1} . We define the following matrix for each clause C_i

$$\mathbf{C}_i := \begin{bmatrix} l_i^1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & l_i^2 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & l_i^3 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 & c_i \end{bmatrix}.$$

Here we use l_i^1, l_i^2, l_i^3 , and c_i to represent two missing entries for notational purposes. Note that the matrices \mathbf{C}_i are identical for all $i \in [0, m-1]$. We will prove that ϕ is satisfiable if and only if it is possible to complete matrices $\mathcal{C} := \{\mathbf{C}_i \mid i \in [0, m-1]\}$ satisfying the following constraints:

1. The missing entries l_i^j are filled by 10 or 01 for each $i \in [0, m-1]$ and $j \in [3]$.
2. The missing entries c_i are filled by 00, 01, or 10 for each $i \in [0, m-1]$.
3. If the missing entries c_i are filled by 00 (01, 10), then l_i^1 (l_i^2, l_i^3 , respectively) are filled by 10 for each $i \in [0, m-1]$.
4. Let \mathcal{Z} be the set such that $(i, j, i', j') \in \mathcal{Z}$ if and only if the j -th literal in C_i and the j' -th literal in $C_{i'}$ correspond to the same variable and one is the negation of the other for each $i < i' \in [0, m-1]$ and $j, j' \in [3]$. If $(i, j, i', j') \in \mathcal{Z}$, then either l_i^j or $l_{i'}^{j'}$ is filled by 01.

Note that there are three choices for the completion of c_i by Constraint 2. The intuitive idea is that the completion of c_i dictates which literal (in binary encoding) in the clause C_i is satisfied. We then obtain a satisfying truth assignment for ϕ , as we shall see in the following claim.

► **Claim 29.** The formula ϕ is satisfiable if and only if the matrices \mathcal{C} can be completed according to Constraints 1 to 4.

Proof. (\Rightarrow) If there exists a truth assignment τ satisfying ϕ , then at least one literal in the clause C_i evaluates to true for each $i \in [0, m-1]$. We choose an arbitrary number $l_i \in [3]$ such that the l_i -th literal of C_i is satisfied in τ for each $i \in [0, m-1]$. For each $i \in [0, m-1]$ we complete the matrix C_i as follows:

- If $l_i = 1$, then the missing entries c_i, l_i^1, l_i^2, l_i^3 are filled by 00, 10, 01, 01, respectively.
- If $l_i = 2$, then the missing entries c_i, l_i^1, l_i^2, l_i^3 are filled by 01, 01, 10, 01, respectively.
- If $l_i = 3$, then the missing entries c_i, l_i^1, l_i^2, l_i^3 are filled by 10, 01, 01, 10, respectively.

$$\begin{array}{c}
\left[\begin{array}{cccccccc}
l_i^1 & 0 & 0 & 0 & 0 & 1 & 1 \\
0 & 0 & l_i^2 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & l_i^3 & 0 & 1 \\
1 & 0 & 1 & 0 & 1 & 0 & c_i \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\
1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\
0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\
1 & 0 & 1 & 0 & 1 & 0 & 0 & 0
\end{array} \right]
\end{array}
\quad
\begin{array}{cccccccc}
\vdots & & & & & & & \vdots \\
\cdots & l_i^j & \cdots & 00 & \cdots & 00 & \cdots & 01 & \cdots \\
\vdots & & & & & & & \vdots & \\
\vdots & & & & & & & \vdots & \\
00 & & & & & & & & 00 \\
\vdots & & & & \ddots & & & \vdots & \\
00 & & & & & & & & 00 \\
\vdots & & & & & & & \vdots & \\
\vdots & & & & & & & \vdots & \\
\cdots & 01 & \cdots & 00 & \cdots & 00 & \cdots & l_{i'}^{j'} & \cdots \\
\vdots & & & & & & & \vdots &
\end{array}$$

■ **Figure 7** The matrix $\mathbf{C}'_i \in \{0, 1, \square\}^{11 \times 8}$ (left). The rows $\{11i + j, 11i' + j'\}$ and the columns $\{8i + 2j - 1, 8i + 2j, 8j' + 2j' - 1, 8j' + 2j'\}$ of \mathbf{C} for $(i, j, i', j') \in \mathcal{Z}$ (right).

It is easy to verify that Constraints 1 to 3 are satisfied. We claim that Constraint 4 is also satisfied. Suppose to the contrary that there exists an $(i, j, i', j') \in \mathcal{Z}$ such that the missing entries l_i^j and $l_{i'}^{j'}$ are both filled by 10. Then, we have $l_i = j$ and $l_{i'} = j'$, meaning that τ satisfies both x and $\neg x$ (a contradiction).

(\Leftarrow) For each $i \in [0, m - 1]$ and $j \in [3]$ where l_i^j is filled by 10, we construct a truth assignment such that the j -th literal of C_i is satisfied. No variable is given opposing truth values by such a truth assignment because of Constraint 4. It also satisfies every clause: Otherwise, there exists an integer $i \in [0, m - 1]$ such that all l_i^1, l_i^2, l_i^3 are completed by 01 due to Constraint 1. Now we have a contradiction because Constraints 2 and 3 imply that at least one of l_i^1, l_i^2, l_i^3 is filled by 10. \triangleleft

Part II.

We provided matrices \mathcal{C} as well as the constraints on the completion of \mathcal{C} in Part I. Now, we describe how to construct a matrix \mathbf{S} that admits a completion \mathbf{T} with $\gamma(\mathbf{T}) \geq \alpha$ and $\delta(\mathbf{T}) \leq \beta$ if and only if \mathcal{C} can be completed fulfilling Constraints 1 to 4.

First, we introduce a matrix $\mathbf{C}'_i \in \{0, 1, \square\}^{11 \times 8}$ obtained from \mathbf{C}_i by adding row vectors as follows (see Figure 7):

- The first four row vectors of \mathbf{C}'_i are identical to the row vectors of \mathbf{C}_i .
- The row vectors $\mathbf{C}'_i[5]$, $\mathbf{C}'_i[6]$, and $\mathbf{C}'_i[7]$ are obtained by completing the missing entries in $\mathbf{C}_i[1]$, $\mathbf{C}_i[2]$, and $\mathbf{C}_i[3]$, respectively, with 00.
- The row vectors $\mathbf{C}'_i[8]$, $\mathbf{C}'_i[9]$, and $\mathbf{C}'_i[10]$ are obtained by completing the missing entries in $\mathbf{C}_i[1]$, $\mathbf{C}_i[2]$, and $\mathbf{C}_i[3]$, respectively, with 11.
- The row vector $\mathbf{C}'_i[11]$ is obtained by completing the missing entries in $\mathbf{C}_i[4]$ with 00.

Next, we construct a matrix $\mathbf{C} \in \{0, 1, \square\}^{11m \times 8m}$ from the matrices \mathbf{C}'_i as follows (see also Figure 7): We start with an empty matrix of size $11m \times 8m$. We first place $\mathbf{C}'_0, \dots, \mathbf{C}'_{m-1}$ on the diagonal. Then, we place 01 at the intersection of the row containing l_i^j ($l_{i'}^{j'}$) and the columns containing $l_{i'}^{j'}$ (l_i^j , respectively), for each $(i, j, i', j') \in \mathcal{Z}$. Finally, let the remaining entries be all 0. The formal definition is given as follows:

- $\mathbf{C}[[11i + 1, 11i + 1], [8i + 1, 8i + 8]] = \mathbf{C}'_i$ for each $i \in [0, m - 1]$.
- $\mathbf{C}[11i + j, 8i' + 2j'] = 1$ and $\mathbf{C}[11i' + j', 8i + 2j] = 1$ for each $(i, j, i', j') \in \mathcal{Z}$.
- All other entries are 0.

Let $n = 11m$ be the number of rows in \mathbf{C} . We show in the next claim that $\delta(\mathbf{C}) \leq 8$.

▷ **Claim 30.** $d(\mathbf{C}[h], \mathbf{C}[h']) \leq 8$ for each $h < h' \in [n]$.

Proof. It suffices to show that $d(\mathbf{C}[h], 0^{8m}) \leq 4$ for each $h \in [n]$, because we then have $d(\mathbf{C}[h], \mathbf{C}[h']) \leq d(\mathbf{C}[h], 0^{8m}) + d(\mathbf{C}[h'], 0^{8m}) \leq 8$ by the triangle inequality. Suppose that $h = 11i + j$ for $i \in [0, m-1]$ and $j \in [3]$. Then, the row vector $\mathbf{C}[h]$ contains at most two 1's in \mathbf{C}'_i , and exactly two 1's elsewhere because each literal appears in the formula ϕ exactly twice. It follows that $\mathbf{C}[h]$ contains at most four 1's. Hence, we can assume that $h = 11i + j$ for $i \in [0, m-1]$ and $j \in [4, 11]$. Then, the row vector $\mathbf{C}[h]$ contains at most four 1's, because all 1's appear in \mathbf{C}'_i . This shows the claim. ◁

Now we define seven “types” H_1, \dots, H_7 of row index pairs. For each $h < h' \in [n]$,

- $(h, h') \in H_1$ if $h = 11i + j$ and $h' = 11i + j'$ for some $i \in [0, m-1]$ and $(j, j') \in \{(1, 5), (2, 6), (3, 7), (1, 8), (2, 9), (3, 10)\}$.
- $(h, h') \in H_2$ if $h = 11i + j$ and $h' = 11i + j'$ for some $i \in [0, m-1]$ and $(j, j') = (4, 11)$.
- $(h, h') \in H_3$ if $h = 11i + j$ and $h' = 11i + j'$ for some $i \in [0, m-1]$ and $(j, j') \in \{(1, 4), (2, 4), (3, 4)\}$.
- $(h, h') \in H_4$ if $h = 11i + j$ and $h' = 11i' + j'$ for $(i, j, i', j') \in \mathcal{Z}$.
- For each $h < h' \in [n]$ with $(h, h') \notin H_1, \dots, H_4$,
 - $(h, h') \in H_5$ if both $\mathbf{C}[h]$ and $\mathbf{C}[h']$ have missing entries.
 - $(h, h') \in H_6$ if exactly one of $\mathbf{C}[h]$ and $\mathbf{C}[h']$ have missing entries.
 - $(h, h') \in H_7$ if neither of $\mathbf{C}[h]$ and $\mathbf{C}[h']$ has missing entries.

For each type of row index pairs, we adjust the pairwise distances using Lemma 27.

▷ **Claim 31.** There exists a $\beta \in \mathbb{N}$ and a complete matrix \mathbf{D} over $\{0, 1\}$ with n rows such that all of the following hold for $\mathbf{S} = [\mathbf{C} \ \mathbf{D}]$:

- $d(\mathbf{S}[h], \mathbf{S}[h']) = \beta - 1$ for each $(h, h') \in H_1$ (cf. Constraint 1).
- $d(\mathbf{S}[h], \mathbf{S}[h']) = \beta - 1$ for each $(h, h') \in H_2$ (cf. Constraint 2).
- $d(\mathbf{S}[h], \mathbf{S}[h']) = \beta - 3$ for each $(h, h') \in H_3$ (cf. Constraint 3).
- $d(\mathbf{S}[h], \mathbf{S}[h']) \in \{\beta - 3, \beta - 2\}$ for each $(h, h') \in H_4$ (cf. Constraint 4).
- $d(\mathbf{S}[h], \mathbf{S}[h']) \in \{\beta - 3, \beta - 2\}$ for each $(h, h') \in H_5$.
- $d(\mathbf{S}[h], \mathbf{S}[h']) \in \{\beta - 2, \beta - 1\}$ for each $(h, h') \in H_6$.
- $d(\mathbf{S}[h], \mathbf{S}[h']) \in \{\beta - 1, \beta\}$ for each $(h, h') \in H_7$.

Proof. We obtain the matrix \mathbf{D} by horizontally stacking $\mathbf{B}_{h,h'}^n$ of Lemma 27 $c_{h,h'}$ times (where $c_{h,h'} \in \mathbb{N}$ is to be defined) for each $h < h' \in [n]$. Recall that $d(\mathbf{B}_{h,h'}^n[i], \mathbf{B}_{h,h'}^n[i'])$ equals $\gamma(\mathbf{B}_{h,h'}^n)$ if $(h, h') = (i, i')$ and $\gamma(\mathbf{B}_{h,h'}^n) + 2$ otherwise) and let

$$\beta = \sum_{h < h' \in [n]} c_{h,h'} \cdot \gamma(\mathbf{B}_{h,h'}^n) + 11.$$

Observe that the pairwise row distance can be rewritten as follows for each $h < h' \in [n]$:

$$\begin{aligned} d(\mathbf{S}[h], \mathbf{S}[h']) &= d(\mathbf{C}[h], \mathbf{C}[h']) + c_{h,h'} \cdot (\gamma(\mathbf{B}_{h,h'}^n) + 2) + \sum_{\substack{i < i' \in [n], \\ (i, i') \neq (h, h')}} c_{h,h'} \cdot \gamma(\mathbf{B}_{h,h'}^n) \\ &= d(\mathbf{C}[h], \mathbf{C}[h']) + 2c_{h,h'} + \beta - 11. \end{aligned}$$

We define $c_{h,h'}$ for each $(h, h') \in H_1 \cup H_2 \cup H_3$ as follows.

- Let $c_{h,h'} = 4$ for each $(h, h') \in H_1$. Then, we have $d(\mathbf{S}[h], \mathbf{S}[h']) = 2 + 2 \cdot 4 - \beta - 11 = \beta - 1$.

- Let $c_{h,h'} = 5$ for each $(h, h') \in H_2$. Then, we have $d(\mathbf{S}[h], \mathbf{S}[h']) = 0 + 2 \cdot 5 - \beta - 11 = \beta - 1$.
- Let $c_{h,h'} = 2$ for each $(h, h') \in H_3$. Then, we have $d(\mathbf{S}[h], \mathbf{S}[h']) = 4 + 2 \cdot 2 - \beta - 11 = \beta - 3$.

For the remainder (that is, $(h, h') \in H_4 \cup \dots \cup H_7$), it has to be shown that there exists $c_{h,h'} \in \mathbb{N}$ such that $d(\mathbf{S}[h], \mathbf{S}[h']) \in \{x, x + 1\}$ holds for $x \in \mathbb{N}$ with $x \geq \beta - 3$. Let $c_{h,h'} = \lceil (11 + x - \beta - d(\mathbf{C}[h], \mathbf{C}[h']))/2 \rceil$. Clearly, $c_{h,h'}$ is an integer and it holds that $c_{h,h'} \geq 0$ because $x - \beta \geq -3$ and $d(\mathbf{C}[h], \mathbf{C}[h']) \leq 8$ by Claim 30. Moreover, we have $d(\mathbf{S}[h], \mathbf{S}[h']) = x$ if $11 + x - \beta - d(\mathbf{C}[h], \mathbf{C}[h'])$ is even and $d(\mathbf{S}[h], \mathbf{S}[h']) = x + 1$ otherwise. \triangleleft

Finally, we show that Constraints 1 to 4 are essentially the same as the pairwise row distance constraints on the matrix \mathbf{S} of Claim 31.

\triangleright **Claim 32.** The matrices \mathcal{C} can be completed according to Constraints 1 to 4, if and only if \mathbf{S} admits a completion \mathbf{T} with $\delta(\mathbf{T}) \leq \beta$.

Proof. (\Rightarrow) Let \mathbf{T} be the matrix where the missing entries of \mathbf{S} are filled as in the completion of \mathcal{C} . We show that $d(\mathbf{T}[h], \mathbf{T}[h']) \leq \beta$ holds for each $h < h' \in [n]$.

- Suppose that $(h, h') \in H_1$. Then, the missing entries in $\mathbf{S}[h]$ are filled by 10 or 01 by Constraint 1 and $\mathbf{S}[h']$ (which has no missing entries) has 00 or 11 in the corresponding positions. Hence, $d(\mathbf{T}[h], \mathbf{T}[h']) = d(\mathbf{S}[h], \mathbf{S}[h']) + 1 = \beta$.
- Suppose that $(h, h') \in H_2$. Then, the missing entries in $\mathbf{S}[h]$ are filled by 00, 01, or 10 by Constraint 2 and $\mathbf{S}[h']$ (which has no missing entries) has 00 in the corresponding positions. Hence, $d(\mathbf{T}[h], \mathbf{T}[h']) = d(\mathbf{S}[h], \mathbf{S}[h']) + 1 = \beta$.
- Suppose that $(h, h') \in H_3$. Note that $\mathbf{S}[h]$ has missing entries l_i^j and $\mathbf{S}[h']$ has missing entries c_i for $i \in [0, m - 1]$ and $j \in [3]$. Let $c_i^1 c_i^2$ be the completion of c_i for $c_i^1, c_i^2 \in \{0, 1\}$. If $\mathbf{T}[h]$ has $1 - c_i^1$ and $1 - c_i^2$ in the corresponding positions, then $d(\mathbf{T}[h], \mathbf{T}[h']) = d(\mathbf{S}[h], \mathbf{S}[h']) + 2 = \beta - 1$. Otherwise, $\mathbf{T}[h]$ matches in at least one position where $\mathbf{T}[h']$ has missing entries c_i . Therefore, $d(\mathbf{T}[h], \mathbf{T}[h']) \leq d(\mathbf{S}[h], \mathbf{S}[h']) + 3 = \beta$.
- Suppose that $(h, h') \in H_4$. Note that $\mathbf{S}[h]$ has missing entries l_i^j and $\mathbf{S}[h']$ has missing entries $l_{i'}^{j'}$. Also note that $\mathbf{S}[h]$ and $\mathbf{S}[h']$ have 01 where the other row vector has missing entries. Since either l_i^j or $l_{i'}^{j'}$ must be completed by 01 due to Constraint 4, we have $d(\mathbf{T}[h], \mathbf{T}[h']) = d(\mathbf{S}[h], \mathbf{S}[h']) \leq \beta - 2 + 2 = \beta$.
- Suppose that $(h, h') \in H_5 \cup H_6 \cup H_7$. Let $x \in \{0, 1, 2\}$ be the number of row vectors with missing entries in $\{\mathbf{S}[h], \mathbf{S}[h']\}$. Then, we have $d(\mathbf{S}[h], \mathbf{S}[h']) \in \{\beta - x - 1, \beta - x\}$. If $\mathbf{S}[h]$ has missing entries, then $\mathbf{S}[h']$ has 00 in the corresponding positions, and vice versa. Since the missing entries are filled by 00, 01, or 10 according to Constraint 1 and 2, we have $d(\mathbf{T}[h], \mathbf{T}[h']) \leq d(\mathbf{S}[h], \mathbf{S}[h']) + x = \beta$.

(\Leftarrow) We complete the matrices in \mathcal{C} in the same way as in the completion of \mathbf{S} . It is easy to verify all Constraints 1 to 4 are satisfied. \triangleleft

Note that $\gamma(\mathbf{T}) \geq \gamma(\mathbf{S}) \geq \beta - 3$ for any completion \mathbf{T} of \mathbf{S} . Hence, it follows from Claims 29 and 32 that ϕ is satisfiable if and only if the DMC instance $(\mathbf{S}, \alpha, \beta)$ is a **Yes**-instance, for any $\alpha \leq \beta - 3$. This concludes the proof of Theorem 28. \blacktriangleleft

To prove that DMC is NP-hard for $\alpha = \beta$ and $k = 3$, we provide a polynomial-time reduction from another NP-hard variant of 3-SAT [17]:

CUBIC MONOTONE 1-IN-3 SAT

Input: A Boolean formula in conjunctive normal form, in which each variable appears exactly three times and each clause contains exactly three distinct positive literals.

Question: Is there a truth assignment that satisfies exactly one literal in each clause?

$$\mathbf{C} = \left[\begin{array}{cccc|cccc|c} a_1 & 00 & 00 & 00 & 011 & 011 & 011 & 000 & 1 \\ 00 & a_2 & 00 & 00 & 101 & 101 & 000 & 011 & 1 \\ 00 & 00 & a_3 & 00 & 110 & 000 & 101 & 101 & 1 \\ 00 & 00 & 00 & a_4 & 000 & 110 & 110 & 110 & 1 \\ \hline 10 & 10 & 10 & 00 & b_1 & 000 & 000 & 000 & 0 \\ 10 & 10 & 00 & 10 & 000 & b_2 & 000 & 000 & 0 \\ 10 & 00 & 10 & 10 & 000 & 000 & b_3 & 000 & 0 \\ 00 & 10 & 10 & 10 & 000 & 000 & 000 & b_4 & 0 \\ \hline 00 & 00 & 00 & 00 & 000 & 000 & 000 & 000 & 0 \end{array} \right]$$

■ **Figure 8** An example of \mathbf{C} for $\phi = (x_1 \vee x_2 \vee x_3) \wedge (x_1 \vee x_2 \vee x_4) \wedge (x_1 \vee x_3 \vee x_4) \wedge (x_2 \vee x_3 \vee x_4)$.

Our reduction heavily depends on the fact that $\alpha = \beta$. This is contrary to the reduction in the proof of Theorem 28, which in fact works for any $\alpha \leq \beta - 3$.

► **Theorem 33.** *DMC is NP-hard for $k = 3$ and $\alpha = \beta$.*

Proof. Let ϕ be an instance of CUBIC MONOTONE 1-IN-3 SAT. Our proof has two parts: First, we provide an incomplete matrix \mathbf{C} and we show that ϕ is a **Yes**-instance if and only if \mathbf{C} can be completed under certain constraints. Then, we obtain an instance $(\mathbf{S}, \alpha, \alpha)$ of DMC by adjusting the pairwise row distances with the help of Lemma 27.

Suppose that ϕ contains variables x_1, \dots, x_m and clauses C_1, \dots, C_m , where $C_i = (C_i^1 \vee C_i^2 \vee C_i^3)$ for each $i \in [m]$. First, we define matrices $\mathbf{C}_1, \mathbf{C}_3 \in \{0, 1, \square\}^{m \times 2m}$ and $\mathbf{C}_2, \mathbf{C}_4 \in \{0, 1, \square\}^{m \times 3m}$. We use a_i (and b_i) to represent two (three, respectively) missing entries in \mathbf{C}_1 (\mathbf{C}_4 , respectively) for each $i \in [m]$. For each $i \in [m]$ and $j \in [m]$, let

$$\mathbf{C}_1[i, \{2j-1, 2j\}] = \begin{cases} a_i & \text{if } i = j, \\ 00 & \text{otherwise.} \end{cases} \quad \mathbf{C}_2[i, \{3j-2, 3j\}] = \begin{cases} 011 & \text{if } x_i = C_j^1, \\ 101 & \text{if } x_i = C_j^2, \\ 110 & \text{if } x_i = C_j^3, \\ 000 & \text{otherwise.} \end{cases}$$

$$\mathbf{C}_3[i, \{2j-1, 2j\}] = \begin{cases} 10 & \text{if } x_i \text{ is in } C_j, \\ 00 & \text{otherwise.} \end{cases} \quad \mathbf{C}_4[i, \{3j-2, 3j\}] = \begin{cases} b_i & \text{if } i = j, \\ 000 & \text{otherwise.} \end{cases}$$

We obtain an incomplete matrix $\mathbf{C} \in \{0, 1, \square\}^{(2m+1) \times (5m+1)}$ by appending a column vector $(0^m 1^m)^T$ and a row vector 0^{5m+1} to the following matrix

$$\begin{bmatrix} \mathbf{C}_1 & \mathbf{C}_2 \\ \mathbf{C}_3 & \mathbf{C}_4 \end{bmatrix}.$$

Refer to Figure 8 for an illustration.

Intuitively speaking, we will use the first m rows to encode the variables and the following m rows to encode the clauses.

► **Claim 34.** There is a truth assignment that satisfies exactly one literal in clause C_i for each $i \in [m]$ if and only if there is a completion \mathbf{C}' of \mathbf{C} such that

1. $d(\mathbf{C}'[i], \mathbf{C}[2m+1]) = d(\mathbf{C}[i], \mathbf{C}[2m+1]) + 1$ for each $i \in [2m]$.
2. $d(\mathbf{C}'[i], \mathbf{C}'[m+i']) = d(\mathbf{C}[i], \mathbf{C}[m+i']) + 3$ for each $i, i' \in [m]$ such that x_i is in $C_{i'}$.

Proof. (\Rightarrow) Let τ be a truth assignment satisfying exactly one literal in each clause of ϕ . Consider the matrix \mathbf{C}' obtained by completing \mathbf{C} as follows for each $i \in [m]$:

- The missing entries a_i are filled by 10 if x_i is true in τ and by 01 otherwise.
- The missing entries b_i are filled by 100 if C_i^1 is true in τ .
- The missing entries b_i are filled by 010 if C_i^2 is true in τ .
- The missing entries b_i are filled by 001 if C_i^3 is true in τ .

It is easy to see that the first constraint of the claim is indeed fulfilled. For the other constraint, consider $i, i' \in [m]$ such that x_i is in $C_{i'}$. We prove that $d(\mathbf{C}'[i], \mathbf{C}'[m+i']) - d(\mathbf{C}[i], \mathbf{C}[m+i']) = 3$ or equivalently,

$$d(a'_i, \mathbf{C}[m+i', [2i'-1, 2i']]) + d(\mathbf{C}[i, [2m+3i'-2, 2m+3i']], b'_{i'}) = 3,$$

where a'_i and $b'_{i'}$ are the completion of a_i and $b_{i'}$ in \mathbf{C}' . We show that it holds for the case $x_i = C_i^1$. It can be proven analogously for the cases of $x_i = C_i^2$ and $x_i = C_i^3$ as well.

Note that $\mathbf{C}[m+i', [2i'-1, 2i']] = 10$ and $\mathbf{C}[i, [2m+3i'-2, 2m+3i']] = 011$. If x_i is true in τ , then a'_i and $b'_{i'}$ are 10 and 100, respectively. Thus, the equality above holds. If x_i is false in τ , then $a'_i = 01$ and $b'_{i'} \in \{010, 001\}$. Again the equality above holds.

(\Leftarrow) Let a'_i and b'_i be the completion of a_i and b_i in \mathbf{C}' for each $i \in [m]$. Due to the first constraint, exactly one entry in a'_i and b'_i must be 1 for each $i \in [m]$. Now consider the truth assignment τ that assigns x_i to true if $a'_i = 10$ and false if $a'_i = 01$ for each $i \in [m]$. We show that τ satisfies exactly one literal in each clause of ϕ . Consider $i \in [m]$ with $C_i = (x_{i_1} \vee x_{i_2} \vee x_{i_3})$. By the second constraint, we have

$$\sum_{j=1}^3 d(\mathbf{C}'[i_j], \mathbf{C}'[m+i]) - d(\mathbf{C}[i_j], \mathbf{C}[m+i]) = 9.$$

Rewriting the left-hand side in terms of $a'_{i_1}, a'_{i_2}, a'_{i_3}, b'_i$, we obtain

$$d(011, b'_i) + d(101, b'_i) + d(110, b'_i) + d(10, a'_{i_1}) + d(10, a'_{i_2}) + d(10, a'_{i_3}) = 9.$$

Since $b'_i \in \{100, 010, 001\}$, it follows that the first three terms sum up to exactly 5 and hence $d(10, a'_{i_1}) + d(10, a'_{i_2}) + d(10, a'_{i_3}) = 4$. This means that exactly one of $a'_{i_1}, a'_{i_2}, a'_{i_3}$ is 10 and the remaining two are 01. Thus, exactly one literal in C_i is satisfied. \triangleleft

Now let us note some observations on the pairwise distances in \mathbf{C} (see Figure 8):

- For each $i \in [m]$, $d(\mathbf{C}[i], \mathbf{C}[2m+1]) = 7$ and $d(\mathbf{C}[m+i], \mathbf{C}[2m+1]) = 3$.
- For each $i \neq i' \in [m]$, $d(\mathbf{C}[i], \mathbf{C}[i']) = 12 - 2c_{i,i'}$ and $d(\mathbf{C}[m+i], \mathbf{C}[m+i']) = 6 - 2c'_{i,i'}$. Here $c_{i,i'} \in \{0, 1, 2, 3\}$ is the number of clauses that contain both x_i and $x_{i'}$, and $c'_{i,i'} \in \{0, 1, 2, 3\}$ is the number of variables that are both in C_i and $C_{i'}$.
- For each $i, i' \in [m]$ with x_i in $C_{i'}$, $d(\mathbf{C}[i], \mathbf{C}[m+i']) = 7$.
- For each $i, i' \in [m]$ with x_i not in $C_{i'}$, $d(\mathbf{C}[i], \mathbf{C}[m+i']) = 10$.

We build an incomplete matrix \mathbf{S} with $2m+1$ rows from \mathbf{C} by horizontally appending matrices $\mathbf{B}_{i,i'}^{2m+1}$ of Lemma 27. Recall that $d(\mathbf{B}_{i,i'}^{2m+1}[i], \mathbf{B}_{i,i'}^{2m+1}[i']) = \gamma(\mathbf{B}_{i,i'}^{2m+1}) + 2$ and $d(\mathbf{B}_{i,i'}^{2m+1}[h], \mathbf{B}_{i,i'}^{2m+1}[h']) = \gamma(\mathbf{B}_{i,i'}^{2m+1})$ for all $h < h' \in [2m+1]$ with $(h, h') \neq (i, i')$. For $c \in \mathbb{N}$, let $c\mathbf{B}_{i,i'}$ be the matrix obtained by horizontally stacking $\mathbf{B}_{i,i'}^{2m+1}$ c times. We also compute a value for $\alpha \in \mathbb{N}$ as follows: We start with $\alpha = 14$ and we increase α by $c \cdot \gamma(\mathbf{B}_{i,i'}^{2m+1})$ each time $c\mathbf{B}_{i,i'}$ is appended to \mathbf{C} . We horizontally append the following matrices:

- $3\mathbf{B}_{i,2m+1}$ and $5\mathbf{B}_{m+i,2m+1}$ for each $i \in [m]$.
- $c_{i,i'}\mathbf{B}_{i,i'}$ and $(c'_{i,i'} + 3)\mathbf{B}_{m+i,2m+1}$ for each $i < i' \in [m]$.
- $2\mathbf{B}_{i,m+i'}$ for each $i, i' \in [m]$ with x_i in $C_{i'}$.
- $1\mathbf{B}_{i,m+i'}$ for each $i, i' \in [m]$ with x_i not in $C_{i'}$.

Note that for each $i, i' \in [2m+1]$, $d(\mathbf{S}[i], \mathbf{S}[i']) = d(\mathbf{C}[i], \mathbf{C}[i']) + 2 \cdot n_{i,i'} + \alpha - 14$, where $n_{i,i'}$ is the number of appended $\mathbf{B}_{i,m+i'}$'s. Thus, the pairwise row distances in \mathbf{S} are given as follows:

- For each $i \in [m]$, $d(\mathbf{S}[i], \mathbf{S}[2m+1]) = 7 + 2 \cdot 3 + \alpha - 14 = \alpha - 1$ and $d(\mathbf{S}[m+i], \mathbf{S}[2m+1]) = 3 + 2 \cdot 5 + \alpha - 14 = \alpha - 1$.
- For each $i \neq i' \in [m]$, $d(\mathbf{S}[i], \mathbf{S}[i']) = (12 - 2c_{i,i'}) + 2 \cdot c_{i,i'} + \alpha - 14 = \alpha - 2$ and $d(\mathbf{S}[i], \mathbf{S}[i']) = (6 - 2c'_{i,i'}) + 2 \cdot (c'_{i,i'} + 3) + \alpha - 14 = \alpha - 2$.
- For each $i, i' \in [m]$ with x_i in $C_{i'}$, $d(\mathbf{S}[i], \mathbf{S}[m+i']) = 7 + 2 \cdot 2 + \alpha - 14 = \alpha - 3$.
- For each $i, i' \in [m]$ with x_i not in $C_{i'}$, $d(\mathbf{S}[i], \mathbf{S}[m+i']) = 10 + 2 \cdot 1 + \alpha - 14 = \alpha - 2$.

Finally, we prove that one can complete \mathbf{C} as specified in Claim 34 if and only if one can complete \mathbf{S} into a matrix \mathbf{T} with $\gamma(\mathbf{T}) = \delta(\mathbf{T}) = \alpha$.

▷ **Claim 35.** There is a completion \mathbf{C}' of \mathbf{C} such that

1. $d(\mathbf{C}'[i], \mathbf{C}[2m+1]) = d(\mathbf{C}[i], \mathbf{C}[2m+1]) + 1$ for each $i \in [2m]$.
 2. $d(\mathbf{C}'[i], \mathbf{C}'[m+i']) = d(\mathbf{C}[i], \mathbf{C}[m+i']) + 3$ for each $i, i' \in [m]$ such that x_i is in $C_{i'}$.
- if and only if there is a completion \mathbf{T} of \mathbf{S} with $d(\mathbf{T}[i], \mathbf{T}[i']) = \alpha$ for all $i \neq i' \in [2m+1]$.

Proof. (\Rightarrow) Consider the completion \mathbf{T} of \mathbf{S} in which each missing entry is completed as in \mathbf{C}' . Let a'_i and b'_i be the completion of a_i and b_i for each $i \in [m]$. We have $a'_i \in \{10, 01\}$ and $b'_i \in \{100, 010, 001\}$ due to the first constraint. Now we examine each pairwise row distance.

- For each $i \in [m]$, $d(\mathbf{T}[i], \mathbf{T}[2m+1]) = d(\mathbf{S}[i], \mathbf{S}[2m+1]) + d(a'_i, 00) = (\alpha - 1) + 1 = \alpha$ and $d(\mathbf{T}[m+i], \mathbf{T}[2m+1]) = d(\mathbf{S}[m+i], \mathbf{S}[2m+1]) + d(b'_i, 00) = (\alpha - 1) + 1 = \alpha$.
- For each $i \neq i' \in [m]$, $d(\mathbf{T}[i], \mathbf{T}[i']) = d(\mathbf{S}[i], \mathbf{S}[i']) + d(a_i, 00) + d(a_{i'}, 00) = (\alpha - 2) + 1 + 1 = \alpha$ and $d(\mathbf{T}[m+i], \mathbf{T}[m+i']) = d(\mathbf{S}[m+i], \mathbf{S}[m+i']) + d(b_i, 000) + d(b_{i'}, 000) = (\alpha - 2) + 1 + 1 = \alpha$.
- For each $i, i' \in [m]$ with x_i in $C_{i'}$, $d(\mathbf{T}[i], \mathbf{T}[m+i']) = d(\mathbf{S}[i], \mathbf{S}[m+i']) + (d(\mathbf{C}'[i], \mathbf{C}'[m+i']) - d(\mathbf{C}[i], \mathbf{C}[m+i'])) = (\alpha - 3) + 3 = \alpha$ because of the second constraint.
- For each $i, i' \in [m]$ with x_i not in $C_{i'}$, $d(\mathbf{T}[i], \mathbf{T}[m+i']) = d(\mathbf{S}[i], \mathbf{S}[m+i']) + d(a_i, 00) + d(b'_i, 000) = (\alpha - 2) + 1 + 1 = \alpha$.

Hence, all pairwise row distances are equal to α .

(\Leftarrow) Consider the completion \mathbf{C}' of \mathbf{C} in which each missing entry is completed as in \mathbf{T} . For each $i \in [2m]$, we have $d(\mathbf{C}'[i], \mathbf{C}[2m+1]) - d(\mathbf{C}[i], \mathbf{C}[2m+1]) = d(\mathbf{T}[i], \mathbf{T}[2m+1]) - d(\mathbf{S}[i], \mathbf{S}[2m+1]) = \alpha - (\alpha - 1) = 1$. Moreover, it holds for each $i, i' \in [m]$ with x_i in $C_{i'}$ that $d(\mathbf{C}'[i], \mathbf{C}'[m+i']) - d(\mathbf{C}[i], \mathbf{C}[m+i']) = d(\mathbf{T}[i], \mathbf{T}[m+i']) - d(\mathbf{S}[i], \mathbf{S}[m+i']) = \alpha - (\alpha - 3) = 3$. This concludes the proof of the claim. ◁

Combining Claim 34 and Claim 35, we have that ϕ is a **Yes**-instance if and only if the DMC instance $(\mathbf{S}, \alpha, \alpha)$ is a **Yes**-instance. ◀

To close this section, we conjecture that DMC with $k = 3$ is actually NP-hard for every value of $\beta - \alpha$. Similar reductions from 3-SAT variants might work here. The case $k = 2$ and $\beta - \alpha = 1$ is probably the most promising candidate for being solvable in polynomial time.

5 Conclusion

Together with the recent work of Eiben et al. [7], our work is seemingly among the first in the context of stringology that makes extensive use of Deza's theorem and sunflowers. While Eiben et al. [7] achieved classification results in terms of parameterized (in)tractability, we conducted a detailed complexity analysis in terms of polynomial-time solvable versus

NP-hard cases. Figure 2 provides a visual overview on our results for DIAMETER MATRIX COMPLETION (DMC), also spotting concrete open questions.

Going beyond open questions directly arising from Figure 2, we remark that it is known that the clustering variant of DMC can be solved in polynomial time when the number of clusters is two and the matrix is complete [13]. Hence, a natural question arises whether our tractability results can be extended to this variant as well. Furthermore, we proved that there is a linear-time algorithm solving DMC when $\beta = 2$ and $\alpha = 0$ (Theorem 12). This leads to the question whether linear time is also possible for arbitrary (non-constant) alphabet size. Next, we are curious whether the phenomenon we observed in Theorem 22 concerning the exponential dependence of the running time for $(\alpha, \alpha + 1)$ -DMC when α is even but independence of α when it is odd can be further substantiated or whether one can get rid off the “ α -dependence” in the even case. In terms of standard parameterized complexity analysis, we wonder whether DMC is fixed-parameter tractable with respect to $\beta + k$. Note that $k \in \theta(\ell)$ in our NP-hardness proof for the case $\beta = 4$ (Theorem 23).

Finally, performing a multivariate fine-grained complexity analysis in the same spirit as in recent work for LONGEST COMMON SUBSEQUENCE [3] would be another natural next step.

References

- 1 Bengt Aspvall, Michael F Plass, and Robert Endre Tarjan. A linear-time algorithm for testing the truth of certain quantified Boolean formulas. *Information Processing Letters*, 8(3):121–123, 1979. 16, 19
- 2 Piotr Berman, Marek Karpinski, and Alex D. Scott. Approximation hardness of short symmetric instances of MAX-3SAT. *Electronic Colloquium on Computational Complexity (ECCC)*, (049), 2003. 19
- 3 Karl Bringmann and Marvin Künnemann. Multivariate fine-grained complexity of longest common subsequence. In *29th Annual ACM-SIAM Symposium on Discrete Algorithms, (SODA '18)*, pages 1216–1235, 2018. 27
- 4 Laurent Bulteau, Falk Hüffner, Christian Komusiewicz, and Rolf Niedermeier. Multivariate algorithmics for NP-hard string problems. *Bulletin of the EATCS*, 114, 2014. 3
- 5 Michel Deza. Une propriété extrême des plans projectifs finis dans une classe de codes équi-distants. *Discrete Mathematics*, 6(4):343–352, 1973. 5
- 6 Michel Deza. Solution d'un problème de Erdős-Lovász. *Journal of Combinatorial Theory, Series B*, 16(4):166–167, 1974. 3, 5
- 7 Eduard Eiben, Robert Ganian, Iyad Kanj, Sebastian Ordyniak, and Stefan Szeider. On clustering incomplete data. *CoRR*, abs/1911.01465, 2019. URL: <http://arxiv.org/abs/1911.01465>. 1, 2, 3, 5, 13, 26
- 8 Jörg Flum and Martin Grohe. *Parameterized Complexity Theory*. Texts in Theoretical Computer Science. An EATCS Series. Springer, 2006. 5
- 9 Vincent Froese, René van Bevern, Rolf Niedermeier, and Manuel Sorge. Exploiting hidden structure in selecting dimensions that distinguish vectors. *Journal of Computer and System Sciences*, 82(3):521–535, 2016. 5, 7, 13
- 10 Harold N. Gabow. An efficient reduction technique for degree-constrained subgraph and bidirected network flow problems. In *15th Annual ACM Symposium on Theory of Computing, (STOC '83)*, pages 448–456, 1983. 11, 12
- 11 Robert Ganian, Iyad A. Kanj, Sebastian Ordyniak, and Stefan Szeider. Parameterized algorithms for the matrix completion problem. In *35th International Conference on Machine Learning, (ICML '18)*, pages 1642–1651, 2018. 2
- 12 Jiawei Gao, Russell Impagliazzo, Antonina Kolokolova, and Ryan Williams. Completeness for first-order properties on sparse structures with algorithmic applications. *ACM Transactions on Algorithms*, 15(2):23:1–23:35, 2019. 16
- 13 Leszek Gąsieniec, Jesper Jansson, and Andrzej Lingas. Approximation algorithms for Hamming clustering problems. *Journal of Discrete Algorithms*, 2(2):289–301, 2004. 27
- 14 Danny Hermelin and Liat Rozenberg. Parameterized complexity analysis for the closest string with wildcards problem. *Theoretical Computer Science*, 600:11–18, 2015. 3, 15
- 15 Stasys Jukna. *Extremal Combinatorics: With Applications in Computer Science*. Springer Science & Business Media, 2011. 5
- 16 Tomohiro Koana, Vincent Froese, and Rolf Niedermeier. Parameterized algorithms for matrix completion with radius constraints. *CoRR*, abs/2002.00645, 2020. 2, 3, 7, 8, 11
- 17 Christopher Moore and J. M. Robson. Hard tiling problems with simple tiles. *Discrete & Computational Geometry*, 26(4):573–590, 2001. 23