

Anomalous solutions to nonlinear hyperbolic equations

Michael Oberguggenberger*

Abstract

The behavior of sufficiently regular solutions to semilinear hyperbolic equations has attracted a great deal of attention in the past decades, concerning local/global existence, finite time blow-up, critical exponents, and propagation of singularities. Solutions of lower regularity may exhibit unexpected (anomalous) propagation of singularities. The purpose of this paper is to present various striking examples that seemingly have not been addressed in the literature so far. The key issue is the interpretation of the nonlinear operations.

1 Introduction

This paper serves to display various unusual, or *anomalous* solutions to semilinear wave equations

$$\frac{1}{c^2} \partial_t^2 u - \Delta u = f(x, t, u), \quad u(x, 0) = u_0(x), \quad \partial_t u(x, 0) = u_1(x) \quad (1)$$

in space dimension $n \geq 1$, and to advection-reaction equations

$$\frac{1}{c} \partial_t u + \partial_x u = f(x, t, u), \quad u(x, 0) = u_0(x) \quad (2)$$

in one space dimension as prototypical hyperbolic partial differential equations. The nonlinear function f is assumed to be smooth. Here typically $f(x, t, u) = \pm |u|^p$ or $|u|^{p-1}u$. The main research direction in the past decades has been to find bounds on the exponent p and the regularity of the initial data, asking about the existence of global solutions with small or large initial data, local solutions, self-similar solutions, blow-up in finite time or stability of blow-up. The reader is referred to the discussion in the monograph [13], the survey article from the 1990s [35], a collection of currently known critical exponents [23] and some of the papers discussing the development of the field [16, 36]. Relevant literature on self-similar solutions and stationary solutions as building blocks will be quoted at the appropriate place in Section 4.

In the 1980s and 1990s, a central question has been propagation of singularities, which started with the discovery of Jeffrey Rauch and Michael Reed [28, 29] that in semilinear hyperbolic equations and systems, singularities do not only propagate out from initial singularities along characteristics or bicharacteristics as in the linear case, but may be created at later times by the interaction of previous singularity bearing (bi-)characteristics. For example, an initial singularity at the origin in problem (1) may lead to singularities in the solution that fill up the solid light cone [2]. For a survey of the vast literature up to around 1990 we refer to the monograph [3]. Rauch and Reed coined the term *anomalous singularities* for this phenomenon.

The results on anomalous singularities required sufficient overall regularity of the solution, for example H_{loc}^s -regularity with $s > (n + 1)/2$, and the mechanism for creating the anomalous singularities was still based on characteristics, bicharacteristics and their interaction.

The anomalous solutions presented in this paper are distinguished by (a) lower regularity than in the previous literature and (b) propagation along non-characteristic curves. The majority of examples is based on non-regular solutions to the corresponding stationary elliptic equation. Derivatives are always understood in the sense of distributions. In an attempt to categorize the solutions, four types will be singled out:

- Type I:* products defined by Hörmander's wave front set criterion;
- Type II:* products and powers evaluated by Nemytskii operators;
- Type III:* limits of weak asymptotic solutions;
- Type IV:* sequential solutions, especially very weak solutions in the sense of Ruzhansky.

*Unit of Engineering Mathematics, University of Innsbruck, Technikerstraße 13, 6020 Innsbruck, Austria, (michael.oberguggenberger@uibk.ac.at)

It is worth noting that all constructed solutions come with a certain assertion of uniqueness.

The plan of the paper is as follows. Section 2 serves to recall results on anomalous propagation of singularities for sufficiently regular solutions, for reasons of comparison. Section 3 addresses Type I solutions, introducing the employed multiplication of distributions and discussing the question of regularization. Section 4 will exhibit seemingly harmless solutions lying in an L^p -space on which the nonlinear operations are defined and continuous (Type II). In Section 5 it will be shown that the solutions from Section 4 arise as limits of nets of asymptotic solutions (satisfying the equations up to an error term converging weakly to zero, Type III). In Section 6 nets of smooth functions $(u_\varepsilon)_{\varepsilon>0}$ will be constructed that solve the equations at each fixed $\varepsilon > 0$, but need not necessarily converge as $\varepsilon \rightarrow 0$ (Type IV). Nevertheless, their regularity properties can be characterized by suitable estimates on their growth in terms of negative powers of ε as $\varepsilon \rightarrow 0$. The appendix serves to recall some notions required to define the products arising in Type I solutions.

The author has been aware of the existence of these anomalous solutions since the early 1980s, but due to a lack of explanation, hesitated to publish them so far. It is hoped that this publication will arouse interest in these types of solutions among the community. Many more examples of similar nature are known, collected by the author and in joint work with Hideo Deguchi [9].

What concerns notation, H^s denotes the usual Sobolev space based on L^2 ; \mathcal{C}^k denotes the space of k -times differentiable functions, \mathcal{C}_b^k the subspace of functions with bounded derivatives up to order k . The notation for spaces of test functions and distributions follows [34]. The Fourier transform is defined as $\mathcal{F}\varphi(\xi) = \int e^{-2\pi i x \xi} \varphi(x) dx$.

2 Propagation of singularities for regular solutions

This section serves to recall results from the 1980s on propagation of singularities for solutions to semilinear hyperbolic systems. These results hold for sufficiently regular solutions (L_{loc}^∞ in one space dimension, H_{loc}^s for $s > (n+1)/2$ in space dimension n). We do not strive for full generality – the quoted results will be contrasted with the much less regular solutions to be constructed in the following sections.

We start with $(m \times m)$ -systems of first order hyperbolic equations in one space dimension, considering the initial value problem

$$\begin{aligned} (\partial_t + \Lambda \partial_x)u(x, t) &= f(x, t, u(x, t)), \quad (x, t) \in R \\ u(x, 0) &= u_0(x), \quad x \in R_0 \end{aligned} \tag{3}$$

where $R_0 \subset \mathbb{R}$ is an interval and $R \subset \mathbb{R} \times [0, \infty)$ is its domain of determinacy. Here $u = (u_1, \dots, u_m)$, $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_m)$ with real and constant entries λ_i , and $f = (f_1, \dots, f_m)$ is smooth. Let $x_1, \dots, x_k \in R_0$ and denote by S_0 the union of characteristic lines emanating from x_1, \dots, x_k . Following [29], construct the forward characteristic lines starting at the intersection points of S_0 and call this set S_1 . Let S_2 be the set of forward characteristic lines starting from the intersection points of S_1 . Continue recursively to construct a sequence of sets S_j . Let S be the closure of $\bigcup_{j=0}^\infty S_j$ intersected with R .

Proposition 2.1. *Let $u \in (L^\infty(R))^m$ satisfy (3) in the sense of distributions and take on the initial data $u_0 \in (L^\infty(R_0))^m$. Suppose that u_0 is \mathcal{C}^∞ with each derivative uniformly bounded on the complement of the finitely many points x_1, \dots, x_k . Then u is \mathcal{C}^∞ on $R \setminus S$ and all derivatives of u have continuous extensions from each connected component of $R \setminus S$ to its closure.*

Proof. This is Theorem 1 from [29]. □

Remark 2.2. (a) *If the function f is linear, then the solution u is in \mathcal{C}^∞ on $R \setminus S_0$ – singularities can only lie on characteristic curves tracing back to the singularities of the initial data. In the nonlinear case, the solution is not \mathcal{C}^∞ on $S \setminus S_0$, in general. The singularities belonging to $S \setminus S_0$ in the nonlinear case have been termed anomalous singularities by the authors.*

(b) *In the scalar case and in the case of (2×2) -systems (thus $m = 1$ or $m = 2$), $S = S_0$, so no anomalous singularities arise.*

Next we recall a result of [27] on propagation of singularities for semilinear wave equations. Consider the initial value problem

$$\begin{aligned} (\partial_t^2 - \Delta)v(x, t) &= f(v(x, t)), \quad (x, t) \in \mathbb{R}^n \times \mathbb{R}, \\ v(x, 0) &= v_0(x), \quad \partial_t v(x, 0) = v_1(x), \quad x \in \mathbb{R}^n, \end{aligned} \tag{4}$$

where f is a polynomial with $f(0) = 0$, Δ denotes the n -dimensional Laplace operator, and $u_0 \in H_{loc}^s(\mathbb{R}^n)$, $u_1 \in H_{loc}^{s-1}(\mathbb{R}^n)$ with $s > (n+1)/2$. Note that $H_{loc}^s(\mathbb{R}^n \times \mathbb{R})$ is an algebra in this case, even contained in the space of continuous functions, so $f(u)$ is classically defined.

Proposition 2.3. *Let $s > (n+1)/2$ and $v \in H_{\text{loc}}^s(\mathbb{R}^n \times \mathbb{R})$ satisfy (4) in the sense of distributions. Suppose that v_0 and v_1 belong to $C^\infty(\mathbb{R}^n \setminus \{0\})$. Then v is C^∞ on $\{|x| > |t|\}$, and it belongs to $H_{\text{loc}}^{s+1+\sigma}(\mathbb{R}^n \times \mathbb{R})$ on $\{|x| < |t|\}$ for all $\sigma < s - (n+1)/2$.*

Proof. This follows from Theorem 3.1, together with Theorem 1.1 of [27]. \square

Remark 2.4. *In space dimension $n = 1$, the solution v is actually C^∞ in $\{|x| < |t|\}$, as follows from the Corollary to Theorem 2 in [28] as well as the earlier paper [30].*

It is known that the solution is not necessarily better than $H^{s+1+\sigma}$ in $\{|x| < |t|\}$ in space dimension $n \geq 2$. For a survey of the state of the art around 1990, see [3].

3 Type I solutions – multiplication of distributions

In this section, we address weak solutions to nonlinear equations where the involved products or powers exist in the sense of Hörmander’s wave front set criterion [18]. The examples will be based on the one-dimensional distribution

$$u_0(x) = \frac{1}{x + i0} = \lim_{\varepsilon \rightarrow 0} \frac{1}{x + i\varepsilon} = \text{vp} \frac{1}{x} - i\pi\delta(x) \quad (5)$$

also denoted by $\delta_+(x)$ in the physics literature. Here $\text{vp} \frac{1}{x}$ denotes the principal value distribution $\text{vp} \frac{1}{x} = \partial_x \log|x|$ and $\delta(x)$ is the Dirac measure. The Fourier transform of $u_0(x)$ and its auto-convolution are

$$(\mathcal{F}u_0)(\xi) = -2\pi i H(\xi) \quad \text{and} \quad (\mathcal{F}u_0 * \mathcal{F}u_0)(\xi) = -4\pi^2 \xi H(\xi)$$

where H denotes the Heaviside function. In particular, the wavefront set of u_0 is $\{(0, \xi) : \xi > 0\}$, thus u_0^2 exists according to Hörmander’s criterion. Actually, it can simply be computed as Fourier product (see Appendix),

$$u_0^2 = \mathcal{F}^{-1}(\mathcal{F}u_0 * \mathcal{F}u_0),$$

as well as all its powers. It holds that

$$u_0^2(x) = \left(\frac{1}{x + i0}\right)^2 = -\left(\frac{1}{x + i0}\right)' = \text{Pf} \frac{1}{x^2} + i\pi\delta'(x) = -u_0'(x) \quad (6)$$

where $\text{Pf} \frac{1}{x^2}$ is the Hadamard finite part distribution, and

$$2u_0^3(x) = 2\left(\frac{1}{x + i0}\right)^3 = \left(\frac{1}{x + i0}\right)'' = u_0''(x). \quad (7)$$

3.1 A nonlinear advection-reaction equation

Proposition 3.1. *The distribution $u(x, t) \equiv u_0(x)$ given by (5) is a weak solution to the initial value problem*

$$\frac{1}{c} \partial_t u + \partial_x u + u^2 = 0, \quad u(x, 0) = u_0(x) \quad (8)$$

for whatever $c \in \mathbb{R}, c \neq 0$, where the square is understood in the sense of Hörmander’s product.

Proof. It is clear from (6) that $\partial_x u + u^2 = 0$ and that $\partial_t u = 0$. \square

Clearly, the mechanism producing this result is that the stationary solution satisfies the nonlinear differential relation $u_0' = -u_0^2$. Further reasons why a genuine distribution can satisfy such a relation will be discussed below. At first we wish to point out that the solution given in Proposition 3.1 exhibits anomalous propagation of singularities. Indeed,

$$\text{singsupp } u = \{(x, t) : x = 0, t \geq 0\}$$

while the expected singular support from Proposition 2.1 or Remark 2.2(b) should be $\{(x, t) : x = ct\}$. To be sure, u_0 does not belong to L^∞ as required in Proposition 2.1.

Remark 3.2. *It should be noted that anomalous propagation of singularities is not confined to stationary solutions. The following example, due to Deguchi [9], shows that any anomalous propagation speed is possible. Indeed,*

$$u(x, t) = \frac{1}{ax + bct + i0} \quad \text{with } a + b = 1$$

solves equation (8) with initial data $u_0(x) = (ax + i0)^{-1}$.

Remark 3.3. One possible explanation why the mentioned nonlinear differential relation, as well as similar relations for the higher derivatives, hold for the specific distribution (5) can be obtained by studying its representation as a boundary value of an analytic function. Indeed, every distribution $v \in \mathcal{D}'(\mathbb{R})$ can be represented as the boundary value of a function $\widehat{v}(z)$, analytic in $\mathbb{C} \setminus \text{supp}(u)$, in the sense

$$v(x) = \lim_{\varepsilon \rightarrow 0} (\widehat{v}(x + i\varepsilon) - \widehat{v}(x - i\varepsilon)), \quad (9)$$

see e.g. [37]. If v is a distribution of compact support, $\widehat{v}(z)$ is given by the Fantappiè indicatrix

$$\widehat{v}(z) = \frac{1}{2\pi i} \langle v(x), \frac{1}{x-z} \rangle$$

and in general by a partition of unity procedure. Further, $|\widehat{v}(z)|$ grows at most like a negative power of $|\text{Im } z|$ as $\text{Im } z \rightarrow 0$, locally uniformly in $\text{Re } z$. The representation $\widehat{v}(z)$ is unique up to a function analytic on \mathbb{C} . Further, every function $\widehat{v}(z)$, analytic in $\mathbb{C} \setminus \mathbb{R}$ and satisfying the growth condition has a distributional boundary value in the sense of (9).

If the support of $\widehat{v}(z)$ is contained in $\{\text{Im } z > 0\}$, the representation is unique. Thus the space of distributions $\mathcal{H}_+(\mathbb{R})$ whose Fantappiè parametrix has support in the upper complex half plane is isomorphic to the space of analytic functions in the upper complex half plane satisfying the mentioned growth condition. However, the latter space is a differential algebra, the differential-algebraic structure of which can be transported to $\mathcal{H}_+(\mathbb{R})$, rendering it a differential algebra [37].

This is exactly the case with $u_0(x)$ given by (5) for which

$$\widehat{u}_0(z) = \begin{cases} \frac{1}{z}, & \text{Im } z > 0, \\ 0, & \text{Im } z < 0. \end{cases}$$

In the algebra of analytic functions in the upper half plane, the functional relation

$$\frac{d^k}{dz^k} \left(\frac{1}{z} \right) = (-1)^k k! \left(\frac{1}{z} \right)^{k+1}, \quad z \neq 0$$

obtains. In this way, formulas (6) and (7) are explained. The differential-algebraic relations persist in the boundary values.

3.1.1 Analytic regularization

It will be instructive to study the behavior of approximate solutions when the initial data are regularized. The first obvious possibility is to consider the analytic regularization defining the distribution $u_0(x) = 1/(1+i0)$. We wish to solve the regularized problem

$$\frac{1}{c} \partial_t u_\varepsilon + \partial_x u_\varepsilon + u_\varepsilon^2 = 0, \quad u_\varepsilon(x, 0) = u_{0\varepsilon}(x) = \frac{1}{x + i\varepsilon}. \quad (10)$$

Solving (10) by the method of characteristics results in the unique classical solution

$$u_\varepsilon(x, t) = \frac{u_{0\varepsilon}(x - ct)}{1 + ct u_{0\varepsilon}(x - ct)} = \frac{\frac{1}{x - ct + i\varepsilon}}{1 + ct \frac{1}{x - ct + i\varepsilon}} = \frac{1}{x + i\varepsilon}.$$

Thus, by simple arithmetic, $u_\varepsilon(x, t) \equiv u_{0\varepsilon}(x)$ and so the solution given in Proposition 3.1 coincides with the weak limit of approximate solutions when the initial data are replaced by their analytic regularization.

3.1.2 Regularization by convolution with a mollifier

The purpose of this subsection is to show that the convergence of the approximate solution is a peculiarity of the analytic regularization and does not hold if the initial data are regularized by convolution with a standard Friedrichs mollifier $\varphi_\varepsilon(x) = \varepsilon^{-1} \varphi(x/\varepsilon)$ with $\varphi \in \mathcal{D}(\mathbb{R})$, $\int \varphi(x) dx = 1$. For the sake of the argument, we take $\varphi \geq 0$ symmetric, $\text{supp } \varphi \subset (-1, 1)$. Thus let

$$U_{0\varepsilon}(x) = (u_0 * \varphi_\varepsilon)(x)$$

and let $U_\varepsilon(x, t)$ be the corresponding classical solution to (10) with initial data $U_{0\varepsilon}$. By the method of characteristics,

$$U_\varepsilon(x, t) = \frac{(\text{vp} \frac{1}{x} * \varphi_\varepsilon)(x - ct) - i\pi \varphi_\varepsilon(x - ct)}{1 + ct((\text{vp} \frac{1}{x} * \varphi_\varepsilon)(x - ct) - i\pi \varphi_\varepsilon(x - ct))}.$$

In particular,

$$U_\varepsilon(ct - \varepsilon, t) = \frac{(\text{vp}_{\frac{1}{x}} * \varphi_\varepsilon)(-\varepsilon)}{1 + ct(\text{vp}_{\frac{1}{x}} * \varphi_\varepsilon)(-\varepsilon)} \quad (11)$$

We show that the solution $U_\varepsilon(x, t)$ blows up at latest at

$$t_\varepsilon = \frac{-1/c}{(\text{vp}_{\frac{1}{x}} * \varphi_\varepsilon)(-\varepsilon)} = \frac{1/c}{(\text{vp}_{\frac{1}{x}} * \varphi_\varepsilon)(\varepsilon)}$$

and that this number is of order ε as $\varepsilon \rightarrow 0$. Thus there is no global solution, when Friedrichs regularization is used.

Indeed, starting from the defining formula

$$(\text{vp}_{\frac{1}{x}} * \varphi_\varepsilon)(x) = \lim_{\eta \rightarrow 0} \int_{|x-y| \geq \eta} \frac{\varphi_\varepsilon(y)}{x-y} dy,$$

some simple manipulations using the support properties of φ lead to

$$(\text{vp}_{\frac{1}{x}} * \varphi_\varepsilon)(-\varepsilon) = \lim_{\eta \rightarrow 0} \int_{-1+\eta/\varepsilon}^{\infty} \frac{\varphi(y)}{-\varepsilon(1+y)} dy = -\frac{1}{\varepsilon} \int_{\text{supp } \varphi} \frac{\varphi(y)}{1+y} dy = -\frac{1}{\varepsilon} C_\varphi$$

where C_φ is a positive constant. This shows that the denominator in (11) is indeed zero at $t_\varepsilon = \varepsilon/cC_\varphi$, while the numerator is nonzero.

3.1.3 Separation in real and imaginary part

One might argue that the complex valued initial value problem (10) is actually a real valued, nonstrictly hyperbolic system. This is indeed the case; the real and imaginary part of the analytically regularized solution are

$$u_\varepsilon(x, t) = \frac{1}{x + i\varepsilon} = v_\varepsilon(x, t) + iw_\varepsilon(x, t) = \frac{x}{x^2 + \varepsilon^2} - i \frac{\varepsilon}{x^2 + \varepsilon^2}.$$

The hyperbolic system for the real and imaginary part is

$$\begin{aligned} \partial_t v_\varepsilon + \partial_x v_\varepsilon &= -v_\varepsilon^2 + w_\varepsilon^2, \\ \partial_t w_\varepsilon + \partial_x w_\varepsilon &= -2v_\varepsilon w_\varepsilon. \end{aligned}$$

Here $v_\varepsilon(x, t) \rightarrow \text{vp}_{\frac{1}{x}}$ and $w_\varepsilon(x, t) \rightarrow -\pi\delta(x)$ as $\varepsilon \rightarrow 0$. However, it is well-known (and rather immediate) that v_ε^2 and w_ε^2 do not converge in $\mathcal{D}'(\mathbb{R})$ as $\varepsilon \rightarrow 0$. Thus the individual terms in the first line make no sense in the limit. (By purely arithmetic manipulations involving $1/(x + i\varepsilon)$ and $1/(x - i\varepsilon)$ and their limits, the limit in the right-hand side of the second line is seen to exist and to equal $-\pi\delta'(x)$.)

3.2 A nonlinear wave equation

In the same vein, the distribution $u_0(x)$ can serve to produce a solution to a semilinear wave equation in one space dimension.

Proposition 3.4. *The distribution $u(x, t) = u_0(x)$ given by (5) is a weak solution to the initial value problem*

$$\frac{1}{c^2} \partial_t^2 u - \partial_x^2 u + 2u^3 = 0, \quad u(x, 0) = u_0(x), \quad \partial_t u(x, 0) = 0 \quad (12)$$

for whatever $c > 0$, where the cubic term is understood in the sense of Hörmander's product.

Proof. It is clear from (7) that $-\partial_x^2 u + 2u^3 = 0$ and that $\partial_t u = 0$. □

In the real-valued case, the wave equation (12) has a so-called defocusing nonlinearity. For initial data (u_0, u_1) in $H^1(\mathbb{R}) \times L^2(\mathbb{R})$, it would have a unique global finite energy solution [35], belonging to $\mathcal{C}([0, \infty) : H^1(\mathbb{R})) \cap \mathcal{C}^1([0, \infty) : L^2(\mathbb{R}))$. By local existence theory, it could also be extended to small negative times, and hence would belong to L_{loc}^∞ in an open neighborhood of the half plane. As in Remark 2.4, the Corollary to Theorem 2 in [28] would imply that a singularity in the initial data at $x = 0$ can only spread along the characteristic lines $x = \pm ct$. Clearly, the solution given in Proposition 12 neither has the required regularity properties nor does it show the expected singularity propagation.

Remark 3.5. *The distribution $u_0(x)$ is homogeneous of degree -1 . Thus $u(x, t) = u_0(x)$ is a self-similar solution to (12), satisfying $\mu u(\mu x, \mu t) = u(x, t)$ for all $\mu > 0$.*

4 Type II solutions – Nemytskii operators

This section addresses weak solutions, whereby the nonlinear terms are defined by Nemytskii operators. We recall the pseudofunctions R_λ , meromorphic functions of $\lambda \in \mathbb{C}$ with values in the space of tempered distributions $\mathcal{S}'(\mathbb{R}^n)$ [11, Chapter 17]. For $\operatorname{Re} \lambda > -n$ they are given by

$$\langle R_\lambda, \varphi \rangle = \int |x|^\lambda \varphi(x) dx$$

and can be analytically continued to $\mathbb{C} \setminus \{-n - 2k : k \in \mathbb{N}\}$. Outside the poles, they satisfy

$$\Delta R_\lambda = \lambda(\lambda + n - 2)R_{\lambda-2}$$

In particular, when $\lambda > 2 - n$ and $p = 1 - 2/\lambda$, R_λ belongs to $L_{\text{loc}}^p(\mathbb{R}^n)$, $(R_\lambda)^p = R_{\lambda p}$ and it satisfies the elliptic equation

$$\Delta R_\lambda = \lambda(\lambda + n - 2)(R_\lambda)^p,$$

where the derivatives are understood in the weak sense and the p th power as the evaluation of the Nemytskii operator $L_{\text{loc}}^p(\mathbb{R}^n) \rightarrow L_{\text{loc}}^1(\mathbb{R}^n)$.

We note that for $\lambda \in \mathbb{R} \setminus \{-n - 2k : k \in \mathbb{N}\}$, R_λ is homogeneous of degree λ , and $R_\lambda \in H_{\text{loc}}^1(\mathbb{R}^n)$, if $\lambda > (2 - n)/2$.

As examples to be discussed further, we only consider two cases in which p is a positive integer. In the context of propagation of singularities, fractional powers are not interesting for our purpose, because they represent non-smooth nonlinearities. We use the solutions R_λ as examples of peculiar rotationally symmetric stationary solutions to nonlinear wave equations.

Example 4.1. Let $n = 3$ and $\lambda = -1/2$ (then $\lambda(\lambda + n - 2) = -1/4$). Let $u_0(x) = |x|^{-1/2}$. Then $u_0 \in L_{\text{loc}}^5(\mathbb{R}^3)$, and $u(x, t) \equiv u_0(x)$ satisfies the nonlinear wave equation

$$\frac{1}{c^2} \partial_t^2 u - \Delta u - \frac{1}{4} u^5 = 0, \quad u(x, 0) = u_0(x), \quad \partial_t u(x, 0) = 0 \quad (13)$$

for whatever $c > 0$.

Example 4.2. Let $n = 4$ and $\lambda = -1$ (then $\lambda(\lambda + n - 2) = -1$). Let $u_0(x) = |x|^{-1}$. Then $u_0 \in L_{\text{loc}}^3(\mathbb{R}^4)$, and $u(x, t) \equiv u_0(x)$ satisfies the nonlinear wave equation

$$\frac{1}{c^2} \partial_t^2 u - \Delta u - u^3 = 0, \quad u(x, 0) = u_0(x), \quad \partial_t u(x, 0) = 0 \quad (14)$$

for whatever $c > 0$.

In all these cases, derivatives are understood in the weak sense and the powers of u exist as locally integrable functions, actually as evaluations of the continuous map $u \rightarrow u^p$ from $L_{\text{loc}}^p \rightarrow L_{\text{loc}}^1$. Note that the nonlinear operation is taken outside the space of distributions, and the result is embedded afterwards.

Remark 4.3. (a) As u_0 is nonnegative, we might replace u^5 by $|u|^5$ or $|u|^4 u$. In any case, we are dealing with so-called focusing nonlinearities.

(b) Recall that $u(x, t)$ is a self-similar solution to the nonlinear wave equation

$$\frac{1}{c^2} \partial_t^2 u - \Delta u \pm |u|^p = 0, \quad (15)$$

if $u(x, t) = \mu^\alpha u(\mu t, \mu x)$ for all $\mu > 0$, where necessarily $\alpha = 2/(p - 1)$. On the other hand, $u_0 = R_\lambda$ is homogeneous of degree λ , that is, $u_0(sx) = s^\lambda u_0(x)$. It also satisfies equation (15) when $\lambda - 2 = \lambda p$, i.e., $\lambda = -2/(p - 1)$. Thus the special solutions exhibited here are self-similar solutions to the nonlinear wave equation. However, they do not fall into the classes of functions considered e.g. in [4, 19, 25, 26, 31]. It should be noted that solutions to nonlinear elliptic equations have also been used in the literature. They can serve for constructing solutions of finite life span, but also for proving the existence of (time-dependent) self-similar solutions [7, 12, 20, 21].

5 Type III – weak asymptotic solutions

A net of smooth functions $(u_\varepsilon)_{\varepsilon>0}$ is called a *weak asymptotic solution* [8] to a nonlinear partial differential equation, such as equation (15), if it has a limit in the space of distributions and if it satisfies the equation up to an error term which tends to zero weakly as $\varepsilon \rightarrow 0$.

The basic example derives again from a nonlinear elliptic equation. Indeed, in \mathbb{R}^n , we start from the relation

$$\Delta(|x|^2 + \varepsilon^2)^q = ((2qn + 4q(q-1))|x|^2 + 2qn\varepsilon^2)(|x|^2 + \varepsilon^2)^{q-2}.$$

We will simply work out two special cases that correspond to the ones in Examples 4.1 and 4.2.

Example 5.1. Let $n = 3$ and $q = -1/4$. By simple arithmetic,

$$(2qn + 4q(q-1))|x|^2 + 2qn\varepsilon^2 = -\frac{1}{4}(|x|^2 - \varepsilon^2) - \frac{5}{4}\varepsilon^2$$

and so

$$\Delta(|x|^2 + \varepsilon^2)^{-1/4} = -\frac{1}{4}(|x|^2 + \varepsilon^2)^{-5/4} - \frac{5}{4}\varepsilon^2(|x|^2 + \varepsilon^2)^{-9/4}.$$

Thus

$$u_\varepsilon(x, t) = (|x|^2 + \varepsilon^2)^{-1/4}$$

satisfies the nonlinear wave equation

$$\frac{1}{c^2}\partial_t^2 u_\varepsilon - \Delta u_\varepsilon - \frac{1}{4}u_\varepsilon^5 - \frac{5}{4}\varepsilon^2 u_\varepsilon^9 = 0 \quad (16)$$

for whatever $c > 0$. An easy calculation shows that $\varepsilon^2 u_\varepsilon^9$ converges to zero in $\mathcal{D}'(\mathbb{R}^3)$ as $\varepsilon \rightarrow 0$. Thus u_ε is a weak asymptotic solution to the nonlinear wave equation (13) with initial data converging to $u_0(x) = |x|^{-1/2}$. As in Example 4.1 we set $u(x, t) = u_0(x)$. By the continuity assertions for Type II solutions,

$$u_\varepsilon \rightarrow u, \quad u_\varepsilon^5 \rightarrow u^5 \quad \text{in } L_{\text{loc}}^1(\mathbb{R}^3) \quad \text{as } \varepsilon \rightarrow 0,$$

thus each term in equation (16) converges to the corresponding term in equation (13). Further, u_ε is a smooth approximation to u ; as $\varepsilon \rightarrow 0$, a singularity emerges at $x = 0$.

It is of interest to note that the solution to the regularized equation (16) is unique. This emphasizes again the anomaly in the propagation of singularities in the initial value problem (13).

Lemma 5.2. Let $n = 1, n = 2$ or $n = 3$. Assume that $u_0 \in C_b^1(\mathbb{R}^n)$, $u_1 \in C_b^0(\mathbb{R}^n)$ and let f be smooth. Given any $T > 0$, the initial value problem

$$\frac{1}{c^2}\partial_t^2 u - \Delta u = f(u), \quad u(x, 0) = u_0(x), \quad \partial_t u(x, 0) = u_1(x) \quad (17)$$

has at most one weak solution in $C_b^0(\mathbb{R}^n \times [0, T])$.

Proof. Let $S(t)$ be the fundamental solution of the Cauchy problem, that is, $S(t)$ is the inverse Fourier transform of $\sin(c|\xi|t)/|\xi|$. In space dimensions $n = 1, 2, 3$, $S(t)$ is a finite measure of total mass ct . The solution is given by

$$u(\cdot, t) = \frac{d}{dt}S(t) * u_0 + S(t) * u_1 + \int_0^t S(t-s) * f(u(\cdot, s)) ds.$$

By Young's inequality, the L^∞ -estimate

$$\|u(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} \leq C(t)\|u_0, \nabla u_0, u_1\|_{L^\infty(\mathbb{R}^n)} + \int_0^t (ct - cs)\|u(\cdot, s)\|_{L^\infty(\mathbb{R}^n)} ds$$

holds, where $C(t)$ is a constant depending linearly on t . Applying this estimate to the difference $u - v$ of two solutions with the same initial data, writing $f(u) - f(v) = (u - v)g(u, v)$ with g smooth and applying Gronwall's inequality shows that $u = v$. \square

Example 5.3. Let $n = 4$ and $q = -1/2$ and let

$$u_\varepsilon(x, t) = (|x|^2 + \varepsilon^2)^{-1/2}.$$

By the same arguments as in Example 5.1 one sees that u_ε satisfies the nonlinear wave equation

$$\frac{1}{c^2} \partial_t^2 u_\varepsilon - \Delta u_\varepsilon - u_\varepsilon^3 - 3\varepsilon^2 u_\varepsilon^5 = 0 \quad (18)$$

for whatever $c > 0$. Again, one shows that $\varepsilon^2 u_\varepsilon^5$ converges to zero in $\mathcal{D}'(\mathbb{R}^4)$ as $\varepsilon \rightarrow 0$, and u_ε is a weak asymptotic solution to the nonlinear wave equation (14) with initial data converging to $u_0(x) = |x|^{-1}$. With $u(x, t) \equiv u_0(x)$, one has again

$$u_\varepsilon \rightarrow u, \quad u_\varepsilon^3 \rightarrow u^3 \quad \text{in } L^1_{\text{loc}}(\mathbb{R}^4) \quad \text{as } \varepsilon \rightarrow 0,$$

thus each term in equation (18) converges to the corresponding term in equation (14). The same behavior as in Example 5.1 is observed.

Due to the continuity of the Nemytskii operators, the weak asymptotic solutions constructed here are consistent with the solutions presented in Section 4.

6 Type IV – sequential solutions

In this section, we address solutions defined by nets of smooth functions which do not necessarily converge. To introduce the concept, let Ω be an open subset of \mathbb{R}^n and let P be a possibly nonlinear partial differential operator which is a smooth function of its arguments, $Pu = P(x, u, \partial u, \dots)$. Let $(u_\varepsilon)_{\varepsilon > 0}$ a net of functions belonging to $\mathcal{C}^\infty(\Omega)$. If $Pu_\varepsilon = 0$ for all sufficiently small $\varepsilon > 0$, then the net $(u_\varepsilon)_{\varepsilon > 0}$ is called a *sequential solution* of the equation $Pu = 0$, following e.g. [32]. The net $(u_\varepsilon)_{\varepsilon > 0}$ may or may not converge. Even if $(u_\varepsilon)_{\varepsilon > 0}$ converges, individual terms in $P(x, u, \partial u, \dots)$ may or may not converge. However, if $(u_\varepsilon)_{\varepsilon > 0}$ converges to a distribution u , together with all individual terms in $P(x, u, \partial u, \dots)$, then u can be called a *proper weak solution* to $Pu = 0$ [22].

Restricting the class of sequential solutions to *moderate nets* allows one to establish a regularity theory for sequential solutions, even if they diverge. A net of smooth functions $(u_\varepsilon)_{\varepsilon > 0}$ on Ω is called *moderate*, if for all compact subsets K of Ω and all multi-indices $\alpha \in \mathbb{N}_0^n$ there exists $b \geq 0$ such that

$$\sup_{x \in K} |\partial^\alpha u_\varepsilon(x)| = O(\varepsilon^{-b}) \quad \text{as } \varepsilon \rightarrow 0.$$

The net of smooth functions $(u_\varepsilon)_{\varepsilon > 0}$ on Ω is called *negligible*, if for all compact subsets K of Ω , all multi-indices $\alpha \in \mathbb{N}_0^n$ and all $a \geq 0$,

$$\sup_{x \in K} |\partial^\alpha u_\varepsilon(x)| = O(\varepsilon^a) \quad \text{as } \varepsilon \rightarrow 0.$$

Following [15, 33], a moderate net satisfying $Pu_\varepsilon = 0$ for all sufficiently small $\varepsilon > 0$ is called a *very weak solution* to the equation $Pu = 0$. If $(u_\varepsilon)_{\varepsilon > 0}$ is moderate and $Pu_\varepsilon = n_\varepsilon$ where $(n_\varepsilon)_{\varepsilon > 0}$ is a negligible net, then $(u_\varepsilon)_{\varepsilon > 0}$ is a *Colombeau solution* to the equation $Pu = 0$. (As a matter of fact, its equivalence class in the Colombeau algebra $\mathcal{G}(\Omega)$ is a solution in the differential-algebraic sense [6, 17, 24].)

Finally, a net $(u_\varepsilon)_{\varepsilon > 0}$ is said to possess the \mathcal{G}^∞ -property, if for all compact subsets K of Ω there is $b \geq 0$ such that for all multi-indices $\alpha \in \mathbb{N}_0^n$,

$$\sup_{x \in K} |\partial^\alpha u_\varepsilon(x)| = O(\varepsilon^{-b}) \quad \text{as } \varepsilon \rightarrow 0.$$

(Note the change in quantifiers: the local order of growth is the same for all derivatives.) The significance of this notion is that it generalizes \mathcal{C}^∞ -smoothness from distributions to moderate nets. In fact, if $w \in \mathcal{E}'(\Omega)$ is a compactly supported distribution and φ_ε is a mollifier ($\varphi_\varepsilon(x) = \varepsilon^{-n} \varphi(x/\varepsilon)$ with φ smooth, rapidly decaying and $\int \varphi(x) dx = 1$), then

- $w_\varepsilon = w * \varphi_\varepsilon|_\Omega$ defines a moderate net;
- $(w_\varepsilon)_{\varepsilon > 0}$ has the \mathcal{G}^∞ -property if and only if $w \in \mathcal{C}^\infty(\Omega)$.

The \mathcal{G}^∞ -singular support of a moderate net $(u_\varepsilon)_{\varepsilon > 0}$ is defined as the complement of the largest open subset $\omega \subset \Omega$ such that $(u_\varepsilon|_\omega)_{\varepsilon > 0}$ has the \mathcal{G}^∞ -property on ω . With this notion, \mathcal{C}^∞ -regularity theory and propagation of singularities for linear partial differential equations can be literally transferred to the setting of moderate nets in the case of linear equations (with possibly non-smooth coefficients) see e.g. [10, 14, 24].

The same notions can be introduced for nets of smooth functions defined on the closure of an open subset of \mathbb{R}^n , thereby enabling the study of initial value problems or boundary value problems.

6.1 Moderate sequential solutions to an advection-reaction equation

We are going to construct moderate sequential solutions to the advection-reaction equation in one space dimension

$$\frac{1}{c}\partial_t u + \partial_x u + \frac{2}{p} x u^{p+1} = 0, \quad u(x, 0) = u_0(x) \quad (19)$$

where – for simplicity – p is a positive integer. We first note that for continuous initial data, there is at most one solution.

Lemma 6.1. *Assume that $u_0 \in C_b^0(\mathbb{R})$, $c \neq 0$ and let f be smooth. Given any $T > 0$, the initial value problem*

$$\frac{1}{c}\partial_t u + \partial_x u = f(x, t, u), \quad u(x, 0) = u_0(x) \quad (20)$$

has at most one weak solution in $C_b^0(\mathbb{R}^n \times [0, T])$.

Proof. Indeed, if u is a solution, it solves the integral equation

$$u(x, t) = u_0(x - ct) + \int_0^t f(x - ct + cs, s, u(x - ct + cs, s)) ds.$$

Uniqueness follows by the same argument as in the proof of Lemma 5.2. \square

It is immediately checked that, for each $\varepsilon > 0$, the smooth function

$$u_\varepsilon(x, t) \equiv u_{0\varepsilon}(x) = (x^2 + \varepsilon^2)^{-1/p} \quad (21)$$

is a solution to the initial value problem

$$\frac{1}{c}\partial_t u_\varepsilon + \partial_x u_\varepsilon + \frac{2}{p} x u_\varepsilon^{p+1} = 0, \quad u_\varepsilon(x, 0) = (x^2 + \varepsilon^2)^{-1/p}. \quad (22)$$

According to Lemma 6.1, the solution is unique. It is clear that the net $(u_\varepsilon)_{\varepsilon>0}$ is moderate, hence it defines a moderate sequential solution to (19).

Lemma 6.2. *The net $(u_{0\varepsilon})_{\varepsilon>0}$ converges for $p \geq 3$ and diverges for $p = 1, 2$. In particular, $(u_{0\varepsilon}^{p+1})_{\varepsilon>0}$ diverges for every p .*

Proof. For $p \geq 3$, $u_0(x) = |x|^{-2/p}$ belongs to the space of locally integrable functions, and $u_{0\varepsilon}(x) = (x^2 + \varepsilon^2)^{-1/p}$ converges to it in that space.

Let $p = 2$ and take a test function $\varphi \geq 0$ such that $\varphi(x) = 1$ on $[-1, 1]$. Then

$$\langle u_{0\varepsilon}, \varphi \rangle = \int_{-\infty}^{\infty} \frac{\varphi(x)}{\sqrt{x^2 + \varepsilon^2}} dx \geq \int_{-1}^1 \frac{1}{\sqrt{x^2 + \varepsilon^2}} dx = \int_{-1/\varepsilon}^{1/\varepsilon} \frac{1}{\sqrt{y^2 + 1}} dy \rightarrow \infty$$

as $\varepsilon \rightarrow 0$. A similar argument shows that $(x^2 + \varepsilon^2)^{-q}$ diverges for $q > 1/2$. Thus $u_{0\varepsilon}(x) = (x^2 + \varepsilon^2)^{-1/p}$ diverges when $p < 2$ as well, in particular, for $p = 1$. Further, $u_{0\varepsilon}^{p+1}(x) = (x^2 + \varepsilon^2)^{-1-1/p}$ diverges for every nonnegative p . \square

This shows that even in the convergent case $p \geq 2$, the limit $u = \lim_{\varepsilon \rightarrow 0} u_\varepsilon$ is not a proper solution of equation (19).

6.1.1 The special case $p = 2$

Let us have a more detailed look at the (divergent) case $p = 2$. Then the function

$$u_\varepsilon(x, t) = (x^2 + \varepsilon^2)^{-1/2}, \quad (23)$$

at fixed $\varepsilon > 0$, is a solution to the advection-reaction equation

$$\frac{1}{c}\partial_t u_\varepsilon + \partial_x u_\varepsilon + x u_\varepsilon^3 = 0, \quad u_\varepsilon(x, 0) = (x^2 + \varepsilon^2)^{-1/2}. \quad (24)$$

According to Lemma 6.1, this solution is unique. We may study its \mathcal{G}^∞ -regularity properties.

Proposition 6.3. *The \mathcal{G}^∞ -singular support of $(u_\varepsilon)_{\varepsilon>0}$ is $\{(0, t) : t \geq 0\}$.*

Proof. Let $\chi(x) = (x^2 + 1)^{-1/2}$. Then $(u_\varepsilon(x, t) = (x^2 + \varepsilon^2)^{-1/2} = \chi_\varepsilon(x) = \varepsilon^{-1}\chi(x/\varepsilon)$. It is straightforward to show that the k th derivative of χ is of the form

$$\chi^{(k)}(x) = P_k(x)(x^2 + 1)^{-k-1/2}$$

where P_k is a polynomial of degree k . Therefore,

$$\chi_\varepsilon^{(k)}(x) = \varepsilon^{-k-1}P_k\left(\frac{x}{\varepsilon}\right)\left(\frac{x^2}{\varepsilon^2} + 1\right)^{-k-1/2} = \varepsilon^k P_k\left(\frac{x}{\varepsilon}\right)(x^2 + \varepsilon^2)^{-k-1/2}.$$

When $|x| \geq x_0 > 0$, the latter expression is bounded independently of $\varepsilon > 0$. Thus $(u_\varepsilon)_{\varepsilon>0}$ has the \mathcal{G}^∞ -property in the region $\{(x, t) : |x| > 0, t \geq 0\}$.

On the other hand, $\chi(x)$ is the derivative of $\operatorname{arsinh} x$, whose Taylor expansion shows that $\chi^{(k)}(x) \neq 0$ when k is an even integer. Thus

$$\chi_\varepsilon^{(k)}(0) = \varepsilon^{-k-1}\chi^{(k)}(0)$$

does not have the \mathcal{G}^∞ -property: the line $x = 0$ is contained in the \mathcal{G}^∞ -singular support. \square

This shows that the moderate sequential solution to (24) exhibits anomalous propagation of singularities. The initial \mathcal{G}^∞ -singularity at $x = 0$ is not propagated along the line $x = ct$ as in the linear case, but rather remains at $x = 0$ for all times.

Remark 6.4. *Actually, the classical initial value problem $\frac{1}{c}\partial_t v + \partial_x v + x v^3 = 0$, $v(x, 0) = v_0(x)$ can be solved explicitly. Transformation to characteristic coordinates $s = t$, $y = x - ct$ leads to an ordinary differential equation and to the solution*

$$v(x, t) = \frac{v_0(x - ct)}{\sqrt{(x^2 - (x - ct)^2)v_0^2(x - ct) + 1}}.$$

Inserting $v_0(x) = (x^2 + \varepsilon^2)^{-1/2}$ it turns out that by simple arithmetic, $v(x, t) = (x^2 + \varepsilon^2)^{-1/2}$, supporting the fact that $u_\varepsilon(x, t)$ as given above by (23) is indeed the solution. The same phenomenon also happens for $p \neq 2$ in (21) and (22).

6.2 Moderate sequential solutions to a nonlinear wave equation

Taking a further x -derivative, it is seen that $u_\varepsilon(x, t)$ given by (23) also solves the one-dimensional nonlinear wave equation

$$\frac{1}{c^2}\partial_t^2 u_\varepsilon - \partial_x^2 u_\varepsilon + u_\varepsilon^3 + 3x^2 u_\varepsilon^5 = 0, \quad u_\varepsilon(x, 0) = (x^2 + \varepsilon^2)^{-1/2}, \quad \partial_t u_\varepsilon(x, 0) = 0$$

for every $c > 0$. In this case, standard energy estimates can be used to show that the solution is unique.

Lemma 6.5. *Given $v_0 \in H^1(\mathbb{R})$, $v_1 \in L^2(\mathbb{R})$ of finite energy (defined by (26) below), the equation*

$$\frac{1}{c^2}\partial_t^2 v - \partial_x^2 v + v^3 + 3x^2 v^5 = 0, \quad v(x, 0) = v_0(x), \quad \partial_t v(x, 0) = v_1(x) \quad (25)$$

has a unique solution $v \in \mathcal{C}([0, \infty) : H^1(\mathbb{R})) \cap \mathcal{C}^1([0, \infty) : L^2(\mathbb{R}))$ of finite energy, where $c > 0$.

Proof. It is quite obvious that the energy

$$E(t) = \frac{1}{2} \int_{-\infty}^{\infty} \left(|\partial_t v|^2 + c^2 |\partial_x v|^2 + |v|^4 + 3x^2 |v|^6 \right) dx \quad (26)$$

is conserved. The proof follows standard arguments (see e.g. [35]). \square

At fixed $\varepsilon > 0$, $u_\varepsilon(x, 0) = (x^2 + \varepsilon^2)^{-1/2}$ belongs to $H^1(\mathbb{R})$ and, together with $\partial_t u_\varepsilon(x, 0) = 0$, forms initial data of finite energy. Thus the stationary solution $u_\varepsilon(x, t) = u_\varepsilon(x, 0)$ is the unique solution in this sense. The net $(u_\varepsilon)_{\varepsilon>0}$ provides a moderate sequential solution to the nonlinear wave equation (25). Its \mathcal{G}^∞ -singular support $\{(x, t), x = 0, t \geq 0\}$ has been computed in Proposition 6.3. Again, this differs from the linear case [14] and the nonlinear, classical case (Propositions 2.1, 2.3), according to which the singular support should be $\{(x, t), |x| = ct, t \geq 0\}$.

Anomalous propagation of singularities persists for sequential solutions.

Acknowledgements: I wish to thank the organizers of the INdAM Workshop *Anomalies in Partial Differential Equations* for providing an attractive environment for presenting the results of the paper. Discussions with various participants led to further insight. In particular, I would like to thank Lavi Karp, Sandra Lucente, Alberto Parmeggiani, Michael Reissig, Luigi Rodino and Michael Ruzhansky for helpful remarks. Thanks go to Hideo Deguchi for joint work on the topic since 2016.

Appendix: On multiplication of distributions

Let $S, T \in \mathcal{S}'(\mathbb{R}^n)$. The \mathcal{S}' -convolution of S and T is said to exist, if

$$(\varphi * \check{S})T \in \mathcal{D}'_{L^1}(\mathbb{R}^n), \quad \text{for all } \varphi \in \mathcal{S}(\mathbb{R}^n).$$

where $\check{S}(x) = S(-x)$. In this case, the convolution is defined by $\langle S * T, \varphi \rangle = \langle (\varphi * \check{S})T, 1 \rangle$, and $S * T$ belongs to $\mathcal{S}'(\mathbb{R}^n)$.

Let $u, v \in \mathcal{S}'(\mathbb{R}^n)$. If the \mathcal{S}' -convolution of $\mathcal{F}u$ and $\mathcal{F}v$ exists, one may define the *Fourier product*

$$u \cdot v = \mathcal{F}^{-1}(\mathcal{F}u * \mathcal{F}v).$$

The definition can be localized [1] as follows. Assume that for every $x \in \mathbb{R}^n$ there is a neighborhood Ω_x and $\chi_x \in \mathcal{D}(\mathbb{R}^n)$, $\chi_x \equiv 1$ on Ω_x , such that the \mathcal{S}' -convolution of $\mathcal{F}(\chi_x u)$ and $\mathcal{F}(\chi_x v)$ exists. Locally near x , the product $u \cdot v$ is defined to be $\mathcal{F}^{-1}(\mathcal{F}(\chi_x u) * \mathcal{F}(\chi_x v))$. Globally, it is defined by a partition of unity argument.

A special case arises when the distributions satisfy Hörmander's wave front set criterion [18], requiring that for every $(x, \xi) \in \mathbb{R}^n \times \mathbb{R}^n \setminus \{0\}$, $(x, \xi) \in \text{WF}(u)$ implies $(x, -\xi) \notin \text{WF}(v)$.

In space dimension $n = 1$, a very convenient case arises when $\text{supp } \mathcal{F}u$ and $\text{supp } \mathcal{F}v$ are contained in $[0, \infty)$. (In particular, Hörmander's criterion is fulfilled.) The basic example used in Section 3 is

$$u_0(x) = \frac{1}{x + i0} = \lim_{\varepsilon \rightarrow 0} \frac{1}{x + i\varepsilon} = \text{vp} \frac{1}{x} - i\pi\delta(x)$$

whose Fourier transform is $(\mathcal{F}u_0)(\xi) = -2\pi i H(\xi)$. The auto-convolution results in $(\mathcal{F}u_0 * \mathcal{F}u_0)(\xi) = -4\pi^2 \xi H(\xi)$. Thus $u_0^2 = \mathcal{F}^{-1}(\mathcal{F}u_0 * \mathcal{F}u_0)$ exists as Fourier product, and the formula shows that $u_0^2(x) = -u'_0(x)$. The remaining formulas used in Section 3 follow in the same way.

A more general definition of the product of distributions on \mathbb{R}^n can be obtained by regularization and passage to the limit. The *model product* of u and v is defined as

$$[u \cdot v] = \lim_{\varepsilon \rightarrow 0} (u * \varphi_\varepsilon)(v * \varphi_\varepsilon)$$

provided the limit exists for all mollifiers φ_ε of the form $\varphi_\varepsilon(x) = \varepsilon^{-n} \varphi(x/\varepsilon)$ with $\varphi \in \mathcal{D}(\mathbb{R}^n)$, $\int \varphi(x) dx = 1$, and is independent of the chosen mollifier. If the Fourier product exists, so does the model product.

In the one-dimensional case ($n = 1$), a yet more general definition is obtained by using the representation by boundary values of analytic functions, which was discussed in Section 3. Let $u \in \mathcal{D}'(\mathbb{R})$. Let

$$\tilde{u}_\varepsilon(x) = \hat{u}(x + i\varepsilon) - \hat{u}(x - i\varepsilon),$$

with the right-hand side as in (9). It was seen in Section 3 that $u(x) = \lim_{\varepsilon \rightarrow 0} \tilde{u}_\varepsilon(x)$. If $u \in \mathcal{D}'_{L^1}(\mathbb{R})$, \tilde{u}_ε is obtained by convolving u with the special mollifier $\psi_\varepsilon(x) = \varepsilon/(\pi(x^2 + \varepsilon^2))$. The *Tillmann product* [37] of two distributions u, v is defined by

$$u \circ v = \lim_{\varepsilon \rightarrow 0} \tilde{u}_\varepsilon \cdot \tilde{v}_\varepsilon$$

provided the limit exists. The definition does not work in higher space dimensions; there, harmonic regularization should be used [5]. In any case, the powers in (6) and (7) can also be understood in the sense of the Tillmann product.

Hörmander's criterion implies the existence of the Fourier product, which implies the existence of the model product and in turn also the existence of the Tillmann product. None of the implications can be reversed.

The other products used in this paper enter at different levels. For example, the most basic product of a smooth function with a distributions enters below Hörmander's criterion. The product in $H^s_{\text{loc}}(\mathbb{R}^n)$ when this space is an algebra ($s > n/2$) enters as a subcase of the Fourier product, but is independent of Hörmander's criterion. The Nemytskii operators in the form of a continuous map $L^p_{\text{loc}} \times L^q_{\text{loc}} \rightarrow L^1_{\text{loc}}$, $1/p + 1/q = 1$, enter at the level of the model product, but are independent of the Fourier product criterion. For more details on these circle of ideas, see [24].

References

- [1] Warren Ambrose. Products of distributions with values in distributions. *J. Reine Angew. Math.*, 315:73–91, 1980.
- [2] Michael Beals. Self-spreading and strength of singularities for solutions to semilinear wave equations. *Ann. of Math. (2)*, 118(1):187–214, 1983.

- [3] Michael Beals. *Propagation and interaction of singularities in nonlinear hyperbolic problems*, volume 3 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 1989.
- [4] Piotr Bizoń, Dieter Maison, and Arthur Wasserman. Self-similar solutions of semilinear wave equations with a focusing nonlinearity. *Nonlinearity*, 20(9):2061–2074, 2007.
- [5] Volker Boie. Multiplication of distributions. *Comment. Math. Univ. Carolin.*, 39(2):309–321, 1998.
- [6] Jean-François Colombeau. *Elementary introduction to new generalized functions*, volume 113 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1985.
- [7] Raphaël Côte, Carlos E. Kenig, Andrew Lawrie, and Wilhelm Schlag. Profiles for the radial focusing 4d energy-critical wave equation. *Comm. Math. Phys.*, 357(3):943–1008, 2018.
- [8] Vladimir G. Danilov and Georgii A. Omel’yanov. Weak asymptotics method and the interaction of infinitely narrow δ -solitons. *Nonlinear Anal.*, 54(4):773–799, 2003.
- [9] Hideo Deguchi. Personal communication and joint research. 2016 – 2019.
- [10] Hideo Deguchi and Michael Oberguggenberger. Propagation of singularities for generalized solutions to wave equations with discontinuous coefficients. *SIAM J. Math. Anal.*, 48(1):397–442, 2016.
- [11] Jean Dieudonné. *Éléments d’analyse. Tome III: Chapitres XVI et XVII*. Cahiers Scientifiques, Fasc. XXXIII. Gauthier-Villars Éditeur, Paris, 1970.
- [12] Roland Donniger. Strichartz estimates in similarity coordinates and stable blowup for the critical wave equation. *Duke Math. J.*, 166(9):1627–1683, 2017.
- [13] Marcelo R. Ebert and Michael Reissig. *Methods for partial differential equations*. Birkhäuser/Springer, Cham, 2018. Qualitative properties of solutions, phase space analysis, semilinear models.
- [14] Claudia Garetto and Michael Oberguggenberger. Generalized Fourier integral operator methods for hyperbolic equations with singularities. *Proc. Edinb. Math. Soc. (2)*, 57(2):423–463, 2014.
- [15] Claudia Garetto and Michael Ruzhansky. Hyperbolic second order equations with non-regular time dependent coefficients. *Arch. Ration. Mech. Anal.*, 217(1):113–154, 2015.
- [16] Vladimir Georgiev, Hans Lindblad, and Christopher D. Sogge. Weighted Strichartz estimates and global existence for semilinear wave equations. *Amer. J. Math.*, 119(6):1291–1319, 1997.
- [17] Michael Grosser, Michael Kunzinger, Michael Oberguggenberger, and Roland Steinbauer. *Geometric theory of generalized functions with applications to general relativity*, volume 537 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 2001.
- [18] Lars Hörmander. Fourier integral operators. I. *Acta Math.*, 127(1-2):79–183, 1971.
- [19] Jun Kato, Makoto Nakamura, and Tohru Ozawa. A generalization of the weighted Strichartz estimates for wave equations and an application to self-similar solutions. *Comm. Pure Appl. Math.*, 60(2):164–186, 2007.
- [20] Carlos E. Kenig and Frank Merle. Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation. *Acta Math.*, 201(2):147–212, 2008.
- [21] Joachim Krieger, Wilhelm Schlag, and Daniel Tataru. Slow blow-up solutions for the $H^1(\mathbb{R}^3)$ critical focusing semilinear wave equation. *Duke Math. J.*, 147(1):1–53, 2009.
- [22] Hans Lindblad. Counterexamples to local existence for semi-linear wave equations. *Amer. J. Math.*, 118(1):1–16, 1996.
- [23] Sandra Lucente. Critical exponents and where to find them. *Bruno Pini Math. Anal. Semin.*, 9:102–114, 2018.
- [24] Michael Oberguggenberger. *Multiplication of distributions and applications to partial differential equations*, volume 259 of *Pitman Research Notes in Mathematics Series*. Longman Scientific & Technical, Harlow, 1992.
- [25] Hartmut Pecher. Self-similar and asymptotically self-similar solutions of nonlinear wave equations. *Math. Ann.*, 316(2):259–281, 2000.
- [26] Hartmut Pecher. Sharp existence results for self-similar solutions of semilinear wave equations. *NoDEA Nonlinear Differential Equations Appl.*, 7(3):323–341, 2000.
- [27] Jeffrey Rauch. Singularities of solutions to semilinear wave equations. *J. Math. Pures Appl. (9)*, 58(3):299–308, 1979.

- [28] Jeffrey Rauch and Michael Reed. Propagation of singularities for semilinear hyperbolic equations in one space variable. *Ann. of Math. (2)*, 111(3):531–552, 1980.
- [29] Jeffrey Rauch and Michael Reed. Jump discontinuities of semilinear, strictly hyperbolic systems in two variables: creation and propagation. *Comm. Math. Phys.*, 81(2):203–227, 1981.
- [30] Michael Reed. Propagation of singularities for non-linear wave equations in one dimension. *Comm. Partial Differential Equations*, 3(2):153–199, 1978.
- [31] Francis Ribaud and Abdellah Youssfi. Global solutions and self-similar solutions of semilinear wave equation. *Math. Z.*, 239(2):231–262, 2002.
- [32] Elemer E. Rosinger. *Nonlinear partial differential equations*, volume 44 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam-New York, 1980. Sequential and weak solutions.
- [33] Michael Ruzhansky and Niyaz Tokmagambetov. Wave equation for operators with discrete spectrum and irregular propagation speed. *Arch. Ration. Mech. Anal.*, 226(3):1161–1207, 2017.
- [34] Laurent Schwartz. *Théorie des distributions*. Publications de l’Institut de Mathématique de l’Université de Strasbourg, No. IX-X. Nouvelle édition, entièrement corrigée, refondue et augmentée. Hermann, Paris, 1966.
- [35] Michael Struwe. Semi-linear wave equations. *Bull. Amer. Math. Soc. (N.S.)*, 26(1):53–85, 1992.
- [36] Hiroyuki Takamura and Kyouhei Wakasa. Global existence for semilinear wave equations with the critical blow-up term in high dimensions. *J. Differential Equations*, 261(2):1046–1067, 2016.
- [37] Heinz Günther Tillmann. Darstellung der Schwartzschen Distributionen durch analytische Funktionen. *Math. Z.*, 77:106–124, 1961.