

A Reflection Formula for the Gaussian Hypergeometric Function of Matrix Argument

Donald Richards* and Qifu Zheng†

February 14, 2020

Abstract

We obtain a reflection formula for the Gaussian hypergeometric function of real symmetric matrix argument. We also show that this result extends to the Gaussian hypergeometric function defined over the symmetric cones, and even to generalizations of the Gaussian hypergeometric function defined in terms of series of Jack polynomials. We also obtain a quadratic transformation formula for the 2×2 Gaussian hypergeometric function of Hermitian matrix argument.

1 Introduction

The Gaussian hypergeometric series is well-known to satisfy a large number of linear (contiguous), quadratic, and nonlinear transformation formulas. We refer to Andrews, *et al.* [1, Chapter 3], Erdélyi, *et al.* [3, Chapter II], or Olde Daalhuis [15, Chapter 15] for extensive treatments of these formulas.

In this paper, we are concerned with a reflection formula,

$${}_2F_1(a, b; a + b + 1 - c; 1 - x) = \frac{\Gamma(c - a)\Gamma(c - b)}{\Gamma(c)\Gamma(c - a - b)} {}_2F_1(a, b; c; x), \quad (1.1)$$

valid for $0 < x < 1$, $-a \in \mathbb{N}$, and for b, c satisfying various conditions. We extend the identity (1.1) to the Gaussian hypergeometric function of matrix argument [5, 9, 10, 14], to the Gaussian hypergeometric function defined on symmetric cones, and even to the Gaussian hypergeometric function defined in terms of Jack polynomials [4, 13].

*Department of Statistics, Pennsylvania State University, University Park, PA 16802, U.S.A. E-mail address: richards@stat.psu.edu.

†Department of Mathematics & Statistics, The College of New Jersey, Ewing, NJ 08628, U.S.A. E-mail address: zheng@tcnj.edu.

MSC 2010 subject classifications: Primary 33C80; Secondary 33C15.

Key words and phrases. Hypergeometric function of matrix argument, Jack polynomial, Schur function, zonal polynomial.

We also consider the problem of deriving quadratic transformations for the Gaussian hypergeometric function of matrix argument, and we derive such a transformation for the hypergeometric functions defined on the space of Hermitian 2×2 matrices.

2 Remarks on the classical reflection formula (1.1)

The reflection formula (1.1) is a special case of a well-known linear transformation of the Gaussian hypergeometric function; see [15, Eq. (15.8.7)]. The formula is also stated indirectly by Whittaker and Watson [17, p. 296, Miscellaneous Example 2], who posed the problem of proving that if $-a \in \mathbb{N}$ and $b, c \notin \mathbb{N}$ then the ratio

$$\frac{{}_2F_1(a, b; a + b + 1 - c; 1 - x)}{{}_2F_1(a, b; c; x)}, \quad (2.1)$$

is independent of x ; further, the reader is asked to calculate the corresponding constant. Once it has been proved that this ratio is independent of x , the constant of proportionality can be found by setting $x = 1$ and applying Gauss' formula,

$${}_2F_1(a, b; c; 1) = \frac{\Gamma(c) \Gamma(c - a - b)}{\Gamma(c - a) \Gamma(c - b)}, \quad (2.2)$$

for $\operatorname{Re}(c - a - b) > 0$. We remark that we were unable to infer, despite a close reading of [17, Chapter XIV], whether Whittaker and Watson had expected readers to solve this problem using a method alternative to the previously-mentioned linear transformation, and this motivated us to investigate several approaches to proving (1.1).

For $a \in (0, 1)$, $b = 1 - a$, and $c = 1$, Berndt and Chan [2] showed that certain elliptic modular functions are expressible in terms of the inverse functions of ratios of the type (2.1). For $n \in \mathbb{N}$, Hannah [8, p. 88] stated (1.1) in the form,

$${}_2F_1(-n, b; c; 1 - x) = \frac{(c - b)_n}{(c)_n} {}_2F_1(-n, b; -n + b + 1 - c; x),$$

$n \in \mathbb{N}$, where $c, 1 + b - c - n \notin \{0, -1, -2, \dots, -n + 1\}$, and remarked that a proof can be obtained by induction on n .

3 A reflection formula for the Gaussian hypergeometric function of matrix argument

Another approach to establishing (1.1) is by means of Euler's hypergeometric differential equation for the ${}_2F_1$ function. As we now show, this approach generalizes to the Gaussian hypergeometric function of matrix argument.

First, we consider the case of $m \times m$ real symmetric matrix arguments, X . For $\operatorname{Re}(a) > \frac{1}{2}(m-1)$, let

$$\Gamma_m(a) = \pi^{m(m-1)/4} \prod_{j=1}^m \Gamma\left(a - \frac{1}{2}(j-1)\right)$$

denote the *multivariate gamma function*. A *partition* $\kappa = (k_1, \dots, k_m)$ is a m -tuple of nonnegative integers k_1, \dots, k_m such that $k_1 \geq \dots \geq k_m$; the *weight* of κ is $|\kappa| = k_1 + \dots + k_m$; the *partitional rising factorial* is

$$[a]_\kappa = \prod_{j=1}^m \left(a - \frac{1}{2}(j-1)\right)_{k_j},$$

where $(a)_k = a(a+1)\cdots(a+k-1)$ is the classical rising factorial; and we denote by $Z_\kappa(X)$ the corresponding *zonal polynomial* [4, 5, 10].

For $a, b, c \in \mathbb{C}$ and for X , a $m \times m$ real symmetric matrix, the *Gaussian hypergeometric function of matrix argument* is

$${}_2F_1(a, b; c; X) = \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{|\kappa|=k} \frac{[a]_\kappa [b]_\kappa}{[c]_\kappa} Z_\kappa(X), \quad (3.1)$$

where the inner sum is over all partitions κ of weight k . The existence of this series requires that $[c]_\kappa \neq 0$ for all partitions κ , equivalently, $-c + 1 + \frac{1}{2}(j-1) \notin \mathbb{N}$ for all $j = 1, \dots, m$. Denote by x_1, \dots, x_m the eigenvalues of X ; if the series (3.1) is non-terminating then it converges for all $m \times m$ real symmetric matrices X such that $\|X\| := \max\{|x_1|, \dots, |x_m|\} < 1$, and otherwise it converges for all such X [5].

The Gaussian hypergeometric function of matrix argument can also be defined by a generalized Euler-type integral over the cone of positive definite matrices [5, 9]. Letting I_m denote the $m \times m$ identity matrix then, as a consequence of those generalized Euler-type integrals, Herz [9] generalized (2.2) to:

$${}_2F_1(a, b; c; I_m) = \frac{\Gamma_m(c) \Gamma_m(c-a-b)}{\Gamma_m(c-a) \Gamma_m(c-b)}, \quad (3.2)$$

for $\operatorname{Re}(c-a-b) > \frac{1}{2}(m-1)$.

The zonal polynomials $Z_\kappa(X)$ depend only on x_1, \dots, x_m , the eigenvalues of X , and are symmetric functions of x_1, \dots, x_m ; therefore, the same holds for the hypergeometric function of matrix argument. Muirhead [14, p. 274, Theorem 7.5.5] proved that the function ${}_2F_1(a, b; c; X)$ in (3.1) is the unique solution of each of the partial differential

equations

$$x_i(1-x_i)\frac{\partial^2 F}{\partial x_i^2} + \left[c - \frac{1}{2}(m-1) - (a+b+1 - \frac{1}{2}(m-1))x_i + \frac{1}{2} \sum_{\substack{j=1 \\ j \neq i}}^m \frac{x_i(1-x_i)}{x_i-x_j} \right] \frac{\partial F}{\partial x_i} - \frac{1}{2} \sum_{\substack{j=1 \\ j \neq i}}^m \frac{x_j(1-x_j)}{x_i-x_j} \frac{\partial F}{\partial x_j} = abF, \quad (3.3)$$

$i = 1, \dots, m$, subject to the conditions that:

- (i) F is a symmetric function of x_1, \dots, x_m ;
- (ii) F is analytic at $X = 0$, i.e., F is expressible in a neighborhood of 0 as an infinite series of zonal polynomials $F(X) = \sum_{k=0}^{\infty} \sum_{|\kappa|=k} c_{\kappa} Z_{\kappa}(X)$, where the coefficients c_{κ} do not depend on m ; and
- (iii) $F(0) = 1$.

The following result establishes a statement in [15, Eq. (35.7.9)] and also provides a correction to the hypotheses state there on the parameters a, b, c .

Proposition 3.1. *Suppose that $-a+1+\frac{1}{2}(j-1) \in \mathbb{N}$ for some $j = 1, \dots, m$. Further, suppose that $-c+1+\frac{1}{2}(j-1) \notin \mathbb{N}$ and $-a-b+c-\frac{1}{2}(m-j) \notin \mathbb{N}$ for all $j = 1, \dots, m$. If $0 < X < I_m$, i.e., all eigenvalues of X are in the interval $(0, 1)$, then*

$${}_2F_1(a, b; a+b+1-c+\frac{1}{2}(m-1); I_m - X) = \frac{\Gamma_m(c-a)\Gamma_m(c-b)}{\Gamma_m(c)\Gamma_m(c-a-b)} {}_2F_1(a, b; c; X). \quad (3.4)$$

Proof. Under the stated hypotheses on b and c , the partitional rising factorials $[c]_{\kappa}$ and $[a+b+1-c+\frac{1}{2}(m-1)]_{\kappa}$ are non-zero for all partitions κ ; therefore, both sides of (3.4) are well-defined for all symmetric $m \times m$ matrices X such that $\|X\| < 1$ and $\|I_m - X\| < 1$. Moreover, under the assumption on a , both sides of (3.4) are terminating series, i.e., polynomials in X , hence are analytic at $X = 0$. Moreover, by (3.2), both sides are equal at $X = 0$.

Replacing x_i by $1-x_i$, $i = 1, \dots, m$, in each of the partial differential equations in the system (3.3), we see that the outcome of this transformation is that c is replaced by $a+b+1-c+\frac{1}{2}(m-1)$. Finally, applying the uniqueness result of Muirhead [14, Theorem 7.5.5], we find that the left- and right-hand sides of (3.4) coincide. \square

The reflection formula (3.4) extends to the hypergeometric function defined on the symmetric cones [4, 5] or in terms of series of Jack polynomials [13]. These generalizations satisfy a system of partial differential equations defined by means of the

generalized Muirhead operators; *viz.*, for an arbitrary parameter $d > 0$, the system of partial differential equations is:

$$x_i(1-x_i)\frac{\partial^2 F}{\partial x_i^2} + \left[c - (m-1)d - (a+b+1 - (m-1)d)x_i + d \sum_{\substack{j=1 \\ j \neq i}}^m \frac{x_i(1-x_i)}{x_i-x_j} \right] \frac{\partial F}{\partial x_i} - d \sum_{\substack{j=1 \\ j \neq i}}^m \frac{x_j(1-x_j)}{x_i-x_j} \frac{\partial F}{\partial x_j} = abF, \quad (3.5)$$

$i = 1, \dots, m$.

Denote by $J_\kappa(x_1, \dots, x_m; d)$ the corresponding Jack polynomials. Then each of the equations (3.5) has a common unique solution, denoted by ${}_2F_1(a, b; c; x_1, \dots, x_m; d)$, subject to the conditions:

- (iv) F is a symmetric function of x_1, \dots, x_m ;
- (v) F is analytic at $(0, \dots, 0)$, i.e., F is expressible in a neighborhood of the origin as a series of Jack polynomials $F(x_1, \dots, x_m) = \sum_{k=0}^{\infty} \sum_{|\kappa|=k} c_\kappa J_\kappa(x_1, \dots, x_m; d)$, where the coefficients c_κ do not depend on m ; and
- (vi) $F(0, \dots, 0) = 1$.

For $\text{Re}(a) > (m-1)d$, let

$$\Gamma_m(a; d) = \pi^{m(m-1)d/2} \prod_{j=1}^m \Gamma(a - (j-1)d).$$

denote the corresponding multivariate gamma function. By applying the same argument used to establish Proposition 3.1, we obtain the reflection formula,

$$\begin{aligned} & {}_2F_1(a, b; a+b+1-c+(m-1)d; 1-x_1, \dots, 1-x_m; d) \\ &= \frac{\Gamma_m(c-a; d)\Gamma_m(c-b; d)}{\Gamma_m(c; d)\Gamma_m(c-a-b; d)} {}_2F_1(a, b; c; x_1, \dots, x_m; d), \end{aligned}$$

subject to the conditions that $-a+1+(j-1)d \in \mathbb{N}$ for some $j = 1, \dots, m$, and $-c+1+(j-1)d \notin \mathbb{N}$ and $-a-b+c-(m-j)d \notin \mathbb{N}$ for all $j = 1, \dots, m$.

4 A quadratic transformation in the 2×2 Hermitian case

The problem of deriving nonlinear transformation formulas for the special functions of matrix argument was raised first by Herz [9, p. 488], who noted the difficulty of deriving

a quadratic transformation between the Bessel and confluent hypergeometric functions of matrix argument. Such transformations are still generally unexplored and do not appear to follow from Euler integral representations, or manipulation of zonal polynomial or Jack polynomial series expansions. It appears that such quadratic transformations will require analysis of the full monodromy group of the system of hypergeometric differential equations (3.5), and we note that Korányi [12] studied a “diagonal” subgroup of that monodromy group and obtained a generalization of Kummer’s twenty-four solutions for the classical Gaussian hypergeometric differential equation.

A well-known quadratic transformation for the classical Gaussian hypergeometric function is

$${}_2F_1\left(\alpha, \alpha - \beta + \frac{1}{2}; \beta + \frac{1}{2}; t^2\right) = (1+t)^{-2\alpha} {}_2F_1\left(\alpha, \beta; 2\beta; \frac{4t}{(1+t)^2}\right); \quad (4.1)$$

see [1, p. 176, Exercise 1(d)] or [15, Eq. (15.8.21)]. We now illuminate the difficulties of deriving quadratic transformations in the matrix case by extending (4.1) to the hypergeometric function of *Hermitian* matrix argument, i.e., defined on the space of $m \times m$ Hermitian matrices and corresponding to $d = 2$ in (3.5).

In the Hermitian case, the multivariate gamma function is

$$\Gamma_m(a) = \pi^{m(m-1)/2} \prod_{j=1}^m \Gamma(a - j + 1),$$

$\operatorname{Re}(a) > m - 1$; the partitional rising factorial is

$$[a]_{\kappa} = \prod_{j=1}^m (a - j + 1)_{k_j};$$

the zonal polynomial, Z_{κ} , is a multiple of the well-known Schur function s_{κ} [6, 10]; and the Gaussian hypergeometric function of X , a $m \times m$ Hermitian matrix, is

$${}_2F_1(a, b; c; X) = \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{|\kappa|=k} \frac{[a]_{\kappa} [b]_{\kappa}}{[c]_{\kappa}} Z_{\kappa}(X). \quad (4.2)$$

The Gaussian hypergeometric function of $m \times m$ Hermitian matrix arguments, X and Y , is defined as

$${}_2F_1(a, b; c; X, Y) = \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{|\kappa|=k} \frac{[a]_{\kappa} [b]_{\kappa}}{[c]_{\kappa}} \frac{Z_{\kappa}(X) Z_{\kappa}(Y)}{Z_{\kappa}(I_m)}, \quad (4.3)$$

In both series (4.2) and (4.3), the parameter c is such that $-c + j \notin \mathbb{N}$ for all $j = 1, \dots, m$. If the series (4.2) is non-terminating then it converges for all X such that $\|X\| < 1$; further, if (4.3) is non-terminating then it converges if $\|X\| \cdot \|Y\| < 1$; see [5, Theorem 6.3].

Since the zonal polynomials depend only on the eigenvalues of their matrix arguments then we may assume, with no loss of generality, that $X = \text{diag}(x_1, \dots, x_m)$ and $Y = \text{diag}(y_1, \dots, y_m)$. Define

$$V(X) = \prod_{1 \leq i < j \leq m} (x_i - x_j),$$

and

$$c_{2,1} = \beta_m^{-1} \frac{\prod_{i=1}^m (c - m + 1)_{m-i}}{\prod_{i=1}^m (i-1)! (a - m + 1)_{m-i} (b - m + 1)_{m-i}}.$$

The hypergeometric function of Hermitian matrix arguments can be expressed in terms of a determinant of classical hypergeometric functions. Denote by $\det(a_{ij})$ the determinant of a $m \times m$ matrix with (i, j) th entry a_{ij} ; then by [6, Theorem 4.2], [11],

$${}_2F_1(a, b; c; X, Y) = c_{2,1} \frac{\det({}_2F_1(a - m + 1, b - m + 1; c - m + 1; x_i y_j))}{V(X)V(Y)}, \quad (4.4)$$

where the ${}_2F_1$ functions on the right-hand side are the classical Gaussian hypergeometric functions, and L'Hospital's rule is to be applied if any of x_1, \dots, x_m , or y_1, \dots, y_m , coincide. In particular, as in [7, Eq. 5], by evaluating the limit as $Y \rightarrow I_m$, i.e., $y_1, \dots, y_m \rightarrow 1$, it follows from (4.3) and (4.4) that

$${}_2F_1(a, b; c; X) = \frac{\det(x_i^{m-j} {}_2F_1(a - j + 1, b - j + 1; c - j + 1; x_i))}{V(X)}. \quad (4.5)$$

Setting $(a, b; c) = (\alpha + m - 1, \alpha - \beta + m - \frac{1}{2}; \beta + m - \frac{1}{2})$, equivalently

$$(a - m + 1, b - m + 1; c - m + 1) = (\alpha, \alpha - \beta + \frac{1}{2}; \beta + \frac{1}{2}), \quad (4.6)$$

then we obtain

$$\begin{aligned} c_{2,1}^{-1} V(X)V(Y) {}_2F_1(a, b; c; X, Y) &= \det({}_2F_1(a - m + 1, b - m + 1; c - m + 1; x_i y_j)) \\ &= \det({}_2F_1(\alpha, \alpha - \beta + \frac{1}{2}; \beta + \frac{1}{2}; x_i y_j)). \end{aligned}$$

Applying (4.1), we obtain

$$\begin{aligned} c_{2,1}^{-1} V(X^2)V(Y^2) {}_2F_1(a, b; c; X^2, Y^2) \\ = \det\left(\left(1 + x_i y_j\right)^{-2\alpha} {}_2F_1\left(\alpha, \beta; 2\beta; \frac{4x_i y_j}{(1 + x_i y_j)^2}\right)\right). \end{aligned} \quad (4.7)$$

From now on, we set $m = 2$. Applying to the right-hand side of (4.7) the 2×2 determinantal identity,

$$\det(a_{ij} b_{ij}) = b_{11} b_{22} \det(a_{ij}) + a_{12} a_{21} \det(b_{ij}), \quad (4.8)$$

we obtain

$$\begin{aligned}
& c_{2,1}^{-1} V(X^2) V(Y^2) {}_2F_1(a, b; c; X^2, Y^2) \\
&= \det \left((1 + x_i y_j)^{-2\alpha} \right) \cdot \prod_{j=1}^2 {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_j y_j}{(1 + x_j y_j)^2} \right) \\
&\quad + (1 + x_1 y_2)^{-2\alpha} (1 + x_2 y_1)^{-2\alpha} \cdot \det \left({}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_i y_j}{(1 + x_i y_j)^2} \right) \right). \quad (4.9)
\end{aligned}$$

Next, we divide both sides of the latter equation by $V(Y^2) \equiv y_1^2 - y_2^2$, and let $Y \rightarrow I_2$, i.e., $y_1, y_2 \rightarrow 1$. By applying L'Hospital's rule, we obtain

$$\lim_{Y \rightarrow I_2} \frac{\det \left((1 + x_i y_j)^{-2\alpha} \right)}{V(Y^2)} = \alpha (1 + x_1)^{-2\alpha-1} (1 + x_2)^{-2\alpha-1} (x_2 - x_1).$$

Also, using the fact that

$$\left. \frac{\partial}{\partial y_1} \frac{4xy_1}{(1 + xy_1)^2} \right|_{y_1=1} = \frac{4x(1-x)}{(1+x)^3},$$

and the well-known formula,

$$\frac{d}{dy} {}_2F_1(a, b; c; y) = \frac{ab}{c} {}_2F_1(a+1, b+1; c+1; y),$$

we obtain

$$\begin{aligned}
& \lim_{Y \rightarrow I_2} \frac{\det \left({}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_i y_j}{(1 + x_i y_j)^2} \right) \right)}{V(Y^2)} \\
&= \frac{1}{2} \lim_{y_1 \rightarrow 1} \frac{1}{y_1 - 1} \left| \begin{array}{cc} {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_1 y_1}{(1 + x_1 y_1)^2} \right) & {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_1}{(1 + x_1)^2} \right) \\ {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_2 y_1}{(1 + x_2 y_1)^2} \right) & {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_2}{(1 + x_2)^2} \right) \end{array} \right| \\
&= \alpha \det(a_{ij} b_{ij}), \quad (4.10)
\end{aligned}$$

where, for $i, j = 1, 2$,

$$\begin{aligned}
a_{ij} &= \left(\frac{1 - x_i}{4(1 + x_i)^2} \right)^{2-j}, \\
b_{ij} &= \left(\frac{4x_i}{(1 + x_i)^2} \right)^{2-j} {}_2F_1 \left(\alpha + 2 - j, \beta + 2 - j; 2\beta + 2 - j; \frac{4x_i}{(1 + x_i)^2} \right).
\end{aligned}$$

Applying (4.8), we find that the determinant $\det(a_{ij}b_{ij})$ in (4.10) equals

$$\begin{aligned} & \frac{4x_1}{(1+x_1)^2} \cdot \det \left(\left(\frac{1-x_i}{4(1+x_i)^2} \right)^{2-j} \right) \cdot \prod_{j=1}^2 {}_2F_1 \left(\alpha+2-j, \beta+2-j; 2\beta+2-j; \frac{4x_j}{(1+x_j)^2} \right) \\ & + \frac{1-x_2}{4(1+x_2)^2} \cdot \det \left(\left(\frac{4x_i}{(1+x_i)^2} \right)^{2-j} {}_2F_1 \left(\alpha+2-j, \beta+2-j; 2\beta+2-j; \frac{4x_i}{(1+x_i)^2} \right) \right). \end{aligned}$$

Note that

$$\det \left(\left(\frac{1-x_i}{4(1+x_i)^2} \right)^{2-j} \right) = \frac{x_2-x_1}{4(1+x_1)(1+x_2)}$$

and, by (4.5),

$$\begin{aligned} & \det \left(\left(\frac{4x_i}{(1+x_i)^2} \right)^{2-j} {}_2F_1 \left(\alpha+2-j, \beta+2-j; 2\beta+2-j; \frac{4x_i}{(1+x_i)^2} \right) \right) \\ & = V(4X(I_2+X)^{-2}) {}_2F_1(\alpha+1, \beta+1; 2\beta+1; 4X(I_2+X)^{-2}). \end{aligned}$$

The conclusion is that when both sides of (4.9) are divided by $V(Y^2)$ and then $Y \rightarrow I_2$, we obtain after some simplifications the result,

$$\begin{aligned} & c_{2,1}^{-1} V(X^2) {}_2F_1(a, b; c; X^2) \\ & = \alpha(x_2-x_1) \prod_{j=1}^2 (1+x_j)^{-2\alpha-1} {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_j}{(1+x_j)^2} \right) \\ & \quad + \alpha(1+x_1)^{-2\alpha-2} (1+x_2)^{-2\alpha-1} (x_2-x_1) \\ & \quad \times \left[\frac{x_1}{(1+x_1)} \prod_{j=1}^2 {}_2F_1 \left(\alpha+2-j, \beta+2-j; 2\beta+2-j; \frac{4x_j}{(1+x_j)^2} \right) \right. \\ & \quad \left. - \frac{(1-x_2)(1-x_1x_2)}{(1+x_2)^3} {}_2F_1(\alpha+1, \beta+1; 2\beta+1; 4X(I_2+X)^{-2}) \right]. \end{aligned}$$

Dividing both sides of the latter equation by x_1-x_2 and using (4.6) to substitute for (a, b, c) in terms of (α, β) , we obtain

$$\begin{aligned} & c_{2,1}^{-1} (x_1+x_2) {}_2F_1(\alpha+1, \alpha-\beta+\frac{3}{2}; \beta+\frac{3}{2}; X^2) \\ & = -\alpha \prod_{j=1}^2 (1+x_j)^{-2\alpha-1} {}_2F_1 \left(\alpha, \beta; 2\beta; \frac{4x_j}{(1+x_j)^2} \right) \\ & \quad + \alpha(1+x_1)^{-2\alpha-2} (1+x_2)^{-2\alpha-1} \\ & \quad \times \left[-\frac{x_1}{(1+x_1)} \prod_{j=1}^2 {}_2F_1 \left(\alpha+2-j, \beta+2-j; 2\beta+2-j; \frac{4x_j}{(1+x_j)^2} \right) \right. \\ & \quad \left. + \frac{(1-x_2)(1-x_1x_2)}{(1+x_2)^3} {}_2F_1(\alpha+1, \beta+1; 2\beta+1; 4X(I_2+X)^{-2}) \right]. \end{aligned}$$

References

- [1] Andrews, G. E., Askey, R., and Roy, R. (1999). *Special Functions*. Cambridge University Press, New York.
- [2] Berndt, B. C., and Chan, H. H. (1999). Ramanujan and the modular j -invariant. *Canad. Math. Bull.*, **42**, 427–440.
- [3] Erdélyi, A., Magnus, W., Oberhettinger, F., and Tricomi, F. G. (1953). *Higher Transcendental Functions*, Volume 1. McGraw-Hill, New York, NY.
- [4] Faraut, J., and Korányi, A. (1994). *Analysis on Symmetric Cones*, Oxford University Press, Oxford, UK.
- [5] Gross, K. I., and Richards, D. St. P. (1987). Special functions of matrix argument. I. Algebraic induction, zonal polynomials, and hypergeometric functions. *Trans. Amer. Math. Soc.*, **301**, 781–811.
- [6] Gross, K. I., and Richards, D. St. P. (1989). Total positivity, spherical series, and hypergeometric functions of matrix argument. *J. Approx. Theory*, **59**, 224–246.
- [7] Gross, K. I., and Richards, D. St. P. (1991). Hypergeometric functions on complex matrix space. *Bull. Amer. Math. Soc.*, **24**, 349–355.
- [8] Hannah, J. P. (2013). *Identities for the Gamma and Hypergeometric Functions: An Overview from Euler to the Present*. M.Sc. Dissertation, University of the Witwatersrand.
- [9] Herz, C. S. (1955). Bessel functions of matrix argument. *Ann. Math.*, **64**, 474–523.
- [10] James, A. T. (1964). Distributions of matrix variates and latent roots derived from normal samples. *Ann. Math. Statist.*, **35**, 475–501.
- [11] Khatri, C. G. (1970). On the moments of traces of two matrices in three situations for complex multivariate normal populations. *Sankhyā Ser. A*, **32**, 65–80.
- [12] Korányi, A. (1990). Transformation properties of the generalized Muirhead operators. *Colloq. Math.*, **60/61**, 665–669.
- [13] Macdonald, I. G. (2013). Hypergeometric functions, I. Unpublished manuscript, arXiv: <http://arxiv.org/abs/1309.4568>
- [14] Muirhead, R. J. (1982). *Aspects of Multivariate Statistical Analysis*. Wiley, New York.

- [15] Olde Daalhuis, A. B. (2010). Chapter 15: Hypergeometric Function. *NIST Handbook of Mathematical Functions*, pp. 321–349, Cambridge University Press, New York.
- [16] Richards, D. St. P. (2010). Chapter 35: Functions of Matrix Argument. *NIST Handbook of Mathematical Functions*, pp. 767–774, Cambridge University Press, New York.
- [17] Whittaker, E. T., and Watson, G. N. (1950). *A Course of Modern Analysis*, fourth edition, reprinted. Cambridge University Press, Cambridge, England.