

Rigidity of matrix group actions on CAT(0) spaces with possible parabolic isometries and uniquely arcwise connected spaces

Shengkui Ye

February 14, 2020

Abstract

It is well-known that $\mathrm{SL}_n(\mathbf{Q}_p)$ acts without fixed points on an $(n-1)$ -dimensional CAT(0) space (the affine building). We prove that $n-1$ is the smallest dimension of CAT(0) spaces on which matrix groups act without fixed points. Explicitly, let R be an associative ring with identity and $E'_n(R)$ the extended elementary subgroup. Any isometric action of $E'_n(R)$ on a complete CAT(0) space X^d of dimension $d < n-1$ has a fixed point. Similar results are discussed for automorphism groups of free groups. Furthermore, we prove that any action of $\mathrm{Aut}(F_n)$, $n \geq 3$, on a uniquely arcwise connected space by homeomorphisms has a fixed point.

1 Introduction

Let p be a prime, \mathbf{Q}_p the p -adic number field and $\mathrm{SL}_n(\mathbf{Q}_p)$ the special linear group. It is well-known that $\mathrm{SL}_n(\mathbf{Q}_p)$ acts without fixed points on the affine building X , which is an $(n-1)$ -dimensional CAT(0) space. In the article, we prove that $n-1$ is the smallest dimension of CAT(0) spaces on which $\mathrm{SL}_n(\mathbf{Q}_p)$ acts without fixed points. More generally, let R be an associative ring with identity and $E'_n(R) < \mathrm{GL}_n(R)$ the extended elementary subgroup (cf. Section 2.2), which is an analog of $\mathrm{SL}_n(\mathbf{Q}_p)$ for general rings. Our first result is the following.

Theorem 1.1 *Any isometric action of the extended elementary group $E'_n(R)$ on a complete CAT(0) space X of dimension $d < n-1$ has a fixed point.*

It is a classical result of Serre [22] that any isometric action of $\mathrm{SL}_n(\mathbb{Z})$ ($n \geq 3$) on a simplicial tree has a fixed point. Farb [16] considered a high-dimensional analog. For a reduced, irreducible root system Φ has rank $r \geq 2$ and a finitely generated commutative ring R , let $E(\Phi, R)$ be the elementary subgroup of the Chevalley group $G(\Phi, R)$. Farb [16] proves that any action of $E(\Phi, R)$ on a complete CAT(0) space X of dimension $d < r-1$ by *semisimple* isometries has

a fixed point. This gives also a generalization of a result obtained by Fukunaga [17] concerning groups acting on trees. Let Σ_g be the orientable closed surface and $\text{MCG}(\Sigma_g)$ the mapping class group. Bridson [6, 7] proves any action of $\text{MCG}(\Sigma_g)$ on a complete $\text{CAT}(0)$ space X of dimension $d < g - 1$ by *semisimple* isometries has a fixed point. The author [23] obtains similar results for actions of the elementary subgroup $E_n(R)$ and the quadratic elementary subgroup $EU_{2n}(R, \Lambda)$ for general rings. However, most of results are proved for semisimple actions. The group action on $\text{CAT}(0)$ spaces of automorphism groups of free groups is studied by Bridson [8, 9] and Varghese [24]. Barnhill [1] considers the property FA_n for Coxeter groups.

In this article, we study group actions on $\text{CAT}(0)$ spaces with possible parabolic isometries. Note that the semisimple isometries could be very different from generic isometries. For example, any action of the special linear group $\text{SL}_n(\mathbb{Z})$ ($n \geq 3$) on a complete $\text{CAT}(0)$ space X by semisimple isometries has a fixed point, while $\text{SL}_n(\mathbb{Z})$ acts on the symmetric space $\text{SL}_n(\mathbb{R})/\text{SO}(n)$ properly and thus without fixed points. It is an open problem in geometric group theory to study the minimal dimensions of $\text{CAT}(0)$ spaces on which matrix groups or automorphism groups of free groups act without fixed points (see Bridson [4], Question 9.2).

In order to state our results easily, it is better to define the following.

Definition 1.2 (*relative property \mathcal{FA}_n*) *Let H be a subgroup of a group G . The pair (G, H) has the relative property \mathcal{FA}_n (resp. $s\mathcal{FA}_n$) if whenever G acts on an n -dimensional complete $\text{CAT}(0)$ space X by isometries (resp. semisimple isometries), the subgroup H has a fixed point.*

We call a group G has property \mathcal{FA}_n if (G, G) has the relative property \mathcal{FA}_n . It should be pointed that we don't require the isometries to be semisimple in the definition of \mathcal{FA}_n . Note that the property $s\mathcal{FA}_n$ is the same as the strong property FA_n considered by Farb [16].

Theorem 1.3 *Let R be an associative ring and $E'_n(R)$ the extended elementary subgroup. Then $(R^n \rtimes E'_n(R), R^n)$ has the relative property \mathcal{FA}_{n-1} .*

Since $E_n(\mathbb{Z}[\frac{1}{p}]) \rtimes \mathbb{Z}[\frac{1}{p}]^n$ as a subgroup of $\text{GL}_{n+1}(\mathbb{Q}_p)$ acts on an affine building X of dimension n without fixed points, the dimension in Theorem 1.3 is sharp.

Let $\text{Aut}(F_n)$ be the automorphism group of the rank- n free group F_n .

Theorem 1.4 *$(\text{Aut}(F_n), \text{Aut}(F_2))$ has the relative property \mathcal{FA}_{n-3} for any $n \geq 4$.*

Corollary 1.5 *Any isometric action of $\text{Aut}(F_n)$ ($n \geq 4$) on a complete $\text{CAT}(0)$ space X of dimension $d \leq 2[\frac{n}{3}]$ has a fixed point, i.e. $\text{Aut}(F_n)$ has property $\mathcal{FA}_{2[n/3]}$. Here $[n/3]$ is the integer part.*

The previous result was also observed by Bridson [8]. The case of $d \leq 2[n/2]$ is already known by Varghese [24].

The proofs of the above theorems are based on Helly’s theorem and the theory of Coxeter groups. The method has further applications to group actions on 1-dimensional spaces, like dendrites and unique arcwise connected spaces. Recall that a dendrite X is a connected compact metrizable space (continuum) such that any two points are the extremities of a unique arc in X (cf. [15]). Let Γ be a lattice in a semisimple algebraic group of rank at least two. Buchesne and Monod [15] prove that any action of Γ on a dendrite X fixes a fixed point or a pair of two points. An \mathbb{R} -tree is a geodesic metric space in which there is a unique arc connecting each pair of points. Bogopolski [2] prove that any isometric action of $\text{Aut}(F_n)$ on a simplicial tree has a fixed point. Culler and Vogtmann [14] give a short proof based on their idea of “minipotent” elements. Bridson [5] proves a similar result for group actions on \mathbb{R} -trees using the triangle condition. We prove the following.

Theorem 1.6 *Let X be a unique arcwise connected space (eg. a tree or dendrite). Any action of $\text{Aut}(F_n)$ (or $\text{SL}_n(\mathbb{Z}), n \geq 3$) on X by homeomorphisms has a fixed point.*

This gives a simultaneous generalization of the fixed point property for both group actions on simplicial trees, \mathbb{R} -trees and those on dendrites.

2 Notations and basic facts

2.1 CAT(0) spaces

Let (X, d_X) be a geodesic metric space. For three points $x, y, z \in X$, the geodesic triangle $\Delta(x, y, z)$ consists of the three vertices x, y, z and the three geodesics $[x, y], [y, z]$ and $[z, x]$. Let \mathbb{R}^2 be the Euclidean plane with the standard distance $d_{\mathbb{R}^2}$ and $\bar{\Delta}$ a triangle in \mathbb{R}^2 with the same edge lengths as Δ . Denote by $\varphi : \Delta \rightarrow \bar{\Delta}$ the map sending each edge of Δ to the corresponding edge of $\bar{\Delta}$. The space X is called a CAT(0) space if for any triangle Δ and two elements $a, b \in \Delta$, we have the inequality

$$d_X(a, b) \leq d_{\mathbb{R}^2}(\varphi(a), \varphi(b)).$$

The typical examples of CAT(0) spaces include simplicial trees, hyperbolic spaces, products of CAT(0) spaces and so on. From now on, we assume that X is a complete CAT(0) space. Denote by $\text{Isom}(X)$ the isometry group of X . For any $g \in \text{Isom}(X)$, let

$$\text{Minset}(g) = \{x \in X : d(x, gx) \leq d(y, gy) \text{ for any } y \in X\}$$

and let $\tau(g) = \inf_{x \in X} d(x, gx)$ be the translation length of g . When the fixed-point set $\text{Fix}(g) \neq \emptyset$, we call g elliptic. When $\text{Minset}(g) \neq \emptyset$ and $d_X(x, gx) = \tau(g) > 0$ for any $x \in \text{Minset}(g)$, we call g hyperbolic. The group element g is called semisimple if the minimal set $\text{Minset}(g)$ is not empty, i.e. it is either elliptic or hyperbolic. A subset C of a CAT(0) space is convex, if any two points

$x, y \in C$ can be connected by the geodesic segment $[x, y] \subset C$. For more details on CAT(0) spaces, see the book of Bridson and Haefliger [10].

The following lemma is from [10] (II.2.4).

Lemma 2.1 *Let $\gamma : X \rightarrow X$ be an isometry of a CAT(0) space X and C an γ -invariant convex, complete subspace.*

(1) *For any $x \in X$, there exists a unique $p(x) \in C$ such that $d(x, p(x)) = d(x, C) := \inf_{c \in C} d(x, c)$. This gives a projection $p : X \rightarrow C$, which is distance-non-increasing.*

(2) *We have $p(\gamma x) = \gamma p(x)$ for any $x \in X$. Moreover, $\text{Min}(\gamma|_C) = \text{Min}(\gamma) \cap C$.*

Proof. The first part (1) is [10] (II.2.4). The second part is in the proof II.6.2(4) of [10]. Actually, we have that

$$d(\gamma x, \gamma p x) = d(x, p x) = d(x, C) = d(\gamma x, C)$$

and thus $\gamma p x = p(\gamma x)$ by the uniqueness of projection points. Therefore, $p \text{Min}(X) = \text{Min}(\gamma) \cap C = \text{Min}(\gamma|_C)$. ■

Lemma 2.2 *Let $G = A \rtimes H$ be a semi-direct product of two groups A and H . Suppose that both A and H have non-empty fixed point sets X^A and X^H . Then G has non-empty fixed point set X^G .*

Proof. For any $x \in X^A$ and any $h \in H, a \in A$, we have that

$$ahx = h(h^{-1}ah)x = hx.$$

This shows that $hX^A = X^A$. Moreover, the fixed point X^A is a convex closed complete subspace of X . Let $p : X \rightarrow X^A$ be the projection as defined in Lemma 2.1. Since $p(hx) = hp(x)$ for any $x \in X$, we have that $p(X^H) = X^H \cap X^A = X^G \neq \emptyset$. ■

2.2 Matrix groups

In this subsection, we briefly recall the definitions of the elementary subgroups $E_n(R)$ of the general linear group $\text{GL}_n(R)$. Let R be an associative ring with identity and $n \geq 2$ be an integer. The general linear group $\text{GL}_n(R)$ is the group of all $n \times n$ invertible matrices with entries in R . For an element $r \in R$ and any integers i, j such that $1 \leq i \neq j \leq n$, denote by $e_{ij}(r)$ the elementary $n \times n$ matrix with 1s in the diagonal positions and r in the (i, j) -th position and zeros elsewhere. The group $E_n(R)$ is generated by all such $e_{ij}(r)$, i.e.

$$E_n(R) = \langle e_{ij}(r) \mid 1 \leq i \neq j \leq n, r \in R \rangle.$$

Let $D = \{\text{diag}(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n) \mid \varepsilon_i = \pm 1\}$ be the diagonal subgroup. The extended elementary group $E'_n(R)$ is defined as $E_n(R)$, when n is odd and $\langle E_n(R), D \rangle < \text{GL}_n(R)$ when n is even. Note that

$$\begin{pmatrix} -1 & \\ & -1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}.$$

This implies that $\langle E_n(R), D \rangle$ contains $E_n(R)$ as an index-2 subgroup. When $R = \mathbb{Z}$ and n is even, we have $E'_n(\mathbb{Z}) = \mathrm{GL}_n(\mathbb{Z})$.

Denote by I_n the identity matrix and by $[a, b]$ the commutator $aba^{-1}b^{-1}$. The following lemma displays the commutator formulas for $E_n(R)$ (cf. Lemma 9.4 in [20]).

Lemma 2.3 *Let R be a ring and $r, s \in R$. Then for distinct integers i, j, k, l with $1 \leq i, j, k, l \leq n$, the following hold:*

- (1) $e_{ij}(r + s) = e_{ij}(r)e_{ij}(s)$;
- (2) $[e_{ij}(r), e_{jk}(s)] = e_{ik}(rs)$;
- (3) $[e_{ij}(r), e_{kl}(s)] = I_n$.

By Lemma 2.3, the group $E_n(R)$ ($n \geq 3$) is finitely generated when the ring R is finitely generated.

3 Basic facts on the relative property \mathcal{FA}_n

Definition 3.1 *Let H be a subgroup of a group G . The pair (G, H) has the relative property \mathcal{FA}_n (resp. $s\mathcal{FA}_n$) if whenever G acts isometrically (resp. semi-simply) on a complete n -dimensional CAT(0) space X , the subgroup H has a fixed point.*

We call that G has property \mathcal{FA}_n if (G, G) has the relative property \mathcal{FA}_n . The following are some basic properties.

Lemma 3.2 (1) *For groups $H < K < G$, if (K, H) has \mathcal{FA}_n (resp. $s\mathcal{FA}_n$), then so does (G, H) . If (G, K) has \mathcal{FA}_n (resp. $s\mathcal{FA}_n$), then so does (G, H) .*

(2) *Let $H_2 < H_1$ be two subgroups of G . If H_1 is bounded generated by finitely many conjugates of H_2 , then (G, H_1) has \mathcal{FA}_n (resp. $s\mathcal{FA}_n$) if and only if (G, H_2) has \mathcal{FA}_n (resp. $s\mathcal{FA}_n$).*

(3) *If H is of finite index in G , then (G, H) has \mathcal{FA}_n (resp. $s\mathcal{FA}_n$) if and only if G has \mathcal{FA}_n (resp. $s\mathcal{FA}_n$).*

(4) *If (G, H) has the relative property \mathcal{FA}_{n+1} (resp. $s\mathcal{FA}_n$), then (G, H) has the relative property \mathcal{FA}_n (resp. $s\mathcal{FA}_n$).*

Proof. (1) is obvious from the definition.

For (2), let H_1 be bounded generated by K_1, K_2, \dots, K_k (i.e. there exists $N > 0$ such that each element $g \in G$ is a product of at most N elements in $\cup_{i=1}^k K_i$), where each K_i is a conjugation of H_2 . When H_2 has a fixed point x , each K_i has a fixed point. Since each orbit $H_i x$ is bounded, the orbit $H_1 x$ is bounded and thus H_1 has a fixed point. When (G, H_1) has \mathcal{FA}_n , it follows (1) that (G, H_2) has \mathcal{FA}_n .

For (3), when H has a fixed point x , the orbit Gx is bounded and thus G also has a fixed point.

For (4), when G acts on a n -dimensional complete CAT(0) space X by isometries, the group G can also act on the product $X \times \mathbb{R}$ by trivial action on the second component. When H has a fixed point in $X \times \mathbb{R}$, it has also a fixed point in X . ■

Lemma 3.3 *Let H be a nilpotent group subgroup of a finitely generated group G . Suppose that each element in H is distorted in G . Then (G, H) has the relative property $s\mathcal{FA}_n$ for any n .*

Proof. When G acts on a complete CAT(0) space X by semisimple isometries, each element in H is either hyperbolic or elliptic. Suppose that some element $h \in H$ is hyperbolic and thus has a translation axis l . Fix a finite generating set $S < G$ and denote by l_S the word length function. For $x \in l$, the translation

$$\begin{aligned} |h| &= d(x, hx) = \lim_{n \rightarrow \infty} \frac{d(x, h^n x)}{n} \\ &\leq \lim_{n \rightarrow \infty} \frac{l_S(h^n) \text{Max}\{d(x, sx) : s \in S\}}{n} = 0, \end{aligned}$$

which is impossible. This shows that each element in H is elliptic. Since H is nilpotent, there is an upper central series

$$\{1\} = Z_0 \trianglelefteq Z_1 \trianglelefteq \cdots \trianglelefteq Z_n = H$$

such that each Z_{i+1}/Z_i is the center of G/Z_i . Note that Z_1 is abelian. Any finitely generated subgroup K of Z_1 has a fixed point by Lemma 2.2. Since the countable abelian group Z_1 is a direct limit of finitely generated subgroups, it also has a fixed point. Inductively suppose that Z_i has a fixed point. The quotient (abelian) group Z_{i+1}/Z_i acts invariantly on the fixed point set $\text{Fix}(Z_i)$ and also thus has a fixed point. This proves that Z_{i+1} and thus H has a fixed point. ■

Remark 3.4 *The previous lemma does not hold for \mathcal{FA}_n . For example, the subgroup $\langle e_{12}(1) \rangle \cong \mathbb{Z}$ generated by the $n \times n$ matrix with 1 in the (1,1)-th position and the diagonals, 0s elsewhere, is distorted in $\text{SL}_n(\mathbb{Z})$ ($n \geq 3$). But $\text{SL}_n(\mathbb{Z})$ acts properly on the symmetric space $\text{SL}_n(\mathbb{R})/\text{SO}(n)$.*

4 Helly's theorem and Coxeter groups

The classical Helly's theorem says that for m non-empty convex subsets $\{F_i\}_{i=1}^m$ ($m \geq n + 2$) in the Euclidean space \mathbb{R}^n satisfying the intersection of any $n + 1$ sets $\cap_{j=1}^n F_{i_j} \neq \emptyset$, the whole intersection $\cap_{i=1}^m F_i \neq \emptyset$. The following is the Helly theorem for CAT(0) spaces (cf. Bridson [7], Theorem 3.2; Varghese [24], Theorem 5.10.)

Lemma 4.1 *(the Helly theorem) Let X be a d -dimensional (i.e. covering dimension) complete CAT(0) space and S a finite family of non-empty closed convex subspaces. If the intersection of each $(d + 1)$ -elements of S is non-empty, then $\cap S$ is non-empty.*

Corollary 4.2 ([24], 5.1.) *Let G be a group, Y a finite generating set of G and X d -dimensional complete CAT(0) space. Let $\Phi : G \rightarrow \text{Isom}(X)$ be a homomorphism. If each $(d+1)$ -element subset of Y has a fixed point in X , then G has a fixed point in X .*

The following was firstly proved by Bridson [7] (Proposition 3.4):

Lemma 4.3 *Let X^d be a complete CAT(0) space and G a group acting on X by isometries. Suppose that there are integers k_1, k_2, \dots, k_m and subset $S_i, i = 1, 2, \dots, m$ satisfying*

- (1) *every k_i -element of S_i has a common fixed point;*
- (2) *$[s_i, s_j] = 1$ for any $s_i \in S_i, s_j \in S_j, i \neq j$;*
- (3) *$d < k_1 + k_2 + \dots + k_m$.*

Then for some i every finitely many elements in S_i have a common fixed point.

4.1 The action of Coxeter groups

A Coxeter group (W, S) is a group W with the presentation

$$\langle s_1, s_2, \dots, s_n \mid (s_i s_j)^{m_{ij}} = 1 \rangle$$

where $S = \{s_1, s_2, \dots, s_n\}$, $m_{ii} = 1$ and $m_{ij} \geq 2$ for $i \neq j$. It is possible that $m_{ij} = \infty$, which means no relations of the form $(s_i s_j)^\infty$. There is a canonical group homomorphism $\rho : W \rightarrow \mathbb{Z}/2$ defined by mapping each s_i to the generator $1 \in \mathbb{Z}/2$. The kernel is denoted by $SW = \ker \rho$, called the alternating Coxeter subgroup. For a subset $T \subseteq S$, the subgroup $W_T < W$ generated by elements in T is called a parabolic subgroup (or a special subgroup in the literature). The Coxeter matrix is the matrix $M = (m_{ij})_{n \times n}$ and the Schläfli matrix is $C = (-\cos(\frac{\pi}{m_{ij}}))_{n \times n}$. The Coxeter graph of (W, S) is a graph defined as the following. The vertices of the graph are labelled by generators in S . Vertices (s_i, s_j) are adjacent if and only if $m_{ij} \geq 3$. An edge is labelled with the value of m_{ij} whenever the value is ≥ 4 . The following is well-known (cf. [13]).

Lemma 4.4 *The Coxeter group is finite if and only the eigenvalues of the Schläfli matrix are all positive. The finite Coxeter groups consist of three one-parameter families of increasing rank A_n, B_n, D_n , one one-parameter family of dimension two $I_2(p)$, and six exceptional groups: E_6, E_7, E_8, F_4, H_3 and H_4 .*

Figure 1: Coxeter graphs of the finite Coxeter groups

Definition 4.5 Let $n > 0$ be an integer. A Coxeter group (W, S) is n -spherical if any $n + 1$ elements in S generates a finite group.

Lemma 4.6 Let X be a complete CAT(0) space of dimension $n \geq 1$. Let (W, S) be an n -spherical Coxeter group. Then any action of W on X has a fixed point.

Proof. By the assumption, the fixed point set X^s is closed for each generator $s \in S$. The intersection $\bigcap_{i=1}^k X^{s_i} = X^{W_{\{s_1, s_2, \dots, s_k\}}}$ is not empty for $k \leq n + 1$, since $W_{\{s_1, s_2, \dots, s_k\}}$ is finite. The Helly theorem (cf. Lemma 4.1) implies that the fixed point set X^W is not empty. ■

5 Actions of matrix groups on CAT(0) spaces

In this section, we always suppose that X is a complete CAT(0) space and $\text{Isom}(X)$ is the group of isometries. Let R be a finitely generated ring and $E_n(R)$ (resp. $E'_n(R)$) the (resp. extended) elementary subgroup. In this section, we will prove Theorem 1.1.

Lemma 5.1 Let $1 \leq j \neq i \leq n$. The elementary subgroup $E_n(R)$ has property \mathcal{FA}_{n-2} if and only if $(E_n(R), e_{ij}(x))$ has property \mathcal{FA}_{n-2} for any $x \in R$.

Proof. For any $y \in R$, the two matrices $e_{ij}(x)$ and $e_{ij}(y)$ commute with each other. Lemma 2.2 implies that they have a common fixed point. Therefore, the subgroup $e_{ij} := \langle e_{ij}(x) : x \in R \rangle$ has a fixed point. Since all e_{ij} ($i \neq j$) are conjugate, e_{12} has a fixed point. Apply Lemma 2.2 again to get that the subgroup $\langle e_{1i}(x) \mid 2 \leq i \leq n, x \in R \rangle$ has a fixed point and the upper triangular subgroup $H = \langle e_{ij}(x) \mid 1 \leq i < j \leq n, x \in R \rangle$ has a fixed point. Note that $E_n(R)$ is generated by $S = \{e_{12}, e_{23}, \dots, e_{n-1,n}, e_{n,1}\}$ and any $n - 1$ elements of S generate a subgroup isomorphic to H . The Helly theorem implies that $E_n(R)$ has a global fixed point. ■

Denote by $\{a_i\}_{i=1}^n$ the standard basis of the finitely generated free abelian group \mathbb{Z}^n , by $\{\sigma_i\}_{i=1}^n$ the standard basis of the elementary abelian group $(\mathbb{Z}/2)^n$. Let $G = \mathbb{Z}^n \rtimes ((\mathbb{Z}/2)^n \rtimes S_n)$ be the semi-direct product, where $\mathbb{Z}/2$ acts on \mathbb{Z} by reflection and S_n acts on $(\mathbb{Z}/2)^n$ and \mathbb{Z}^n by permuting the standard basis. Let $\rho : (\mathbb{Z}/2)^n \rtimes S_n \rightarrow \{+1, -1\}$ be the group homomorphism mapping each σ_i and each permutation (ij) to -1 . Denote by $A_n^+ = \ker \rho$ and $G^+ = \mathbb{Z}^n \rtimes A_n^+$.

Proposition 5.2 Let $n \geq 2$. Any isometric action of G on a complete CAT(0) space X^d ($d < n$) has a global fixed point. In other words, the group G has the property \mathcal{FA}_{n-1} .

Proof. Choose

$$S = \{a_1\sigma_1, (12), (23), \dots, (n-1, n), \sigma_n\}.$$

Note that each element of S is of order two. Let $H = (W, S)$ be the Coxeter group assigned to the Coxeter matrix (m_{ij}) defined by $m_{ii} = 2, m_{12} = 4, m_{i,i+1} = 3$ for $2 \leq i \leq n$ and $m_{n,n+1} = 4$ and all other entries $m_{ij} = \infty$. Recall that

$$H = \langle s_i \in S \mid (s_i s_j)^{m_{ij}} = 1, 1 \leq i, j \leq n \rangle.$$

Let $f : H \rightarrow \langle S \rangle$ be defined as

$$\begin{aligned} s_1 &\rightarrow a_1 \sigma_1, \\ s_{i+1} &\rightarrow (i, i+1), 1 \leq i \leq n-1, \\ s_{n+1} &\rightarrow \sigma_n. \end{aligned}$$

It could be directly checked that f is a group homomorphism. The Coxeter graph of (W, S) is a path consisting of $n+1$ vertices with the edges $(s_1, s_2), (s_n, s_{n+1})$ labeled by 4. The subgraph spanned by any n vertices is either a path of type B_n or disjoint union of two paths of type B_k . By the classification of finite Coxeter groups (cf. Lemma 4.4), any spherical subgroup of rank n is finite in H . This proves that the subgroup generated by any n elements is finite. The Helly theorem (cf. Lemma 4.1) implies that S has a common fixed point x . Note that $a_1 \in \langle S \rangle$ and thus $\mathbb{Z}^n < \langle S \rangle$. Therefore, the action of G on X has a bounded orbit Gx and thus has a global fixed point by Lemma 3.2 (3). ■

Note that $\mathbb{Z}^n \rtimes ((\mathbb{Z}/2)^n \rtimes S_n)$ could act on the Euclidean space \mathbb{R}^n in the standard way. This shows that the bound in the previous proposition is sharp.

Lemma 5.3 *Let $E_n(R)$ act isometrically on a complete CAT(0) space X^d ($d < n-1$). Assume n is odd. For any element $x \in R$, the subgroup $\langle e_{1i}(x) \mid i = 2, 3, \dots, n \rangle$ has a fixed point.*

Proof. When n is odd, let H be the subgroup generated by

$$\{-I_n(12), -I_n(23), \dots, -I_n(n-1, n), -I_n \sigma_n\}$$

and $\langle e_{1i}(x) \mid i = 2, 3, \dots, n \rangle$. Define a group homomorphism

$$\alpha : \mathbb{Z}^n \rtimes ((\mathbb{Z}/2)^n \rtimes S_n) \rightarrow H$$

by

$$a_i \mapsto e_{1,i+1}(x), \sigma_i \mapsto -I_n \varepsilon_i$$

and

$$(i, i+1) \mapsto -I_n(i, i+1).$$

Proposition 5.2 implies that H has a fixed point. ■

Proof of Theorem 1.3. Suppose that $R^n \rtimes E'_n(R)$ acts on a complete $n-1$ dimensional CAT(0) space X . When n is odd, we view $(\mathbb{Z}/2)^n \rtimes S_n$ as a

subgroup of $\mathrm{SL}_n(\mathbb{Z}) = E_n(\mathbb{Z})$ through the group homomorphism α in the proof of Lemma 5.3. The ring homomorphism $\mathbb{Z} \rightarrow R$ preserving the identity induces a group homomorphism $\mathrm{GL}_n(\mathbb{Z}) \rightarrow E'_n(R)$. Without confusions, we still denote the image by $(\mathbb{Z}/2)^n \rtimes S_n$. For any $x \in R$, let

$$\begin{aligned} S_x &= \{(x, \sigma_1), (12), (23), \dots, (n-1, n), \sigma_n\} \\ &\subseteq R^n \rtimes E'_n(R). \end{aligned}$$

The obvious map $S \rightarrow S_x$ by $\sigma_1 a_1 \rightarrow ((x, 0, \dots, 0), \sigma_1)$ induces a group homomorphism

$$\mathbb{Z}^n \rtimes ((\mathbb{Z}/2)^n \rtimes S_n) \rightarrow R^n \rtimes ((\mathbb{Z}/2)^n \rtimes S_n).$$

Proposition 5.2 implies that the subgroup xR^n (for any x) and thus R^n has a fixed point. ■

Proof of Theorem 1.1. When n is odd, we have $E'_n(R) = E_n(R)$. Theorem 1.1 follows Lemma 5.3 and Lemma 5.1. When n is even, the same group homomorphism α as in Lemma 5.3 and Proposition 5.2 imply that the subgroup $\langle e_{1i}(x) \mid i = 2, 3, \dots, n \rangle$ has a fixed point. Theorem 1.1 is proved by Lemma 1.3. ■

6 Actions of $\mathrm{Aut}(F_n)$ on $\mathrm{CAT}(0)$ spaces

Suppose that the free $F_n = \langle a_1, \dots, a_n \rangle$ and $\mathrm{Aut}(F_n)$ the group of automorphisms. For any $1 \leq i \leq n$, let $\varepsilon_i : F_n \rightarrow F_n$ be the involution defined by $a_i \rightarrow a_i^{-1}$ and $a_j \rightarrow a_j$ for any $j \neq i$. The Nielsen transformations are defined by

$$\rho_{ij} : a_i \rightarrow a_i a_j, a_k \rightarrow a_k, k \neq i$$

and

$$\lambda_{ij} : a_i \rightarrow a_j a_i, a_k \rightarrow a_k, k \neq i.$$

For $1 \leq i \neq j \leq n$, denote by (ij) the permutation of basis. It is well-known that $\mathrm{Aut}(F_n)$ is generated by the elements ρ_{ij} , λ_{ij} and ε_i (see Gersten [18]).

The following lemma is a variation on the argument that Gersten [19] used to show that $\mathrm{Aut}(F_n)$ cannot act properly and cocompactly on a $\mathrm{CAT}(0)$ space if $n \geq 3$. (cf. [10] p. 253).

Lemma 6.1 *Let $G = \langle \alpha, \beta, t \mid t^p \beta t^{-p} = \beta \alpha^p$ for any integer p ⟩. If $(G, \langle t \rangle)$ has the relative property \mathcal{FA}_n , then $(G, \langle \alpha \rangle)$ has \mathcal{FA}_n for any n .*

Proof. Suppose that G acts on an n -dimensional complete $\mathrm{CAT}(0)$ space X and t has a fixed point $x \in X$. For any p we have

$$\begin{aligned} d(\alpha^p x, x) &= d(\beta^{-1} t^p \beta t^{-p} x, x) \leq d(\beta^{-1} t^p \beta x, \beta^{-1} t^p x) + d(\beta^{-1} t^p x, x) \\ &\leq d(\beta x, x) + d(\beta^{-1} x, x). \end{aligned}$$

Therefore, the orbit $\{\alpha^p x : p \in \mathbb{Z}\}$ is bounded and thus α has a fixed point. ■

Theorem 6.2 *Let X^d be a complete CAT(0) space and $\text{Aut}(F_n)$ ($n \geq 4$) act on X by isometries. When $d < n - 2$, each Nielsen transformation of $\text{Aut}(F_n)$ has a fixed point.*

Proof. Let H be the subgroup generated by

$$S = \{\rho_{21}^{-1}\rho_{31}(23), (34), \dots, (n-1, n), (2, n)\}.$$

It can be directly checked that the product of any successive two elements (including that of the first element and the last one) is of order 3. Let H be the coxeter group assigned to the Coxeter graph K a loop of length $n - 1$. Explicitly, the group

$$H = \langle s_1, \dots, s_{n-1} \mid s_i^2 = (s_i s_{i+1})^3 = 1, 1 \leq i \leq n-1 \rangle,$$

where $s_n = s_1$. Let $f : H \rightarrow \langle S \rangle$ be defined as

$$\begin{aligned} s_1 &\rightarrow \rho_{21}^{-1}\rho_{31}(23), \\ s_i &\rightarrow (i+1, i+2), 2 \leq i \leq n-2, \\ s_{n-1} &\rightarrow (2, n). \end{aligned}$$

It can be directly checked that f is a group homomorphism. Since the subgraph spanned by $n - 2$ vertices in the Coxeter graph K is of type A , any $n - 2$ elements in S generate a finite group (cf. Lemma 4.4). By the Helley theorem, S has a common fixed point. Therefore, the element $\rho_{21}^{-1}\rho_{31} \in \langle S \rangle$ has a fixed point. Thus $\rho_{ij}^{-1}\rho_{i'j}$ has fixed point for any i, i', j after conjugating by a permutation.

Let $T = \rho_{32}^{-1}\rho_{42}$. It can be directly checked that $T^{-p}\rho_{13}T^p = \rho_{13}\rho_{12}^p$ for any integer p . Let $G = \langle \alpha, \beta, t \mid t^p\beta t^{-p} = \beta\alpha^p \text{ for any integer } p \rangle$. Define a group homomorphism

$$f : G \rightarrow \text{Aut}(F_n)$$

by $f(t) = T$, $f(\alpha) = \rho_{12}$, $f(\beta) = \rho_{13}$. Lemma 6.1 implies that ρ_{12} has a fixed point. Since any Nielsen transformation is conjugate to ρ_{12} , the statement is proved. ■

Theorem 1.4 is the same as the following theorem.

Theorem 6.3 *Let X^d be a complete CAT(0) space and $\text{Aut}(F_n)$ ($n \geq 4$) act on X by isometries. When $d < n - 2$, the subgroup $\text{Aut}(F_2)$ has a fixed point.*

Proof. It is already proved in Theorem 6.2 that each Nielsen transformation has a fixed point. Choose

$$S = \{\rho_{12}\varepsilon_2, \varepsilon_1, \varepsilon_2, \varepsilon_1\varepsilon_2(12)\}.$$

We check that every two elements on S has a common fixed in the following:

The elements $\rho_{12}\varepsilon_2$ and ε_1 lie in $\langle \rho_{12}, \lambda_{12} \rangle \rtimes \langle \varepsilon_2, \varepsilon_1 \rangle \cong \mathbb{Z}^2 \rtimes (\mathbb{Z}/2)^2$. Since ρ_{12} commutes with λ_{12} , they have a common fixed point. Therefore, the elements

$\rho_{12}\varepsilon_2$ and ε_1 have a common fixed point. The elements $\rho_{12}\varepsilon_2$ and ε_2 generate an infinite Dihedral group and thus have a common fixed point. The elements $\rho_{12}\varepsilon_2$ and $\varepsilon_1\varepsilon_2(12)$ generate a finite group and thus have a common fixed point.

Note that $\text{Aut}(F_n)$ contains $2\lfloor \frac{n}{2} \rfloor$ copies of $\text{Aut}(F_2)$ along the diagonal. The Bridson's lemma 4.3 implies that when $d < 2\lfloor \frac{n}{2} \rfloor$, the set S has a common fixed point. Since S generates $\text{Aut}(F_2)$, the proof is finished. ■

Lemma 6.4 *Let X^d be a complete CAT(0) space and $\text{Aut}(F_n)$ ($n \geq 5$) act on X by isometries. When $d < n - 3$, the subgroup $\text{Aut}(F_2) \times F_2$ has a fixed point.*

Proof. Let $T : F_n \rightarrow F_n$ be the transformation given by

$$x_4 \rightarrow x_4w^{-1}, x_5 \rightarrow x_5w,$$

for a word $w \in \langle x_1, x_3 \rangle < F_n$. Then $T^{-1}\rho_{24}T = \rho_{24}\rho_{2,w}$, where $\rho_{2,w}(x_2) = x_2w$ and $\rho_{2,w}(x_i) = x_i$ for any $i \neq 2$. Consider the set

$$S = \{T(45), (56), \dots, (n-1, n), (2, n), (4, n)\}.$$

A similar argument using Coxeter groups as in the proof of Theorem 6.2 shows that any $n-4$ elements of S generate a finite group. This implies that when $d < n-3$, the whole group has a fixed point by the Helly's theorem. Therefore, the element T has a fixed point. Note that for any integer p , we have $T^{-p}\rho_{24}T^p = \rho_{24}\rho_{2,w}^p$. Let $G = \langle \alpha, \beta, t \mid t^p\beta t^{-p} = \beta\alpha^p \text{ for any integer } p \rangle$. Define a group homomorphism

$$f : G \rightarrow \text{Aut}(F_n)$$

by $f(t) = T$, $f(\alpha) = \rho_{2,w}$, $f(\beta) = \rho_{24}$. Lemma 6.1 implies that $\rho_{2,w}$ has a fixed point. Since $(12)\rho_{2,w}(12) = \rho_{1,v}$, $v \in \langle x_2, x_3 \rangle$, we see that $F_2 = \langle \rho_{1v} : v \in \langle x_2, x_3 \rangle \rangle$ has a fixed point. Note that $\text{Aut}(F_2)$ normalizes F_2 . Theorem 1.4 shows that $\text{Aut}(F_2)$ has a fixed. The semi-direct product $\text{Aut}(F_2) \times F_2$ has a fixed point by Lemma 2.2. ■

Proof of Corollary 1.5. Let

$$S = \{\varepsilon_2\rho_{12}, \varepsilon_1(23), \varepsilon_1\varepsilon_2(12), (i, i+1), i = 3, \dots, n-1, \varepsilon_n\},$$

$$S_1 = \{\varepsilon_2\rho_{12}, \varepsilon_3, \varepsilon_1(23), \varepsilon_1\varepsilon_2(12)\}.$$

It is direct (see [24]) to check that any 2 elements in S or S_1 generate a finite group and thus have a common fixed point. Note that $\text{Aut}(F_n)$ contains $\lfloor \frac{n}{3} \rfloor$ copies of $\text{Aut}(F_3)$ along the diagonal. Since S_1 lies in $\text{Aut}(F_3)$, we know that $\text{Aut}(F_n)$ contains $\lfloor \frac{n}{3} \rfloor$ copies of S_1 , any two of which commute. Bridson's result (cf. Lemma 4.3) implies that when $d < 2\lfloor \frac{n}{3} \rfloor$, the set S_1 has a global fixed point.

We will prove that any $(d+1)$ -element subset of S has a common fixed point. Inductively, assume that it is already proved for any $(k-1)$ -element subset ($k \geq 4$). Let

$$S_k = \{\varepsilon_2\rho_{12}, \varepsilon_3, \varepsilon_1(23), \varepsilon_1\varepsilon_2(12), (i, i+1), i = 3, \dots, k-1\}.$$

Note that any k -element subset of S generates a finite group, except possibly S_k . Note also that S_k is subset of $\text{Aut}(F_k)$ and $\text{Aut}(F_n)$ contains $\lfloor \frac{n}{k} \rfloor$ copies of $\text{Aut}(F_k)$ and thus of S_k . Lemma 4.3 implies that when $d < (k-1)\lfloor \frac{n}{k} \rfloor$ any finitely many elements of S_k has a fixed point. Note that

$$d < 2\lfloor \frac{n}{3} \rfloor \leq (k-1)\lfloor \frac{n}{k} \rfloor$$

for any $k \geq 4$. Therefore, any k -element subset S_k has a fixed point. The Helly theorem (cf. Lemma 4.1) implies that $\langle S \rangle = \text{Aut}(F_n)$ has a fixed point. ■

7 Generalizations

In this section, we generalize the theorems proved in previous sections to a more general setting. For this purpose, we give the following definition.

Definition 7.1 *Let X be a topological space and G be a subgroup of its homeomorphism group $\text{Homeo}(X)$. For an integer $n > 0$, the pair (transformation group) (X, G) is n -Helly good if*

- (i) *any finite subgroup $H < G$ has a non-empty fixed point set $\text{Fix}(H)$; and*
- (ii) *any collection of finitely many finite subgroups $\{H_i\}_{i \in I}$ ($H_i < G$) has a non-empty intersection $\cap \text{Fix}(H_i)$, whenever $n+1$ of them has a nonempty intersection $\cap_{j=1}^{n+1} \text{Fix}(H_{i_j})$.*

In such a case, we call X n -Helly good with respect to the group G .

Theorem 7.2 *Let (X, G) be an m -Helly good pair. Suppose that $n > m$ and R is a ring. When the semi-direct product $R^n \rtimes E'_n(R)$ acts on X by homeomorphisms in G , the abelian subgroup $\{(x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{Z}x \subset R\}$ has a fixed point for any $x \in R$. In particular, any elementary matrix $e_{ij}(x)$ has a fixed point when $E'_{n+1}(R)$ acts on X by homeomorphisms in G , for any $x \in R$ and $1 \leq i \neq j \leq n$.*

Proof. The proof is similar to that of Proposition 5.2. For any $x \in R$, let

$$\begin{aligned} S_x &= \{(x, \sigma_1), (12), (23), \dots, (n-1, n), \sigma_n\} \\ &\subseteq R^n \rtimes E'_n(R). \end{aligned}$$

Let $H = (W, S)$ be the coxeter group assigned to the Coxeter matrix (m_{ij}) defined by $m_{ii} = 2, m_{12} = 4, m_{i, i+1} = 3$ for $2 \leq i \leq n$ and $m_{n, n+1} = 4$ and all other entries $m_{ij} = \infty$. Recall that

$$H = \langle s_i \in S \mid (s_i s_j)^{m_{ij}} = 1, 1 \leq i, j \leq n \rangle.$$

Let $f : H \rightarrow \langle S_x \rangle$ be defined as

$$\begin{aligned} s_1 &\rightarrow ((x, 0, \dots, 0), \sigma_1), \\ s_{i+1} &\rightarrow (i, i+1), 1 \leq i \leq n-1, \\ s_{n+1} &\rightarrow \sigma_n. \end{aligned}$$

It could be directly checked that f is a group homomorphism. The Coxeter graph of (W, S) is a path consisting of $n + 1$ vertices with the edges $(s_1, s_2), (s_n, s_{n+1})$ labeled by 4. The subgraph spanned by any n vertices is either a path of type B_n or disjoint union of two paths of type B_k . By the classification of finite Coxeter groups (cf. Lemma 4.4), any spherical subgroup of rank n is finite in H . This proves that the subgroup generated by any n elements is finite. The definition of m -Helly good spaces implies that S has a common fixed point $x_0 \in X$. Note that $\{(x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{Z}x \subset R\} < \langle S \rangle$.

We view $R^n = \langle e_{12}(x_1), e_{13}(x_2), \dots, e_{1n}(x_{n-1}) \mid x_i \in R \rangle < E'_{n+1}(R)$. Therefore, $e_{12}(x)$ has a fixed point. Since any elementary matrix $e_{ij}(x)$ is conjugate to $e_{12}(x)$, the last claim is proved. ■

Bridson [5] obtains ‘The Triangle Criterion’ for group actions on \mathbb{R} -trees: If Γ is generated by $A_1 \cup A_2 \cup A_3$ and $H_{ij} = \langle A_i, A_j \rangle$ is finite for $i, j = 1, 2, 3$, then Γ has property $F\mathbb{R}$ (i.e., any isometric action of Γ on an \mathbb{R} -tree has a fixed point). In particular, the automorphism group of free group $\text{Aut}(F_n)$ and the special linear group $\text{SL}_n(\mathbb{Z}), n \geq 3$, satisfy the triangle conditions and thus have property $F\mathbb{R}$. We prove a similar result for 1-Helly good transformation group (X, G) :

Theorem 7.3 *Let (X, G) be a 1-Helly good pair. If Γ is generated by $A_1 \cup A_2 \cup A_3$ and $H_{ij} = \langle A_i, A_j \rangle$ is finite for $i, j = 1, 2, 3$, then any G -action of Γ on X has a fixed point. In other words, any group homomorphism $f : \Gamma \rightarrow X$ has a fixed point. In particular, any G -action of $\text{Aut}(F_n)$ (or $\text{SL}_n(\mathbb{Z}), n \geq 3$) on X has a fixed point.*

Proof. Since each H_{ij} is finite, it has a fixed point by condition (i) in the definition of n -Helly good pair. The condition (ii) implies the existence of a global fixed point. ■

Now we discuss some examples, which are n -Helly good.

Example 7.4 *A CAT(0) space X of dimension n is n -Helly good, with respect to the isometry group $G = \text{Isom}(X)$.*

Proof. This follows easily the Helly’s theorem (see Lemma 4.1) and the Bruhat-Tits fixed point theorem (see [10], Corollary 2.8, p.179). ■

Recall that a dendrite is a connected compact metrizable space X satisfying one of the following equivalent conditions:

- (1) Any two distinct points of X can be separated by a point.
- (2) X is locally connected and contains no simple closed curve.
- (3) The intersection of any two connected subsets of X remains connected.
- (4) X is a one-dimensional absolute retract.
- (5) $C(X)$ is projective in the category of unital C^* -algebras.
- (6) Any two points are extremities of a unique arc (i.e. a homeomorphic image of a compact interval in the real line) in X .

For more details, see [11][12][21]. We need the following result.

Lemma 7.5 *Let X be a dendrite. Let G be a finite group. Then any action of G on X has a fixed point.*

Proof. When G is finite, this is well-known (for example, see [15], Corollary 5.1). ■

A dendrite X is actually 1-Helly good, with respect to the homeomorphism group $\text{Homeo}(X)$. This is a special case of the following result. A uniquely arcwise connected space X is a hausdorff topological space in which every pair of distinct points x, y are joined by a unique arc $[x, y]$, which is a subset homeomorphic to a closed real interval. For more information, see Bowditch [3].

Lemma 7.6 *Let X be a uniquely arcwise connected space. Then X is 1-Helly good, with respect to the homeomorphism group $\text{Homeo}(X)$.*

Proof. Let $H < \text{Homeo}(X)$ be a finite subgroup. Fix $x \in X$. Since the orbit Hx is finite and there is a unique arc $[x, hx]$ connecting x and hx for each $h \in H$, the union $\cup_{h \in H} [x, hx]$ is a compact tree and thus a dendrite (cf. [3], Lemma 1.2). Note that the union is invariant under the action of H . Lemma 7.5 says that any action of H has a fixed point. Therefore, the action of H on X has a non-empty fixed point set $\text{Fix}(H)$. This proves the condition (i).

For the condition (ii), let H_1, H_2, \dots, H_n be a collection of finitely many finite subgroups. For any two points $x, y \in \text{Fix}(H_i)$, the unique arc $[x, y]$ is invariant under the action of H_i . Since a finite group cannot act freely on the real line \mathbb{R} , every point in $[x, y]$ is a fixed point of H_i . This proves that the fixed point set $\text{Fix}(H_i)$ is path connected. For each $i, j = 1, 2, \dots, n$, choose $x_{i,j} \in \text{Fix}(H_i) \cap \text{Fix}(H_j)$. Denote by $A = \{x_{i,j} : i, j = 1, 2, \dots, n\}$ and $Y = \cup_{x,y \in A} [x, y]$, which is a compact tree. The result follows from Helly's theorem for trees where connectedness coincides with convexity. ■

Theorem 1.6 is a corollary of Lemma 7.6 and Theorem 7.3.

References

- [1] A. Barnhill, The FA_n conjecture for Coxeter groups, *Algebraic & Geometric topology* **6** (2006), 2117-2150.
- [2] O V Bogopol'skii, Arboreal decomposability of groups of automorphisms of free groups, *Algebra i Logika* **26** (1987) 131–149.
- [3] B H Bowditch, Treelike structures arising from continua and convergence groups, *Mem. Amer. Math. Soc.* **139** (1999) viii+86.
- [4] M. R. Bridson, Problems concerning hyperbolic and $\text{CAT}(0)$ groups (2007), available at: <https://docs.google.com/file/d/0B-tup63120-GVVZqNFITcEJmMmc/edit>.

- [5] M.R. Bridson, A condition that prevents groups from acting nontrivially on trees, *Geometry & Topology Monographs* 14 (2008) 129–133.
- [6] M.R. Bridson, Semisimple actions of mapping class groups on CAT(0) spaces, “*The Geometry of Riemann Surfaces*” (F. P. Gardiner, G. Gonzalez-Diez and C. Kourouniotis, eds.), LMS Lecture Notes 368, Cambridge Univ. Press, Cambridge, 2010, pp. 1–14.
- [7] M.R. Bridson, On the dimension of CAT(0) spaces where mapping class groups act, *Journal für reine und angewandte Mathematik*, **673** (2012), 55-68.
- [8] M.R. Bridson, Helly’s theorem, CAT(0) spaces, and actions of automorphism groups of free groups, preprint.
- [9] M.R. Bridson, The rhombic dodecahedron and semisimple actions of $\text{Aut}(F_n)$ on CAT(0) spaces, *Fundamenta mathematicae*, **214** (2011), 13-25.
- [10] M.R. Bridson and A. Haefliger, *Metric spaces of nonpositive curvature*, Grundlehren der Math. Wiss. 319, Springer-Verlag, Berlin, 1999.
- [11] J. J. Charatonik & W. J. Charatonik, “Dendrites”, in XXX National Congress of the Mexican Mathematical Society (Aguascalientes, 1997), *Aportaciones Matemáticas. Comunicaciones*, vol. 22, Sociedad Matemática Mexicana, 1998, p. 227-253.
- [12] A. Chigogidze & A. N. Dranishnikov, “Which compacta are noncommutative ARs?”, *Topology Appl.* 157 (2010), no. 4, p. 774-778.
- [13] H. Coxeter, The complete enumeration of finite groups of the form $r_i^2 = (r_i r_j)^{k_{ij}} = 1$, *J. London Math. Soc.*, 1, 10 (1935): 21–25.
- [14] M Culler, K Vogtmann, A group-theoretic criterion for property FA, *Proc. Amer. Math. Soc.* 124 (1996) 677–683.
- [15] B. Duchesne and N. Monod, Group actions on Dendrites and curves, *Ann. Inst. Fourier, Grenoble* 68, 5 (2018) 2277-2309.
- [16] B. Farb, Group actions and Helly’s theorem, *Advances in Mathematics* **222** (2009), 1574-1588.
- [17] M. Fukunaga, Fixed points of elementary subgroups of Chevalley groups acting on trees, *Tsukuba J. Math.*, **3** (1979), 7-16.
- [18] S.M. Gersten, A presentation for the special automorphism group of a free group, *J. Pure Appl. Algebra*, 33 (1984) 269-279.
- [19] S.M. Gersten, The automorphism group of a free group is not a CAT(0) group, *Proc. Amer. Math. Soc.* 121 (1994), 999-1002.

- [20] B.A. Magurn. *An algebraic introduction to K-theory*, Cambridge University Press, 2002.
- [21] S. B. Nadler, Jr., *Continuum theory. An introduction*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 158, John Wiley & Sons, 1992, xiv+328 pages.
- [22] J-P Serre, *Trees*, Springer Monographs in Mathematics, Springer, Berlin (2003). Translated from the French original by John Stillwell, Corrected 2nd printing of the 1980 English translation.
- [23] S. Ye, *Low-dimensional representations of matrix groups and group actions on CAT(0) spaces and manifolds*, J. Algebra 409 (2014), 219-243.
- [24] O. Varghese, *Fixed points for actions of $\text{Aut}(F_n)$ on CAT(0) spaces*. Münster Journal of Mathematics, vol. 7, (2014), 439-462.

Department of Mathematical Sciences, Xi'an Jiaotong-Liverpool University,
111 Ren Ai Road, Suzhou, Jiangsu 215123, China.
E-mail: Shengkui.Ye@xjtlu.edu.cn