

Breather wave and lump-type solutions of new (3+1)-dimensional Boiti-Leon-Manna-Pempinelli equation in incompressible fluid

Jian-Guo Liu*, Abdul-Majid Wazwaz

Received: date / Accepted: date

Abstract Under investigation is a new (3+1)-dimensional Boiti-Leon-Manna-Pempinelli equation. The main results are listed as follows: (i) lump solutions; (ii) Interaction solutions between lump wave and solitary waves; (iii) Interaction solutions between lump wave and periodic waves; (iv) Breather wave solutions. Furthermore, graphical representation of all solutions is studied and shown in some 3D- and contour plots.

Keywords lump solutions, breather wave solutions, solitary waves, periodic waves.

Mathematics Subject Classification (2000) 35C08 · 68M07 · 33F10

1 Introduction

The (3+1)-dimensional Boiti-Leon-Manna-Pempinelli (BLMP) equation can be used to describe the wave propagation in incompressible fluid [1] and the interaction of the Riemann wave [2]. Many results about BLMP equation have been obtained. Darvishi [2] presented the stair and step Soliton Solutions. Zuo [3] derived the bilinear form, Lax pairs and Bäcklund transformations. Tang [4] obtained the new periodic-wave solutions. Liu [5,6] given the new three-wave solutions and new non-traveling wave solutions. Mabrouk [7] obtained some new analytical solutions. Li [8] presented the multiple-lump waves solutions.

Jian-Guo Liu(*Corresponding author)
College of Computer, Jiangxi University of Traditional Chinese Medicine, Jiangxi 330004,
China Tel.: +8613970042436
Fax: +86079187119019
E-mail: 20101059@jxutcm.edu.cn

Abdul-Majid Wazwaz
Department of Mathematics, Saint Xavier University, Chicago, IL 60655, USA
E-mail: wazwaz@sxu.edu

Osman [9] derived lump waves, breather waves, mixed waves, and multi-soliton wave solutions. Peng [10] studied the breather waves and rational solutions. Xu [11] given the Painlevé analysis, lump-kink solutions and localized excitation solutions.

In this paper, a new (3+1)-dimensional BLMP equation is studied as follows [12]

$$(u_x + u_y + u_z)_t + (u_x + u_y + u_z)_{xxx} + (u_x(u_x + u_y + u_z))_x = 0, \quad (1)$$

where $u = u(x, y, z, t)$. The new BLMP equation was first proposed by Wazwaz [12]. The integrability, compatibility conditions, multiple soliton solutions and multiple complex soliton solutions were discussed by Painlevé test and Hirota's direct method.

Based on the transformation

$$u = -2 [\ln \xi(x, y, z, t)]_x, \quad (2)$$

the new BLMP equation has the bilinear form

$$[D_x^4 + D_y D_x^3 + D_z D_x^3 + D_t D_x + D_t D_y + D_t D_z] \xi \cdot \xi = 0. \quad (3)$$

The organization of this paper is as follows. Section 2 gives the lump solutions; Section 3 obtains the interaction solutions between lump wave and solitary waves; Section 4 derives the interaction solutions between lump wave and periodic waves; Section 5 presents the breather wave solutions. All results have been verified to be correct by using Mathematica software [13-25]. Section 6 gives the conclusion.

2 Lump solutions

Generally speaking, the lump solutions of nonlinear integrable equations can be assumed as follows

$$\xi = \gamma_3 + (\alpha_4 t + \alpha_1 x + \alpha_2 y + \alpha_3 z)^2 + (\beta_4 t + \beta_1 x + \beta_2 y + \beta_3 z)^2, \quad (4)$$

where $\alpha_i (1 \leq i \leq 4)$, $\beta_i (1 \leq i \leq 4)$ and γ_3 are undetermined real parameters. Substituting Eq. (4) into Eq. (3) and making the coefficients of x^2 , y^2 , xy , xz et al. be zero, undetermined real parameters in Eq. (4) have the following results

$$(I) \quad \alpha_3 = -\alpha_1 - \alpha_2, \beta_2 = -\beta_1 - \beta_3. \quad (5)$$

$$(II) \quad \beta_1 = -\frac{\alpha_1 \alpha_4}{\beta_4}, \alpha_3 = \frac{\alpha_1 \beta_3}{\beta_1}, \beta_2 = -\frac{\alpha_2 \beta_3}{\alpha_1 + \alpha_2}, \beta_4 = \frac{(\alpha_1 + \alpha_2) \alpha_4}{\beta_3}. \quad (6)$$

$$(III) \quad \alpha_3 = \frac{\alpha_1 \beta_3}{\beta_1}, \beta_1 = -\frac{\alpha_1 \beta_3}{\alpha_1 + \alpha_2}, \beta_2 = -\frac{\alpha_2 \beta_3}{\alpha_1 + \alpha_2}. \quad (7)$$

$$(IV) \quad \alpha_3 = -\frac{\beta_1 \beta_3}{\alpha_1}, \beta_3 = \frac{\alpha_1^2 + \alpha_2 \alpha_1}{\beta_1}, \beta_2 = -\frac{\alpha_1 (\alpha_1 + \alpha_2) + \beta_1^2}{\beta_1}. \quad (8)$$

$$(V) \quad \alpha_2 = -\alpha_1, \beta_2 = -\beta_1, \alpha_3 = \beta_3 = 0. \quad (9)$$

$$(VI) \quad \alpha_4 = \beta_4 = 0, \beta_3 = -\frac{\alpha_1(\alpha_1 + \alpha_2 + \alpha_3) + \beta_1(\beta_1 + \beta_2)}{\beta_1}. \quad (10)$$

$$(VII) \quad \alpha_3 = -\alpha_1 - \alpha_2, \beta_1 = -\beta_2 - \beta_3, \beta_4 = \frac{2\alpha_1\alpha_4\beta_1}{\alpha_1^2 - \beta_1^2}. \quad (11)$$

$$(VIII) \quad \alpha_3 = -\alpha_1 - \alpha_2, \beta_1 = \frac{\alpha_1\beta_2}{\alpha_2},$$

$$\beta_3 = -\frac{(\alpha_1 + \alpha_2)\beta_2}{\alpha_2}, \beta_4 = \frac{2\alpha_1\alpha_4\beta_1}{\alpha_1^2 - \beta_1^2}. \quad (12)$$

Substituting Eqs. (5)-(12) into Eq. (2) and Eq. (4), eight different types of lump solutions are derived. As an example, substituting Eq. (5) into Eq. (2) and Eq. (4), we have

$$u = -[2[2\alpha_1[\alpha_4 t + \alpha_1 x + \alpha_2 y - (\alpha_1 + \alpha_2)z] + 2\beta_1[\beta_4 t + \beta_1 x - (\beta_1 + \beta_3)y + \beta_3 z]]/[\gamma_3 + [\alpha_4 t + \alpha_1 x + \alpha_2 y - (\alpha_1 + \alpha_2)z]^2 + [\beta_4 t + \beta_1 x - (\beta_1 + \beta_3)y + \beta_3 z]^2]. \quad (13)$$

Graphical representation of lump solution (13) is shown in Fig. 1. It's obvious that there is a peak and a bottom in Fig. 1, and they are symmetric. In the peak, solution (13) has a maximum $2\sqrt{5}$ at $x = -1/\sqrt{5}$ and $y = 0$. In the bottom, solution (13) has a minimum $-2\sqrt{5}$ at $x = 1/\sqrt{5}$ and $y = 0$.




Fig. 1. Lump solution (13) with $\alpha_1 = \beta_3 = -2$, $\beta_4 = -3$, $\alpha_4 = 2$, $t = z = 0$, $\alpha_2 = \beta_1 = \gamma_3 = 1$.

3 Interaction solutions between lump wave and solitary waves

In this section, we will discuss the interaction phenomenon between lump wave and solitary waves. Considering the following mixed functions

$$\xi = \gamma_3 + (\alpha_4 t + \alpha_1 x + \alpha_2 y + \alpha_3 z)^2 + (\beta_4 t + \beta_1 x + \beta_2 y + \beta_3 z)^2$$

$$+ \gamma_1 e^{t\varphi_4 + \varphi_1 x + \varphi_2 y + \varphi_3 z} + \gamma_2 e^{-t\varphi_4 - \varphi_1 x - \varphi_2 y - \varphi_3 z}, \quad (14)$$

where $\varphi_i (1 \leq i \leq 4)$ and $\gamma_i (i = 1, 2)$ are undetermined real parameters. As an example, substituting Eq. (14) and Eq. (5) into Eq. (3) and making the coefficients of $e^{t\varphi_4 + \varphi_1 x + \varphi_2 y + \varphi_3 z} x^2$, $e^{t\varphi_4 + \varphi_1 x + \varphi_2 y + \varphi_3 z} y^2$ et al. be zero, undetermined real parameters in Eq. (14) have the following results

$$\alpha_3 = -\alpha_1 - \alpha_2, \beta_2 = -\beta_1 - \beta_3, \varphi_3 = -\varphi_1 - \varphi_2. \quad (15)$$

Substituting Eq. (15) into Eq. (2) and Eq. (14), the interaction solutions between lump wave and solitary waves can be read as

$$\begin{aligned} u = & -[2[-\gamma_2 \varphi_1 \exp[-t\varphi_4 - \varphi_1 x - \varphi_2 y + (\varphi_1 + \varphi_2)z] + 2\alpha_1[\alpha_4 t + \alpha_1 x \\ & + \alpha_2 y - (\alpha_1 + \alpha_2)z] + 2\beta_1[\beta_4 t + \beta_1 x - (\beta_1 + \beta_3)y + \beta_3 z] \\ & + \gamma_1 \varphi_1 e^{t\varphi_4 + \varphi_1 x + \varphi_2 y - (\varphi_1 + \varphi_2)z}]/[\gamma_3 + \gamma_2 \exp[-t\varphi_4 - \varphi_1 x - \varphi_2 y \\ & + (\varphi_1 + \varphi_2)z] + [\alpha_4 t + \alpha_1 x + \alpha_2 y - (\alpha_1 + \alpha_2)z]^2 + [\beta_4 t + \beta_1 x \\ & - (\beta_1 + \beta_3)y + \beta_3 z]^2 + \gamma_1 e^{t\varphi_4 + \varphi_1 x + \varphi_2 y - (\varphi_1 + \varphi_2)z}]. \end{aligned} \quad (16)$$

Graphical representation of the interaction solutions (16) is shown in Fig. 2, Fig. 3 and Fig. 4. Fig. 2 describes the interaction phenomenon between lump wave and one solitary wave. Fig. 3 and Fig. 4 represent the interaction phenomenon between lump wave and two solitary waves.




Fig. 2. Solution (16) with $\alpha_1 = \beta_3 = \gamma_1 = -2$, $\beta_4 = \varphi_1 = -3$, $\alpha_4 = \varphi_2 = 2$, $\gamma_2 = z = 0$, $\alpha_2 = \beta_1 = \gamma_3 = \varphi_4 = 1$, when $t = -3$ in (a,d), $t = 0$ in (b,e), $t = 3$ in (c,f).

4 Interaction solutions between lump wave and periodic waves

In this section, we will investigate the interaction phenomenon between lump wave and periodic waves. Assuming the following mixed functions

$$\xi = \gamma_3 + (\alpha_4 t + \alpha_1 x + \alpha_2 y + \alpha_3 z)^2 + (\beta_4 t + \beta_1 x + \beta_2 y + \beta_3 z)^2 + \gamma_2 \sin(\mu_4 t + \mu_1 x + \mu_2 y + \mu_3 z) + \gamma_1 \cos(t\varphi_4 + \varphi_1 x + \varphi_2 y + \varphi_3 z), \quad (17)$$

where $\mu_i (1 \leq i \leq 4)$ is undetermined real parameter.




Fig. 3. Solution (16) with $\alpha_1 = \beta_3 = \gamma_1 = \gamma_2 = -2$, $\beta_4 = \varphi_1 = -3$, $z = 0$, $\alpha_4 = \varphi_2 = 2$, $\alpha_2 = \beta_1 = \gamma_3 = \varphi_4 = 1$, when $t = -4$ in (a), $t = -2$ in (b), $t = 0$ in (c), $t = 2$ in (d), $t = 4$ in (e).




Fig. 4. The corresponding contour plots of Fig. 3.

Substituting Eq. (17) and Eq. (5) into Eq. (3) and making the coefficients of $\cos(t\varphi_4 + \varphi_1x + \varphi_2y + \varphi_3z)x^2$, $\sin(\mu_4t + \mu_1x + \mu_2y + \mu_3z)x^2$ et al. be zero, undetermined real parameters in Eq. (17) have the following results

$$\alpha_3 = -\alpha_1 - \alpha_2, \beta_2 = -\beta_1 - \beta_3, \varphi_3 = -\varphi_1 - \varphi_2, \mu_3 = -\mu_1 - \mu_2. \quad (18)$$

Substituting Eq. (18) into Eq. (2) and Eq. (17), the interaction solutions between lump wave and periodic waves can be written as

$$\begin{aligned} u = & -[2\alpha_1[\alpha_4t + \alpha_1x + \alpha_2y - (\alpha_1 + \alpha_2)z] + 2\beta_1[\beta_4t + \beta_1x - (\beta_1 + \beta_3)y \\ & + \beta_3z] + \gamma_2\mu_1 \cos[\mu_4t + \mu_1x + \mu_2y - (\mu_1 + \mu_2)z] - \gamma_1\varphi_1 \sin[t\varphi_4 + \varphi_1x \\ & + \varphi_2y - (\varphi_1 + \varphi_2)z]] / [\gamma_3 + [\alpha_4t + \alpha_1x + \alpha_2y - (\alpha_1 + \alpha_2)z]^2 + [\beta_4t \\ & + \beta_1x - (\beta_1 + \beta_3)y + \beta_3z]^2 + \gamma_2 \sin[\mu_4t + \mu_1x + \mu_2y - (\mu_1 + \mu_2)z] \\ & + \gamma_1 \cos[t\varphi_4 + \varphi_1x + \varphi_2y - (\varphi_1 + \varphi_2)z]]. \end{aligned} \quad (19)$$



**Fig. 5.** Solution (19) with $\alpha_1 = \beta_3 = \gamma_1 = -2$, $\beta_4 = \varphi_1 = -3$, $z = t = \gamma_2 = 0$, $\alpha_4 = \varphi_2 = \mu_1 = \mu_4 = 2$, $\alpha_2 = \beta_1 = \gamma_3 = \varphi_4 = 1$, $\mu_2 = -1$.

Fig. 6. Solution (19) with $\alpha_1 = \beta_3 = \gamma_1 = \gamma_2 = -2$, $\beta_4 = -3$, $z = t = 0$, $\alpha_4 = \mu_1 = \mu_4 = \varphi_1 = \varphi_4 = 2$, $\alpha_2 = \beta_1 = \gamma_3 = 1$, $\mu_2 = \varphi_2 = -1$.

Graphical representation of the interaction solutions (19) is shown in Fig. 5 and Fig. 6. Fig. 5 describes the interaction phenomenon between lump wave and one periodic wave. Fig. 6 represents the interaction phenomenon between lump wave and two periodic waves.

5 Breather wave solutions

In this section, we will study the breather wave solutions. Choosing the following mixed functions

$$\xi = k_1 e^{t\varphi_4 + \varphi_1 x + \varphi_2 y + \varphi_3 z} + \gamma_2 \sin(\mu_4 t + \mu_1 x + \mu_2 y + \mu_3 z) + \gamma_1 \cos(\nu_4 t + \nu_1 x + \nu_2 y + \nu_3 z) + e^{-t\varphi_4 - \varphi_1 x - \varphi_2 y - \varphi_3 z}, \quad (20)$$

where $\nu_i (1 \leq i \leq 6)$ and k_1 are unknown real parameters. Substituting Eq. (20) into Eq. (3), we have the following results

$$\nu_3 = -\nu_1 - \nu_2, \varphi_3 = -\varphi_1 - \varphi_2, \mu_3 = -\mu_1 - \mu_2. \quad (21)$$

Substituting Eq. (21) into Eq. (2) and Eq. (20), the breather wave solutions can be presented as

$$u = -[2[-\varphi_1 \exp[-t\varphi_4 - \varphi_1 x - \varphi_2 y + (\varphi_1 + \varphi_2)z] + \gamma_2 \mu_1 \cos[\mu_4 t + \mu_1 x + \mu_2 y - (\mu_1 + \mu_2)z] + k_1 \varphi_1 e^{t\varphi_4 + \varphi_1 x + \varphi_2 y - (\varphi_1 + \varphi_2)z} - \gamma_1 \nu_1 \sin[\nu_4 t + \nu_1 x + \nu_2 y - (\nu_1 + \nu_2)z]] / [\exp[-t\varphi_4 - \varphi_1 x - \varphi_2 y + (\varphi_1 + \varphi_2)z] + k_1 e^{t\varphi_4 + \varphi_1 x + \varphi_2 y - (\varphi_1 + \varphi_2)z} + \gamma_2 \sin[\mu_4 t + \mu_1 x + \mu_2 y - (\mu_1 + \mu_2)z] + \gamma_1 \cos[\nu_4 t + \nu_1 x + \nu_2 y - (\nu_1 + \nu_2)z]]. \quad (22)$$

Graphical representation of the breather wave solutions (22) is displayed in Fig. 7.




Fig. 7. Breather wave solutions (22) with $\gamma_1 = -2$, $\varphi_1 = \nu_2 = -3$, $z = t = 0$, $\gamma_2 = \mu_1 = \mu_4 = \varphi_2 = \nu_1 = 2$, $\nu_4 = \varphi_4 = 1$, $\mu_2 = k_1 = -1$.

6 Conclusion

Recently, a new $(3+1)$ -dimensional BLMP equation is introduced by Wazwaz. At present, there is no literature on the lump solution of this equation. In this paper, breather wave and lump-type solutions of the new $(3+1)$ -dimensional BLMP equation are presented. Lump-type solutions contain the interaction solutions between lump and solitary waves, and the interaction solutions between lump and periodic waves, which have not been studied in any literature. Furthermore, graphical representation for lump solution is displayed in Fig. 1. Interaction phenomenon of lump-type solutions are demonstrated in Figs. 2-7.

Compliance with ethical standards

Conflict of interests The authors declare that there is no conflict of interests regarding the publication of this article.

Ethical standard The authors state that this research complies with ethical standards. This research does not involve either human participants or animals.

References

1. Boiti, M., Leon, J., Pempinelli, F.: Integrable two-dimensional generalization of the sine- and sinh-Gordon equations. *Inverse Probl.*, 3(1), 37-49 (1987)
2. Darvishi, M.T., Najafi, M., Kavitha, L., Venkatesh, M.: Stair and step soliton solutions of the integrable $(2+1)$ and $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equations. *Commun. Theor. Phys.*, 58(6), 785-794 (2012)
3. Zuo, D.W., Gao, Y.T., Yu, X., Sun, Y.H., Xue, L.: On a $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli Equation. *Z. Naturforsch. A.*, 70, 309-316 (2015)
4. Tang, Y.N., Zai, W.J.: New periodic-wave solutions for $(2+1)$ - and $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equations. *Nonlinear Dyn.*, 81(1-2), 249-255 (2015)
5. Liu, J.G., Du, J.Q., Zeng, Z.F., Nie, B.: New three-wave solutions for the $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation. *Nonlinear Dyn.*, 88(1), 655-661 (2017)
6. Liu, J.G., Tian, Y., Hu, J.G.: New non-traveling wave solutions for the $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation. *Appl. Math. Lett.*, 79, 162-168 (2018)
7. Mabrouk, S.M., Rashed, A.S.: Analysis of $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation via Lax pair investigation and group transformation method. *Comput. Math. Appl.*, 74, 2546-2556 (2017)
8. Li, B.Q., Ma, Y.L.: Multiple-lump waves for a $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation arising from incompressible fluid. *Comput. Math. Appl.*, 76, 204-214 (2018)
9. Osman, M.S., Wazwaz, A.M.: A general bilinear form to generate different wave structures of solitons for a $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation. *Math. Method. Appl. Sci.*, DOI: 10.1002/mma.5721 (2019)
10. Peng, W.Q., Tian, S.F., Zhang, T.T.: Breather waves and rational solutions in the $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation. *Comput. Math. Appl.*, 77, 715-723 (2019)
11. Xu, G.Q.: Painlevé analysis, lump-kink solutions and localized excitation solutions for the $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equation. *Appl. Math. Lett.*, 97, 81-87 (2019)
12. Wazwaz, A.M.: Painlevé analysis for new $(3+1)$ -dimensional Boiti-Leon-Manna-Pempinelli equations with constant and time-dependent coefficients. In press

13. Yang, X.J., Machado, J.A.T: A new fractal nonlinear Burgers' equation arising in the acoustic signals propagation. *Math. Method. Appl. Sci.*, DOI: 10.1002/mma.5904 (2019)
14. Yang, X.J., Feng, Y.Y., Cattani, C., Inc, M.: Fundamental solutions of anomalous diffusion equations with the decay exponential kernel. *Math. Method. Appl. Sci.*, 42(11), 4054-4060 (2019)
15. Gao, X.Y.: Mathematical view with observational/experimental consideration on certain (2+1)-dimensional waves in the cosmic/laboratory dusty plasmas. *Appl. Math. Lett.*, 91, 165-172 (2019)
16. Ma, W.X.: A generating scheme for conservation laws of discrete zero curvature equations and its application. *Comput. Math. Appl.*, 78(10), 3422-3428 (2019)
17. Liu, J.G., Yang, X.J., Feng, Y.Y.,: Resonant multiple wave solutions to some integrable soliton equations. *Chinese Phys. B*, DOI: 10.1088/1674-1056/ab4d47 (2019)
18. Wazwaz, A.M., Kaur, L.: Optical solitons for nonlinear Schrödinger (NLS) equation in normal dispersive regimes. *Optik*, 184, 428-435 (2019)
19. Lü, X., Lin, F.H., Qi, F.H.: Analytical study on a two-dimensional Korteweg-de Vries model with bilinear representation, Bäcklund transformation and soliton solutions. *Appl. Math. Model.*, 39, 3221-3226 (2015)
20. W.X. Ma, J. Li, C.M. Khalique, A Study on Lump Solutions to a Generalized Hirota-Satsuma-Ito Equation in (2+1)-Dimensions, *Complexity*, 11 (2018) 9059858
21. Lan, Z.Z., Su, J.J.: Solitary and rogue waves with controllable backgrounds for the non-autonomous generalized AB system. *Nonlinear Dyn.*, 96, 2535-2546 (2019)
22. Xie, X.Y., Meng, G.Q.: Dark solitons for the (2+1)-dimensional Davey-Stewartson-like equations in the electrostatic wave packets. *Nonlinear Dyn.*, 93, 779-783 (2018)
23. Wazwaz, A.M.: The integrable time-dependent sine-Gordon equation with multiple optical kink solutions. *Optik*, DOI: 10.1016/j.ijleo.2019.01.018 (2019)
24. Lü, X., and Ling, L.M.: Vector bright solitons associated with positive coherent coupling via Darboux transformation. *Chaos*, 25, 123103 (2015)
25. Xu, T., Chen, Y.: Semirational solutions to the coupled Fokas-CLenells equations. *Nonlinear Dyn.*, 95, 87-99 (2019)