

On the Edge Crossings of the Greedy Spanner

David Eppstein ✉

Department of Computer Science, University of California, Irvine

Hadi Khodabandeh ✉ 

Department of Computer Science, University of California, Irvine

Abstract

The greedy t -spanner of a set of points in the plane is an undirected graph constructed by considering pairs of points in order by distance, and connecting a pair by an edge when there does not already exist a path connecting that pair with length at most t times the Euclidean distance. We prove that, for any $t > 1$, these graphs have at most a linear number of crossings, and more strongly that the intersection graph of edges in a greedy t -spanner has bounded degeneracy. As a consequence, we prove a separator theorem for greedy spanners: any k -vertex subgraph of a greedy spanner can be partitioned into sub-subgraphs of size a constant fraction smaller, by the removal of $O(\sqrt{k})$ vertices. A recursive separator hierarchy for these graphs can be constructed from their planarizations in linear time, or in near-linear time if the planarization is unknown.

2012 ACM Subject Classification Theory of computation → Sparsification and spanners; Theory of computation → Computational geometry; Theory of computation → Design and analysis of algorithms

Keywords and phrases Geometric Spanners, Greedy Spanners, Separators, Crossing Graph, Sparsity

Funding This work was supported in part by the US National Science Foundation under grant CCF-1616248.


1 Introduction

Geometric spanners are geometric graphs whose distances approximate distances in complete graphs, while having fewer edges than complete graphs. Given a set of points V on the Euclidean plane (or in any other metric space), a t -spanner on V can be defined as a graph S having V as its set of vertices V and satisfying the following inequality for every pair of points (P, Q) :


$$d_S(P, Q) \leq t \cdot d(P, Q) \tag{1}$$

where $d_S(P, Q)$ is the length of the shortest path between P and Q using the edges in S , and $d(P, Q)$ is the Euclidean distance of P and Q . We call Equation 1 the *bounded stretch property*. Because of this inequality, t -spanners provide a t -approximation for the pairwise distances between the set of points in V . The parameter $t > 1$ is called the *stretch factor* or *spanning ratio* of the spanner and determines how accurate the approximate distances are; spanners having smaller stretch factors are more accurate.

Spanners can be defined in any metric space, but they are often located in a geometric space, where a heavy or undesirable network is given and finding a sparse and light-weight spanner and working with it instead of the actual network makes the computation easier and faster. Finding light-weight geometric spanners has been a topic of interest in many areas of computer science, including communication network design and distributed computing. These subgraphs have few edges and are easy to construct, leading them to appear in a wide range of applications since they were introduced [14, 38, 45]. In wireless ad hoc networks t -spanners are used to design sparse networks with guaranteed connectivity and guaranteed bounds on routing length [5]. In distributed computing spanners provide communication-efficiency and time-efficiency through the sparsity and the bounded stretch property [9, 23, 7, 24]. There



■ **Figure 1** Greedy spanners of 128 random points with stretch factor 2 (left) and 1.1 (right)



■ **Figure 2** Nonplanar greedy spanner with stretch factor 11.3

has also been extensive use of geometric spanners in the analysis of road networks [25, 2, 13]. In robotics, geometric spanners helped motion planners to design near-optimal plans on a sparse and light subgraph of the actual network [19, 43, 16]. Spanners have many other applications including computing almost shortest paths [22, 15, 46, 30], and overlay networks [12, 47, 37].

Researchers have developed various construction techniques for spanners, depending on the specific additional properties needed in these applications. Well-separated pair decomposition, θ -graphs, and greedy spanners are among the most well-known of these geometric spanner constructions. Here, we focus on the greedy spanner. It was first introduced by Althöfer [3, 4] and Bern, generalizing a pruning strategy used by Das and Joseph [17] on a triangulation of the planar graph [25].

A greedy spanner can be constructed by running the greedy spanner algorithm (Algorithm 1) on a set of points on the Euclidean plane. This short procedure adds edges one at a time to the spanner it constructs, in ascending order by length. For each pair of vertices, in this order, it checks whether that pair already satisfies the bounded stretch inequality

using the edges already added. If not, it adds a new edge connecting the pair. Therefore, by construction, each pair of vertices satisfies the inequality, either through previous edges or (if not) through the newly added edge. The resulting graph is therefore a t -spanner. Examples of the results of this algorithm, for two different stretch factors, are shown in Figure 1. Although the 2-spanner in the figure is planar, this is not true for 2-spanners in general: there exist point sets with non-planar greedy t -spanners for arbitrarily large values of t (Figure 2), and by placing widely-spaced copies of the same construction within a single point set, one can construct point sets whose greedy t -spanners have linearly many crossings, for arbitrarily large values of t .

■ **Algorithm 1** The naive greedy spanner algorithm.

```

1: procedure NAIVE-GREEDY( $V$ )
2:   Let  $S$  be a graph with vertices  $V$  and edges  $E = \{\}$ 
3:   for each pair  $(P, Q) \in V^2$  in increasing order of  $d(P, Q)$  do
4:     if  $d_S(P, Q) > t \cdot d(P, Q)$  then
5:       Add edge  $PQ$  to  $E$ 
6:   return  $S$ 

```

A naïve implementation of the greedy spanner algorithm runs in time $\mathcal{O}(n^3 \log n)$, where n is the number of given points [11]. Bose et al. [11] improved the running time of Algorithm 1 to near-quadratic time using a bounded version of Dijkstra’s algorithm. Narasimhan et al. proposed an approximate version of the greedy spanner algorithm that reached a running time of $\mathcal{O}(n \log n)$, based on the use of approximate shortest path queries [18, 36, 44].

Despite the simplicity of Algorithm 1, Farshi and Gudmundsson [29] observed that in practice, greedy spanners are surprisingly good in terms of the number of edges, weight, maximum vertex degree, and also the number of edge crossings. Many of these properties have been proven rigorously. Filster and Solomon [31] proved that greedy spanners have size and lightness that is optimal to within a constant factor for worst-case instances. They also achieved a near-optimality result for greedy spanners in spaces of bounded doubling dimension. Borradaile, Le, and Wulff-Nilsen [10] recently proved optimality for doubling metrics, generalizing a result of Narasimhan and Smid [44], and resolving an open question posed by Gottlieb [35], and Le and Solomon showed that no geometric t -spanner can do asymptotically better than the greedy spanner in terms of number of edges and lightness [41]. However, past work has not proven rigorous bounds on the number of crossings of greedy spanners.

One reason for particular interest in bounds on the number of crossings is the close relation, for geometric graphs in the plane, between crossings and *separators*. The well-known planar separator theorem of Lipton and Tarjan [42] states that any planar graph (that is, a geometric graph with no crossings) can be partitioned into subgraphs whose size is at most a constant fraction of the total by the removal of $\mathcal{O}(\sqrt{n})$ vertices. This property is central to the efficiency of many algorithms on planar graphs [34, 21, 27, 26, 40], and applied as well in multiple computational geometry problems [32, 6, 39]. Analogous separator theorems have been extended from planar graphs to graphs with few crossings per edge [20], or more generally to graphs with sparse patterns of crossings [28, 8]. Past work has not shown that greedy spanners have small separators, but as we will show, bounds on their crossings can be used to show that they do.

1.1 Our Contribution

In this paper we prove that greedy t -spanners in the Euclidean plane have few crossings, for any $t > 1$, and we use this result (together with a result of Eppstein and Gupta [28] on graphs with sparse patterns of crossings) to prove that greedy spanners in the Euclidean plane have small separators. In particular, we prove:

- **Claim 1.** Each edge in a greedy spanner can be crossed by only $O(1)$ edges of equal or greater length, where the constant in the $O(1)$ depends only on t , the stretch factor of the spanner. More precisely as $t \rightarrow 1$ there are $O(1/(t-1)^2)$ edges that cross the given edge and are longer than it by a factor of $\Omega(1/(t-1))$ (Theorem 14), and $1/(t-1)^{O(1)}$ edges that cross the given edge and have length at least ϵ times it, for any constant $\epsilon > 0$ (Theorem 17).
- **Claim 2.** For some choices of t , there exist greedy spanners in which some edges are crossed by a linear number of (significantly shorter) edges (Theorem 24).
- **Claim 3.** Every n -vertex greedy spanner, and every n -vertex subgraph of a greedy spanner, can be partitioned into connected components of size at most cn for a constant $c < 1$ by the removal of $O(\sqrt{n})$ vertices. Again, the constant factor in the $O(\sqrt{n})$ term depends only on the stretch factor of the spanner. Moreover, a separator hierarchy for the greedy spanner can be constructed from its planarization in near-linear time (Theorem 20).

It is known that the spanners that are constructed by some other methods, i.e. semi-separated pair decomposition [1] and hierarchical decomposition [33], have small $\mathcal{O}(\sqrt{n})$ -separators in two dimensions. Although experimental results of Farshi and Gudmundsson on greedy spanners of random point sets had shown the number of crossings to be small in practice [29] our results are the first theoretical results on this property, the first to study crossings for worst-case and not just random instances, and the first to prove that greedy spanners have small $\mathcal{O}(\sqrt{n})$ -separators.

1.2 Intuition

Our proof that edges can be crossed by only a bounded number of edges of greater or equal length splits into two cases, one for crossings by edges of significantly greater length and another for crossings by edges of similar length.

For edges of significantly greater length, we divide the greedy spanner edges that might cross the given edge into a constant number of nearly-parallel sets of edges, and prove the bound separately within each such set. We show that, within a set of nearly-parallel long edges that all cross the given edge, the edges can be totally ordered by their projections onto a base line, because edges whose endpoints project to nested intervals would contradict the greedy property of the spanner (the inner of two nested edges could be used to shortcut the outer one). By similar reasoning, the endpoints of any two nearly-parallel long crossing edges are separated by a distance that is at least a constant fraction of the length of the smaller edge. This geometric growth in the separation of the endpoints leads to a system of inequalities on the lengths of the edges that can only be satisfied when the number of crossing edges is bounded by a constant.

For edges of comparable length to the crossed edge, we use a grid to partition the crossing edges into a constant number of subsets of edges, such that within each subset all edges have pairs of endpoints that are close to each other relative to the length of the edge, and we show that each of these subsets can contain only a unique edge.

Our construction showing that a single edge can be crossed linearly many times is based on the combination of three “zig-zag” sets of points, evenly spaced in their x -coordinates and alternating between two different y -coordinates. In the top and bottom zig-zag, the distance along the zigzag between two consecutive points with the same y -coordinates is exactly t times the difference between their x -coordinates, while in the middle zig-zag it is slightly greater. The greedy spanner for this point set contains the zig-zag edges, plus a single long edge crossing all of the middle edges, for a pair of points that are far enough from each other along the middle zig-zag for their Euclidean distance to be almost the same as their difference in x -coordinates (differing by a number smaller than the amount by which a single edge of the middle zig-zag exceeds t times its difference in x -coordinates).

The results on separators follow from previous results on the existence of separators in graphs whose edge intersection graphs have bounded degeneracy [28].

2 Preliminaries

As we mentioned earlier, t -spanners can be defined in any metric space. For a given graph G , a t -spanner is defined in the following way,

► **Definition 1** (t -spanner). *Given a metric graph $G = (V, E, d)$, i.e. weighted graph with distances as weights, a t -spanner is a spanning subgraph S of G such that for any pair of vertices $u, w \in V$,*

$$d_G(u, w) \leq t \cdot d(u, w)$$

where $d_G(u, w)$ is the length of the shortest path in G between u and w .

Then the greedy spanner on a given set of points V can be defined in the following way,

► **Definition 2** (greedy spanner). *Given a set of points V in any metric space, a greedy spanner on V is a t -spanner that is an output of Algorithm 1.*

Here we restrict the problem to geometric graphs and we take advantage of inequalities that hold in geometric space.

We consider the natural embedding that the greedy spanner inherits from its vertices. Edges are drawn as straight segments between the two points corresponding to the two endpoints of the edge. We say two edges of the spanner cross or intersect if their corresponding segments intersect at some interior point. The crossing graph of a given embedding can be defined in this way,

► **Definition 3** (crossing graph). *Given a graph $G(V, E)$ and its Euclidean embedding, the crossing graph $Cr(G)$ is a graph $G'(E, C)$ whose vertices are the edges of the original graph and for each two vertices $e, f \in E$ there is an edge between them if and only if they intersect with each other in the embedding given for G .*

Most of the proofs here use a lemma that we call the *short-cutting lemma*, which is simple but very useful in greedy spanners. The lemma is proven in [44] and it states that a t -spanner edge cannot be shortcut by some other edges of the spanner by a factor of t . Formally,

► **Lemma 4** (short-cutting). *An edge AB of a greedy t -spanner cannot be shortcut by some other spanner edges by a factor of t , i.e. there is no constant k and points $A = P_0, P_1, \dots, P_k = B$ that $P_0P_1, P_1P_2, \dots, P_{k-1}P_k$ are all spanner edges distinct from AB , and*

$$\sum_{i=0}^{k-1} |P_i P_{i+1}| \leq t \cdot |AB|$$

If some of the segments $P_i P_{i+1}$ are not included in the spanner, the same argument still works but a factor t appears before the term $|P_i P_{i+1}|$ in the summation. So

► **Corollary 5** (Extended short-cutting). *Given a greedy t -spanner S and an edge AB of S , there cannot be a constant k and points $A = P_0, P_1, \dots, P_k = B$ such that*

$$\sum_{P_i P_{i+1} \in S} |P_i P_{i+1}| + t \cdot \sum_{P_i P_{i+1} \notin S} |P_i P_{i+1}| \leq t \cdot |AB|$$

The proof of Lemma 4 and Corollary 5 are included in Appendix A for reference. In the following section we consider intersections between an arbitrary edge of a greedy spanner and sufficiently larger edges, and we show a constant bound on the number of intersections per edge. In subsection 3.5 we again prove a constant bound for the number of intersections between a spanner edge and other edges of almost the same length. Finally, in Appendix B we introduce an example in which the number of intersections with smaller edges can be more than any constant bound, completing our analysis. In section 4 we introduce some new results and improvements based on the constant bound we provided earlier.

3 Few intersections with long edges

In this section, we prove an upper bound on the number of intersections of an edge with sufficiently larger edges. We will specifically show that the number of intersections, in this case, has a constant bound that only depends on t . Later in subsection 3.5 we prove a constant bound also exists for the intersections with the edges that have almost the same length of the intersecting edge. Hence we prove our first claim.

In this setting, we consider an arbitrary edge AB of the spanner, and we are interested in counting the number of intersections that AB may have with sufficiently larger edges, i.e. edges PQ that intersect AB at some interior point with $|PQ| > c \cdot |AB|$ for some constant $c > 1$ which we will specify later.

First, we only consider a set of *almost-parallel* spanner segments that cross AB , where we define the term *almost-parallel* below, and we put a bound on the number of these segments. Then we generalize the bound to hold for all large spanner segments that cross AB .


3.1 Definitions

► **Definition 6** (almost-parallel). *We say a pair of arbitrary segments PQ and RS in the plane are almost-parallel or θ -parallel if there is an angle of at most θ between them. We say a set of segments are almost-parallel if every pair of segments chosen from the set are almost-parallel.*

For any set of almost-parallel segments, we define a baseline to measure the angles and distances with respect to that line.

► **Definition 7** (baseline). *Given a set of almost-parallel (or θ -parallel) segments in the plane, denoted by S , the baseline $b(S)$ of the set of segments S is the segment with the smallest slope.*

We use the uniqueness of the segment chosen in Definition 7 and we emphasize that any other definition works if it determines a unique segment for any almost-parallel set of segments.



■ **Figure 3** Ordering segments by projecting on the baseline l , here $P_i Q_i <_{\mathcal{R}} P_j Q_j$.

In subsection 3.2, we define a total ordering on a set of almost-parallel segments that cross a spanner segment AB . Once we have sorted these segments based on the ordering, in subsection 3.3 we prove the distance between the endpoints of two consecutive segments is at least a constant fraction of the length of the smaller segment. Putting together these two parts, in subsection 3.4 we prove there cannot be more than a constant number of segments in the sequence.

3.2 A total ordering on almost-parallel intersecting segments

In this section, we define an ordering on a set of almost-parallel segments of the t -spanner. The ordering is based on the order of the projections of the endpoints of the segments on the baseline corresponding to the segments. We first define the ordering and then we use Lemma 9 and Lemma 10 to prove that it is a total ordering when the set of almost-parallel segments are all crossing a given segment of the spanner.

Consider a set of almost-parallel spanner segments that cross some spanner segment. One can define an ordering on this set of almost-parallel segments, which we call the *endpoint-ordering*, based on how their endpoints are ordered along the direction they are aligned to. We formulate the definition in the following way,


► **Definition 8** (endpoint-ordering). *Let $S = \{P_i Q_i : i = 1, 2, \dots, k\}$ be a set of almost-parallel segments. Also let l be the baseline of S , $b(S)$. Define the endpoint-ordering \mathcal{R} between two segments $P_i Q_i$ and $P_j Q_j$ by projecting the endpoints P_i, P_j, Q_i, Q_j to the baseline l and comparing the order of the projected points P'_i, P'_j, Q'_i, Q'_j along an arbitrary direction of the baseline l ,*

- $P_i Q_i <_{\mathcal{R}} P_j Q_j$ if the projections are ordered as $P'_i P'_j Q'_i Q'_j$ or $P'_i Q'_i P'_j Q'_j$.
- $P_i Q_i >_{\mathcal{R}} P_j Q_j$ if they are ordered as $P'_j P'_i Q'_j Q'_i$ or $P'_j Q'_j P'_i Q'_i$. (Figure 3)

We claim that the endpoint-ordering is a total ordering on the set of almost-parallel segments. This basically means that after projecting two almost-parallel segments on the baseline, none of the resulting projections would lie completely inside the other one. Other cases correspond to a valid endpoint-ordering.

In order to prove this, first, we prove a simpler case when the two segments intersect with each other. This assumption will help to significantly simplify the proof. Later we use this lemma to show the original claim is also true.

► **Lemma 9.** *Let MN and PQ be two intersecting segments from a set of θ -parallel spanner segments. Also assume that $\theta < \frac{t-1}{2t}$ where t is the stretch factor of the spanner. Then MN*



■ **Figure 4** Proof of Lemma 9.

and PQ are endpoint-ordered, i.e. the projection of one of the segments on the baseline of the set cannot be included in the projection of the other one.

Proof. We prove the lemma by contradiction. Without loss of generality suppose that the projections of P and Q on some baseline l are both between the projections of M and N (on the same baseline). We show that MN can be shortcut by PQ by a factor of t , i.e.

$$t \cdot |MP| + |PQ| + t \cdot |QN| \leq t \cdot |MN|$$

Let P', Q', M' , and N' be the corresponding projections of P, Q, M , and N on l , respectively (Figure 4). Also let I be the intersection point and $\alpha = \angle PMI$, and also γ to be the angle between MN and the baseline, according to the figure. By the assumption P' is between M' and N' , so $\alpha \leq \pi/2 + \gamma \leq \pi/2 + \theta$. Let also P'' be the point on MN s.t. $|MP''| = |MP|$ and $\beta = \angle MPP'' = \angle MP''P$. Then by sine law,

$$\begin{aligned} \frac{|MI| - |MP|}{|PI|} &= \frac{|P''I|}{|PI|} = \frac{\sin(\beta - \theta)}{\sin \beta} = \frac{\sin(\pi/2 - \alpha/2 - \theta)}{\sin(\pi/2 - \alpha/2)} = \frac{\cos(\alpha/2 + \theta)}{\cos(\alpha/2)} \\ &= \cos \theta - \sin \theta \tan(\alpha/2) \end{aligned} \quad (2)$$

but we have,

$$\cos \theta \geq 1 - \theta^2/2 \geq 1 - \theta/4 \quad (3)$$

as $\theta < \frac{t-1}{2t} < 1/2$. Also,

$$\tan(\alpha/2) \leq \tan(\pi/4 + \theta/2) = \tan(\pi/4 + 1/4) < \frac{7}{4} \quad (4)$$

Putting together Equation 2, Equation 3, and Equation 4, also using $\sin \theta \leq \theta$,

$$\frac{|MI| - |MP|}{|PI|} \geq (1 - \theta/4) - \left(\frac{7}{4}\right)\theta = 1 - 2\theta > \frac{1}{t}$$

which is equivalent to $t \cdot |MI| - t \cdot |MP| \geq |PI|$. Similarly, $t \cdot |NI| - t \cdot |NQ| \geq |QI|$. Adding together,

$$t \cdot |MN| - t \cdot |MP| - t \cdot |NQ| \geq |PQ|$$

which is what we are looking for. ◀

Lemma 9 assumes that segments intersect at some interior point. In order to prove the totality of the ordering, we also need to prove the claim when the segments do not intersect with each other. Instead, in this case, both segments intersect some spanner edge. We use Lemma 9 to prove this in the Lemma 10.

► **Lemma 10.** *Let MN and PQ be two segments chosen from a set of θ -parallel spanner segments that cross a spanner edge AB . Also assume that $\theta < \frac{t-1}{2(t+1)}$, and $\min(|MN|, |PQ|) \geq \frac{3t(t+1)}{t-1}|AB|$, where t is the spanner parameter. Then MN and PQ are endpoint-ordered.*

The proof of this lemma is included in Appendix A. Based on Lemma 10 it is easy to prove the main result of this section, Proposition 11.

► **Proposition 11.** *Given an arbitrary edge AB of a t -spanner, for a set of sufficiently large almost-parallel spanner edges that intersect AB , the endpoint-ordering we defined in Definition 8 is a total ordering.*

Proof. Totality requires reflexivity, anti-symmetry, transitivity, and comparability. Reflexivity and transitivity are trivial because of the projection. Anti-symmetry and comparability follow directly from Lemma 10. ◀

Now that we have ordered the set of almost-parallel spanner segments, we can prove a lower bound on the distance of two ordered segments. Later we prove a bound on the number of these segments based on the resulting distance lower bound.

3.3 Lower bounding the distance of endpoints of two crossing segments

In subsection 3.2 we restricted the problem to a set of almost-parallel spanner segments that intersect another spanner segment, and we defined an ordering on these segments. The next step is to find a lower bound on the distance of two almost-parallel segments that intersect some spanner segment AB . The idea is to show that both endpoints of two ordered segments cannot be arbitrarily close, and hence there cannot be more than a constant number of them in a sequence.


More specifically, we show in Proposition 13 that the corresponding endpoints of two almost-parallel spanner segments that both cross the same spanner segment should have a distance of at least a constant fraction of the length of the smaller segment, otherwise the longer segment could be shortcut by the smaller one, which is indeed a contradiction. A weaker version of this lemma is proven in [44] and it is called “gap property”, but the inequality we show here is actually stronger.

First we propose a geometric inequality in Lemma 12 that helps to prove the proposition. Then we complete the proof of the proposition at the end of this section.

► **Lemma 12.** *Let MN and PQ be two segments in the plane with angle θ . Then*

$$||MN| - |PQ|| > ||MP| - |NQ|| - 2 \sin(\theta/2) \cdot |PQ|$$

Proof. By swapping MN and PQ , it turns out that the case where $|MN| \geq |PQ|$ is stronger than $|MN| \leq |PQ|$. So without loss of generality, let $|MN| \geq |PQ|$ and by symmetry $|MP| \geq |NQ|$. Let Q' be the rotation of Q around P by θ , so that PQ' and MN are parallel, and $|PQ'| = |PQ|$ (Figure 5). Let Q'' be the point on the ray PQ' where $|PQ''| = |MN|$.



■ **Figure 5** Proof of Lemma 12.

As a result Q'' and P will be on different sides of Q' . By the triangle inequality,

$$\begin{aligned}
 |MN| - |PQ| &= |PQ''| - |PQ'| = |Q'Q''| \geq |NQ''| - |NQ'| \\
 &= |MP| - |NQ'| \geq |MP| - (|NQ| + |QQ'|) \\
 &= |MP| - |NQ| - 2|PQ| \cdot \sin(\theta/2)
 \end{aligned}$$

◀

Now we state and prove Proposition 13. As we mentioned earlier, the idea is to show one of the segments can be shortcut by the other one if one of the matching endpoints is very close. In the simplest case when the segments are two opposite sides of a rectangle, it is easy to see that a distance of $\frac{t-1}{2}|PQ|$ on both sides is required to prevent short-cutting. In the general case, when the segments are placed arbitrarily, Proposition 13 holds.

► **Proposition 13.** *Let MN and PQ be two θ -parallel spanner segments. The matching endpoints of these two segments cannot be closer than a constant fraction of the length of the smaller segment. More specifically,*

$$\min(|MP|, |NQ|) \geq \frac{t-1-2\sin(\theta/2)}{2t} \min(|MN|, |PQ|)$$

Proof. Without loss of generality and by symmetry, let $|NQ| \leq |MP|$. Suppose, on the contrary, that $|NQ| < \frac{t-1-2\sin(\theta/2)}{2t} |PQ|$. Then,

$$\begin{aligned}
 t \cdot |MP| + |PQ| + t \cdot |NQ| &\leq t \cdot (|MN| - |PQ| + |NQ| + 2\sin(\theta/2) \cdot |PQ|) + |PQ| + t \cdot |NQ| \\
 &= t \cdot |MN| - (t-1-2\sin(\theta/2))|PQ| + 2t \cdot |NQ| \\
 &< t \cdot |MN|
 \end{aligned}$$

So MN can be shortcut by PQ within a factor of t which contradicts the extended short-cutting lemma for the edge MN and the path $MPQN$. ◀

So far, in Proposition 13 we proposed an ordering on the set of almost-parallel spanner segments that cross a given edge and we proved each of these segments has a significant distance from the other ones. In the next section we put together these results and we find a constant upper bound on the number of these segments.

3.4 Putting together

Based on the ordering proposed in subsection 3.2, and the lower bound we proved in subsection 3.3, we can show that the following constant upper bound on the number of intersections with sufficiently large edges holds.

If we look at one of the endpoints of the endpoint-ordered sequence of almost-parallel spanner segments, and we project them on the baseline, the distance of every two consecutive projected points cannot be smaller than a constant fraction of the length of the smaller segment, i.e. $|P'_i P'_{i+1}| \geq C \cdot \min(|P_i Q_i|, |P_{i+1} Q_{i+1}|)$ for all values of $i = 0, 1, \dots, k-1$. Summing up these inequalities leads to a bound on k , the number of segments.

► **Theorem 14.** *For sufficiently small θ , the number of sufficiently large θ -parallel segments that intersect a given edge AB of a t -spanner is limited by*

$$\frac{4t}{(t-1-2\sin(\theta/2))\cos\theta} + 1$$

By sufficiently large we specifically mean larger than $\frac{3t(t+1)}{t-1}|AB|$.

Proof. Let $P_i Q_i$ s be the segments larger than AB that intersect AB at some angle in $[\alpha, \alpha + \theta)$. Let $P_0 Q_0$ be the shortest edge among $P_i Q_i$ s. Because of the total ordering, at least half of the segments are larger than $P_0 Q_0$ with respect to the ordering \mathcal{R} , or at least half of them are smaller than $P_0 Q_0$ with respect to \mathcal{R} . Without loss of generality, assume that half of the segments are larger than $P_0 Q_0$ with respect to \mathcal{R} , and they are indexed by $i = 1, 2, \dots, (k-1)/2$. Also let P'_i s and Q'_i s be the projections of P_i s and Q_i s on the base line l . By Proposition 13, for all i , P_{i+1} is farther than P_i by a constant fraction of $\min(|P_i Q_i|, |P_{i+1} Q_{i+1}|)$, so

$$\begin{aligned} \sum_{i=0}^{(k-3)/2} |P_i P_{i+1}| &> \frac{t-1-2\sin(\theta/2)}{2t} \sum_{i=0}^{(k-3)/2} \min(|P_i Q_i|, |P_{i+1} Q_{i+1}|) \\ &\geq \frac{t-1-2\sin(\theta/2)}{2t} \cdot \frac{k-1}{2} |P_0 Q_0| \end{aligned}$$

$$\text{If } k \geq \frac{4t}{(t-1-2\sin(\theta/2))\cos\theta} + 1,$$

$$\sum_{i=0}^{(k-3)/2} |P_i P_{i+1}| > \frac{1}{\cos\theta} |P_0 Q_0|$$

or equivalently

$$|P_0 Q_0| < \sum_{i=0}^{(k-3)/2} |P_i P_{i+1}| \cos\theta \leq \sum_{i=0}^{(k-3)/2} |P'_i P'_{i+1}| = |P'_0 P'_{\frac{k-1}{2}}|$$

which is not possible, because $P'_0 P'_{\frac{k-1}{2}}$ lies inside $P'_0 Q'_0$ and so $|P'_0 P'_{\frac{k-1}{2}}| \leq |P'_0 Q'_0| \leq |P_0 Q_0|$ which contradicts the last inequality above. ◀

The constraints on θ imposed by our earlier lemmas imply that, as $t \rightarrow 1$, we should choose θ proportional to $t-1$. Asymptotically, as $t \rightarrow 1$, the number of large segments of all angles that intersect AB is $O(1/(t-1)^2)$, with one factor of $1/(t-1)$ coming from the bound in the theorem and the second factor coming from the number of different classes of nearly-parallel segments.

3.5 Almost-equal length edges

In the previous subsections, we proved a bound on the number of intersections with relatively larger edges. Here we prove a constant bound on the number of intersections with edges that are nearly the same length as the length of the intersecting edge. Later in Appendix B we consider intersections with relatively smaller edges, which completes our analysis for this problem.

For same-length intersections Lemma 10 does not hold anymore, hence the endpoint-ordering is not necessarily a total ordering in this case. Since totality is a key requirement for the rest of the proof the same proof will not work anymore. But Proposition 13 still holds as it has no assumption on the ordering of the segments.

Our idea is to partition the neighborhood of AB into a square network, such that no two spanner segments can have both endpoints in the same squares (Figure 6). If this happens, then by Proposition 13 one of the segments should be shortcut by the other one, leading to a contradiction because both segments are already included in the spanner.

We first prove a simpler version of Proposition 13 that does not include θ in the inequality, as we are not using the almost-parallel assumption and the value of θ can be large enough to make the inequality in Proposition 13 trivial. We will use this modified version to prove our claim.

► **Lemma 15.** *Given a greedy spanner with parameter t and two spanner segments MN and PQ ,*

$$\max(|MP|, |NQ|) \geq \frac{t-1}{2t} \min(|MN|, |PQ|)$$

Proof. Suppose on the contrary that

$$\max(|MP|, |NQ|) < \frac{t-1}{2t} \min(|MN|, |PQ|)$$

Also, without loss of generality assume that $|MN| \geq |PQ|$. Then,

$$t \cdot |MP| + |PQ| + t \cdot |NQ| \leq (t-1) \min(|MN|, |PQ|) + |PQ| = t \cdot |PQ| \leq t \cdot |MN|$$

which contradicts the extended short-cutting lemma for the edge MN and the path $MPQN$. ◀


► **Proposition 16.** *The number of spanner segments PQ that cross a segment AB of a t -spanner and that have length within $\alpha \cdot |AB| \leq |PQ| \leq \beta \cdot |AB|$ is limited by*

$$\left[\frac{2\beta(2\beta+1)}{\alpha^2} \cdot \frac{8t^2}{(t-1)^2} \right]^2$$

where t is the spanner parameter.

Proof. Partition the area around AB with squares of edge length $\frac{t-1}{2\sqrt{2}t} \cdot \alpha|AB|$ with edges parallel or perpendicular to AB . The area that an endpoint of a crossing segment can lie in is a rectangle of size $(2\beta+1)|AB|$ by $2\beta|AB|$ (Figure 6). The total number of squares in this area would be

$$\frac{2\beta(2\beta+1)}{\alpha^2} \cdot \frac{8t^2}{(t-1)^2}$$



■ **Figure 6** Partition of the area around AB

But for each crossing segment the pair of squares that contain the two endpoints of the segment is unique. Otherwise two segments, e.g. MN and PQ , will have both endpoints at the same pair, which means

$$\max(|MP|, |NQ|) < (\sqrt{2}) \left(\frac{t-1}{2\sqrt{2}t} \cdot \alpha |AB| \right) = \frac{t-1}{2t} \cdot \alpha |AB| \leq \frac{t-1}{2t} \min(|MN|, |PQ|)$$

which cannot happen due to Lemma 15. So the total number of pairs, and hence the total number of crossing segments, would be

$$\left[\frac{2\beta(2\beta+1)}{\alpha^2} \cdot \frac{8t^2}{(t-1)^2} \right]^2$$

◀

In Proposition 16 both α and β can be chosen arbitrarily, and the bound is a strictly increasing function of β and a strictly decreasing function of α . The bound tends to infinity when β is large enough, and also when α is small enough. So it basically does not prove any constant bound for the cases that edges are very small or very large. But for the edges of almost the same length, it gives a constant upper bound.

Putting together the main results of section 3 and subsection 3.5 we can prove the following bound for the number of intersections with not-relatively-small spanner segments.

► **Theorem 17.** *Given a spanner segment AB in the Euclidean plane and a positive constant ϵ , the number of edges of length at least $\epsilon \cdot |AB|$ of the spanner that intersect AB is $\mathcal{O}(\frac{t^{12}}{\epsilon^4(t-1)^8})$.*

Proof. By Theorem 14 the number of intersections with edges PQ such that $|PQ| \geq \frac{3t(t+1)}{t-1} |AB|$ is bounded by

$$C_1 = \frac{4t}{(t-1-2\sin(\theta/2))\cos\theta} + 1 \in \mathcal{O}\left(\frac{t}{t-1}\right)$$

On the other side, putting $\alpha = \epsilon$ and $\beta = \frac{3t(t+1)}{t-1}$ into Proposition 16 implies that the number of intersections with edges larger than AB and smaller than $\frac{3t(t+1)}{t-1} |AB|$ is at most

$$C_2 = \left[\frac{2}{\epsilon^2} \left(\frac{3t(t+1)}{t-1} \right) \left(2 \frac{3t(t+1)}{t-1} + 1 \right) \left(\frac{8t^2}{(t-1)^2} \right) \right]^2 \in \mathcal{O}\left(\frac{t^{12}}{\epsilon^4(t-1)^8}\right)$$

Hence the number of intersections with edges larger than AB is at most $C_1 + C_2$, which is $\mathcal{O}(\frac{t^{12}}{\epsilon^4(t-1)^8})$. ◀

In section 3 we proved the number of intersections with sufficiently large edges is bounded by a constant and now we completed the proof for all larger edges. In Appendix B, we show that the same argument does not work for intersections with arbitrarily smaller edges, and we provide an example that shows there can be an arbitrarily large number of intersections with smaller edges. This completes our analysis of the problem. In the following section, we show some of the applications of this result, most importantly, the sparsity of the crossing graph of the greedy spanner.

4 Separators

In this section, we use the crossing bound that we proved in Theorem 17 to show that greedy spanners have small separators. First, we start with the definition of *degeneracy*, which is a measure of sparsity of a graph.

► **Definition 18** (degeneracy). *A graph G is called k -degenerate, if each subgraph of G has a vertex of degree at most k . The smallest value of k for which a graph is k -degenerate is called the degeneracy of the graph.*

The first important consequence of Theorem 17 is the constant degeneracy of the crossing graph of the greedy spanner, implying its sparsity and linearity of the number of edges, i.e. crossing.

► **Theorem 19.** *The crossing graph of a greedy t -spanner has a constant degeneracy.*

Proof. In any subgraph of the crossing graph, by Theorem 17 the node corresponding to the smallest edge has at most a constant number of neighbours. ◀

This, together with the result of [28] implies the existence of sublinear separators for greedy spanners. A separator is a subset of vertices whose removal splits the graph into smaller pieces. A sublinear separator is a sublinear number of vertices with the same property. The splitting can be recursively performed on the smaller parts and a separator hierarchy can be constructed in this way, which effectively helps in the design of new recursive algorithms. The planarization of a graph, which is obtained by adding new vertices on the edge intersections of the graph, would help us to find such hierarchy in linear time, otherwise, a near-linear time algorithm would be used.

► **Theorem 20.** *Greedy spanners have separators of size $\mathcal{O}(\sqrt{n})$. Also, a separator hierarchy for them can be constructed from their planarization in linear time.*

Proof. By Theorem 19 the crossing graph of the greedy t -spanner has a constant degeneracy, so by Theorem 6.9 of [28] they have separators of size $\mathcal{O}(\sqrt{n})$. Also by the same theorem, a separator hierarchy for them can be constructed from their planarization in linear time. ◀

One of the basic algorithms that can be improved using the separator hierarchy is Dijkstra's single-source shortest path algorithm, which runs in $\mathcal{O}(n \log n)$ time on a graph with n vertices. As a result of Theorem 20 linear algorithms exist for finding single-source shortest path on greedy spanners. If the planarization has not already been found, it can be constructed in time $\mathcal{O}(n \log^{(i)} n)$ for any constant i , where $\log^{(i)}$ denotes the i -times iterated logarithm, e.g. $\log^{(3)} n = \log \log \log n$ [27].

► **Corollary 21.** *Single source shortest paths can be computed in time $O(n \log^{(i)} n)$ on a greedy spanner.*

Proof. This follows from the planarization algorithm and from the existence and construction of separators from planarizations by Corollary 6.10 of [28]. ◀

5 Conclusions

We have shown that greedy t -spanners in the plane have linearly many crossings, and that the intersection graphs of their edges have bounded degeneracy but can have unbounded (and even linear) degree. As a consequence, we proved that these graphs have small separators.

Given these results, it is natural to ask whether higher-dimensional Euclidean greedy t -spanners also have small separators. This cannot be achieved through bounds on crossings, because in dimensions greater than two, graphs whose vertices are in general position can have no crossings. We leave this question open for future research.

References

- 1 Mohammad A Abam and Sarel Har-Peled. New constructions of sspds and their applications. *Computational Geometry*, 45(5-6):200–214, 2012.
- 2 Mohammad Ali Abam, Mark De Berg, Mohammad Farshi, and Joachim Gudmundsson. Region-fault tolerant geometric spanners. *Discrete & Computational Geometry*, 41(4):556–582, 2009. doi:10.1007/s00454-009-9137-7.
- 3 Ingo Althöfer, Gautam Das, David Dobkin, and Deborah Joseph. Generating sparse spanners for weighted graphs. In John R. Gilbert and Rolf Karlsson, editors, *Proceedings of the 2nd Scandinavian Workshop on Algorithm Theory (SWAT)*, volume 447 of *Lecture Notes in Computer Science*, pages 26–37. Springer, 1990. doi:10.1007/3-540-52846-6_75.
- 4 Ingo Althöfer, Gautam Das, David Dobkin, Deborah Joseph, and José Soares. On sparse spanners of weighted graphs. *Discrete & Computational Geometry*, 9(1):81–100, 1993. doi:10.1007/BF02189308.
- 5 Khaled Alzoubi, Xiang-Yang Li, Yu Wang, Peng-Jun Wan, and Ophir Frieder. Geometric spanners for wireless ad hoc networks. *IEEE Transactions on Parallel and Distributed Systems*, 14(4):408–421, 2003. doi:10.1109/TPDS.2003.1195412.
- 6 Srinivasa Arikati, Danny Z. Chen, L. Paul Chew, Gautam Das, Michiel Smid, and Christos D. Zaroliagis. Planar spanners and approximate shortest path queries among obstacles in the plane. In Josep Diaz and Maria Serna, editors, *Proceedings of the 4th European Symposium on Algorithms (ESA)*, volume 1136 of *Lecture Notes in Computer Science*, pages 514–528. Springer, 1996. doi:10.1007/3-540-61680-2_79.
- 7 Baruch Awerbuch, Bonnie Berger, Lenore Cowen, and David Peleg. Near-linear time construction of sparse neighborhood covers. *SIAM Journal on Computing*, 28(1):263–277, 1998. doi:10.1137/S0097539794271898.
- 8 Sang Won Bae, Jean-Francois Baffier, Jinhee Chun, Peter Eades, Kord Eickmeyer, Luca Grilli, Seok-Hee Hong, Matias Korman, Fabrizio Montecchiani, Ignaz Rutter, and Csaba D. Tóth. Gap-planar graphs. *Theoretical Computer Science*, 745:36–52, 2018. doi:10.1016/j.tcs.2018.05.029.
- 9 Surender Baswana, Telikepalli Kavitha, Kurt Mehlhorn, and Seth Pettie. Additive spanners and (α, β) -spanners. *ACM Transactions on Algorithms*, 7(1):5, 2010. doi:10.1145/1868237.1868242.
- 10 Glencora Borradaile, Hung Le, and Christian Wulff-Nilsen. Greedy spanners are optimal in doubling metrics. In *Proceedings of the 30th ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2371–2379. Society for Industrial and Applied Mathematics, 2019. doi:10.1137/1.9781611975482.145.

- 11 Prosenjit Bose, Paz Carmi, Mohammad Farshi, Anil Maheshwari, and Michiel Smid. Computing the greedy spanner in near-quadratic time. *Algorithmica*, 58(3):711–729, 2010. doi:10.1007/s00453-009-9293-4.
- 12 Rebecca Braynard, Dejan Kostic, Adolfo Rodriguez, Jeffrey Chase, and Amin Vahdat. Opus: an overlay peer utility service. In *Proceedings of the 5th IEEE Conference on Open Architectures and Network Programming (OPENARCH)*, pages 167–178. IEEE, 2002. doi:10.1109/OPNARC.2002.1019237.
- 13 Shiri Chechik, Michael Langberg, David Peleg, and Liam Roditty. Fault tolerant spanners for general graphs. *SIAM Journal on Computing*, 39(7):3403–3423, 2010. doi:10.1137/090758039.
- 14 Paul Chew. There are planar graphs almost as good as the complete graph. *Journal of Computer and System Sciences*, 39(2):205–219, 1989. doi:10.1016/0022-0000(89)90044-5.
- 15 Edith Cohen. Fast algorithms for constructing t -spanners and paths with stretch t . *SIAM Journal on Computing*, 28(1):210–236, 1998. doi:10.1137/S0097539794261295.
- 16 Gautam Das. The visibility graph contains a bounded-degree spanner. In *Proceedings of the 9th Canadian Conference on Computational Geometry (CCCG)*, pages 70–75, 1997. URL: <https://cccg.ca/proceedings/1997/>.
- 17 Gautam Das and Deborah Joseph. Which triangulations approximate the complete graph? In Hristo Djidjev, editor, *Proceedings of the International Symposium on Optimal Algorithms (OA)*, volume 401 of *Lecture Notes in Computer Science*, pages 168–192. Springer, 1989. doi:10.1007/3-540-51859-2_15.
- 18 Gautam Das and Giri Narasimhan. A fast algorithm for constructing sparse Euclidean spanners. *International Journal of Computational Geometry & Applications*, 7(04):297–315, 1997. doi:10.1142/S0218195997000193.
- 19 Andrew Dobson and Kostas E. Bekris. Sparse roadmap spanners for asymptotically near-optimal motion planning. *International Journal of Robotics Research*, 33(1):18–47, 2014. doi:10.1177/0278364913498292.
- 20 Vida Dujmović, David Eppstein, and David R. Wood. Structure of graphs with locally restricted crossings. *SIAM Journal on Discrete Mathematics*, 31(2):805–824, 2017. doi:10.1137/16M1062879.
- 21 Zdenek Dvorak and Sergey Norin. Strongly sublinear separators and polynomial expansion. *SIAM Journal on Discrete Mathematics*, 30(2):1095–1101, 2016. doi:10.1137/15M1017569.
- 22 Michael Elkin. Computing almost shortest paths. *ACM Transactions on Algorithms*, 1(2):283–323, 2005. doi:10.1145/1103963.1103968.
- 23 Michael Elkin and David Peleg. $(1 + \epsilon, \beta)$ -spanner constructions for general graphs. *SIAM Journal on Computing*, 33(3):608–631, 2004. doi:10.1137/S0097539701393384.
- 24 Michael Elkin and Jian Zhang. Efficient algorithms for constructing $(1 + \epsilon, \beta)$ -spanners in the distributed and streaming models. *Distributed Computing*, 18(5):375–385, 2006. doi:10.1007/s00446-005-0147-2.
- 25 David Eppstein. Spanning trees and spanners. In Jörg-Rüdiger Sack and Jorge Urrutia, editors, *Handbook of Computational Geometry*, pages 425–461. North-Holland, 2000. doi:10.1016/B978-044482537-7/50010-3.
- 26 David Eppstein and Michael T. Goodrich. Studying (non-planar) road networks through an algorithmic lens. In *Proceedings of the 16th ACM SIGSPATIAL International Conference on Advances in Geographic Information Systems*, pages A16:1–A16:10. ACM, 2008. doi:10.1145/1463434.1463455.
- 27 David Eppstein, Michael T. Goodrich, and Darren Strash. Linear-time algorithms for geometric graphs with sublinearly many edge crossings. *SIAM Journal on Computing*, 39(8):3814–3829, 2010. doi:10.1137/090759112.
- 28 David Eppstein and Siddharth Gupta. Crossing patterns in nonplanar road networks. In *Proceedings of the 25th ACM SIGSPATIAL International Conference on Advances in Geographic Information Systems*, pages A40:1–A40:9. ACM, 2017. doi:10.1145/3139958.3139999.

- 29 Mohammad Farshi and Joachim Gudmundsson. Experimental study of geometric t -spanners. *ACM Journal of Experimental Algorithmics*, 14:1.3:1–1.3:29, 2009. doi:10.1145/1498698.1564499.
- 30 Joan Feigenbaum, Sampath Kannan, Andrew McGregor, Siddharth Suri, and Jian Zhang. Graph distances in the streaming model: the value of space. In *Proceedings of the 16th ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 745–754. Society for Industrial and Applied Mathematics, 2005.
- 31 Arnold Filtser and Shay Solomon. The greedy spanner is existentially optimal. In *Proceedings of the 35th ACM SIGACT-SIGOPS Symposium on Principles of Distributed Computing (PODC)*, pages 9–17. ACM, 2016. doi:10.1145/2933057.2933114.
- 32 Alan M. Frieze, Gary L. Miller, and Shang-Hua Teng. Separator based parallel divide and conquer in computational geometry. In *Proceedings of the 4th ACM Symposium on Parallel Algorithms and Architectures (SPAA)*, volume 92, pages 420–429, 1992. doi:10.1145/140901.141934.
- 33 Martin Fürer and Shiva Prasad Kasiviswanathan. Spanners for geometric intersection graphs. In *Workshop on Algorithms and Data Structures*, pages 312–324. Springer, 2007.
- 34 Michael T. Goodrich. Planar separators and parallel polygon triangulation. *Journal of Computer and System Sciences*, 51(3):374–389, 1995. doi:10.1006/jcss.1995.1076.
- 35 Lee-Ad Gottlieb. A light metric spanner. In *Proceedings of the 56th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 759–772. IEEE, 2015. doi:10.1109/FOCS.2015.52.
- 36 Joachim Gudmundsson, Christos Levcopoulos, and Giri Narasimhan. Fast greedy algorithms for constructing sparse geometric spanners. *SIAM Journal on Computing*, 31(5):1479–1500, 2002. doi:10.1137/S0097539700382947.
- 37 Lujun Jia, Rajmohan Rajaraman, and Christian Scheideler. On local algorithms for topology control and routing in ad hoc networks. In *Proceedings of the 15th ACM Symposium on Parallel Algorithms and Architectures (SPAA)*, pages 220–229. ACM, 2003. doi:10.1145/777412.777447.
- 38 J. Mark Keil. Approximating the complete Euclidean graph. In Rolf Karlsson and Andrzej Lingas, editors, *Proceedings of the 1st Scandinavian Workshop on Algorithm Theory (SWAT)*, volume 318 of *Lecture Notes in Computer Science*, pages 208–213. Springer, 1988. doi:10.1007/3-540-19487-8_23.
- 39 David Kirkpatrick. Optimal search in planar subdivisions. *SIAM Journal on Computing*, 12(1):28–35, 1983. doi:10.1137/0212002.
- 40 Philip N. Klein, Shay Mozes, and Christian Sommer. Structured recursive separator decompositions for planar graphs in linear time. In *Proceedings of the 45th ACM Symposium on Theory of Computing (STOC)*, pages 505–514. ACM, 2013. doi:10.1145/2488608.2488672.
- 41 Hung Le and Shay Solomon. Truly optimal Euclidean spanners. In *Proceedings of the 60th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 1078–1100. IEEE, 2019. doi:10.1109/FOCS.2019.00069.
- 42 Richard J. Lipton and Robert Endre Tarjan. A separator theorem for planar graphs. *SIAM Journal on Applied Mathematics*, 36(2):177–189, 1979. doi:10.1137/0136016.
- 43 James D. Marble and Kostas E. Bekris. Asymptotically near-optimal planning with probabilistic roadmap spanners. *IEEE Transactions on Robotics*, 29(2):432–444, 2013. doi:10.1109/TRO.2012.2234312.
- 44 Giri Narasimhan and Michiel Smid. *Geometric Spanner Networks*. Cambridge University Press, 2007. doi:10.1017/CB09780511546884.
- 45 David Peleg and Alejandro A. Schäffer. Graph spanners. *Journal of Graph Theory*, 13(1):99–116, 1989. doi:10.1002/jgt.3190130114.
- 46 Liam Roditty and Uri Zwick. On dynamic shortest paths problems. *Algorithmica*, 61(2):389–401, 2011. doi:10.1007/s00453-010-9401-5.

- 47 Wenjie Wang, Cheng Jin, and Sugih Jamin. Network overlay construction under limited end-to-end reachability. In *Proceedings of the 24th Joint Conference of the IEEE Computer and Communications Societies (INFOCOM)*, volume 3, pages 2124–2134. IEEE, 2005.

A

 Proof of some lemmas

In this section we prove Lemma 4 (short-cutting lemma) and Corollary 5 (extended short-cutting) which are known but useful results that we used multiple times in this paper. We also include the proof of Lemma 10 which uses some geometric arguments similar to the proof of Lemma 9.

Proof of Lemma 4. Suppose on the contrary that such points exist. If AB is larger than all other segments $P_i P_{i+1}$, then it should be added the last by the greedy algorithm, so when AB is being added all $P_i P_{i+1}$ s are already included in the spanner, and

$$\sum_{i=0}^{k-1} |P_i P_{i+1}| \leq t \cdot AB$$

By the definition AB should not be added to the graph because there is a path in the spanner with length at most $t \cdot AB$, which contradicts the assumption.

So assume that AB is not larger than all $P_i P_{i+1}$ s. Denote the largest among $P_i P_{i+1}$ s by $P_{i_0} P_{i_0+1}$. Then by the assumption

$$\begin{aligned} \sum_{i \neq i_0} |P_i P_{i+1}| + |AB| &\leq \sum_{i \neq i_0} |P_i P_{i+1}| + |P_{i_0} P_{i_0+1}| \\ &= \sum_i |P_i P_{i+1}| \leq t \cdot |AB| \leq t \cdot |P_{i_0} P_{i_0+1}| \end{aligned}$$

which shows that $P_{i_0} P_{i_0+1}$ can be shortcut by some smaller segments by a factor of t , which is impossible according to what we proved earlier in this lemma. \blacktriangleleft

Proof of Corollary 5. Assume to the contrary that such points exist. For any non-spanner segment $P_i P_{i+1}$ there exists a path $\mathcal{P}(P_i P_{i+1})$ from P_i to P_{i+1} that has length at most $t \cdot |P_i P_{i+1}|$. So by replacing each non-spanner segment $P_i P_{i+1}$ by its own path $\mathcal{P}(P_i P_{i+1})$ in the shortcut path $P_0 P_1 \dots P_k$ the length of the resulting path would be

$$\sum_{P_i P_{i+1} \in S} |P_i P_{i+1}| + \sum_{P_i P_{i+1} \notin S} |\mathcal{P}(P_i P_{i+1})| \leq \sum_{P_i P_{i+1} \in S} |P_i P_{i+1}| + t \cdot \sum_{P_i P_{i+1} \notin S} |P_i P_{i+1}| \leq t \cdot |AB|$$


which shows that the new path is also a shortcut for AB by a factor of t . But the new path only consists of the spanner segments, which is impossible by Lemma 4 and leads to a contradiction. \blacktriangleleft

Proof of Lemma 10. Again, the proof goes by contradiction. Without loss of generality suppose that the projections of P and Q on some baseline l are both between the projections of M and N (on the baseline). We use Lemma 9 to show that MN can be shortcut by PQ by a factor of t , i.e.

$$t \cdot |MP| + |PQ| + t \cdot |QN| \leq t \cdot |MN|$$

The idea is to move PQ by a small amount with respect to its length, so that the new segment intersects MN , and then use Lemma 9. We also keep track of the changes in both sides of the inequality during this movement to show the inequality holds for original points.

Let the segments MN and PQ intersect AB at S and T , respectively. One can move PQ by vector \overrightarrow{TS} in order to intersect MN . Let the new segment be $P'Q'$. But the projections of P' and Q' on the baseline may not be between M and N anymore. In order to preserve



■ **Figure 7** Proof of Lemma 10.

this, we can extend MN on one side by $|\overrightarrow{TS}|$ to get a new segment $M'N'$. Extending by this amount is enough to preserve the betweenness. For example, in Figure Figure 7), we moved PQ by \overrightarrow{TS} to get $P'Q'$. Now $P'Q'$ intersects MN (at S), but the projection of P' on the baseline is not between the projections of M and N anymore. So we extend MN from M by $|\overrightarrow{TS}|$ to get M' . Now the projection of P' on the baseline is between the projections of M' and N . Before the movement the projections of P and Q are both between the projections of M and N , so after movement at most one of the projections of P' or Q' can be outside of the projections of M and N . So extending on one side will be sufficient.

Now $P'Q'$ and $M'N'$ intersect each other and the projections of P' and Q' are between the projections of M' and N' , we can use Lemma 9. By the assumption $\theta = \frac{t'-1}{2t'}$ where $t' = (t+1)/2$, so Lemma 9 implies that,

$$t' \cdot |M'P'| + |P'Q'| + t' \cdot |Q'N'| \leq t' \cdot |M'N'| \quad (5)$$

By the triangle inequality after this movement MP and NQ each will decrease by at most $|\overrightarrow{TS}| \leq |AB|$. So,

$$|M'P'| \geq |MP| - |AB|, |N'Q'| \geq |NQ| - |AB| \quad (6)$$

Also length of MN will increase by at most $|\overrightarrow{TS}| \leq |AB|$, so

$$|M'N'| \leq |MN| + |AB| \quad (7)$$


The length of PQ does not change though. Putting together Equation 5, Equation 6, and Equation 7,

$$\begin{aligned} |PQ| = |P'Q'| &\leq t' \cdot (|M'N'| - |M'P'| - |N'Q'|) \\ &\leq \frac{t+1}{2} \cdot (|MN| - |MP| - |NQ| + 3|AB|) \\ &\leq \frac{t+1}{2} \cdot (|MN| - |MP| - |NQ|) + \frac{t+1}{2} \cdot \left(\frac{t-1}{t(t+1)}|PQ|\right) \end{aligned}$$

So

$$|PQ| \leq t \cdot (|MN| - |MP| - |NQ|)$$

which is the result. ◀



■ **Figure 8** A horizontal zig-zag, and its stretch factor $\Delta y/\Delta x$

B Many intersections with short edges

We proved in sections 3 and 3.5 that, in greedy spanners, each edge has $O(1)$ crossings with edges of greater or equal length. It is natural to ask whether this holds more generally for all crossings, regardless of length. That is, is the total number of crossings for each edge bounded by a constant, depending only on t ? In this section we will show that this is not true, by constructing a family of arrangements of points in the plane that have arbitrarily many intersections between a long edge and a set of smaller edges.

B.1 Zig-zags

The building block of our construction is an arrangement of points which form a zig-zag shape, as in Figure 8. After running the greedy spanner algorithm on a horizontal zig-zag like this, denoted by Z , if Z is not stretched too much along the vertical axis, the first set of edges that will be added to the graph by the greedy algorithm are actually the zig-zag edges that are drawn in Figure 8. Then, depending on the shape of the zig-zag and parameter t , other edges may or may not be added in the future iterations. More specifically, we will show that this only depends on a parameter we call the *stretch-factor* of the zig-zag.

► **Definition 22** (zig-zag). Let $Z = P_0P_1 \dots P_k$ be a sequence of points on the Euclidean plane. We say Z forms a Zig-Zag if there exist two perpendicular vectors $\vec{\Delta x}$ and $\vec{\Delta y}$ that

$$P_i = P_0 + i\vec{\Delta x} + (i \bmod 2)\vec{\Delta y}$$

The direction of the vector $\vec{\Delta x}$ is called the direction of the zigzag and the ratio $|\vec{\Delta y}|/|\vec{\Delta x}|$ is called the stretch factor of the zig-zag, and is denoted by $s(Z)$. (Figure 8)

Hence a zig-zag which is more stretched toward the $\vec{\Delta y}$ vector will have a larger stretch-factor, and a zig-zag which is more stretched along the $\vec{\Delta x}$ vector will have a smaller stretch-factor.


► **Lemma 23** (zig-zag spanner). Consider a zig-zag $Z = P_0P_1 \dots P_k$ with more than two vertices ($k > 2$) in which the consecutive pairs P_iP_{i+1} are connected to each other ($0 \leq i < k$). For any $t > 1$, the zig-zag forms a t -spanner if and only if $s(Z) \leq \sqrt{t^2 - 1}$.

Proof. For $i < j$, the length of the path between P_i and P_j is

$$d_Z(P_i, P_j) = (j - i)|\vec{\Delta x} + \vec{\Delta y}|$$

while the Euclidean distance between P_i and P_j is

$$d(P_i, P_j) = |(j - i)\vec{\Delta x} + (j - i \bmod 2)\vec{\Delta y}|$$



■ **Figure 9** Example with more than constant intersections with smaller edges

The zig-zag forms a t -spanner if and only if $d_Z(P_i, P_j) \leq t \cdot d(P_i, P_j)$ for all $i < j$. Assume that $(j - i) \bmod 2 = 0$, this inequality turns into

$$(j - i)|\vec{\Delta x} + \vec{\Delta y}| \leq t \cdot (j - i)|\vec{\Delta x}|$$

which is equivalent to $s(Z) \leq \sqrt{t^2 - 1}$. So this is a necessary condition, and it can be shown that it is a sufficient condition too. Because assuming $s(Z) \leq \sqrt{t^2 - 1}$, in a similar way,

$$(j - i)|\vec{\Delta x} + \vec{\Delta y}| \leq t \cdot (j - i)|\vec{\Delta x}|$$


The left side of the inequality is $d_Z(P_i, P_j)$ and the right side is no more than $t \cdot d(P_i, P_j)$ because it is missing the term $(j - i \bmod 2)\vec{\Delta y}$, so $d_Z(P_i, P_j) \leq t \cdot d(P_i, P_j)$. ◀

B.2 Introducing the arrangement


Now we introduce the arrangement. Consider two horizontal zig-zags U on the top and B on the bottom which are connected together using a middle zig-zag M (Figure 9). U is colored by green, B is colored by blue, and M is colored by red. So there are four rows of points and three zig-zags U , M , and B , which connect these points together. U and M share the second row, while M and B share the third row. The first row is only included in U , and the last row is only included in B . For now, suppose that there are enough points in each row. Later we will see that if the number of points is larger than a specific amount, then a large edge appears at some point in the greedy algorithm, intersecting many edges in between.

All of the zig-zags U , M , and B can have arbitrary stretch-factors as we can move the rows up or down to adjust the stretch-factor of each zig-zag independently. So assume that $s(U) = s(B) = \sqrt{t^2 - 1}$ and $s(M) = \sqrt{(t + \delta)^2 - 1}$, for some small positive δ which will be specified later. In other words, U and B are the most stretched zig-zags that form a t -spanner and M is a slightly more stretched zig-zag, which is not a t -spanner by itself anymore.

With this choice of stretch-factors, it is not hard to see, by the Pythagorean theorem, that the length of the zig-zag path between two points on U , say a and b , is exactly $t \cdot |x_a - x_b|$. And the length of the path between two points on B is also the same expression. But in a similar way, the length of the zig-zag path between two points on M would be slightly more, $(t + \delta) \cdot |x_a - x_b|$.



■ **Figure 10** Vertical dashed segments are included in the graph but not horizontal ones.



■ **Figure 11** The big dashed zig-zag might be included in the graph or might not.


B.3 Simulating the greedy algorithm on the arrangement

For an appropriate choice of t (one causing the angles of all zig-zags to lie strictly between 60° and 120°), the greedy spanner algorithm will first add the zig-zag edges in U , B , and M , as they are the closest pairs of vertices. According to the chosen stretch-factors, no edges will be added to U and B in the future. For example, the horizontal dashed blue edges in Figure 10 will not be added as the endpoints of these segments both belong to U or B , which are t -spanners by themselves. So any potential edge must be between U and B .

The next set of edges that may be added by the algorithm are the vertical edges between rows 1 and 3, and 2 and 4 (red dashed segments in Figure 10). These are the closest pairs across U and B which are not connected, so they will be included first. The edges between rows 1 and 4 which connect the points in consecutive columns (dashed blue segments in Figure 11) may also be added in the next iteration, depending on how small the value of t is, but we will see that they do not affect the length of the shortest paths between pairs of points in U and B that much.

B.4 Sufficiency of small edges for close pairs

Now we claim that the edges we found until now are the only local, i.e. small, edges between these points, and the next edge that is going to be added by the greedy algorithm, would be a large one which intersects many of the zig-zag edges in M . The greedy algorithm may stop here and do not add any edges, but we will prove later that this is not possible. We are not



■ **Figure 12** $P(u, b)$, which uses some of the edges in U and B and only one edge in M . Here $i = 1$ and $j = 7$.

going to address this issue in this section.

Intuitively, one can use edges in U and B , and only one edge in M to build a path from any point in U to any point in B (see Figure 12). Again, intuitively, zig-zags are defined in a way that the length of this path is more than $t \cdot |x_u - x_b|$ by a small constant. But when u and b are not far away $|x_u - x_b|$ is much less than $d(u, b)$ and hence the length of the path is no more than $t \cdot d(u, b)$. On the other hand, when u and b are far away, $|x_u - x_b|$ is closer than any constant to $d(u, b)$ (because here $|y_u - y_b|$ is bounded), hence the length of the path becomes more than $t \cdot d(u, b)$ and a long edge appears.

In order to prove this formally, as stated above, any potential edge must be between U and B . So let $u \in U$ and $b \in B$ be two arbitrary points in the top and the bottom zig-zags, respectively. Also assume that u is the i -th point in U ($i = 0, 1, \dots$), and b is the j -th point in B ($j = 0, 1, \dots$), counting from left (Figure 12).

We assume that no edges other than the ones we stated above have been added so far, and we compute the length of a path we propose between u and b that uses these edges and we show that it is less than $t \cdot d(u, b)$ if $d(u, b)$ is not very large. In this way, we prove that the next edge which is going to be added would be a large one.

Without loss of generality, assume that $i \leq j$. Consider a path that uses zig-zag edges of U and B and only one of the edges in M to reach from u to b . Denote this path by $P(u, b)$. Such a path is drawn by a red dashed line for two sample points in Figure 12. Clearly, we do not use any edge twice and we only use zig-zag edges in U , B , or M .

We will show that $|P(u, b)|$, the length of the red path, is not more than $t \cdot d(u, b)$ when $d(u, b)$ is not very large. By the definition, $P(u, b)$ uses $j - i - 1$ edges of U and B , and one edge in M , so

$$|P(u, b)| = (j - i - 1)l + l' \quad (8)$$

where l is the edge length in U (and B), and l' is the edge length in M . On the other side, the distance along the x -axis between u and b is $(i - j)\Delta x$, where Δx is defined in Definition 22. The distance along the y -axis between u and b is at least the height of the zig-zag M , which is by the definition $s(M)\Delta x$. This distance can be strictly more than $s(M)\Delta x$ when u is in the first row or b is in the last row. So,

$$d(u, b) \geq \sqrt{(j - i)^2(\Delta x)^2 + s(M)^2(\Delta x)^2} = \sqrt{(j - i)^2 + (t + \delta)^2 - 1}\Delta x \quad (9)$$

In order to show $|P(u, b)| \leq t \cdot d(u, b)$, we use Equation 8 and Equation 9 to show $|P(u, b)|^2 -$

$t^2 \cdot d(u, b)^2$ is non-positive,

$$\begin{aligned} |P(u, b)|^2 - t^2 \cdot d(u, b)^2 &\leq [(j-i-1)l + l']^2 - [(j-i)^2 + (t+\delta)^2 - 1] (t\Delta x)^2 \\ &= [(j-i) + (l'/l - 1)]^2 l^2 - [(j-i)^2 + (t+\delta)^2 - 1] l^2 \\ &= [2(j-i)(l'/l - 1) + (l'/l - 1)^2 - (t+\delta)^2 + 1] l^2 \end{aligned}$$

We used $t\Delta x = l$ in the first equality. Now by putting $l'/l = \frac{t+\delta}{t}$, when $j-i \leq t(t^2-1)/(2\delta)$,

$$\begin{aligned} |P(u, b)|^2 - t^2 \cdot d(u, b)^2 &\leq \left[2(j-i)\frac{\delta}{t} + \left(\frac{\delta}{t}\right)^2 - (t+\delta)^2 + 1 \right] l^2 \\ &\leq [(t^2-1) + \delta^2 - (t+\delta)^2 + 1] l^2 \leq 0 \end{aligned}$$

So no edge is required between u and b and if there is any edge between them, it must be the case that $j-i > t(t^2-1)/(2\delta)$. On the other side, the edge (u, b) , if exists, will intersect at least $j-i-2$ of the zig-zag edges which separate u and b . So one can choose δ to be sufficiently small to increase the number of intersections.

B.5 Existence of a large edge

Now we address the issue we mentioned earlier, that the greedy algorithm may stop after adding the small edges we discussed in section 3.3 and never add any large edges. We need to prove the existence of such a large edge to complete the proof.

Again, let u be the i -th point in U and b be the j -th point in B , counting from left. We will show that when $j-i$ is large enough an edge is required between u and b . None of the edges that we mentioned so far connects two points whose x -distance is more than Δx . So the shortest path between u and b , denoted by $P^*(u, b)$, needs at least $j-i$ edges to reach from u to b . At least one of these edges should be across U and B , hence having a length at least l' . The other edges have lengths of at least l , as it is the smallest edge in the graph. Thus

$$|P^*(u, b)| \geq (j-i-1)l + l' \quad (10)$$

Again, the x -distance of u and b is $(i-j)\Delta x$, and the y -distance of them is at most the height of the whole figure, which is the sum of the height of the three zig-zags, $(s(U) + s(M) + s(B))\Delta x$. So,

$$\begin{aligned} d(u, b) &\leq \sqrt{(j-i)^2(\Delta x)^2 + (s(U) + s(M) + s(B))^2(\Delta x)^2} \\ &\leq \sqrt{(j-i)^2 + (3s(M))^2} \Delta x = \sqrt{(j-i)^2 + 9(t+\delta)^2 - 9} \Delta x \end{aligned} \quad (11)$$

The second inequality follows from the fact that $s(M)$ is the maximum among $s(U)$, $s(M)$, and $s(B)$. Similarly, we use Equation 10 and Equation 11 to show that $|P(u, b)|^2 - t^2 \cdot d(u, b)^2$ is positive,

$$|P^*(u, b)|^2 - t^2 \cdot d(u, b)^2 \geq \left[2(j-i)\frac{\delta}{t} + \left(\frac{\delta}{t}\right)^2 - 9((t+\delta)^2 - 1) \right] l^2$$

When $j-i \geq 9t((t+\delta)^2 - 1)/(2\delta)$,

$$|P^*(u, b)|^2 - t^2 \cdot d(u, b)^2 \geq \left[9((t+\delta)^2 - 1) + \left(\frac{\delta}{t}\right)^2 - 9((t+\delta)^2 - 1) \right] l^2 > 0$$

Hence the result.

► **Theorem 24.** *For some values of t , there is no constant bound (depending only on t) on the number of crossings between an edge of a greedy t -spanner and other smaller edges.*

Proof. This follows from the existence of the example above. ◀