

PPN rotation curves in static distributions with spherical symmetry

Henrique Matheus Gaury^{1,*} and Javier Ramos-Caro^{1,†}

¹*Departamento de Física, Universidade Federal de São Carlos, São Carlos, 13565-905 SP, Brazil*

(Dated: January 1, 2022)

From a Parametrized Post-Newtonian (PPN) [46–51] perspective, we address the question of whether or not the new degrees of freedom, represented by the PPN potentials, can lead to significant modifications in the dynamics of galaxies in the direction of rendering dark matter obsolete. Here, we focus on the study of rotation curves associated with spherically symmetric configurations.

The values for the post-Newtonian parameters, which help us to classify the different metric theories of gravity, are tightly constrained mainly by solar system experiments [47, 48]. Such restrictions renders the modifications of gravitational effects, with respect to General Relativity (GR), to be insignificant, making attempts to find alternative metrical theories rather fruitless. However, in recent years, metric theories characterized by screening mechanisms [4, 10, 22, 25] have become popular, due to the fact that they lead to the possibility of modifications in larger scales than the solar system while retaining the success of GR on it, allowing for violations of the constraints of the Post-Newtonian parameters.

In such a context, we consider here two kinds of solutions for field equations: (i) Vacuum solutions (i.e. when no matter fields are present) and (ii) fields in the presence of a polytropic distribution of matter. For the case (i) we find that the post-Newtonian corrections do not lead to modifications significant enough to be considered an alternative to the dark matter hypothesis. In the case (ii) we find that for a wide range of values for the PPN parameters γ , β , ξ , α_3 , ζ_1 and ζ_2 , the need for dark matter is unavoidable, in order to find flat rotation curves. It is only for theories in which $\zeta_3 > 0$ that some resemblance of flat rotation curves is found. The latter suggests, at least for the models considered, that these are the only theories capable of replacing dark matter as a possible explanation for the dynamics of galaxies.

PACS numbers: 04.20.-q, 95.30.Sf, 98.20.-d

I. INTRODUCTION

The success of the Theory of General Relativity (TGR) lies both in its conceptual beauty and in the fact of adjusting satisfactorily to a considerable amount of observations. In fact, it is widely known that TGR has among the best comparison with experiment of modern science. In the solar system, mediated by the Parametrized Post-Newtonian formalism (PPN), it agrees significantly with observations [46–51], and, in more recent years, it has been shown to be in agreement with gravitational radiation experiments, as measured by Ligo [27]. However, despite the great success of TGR, there are still big questions to be solved as, for example: Is General Relativity valid in galactic scales?

Vera Rubin et al. introduced the idea, via galactic rotation curves, that visible matter is insufficient to explain the internal dynamics of a number of galaxies [8, 9]. This is known as the “missing mass problem”, the central evidence that some form of dark matter must exist in galaxies. Some other indirect evidences such as the stability of galaxies, galaxy cluster dynamics, cosmological structure formation and cosmic microwave background anisotropies all point out to the existence of such dark component [1, 7, 26, 45],

However, no direct evidence about the existence of dark matter is known to this day [15–18]. This fact, on the other hand, motivates the searching for different explanations to the missing mass problem, as the formulation of alternative theories of gravity capable of making obsolete the dark matter hypothesis. In the literature there are several proposals for modifying gravity, most with the intent of explaining the phenomenology of cosmological regimes, mainly the accelerated expansion (see for example [4, 12, 48]). There exist theories with some success as alternatives to dark matter at galactic scales. Perhaps the most notorious is Mond (Modified Newton Dynamics)[28–30, 32, 33, 35], which proposes that the gravitational force behaves differently below a fundamental acceleration. But as successful as it is at explaining the rotation curves of various galaxies and the Tully-Fisher relation [24, 41, 43, 44], recent developments have made this interesting possibility highly unlikely [2].

Changing the foundations of a fundamental theory such as gravity is no easy feat. The vast space of possible competing theories makes treating this problem, case by case, an unsurmountable task. A more economic way is to have on hand a formalism containing, as particular cases, a large amount of theories. Based on this idea, several general schemes have been constructed, and the Parameterized Post-Newtonian (PPN) formalism, by Will et al [46–51], is probably the most used. It generalizes the Post-Newtonian expansion for metrical theories of gravity by the introduction of 10 parameters (i.e. the so called post-Newtonian parameters) which help us to

*henmgaury@df.ufscar.br

†javier@ufscar.br

differentiate between the competing theories. In the last decades, highly precise experiments in the solar system have lead to ever increasingly smaller bounds for the PPN parameters (some being at most 10^{-20} [48]), leading to discard most alternative theories.

However, recent developments suggest that this may not be the final word for a number of alternative metrical theories. The existence of *screening mechanisms* (SM) allows these theories to pass experiments in solar scales without maiming its deviations from General Relativity at larger scales [4, 10, 22, 25]. These theories, with some SM, cannot be perturbatively expanded from infinity all the way to the Schwarzschild radius [10], a basic tenet of the classical perturbation theory (and, therefore, of the PPN approach). This fact leads to different predictions at solar system scales, than the ones by the PPN formalism. The reason for this is that the failures of the perturbative expansion only happen within a screening radius r_V [10], outside of which the classical perturbation theory becomes applicable. Therefore, in accordance with [10], at least as long we are far from r_V , the linear theory should be the PPN approach.

Not all theories with some SM can be used to justify modifications at the astrophysical scale. Take for instance the cubic galileon [10], which predicts the screening for all scales smaller than galaxy clusters, therefore excluding astrophysical modifications. On the other hand, the problem on the existence of SMs that allows modifications at the astrophysical scale, is a question addressed by [14, 40], showing that some SM mimic the effects of dark matter in galaxy clusters and in our own galaxy.

In this paper we assume the existence of some SM capable of hiding the modifications to TGR in some scale (say, of the order of the solar system), but keeping the modifications to TGR at the scale of galaxies or galaxy clusters (i.e. the existence of a theory decoupling solar system from astrophysical scales). As long as the analysis is restricted to outside and far from r_V , the perturbative expansion should be the PPN scheme [10]. In the face of such prerogative it is pertinent to raise some questions: what are the corrections to the rotation curves that arise as consequences of the PPN potentials? Are these corrections enough to explain the missing mass problem in galaxies?

For the sake of simplicity, one can start to address this problem by obtaining the PPN potentials generated by static configurations with spherical symmetry, in order to obtain an expression for the corresponding circular velocity, as done in section IV. There we will find that the existence of circular motion (and, therefore, the existence of rotation curves) depends strongly of the vanishing of two PPN parameters. In particular, it is required that $\alpha_1 = \alpha_2 = 0$. In the same section we also study the behavior of circular velocity outside a sphere with finite radius, finding that no choice of the PPN parameters can lead to the flattening of the rotation curves. Away from the spherical distribution, rotation curves exhibit

the usual Keplerian fall off.

In section V we analyze the behavior of circular velocity inside static spherical distributions, focusing on the case of the so-called polytropes, one of the simplest models with some relevance in galactic dynamics. To construct polytropic models in the PPN approximation we use the “ f to ρ ” approach of galactic dynamics [11, 38, 39], in which the matter distribution (given by mass density, ρ , in Newtonian gravity) is obtained from a known distribution function f . In the framework of metrical theories, and thereof the PPN scheme, we assume that the distribution function (DF) satisfies a generalized version of the collisionless Boltzmann equation (CBE) [53], also called the Vlasov equation. This assumption is valid whenever the system in study is sufficiently smooth and encounters can be disregarded, as in the case of galaxies [11]. Then, as a first step we will derive a version of the CBE that accounts for the first PPN corrections, in a similar fashion as in [38], focusing on stationary solutions. To construct the Polytropes, by extension, we will provide an ergodic DF (i.e. depending on energy) with the same form as in the Newtonian case, determining all the corresponding matter fields.

In section V A we show that, for polytropic models, only one of the PPN parameters ($\zeta_3 > 0$) can effectively lead to flattened rotation curves within a certain radius ($\tilde{r} < 10$). Outside such region, the usual Keplerian fall off is still observed, even for the most exotic theories. Everything seems to indicate that, in static spherical distributions, the modifications encompassed by the PPN scheme are not enough to explain the flatness of rotation curves without the introduction of additional matter.

The rest of the paper is organized as follows. In Sec. II we present a brief overview of the PPN formalism, pointing out it’s most notable features. In Subsec. II A we introduce the material content we will consider in the modeling. In Sec. III we finally specialize to the static spherically symmetric case presenting the field equations. In Sec. V A we construct the polytropic models and their respective rotations curves for various theories.

Throughout the paper, we will regard Latin indices to run from 1 to 3, i.e. $i = \{1, 2, 3\}$, and Greek indices to run from 0 to 3, i.e. $\mu = \{0, 1, 2, 3\}$. We will also use, whenever suited, the notation $\partial f / \partial x = f_{,x}$. Terms of different orders of c will carry an indice as follows,

$${}^n A \equiv {}^n A \sim c^{-n},$$

this means, ${}^n A$ is of the order of c^{-n} .

II. THE PPN FORMALISM

There is a well known approximation scheme for general relativity when we consider weak fields and slow moving matter, known as the post-Newtonian approximation. In the context of alternative metric theories of

gravity there is a general formalism developed by Will and Nordtvedt [46–51], known as the Parametrized Post-Newtonian (PPN) approximation, that find a similar approximation scheme, but introducing parameters that are different for each theory.

In this formalism the metric is determined by perturbations of a flat background in terms of a parameter $\epsilon \ll 1$, such that

$$\epsilon \sim v^2/c^2 \sim GM/rc^2 \sim p/\rho c^2,$$

where v , M , r , p and ρ are the characteristic velocity, mass, length (or separation), pressure and density in the system, and G is the (Newtonian) gravitational constant. At first approximation, the metric can be written as [47]

$$g_{00} = -1 + \frac{2U}{c^2} + \frac{\mathcal{W}}{c^4} + O(c^{-6}), \quad (1)$$

$$g_{0j} = \frac{\mathcal{Q}_j}{c^3} + O(c^{-5}), \quad (2)$$

$$g_{jk} = \left(1 + \frac{2\gamma U}{c^2}\right) \delta_{jk} + O(c^{-4}), \quad (3)$$

where U reduces to the Newtonian potential in the limit $c \rightarrow \infty$ and functions \mathcal{W} , \mathcal{Q}_j are defined in terms of post-Newtonian potentials Ψ , X , U_j ($j = 1, 2, 3$), ϕ_k ($k = 1, \dots, 6$), ϕ_w , Φ^{PF} , Φ_j^{PF} , as

$$\mathcal{W} \equiv 2(\Psi - \beta U^2) + \Phi^{\text{PF}}, \quad (4)$$

$$\begin{aligned} \mathcal{Q}_j &\equiv -\left[2(1 + \gamma) + \frac{\alpha_1}{2}\right] U_j \\ &\quad - \frac{1}{2}[1 + \alpha_2 - \zeta_1 + 2\xi] X_{,tj} + \Phi_j^{\text{PF}}, \end{aligned} \quad (5)$$

$$\begin{aligned} \Psi &= \frac{1}{2}(2\gamma + 1 + \alpha_3 + \zeta_1 - 2\xi) \phi_1 \\ &\quad - (2\beta - 1 - \zeta_2 - \xi) \phi_2 \\ &\quad + (1 + \zeta_3) \phi_3 + (3\gamma + 3\zeta_4 - 2\xi) \phi_4 \\ &\quad - \frac{1}{2}(\zeta_1 - 2\xi) \phi_6 - \xi \phi_w, \end{aligned} \quad (6)$$

and γ , β , ξ , α_1 , α_2 , α_3 , ζ_1 , ζ_2 , ζ_3 , ζ_4 are the post-Newtonian parameters.

Usually, in the PPN scheme the material content is assumed to be a perfect fluid described by an energy momentum tensor of the form [47]

$$c^{-2}T^{00} = \rho^* \left\{ 1 + \frac{1}{c^2} \left[\frac{1}{2}u^2 - (3\gamma - 2)U + \Pi \right] \right\}, \quad (7)$$

$$c^{-1}T^{0j} = \rho^* u^j \left\{ 1 + \frac{1}{c^2} \left[\frac{1}{2}u^2 - (3\gamma - 2)U + \Pi + \frac{p}{\rho^*} \right] \right\}, \quad (8)$$

$$\begin{aligned} T^{ij} &= \rho^* u^i u^j \left\{ 1 + \frac{1}{c^2} \left[\frac{1}{2}u^2 - (3\gamma - 2)U + \Pi + \frac{p}{\rho^*} \right] \right\} \\ &\quad + p \left(1 - \frac{2\gamma U}{c^2} \right) \delta^{ij}, \end{aligned} \quad (9)$$

where p is the pressure field, u^k is the velocity field, $\Pi = \varepsilon/\rho^*$ (ε is the internal energy) and ρ^* is the conserved density, which is related to the proper mass density ρ through the relation

$$\rho = \left[1 - \frac{1}{c^2} \left(\frac{1}{2}u^2 + 3\gamma U \right) \right] \rho^*.$$

Such an assumption implies that potentials U , ϕ_1 , ϕ_2 , ϕ_3 , ϕ_4 , ϕ_6 , X , U_j and ϕ_w are determined by the field equations,

$$\nabla^2 \{U, \phi_1, \phi_2, \phi_3, U_j\} = -4\pi G \rho^* \{1, u^2, U, \Pi, u^j\} \quad (10)$$

$$\nabla^2 \phi_4 = -4\pi G p, \quad \nabla^2 X = 2U \quad (11)$$

$$\nabla^4 (\phi_6 - 3\phi_1) = -2G (\rho^* u^i u^j)_{,ij} \quad (12)$$

$$\nabla^2 (\phi_w + 2U^2 - 3\phi_2) = -2X_{,ij} U_{,ij}, \quad (13)$$

whereas the preferred-frame potentials, Φ^{PF} and Φ_j^{PF} , can be written in terms of the velocity of the PPN co-ordinate frame relative to an (hypotetical) universal preferred frame, denoted by w^j :

$$\begin{aligned} \Phi^{\text{PF}} &= (\alpha_3 - \alpha_1) w^2 U + \alpha_2 w^j w^k X_{,kj} \\ &\quad + (2\alpha_3 - \alpha_1) w^j U_j, \end{aligned} \quad (14)$$

$$\Phi_j^{\text{PF}} = -\frac{1}{2} \alpha_1 w_j U + \alpha_2 w^k X_{,kj}. \quad (15)$$

All of the above fields are defined in a 3-dimensional Euclidean space and, in particular, equations (12)-(15) are stated in (Quasi-) Cartesian coordinates [34, 47]. Later on, when we address situations involving spherical symmetry, it will be useful to represent the previous equations in non-Cartesian coordinates. As long as the coordinate transformation does not involve time or velocities (changes of reference frame), we can regard it as a diffeomorphism in Euclidean space which does not require to consider the PPN metric (1)-(3). For this reason it is not difficult to find the covariant form of relations (12)-(15), by switching every common derivative by a covariant derivative and by introducing the Euclidean metric γ_{ab} at every inner product:

$$\nabla^2 (\phi_w + 2U^2 - 3\phi_2) = -2\gamma^{ad} \gamma^{be} X_{,ab} U_{;de}, \quad (16)$$

$$\begin{aligned} \Phi^{\text{PF}} &= (\alpha_3 - \alpha_1) w^2 U + \alpha_2 w^j w^k X_{;kj} \\ &\quad + (2\alpha_3 - \alpha_1) w^j U_j, \end{aligned} \quad (17)$$

$$\Phi_j^{\text{PF}} = -\frac{1}{2} \alpha_1 w_j U + \alpha_2 w^k X_{;kj}, \quad (18)$$

$$\nabla^4(\phi_6 - 3\phi_1) = -\frac{2G}{\mathcal{R}}(\mathcal{R}\rho^*u^iu^j)_{,ij} - \frac{2G}{\mathcal{R}}(\mathcal{R}\{_{ik}^j\rho^*u^iu^k\}_{,j}, \quad (19)$$

where $\mathcal{R} = \sqrt{\det(\gamma_{ij})}$ and

$$X_{;ab} \equiv X_{,ab} - \{_{ab}^c\} X_c, \quad (20)$$

$$\{_{ab}^c\} \equiv \frac{1}{2}\gamma^{c\mu}(\gamma_{a\mu,b} + \gamma_{b\mu,a} - \gamma_{ab,\mu}) \quad (21)$$

On the other hand, the equations of motion for a free-falling test particle, can be written in terms of the fields appearing in the above expressions:

$$\begin{aligned} \frac{dv^i}{dt} + \{_{jk}^i\} v^j v^k = \gamma^{ik} & \left\{ U_{,k} + \frac{1}{c^2} \left[\frac{1}{2} \mathcal{W}_{,k} - \mathcal{Q}_{k,t} \right. \right. \\ & - 2\gamma U U_{,k} - (\mathcal{Q}_{k,j} - \mathcal{Q}_{j,k}) v^j - (2\gamma + 1) v_k U_{,t} \\ & \left. \left. - 2(\gamma + 1) U_{,j} v^j v_k + \gamma U_{,k} v^2 \right] \right\}. \end{aligned} \quad (22)$$

By choosing the material fields ρ^* , $\rho^* \Pi$, P , $\rho^* u^2$, $\rho^* u^i$ and $\rho^* u^i u^j$ it should be possible, in principle, to solve (10), (11) and (16)-(19) and, in consequence, determine the test particle motion through equations (22).

A. Statistical description of collisionless systems

The study of huge astrophysical ensembles such as globular clusters, galaxies and galaxy clusters, is simplified by adopting a statistical description and introducing some assumptions about the configurations: (i) the particles of the system have the same mass; (ii) collisions or encounters between particles are insignificant; (iii) the gravitational fields are all regarded to be smooth and continuous throughout space. Thus the system can be described entirely by a probability density or distribution function (DF), that represents the number density of particles of a point (x^i, V^i) in the phase-space (here V^i represents the four-velocity). Assumptions (i) and (ii) lead us to consider that the DF, denoted here by $f(x^i, V^i)$, satisfy the general-relativistic Vlasov equation,

$$V^\mu \frac{\partial f}{\partial x^\mu} - \Gamma_{\mu\nu}^i V^\mu V^\nu \frac{\partial f}{\partial V^i} = 0. \quad (23)$$

In order to find equation (23) to PPN order we follow a similar procedure as in [38], which take into account the map $(x^\mu, V^i) \rightarrow (x^\mu, v^i(x^\mu, V^i))$ and perform an expansion of the left hand side of (23) to post-Newtonian order. After some calculations, we find the Vlasov equation in

the PPN approach (see Appendix B):

$$\begin{aligned} & \left[1 + \frac{U}{c^2} + \frac{v^2}{2c^2} \right] \left(\frac{\partial f}{\partial t} + v^i \frac{\partial f}{\partial x^i} \right) + U_{,i} \frac{\partial f}{\partial v^i} \\ & - \frac{\partial f}{\partial v^i} \frac{v^i}{c^2} U_{,t} (1 + 2\gamma) - \frac{\partial f}{\partial v^i} \frac{v^i v^j}{c^2} U_{,j} (2 + 2\gamma) \\ & - \frac{\partial f}{\partial v^i} \left\{ -\frac{1}{2} \frac{\mathcal{W}_{,i}}{c^2} + \frac{\mathcal{Q}_{i,t}}{c^2} + \frac{2\gamma - 1}{c^2} U U_{,i} \right\} \\ & - [\mathcal{Q}_{j,l} - \mathcal{Q}_{l,j}] \frac{v^l}{c^2} \frac{\partial f}{\partial v^j} + \left(\gamma + \frac{1}{2} \right) U_{,j} \frac{v^2}{c^2} \frac{\partial f}{\partial v^j} = 0. \end{aligned} \quad (24)$$

From (24) one can prove that

$$\frac{df}{dt} = 0,$$

which means that any stationary solution of equation (24) must be a function only of the integrals of motion of the system and vice versa: any function of the integrals of motion is a solution of (24).

From the DF we can build its moments, being the most relevant ones, for this work, the components of the stress-energy tensor [52]:

$$T^{\mu\nu} = c \int \frac{V^\mu V^\nu}{-V_0} f \sqrt{-g} d^3V.$$

Since we want the above expression in PPN approximation, we again consider the map $(x^\mu, V^i) \rightarrow (x^\mu, v^i(x^\mu, V^i))$ and the expansion

$$f = f^0 + f^2 + \dots,$$

which lead us to the following relations:

$$c^{-2} T^{00} = \int f d^3v + \int \left(\frac{\mathcal{K}^0}{c^2} f + f^2 \right) d^3v, \quad (25)$$

$$c^{-1} T^{0k} = \int v^k f d^3v + \int v^k \left(\frac{\mathcal{K}^0}{c^2} f + f^2 \right) d^3v, \quad (26)$$

$$T^{kj} = \int v^k v^j f d^3v + \int v^k v^j \left(\frac{\mathcal{K}^0}{c^2} f + f^2 \right) d^3v, \quad (27)$$

where

$$\mathcal{K} = U(3\gamma + 5) + 3v^2.$$

Comparing (25)-(27) with (7)-(9) we can, at least in principle, find all the matter fields (ρ^* , $\rho^* \Pi$, P , $\rho^* u^2$, $\rho^* u^i$ and $\rho^* u^i u^j$) of equations (10)-(19), once we know f .

III. PPN POTENTIALS FOR STATIC CONFIGURATIONS WITH SPHERICAL SYMMETRY

When we are dealing with static spherically symmetric systems, several assumptions are needed. First we require matter to be static, i.e. with a velocity field $u^i = 0$

or, in other words, with a DF such that $T^{0k} = 0$. This means that equations (7)-(9) are simplified:

$$c^{-2}T^{00} = \rho^* \left\{ 1 + \frac{1}{c^2} [\Pi - (3\gamma - 2)U] \right\},$$

$$c^{-1}T^{0j} = 0,$$

$$2T^{ij} = P\delta^{ij}.$$

By comparing the previous equations with (25)-(27), we find the expression of the matter fields as functionals of the DF,

$$\rho^* = 4\pi \int_0^0 f v^2 d^3v, \quad (28)$$

$$P = \frac{4\pi}{3} \int_0^0 f v^4 d^3v, \quad (29)$$

$$\rho^* \Pi = (3\gamma - 2) \rho^* U + 4\pi \int \left(f k + \frac{2}{c^2} \right) v^2 d^3v. \quad (30)$$

We also require that the fields U , ϕ_1 , ϕ_2 , ϕ_3 , ϕ_4 , ϕ_6 , X , U^j and ϕ_w are all static, spherically symmetric and well behaved throughout the space. As a consequence of eqs. (10) and the assumption that the space-time is asymptotically flat, the fields U^j and ϕ_1 can be written as

$$U^j = \frac{A^j}{r}, \quad \phi_1 = \frac{B}{r},$$

for A^j and B constants (we have introduced spherical coordinates (r, θ, φ)). Since we require that the fields are well behaved for all r , we have to choose $A^j = 0$ and $B = 0$, in order to avoid the singularity at $r = 0$, which means that $U^j = \phi_1 = 0$. Now, from equation (12), we have $\phi_6 = 3\phi_1$, leading to $\phi_6 = 0$.

Under such assumptions the field equations take a much simpler form:

$$\nabla^2 U = -4\pi G \rho^*, \quad (31)$$

$$\nabla^2 X = 2U, \quad (32)$$

$$\begin{aligned} \nabla^2 \Psi &= (2\beta - 1 - \zeta_2 - 6\xi) 4\pi G \rho^* U \\ &\quad - (1 + \zeta_3) 4\pi G \rho^* \Pi - (3\gamma + 3\zeta_4 - 2\xi) 4\pi G P \\ &\quad + 4\xi \left[4\pi G \rho^* \frac{X_{,r}}{r} + U_{,r} \left(U_{,r} + \frac{3X_{,r}}{r^2} - \frac{2U}{r} \right) \right], \end{aligned} \quad (33)$$

$$\begin{aligned} \Phi^{\text{PF}} &= w^2 \left[(\alpha_3 - \alpha_1) U + \alpha_2 \frac{X_{,r}}{r} \right] \\ &\quad + \alpha_2 (w^r)^2 \left(X_{,rr} - \frac{X_{,r}}{r} \right), \end{aligned} \quad (34)$$

$$\Phi_r^{\text{PF}} = -\frac{1}{2} \alpha_1 w_r U + \alpha_2 w^r X_{,rr}, \quad (35)$$

$$\Phi_\theta^{\text{PF}} = -\frac{1}{2} \alpha_1 w_\theta U + r \alpha_2 w^\theta X_{,r}, \quad (36)$$

$$\Phi_\varphi^{\text{PF}} = -\frac{1}{2} \alpha_1 w_\varphi U + \alpha_2 w^\varphi r \sin^2 \theta X_{,r}. \quad (37)$$

Note that the assumptions about symmetry's configuration (spherical, in our case) must be the same for the fields U , X and Ψ , as a consequence of eqs. (31)-(33). On the other hand, according to (34)-(37), the preferred frame potentials, $(\Phi^{\text{PF}}, \Phi_r^{\text{PF}}, \Phi_\theta^{\text{PF}}, \Phi_\varphi^{\text{PF}})$, are not necessarily constrained to satisfy the same symmetry assumptions of configuration. In fact, in our case all of them would depend on the angular variables (θ, φ) for the case in which α_1 and α_2 are non-vanishing and assumptions are not made about the preferred frame velocity \vec{w} .

In principle, by introducing a particular DF in (28)-(30), we can determine the matter fields and, consequently, solve the above equations to obtain the fields. However it is convenient, at first, to take into account some fundamental concepts about the orbits in spherical models.

IV. ROTATION CURVES OUTSIDE A STATIC SPHERICAL CONFIGURATION

Here we focus on the problem of circular motion restricted to the equatorial plane (i.e. equatorial circular orbits) of the configuration, in order to study the behavior of rotation curves. In spherical coordinates, equatorial circular orbits satisfy $\theta = \pi/2$, $\ddot{r} = \ddot{\theta} = \dot{r} = \dot{\theta} = 0$ and the equations of motion (22) can be cast as

$$\begin{aligned} (\dot{\varphi})^2 \Big|_{\theta=\frac{\pi}{2}} &= -\frac{1}{r} \left[U_{,r} + \frac{\mathcal{W}_{,r}}{2c^2} - \frac{2\gamma U U_{,r}}{c^2} \right. \\ &\quad \left. - (\Phi_{r,\varphi}^{\text{PF}} - \Phi_{\varphi,r}^{\text{PF}}) \frac{\sqrt{-rU_{,r}}}{rc^2} - \frac{\gamma r (U_{,r})^2}{c^2} \right] \Big|_{\theta=\frac{\pi}{2}}, \end{aligned} \quad (38)$$

$$\frac{1}{c^2 r^2} \left[\frac{\mathcal{W}_{,\theta}}{2} - (\Phi_{\theta,\varphi}^{\text{PF}} - \Phi_{\varphi,\theta}^{\text{PF}}) \dot{\varphi} \right] \Big|_{\theta=\frac{\pi}{2}} = 0, \quad (39)$$

$$\ddot{\varphi} \Big|_{\theta=\frac{\pi}{2}} = \frac{\mathcal{W}_{,\varphi} \Big|_{\theta=\frac{\pi}{2}}}{2c^2 r^2}. \quad (40)$$

In the derivation of the above relations we use the fact that whenever $(r\dot{\varphi})^2$ or $r\dot{\varphi}$ is accompanied by c^{-2} we can substitute it, in accordance with the PPN order, by the Newtonian value [38], i.e. $(r\dot{\varphi})^2 c^{-2} = -rU_{,r} c^{-2}$.

It is important to note that equations (39)-(40) lead to conditions on the existence of circular orbits, which can be expressed by the following relations:

$$\frac{\alpha_2}{c^2 r^2} w^z w^r \Big|_{\theta=\pi/2} \left(\frac{X_{,r}}{r} - X_{,rr} \right) = 0, \quad (41)$$

$$\alpha_2 w^\varphi|_{\theta=\pi/2} X_{,rrr} = -\alpha_1 U_{,r} \sqrt{-rU_{,r}} \quad (42)$$

The first one of the above relations comes from (39), after introducing the expressions for \mathcal{W} and Φ^{PF} in terms of X . The second one comes from (40), in a similar fashion. An interesting consequence of condition (41) is the restriction of the values of some post-Newtonian parameters. Since the term $X_{,r}/r - X_{,rr} \neq 0$ for all asymptotically flat models (i.e. solutions so that $\lim_{r \rightarrow \infty} X_{,r} = 0$) and assuming that conditions (41) and (42) must be satisfied for all choices of \vec{w} , we conclude that *only theories with $\alpha_1 = \alpha_2 = 0$ can guarantee the existence of equatorial circular orbits in static spherically symmetric configurations with no divergent fields*. In other words, only theories in which the angular momentum is conserved globally (i.e. so that $\alpha_1 = \alpha_2 = 0$ [46–48]) can admit equatorial circular orbits in spherical distributions. From here on we focus on these cases.

Introducing the restriction $\alpha_1 = \alpha_2 = 0$ in the equation (38) we obtain the following expression for the circular velocity (i.e. the component $v^\varphi = r\dot{\phi}$ for circular orbits with $\theta = \frac{\pi}{2}$, which we denote shortly as v_c):

$$v_c = \sqrt{-rU_{,r} \left[1 + \frac{\alpha_3 w^2}{2c^2} + \frac{\Psi_{,r}}{c^2 U_{,r}} - \frac{\gamma r U_{,r}}{c^2} - \frac{2(\beta + \gamma)U}{c^2} \right]}, \quad (43)$$

which will be used later to sketch the rotation curves for different theories.

In order to investigate the behavior of circular velocity away from a finite distribution, we first consider the case in which the matter is concentrated inside a sphere of radius a and mass M , and solve the field equations (31)–(33) in the vacuum, i.e. when $\rho^* = P = \rho^* \Pi = 0$ ($r > a$). The expression for the fields U , X and Ψ can be cast as

$$U = \frac{GM}{r}, \quad X = GMr + \frac{C_1}{r} + C_2,$$

$$\Psi = -\frac{C_3}{r} + \frac{\xi G M C_1}{r^4},$$

where C_1 , C_2 and C_3 are constants to be determined from boundary conditions. Introducing the above relations in (43), we obtain the corresponding expression for circular velocity:

$$v_c = \sqrt{\frac{GM}{r} \left[1 + \frac{1}{c^2} \left(A + \frac{B}{r} + \frac{C}{r^3} \right) \right]}, \quad (44)$$

where

$$A = \frac{\alpha_3 w^2}{2} - \frac{C_3}{GM}, \quad B = -(\gamma + 2\beta)GM, \quad C = 4\xi C_1. \quad (45)$$

According to (44), circular velocity, for large values of r , exhibits the usual Keplerian fall off, as in Newtonian theory (i.e. proportional to $1/\sqrt{r}$). Since most solutions,

when far away from the center, could be well approximated to vacuum, therefore (44) should be a good representative of the asymptotic behavior of circular velocities in the PPN approach.

No choice of the PPN parameters can lead to the flattening of the rotation curves. Indeed, such statement is also true for values of r close to the radius a of the mass distribution. This can be easily verified by finding the critic value of r where v_c reaches its maximum. So, the values of r for which v_c is extreme (maximum or minimum) satisfy the cubic equation

$$\left(1 + \frac{A}{c^2} \right) r^3 + \frac{2B}{c^2} r^2 + \frac{4C}{c^2} = 0. \quad (46)$$

The value $r = r_e$ for which v_c reaches its maximum can be cast as

$$r = r_e^0 + r_e^l + \dots,$$

where l is the highest order of c and is determined when substituting in eq. (46). From eq. (46) one may verify that

$$r_e^0 = 0, \quad r_e^{\frac{2}{3}} = \left(\frac{4C}{c^2} \right)^{1/3},$$

which means that $r_e = (4C/c^2)^{1/3}$. Remembering that $C = 4\xi C_1 \sim 4\xi a^3 \epsilon c^2$, we have,

$$r_e \sim (-16\xi\epsilon)^{1/3} a \ll a.$$

This means that there are no solutions of (46) in the region $r > a$. Therefore $dv_c/dr < 0$ for all $r > a$ since v_c and its derivatives are continuous functions and the latter is true for large r .

Now we will abandon the void and consider situations in which some material content permeates the space, in order to allow the possibility of obtaining a different behavior from the one captured by (44).

V. ROTATION CURVES INSIDE A STATIC SPHERICAL CONFIGURATION

Let's go back to the metric of equations (1)–(3). Note that $\alpha_1 = \alpha_2 = 0$ if and only if $g_{\mu\nu}$ is spherically symmetric, i.e. $g_{0j} = 0$, which is necessary for the existence of circular orbits. Therefore we can use the results of reference [3] to simplify the possible choices for a DF. In such paper it was shown that when the space-time is static, spherically symmetric, asymptotically flat and the selfgravitating configuration is made of collisionless identical particles, the DF has the form $f = \xi(E)L^{2(k-1)}$, where L is the angular momentum, k is an integer and $\xi(E)$ is some function of the energy E . This implies that $T^\theta_\theta = kT^r_r$, which can be applied here, since we are adopting the same assumptions about space-time and matter distribution. If we also assume that the system

constitutes a perfect fluid (a basic assumption of the PPN formalism), i.e. $T^\theta_\theta = T^r_r$, we have to chose $k = 1$ and, in consequence, $f = f(E)$. This means that only ergodic DFs are allowed when dealing with static spherical configurations in the PPN formalism.

Here we consider the so-called Polytropes [11, 38], which are, from the perspective of the mathematical form of the DF, the simplest ergodic models of astrophysical interest in galactic dynamics.

A. Polytropes in PPN approach

In Newtonian gravity, polytropes are spherical configurations defined by a DF of the form [11, 38]

$$f(E) = \begin{cases} A_n (-E)^{n-3/2} & \text{for } E < 0 \\ 0 & \text{for } E \geq 0 \end{cases}, \quad (47)$$

where A_n is a constant and n is a real number characterizing the different models, usually known as polytropic index, which is constrained by the condition $n > 1/2$.

Here we assume the same form (47) for selfgravitating configurations in metric theories, taking into account that energy can be cast as $E = E_N + E_{\text{PPN}}$, where E_N is the Newtonian contribution and E_{PPN} represents the PPN correction, so that $E_N \gg E_{\text{PPN}}$ (see appendix D). These distributions will be called here as *PPN polytropes*. The corresponding DF can be expanded (in a similar fashion as done in [38], for the case of 1PN approximation) as the sum of a Newtonian contribution, of order 0 in ϵ , and a PPN contribution, of order 2 in ϵ :

$$f = f^0 + f^2,$$

where we have introduced,

$$f^0 = A_n (-E_N)^{n-3/2}, \quad (48)$$

$$f^2 = -A_n \left(n - \frac{3}{2} \right) E_{\text{PPN}} (-E_N)^{n-5/2}, \quad (49)$$

for $E_N < 0$ and $E_{\text{PPN}} < 0$.

The above DF contributions determine ρ^* , P and Π through equations (28)-(30). The integrals at the right hand side of such equations must be limited to values of velocity such that $E < 0$. This escape velocity, determined by condition $E = 0$, is given by

$$v_e = \sqrt{2U} + \frac{[2\Psi + \Phi^{\text{PF}} - (4\gamma + 5)U^2]}{2c^2\sqrt{2U}}, \quad (50)$$

but, in order to obtain expressions up to first order in ϵ , it is sufficient to take $v_e = \sqrt{2U}$ in the limits of integration. So, by introducing (48), (49) and (50) in (28)-(30), we find the matter fields for the PPN polytropes, as functions of U , ψ and Φ^{PF} :

$$\rho^* = c_n U^n, \quad (51)$$

$$P = c_n \frac{U^{n+1}}{n+1}, \quad (52)$$

$$\rho^* \Pi = c_n U^{n+1} \left[3\gamma + \frac{3}{2} + \frac{27}{8(n+1)} + n \left(\frac{1}{2} - \beta \right) \right] + n c_n \left(\Psi + \frac{\Phi^{\text{PF}}}{2} \right) U^{n-1}, \quad (53)$$

where

$$c_n = A_n (2\pi)^{\frac{3}{2}} \frac{\Gamma(n-1/2)}{\Gamma(1+n)}.$$

Note that density ρ^* and pressure P follow exactly the same equations as their Newtonian counterparts, ensuring that they satisfy the polytropic equation of state [11].

B. Field Equations for PPN Polytropes

Introducing (51)-(53) in the field equations (31)-(33), we obtain the following system of second order differential equations:

$$\frac{1}{\tilde{r}^2} \frac{d}{d\tilde{r}} \left(\tilde{r}^2 \frac{d\tilde{U}}{d\tilde{r}} \right) = -\tilde{U}^n, \quad (54)$$

$$\frac{1}{\tilde{r}^2} \frac{d}{d\tilde{r}} \left(\tilde{r}^2 \frac{d\tilde{X}}{d\tilde{r}} \right) = 2\tilde{U}, \quad (55)$$

$$\begin{aligned} \frac{1}{\tilde{r}^2} \frac{d}{d\tilde{r}} \left(\tilde{r}^2 \frac{d\tilde{\Psi}}{d\tilde{r}} \right) &= b_n \frac{U_0^2}{\Psi_0} \tilde{U}^{n+1} - e_n \frac{U_0}{\Psi_0} \tilde{U}^n - d_n \tilde{\Psi} \tilde{U}^{n-1} \\ &+ 4 \frac{U_0^2}{\Psi_0} \xi \left[\frac{\tilde{U}^n}{\tilde{r}} \frac{d\tilde{X}}{d\tilde{r}} + \frac{d\tilde{U}}{d\tilde{r}} \left(\frac{d\tilde{U}}{d\tilde{r}} + \frac{3}{\tilde{r}^2} \frac{d\tilde{X}}{d\tilde{r}} - \frac{2\tilde{U}}{\tilde{r}} \right) \right]. \end{aligned} \quad (56)$$

where U_0 and Ψ_0 are the values of U and Ψ at the center of the configuration, respectively, and we have defined,

$$\tilde{U} := U/U_0, \quad \tilde{\Psi} := \Psi/\Psi_0, \quad \tilde{X} := C_n U_0^{n-2} X, \quad (57)$$

$$\tilde{r} := r \sqrt{C_n U_0^{n-1}}, \quad (58)$$

along with the constants

$$C_n = 4\pi G c_n, \quad d_n = (1 + \zeta_3) n = \frac{2e_n}{\alpha_3 w^2}. \quad (59)$$

$$\begin{aligned} b_n &= (2\beta - 1 - \zeta_2 - 6\xi) - \frac{(3\gamma + 3\zeta_4 - 2\xi)}{(n+1)} \\ &- (1 + \zeta_3) \left[3\gamma + \frac{3}{2} + \frac{27}{8(n+1)} + \frac{n}{2} - \beta n \right] \end{aligned} \quad (60)$$

Equation (54) has simple analytic solutions for the cases $n = 0, 1, 5$ [11, 38], being $n = 5$ the well known case of Plummer model [11, 36–38]. For other values of n , Eq. (54) requires a numerical solution. On the other hand, equations (55) and (56) are much more complicated, even in the cases $n = 0, 1, 5$, where U has analytical expression. However, as we will show in the following subsection, one can find an asymptotic solution for large r . In subsection V D we will obtain numerical solutions for the system (54)-(56), by using the initial conditions

$$\tilde{U}(0) = \tilde{X}(0) = \tilde{\Psi}(0) = 1, \quad \tilde{U}'(0) = \tilde{X}'(0) = \tilde{\Psi}'(0) = 0,$$

reflecting the assumption that \tilde{U} , \tilde{X} and $\tilde{\Psi}$ have maximum value in the center of the configuration.

Since parameter ϵ can be regarded a representative of the gravitational strength, we also assume that $U_0 = \epsilon c^2$, $\Psi_0 = (U_0)^2$ and $w = \epsilon c$, for numerical and asymptotic solutions. For numerical solutions we consider only $0 \leq \epsilon \leq 0.15$. Larger values of ϵ may exceed the validity limit of the PPN approximation.

On the other hand, we only accept solutions that result in an asymptotically flat space-time. Therefore any solution leading to something different from

$$\lim_{r \rightarrow \infty} g_{00} = -1, \quad \lim_{r \rightarrow \infty} g_{ij} = 1,$$

will be discarded, since it does not satisfy the basic assumptions of the PPN approximation.

C. The Approximated Solution

In order to understand the importance of the contribution of each of the parameters appearing in field equations, specially in relation with the behavior of solutions far from the center of configuration, we consider here an asymptotic solution of (55)-(56), based on the Plummer's polytrope (i.e. the case $n = 5$ of eq. (54) [11, 36, 37]),

$$\tilde{U} = \frac{1}{\sqrt{1 + \frac{\tilde{r}^2}{3}}}. \quad (61)$$

For large radius, $\tilde{r} \gg \sqrt{3}$, we have $\tilde{U} \approx \sqrt{3}/\tilde{r}$, which means that, according to (55), we can write

$$\frac{d\tilde{X}}{d\tilde{r}} \approx \sqrt{3}, \quad \text{for } r \gg \sqrt{3},$$

which, in turn, implies that eq. (56) reduces to

$$\frac{d}{d\tilde{r}} \left(\tilde{r}^2 \frac{d\tilde{\Psi}}{d\tilde{r}} \right) + 9d_5 \frac{\tilde{\Psi}}{\tilde{r}^2} = \frac{27(b_5 - 4\xi)}{\tilde{r}^4} - \frac{9\sqrt{3}\alpha_3 d_5}{2\tilde{r}^3} + \frac{24\xi}{\tilde{r}^2},$$

for $\tilde{r} \gg \sqrt{3}$. The solution of the above equation can be written as

$$\begin{aligned} \tilde{\Psi}(\tilde{r}) = & c_1 \cos\left(\frac{3\sqrt{d_5}}{\tilde{r}}\right) - c_2 \sin\left(\frac{3\sqrt{d_5}}{\tilde{r}}\right) - \frac{\sqrt{3}\alpha_3}{2\tilde{r}} \\ & + \frac{3(b_5 - 4\xi)}{d_5 \tilde{r}^2} - \frac{2(b_5 - 4\xi)}{3d_5^2} + \frac{8\xi}{3d_5}. \end{aligned} \quad (62)$$

Since we are interested in solutions consistent with the requirement of asymptotically flat space-time, we demand that $\lim_{\tilde{r} \rightarrow \infty} \tilde{\Psi}(\tilde{r}) = 0$, leading to the following relation:

$$c_1 = \frac{2b_5 - 8(d_5 + 1)\xi}{3d_5^2}.$$

On the other hand, c_2 has to remain undetermined since no other boundary condition can be applied here. Regardless, one can extract additional information from the corresponding expression for the velocity of circular orbits:

$$\begin{aligned} v_c = & c \sqrt{\epsilon \frac{\sqrt{3}}{\tilde{r}}} \left\{ 1 + \epsilon \frac{\sqrt{3}}{\tilde{r}} \left[\frac{2(b_5 - 4\xi)}{d_5} - (\gamma + 2\beta) \right] \right. \\ & \left. - \epsilon \sqrt{3d_5} \left[c_1 \sin\left(\frac{3\sqrt{d_5}}{\tilde{r}}\right) + c_2 \cos\left(\frac{3\sqrt{d_5}}{\tilde{r}}\right) \right] \right\}^{1/2}. \end{aligned}$$

Note that the second term inside the brackets should be insignificantly small, because it involves terms multiplied by $\epsilon\sqrt{3}/\tilde{r}$. Therefore it is only relevant to the behavior of the curves the third term ($c_1 \sin(3\sqrt{d_5}/\tilde{r}) + c_2 \cos(3\sqrt{d_5}/\tilde{r})$). For large values of d_5 the curves may present significant modifications from the Newtonian ones. One may even find, depending on the constants c_1 and c_2 , curves resembling flat rotation curves. Remembering that $d_5 = 5(1 + \zeta_3)$ we conclude that the only significant contributions will arise from positive values of ζ_3 .

D. The Numerical Solution

To solve the field equations (54)-(56) for the entirety of space we use a fourth order Runge-Kutta method. In figures 1 and 2, we show numerical solutions for \tilde{U} and $\tilde{\Psi}$ for PPN Polytropes with polytropic index $n = 5$, for various values of the PPN parameters.

The behavior of the matter fields is shown in figures 3 and 4, through the effective density $\tilde{\rho}$ for each model, defined as

$$\begin{aligned} \tilde{\rho}^* = & \frac{-4\pi GT}{(U_0)^n C_n} = \frac{-4\pi GT^\mu{}_\mu}{(U_0)^n C_n} \\ = & \frac{4\pi G \rho^*}{(U_0)^n C_n} \left\{ 1 + \frac{1}{c^2} [\Pi - 3\gamma U] \right\} - \frac{12\pi GP}{(U_0)^n C_n} \\ = & \tilde{U}^n \left(1 + \frac{\epsilon n \alpha_3}{2} \right) + \epsilon n \tilde{\Psi} \tilde{U}^{n-1} \\ & + \epsilon \tilde{U}^{n+1} \left[\left(\frac{1}{2} - \beta \right) n + \frac{3}{8(n+1)} + \frac{3}{2} \right]. \end{aligned}$$

The overall behavior turn out to be similar for other polytropic indexes and other combinations of the PPN parameters.




Figure 1: We present the gravitational fields \tilde{U} and $\tilde{\psi}$ for a polytropic model with $n = 5$ for fully conservative theories ($\zeta_1 = \zeta_2 = \zeta_3 = \zeta_4 = \alpha_3 = 0$). The top figures are for β different from 1, the central ones for γ different from 1 and the lowest are for a ξ different from 0. The continuous line represents \tilde{U} and the traced one $\tilde{\psi}$.

E. The PPN corrections to the rotation curves

With the solutions at hand, we sketch the rotation curves for a polytropic model with index $n = 5$ for several values of the PPN parameters (see figures 5 and 6). The rotation curves for $n \neq 5$ are either similar to the ones of $n = 5$ or have unphysical properties.

Although we only present some choices for the PPN parameters (it is not easy to develop a systematic approach to test all possible combination of values for the PPN parameters) it is reasonable to conclude, from the curves we presented, Figs. 5 and 6, that only the ones with $\zeta_3 > 0$ have some resemblance to flattened curves (as expected from the analysis of the asymptotic solution of previous section). The outcome corresponding to other combinations, including other polytropes with $n \neq 5$, are alike Figs. 5 and 6. In particular, the curves with $\zeta_3 = 2$ present some flat behavior, but have lower values of velocity than the Newtonian one (except at some critical value around $\tilde{r} = 6$). So, for this model to be a plausible explanation of the rotation curves of galaxies, it must have less mass than the Newtonian one. This can be seen by comparing the density of mass ρ with the trace of the stress-energy tensor $-T = -T^\mu_\mu$. This is shown in Fig. 4 and we note that $-T$ falls much faster, with the radius, than the density of mass, therefore explaining why the rotation curves for the PPN have a lower first local maximum value of velocity and much sooner than the Newtonian one. This means that a theory with $\zeta_3 > 0$




Figure 2: We present the gravitational fields \tilde{U} and $\tilde{\psi}$ for a polytropic model with $n = 5$ for non-conservative theories. For simplicity, here we consider $\gamma = \beta = 1$ and $\xi = 0$. From top to bottom, we alternate α_3 , ζ_1 , ζ_2 and ζ_3 to be different from 0, respectively. The continuous line represents \tilde{U} and the traced one $\tilde{\psi}$.

does present some interesting corrections to the rotation curves. But, as it should be, the usual Keplerian falloff is still found for $\tilde{r} > 10$.

On the other hand, we note that choices involving $\zeta_3 \leq 0$ (even for high values of the parameter ϵ and other exotic choices), have the same behavior than the Newtonian models far from the center of the configuration.

VI. CONCLUSION

In this paper, the rotation curves of static spherically symmetric models were studied in the framework of the PPN approximation. We first started addressing vacuum solutions and later we focus on polytropic models (see [38] for a similar approach in the post-Newtonian context).

When the matter fields, endowed with spherical symmetry, are assumed static one can readily notice that the PPN metric is not necessarily spherically symmetric (a direct consequence of the preferred frame potentials). Therefore, circular orbits will only exist if and only if the parameters α_1 and α_2 are null. This means, as their




Figure 3: We present the effective density $\tilde{\rho}^*$ for a polytropic model with index $n = 5$ for fully conservative theories ($\zeta_1 = \zeta_2 = \zeta_3 = \zeta_4 = \alpha_3 = 0$). The top figures are for β different from 1, the middle ones for γ different from 1 and the lower ones for ξ different from 0. The continuous line represents the Newtonian curve and the dotted, dashed and dash-dotted are PPN curves with $\epsilon = 0.05$, $\epsilon = 0.1$ and $\epsilon = 0.15$, respectively.

heuristic meaning imply, that we excluded the effects of non-conservation of global angular momenta of the PPN scheme. For all calculations that followed Sec. IV we assumed $\alpha_1 = \alpha_2 = 0$.

As a first setup for testing the circular orbits in the PPN approach, we solved field eqs. (31)-(33) in vacuum outside some static spherically symmetrical distribution of matter of finite radius and presented, to PPN order, the velocity of circular orbits (eq. (44)). We argued that independently of the choice of the PPN parameters such solution can not lead to flat curves. Our argument is based on the simple property that, to PPN order, such solutions have necessarily $dv/dr < 0$ for all $r > a$. Therefore a Keplerian falloff, similar to the ones found in Newtonian gravity, will still be observed in this description of gravity.

To construct astrophysical models, when matter fields are present, we applied a statistical description, following the assumptions of Subsec. II A. In such cases, the system is fully described by a DF that satisfy the CBE, which in the context of the PPN approximation takes the form of eq. (24). In statical spherical models the previous assumptions and the results of Subsec. IV implies (see Subsec. V) that the DF must be a function of only the energy like integral of motion. The Polytropes are a family of models whose DF is a particular case that satisfy the previously stated requirement. In the context of the PNP approach, the Polytropes can be constructed




Figure 4: We present the effective density $\tilde{\rho}^*$ for a polytropic model with index $n = 5$ for non-conservative theories, here we regard the parameters $\gamma = \beta = 1$ and $\xi = 0$. From top to bottom, we take alternatively α_3 , ζ_1 , ζ_2 and ζ_3 different from 0, respectively. The continuous line represents the Newtonian curve and the dotted, dashed and dash-dotted are PPN curves with $\epsilon = 0.05$, $\epsilon = 0.1$ and $\epsilon = 0.15$, respectively.

by a simple extension of their Newtonian DF, this leads to the complete specification of the matter fields. Therefore one can determine all the gravitational fields from (54)-(56).

To solve eqs. (54)-(56) we employed: I) approximated methods (Sec. V C); II) numerical techniques (Sec. V D). In V C, we solved approximately the field equations (for $n = 5$) when the radius is much larger than a characteristic radius of the model ($\tilde{r} \gg \sqrt{3}$). The approximation had the setback of not providing proper boundary conditions, but the velocity of circular orbits did present some terms with behavior different from the Newtonian solution. From the approximated solutions we observed that the larger modifications will generally come from large values of the parameter ζ_3 . But to be certain, since we lack proper boundary conditions, one needs the solution for the entirety of space, which can only be achieved by a numerical route.

Using numerical techniques (fourth order Runge-Kutta method) we were able to find the gravitational and matter fields for all choices of parameters, Figs. 1, 2, 3 and 4.




Figure 5: We present the rotation curves ($\tilde{v}^\varphi = v^\varphi/\sqrt{U_0}$) for a polytropic model with index $n = 5$ for fully conservative theories ($\zeta_1 = \zeta_2 = \zeta_3 = \zeta_4 = \alpha_3 = 0$). The top figures are for β different from 1, the middle ones for γ different from 1 and the lower ones for ξ different from 0. The continuous line represents the Newtonian curve and the dotted, dashed and dash-dotted are PPN curves with $\epsilon = 0.05$, $\epsilon = 0.1$ and $\epsilon = 0.15$, respectively.

In Figs. 5 and 6 we present the rotation curves for several values of the Post-Newtonian parameters, and concluded that many of the choices lead to trivial corrections to the rotation curves, sometimes leading to smaller values of the circular velocities or an even higher descent rate, with the radius, then the Newtonian models.

There was a single choice of the parameters that lead to different behavior. By choosing $\zeta_3 > 0$ we found some flat rotation curves and with the hassle of lower values of circular velocity. But by analyzing the effective density ($-T$) we realize this should be expected, since the amount of matter in this model is significantly less than in its Newtonian counterpart. Therefore a theory with $\zeta_3 > 0$, as far as our analysis goes, does present interesting modifications to the Keplerian behavior. All other choices of the parameters comes short in explaining the rotation curves of galaxies. Since only one of the parameters presents real significance one may conclude that gravity theories compatible with the PPN scheme are not very successful as alternatives to the dark matter hypothesis in galactic dynamics.

Here, we do not claim that the PPN approach contains all the potentials needed to describe theories with SM. In [10, 54, 55] was already shown that the parameterized version of the Vainshteinian screening requires more potentials (which describes the effects of the mechanism). In this paper, we only considered the poten-




Figure 6: We present the rotation curves ($\tilde{v}^\varphi = v^\varphi/\sqrt{U_0}$) for a polytropic model with index $n = 5$ for non-conservative theories, here we regard the parameters $\gamma = \beta = 1$ and $\xi = 0$. From top to bottom, we take alternatively α_3 , ζ_1 , ζ_2 and ζ_3 different from 0, respectively. The continuous line represents the Newtonian curve and the dotted, dashed and dash-dotted are PPN curves with $\epsilon = 0.05$, $\epsilon = 0.1$ and $\epsilon = 0.15$, respectively.

tials present in the standard PPN approach, we excluded the Vainshteinian ones (or from other SMs) that may emerge from a more complete formalism. But it is to be expected that for $r \gg r_v$ the approximation should be equivalent to the PPN approximation. Nonetheless, we recognize that all theories (with or without a SM) have, at least, some of the standard PPN potentials. Even though the work by Avilez-Lopez et al. [10], and the further extensions by [54, 55], does outline the main ideas for constructing the PPNV (Parametrized Post-Newtonian Vainshteinian) formalism, we will leave the complete analysis of this problem for when the Post-Newtonian expansion of theories with SMs are better understood.

Acknowledgments

H.M.G. is grateful for the financial support provided by CNPq (Grant No. 132389/2017-7 and No. 141924/2019-

5).

Appendix A: Derivation of the Field Equations

Expressions (10) and (11) are easily determined from their integral forms (see [47]). But this is not the case for expressions (12) and (13), they require more attention.

In this section we present a derivation of the field equations (12) and (13) from the integral forms found in reference [47].

Derivation of the Field Equation for ϕ_6

Starting from the integral form of ϕ_6 [47]:

$$\begin{aligned}\phi_6 &= G \int \rho^{*'} u'_j u'_k \frac{(\mathbf{x} - \mathbf{x}')^j (\mathbf{x} - \mathbf{x}')^k}{|\mathbf{x} - \mathbf{x}'|^3} d^3 x' \\ &= -G \int \rho^{*'} \left[\mathbf{u}' \cdot (\mathbf{x} - \mathbf{x}') \mathbf{u}' \cdot \nabla |\mathbf{x} - \mathbf{x}'|^{-1} \right] d^3 x'.\end{aligned}$$

We want to reduce this to its differential form, to remove the integral we take the laplacian of the integral form, leading to:

$$\begin{aligned}\nabla^2 \phi_6 &= -G \int \rho^{*'} \left[3u^2 4\pi \delta(\mathbf{x} - \mathbf{x}') + 2(\mathbf{u}' \cdot \nabla)^2 |\mathbf{x} - \mathbf{x}'|^{-1} \right. \\ &\quad \left. - 4\pi (\mathbf{u}' \cdot \nabla) \mathbf{u}' \cdot (\mathbf{x} - \mathbf{x}') \delta(\mathbf{x} - \mathbf{x}') \right] d^3 x',\end{aligned}$$

which follows from

$$\begin{aligned}\nabla^2 \left[\mathbf{u}' \cdot (\mathbf{x} - \mathbf{x}') \mathbf{u}' \cdot \nabla |\mathbf{x} - \mathbf{x}'|^{-1} \right] &= \\ 3u^2 4\pi \delta(\mathbf{x} - \mathbf{x}') + 2(\mathbf{u}' \cdot \nabla)^2 |\mathbf{x} - \mathbf{x}'|^{-1} &- \\ - 4\pi (\mathbf{u}' \cdot \nabla) \mathbf{u}' \cdot (\mathbf{x} - \mathbf{x}') \delta(\mathbf{x} - \mathbf{x}') &.\end{aligned}$$

From this it's straightforward to show that

$$\begin{aligned}\nabla^2 \phi_6 &= 3\nabla^2 \phi_1 - 2G \int \rho^{*'} (\mathbf{u}' \cdot \nabla)^2 \left(\frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) d^3 x' \\ &= 3\nabla^2 \phi_1 - 2G \partial_{ij} \int \frac{\rho^{*'} u^{i'} u^{j'}}{|\mathbf{x} - \mathbf{x}'|} d^3 x' .\end{aligned}$$

The Laplacian of the previous expression results in (12).

Derivation of the Field Equation for ϕ_w

The integral form of ϕ_w can be presented in different forms [47]. Here, we start from the identity, see [47],

$$\phi_w = -U^2 - \phi_2 - \nabla U \cdot \nabla X + G \nabla \cdot \int \frac{\rho^{*'}}{|\mathbf{x} - \mathbf{x}'|} \nabla' X' d^3 x'.$$

Again, we wish to reduce this to a partial differential equation. To achieve this we start from the laplacian of the previous expression, resulting in:

$$\begin{aligned}\nabla^2 \phi_w &= -\nabla^2 U^2 - \nabla^2 \phi_2 - \nabla^2 (\nabla U \cdot \nabla X) \\ &\quad + G \nabla^2 \left(\nabla \cdot \int \frac{\rho^{*'}}{|\mathbf{x} - \mathbf{x}'|} \nabla' X' d^3 x' \right). \quad (A1)\end{aligned}$$

Now the identities,

$$\nabla^2 (\nabla U \cdot \nabla X) = 2U_{,ij} X_{,ij} + 2\nabla U \cdot \nabla U - 4\pi G \nabla X \cdot \nabla \rho^*,$$

$$\nabla^2 U^2 = 2\nabla^2 \phi_2 + 2\nabla U \cdot \nabla U,$$

$$G \nabla^2 \left(\nabla \cdot \int \frac{\rho^{*'}}{|\mathbf{x} - \mathbf{x}'|} \nabla' X' d^3 x' \right) = 2\nabla^2 \phi_2 - 4\pi G \nabla X \cdot \nabla \rho^*,$$

can be substituted in (A1), which implies in:

$$\nabla^2 \phi_w = -2\nabla^2 U^2 + 3\nabla^2 \phi_2 - 2U_{,ij} X_{,ij},$$

this is exactly (13).

Appendix B: A derivation of eq. (24)

Take the map $(x^\mu, V^i) \rightarrow (x^\mu, v^i(x^\mu, V^i))$, where

$$v^i := \frac{dx^i}{dt} = \frac{cV^i}{V^0(x^\mu, V^i)}, \quad (B1)$$

and rewrite (23) in the new variables (x^μ, v^i) , then the partial derivatives of f are expressed as

$$\left(\frac{\partial f}{\partial x^\mu} \right)_{V^i} = \left(\frac{\partial f}{\partial x^\mu} \right)_{v^i} + \left(\frac{\partial f}{\partial v^i} \right)_{x^\mu} \left(\frac{\partial v^i}{\partial x^\mu} \right)_{V^i},$$

$$\left(\frac{\partial f}{\partial V^j} \right)_{x^\mu} = \left(\frac{\partial f}{\partial v^i} \right)_{x^\mu} \left(\frac{\partial v^i}{\partial V^j} \right)_{x^\mu}.$$

A straightforward calculation, using eq. (B1), lead us to express

$$\left(\frac{\partial v^i}{\partial x^\mu} \right)_{V^i} = \frac{-cV^i}{(V^0)^2} \left(\frac{\partial V^0}{\partial x^\mu} \right)_{V^i}, \quad (B2)$$

$$\left(\frac{\partial v^i}{\partial V^j} \right)_{x^\mu} = \frac{c}{V^0} \left[\delta^i_j - \frac{V^i}{V^0} \left(\frac{\partial V^0}{\partial V^j} \right)_{x^\mu} \right]. \quad (B3)$$

Now f , as defined, represent the probability density of a single massive particle, therefore the possible values V^μ can take is restricted to the positive light cone, where the shell condition holds:

$$g_{\mu\nu} V^\mu V^\nu = -c^2, \quad (B4)$$

also $V^0 > 0$. This restriction completely defines V^0 in terms of (x^μ, V^i) or (x^μ, v^i) , which to PPN order can be expressed by

$$V^0 = c \left\{ 1 + \frac{U}{c^2} + \frac{\mathcal{W}}{2c^4} + \frac{\mathcal{Q}_j v^j}{c^4} + \frac{3v^4}{8c^4} + \frac{3U^2}{2c^4} + \left[1 + \frac{(2\gamma + 3)U}{c^2} \right] \frac{v^2}{2c^2} + O(c^{-6}) \right\},$$

also one can determine, from the latter and (B1), V^i as an expression of (x^μ, v^i) .

Some straightforward calculations determines the right-hand side of (B2) and (B3) only in terms of the new variables (x^μ, v^i) and to PPN order, as follows:

$$\left(\frac{\partial v^i}{\partial x^\mu} \right)_{V^k} = -v^i \frac{U_{,\mu}}{c^2},$$

$$\left(\frac{\partial v^i}{\partial V^j} \right)_{x^\mu} = \left(1 - \frac{U}{c^2} - \frac{v^2}{2c^2} \right) \delta_j^i - \frac{v^i v^j}{c^2}.$$

With all these results in mind we can determine each term of the Vlasov equation in terms of (x^μ, v^i) to PPN order:

$$\frac{V^0}{c} \left(\frac{\partial f}{\partial t} + \frac{\partial f}{\partial v^i} \frac{\partial v^i}{\partial t} \right) = \left[1 + \frac{U}{c^2} + \frac{v^2}{2c^2} \right] \frac{\partial f}{\partial t} - v^i \frac{\partial f}{\partial v^i} \frac{U_{,t}}{c^2},$$

$$V^i \left(\frac{\partial f}{\partial x^i} + \frac{\partial f}{\partial v^j} \frac{\partial v^j}{\partial x^i} \right) = v^i \left[1 + \frac{U}{c^2} + \frac{v^2}{2c^2} \right] \frac{\partial f}{\partial x^i} - v^i v^j \frac{\partial f}{\partial v^j} \frac{U_{,i}}{c^2},$$

$$\Gamma_{00}^j V^0 V^0 \frac{\partial f}{\partial v^i} \frac{\partial v^i}{\partial V^j} = \frac{\partial f}{\partial v^i} \left\{ -\frac{1}{2} \frac{\mathcal{W}_{,i}}{c^2} + \frac{\mathcal{Q}_{i,t}}{c^2} + \frac{2\gamma}{c^2} U U_{,i} - U_{,i} \left(1 + \frac{U}{c^2} + \frac{v^2}{2c^2} \right) + U_{,j} \frac{v^i v^j}{c^2} \right\},$$

$$\Gamma_{i0}^j V^i V^0 \frac{\partial f}{\partial v^i} \frac{\partial v^i}{\partial V^j} = \frac{1}{2c^2} [\mathcal{Q}_{j,l} + 2\gamma U_{,t} \delta_{lj} - \mathcal{Q}_{l,j}] v^l \frac{\partial f}{\partial v^j},$$

$$\Gamma_{lp}^j V^l V^p \frac{\partial f}{\partial v^i} \frac{\partial v^i}{\partial V^j} = \gamma [U_{,l} \delta_{jp} + U_{,p} \delta_{jl} - U_{,j} \delta_{lp}] \frac{v^l v^p}{c^2} \frac{\partial f}{\partial v^j}.$$

Substituting each of these expressions in eq. (23) we find the Vlasov equation to PPN order:

$$\begin{aligned} & \left[1 + \frac{U}{c^2} + \frac{v^2}{2c^2} \right] \left(\frac{\partial f}{\partial t} + v^i \frac{\partial f}{\partial x^i} \right) + U_{,i} \frac{\partial f}{\partial v^i} \\ & - \frac{\partial f}{\partial v^i} \frac{v^i}{c^2} U_{,t} (1 + 2\gamma) - \frac{\partial f}{\partial v^i} \frac{v^i v^j}{c^2} U_{,j} (2 + 2\gamma) \\ & - \frac{\partial f}{\partial v^i} \left\{ -\frac{1}{2} \frac{\mathcal{W}_{,i}}{c^2} + \frac{\mathcal{Q}_{i,t}}{c^2} + \frac{2\gamma - 1}{c^2} U U_{,i} \right\} \\ & - [\mathcal{Q}_{j,l} - \mathcal{Q}_{l,j}] \frac{v^l}{c^2} \frac{\partial f}{\partial v^j} + \left(\gamma + \frac{1}{2} \right) U_{,j} \frac{v^2}{c^2} \frac{\partial f}{\partial v^j} = 0, \end{aligned}$$

which is exactly eq. (24).

Appendix C: Derivation of the conditions (41) and (42)

We start from eq. (39) and calculate each term,

$$\begin{aligned} \mathcal{W}_{,\theta}|_{\theta=\frac{\pi}{2}} &= 2\alpha_2 \left(X_{,rr} - \frac{X_{,r}}{r} \right) (w^r w^r_{,\theta})|_{\theta=\frac{\pi}{2}} \\ &= -2\alpha_2 \left(X_{,rr} - \frac{X_{,r}}{r} \right) w^z w^r|_{\theta=\frac{\pi}{2}}, \end{aligned}$$

$$\Phi_{\theta,\varphi}^{\text{PF}}|_{\theta=\frac{\pi}{2}} = \left(\alpha_2 \frac{X_{,r}}{r} - \frac{\alpha_1}{2} U \right) w_{\theta,\varphi}|_{\theta=\frac{\pi}{2}} = 0,$$

$$\Phi_{\varphi,\theta}^{\text{PF}}|_{\theta=\frac{\pi}{2}} = \left(\alpha_2 \frac{X_{,r}}{r} - \frac{\alpha_1}{2} U \right) w_{\varphi,\theta}|_{\theta=\frac{\pi}{2}} = 0.$$

Therefore substituting the previous results in eq. (39) we find condition (41).

The next condition (42) we find by comparing eq. (38) with (40). First we take the derivative with time (t) of eq. (38):

$$\ddot{\varphi}|_{\theta=\frac{\pi}{2}} = \frac{1}{2} \left[(\Phi_{r,\varphi\varphi}^{\text{PF}} - \Phi_{\varphi,r\varphi}^{\text{PF}}) \frac{\sqrt{-rU_{,r}}}{r^2 c^2} - \frac{\mathcal{W}_{,r\varphi}}{2rc^2} \right] \Big|_{\theta=\frac{\pi}{2}}. \quad (\text{C1})$$

Now calculate each term of eq. (C1) and eq. (40), leading to the following:

$$\begin{aligned} \Phi_{r,\varphi\varphi}^{\text{PF}}|_{\theta=\frac{\pi}{2}} &= \left(\alpha_2 \frac{X_{,r}}{r} - \frac{\alpha_1}{2} U \right) w_{r,\varphi\varphi}|_{\theta=\frac{\pi}{2}} \\ &= - \left(\alpha_2 \frac{X_{,r}}{r} - \frac{\alpha_1}{2} U \right) w_r|_{\theta=\frac{\pi}{2}}, \end{aligned}$$

$$\begin{aligned} \Phi_{\varphi,r\varphi}^{\text{PF}}|_{\theta=\frac{\pi}{2}} &= \left(\alpha_2 X_{,r} - \frac{\alpha_1}{2} rU \right)_{,r} \left(\frac{w_\varphi}{r} \right)_{,\varphi} \Big|_{\theta=\frac{\pi}{2}} \\ &= - \left(\alpha_2 X_{,rr} - \frac{\alpha_1}{2} U - \frac{\alpha_1}{2} rU_{,r} \right) w_r|_{\theta=\frac{\pi}{2}}, \end{aligned}$$

$$\begin{aligned} \mathcal{W}_{,\varphi}|_{\theta=\frac{\pi}{2}} &= 2\alpha_2 \left(X_{,rr} - \frac{X_{,r}}{r} \right) (w^r w^r_{,\varphi})|_{\theta=\frac{\pi}{2}} \\ &= 2\alpha_2 \left(X_{,rr} - \frac{X_{,r}}{r} \right) \left(w_r \frac{w_\varphi}{r} \right) \Big|_{\theta=\frac{\pi}{2}}, \end{aligned}$$

$$\mathcal{W}_{,r\varphi}|_{\theta=\frac{\pi}{2}} = 2\alpha_2 \left(X_{,rrr} - \frac{X_{,rr}}{r} + \frac{X_{,r}}{r^2} \right) \left(w_r \frac{w_\varphi}{r} \right) \Big|_{\theta=\frac{\pi}{2}}.$$

Since (C1)=(40) we must have

$$\left[(\Phi_{r,\varphi\varphi}^{\text{PF}} - \Phi_{\varphi,r\varphi}^{\text{PF}}) \sqrt{-rU_{,r}} - \frac{r\mathcal{W}_{,r\varphi}}{2} \right] \Big|_{\theta=\frac{\pi}{2}} = \mathcal{W}_{,\varphi}|_{\theta=\frac{\pi}{2}},$$

now substitute the previous results to find:

$$\begin{aligned} & \left(\alpha_2 X_{,rr} - \alpha_2 \frac{X_{,r}}{r} - \frac{\alpha_1}{2} r U_{,r} \right) \sqrt{-r U_{,r}} w_r \Big|_{\theta=\frac{\pi}{2}} \\ & - \alpha_2 r X_{,rrr} \left(w_r \frac{w_\varphi}{r} \right) \Big|_{\theta=\frac{\pi}{2}} \\ & = \alpha_2 \left(X_{,rr} - \frac{X_{,r}}{r} \right) \left(w_r \frac{w_\varphi}{r} \right) \Big|_{\theta=\frac{\pi}{2}}, \end{aligned}$$

here the choice of \vec{w} shouldn't influence the result, therefore by employing condition (41) in the previous expression we find

$$\alpha_2 w^\varphi \Big|_{\theta=\pi/2} X_{,rrr} = -\alpha_1 U_{,r} \sqrt{-r U_{,r}},$$

which is exactly (42).

Appendix D: Energy like integral of motion

In Sect. V we show that, for the situations analyzed here, it is sufficient to consider DFs depending only on energy. Therefore we need to determine an expression for the conserved energy in the PPN formalism. To do so we follow the same procedure used in [38], which consist of starting from the Lagrangian,

$$\mathcal{L} = \frac{1}{2} g_{\mu\nu} V^\mu V^\nu,$$

which, for the PPN metric (1)-(3), takes the form

$$\begin{aligned} \mathcal{L} = \frac{m^2 c^2}{2} & \left[\left(-1 + \frac{2U}{c^2} + \frac{W}{c^4} \right) \dot{t}^2 \right. \\ & \left. + \left(1 + \frac{2\gamma U}{c^2} \right) \delta_{ij} \frac{V^i V^j}{c^2} \right]. \quad (\text{D1}) \end{aligned}$$

where $\dot{t} = dx^0/d\tau$. Since the metric does not depend on x^0 , then $\partial\mathcal{L}/\partial\dot{t}$ must be a conserved quantity.

By defining the energy to be, [38],

$$E := -\frac{1}{m^2} \frac{\partial\mathcal{L}}{\partial\dot{t}} - c^2,$$

we find, to the order of the PPN, that E can be cast as

$$E = E_N + E_{\text{PPN}},$$

where E_N represents the Newtonian expression for the energy,

$$E_N = -U + \frac{v^2}{2},$$

and E_{PPN} is the PPN correction

$$E_{\text{PPN}} = \left(\gamma + \frac{1}{2} \right) \frac{U v^2}{c^2} - \frac{\Psi}{c^2} - \left(\frac{1}{2} - \beta \right) \frac{U^2}{c^2} - \frac{\Phi^{\text{PF}}}{2c^2} + \frac{3v^4}{8c^2}.$$

As it is usual, one can assume that $E_N \gg E_{\text{PPN}}$.

-
- [1] P. Bull, Y. Akrami, J. Adamek, T. Baker, E. Bellini, J. B. Jiménez, E. Bentivegna, S. Camera, S. Clesse, J. H. Davis, et al., *Physics of the Dark Universe* **12**, 56 (2016), ISSN 2212-6864.
 - [2] D. Rodrigues, V. Marra, A. del Popolo, and Z. Davari, *Nat.Astron.* 2 p. 668-672 (2018), arXiv:1806.06803 [astro-ph.GA].
 - [3] H. M. Gauy and J. Ramos-Caro, *Phys. Rev. D* **97**, 064012 (2018).
 - [4] A. Joyce, B. Jain, J. Khoury, and M. Trodden, *Physics Reports* **568**, 1 (2015), ISSN 0370-1573, beyond the cosmological standard model.
 - [5] A. S. Bolton, S. Rappaport, and S. Burles, *Phys. Rev. D* **74**, 061501 (2006).
 - [6] M. V. d. Santos, J. S. Alcaniz, D. F. Mota, and S. Capozziello, *Phys. Rev. D* **97**, 104010 (2018).
 - [7] D. Psaltis, *Living Reviews in Relativity* **11**, 9 (2008), ISSN 1433-8351.
 - [8] V. C. Rubin, J. W. Kent Ford, and N. Thonnard, *Astrophys. J.* **238**, 471 (1980).
 - [9] V. C. Rubin, D. Burstein, J. W. Kent Ford, and N. Thonnard, *Astrophys. J.* **289**, 81 (1985).
 - [10] A. Avilez-Lopez, A. Padilla, P. M. Saffin, and C. Skordis, *J. Cosmol. Astropart. Phys.* **2015**, 044 (2015).
 - [11] J. Binney and S. Tremaine, *Galactic Dynamics* (Princeton Univ. Press, Princeton, NJ, 2008), 2nd ed.
 - [12] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, *Physics Reports* **513**, 1 (2012), issues 1?3.
 - [13] V. Salzano, D. F. Mota, M. P. Da?browski, and S. Capozziello, *J. Cosmol. Astropart. Phys.* **2016**, 033 (2016).
 - [14] M. H. Chan and H. K. Hui, *Astrophys. J.* **856**, 177 (2018).
 - [15] X. Cui, A. Abdukerim, W. Chen, X. Chen, Y. Chen, B. Dong, D. Fang, C. Fu, K. Giboni, F. Giuliani, et al. (PandaX-II Collaboration), *Phys. Rev. Lett.* **119**, 181302 (2017).
 - [16] G. Aad, T. Abajyan, B. Abbott, J. Abdallah, S. Ab-

- del Khalek, O. Abdinov, R. Aben, B. Abi, M. Abolins, O. S. AbouZeid, et al. (ATLAS Collaboration), *Phys. Rev. D* **90**, 012004 (2014).
- [17] D. S. Akerib, S. Alsum, H. M. Araújo, X. Bai, A. J. Bailey, J. Balajthy, P. Beltrame, E. P. Bernard, A. Bernstein, T. P. Biesiadzinski, et al. (LUX Collaboration), *Phys. Rev. Lett.* **118**, 021303 (2017).
- [18] E. Aprile, J. Aalbers, F. Agostini, M. Alfonsi, F. D. Amaro, M. Anthony, F. Arneodo, P. Barrow, L. Baudis, B. Bauermeister, et al. (XENON Collaboration), *Phys. Rev. Lett.* **119**, 181301 (2017).
- [19] A. Einstein, L. Infeld, and B. Hoffmann, *Ann. Math.* **39**, 65 (1938).
- [20] A. Einstein and L. Infeld, *Ann. Math.* **41**, 455 (1940).
- [21] A. Einstein and L. Infeld, *Canad. J. Math.* **1**, 209 (1949).
- [22] E. Babichev and C. Deffayet, *Classical and Quantum Gravity* **30**, 18 (2013).
- [23] B. Famaey and J. Binney, *Mon. Not. R. Astron. Soc.* **363**, 603 (2005).
- [24] B. Famaey and S. S. McGaugh, *Living Reviews in Relativity* **15**, 10 (2012).
- [25] J. B. Jiménez, A. L. D. Fróes, and D. F. Mota, *Physics Letters B* **725**, 212 (2013), issues 4-5.
- [26] R. Massey, T. Kitching, and J. Richard, *Reports on Progress in Physics* **73**, 086901 (2010).
- [27] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, et al. (LIGO Scientific Collaboration and Virgo Collaboration), *Phys. Rev. Lett.* **116**, 061102 (2016).
- [28] M. Milgrom, *Astrophys. J.* **270**, 365 (1983).
- [29] M. Milgrom, *Astrophys. J.* **270**, 371 (1983).
- [30] M. Milgrom, *Astrophys. J.* **270**, 384 (1983).
- [31] M. Milgrom, *Mon. Not. R. Astron. Soc.* **403**, 886 (2010).
- [32] M. Milgrom, *Mon. Not. R. Astron. Soc.* **437**, 2531 (2014).
- [33] J. Bekenstein and M. Milgrom, *Astrophys. J.* **286**, 7 (1984).
- [34] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman and Company, San Francisco, 1973).
- [35] J. D. Bekenstein, *Phys. Rev. D* **70**, 083509 (2004).
- [36] H. C. Plummer, *Mon. Not. R. Astron. Soc.* **71**, 460 (1911).
- [37] H. Dejonghe, *Mon. Not. R. Astron. Soc.* **224**, 13 (1987).
- [38] C. A. Agón, J. F. Pedraza, and J. Ramos-Caro, *Phys. Rev. D* **83**, 123007 (2011).
- [39] J. Ramos-Caro, C. A. Agón, and J. F. Pedraza, *Phys. Rev. D* **86**, 043008 (2012).
- [40] V. Salzano, D. F. Mota, M. P. Da?browski, and S. Capozziello, *J. Cosmol. Astropart. Phys.* **2016**, 033 (2016).
- [41] R. H. Sanders and S. S. McGaugh, *Annu. Rev. Astron. Astrophys.* **40**, 263 (2002).
- [42] S. McGaugh, *Galaxies* pp. 601–622 (2014).
- [43] S. S. McGaugh, *Astrophys. J.* **143**, 40 (2012).
- [44] R. H. Sanders and S. S. McGaugh, *Annual Review of Astronomy and Astrophysics* **40**, 263 (2002), <https://doi.org/10.1146/annurev.astro.40.060401.093923>.
- [45] T. Baker, D. Psaltis, and C. Skordis, *Astrophys. J.* **802**, 63 (2015).
- [46] C. M. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, Cambridge, 1993).
- [47] E. Poisson and M. W. Clifford, *Gravity Newtonian, Post-Newtonian, Relativistic*. (Cambridge University Press, New York, 2014).
- [48] C. M. Will, *Living Reviews in Relativity* (2014).
- [49] K. S. Thorne and C. M. Will, *Astrophys. J.* **163**, 595 (1971).
- [50] C. M. Will, *Astrophys. J.* **163**, 611-628 (1971).
- [51] C. M. Will and J. K. L. Nordtvedt, *Astrophys. J.* **177**, 757-774 (1972).
- [52] Y. Choquet-Bruhat, *Introduction to General Relativity, Black Holes, and Cosmology* (Oxford University Press, 2015).
- [53] C. Cercignani and G. M. Kremer, *The Relativistic Boltzmann Equation: Theory and Applications* (Birkhäuser, Basel, 2002).
- [54] N. Bolis, C. Skordis, D. B. Thomas, and T. Złóśnik, *Phys. Rev. D* **99**, 084009 (2019), URL <https://link.aps.org/doi/10.1103/PhysRevD.99.084009>.
- [55] R. McManus, L. Lombriser, and J. Peñarrubia, *Journal of Cosmology and Astroparticle Physics* **2017**, 031 (2017), URL <https://doi.org/10.1088%2F1475-7516%2F2017%2F12%2F031>.