

Universal birational invariants and \mathbb{A}^1 -homology

Yuri Shimizu *

February 17, 2020

Abstract

Let k be a field admitting a resolution of singularities. In this paper, we prove that the functor of zeroth \mathbb{A}^1 -homology $\mathbf{H}_0^{\mathbb{A}^1}$ is universal as a functorial birational invariant of smooth proper k -varieties taking values in a category enriched by abelian groups. For a smooth proper k -variety X , we also prove that the dimension of $\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Q})(\mathrm{Spec} k)$ coincides with the number of R -equivalence classes of $X(k)$. We deduce these results as consequences of the structure theorem that for a smooth proper k -variety X , the sheaf $\mathbf{H}_0^{\mathbb{A}^1}(X)$ is the free abelian presheaf generated by the birational \mathbb{A}^1 -connected components $\pi_0^{b\mathbb{A}^1}(X)$ of Asok-Morel.

Contents

1	Birational sheaves and localizations of categories	4
2	Birational \mathbb{A}^1-connected components	8
3	Birational sheaves of modules	13
4	Structure of zeroth \mathbb{A}^1- and Suslin homology	17
5	Universal birational invariance	22
6	Applications to \mathbb{A}^1-homotopy theory	28

Introduction

Let k be a field. In [Mor05], Morel introduced the \mathbb{A}^1 -homology theory of smooth (separated and of finite type) k -schemes. In the unstable \mathbb{A}^1 -homotopy theory introduced by Morel-Voevodsky [MV99], the \mathbb{A}^1 -homology plays a role of the ordinary homology of topological spaces. For each $n \geq 0$, the n -th \mathbb{A}^1 -homology

*Department of Mathematics, Tokyo Institute of Technology
E-mail: qingshuiyouli27@gmail.com

of a smooth k -scheme X is denoted by $\mathbf{H}_n^{\mathbb{A}^1}(X)$. This is a Nisnevich sheaf of abelian groups on the category of smooth k -schemes $\mathcal{S}m_k$. As an application of \mathbb{A}^1 -homology, Asok [Aso12] proved that the zeroth \mathbb{A}^1 -homology is a birational invariant of smooth proper k -varieties if k is infinite. In this paper, we study the structure of the zeroth \mathbb{A}^1 -homology of smooth proper k -varieties and give some applications. In particular, we prove that the zeroth \mathbb{A}^1 -homology functor $\mathbf{H}_0^{\mathbb{A}^1}$ is universal as a functorial birational invariant of smooth proper k -varieties.

First, we state the universal birational invariance of the zeroth \mathbb{A}^1 -homology. Let $\mathcal{S}m_k^{prop}$ be the category of smooth proper k -schemes and $\mathbf{Im} \mathbf{H}_0^{\mathbb{A}^1}$ be the full subcategory of the category of abelian presheaves on $\mathcal{S}m_k$ spanned by objects isomorphic to $\mathbf{H}_0^{\mathbb{A}^1}(X)$ for some $X \in \mathcal{S}m_k^{prop}$. Our first theorem is stated as follows.

Theorem 1 (see Theorem 5.5). *Assume k admits a resolution of singularities. Let \mathcal{A} be an arbitrary category enriched by abelian groups (e.g. an abelian category).*

- (1) *Let $F : \mathcal{S}m_k^{prop} \rightarrow \mathcal{A}$ be an arbitrary functor which sends each birational morphism to an isomorphism. Then there exists one and only one (up to a natural equivalence) additive functor $F_{S_b} : \mathbf{Im} \mathbf{H}_0^{\mathbb{A}^1} \rightarrow \mathcal{A}$ such that the diagram*

$$\begin{array}{ccc} \mathcal{S}m_k^{prop} & \xrightarrow{F} & \mathcal{A} \\ \mathbf{H}_0^{\mathbb{A}^1} \downarrow & \nearrow F_{S_b} & \\ \mathbf{Im} \mathbf{H}_0^{\mathbb{A}^1} & & \end{array}$$

is 2-commutative, i.e., we have a natural equivalence $F \cong F_{S_b} \circ \mathbf{H}_0^{\mathbb{A}^1}$.

- (2) *Let $F' : (\mathcal{S}m_k^{prop})^{op} \rightarrow \mathcal{A}$ be an arbitrary functor which sends each birational morphism to an isomorphism. Then there exists one and only one (up to a natural equivalence) additive functor $F'_{S_b} : (\mathbf{Im} \mathbf{H}_0^{\mathbb{A}^1})^{op} \rightarrow \mathcal{A}$ such that the diagram*

$$\begin{array}{ccc} (\mathcal{S}m_k^{prop})^{op} & \xrightarrow{F'} & \mathcal{A} \\ (\mathbf{H}_0^{\mathbb{A}^1})^{op} \downarrow & \nearrow F'_{S_b} & \\ (\mathbf{Im} \mathbf{H}_0^{\mathbb{A}^1})^{op} & & \end{array}$$

is 2-commutative.

In this theorem, we may replace $\mathcal{S}m_k^{prop}$ with other full subcategory of $\mathcal{S}m_k^{prop}$ like the category of projective varieties (see Definition 5.3 and Example 5.4). A similar result also holds in a motivic situation (see Theorem 5.10). In this case, $\mathcal{S}m_k^{prop}$ is replaced with the full subcategory of the category of finite correspondences consisting of proper schemes and the \mathbb{A}^1 -homology with the zeroth homology of Voevodsky's motives, called Suslin homology sheaves (cf. [Voe00]). The birational invariance theorem of Asok [Aso12] is assumed

k infinite, but this assumption is not necessary for our proof. In general, we prove that a proper birational morphism of (not necessary proper) varieties over an arbitrary field induces an isomorphism of zeroth \mathbb{A}^1 -homology sheaves. (see Proposition 5.12).

Second, we relate the zeroth \mathbb{A}^1 -homology to rational points. We consider the zeroth \mathbb{A}^1 -homology with \mathbb{Q} -coefficients $\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Q})$ for a smooth proper k -variety X .

Theorem 2 (see Theorem 4.7). *Assume k admits a resolution of singularities. For a smooth proper k -variety X , we have*

$$\dim_{\mathbb{Q}} \mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Q})(\mathrm{Spec} k) = \#(X(k)/R).$$

Moreover, $\mathbf{H}_0^{\mathbb{A}^1}(X)(\mathrm{Spec} k) = 0$ if and only if $X(k) = \emptyset$.

Here $X(k)/R$ is the quotient set of $X(k)$ by the R -equivalence introduced by Manin [Man86]. Note that the abelian group $\mathbf{H}_0^{\mathbb{A}^1}(X)(\mathrm{Spec} k)$ can also be expressed in terms of triangulated categories (see Remark 4.10).

Third, we state a structure theorem of zeroth \mathbb{A}^1 -homology sheaves. In [AM11], Asok-Morel constructed a Nisnevich sheaf of sets $\pi_0^{b\mathbb{A}^1}(X)$ on $\mathcal{S}m_k$, called the birational \mathbb{A}^1 -connected components, for each $X \in \mathcal{S}m_k^{prop}$. We generalize this construction for all smooth k -schemes. Our structure theorem is stated as follows.

Theorem 3 (see Theorem 4.1). *Assume k admits a resolution of singularities. For every $X \in \mathcal{S}m_k$, there exists a natural epimorphism of presheaves*

$$\mathbb{Z}_{pre}(\pi_0^{b\mathbb{A}^1}(X)) \twoheadrightarrow \mathbf{H}_0^{\mathbb{A}^1}(X).$$

Moreover, this is an isomorphism if X is proper.

Here $\mathbb{Z}_{pre}(\pi_0^{b\mathbb{A}^1}(X))$ is the free abelian presheaf generated by $\pi_0^{b\mathbb{A}^1}(X)$. A similar result also holds for Suslin homology sheaves (see Theorem 4.11). Theorem 3 has some applications to \mathbb{A}^1 -homotopy theory (see §6). Moreover, the Suslin homology version of Theorem 3 also has an application to zero cycles (see Corollary 4.12). Theorems 1 and 2 are consequences of Theorem 3.

This paper is organized as follows. In §1, we recall and prepare basic results on birational sheaves and the localization of some categories of smooth k -schemes by birational morphisms. In §2, we generalize birational \mathbb{A}^1 -connected components of Asok-Morel for all Nisnevich sheaves on $\mathcal{S}m_k$. In §3, we give a relationship between birational sheaves and strictly \mathbb{A}^1 -invariant sheaves. In §4, we prove Theorems 2 and 3. In §5, we prove Theorem 1. In §6, we give applications to \mathbb{A}^1 -homotopy theory.

Conventions. Throughout this paper, we fix a field k and a commutative unital ring Λ . All k -varieties are assumed irreducible but not assumed geometrically irreducible. The field k is said to admit a resolution of singularities, if k -varieties always have a resolution of singularities. Let $\mathcal{S}m_k$ be the category of separated

and smooth k -schemes of finite type. Objects of $\mathcal{S}m_k$ are simply called smooth k -schemes. We regard every smooth k -scheme X as the presheaf of sets on $\mathcal{S}m_k$ represented by X . For a presheaf \mathcal{F} on $\mathcal{S}m_k$ and an affine scheme $\text{Spec } A$ which is the limit of an inversed system $\{U_\lambda\}_\lambda$ in $\mathcal{S}m_k$, we write

$$\mathcal{F}(A) = \text{colim}_\lambda \mathcal{F}(U_\lambda).$$

For a category \mathcal{C} , we denote $\mathcal{P}resh(\mathcal{C})$ (resp. $\mathcal{P}resh(\mathcal{C}, \Lambda)$) for the category of presheaves of sets (resp. Λ -modules) on \mathcal{C} . A diagram of categories

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ G' \downarrow & & \downarrow G \\ \mathcal{C}' & \xrightarrow{F'} & \mathcal{D}' \end{array}$$

is called 2-commutative, if there exists a natural equivalence $G \circ F \cong F' \circ G'$. For a local ring A , we denote κ_A for its residue field.

Acknowledgments. I would like to thank my adviser Shohei Ma for many useful advices.

1 Birational sheaves and localizations of categories

In this section, we consider localizations of categories in the sense of Gabriel-Zisman [GZ67]. Let \mathcal{C} be a category and S be a family of morphisms in \mathcal{C} . Recall that the localization of \mathcal{C} by S is a category $S^{-1}\mathcal{C}$ with a morphism $\mathcal{L} : \mathcal{C} \rightarrow S^{-1}\mathcal{C}$ such that

- for every $s \in S$, the image $\mathcal{L}(s)$ is an isomorphism in $S^{-1}\mathcal{C}$, and
- for every category \mathcal{D} and every functor $\mathcal{C} \rightarrow \mathcal{D}$ which sends each $s \in S$ to an isomorphism in \mathcal{D} , there exists one and only one (up to a natural equivalence) functor $S^{-1}\mathcal{C} \rightarrow \mathcal{D}$ such that the diagram

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \mathcal{D} \\ \downarrow & & \nearrow \\ S^{-1}\mathcal{C} & & \end{array}$$

is 2-commutative.

Note that the induced functor $\mathcal{P}resh(S^{-1}\mathcal{C}) \rightarrow \mathcal{P}resh(\mathcal{C})$ is fully faithful and its essential image is spanned by presheaves which send each $s \in S$ for a bijection. We especially treat some cases where \mathcal{C} is a category of smooth k -schemes and S consists of all birational morphisms in \mathcal{C} .

1.1 Localizations by birational morphisms

Let $\mathcal{S}m_k^{var}$ (resp. $\mathcal{S}m_k^{prop}$, $\mathcal{S}m_k^{pv}$) be the full subcategory of $\mathcal{S}m_k$ consisting of irreducible (resp. proper, proper and irreducible) k -schemes. For a subcategory $\mathcal{C} \subseteq \mathcal{S}m_k$, we write $S_b^{-1}\mathcal{C}$ for the localization of \mathcal{C} by birational morphisms within \mathcal{C} . In [KS07], Kahn-Sujatha proved that the inclusion $\mathcal{S}m_k^{pv} \hookrightarrow \mathcal{S}m_k^{var}$ induces an equivalence of categories

$$S_b^{-1}\mathcal{S}m_k^{pv} \xrightarrow{\cong} S_b^{-1}\mathcal{S}m_k^{var} \quad (1.1)$$

if k admits a resolution of singularities (see [KS07, Prop. 8.5]).

Let $Cor_{k,\Lambda}$ be the category of finite correspondences of Voevodsky with coefficients in a commutative unital ring Λ (see definition [MVW, Def. 1.5]). We denote Γ for the canonical functor $\mathcal{S}m_k \rightarrow Cor_{k,\Lambda}$ and write

$$Cor_{k,\Lambda}(X, Y) = \text{Hom}_{Cor_{k,\Lambda}}(X, Y)$$

for each $X, Y \in Cor_{k,\Lambda}$. For a subcategory $\mathcal{C} \subseteq Cor_{k,\Lambda}$, we denote $S_b^{-1}\mathcal{C}$ for the localization of \mathcal{C} by finite correspondences associated with a birational morphism. Let $Cor_{k,\Lambda}^{var}$ (resp. $Cor_{k,\Lambda}^{prop}$, $Cor_{k,\Lambda}^{pv}$) be the full subcategory of $Cor_{k,\Lambda}$ consisting of irreducible (resp. proper, proper and irreducible) k -schemes. Next, we give an analogue of the equivalence (1.1) for categories of finite correspondences. For this, we prove the following lemma.

Lemma 1.1. *For all $X_0, X, Y \in Cor_{k,\Lambda}$ and every birational morphism $f : X_0 \rightarrow X$ in $\mathcal{S}m_k$, the induced map*

$$f^* : Cor_{k,\Lambda}(X, Y) \rightarrow Cor_{k,\Lambda}(X_0, Y); c \mapsto c \circ \Gamma(f)$$

is injective.

Proof. Let $i : U \hookrightarrow X$ be a dense open embedding that $f^{-1}(i(U)) \rightarrow i(U)$ is an isomorphism. By the commutativity of the diagram

$$\begin{array}{ccc} Cor_{k,\Lambda}(X, Y) & \xrightarrow{f^*} & Cor_{k,\Lambda}(X_0, Y) \\ i^* \downarrow & & \downarrow \\ Cor_{k,\Lambda}(U, Y) & \xrightarrow{\cong} & Cor_{k,\Lambda}(f^{-1}(U), Y), \end{array}$$

we only need to show that i^* is injective. Note that for every $c \in Cor_{k,\Lambda}(X, Y)$ the finite correspondence $i^*c \in Cor_{k,\Lambda}(U, Y)$ coincides with the pullback of c by the open embedding

$$j : U \times Y \hookrightarrow X \times Y$$

as an algebraic cycle of $U \times Y$. We write $W = (X \times Y) - j(U \times Y)$. Now we obtain an equality

$$\text{Ker}(Z(X \times Y) \xrightarrow{j^*} Z(U \times Y)) = Z(W),$$

where $Z(-)$ means the set of algebraic cycles with Λ -coefficients. Thus we have

$$\begin{aligned} & \text{Ker}(\text{Cor}_{k,\Lambda}(X, Y) \xrightarrow{i^*} \text{Cor}_{k,\Lambda}(U, Y)) \\ &= \text{Cor}_{k,\Lambda}(X, Y) \cap \text{Ker}(Z(X \times Y) \xrightarrow{j^*} Z(U \times Y)) \\ &= \text{Cor}_{k,\Lambda}(X, Y) \cap Z(W). \end{aligned}$$

On the other hand, since $U \times Y$ is non-empty, the map $W \rightarrow X$ is not surjective. Therefore, all non-zero elements of $Z(W)$ are not finite correspondences $X \rightarrow Y$. Thus we have

$$\text{Ker } i^* = \text{Cor}_{k,\Lambda}(X, Y) \cap Z(W) = 0.$$

□

We prove that $S_b^{-1}\text{Cor}_{k,\Lambda}^{pv} \cong S_b^{-1}\text{Cor}_{k,\Lambda}^{var}$.

Proposition 1.2. *Assume k admits a resolution of singularities. Then the inclusion $\text{Cor}_{k,\Lambda}^{pv} \hookrightarrow \text{Cor}_{k,\Lambda}^{var}$ induces an equivalence of categories*

$$S_b^{-1}\text{Cor}_{k,\Lambda}^{pv} \xrightarrow{\cong} S_b^{-1}\text{Cor}_{k,\Lambda}^{var}.$$

Proof. By [KS07, Thm. 2.1 and 4.3], we only need to show that the functors $\text{Cor}_{k,\Lambda}^{pv} \hookrightarrow \text{Cor}_{k,\Lambda}^{var}$ and $\text{Cor}_{k,\Lambda}^{prop} \hookrightarrow \text{Cor}_{k,\Lambda}$ satisfy the conditions (b1)-(b3) in [KS07, Prop. 5.10]. Then (b1) follows from Lemma 1.1 and (b2) and (b3) follow from a similar argument as [KS07, proof of Prop. 8.5] (see also [KS07, proof of Prop. 8.4]). □

Remark 1.3. By a similar proof, we also obtain an equivalence of categories

$$S_b^{-1}\text{Cor}_{k,\Lambda} \xrightarrow{\cong} S_b^{-1}\text{Cor}_{k,\Lambda}^{prop}.$$

1.2 Birational sheaves

Following Asok-Morel [AM11], we call a presheaf \mathcal{F} on $\mathcal{S}m_k$ a *birational sheaf*, if

- B1** the canonical map $\mathcal{F}(U \sqcup V) \rightarrow \mathcal{F}(U) \times \mathcal{F}(V)$ is bijective for all $U, V \in \mathcal{S}m_k$, and
- B2** every open embedding $U \hookrightarrow X$ in $\mathcal{S}m_k$ induces a bijection $\mathcal{F}(X) \xrightarrow{\cong} \mathcal{F}(U)$.

Note that birational sheaves are always Nisnevich (see [AM11, Lem. 6.1.2]). Let $\mathcal{S}hv_k$ be the category of Nisnevich sheaves of sets on $\mathcal{S}m_k$. We denote $\mathcal{S}hv_k^{br}$ for the full subcategory of $\mathcal{S}hv_k$ consisting of birational sheaves. We give a canonical equivalence $\mathcal{S}hv_k^{br} \cong \text{Presh}(S_b^{-1}\mathcal{S}m_k^{var})$.

Lemma 1.4. *There exists an equivalence of categories*

$$\mathcal{S}hv_k^{br} \xrightarrow{\cong} \text{Presh}(S_b^{-1}\mathcal{S}m_k^{var})$$

such that the diagram

$$\begin{array}{ccc} Shv_k^{br} & \xrightarrow{\cong} & Presh(S_b^{-1}Sm_k^{var}) \\ \downarrow & & \downarrow \\ Shv_k & \longrightarrow & Presh(Sm_k^{var}) \end{array}$$

is 2-commutative.

Proof. Let $Presh_{\mathbf{B2}}(Sm_k^{var})$ be the full subcategory of $Presh(Sm_k^{var})$ consisting of presheaves which satisfy **B2**. Then $Presh_{\mathbf{B2}}(Sm_k^{var})$ is equivalent to $Presh(S_b^{-1}Sm_k^{var})$ by the universality of localizations. On the other hand, since every object in Sm_k is a finite coproduct of objects in Sm_k^{var} , the restriction functor $Shv_k^{br} \rightarrow Presh_{\mathbf{B2}}(Sm_k^{var})$ is an equivalence by **B1**. Thus we have a 2-commutative the diagram

$$\begin{array}{ccccc} Shv_k^{br} & \xrightarrow{\cong} & Presh_{\mathbf{B2}}(Sm_k^{var}) & \xleftarrow{\cong} & Presh(S_b^{-1}Sm_k^{var}) \\ \downarrow & & \downarrow & & \downarrow \\ Shv_k & \longrightarrow & Presh(Sm_k^{var}) & \xlongequal{\quad} & Presh(Sm_k^{var}). \end{array}$$

□

For a category \mathcal{C} enriched by Λ -modules, we denote $\mathcal{L}in(\mathcal{C}, \Lambda)$ for the category of Λ -linear presheaves of Λ -modules on \mathcal{C} . We write

$$\mathbf{PST}_k(\Lambda) = \mathcal{L}in(Cor_{k,\Lambda}, \Lambda).$$

Objects in $\mathbf{PST}_k(\Lambda)$ are called a presheaves with transfers. A presheaf with transfers M is called Nisnevich (resp. birational), if so is the direct image Γ_*M by the canonical functor $\Gamma : Sm_k \rightarrow Cor_{k,\Lambda}$. We denote $\mathbf{NST}_k(\Lambda)$ (resp. $\mathbf{NST}_k^{br}(\Lambda)$) for the category of Nisnevich (resp. birational) sheaves with transfers. We prove an analogue of Lemma 1.4 for sheaves with transfers.

Lemma 1.5. *There exists an equivalence of categories*

$$\mathbf{NST}_k^{br}(\Lambda) \xrightarrow{\cong} \mathcal{L}in(S_b^{-1}Cor_{k,\Lambda}^{var}, \Lambda)$$

such that the diagram

$$\begin{array}{ccc} \mathbf{NST}_k^{br}(\Lambda) & \xrightarrow{\cong} & \mathcal{L}in(S_b^{-1}Cor_{k,\Lambda}^{var}, \Lambda) \\ \downarrow & & \downarrow \\ \mathbf{NST}_k(\Lambda) & \longrightarrow & \mathcal{L}in(Cor_{k,\Lambda}^{var}, \Lambda) \end{array}$$

is 2-commutative.

Proof. Let $\mathcal{L}in_{\mathbf{B2}}(\mathcal{C}or_{k,\Lambda}^{var}, \Lambda)$ be the full subcategory of $\mathcal{L}in(\mathcal{C}or_{k,\Lambda}^{var}, \Lambda)$ consisting of presheaves which satisfy **B2**. By a similar argument as the proof of Lemma 1.4, we also have a 2-commutative diagram

$$\begin{array}{ccccc} \mathbf{NST}_k^{br}(\Lambda) & \xrightarrow{\cong} & \mathcal{L}in_{\mathbf{B2}}(\mathcal{C}or_{k,\Lambda}^{var}, \Lambda) & \xleftarrow{\cong} & \mathcal{L}in(S_b^{-1}\mathcal{C}or_{k,\Lambda}^{var}, \Lambda) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbf{NST}_k(\Lambda) & \longrightarrow & \mathcal{L}in(\mathcal{C}or_{k,\Lambda}^{var}, \Lambda) & \xlongequal{\quad} & \mathcal{L}in(\mathcal{C}or_{k,\Lambda}^{var}, \Lambda). \end{array}$$

□

The following lemma says that birational sheaves are \mathbb{A}^1 -invariant.

Lemma 1.6. *For every $\mathcal{F} \in \mathcal{S}hv_k^{br}$ and every $U \in \mathcal{S}m_k^{var}$, the canonical map*

$$\mathcal{F}(U) \rightarrow \mathcal{F}(U \times \mathbb{A}^1)$$

is an isomorphism.

Proof. By [KS15, Thm. 1.7.9], stable birational morphisms are isomorphisms in $S_b^{-1}\mathcal{S}m_k^{var}$. Since the projection $U \times \mathbb{A}^1 \rightarrow U$ is stable biratoyal, we have $\mathcal{F}(U) \cong \mathcal{F}(U \times \mathbb{A}^1)$ by Lemma 1.4. □

2 Birational \mathbb{A}^1 -connected components

In [AM11], Asok-Morel introduced a birational sheaf $\pi_0^{b\mathbb{A}^1}(X)$, called the *birational \mathbb{A}^1 -connected components*, for each $X \in \mathcal{S}m_k^{prop}$. In this section, we generalize this construction for all Nisnevich sheaves on $\mathcal{S}m_k$. We first construct a functor $\pi_0^{br} : \mathcal{S}hv_k \rightarrow \mathcal{S}hv_k^{br}$ and secondly prove that $\pi_0^{br}(X) \cong \pi_0^{b\mathbb{A}^1}(X)$ for every $X \in \mathcal{S}m_k^{prop}$. Moreover, we also prove that the functor π_0^{br} is left adjoint to the inclusion $\mathcal{S}hv_k^{br} \hookrightarrow \mathcal{S}hv_k$. Similarly, we construct a left adjoint functor of the inclusion $\mathbf{NST}_k^{br}(\Lambda) \hookrightarrow \mathbf{NST}_k(\Lambda)$.

2.1 Birational \mathbb{A}^1 -connected components of sheaves

First, we give a functor $\pi_0^{br} : \mathcal{S}hv_k \rightarrow \mathcal{S}hv_k^{br}$.

Definition 2.1. We define a functor $\pi_0^{br} : \mathcal{S}hv_k \rightarrow \mathcal{S}hv_k^{br}$ as the composition

$$\mathcal{S}hv_k \hookrightarrow \mathcal{P}resh(\mathcal{S}m_k) \rightarrow \mathcal{P}resh(\mathcal{S}m_k^{var}) \rightarrow \mathcal{S}hv_k^{br},$$

where the third functor is a left Kan extension of

$$\mathcal{S}m_k^{var} \rightarrow S_b^{-1}\mathcal{S}m_k^{var} \hookrightarrow \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{var}) \cong \mathcal{S}hv_k^{br}.$$

Explicitly, every sheaf \mathcal{F} on $\mathcal{S}m_k$ is canonically isomorphic to the colimit of a direct system of smooth k -varieties $\{X_\lambda\}_\lambda$ (note that every object in $\mathcal{S}m_k$ is

a coproduct of smooth k -varieties). Then by the definition, we have a natural bijection

$$\pi_0^{br}(\mathcal{F})(U) \cong \operatorname{colim}_{\lambda} \operatorname{Hom}_{S_b^{-1}\mathcal{S}m_k^{var}}(U, X_{\lambda})$$

for every $U \in \mathcal{S}m_k$. Thus the map

$$\operatorname{Hom}_{\mathcal{S}m_k}(U, X_{\lambda}) \rightarrow \operatorname{Hom}_{S_b^{-1}\mathcal{S}m_k^{var}}(U, X_{\lambda})$$

induces a natural morphism

$$\mathcal{F} \rightarrow \pi_0^{br}(\mathcal{F})$$

in $\mathcal{S}hv_k$ which is functorial for \mathcal{F} .

Let X be a smooth proper k -variety. Recall that two points in $X(K)$ for a field extension K/k are called R -equivalence (in the sense of Manin [Man86]), if these points are connected by the image of a chain of k -morphisms $\mathbb{P}_K^1 \rightarrow X$. The quotient set of $X(K)$ by the R -equivalence is denoted by $X(K)/R$. The birational \mathbb{A}^1 -connected components $\pi_0^{b\mathbb{A}^1}(X)$ has the property that there exists a canonical bijection

$$X(K)/R \cong \pi_0^{b\mathbb{A}^1}(X)(K)$$

for every finitely generated separable extension K/k (see [AM11, Thm. 6.2.1]). Our sheaf $\pi_0^{br}(X)$ also has the same property.

Lemma 2.2. *Let X be a smooth proper k -scheme and K/k be a finitely generated separable extension. Then the map $X(K) \rightarrow \pi_0^{br}(X)(K)$ induced by the canonical morphism $X \rightarrow \pi_0^{br}(X)$ factors through a bijection*

$$X(K)/R \cong \pi_0^{br}(X)(K).$$

Proof. This follows from the bijection

$$X(K)/R \cong \operatorname{Hom}_{S_b^{-1}\mathcal{S}m_k^{var}}(U, X)$$

in [KS15, Thm. 6.6.3], where U is a smooth model of K . \square

Next, we give an isomorphism $\pi_0^{br}(X) \cong \pi_0^{b\mathbb{A}^1}(X)$ for each $X \in \mathcal{S}m_k^{prop}$.

Proposition 2.3. *For every $X \in \mathcal{S}m_k^{prop}$, there exists an isomorphism of birational sheaves*

$$\pi_0^{br}(X) \cong \pi_0^{b\mathbb{A}^1}(X).$$

Before proving this proposition, we recall the category $\mathcal{F}_k^r - \mathcal{S}et$ introduced by Asok-Morel [AM11]. Let \mathcal{F}_k be the category of finitely generated separable extension fields over k . Each object $\mathcal{S} \in \mathcal{F}_k^r - \mathcal{S}et$ is a covariant functor

$$\mathcal{F}_k \rightarrow \mathcal{S}et$$

together with a map $\mathcal{S}(K) \rightarrow \mathcal{S}(\kappa_A)$ for each $K \in \mathcal{F}_k$ and its discrete valuation ring A with $\kappa_A \in \mathcal{F}_k$. Moreover, a morphism $\mathcal{S} \rightarrow \mathcal{S}'$ in $\mathcal{F}_k^r - \mathcal{S}et$ is a natural transformation $\mathcal{S} \rightarrow \mathcal{S}'$ such that the diagram

$$\begin{array}{ccc} \mathcal{S}(K) & \longrightarrow & \mathcal{S}(\kappa_A) \\ \downarrow & & \downarrow \\ \mathcal{S}'(K) & \longrightarrow & \mathcal{S}'(\kappa_A) \end{array}$$

commutes. The restriction of each $\mathcal{F} \in \mathcal{S}hw_k^{br}$ on \mathcal{F}_k together with the map

$$\mathcal{F}(K) \cong \mathcal{F}(A) \rightarrow \mathcal{F}(\kappa_A)$$

is an object of $\mathcal{F}_k^r - \mathcal{S}et$. Thus we have the restriction functor

$$\text{Res} : \mathcal{S}hw_k^{br} \rightarrow \mathcal{F}_k^r - \mathcal{S}et. \quad (2.1)$$

Proof of Proposition 2.3. By Lemma 1.6 and [AM11, Thm. 6.1.7], the restriction functor (2.1) is fully faithful. Thus we only need to construct an isomorphism

$$\text{Res}(\pi_0^{br}(X)) \cong \text{Res}(\pi_0^{b\mathbb{A}^1}(X)) \quad (2.2)$$

in $\mathcal{F}_k^r - \mathcal{S}et$. Lemma 2.2 and [AM11, Thm. 6.2.1] give a commutative diagram

$$\begin{array}{ccccc} X(K) & \xlongequal{\quad} & X(K) & \xlongequal{\quad} & X(K) \\ \downarrow & & \downarrow & & \downarrow \\ \pi_0^{br}(X)(K) & \xrightarrow{\cong} & X(K)/R & \xleftarrow{\cong} & \pi_0^{b\mathbb{A}^1}(X)(K) \end{array}$$

for all $K \in \mathcal{F}_k$. Then the vertical maps are surjective. Thus for every extension field L/K which is finitely generated and separable over k , the obviously commutative diagram

$$\begin{array}{ccccc} X(K) & \xlongequal{\quad} & X(K) & \xlongequal{\quad} & X(K) \\ \downarrow & & \downarrow & & \downarrow \\ X(L) & \xlongequal{\quad} & X(L) & \xlongequal{\quad} & X(L) \end{array}$$

shows that

$$\begin{array}{ccccc} \pi_0^{br}(X)(K) & \xrightarrow{\cong} & X(K)/R & \xleftarrow{\cong} & \pi_0^{b\mathbb{A}^1}(X)(K) \\ \downarrow & & \downarrow & & \downarrow \\ \pi_0^{br}(X)(L) & \xrightarrow{\cong} & X(L)/R & \xleftarrow{\cong} & \pi_0^{b\mathbb{A}^1}(X)(L) \end{array}$$

commutes. Similarly, for every discrete valuation ring A of K with $\kappa_A \in \mathcal{F}_k$, the diagram

$$\begin{array}{ccccc} X(K) & \xlongequal{\quad} & X(K) & \xlongequal{\quad} & X(K) \\ \downarrow & & \downarrow & & \downarrow \\ X(\kappa_A) & \xlongequal{\quad} & X(\kappa_A) & \xlongequal{\quad} & X(\kappa_A) \end{array}$$

shows that

$$\begin{array}{ccccc}
\pi_0^{br}(X)(K) & \xrightarrow{\cong} & X(K)/R & \xleftarrow{\cong} & \pi_0^{b\mathbb{A}^1}(X)(K) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_0^{br}(X)(\kappa_A) & \xrightarrow{\cong} & X(\kappa_A)/R & \xleftarrow{\cong} & \pi_0^{b\mathbb{A}^1}(X)(\kappa_A)
\end{array}$$

also commutes. Here the map $X(K) \rightarrow X(\kappa_A)$ is the composition of the inverse of the bijection $X(K) \xrightarrow{\cong} X(A)$ (the bijectivity follows from the valuative criterion of properness) and the canonical map $X(A) \rightarrow X(\kappa_A)$. Therefore, the bijection

$$\pi_0^{br}(X)(K) \cong \pi_0^{b\mathbb{A}^1}(X)(K)$$

gives an isomorphism (2.2). \square

From now on, we write $\pi_0^{b\mathbb{A}^1} = \pi_0^{br}$. We have an adjunction $\mathcal{S}hv_k \rightleftarrows \mathcal{S}hv_k^{br}$.

Lemma 2.4. *The functor $\pi_0^{b\mathbb{A}^1}$ is left adjoint to the inclusion $\mathcal{S}hv_k^{br} \hookrightarrow \mathcal{S}hv_k$.*

Proof. Let \mathcal{G} be an arbitrary birational sheaf. For every $X \in \mathcal{S}m_k^{var}$, Yoneda's lemma in $\mathcal{S}m_k$ gives a natural isomorphism

$$\mathrm{Hom}_{\mathcal{S}hv_k}(X, \mathcal{G}) \cong \mathcal{G}(X).$$

Under the identification by the equivalence in Lemma 1.4, the presheaf $\pi_0^{b\mathbb{A}^1}(X)$ is represented by X in $S_b^{-1}\mathcal{S}m_k^{var}$. Thus Yoneda's lemma in $S_b^{-1}\mathcal{S}m_k^{var}$ also gives an isomorphism

$$\mathrm{Hom}_{\mathcal{S}hv_k^{br}}(\pi_0^{b\mathbb{A}^1}(X), \mathcal{G}) \cong \mathcal{G}(X)$$

and we have

$$\mathrm{Hom}_{\mathcal{S}hv_k^{br}}(\pi_0^{b\mathbb{A}^1}(X), \mathcal{G}) \cong \mathrm{Hom}_{\mathcal{S}hv_k}(X, \mathcal{G}). \quad (2.3)$$

Let \mathcal{F} be a Nisnevich sheaf on $\mathcal{S}m_k$ which is the colimit of a direct system $\{X_\lambda\}_\lambda$ in $\mathcal{S}m_k^{var}$. Then (2.3) gives isomorphisms

$$\begin{aligned}
\mathrm{Hom}_{\mathcal{S}hv_k^{br}}(\pi_0^{b\mathbb{A}^1}(\mathcal{F}), \mathcal{G}) &\cong \mathrm{Hom}_{\mathcal{S}hv_k^{br}}(\mathrm{colim}_\lambda \pi_0^{b\mathbb{A}^1}(X_\lambda), \mathcal{G}) \\
&\cong \lim_\lambda \mathrm{Hom}_{\mathcal{S}hv_k^{br}}(\pi_0^{b\mathbb{A}^1}(X_\lambda), \mathcal{G}) \\
&\cong \lim_\lambda \mathrm{Hom}_{\mathcal{S}hv_k}(X_\lambda, \mathcal{G}) \\
&\cong \mathrm{Hom}_{\mathcal{S}hv_k}(\mathrm{colim}_\lambda X_\lambda, \mathcal{G}) \\
&\cong \mathrm{Hom}_{\mathcal{S}hv_k}(\mathcal{F}, \mathcal{G}).
\end{aligned}$$

\square

2.2 Birational \mathbb{A}^1 -connected components with transfers

Next, we define a functor $\Lambda\pi_{0,tr}^{b\mathbb{A}^1} : \mathbf{NST}_k(\Lambda) \rightarrow \mathbf{NST}_k^{br}(\Lambda)$. We denote Λ_{tr} for the Yoneda embedding

$$\mathcal{C}or_{k,\Lambda} \rightarrow \mathbf{NST}_k(\Lambda); X \mapsto \mathcal{C}or_{k,\Lambda}(-, X).$$

Definition 2.5. We define a functor $\Lambda\pi_{0,tr}^{b\mathbb{A}^1} : \mathbf{NST}_k(\Lambda) \rightarrow \mathbf{NST}_k^{br}(\Lambda)$ as the composition

$$\mathbf{NST}_k(\Lambda) \hookrightarrow \mathbf{PST}_k(\Lambda) \rightarrow \mathcal{L}in(\mathcal{C}or_{k,\Lambda}^{var}, \Lambda) \rightarrow \mathbf{NST}_k^{br}(\Lambda),$$

where the third functor is a left Kan extension of

$$\mathcal{C}or_{k,\Lambda}^{var} \rightarrow S_b^{-1}\mathcal{C}or_{k,\Lambda}^{var} \hookrightarrow \mathcal{L}in(S_b^{-1}\mathcal{C}or_{k,\Lambda}^{var}, \Lambda) \cong \mathbf{NST}_k^{br}(\Lambda).$$

We simply write $\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X) = \Lambda\pi_{0,tr}^{b\mathbb{A}^1}(\Lambda_{tr}(X))$.

We have an adjunction $\mathbf{NST}_k(\Lambda) \rightleftarrows \mathbf{NST}_k^{br}(\Lambda)$.

Lemma 2.6. *The functor $\Lambda\pi_{0,tr}^{b\mathbb{A}^1}$ is left adjoint to $\mathbf{NST}_k^{br}(\Lambda) \hookrightarrow \mathbf{NST}_k(\Lambda)$.*

Proof. Let N be an arbitrary birational sheaf with transfers of Λ -modules. For every $X \in \mathcal{C}or_{k,\Lambda}^{var}$, Yoneda's lemma in $\mathcal{C}or_{k,\Lambda}$ gives an isomorphism

$$\mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X), N) \cong N(X).$$

Under the identification by the equivalence in Lemma 1.5, the presheaf $\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X)$ is represented by X in $S_b^{-1}\mathcal{C}or_{k,\Lambda}^{var}$. Thus Yoneda's lemma in $S_b^{-1}\mathcal{C}or_{k,\Lambda}^{var}$ shows that

$$\mathrm{Hom}_{\mathbf{NST}_k(\Lambda)}(\Lambda_{tr}(X), N) \cong N(X) \cong \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X), N). \quad (2.4)$$

Let M be a Nisnevich sheaf with transfers such that

$$M \cong \mathrm{colim}_{\lambda} \Lambda_{tr}(X_{\lambda})$$

for a direct system $\{X_{\lambda}\}_{\lambda}$ in $\mathcal{C}or_{k,\Lambda}^{var}$. Then there exists a canonical isomorphism

$$\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(M) \cong \mathrm{colim}_{\lambda} \Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X_{\lambda})$$

by the definition of the functor $\Lambda\pi_{0,tr}^{b\mathbb{A}^1}$. Thus (2.4) gives isomorphisms

$$\begin{aligned} \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(M, N) &\cong \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\mathrm{colim}_{\lambda} \Lambda_{tr}(X_{\lambda}), N) \\ &\cong \lim_{\lambda} \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\Lambda_{tr}(X_{\lambda}), N) \\ &\cong \lim_{\lambda} \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X_{\lambda}), N) \\ &\cong \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\mathrm{colim}_{\lambda} \Lambda\pi_{0,tr}^{b\mathbb{A}^1}(X_{\lambda}), N) \\ &\cong \mathrm{Hom}_{\mathbf{NST}_k^{br}(\Lambda)}(\Lambda\pi_{0,tr}^{b\mathbb{A}^1}(M), N). \end{aligned}$$

□

Remark 2.7. Lemma 2.6 also gives a left adjoint of the forgetful functor $\mathbf{NST}_k^{br}(\Lambda) \rightarrow \mathcal{S}hv_k$. Indeed, a left Kan extension $\mathcal{S}hv_k \rightarrow \mathbf{NST}_k(\Lambda)$ of the composition $\mathcal{S}m_k \xrightarrow{\Gamma} \mathcal{C}or_{k,\Lambda} \xrightarrow{\Lambda_{tr}} \mathbf{NST}_k(\Lambda)$ is left adjoint to the forgetful functor $\mathbf{NST}_k(\Lambda) \rightarrow \mathcal{S}hv_k$.

3 Birational sheaves of modules

Recall that a Nisnevich sheaf of Λ -modules M on $\mathcal{S}m_k$ is called *strictly \mathbb{A}^1 -invariant*, if the map

$$H_{Nis}^i(U, M) \rightarrow H_{Nis}^i(U \times \mathbb{A}^1, M)$$

induced by the projection $U \times \mathbb{A}^1 \rightarrow U$ is an isomorphism for all $i \geq 0$ and all $U \in \mathcal{S}m_k$. Our aim of this section is to construct a birational sheaf of Λ -modules M^{br} with a monomorphism $\mu^M : M^{br} \hookrightarrow M$ which induces an isomorphism $M^{br}(X) \cong M(X)$ for all $X \in \mathcal{S}m_k^{prop}$. This morphism μ^M plays a key role in the proof of Theorem 3.

3.1 Strictly \mathbb{A}^1 -invariant sheaves and birational invariance

We denote $\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$ (resp. $\mathcal{M}od_k^{br}(\Lambda)$) for the category of strictly \mathbb{A}^1 -invariant (resp. birational) sheaves of Λ -modules. Then we have $\mathcal{M}od_k^{br}(\Lambda) \subseteq \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$. Indeed, all birational sheaves are \mathbb{A}^1 -invariant by Lemma 1.6. On the other hand, \mathbb{A}^1 -invariant birational sheaves of abelian groups are strictly \mathbb{A}^1 -invariant by [AH11, proof of Lem. 2.4]. The following lemma says that every proper birational morphism induces an isomorphism for all strictly \mathbb{A}^1 -invariant sheaves.

Lemma 3.1. *Let M be a strictly \mathbb{A}^1 -invariant sheaf of Λ -modules and $f : X \rightarrow Y$ be a proper birational morphism of smooth k -varieties. Then the induced map $M(Y) \rightarrow M(X)$ is an isomorphism.*

Proof. We write $K = k(X)$ and identify K with $k(Y)$ under the isomorphism induced by f . For each $U \in \mathcal{S}m_k$, the map $M(U) \rightarrow M(k(U))$ gives an isomorphism

$$M(U) \cong \bigcap_{x \in U^{(1)}} \text{Im}(M(\mathcal{O}_{U,x}) \rightarrow M(k(U)))$$

by [Aso12, Lem. 4.2]. Thus we only need to show that

$$\bigcap_{x \in X^{(1)}} \text{Im}(M(\mathcal{O}_{X,x}) \rightarrow M(K)) = \bigcap_{y \in Y^{(1)}} \text{Im}(M(\mathcal{O}_{Y,y}) \rightarrow M(K)).$$

The inclusion \supseteq follows from the commutative diagram

$$\begin{array}{ccc} M(Y) & \longrightarrow & M(k(Y)) \\ f^* \downarrow & & \downarrow \cong \\ M(X) & \longrightarrow & M(k(X)). \end{array}$$

We prove \subseteq . By the valuative criterion of properness, for every $y \in Y^{(1)}$ the commutative diagram

$$\begin{array}{ccc} \mathrm{Spec} K & \longrightarrow & X \\ \downarrow & & \downarrow f \\ \mathrm{Spec} \mathcal{O}_{Y,y} & \longrightarrow & Y \end{array}$$

has a lift $\phi_y : \mathrm{Spec} \mathcal{O}_{Y,y} \rightarrow X$. Then the local ring $\mathcal{O}_{X,\phi_y(y')}$ for the closed point $y' \in \mathrm{Spec} \mathcal{O}_{Y,y}$ coincides with the discrete valuation ring $\mathcal{O}_{Y,y}$ of K . In particular, $\phi_y(y')$ has codimension 1 in X . Therefore, we have

$$\begin{aligned} \bigcap_{x \in X^{(1)}} \mathrm{Im}(M(\mathcal{O}_{X,x}) \rightarrow M(K)) &\subseteq \bigcap_{y \in Y^{(1)}} \mathrm{Im}(M(\mathcal{O}_{X,\phi_y(y')}) \rightarrow M(K)) \\ &= \bigcap_{y \in Y^{(1)}} \mathrm{Im}(M(\mathcal{O}_{Y,y}) \rightarrow M(K)). \end{aligned}$$

□

Remark 3.2. More generally, Lemma 3.1 holds for unramified sheaves of Morel [Mor12] by the same proof.

3.2 Birational construction for strictly \mathbb{A}^1 -invariant sheaves

Assume k admits a resolution of singularities. We define a canonical functor $\mathrm{Mod}_k^{\mathbb{A}^1}(\Lambda) \rightarrow \mathrm{Mod}_k^{br}(\Lambda)$. Lemma 3.1 says that the restriction to $\mathcal{S}m_k^{pv}$ of a strictly \mathbb{A}^1 -invariant sheaf induces a presheaf on $S_b^{-1}\mathcal{S}m_k^{pv}$. Thus we have a functor

$$\mathrm{Mod}_k^{\mathbb{A}^1}(\Lambda) \rightarrow \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{pv}, \Lambda).$$

Moreover, Lemma 1.4 and (1.1) give equivalences of categories

$$\mathrm{Mod}_k^{br}(\Lambda) \xrightarrow{\cong} \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{var}, \Lambda) \xrightarrow{\cong} \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{pv}, \Lambda). \quad (3.1)$$

Definition 3.3. When k admits a resolution of singularities, we define a functor

$$\mathrm{Mod}_k^{\mathbb{A}^1}(\Lambda) \rightarrow \mathrm{Mod}_k^{br}(\Lambda); M \mapsto M^{br}$$

as the composition

$$\mathrm{Mod}_k^{\mathbb{A}^1}(\Lambda) \rightarrow \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{pv}, \Lambda) \xrightarrow{\cong} \mathrm{Mod}_k^{br}(\Lambda).$$

We construct a natural morphism of sheaves $\mu^M : M^{br} \rightarrow M$ for each $M \in \mathrm{Mod}_k^{\mathbb{A}^1}(\Lambda)$. By the construction, we have $M^{br}(X) = M(X)$ for all $X \in \mathcal{S}m_k^{pv}$. For every $U \in \mathcal{S}m_k^{var}$, the map $M(U) \rightarrow M(k(U))$ induces a natural isomorphism

$$M(U) \cong \bigcap_{x \in U^{(1)}} \mathrm{Im}(M(\mathcal{O}_{U,x}) \hookrightarrow M(k(U))) \quad (3.2)$$

by [Aso12, Lem. 4.2]. Thus [CT95, Prop. 2.1.8] shows that

$$\begin{aligned} M^{br}(X) &\cong \bigcap_{x \in X^{(1)}} \text{Im}(M(\mathcal{O}_{X,x}) \hookrightarrow M(k(X))) \\ &= \bigcap_{A \in \mathcal{P}(k(X)/k)} \text{Im}(M(A) \hookrightarrow M(k(X))). \end{aligned}$$

For a smooth proper compactification \bar{U} of $U \in \mathcal{S}m_k^{var}$, we have

$$M^{br}(U) \cong M^{br}(\bar{U}) \cong \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Im}(M(A) \hookrightarrow M(k(U))). \quad (3.3)$$

Thus the sheaf M^{br} can be regarded as the subsheaf

$$U \mapsto \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Im}(M(A) \hookrightarrow M(k(U)))$$

of M under the isomorphism (3.2). Then μ^M is defined as the embedding $M^{br} \hookrightarrow M$. We prove some basic properties of the functor $(-)^{br}$ and the morphism μ^M .

Proposition 3.4. *Assume k admits a resolution of singularities. Let M be a strictly \mathbb{A}^1 -invariant sheaf.*

- (1) *The morphism $\mu^M : M^{br} \rightarrow M$ is injective.*
- (2) *The induced map $\mu_X^M : M^{br}(X) \rightarrow M(X)$ is an isomorphism for every $X \in \mathcal{S}m_k^{prop}$.*
- (3) *The induced map*

$$\text{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(N, M^{br}) \xrightarrow{(\mu^M)_*} \text{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}}(N, M)$$

is an isomorphism for every $N \in \mathcal{M}od_k^{br}(\Lambda)$. In particular, the functor $(-)^{br}$ is right adjoint to the inclusion $\mathcal{M}od_k^{br}(\Lambda) \hookrightarrow \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$.

Proof. The assertions (1) and (2) follow from the construction of M^{br} and μ^M . We prove (3). Since μ^M is injective by (1), so is $(\mu^M)_*$. On the other hand, for a morphism $f : N \rightarrow M$ in $\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$, we have a morphism $f' : N \rightarrow M^{br}$ defined by

$$\begin{aligned} N(U) &\cong \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Im}(N(A) \hookrightarrow N(k(U))) \\ &\rightarrow \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Im}(M(A) \hookrightarrow M(k(U))) \\ &\cong M^{br}(U) \end{aligned}$$

for each $U \in \mathcal{S}m_k^{var}$ by using the isomorphism in (3.3). Then $\mu^M \circ f' = f$ and thus $(\mu^M)_*$ is surjective. \square

3.3 Birational construction for \mathbb{A}^1 -invariant sheaves with transfers

Assume k perfect. Let $\mathbf{HI}_k(\Lambda)$ be the category of \mathbb{A}^1 -invariant sheaves with transfers of Λ -modules. Note that every object in $\mathbf{HI}_k(\Lambda)$ is strictly \mathbb{A}^1 -invariant as a sheaf on $\mathcal{S}m_k$ (see [MVW, Thm. 13.8]). Lemma 1.6 shows that $\mathbf{NST}_k^{br}(\Lambda)$ is a full subcategory of $\mathbf{HI}_k(\Lambda)$. In [KS17, §6], Kahn-Sujatha construct a right adjoint functor

$$\mathbf{NST}_k^{br}(\Lambda) \rightarrow \mathbf{HI}_k(\Lambda); M \mapsto M_{nr}$$

of the inclusion $\mathbf{NST}_k^{br}(\Lambda) \hookrightarrow \mathbf{HI}_k(\Lambda)$ which sends $M \in \mathbf{HI}_k(\Lambda)$ to the presheaf with transfers defined by

$$U \mapsto \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Ker} \left(M(k(U)) \xrightarrow{\partial_A} M_{-1}(\kappa_A) \right).$$

Here M_{-1} is the contraction of M and ∂_A is the residue map associated with A (see definitions [MVW, Lec. 23 and Ex. 24.6]). On the other hand, the sequence

$$0 \rightarrow M(A) \rightarrow M(k(U)) \xrightarrow{\partial_A} M_{-1}(\kappa_A) \rightarrow 0$$

is exact by [MVW, Ex. 24.6]. Thus we have

$$M_{nr}(U) = \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Im}(M(A) \rightarrow M(k(U))). \quad (3.4)$$

Then [KS17, Lem. 6.2.3, 6.2.4 and 6.2.6] show that M_{nr} can be regarded as the subsheaf

$$U \mapsto \bigcap_{A \in \mathcal{P}(k(U)/k)} \text{Im}(M(A) \rightarrow M(k(U)))$$

of M under the isomorphism (3.2). Thus when k admits a resolution of singularities, there exists a canonical isomorphism $\Gamma_*(M_{nr}) \cong (\Gamma_*M)^{br}$ in $\mathcal{M}od_k(\Lambda)$, *i.e.*, the diagram

$$\begin{array}{ccc} \mathbf{HI}_k(\Lambda) & \xrightarrow{(-)_{nr}} & \mathbf{NST}_k^{br}(\Lambda) \\ \Gamma_* \downarrow & & \downarrow \Gamma_* \\ \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda) & \xrightarrow{(-)^{br}} & \mathcal{M}od_k^{br}(\Lambda) \end{array}$$

is 2-commutative, by the construction of M^{br} . Moreover, if we write ν^M for the embedding $M_{nr} \hookrightarrow M$, the diagram

$$\begin{array}{ccc} \Gamma_*(M_{nr}) & \xrightarrow{\Gamma_*(\nu^M)} & \Gamma_*M \\ \cong \downarrow & & \parallel \\ (\Gamma_*M)_{nr} & \xrightarrow{\mu^M} & \Gamma_*M \end{array}$$

commutes. We obtain an analogue of Proposition 3.4 for the functor $(-)_{nr}$ without assuming a resolution of singularities.

Proposition 3.5. *Assume k perfect. Let M be an \mathbb{A}^1 -invariant sheaf with transfers.*

- (1) *The map $\nu^M : M_{nr} \rightarrow M$ is injective.*
- (2) *The induced map $\nu_X^M : M_{nr}(X) \rightarrow M(X)$ is an isomorphism.*

Proof. The assertion (1) follows from the construction of ν^M . By (3.4) and [CT95, Prop. 2.1.8], we have

$$\begin{aligned} M_{nr}(X) &\cong \bigcap_{A \in \mathcal{P}(k(X)/k)} \text{Im}(M(A) \rightarrow M(k(X))) \\ &= \bigcap_{x \in X^{(1)}} \text{Im}(M(\mathcal{O}_{X,x}) \hookrightarrow M(k(X))). \end{aligned}$$

Thus (2) follows from (3.2). \square

4 Structure of zeroth \mathbb{A}^1 - and Suslin homology

4.1 Structure theorem of zeroth \mathbb{A}^1 -homology

Our aim of this subsection is to prove a structure theorem of the \mathbb{A}^1 -homology of smooth proper varieties. In [Mor05], Morel defined the \mathbb{A}^1 -homology sheaf $\mathbf{H}_i^{\mathbb{A}^1}(\mathcal{F})$ for $\mathcal{F} \in \mathcal{S}hv_k$ as a strictly \mathbb{A}^1 -invariant sheaf. Especially, the zeroth \mathbb{A}^1 -homology functor

$$\mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda) : \mathcal{S}hv_k \rightarrow \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$$

can be characterized as a left adjoint of the forgetful functor $\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda) \rightarrow \mathcal{S}hv_k$ (see [Aso12, Lem. 3.3]). For $\mathcal{F} \in \mathcal{S}hv_k$, we denote $\Lambda_{pre}(\mathcal{F})$ for the presheaf of free Λ -modules generated by \mathcal{F} . Note that if \mathcal{F} is birational, then so is $\Lambda_{pre}(\mathcal{F})$. Thus we have the functor

$$\Lambda_{pre} : \mathcal{S}hv_k^{br} \rightarrow \mathcal{M}od_k^{br}(\Lambda).$$

This is left adjoint to the forgetful functor $\mathcal{M}od_k^{br}(\Lambda) \rightarrow \mathcal{S}hv_k^{br}$ by the definition. Our structure theorem of zeroth \mathbb{A}^1 -homology is stated as follows.

Theorem 4.1. *Assume k admits a resolution of singularities. For every $X \in \mathcal{S}m_k$, there exists a natural epimorphism of sheaves*

$$\Lambda_{pre}(\pi_0^{b\mathbb{A}^1}(X)) \twoheadrightarrow \mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda).$$

Moreover, this is an isomorphism if X is proper.

Proof. By Lemmas 2.4 and 3.4, we obtain three adjunctions

$$\mathcal{S}hv_k \rightleftarrows \mathcal{S}hv_k^{br} \rightleftarrows \mathcal{M}od_k^{br}(\Lambda) \rightleftarrows \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda).$$

Hence the composite functor

$$\mathcal{S}h v_k \xrightarrow{\pi_0^{b\mathbb{A}^1}} \mathcal{S}h v_k^{br} \xrightarrow{\Lambda_{pre}} \mathcal{M}od_k^{br}(\Lambda) \hookrightarrow \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$$

is left adjoint to the composition

$$\mathcal{S}h v_k \hookrightarrow \mathcal{S}h v_k^{br} \leftarrow \mathcal{M}od_k^{br}(\Lambda) \xleftarrow{(-)^{br}} \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda).$$

By using this adjunction and Yoneda's lemma in $\mathcal{S}m_k$, we have a natural isomorphism

$$\mathrm{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(\Lambda_{pre}(\pi_0^{b\mathbb{A}^1}(X)), M) \cong \mathrm{Hom}_{\mathcal{S}h v_k}(X, M^{br}) \cong M^{br}(X) \quad (4.1)$$

for all $X \in \mathcal{S}m_k$ and all $M \in \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$. On the other hand, the zeroth \mathbb{A}^1 -homology functor

$$\mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda) : \mathcal{S}h v_k \rightarrow \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$$

is left adjoint to the forgetful functor $\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda) \rightarrow \mathcal{S}h v_k$. Therefore, we also obtain a natural isomorphism

$$\mathrm{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda), M) \cong \mathrm{Hom}_{\mathcal{S}h v_k}(X, M) \cong M(X). \quad (4.2)$$

By (4.1) and (4.2), the natural monomorphism $\mu^M : M^{br} \rightarrow M$ induces a injection

$$\mathrm{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(\Lambda_{pre}(\pi_0^{b\mathbb{A}^1}(X)), M) \rightarrow \mathrm{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda), M).$$

This map is an isomorphism when X is proper by Proposition 3.4. Thus Yoneda's lemma in $\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$ leads an epimorphism of sheaves

$$\Lambda_{pre}(\pi_0^{b\mathbb{A}^1}(X)) \twoheadrightarrow \mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)$$

which is an isomorphism for proper X . □

Corollary 4.2. *Assume k admits a resolution of singularities. For $X \in \mathcal{S}m_k^{prop}$ and $U \in \mathcal{S}m_k^{var}$, the section $\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(U)$ is the free Λ -module generated by $X(k(U))/R$.*

Proof. This follows from Theorem 4.1 and Lemma 2.2. □

We obtain the Künneth formula and the universal coefficient theorem of \mathbb{A}^1 -homology in degree zero.

Corollary 4.3. *Assume k admits a resolution of singularities. For $X, Y \in \mathcal{S}m_k^{prop}$, there exists a natural isomorphism*

$$\mathbf{H}_0^{\mathbb{A}^1}(X \times Y; \Lambda) \cong \mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda) \otimes_{\Lambda} \mathbf{H}_0^{\mathbb{A}^1}(Y; \Lambda),$$

where \otimes_{Λ} means the tensor product of presheaves over Λ .

Proof. This follows from Theorem 4.1 and the natural bijection

$$(X \times Y)(K)/R \cong (X(K)/R) \times (Y(K)/R)$$

for each $K \in \mathcal{F}_k$. □

Corollary 4.4. *Assume k admits a resolution of singularities. For $X \in \mathcal{S}m_k^{prop}$, there exists a natural isomorphism*

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda) \cong \mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z}) \otimes \Lambda,$$

where \otimes means the tensor product of presheaves over \mathbb{Z} .

Proof. This follows from $\Lambda_{pre}(-) \cong \mathbb{Z}_{pre}(-) \otimes \Lambda$ and Theorem 4.1. □

Remark 4.5. Assume k admits a resolution of singularities. By Theorem 4.1, there exists a pairing

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(Y) \times \mathbf{H}_0^{\mathbb{A}^1}(Y; \Lambda)(Z) \rightarrow \mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(Z)$$

defined by the map

$$\mathrm{Hom}_{S_b^{-1}\mathcal{S}m_k}(Y, X) \times \mathrm{Hom}_{S_b^{-1}\mathcal{S}m_k}(Z, Y) \rightarrow \mathrm{Hom}_{S_b^{-1}\mathcal{S}m_k}(Z, X); (f, g) \mapsto f \circ g$$

for all $X, Y, Z \in \mathcal{S}m_k^{prop}$. In particular, this pairing gives a ring structure for $\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(X)$.

4.2 Rational points and finiteness

In this subsection, we give some examples where $\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(k)$ is finitely generated. We first introduce the following notation.

Definition 4.6. We write $b_0^{\mathbb{A}^1}(X) = \dim_{\mathbb{Q}} \mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Q})(k)$ for $X \in \mathcal{S}m_k$. This is called *the zeroth \mathbb{A}^1 -Betti number of X* .

By Corollary 4.2, $b_0^{\mathbb{A}^1}(X)$ coincides with the rank of the free Λ -module $\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(k)$ if X is proper. Our structure theorem of zeroth \mathbb{A}^1 -homology enables to relate the \mathbb{A}^1 -Betti number to k -rational points modulo R -equivalence.

Theorem 4.7. *Assume k admits a resolution of singularities. For a smooth proper k -variety X , we have*

$$b_0^{\mathbb{A}^1}(X) = \#(X(k)/R).$$

In particular, $b_0^{\mathbb{A}^1}(X) = 0$ if and only if $X(k) = \emptyset$.

Proof. This follows from Corollary 4.2. □

The following proposition implies the finiteness of $b_0^{\mathbb{A}^1}(X)$ for a rationally connected smooth proper real variety X .

Proposition 4.8. *Let X be a rationally connected smooth proper variety over \mathbb{R} . Then we have*

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)(\mathbb{R}) \cong H_0(X(\mathbb{R}); \Lambda).$$

In particular, $b_0^{\mathbb{A}^1}(X)$ coincides with the ordinary zeroth Betti number of the real manifold $X(\mathbb{R})$.

Proof. By [Kol99, Cor. 1.7], the set of R -equivalence classes $X(\mathbb{R})/R$ coincides with the connected components of $X(\mathbb{R})$. Thus the assertion follows from Corollary 4.2. \square

By using Theorem 4.7, we also obtain some examples where $b_0^{\mathbb{A}^1}(X)$ is finite.

- Example 4.9.** (1) Let X be a smooth proper variety with finitely many k -rational points. Then $b_0^{\mathbb{A}^1}(X)$ is finite. Thus for example, $b_0^{\mathbb{A}^1}(C)$ is finite for a smooth projective curve C of genus ≥ 2 over a number field by Mordell conjecture proved by Faltings [Fal83].
- (2) Let X be a rationally connected smooth proper variety over either of \mathbb{C} or a p -adic field. Then $b_0^{\mathbb{A}^1}(X)$ is also finite by [Kol99, Cor. 1.5].

Remark 4.10. For X a smooth k -variety, $\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z})(k)$ can be expressed in terms of triangulated categories as follows.

- (1) There exist natural isomorphisms

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z})(k) \cong \mathrm{Hom}_{\mathcal{M}od_k(\mathbb{Z})}(\mathbb{Z}, \mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z})) \cong \mathrm{Hom}_{D_{\mathbb{A}^1}(k)}(\mathbb{Z}, \mathbb{Z}(X)).$$

Here $D_{\mathbb{A}^1}(k)$ is the \mathbb{A}^1 -derived category defined by Morel [Mor05] and $\mathbb{Z}(X)$ is the Nisnevich sheafification of $\mathbb{Z}_{pre}(X)$. This follows from [Aso12, Lem. 3.3] and its proof.

- (2) There also exist natural isomorphisms

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z})(k) \cong \pi_0^S(\Sigma_{S^1}^{\infty}(X_+))(k) \cong \mathrm{Hom}_{\mathcal{SH}_{S^1}(k)}(S^0, \Sigma_{S^1}^{\infty}(X_+)).$$

Here $\Sigma_{S^1}^{\infty}(X_+)$ is the infinity S^1 -suspension of $X_+ = X \sqcup \mathrm{Spec} k$, $\pi_0^S(-)$ is the S^1 -stable zeroth \mathbb{A}^1 -homotopy sheaf, $\mathcal{SH}_{S^1}(k)$ is the \mathbb{A}^1 -homotopy category of S^1 -spectra, and S^0 is the sphere spectrum (see definitions [Mor05]). The first isomorphism is given by [AH11, Prop. 2.1] and the second isomorphism is given by the definition of $\pi_0^S(-)$.

4.3 Zeroth Suslin homology and zero cycles

In this subsection, we prove the Suslin homology version of Theorem 4.1. For $X \in \mathcal{S}m_k$, the Suslin homology sheaf $\mathbf{H}_i^S(X; \Lambda)$ is defined as the i -th homology sheaf with transfers of the motive of X with Λ -coefficients in the sense of Voevodsky [Voe00]. Then the section $\mathbf{H}_i^S(X; \Lambda)(k)$ coincides with the singular homology of Suslin-Voevodsky [SV96], called Suslin homology group. Note that

$\mathbf{H}_i^S(X; \Lambda)$ is \mathbb{A}^1 -invariant by the \mathbb{A}^1 -invariance of the motive of X . Thus we have a functor

$$\mathbf{H}_0^S(-; \Lambda) : \mathcal{C}or_{k, \Lambda} \rightarrow \mathbf{H}\mathbf{I}_k(\Lambda).$$

As a counterpart of Theorem 4.1, we construct an isomorphism of sheaves with transfers $\Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X) \cong \mathbf{H}_0^S(X; \Lambda)$ for $X \in \mathcal{S}m_k^{prop}$.

Theorem 4.11. *Assume k perfect. For every $X \in \mathcal{S}m_k$, there exists a natural epimorphism of sheaves with transfers*

$$\Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X) \twoheadrightarrow \mathbf{H}_0^S(X; \Lambda).$$

Moreover, this is an isomorphism if X is proper.

Proof. By Lemma 2.6, the composite functor

$$\mathbf{NST}_k(\Lambda) \xrightarrow{\Lambda\pi_{0, tr}^{b\mathbb{A}^1}} \mathbf{NST}_k^{br}(\Lambda) \hookrightarrow \mathbf{H}\mathbf{I}_k(\Lambda)$$

is left adjoint to the functor $(-)_nr : \mathbf{H}\mathbf{I}_k(\Lambda) \rightarrow \mathbf{NST}_k(\Lambda)$. For all $X \in \mathcal{S}m_k$ and all $M \in \mathbf{H}\mathbf{I}_k(\Lambda)$, thus we obtain natural isomorphisms

$$\mathrm{Hom}_{\mathbf{H}\mathbf{I}_k(\Lambda)}(\Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X), M) \cong \mathrm{Hom}_{\mathbf{NST}_k(\Lambda)}(\Lambda_{tr}(X), M_{nr}) \cong M_{nr}(X)$$

by Yoneda's lemma in $\mathcal{C}or_{k, \Lambda}$. On the other hand, [Aso12, Lem. 3.3] gives an isomorphism

$$\mathrm{Hom}_{\mathbf{H}\mathbf{I}_k(\Lambda)}(\mathbf{H}_0^S(X; \Lambda), M) \cong M(X).$$

Thus the map

$$\mathrm{Hom}_{\mathbf{H}\mathbf{I}_k(\Lambda)}(\Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X), M_{nr}) \rightarrow \mathrm{Hom}_{\mathbf{H}\mathbf{I}_k(\Lambda)}(\mathbf{H}_0^S(X; \Lambda), M)$$

obtained by $\nu^M : M_{nr} \rightarrow M$ introduces an epimorphism of sheaves with transfers

$$\Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X) \twoheadrightarrow \mathbf{H}_0^S(X; \Lambda)$$

by Yoneda's lemma in $\mathbf{H}\mathbf{I}_k(\Lambda)$. By Proposition 3.5, this morphism is an isomorphism if X is proper. \square

Assume k perfect. For every $X \in \mathcal{S}m_k^{prop}$, there exists a canonical isomorphism

$$\mathbf{H}_0^S(X; \mathbb{Z})(K) \cong \mathrm{CH}_0(X_K)$$

for all $K \in \mathcal{F}_k$ (see [Aso12, Ex. 4.9]). Thus we obtain an application to Chow groups of zero cycles. We simply write

$$S_b^{-1}\mathcal{C}or_k(X, Y) = \mathrm{Hom}_{S_b^{-1}\mathcal{C}or_{k, \mathbb{Z}}}(X, Y)$$

for $X, Y \in \mathcal{S}m_k$.

Corollary 4.12. *Assume k perfect. Let X be a smooth proper k -variety and U be a smooth k -variety. Then we have*

$$S_b^{-1}\mathcal{C}or_k(U, X) \cong \mathrm{CH}_0(X_{k(U)}).$$

Proof. By Theorem 4.11 and the construction of $\Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X)$, we have

$$S_b^{-1}\mathcal{C}or_k(U, X) \cong \mathbb{Z}\pi_{0, tr}^{b\mathbb{A}^1}(X)(U) \cong \mathbf{H}_0^S(X; \mathbb{Z})(U) \cong \mathbf{H}_0^S(X; \mathbb{Z})(k(U)).$$

On the other hand, we also have $\mathbf{H}_0^S(X; \mathbb{Z})(k(U)) \cong \mathrm{CH}_0(X_{k(U)})$ by [Aso12, Ex. 4.9]. \square

Remark 4.13. Assume k perfect. Like Remark 4.5, the composition of morphisms in $S_b^{-1}\mathcal{C}or_{k, \Lambda}$ also induces a pairing

$$\mathbf{H}_0^S(X; \Lambda)(Y) \times \mathbf{H}_0^S(Y; \Lambda)(Z) \rightarrow \mathbf{H}_0^S(X; \Lambda)(Z)$$

and a ring structure of $\mathbf{H}_0^S(X; \Lambda)(X)$ for all $X, Y, Z \in \mathcal{S}m_k^{pv}$. Thus we obtain a pairing

$$\mathrm{CH}_0(X_{k(Y)}) \times \mathrm{CH}_0(Y_{k(Z)}) \rightarrow \mathrm{CH}_0(X_{k(Z)})$$

and a ring structure of $\mathrm{CH}_0(X_{k(X)})$.

5 Universal birational invariance

5.1 Universal birational invariance of \mathbb{A}^1 -homology

Our aim of this subsection is to prove the main theorem of this paper, *i.e.*, the universal birational invariance of zeroth \mathbb{A}^1 -homology. For a category \mathcal{C} , we denote $\Lambda(\mathcal{C})$ for the category whose objects are same with \mathcal{C} and that $\mathrm{Hom}_{\Lambda(\mathcal{C})}(A, B)$ for each $A, B \in \Lambda(\mathcal{C})$ is the free Λ -module generated by $\mathrm{Hom}_{\mathcal{C}}(A, B)$. Note that every functor from \mathcal{C} to an Λ -enriched category uniquely factors through the canonical Λ -linear functor $\mathcal{C} \rightarrow \Lambda(\mathcal{C})$. Thus we have a functor

$$\mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda) : \Lambda(S_b^{-1}\mathcal{S}m_k^{prop}) \rightarrow \mathrm{Mod}_k(\Lambda). \quad (5.1)$$

Proposition 5.1. *Assume k admits a resolution of singularities. Then the functor (5.1) is fully faithful.*

Proof. By [KS07, Thm. 6.4], we only need to show that the functor

$$\mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda) : \Lambda(S_b^{-1}\mathcal{S}m_k^{pv}) \rightarrow \mathrm{Mod}_k(\Lambda) \quad (5.2)$$

is fully faithful. The canonical functor $S_b^{-1}\mathcal{S}m_k^{pv} \rightarrow \Lambda(S_b^{-1}\mathcal{S}m_k^{pv})$ induces an equivalence of categories

$$\mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{pv}, \Lambda) \xrightarrow{\cong} \mathcal{L}in(\Lambda(S_b^{-1}\mathcal{S}m_k^{pv}), \Lambda).$$

On the other hand, we obtain equivalences

$$\mathcal{M}od_k^{br}(\Lambda) \xrightarrow{\cong} \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{var}, \Lambda) \xrightarrow{\cong} \mathcal{P}resh(S_b^{-1}\mathcal{S}m_k^{pv}, \Lambda)$$

by (3.1). Thus we have a functor

$$\mathcal{H} : \Lambda(S_b^{-1}\mathcal{S}m_k^{pv}) \hookrightarrow \mathcal{L}in(\Lambda(S_b^{-1}\mathcal{S}m_k^{pv}), \Lambda) \cong \mathcal{M}od_k^{br}(\Lambda) \hookrightarrow \mathcal{M}od_k(\Lambda),$$

where the first functor is the Yoneda embedding. Therefore, \mathcal{H} is fully faithful by Yoneda's lemma in $\Lambda(S_b^{-1}\mathcal{S}m_k^{pv})$. By Theorem 4.1, there exists a natural isomorphism

$$\mathcal{H}(X) \cong \text{Hom}_{\Lambda(S_b^{-1}\mathcal{S}m_k^{pv})}(-, X) \cong \Lambda_{pre}(\pi_0^{b\mathbb{A}^1}(X)) \cong \mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)$$

for all $X \in \mathcal{S}m_k^{pv}$. Thus the diagram

$$\begin{array}{ccc} \mathcal{S}m_k^{pv} & \xrightarrow{\mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda)} & \mathcal{M}od_k(\Lambda) \\ \downarrow & & \parallel \\ \Lambda(S_b^{-1}\mathcal{S}m_k^{pv}) & \xrightarrow{\mathcal{H}} & \mathcal{M}od_k(\Lambda) \end{array}$$

is 2-commutative. This diagram shows that the functor (5.2) is naturally equivalent to \mathcal{H} . Since \mathcal{H} is fully faithful, so is (5.2). \square

Proposition 5.1 shows that the zeroth \mathbb{A}^1 -homology functor on $S_b^{-1}\mathcal{S}m_k^{prop}$ is conservative and faithful.

Corollary 5.2. *Assume k admits a resolution of singularities. Then the functor*

$$\mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda) : S_b^{-1}\mathcal{S}m_k^{prop} \rightarrow \mathcal{M}od_k(\Lambda). \quad (5.3)$$

is conservative and faithful.

Proof. Since the functors $S_b^{-1}\mathcal{S}m_k^{prop} \rightarrow \Lambda(S_b^{-1}\mathcal{S}m_k^{prop})$ and (5.1) are conservative and faithful by Proposition 5.1, so is (5.3). \square

For describing universal birational invariance, we introduce the following term.

Definition 5.3. Let \mathcal{C} be a full subcategory of $\mathcal{S}m_k^{prop}$. Then \mathcal{C} is called a *birationally fully faithful subcategory* of $\mathcal{S}m_k^{prop}$, if the functor $S_b^{-1}\mathcal{C} \rightarrow S_b^{-1}\mathcal{S}m_k^{prop}$ is fully faithful.

We see some basic examples of birationally fully faithful subcategories of $\mathcal{S}m_k^{prop}$.

Example 5.4. Clearly, $\mathcal{S}m_k^{prop}$ is a birationally fully faithful subcategory of itself. Let $\mathcal{S}m_k^{pjv}$ (resp. $\mathcal{S}m_k^{proj}$) be the full subcategory of $\mathcal{S}m_k^{prop}$ spanned by smooth projective k -varieties (resp. k -schemes). When k admits a resolution of singularities, $\mathcal{S}m_k^{proj}$ is a birationally fully faithful subcategory of $\mathcal{S}m_k^{prop}$ by the equivalence $\mathcal{S}m_k^{proj} \cong \mathcal{S}m_k^{prop}$ in [KS07, Thm. 8.8]. Moreover, $\mathcal{S}m_k^{pjv}$ and $\mathcal{S}m_k^{proj}$ are also birationally fully faithful subcategories of $\mathcal{S}m_k^{prop}$ by [KS07, Thm. 6.4].

We prove the universal birational invariance of the \mathbb{A}^1 -homology functor on birationally fully faithful subcategories of $\mathcal{S}m_k^{prop}$. For a subcategory $\mathcal{C} \subseteq \mathcal{S}m_k^{prop}$, we denote $\mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1}$ for the full subcategory of $\mathcal{M}od_k(\Lambda)$ spanned by sheaves isomorphic to $\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda)$ for some $X \in \mathcal{C}$.

Theorem 5.5. *Assume k admits a resolution of singularities. Let \mathcal{C} be a birationally full subcategory of $\mathcal{S}m_k^{prop}$ and \mathcal{A} be an arbitrary category enriched by Λ -modules.*

- (1) *Let $F : \mathcal{C} \rightarrow \mathcal{A}$ be an arbitrary functor which sends each birational morphism to an isomorphism. Then there exists one and only one (up to a natural equivalence) Λ -linear functor $F_{S_b} : \mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1} \rightarrow \mathcal{A}$ such that the diagram*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{A} \\ \downarrow & \nearrow F_{S_b} & \\ \mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1} & & \end{array}$$

is 2-commutative.

- (2) *Let $F' : \mathcal{C}^{op} \rightarrow \mathcal{A}$ be an arbitrary functor which sends each birational morphism to an isomorphism. Then there exists one and only one (up to a natural equivalence) Λ -linear functor $F'_{S_b} : (\mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1})^{op} \rightarrow \mathcal{A}$ such that the diagram*

$$\begin{array}{ccc} \mathcal{C}^{op} & \xrightarrow{F'} & \mathcal{A} \\ \downarrow & \nearrow F'_{S_b} & \\ (\mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1})^{op} & & \end{array}$$

is 2-commutative.

Proof. Since $S_b^{-1}\mathcal{C} \rightarrow S_b^{-1}\mathcal{S}m_k^{prop}$ is fully faithful, so is the composition

$$\Lambda(S_b^{-1}\mathcal{C}) \rightarrow \Lambda(S_b^{-1}\mathcal{S}m_k^{prop}) \rightarrow \mathcal{M}od_k(\Lambda) \quad (5.4)$$

by Proposition 5.1. On the other hand, the essential image of the functor (5.4) coincides with $\mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1}$. Thus we have an equivalence $\Lambda(S_b^{-1}\mathcal{C}) \cong \mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1}$

such that the diagram

$$\begin{array}{ccc}
\mathcal{C} & \xlongequal{\quad} & \mathcal{C} \\
\downarrow & & \downarrow \mathbf{H}_0^{\mathbb{A}^1}(-; \Lambda) \\
\Lambda(S_b^{-1}\mathcal{C}) & \xrightarrow{\cong} & \mathbf{Im}_{\Lambda}^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1}
\end{array} \tag{5.5}$$

is 2-commutative. Thus (1) follows from the universality on $\mathcal{C} \rightarrow \Lambda(S_b^{-1}\mathcal{C})$. Similarly, we obtain (2) by taking the opposite categories in (5.5). \square

Remark 5.6. The functor F_{S_b} (resp. F'_{S_b}) in Theorem 5.5 is canonically extended to $\mathcal{M}od_k^{br}(\Lambda) \rightarrow \mathcal{A}$ (resp. $(\mathcal{M}od_k^{br}(\Lambda))^{op} \rightarrow \mathcal{A}$), if $\mathcal{S}m_k^{pjv} \subseteq \mathcal{C}$ and \mathcal{A} is cocomplete. Indeed, the equivalences (3.1) and $\mathcal{S}m_k^{pjv} \cong \mathcal{S}m_k^{var}$ in [KS07, Prop. 8.5] induce

$$\mathcal{M}od_k^{br}(\Lambda) \cong \mathcal{L}in(\Lambda(S_b^{-1}\mathcal{S}m_k^{var}), \Lambda) \cong \mathcal{L}in(\Lambda(S_b^{-1}\mathcal{S}m_k^{pjv}), \Lambda).$$

Under the identification by this equivalence, the category $\mathbf{Im}_{\Lambda}^{\mathcal{C}} \mathbf{H}_0^{\mathbb{A}^1}$ contains the full subcategory of $\mathcal{M}od_k^{br}(\Lambda)$ consisting of representable presheaves on $\Lambda(S_b^{-1}\mathcal{S}m_k^{pjv})$ by Theorem 4.1. Thus the purpose extension is obtained as a left Kan extension of $\Lambda(S_b^{-1}\mathcal{S}m_k^{pjv}) \rightarrow \mathcal{A}$ (resp. $(\Lambda(S_b^{-1}\mathcal{S}m_k^{pjv}))^{op} \rightarrow \mathcal{A}$).

5.2 Universal birational invariance of Suslin homology

In this subsection we prove the Suslin homology version of the universal birational invariance property. We have a functor

$$\mathbf{H}_0^S(-; \Lambda) : S_b^{-1}\mathcal{C}or_{k, \Lambda}^{prop} \rightarrow \mathbf{NST}_k(\Lambda). \tag{5.6}$$

Proposition 5.7. *Assume k perfect and admits a resolution of singularities. Then the functor (5.6) is fully faithful.*

Proof. By [KS07, Thm. 6.4], we only need to show that the functor

$$\mathbf{H}_0^S(-; \Lambda) : S_b^{-1}\mathcal{C}or_{k, \Lambda}^{pv} \rightarrow \mathbf{NST}_k(\Lambda) \tag{5.7}$$

is fully faithful. By Lemma 1.5 and Proposition 1.2, we obtain equivalences of categories

$$\mathbf{NST}_k^{br}(\Lambda) \xleftarrow{\cong} \mathcal{L}in(S_b^{-1}\mathcal{C}or_{k, \Lambda}^{var}, \Lambda) \xrightarrow{\cong} \mathcal{L}in(S_b^{-1}\mathcal{C}or_{k, \Lambda}^{pv}, \Lambda).$$

Thus we have a functor

$$\mathcal{H}_{tr} : S_b^{-1}\mathcal{C}or_{k, \Lambda}^{pv} \hookrightarrow \mathcal{L}in(S_b^{-1}\mathcal{C}or_{k, \Lambda}^{pv}, \Lambda) \cong \mathbf{NST}_k^{br}(\Lambda) \hookrightarrow \mathbf{NST}_k(\Lambda)$$

where the first functor is the Yoneda embedding. Hence \mathcal{H}_{tr} is fully faithful by Yoneda's lemma in $S_b^{-1}\mathcal{C}or_{k, \Lambda}^{pv}$. By Theorem 4.11, there exists a natural isomorphism

$$\mathcal{H}_{tr}(X) \cong S_b^{-1}\mathcal{C}or_{k, \Lambda}(-, X) \cong \Lambda\pi_{0, tr}^{b\mathbb{A}^1}(X) \cong \mathbf{H}_0^S(X; \Lambda)$$

for all $X \in \mathcal{C}or_{k,\Lambda}^{pv}$. Thus the diagram

$$\begin{array}{ccc} \mathcal{C}or_{k,\Lambda}^{pv} & \xrightarrow{\mathbf{H}_0^S(-;\Lambda)} & \mathbf{NST}_k(\Lambda) \\ \downarrow & & \parallel \\ S_b^{-1}\mathcal{C}or_{k,\Lambda}^{pv} & \xrightarrow{\mathcal{H}_{tr}} & \mathbf{NST}_k(\Lambda) \end{array}$$

is 2-commutative. Therefore, since \mathcal{H}_{tr} is fully faithful, so is the functor (5.7). \square

We also introduce birationally fully faithful subcategories of $\mathcal{C}or_{k,\Lambda}^{prop}$.

Definition 5.8. Let \mathcal{C} be a full subcategory of $\mathcal{C}or_{k,\Lambda}^{prop}$. Then \mathcal{C} is called a birationally fully faithful subcategory of $\mathcal{C}or_{k,\Lambda}^{prop}$, if the functor $S_b^{-1}\mathcal{C} \rightarrow S_b^{-1}\mathcal{C}or_{k,\Lambda}^{prop}$ is fully faithful.

Example 5.9. Clearly, $\mathcal{C}or_{k,\Lambda}^{prop}$ is a birationally fully faithful subcategory of itself. By [KS07, Thm. 6.4], the subcategory $\mathcal{C}or_{k,\Lambda}^{pv} \subseteq \mathcal{C}or_{k,\Lambda}^{prop}$ is also birationally fully faithful.

For a subcategory $\mathcal{C} \subseteq \mathcal{C}or_{k,\Lambda}^{prop}$, we denote $\mathbf{Im}_{\Lambda}^{\mathcal{C}}\mathbf{H}_0^S$ for the full subcategory of $\mathbf{NST}_k(\Lambda)$ spanned by sheaves with transfers isomorphic to $\mathbf{H}_0^S(X;\Lambda)$ for some $X \in \mathcal{C}$. We prove universal birational invariance of the zeroth Suslin homology.

Theorem 5.10. Assume k perfect and admits a resolution of singularities. Let \mathcal{C} be a birationally fully faithful subcategory of $\mathcal{C}or_{k,\Lambda}^{prop}$ and \mathcal{A} be an arbitrary category enriched by Λ -modules.

- (1) Let $F : \mathcal{C} \rightarrow \mathcal{A}$ be an arbitrary Λ -linear functor which sends each birational morphism to an isomorphism. Then there exists one and only one (up to a natural equivalence) Λ -linear functor $F_{S_b} : \mathbf{Im}_{\Lambda}^{\mathcal{C}}\mathbf{H}_0^S \rightarrow \mathcal{A}$ such that the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{A} \\ \downarrow & \nearrow F_{S_b} & \\ \mathbf{Im}_{\Lambda}^{\mathcal{C}}\mathbf{H}_0^S & & \end{array}$$

is 2-commutative.

- (2) Let $F' : \mathcal{C}^{op} \rightarrow \mathcal{A}$ be an arbitrary Λ -linear functor which sends each birational morphism to an isomorphism. Then there exists one and only one (up to a natural equivalence) Λ -linear functor $F'_{S_b} : \mathbf{Im}_{\Lambda}^{\mathcal{C}}\mathbf{H}_0^S \rightarrow \mathcal{A}$ such that the diagram

$$\begin{array}{ccc} \mathcal{C}^{op} & \xrightarrow{F'} & \mathcal{A} \\ \downarrow & \nearrow F'_{S_b} & \\ (\mathbf{Im}_{\Lambda}^{\mathcal{C}}\mathbf{H}_0^S)^{op} & & \end{array}$$

is 2-commutative.

Proof. By Proposition 5.7, the functors $\mathcal{C} \rightarrow \mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^S$ and $\mathcal{C}^{op} \rightarrow (\mathbf{Im}_\Lambda^{\mathcal{C}} \mathbf{H}_0^S)^{op}$ are equivalences of categories. Thus this theorem follows from the universality of localizations of categories. \square

Remark 5.11. Like Remark 5.6, the functor F_{S_b} (resp. F'_{S_b}) in Theorem 5.10 is canonically extended to $\mathbf{NST}_k^{br}(\Lambda) \rightarrow \mathcal{A}$ (resp. $(\mathbf{NST}_k^{br}(\Lambda))^{op} \rightarrow \mathcal{A}$) if $\mathcal{C}or_{k,\Lambda}^{pr^{op}} \subseteq \mathcal{C}$ and \mathcal{A} is cocomplete.

5.3 Proper birational invariance

In this subsection, we prove that a proper birational morphism of smooth (not necessary proper) k -varieties induces an isomorphism of the zeroth \mathbb{A}^1 - and Suslin homology. This is a refinement of Asok's result [Aso12, Thm. 3.9].

Proposition 5.12. *Let $f : X \rightarrow Y$ be a proper birational morphism of smooth k -varieties.*

(1) *The induced morphisms*

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda) \rightarrow \mathbf{H}_0^{\mathbb{A}^1}(Y; \Lambda)$$

is an isomorphisms of sheaves.

(2) *The induced morphisms*

$$\mathbf{H}_0^S(X; \Lambda) \rightarrow \mathbf{H}_0^S(Y; \Lambda)$$

is an isomorphism of sheaves with transfers.

Proof. By Lemma 3.1, for every $M \in \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$ the induced map $M(Y) \rightarrow M(X)$ is an isomorphism. Thus [Aso12, Lem. 3.3] gives isomorphisms

$$\mathrm{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(\mathbf{H}_0^{\mathbb{A}^1}(Y; \Lambda), M) \cong M(Y) \xrightarrow{\cong} M(X) \cong \mathrm{Hom}_{\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)}(\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda), M)$$

for all $M \in \mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$. Thus Yoneda's lemma in $\mathcal{M}od_k^{\mathbb{A}^1}(\Lambda)$ induces an isomorphism

$$\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda) \xrightarrow{\cong} \mathbf{H}_0^{\mathbb{A}^1}(Y; \Lambda).$$

Similarly, [Aso12, Lem. 3.3] also gives an isomorphism

$$\mathrm{Hom}_{\mathbf{HI}_k(\Lambda)}(\mathbf{H}_0^S(Y; \Lambda), M') \xrightarrow{\cong} \mathrm{Hom}_{\mathbf{HI}_k(\Lambda)}(\mathbf{H}_0^S(X; \Lambda), M')$$

for all $M' \in \mathbf{HI}_k(\Lambda)$. Thus Yoneda's lemma in $\mathbf{HI}_k(\Lambda)$ induces an isomorphism

$$\mathbf{H}_0^S(X; \Lambda) \xrightarrow{\cong} \mathbf{H}_0^S(Y; \Lambda)$$

in $\mathbf{HI}_k(\Lambda)$. \square

6 Applications to \mathbb{A}^1 -homotopy theory

In this section, we give some applications to \mathbb{A}^1 -homotopy theory. A morphism $X \rightarrow Y$ in $\mathcal{S}m_k$ is called a S^1 -stable \mathbb{A}^1 -0-equivalence, if the induced morphism

$$\pi_0^S(\Sigma_{S^1}^\infty(X_+)) \rightarrow \pi_0^S(\Sigma_{S^1}^\infty(Y_+))$$

is an isomorphism. Note that $X \rightarrow Y$ is a S^1 -stable \mathbb{A}^1 -0-equivalence if and only if the induced morphism $\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z}) \rightarrow \mathbf{H}_0^{\mathbb{A}^1}(Y; \mathbb{Z})$ is an isomorphism by [AH11, Prop. 2.1]. When X and Y are proper, we also obtain other equivalent conditions.

Proposition 6.1. *Assume k admits a resolution of singularities. Let $f : X \rightarrow Y$ be a morphism of smooth proper k -schemes. Then the following conditions are equivalent.*

- (1) *The morphism f is a S^1 -stable \mathbb{A}^1 -0-equivalence.*
- (2) *The morphism f is an isomorphism in $S_b^{-1}\mathcal{S}m_k$.*
- (3) *The induced map $X(K)/R \rightarrow Y(K)/R$ is bijective for all $K \in \mathcal{F}_k$.*

Proof. Corollary 5.2 shows that (1) \Rightarrow (2). Moreover, (2) \Rightarrow (3) follows from [KS15, Thm. 6.6.3]. On the other hand, (3) is equivalent to the condition that the induced morphism $\pi_0^{b\mathbb{A}^1}(X) \rightarrow \pi_0^{b\mathbb{A}^1}(Y)$ is an isomorphism by [AM11, Thm. 6.2.1]. Thus Theorem 4.1 shows that the induced map $\mathbf{H}_0^{\mathbb{A}^1}(X; \mathbb{Z}) \rightarrow \mathbf{H}_0^{\mathbb{A}^1}(Y; \mathbb{Z})$ is an isomorphism. Hence, we have (3) \Rightarrow (1). \square

Next, we consider the \mathbb{A}^1 -connectedness of smooth proper varieties (see definition [MV99, p.110]). In [Aso12], Asok proves that a smooth proper k -variety X is \mathbb{A}^1 -connected if and only if the structure morphism $X \rightarrow \text{Spec } k$ induces an isomorphism of zeroth \mathbb{A}^1 -homology sheaves $\mathbf{H}_0^{\mathbb{A}^1}(X; \Lambda) \xrightarrow{\cong} \Lambda$.

Remark 6.2. Proposition 6.1 implies that a smooth proper k -variety X is \mathbb{A}^1 -connected if and only if the structure morphism $X \rightarrow \text{Spec } k$ is an isomorphism in $S_b^{-1}\mathcal{S}m_k$, when k admits a resolution of singularities. However, this assertion holds without the assumption on resolution of singularities. Indeed, X is \mathbb{A}^1 -connected if and only if $\#(X(K)/R) = 1$ for all $K \in \mathcal{F}_k$ by [AM11, Cor. 2.4.4]. On the other hand, by [KS15, Thm. 8.5.1] this is equivalent to the condition that the morphism $X \rightarrow \text{Spec } k$ is an isomorphism in $S_b^{-1}\mathcal{S}m_k$.

The following gives another equivalent condition of the \mathbb{A}^1 -connectedness of X .

Proposition 6.3. *Assume k admits a resolution of singularities. A smooth proper k -variety X is \mathbb{A}^1 -connected if and only if*

$$b_0^{\mathbb{A}^1}(X_{k(X)}) \leq 1 \leq b_0^{\mathbb{A}^1}(X).$$

Proof. “Only if” follows from [Aso12, Thm. 4.14]. We prove “if”. Since $b_0^{\mathbb{A}^1}(X) \neq 0$, the variety X has a k -rational point by Theorem 4.7. Moreover, since $X_{k(X)}$ also has a $k(X)$ -rational point, we have $b_0^{\mathbb{A}^1}(X_{k(X)}) = 1$. This implies

$$\#(X(k(X))/R) = \#(X_{k(X)}(k(X))/R) = 1$$

by Corollary 4.2. Thus [KS15, Thm. 8.5.1.] shows that the structure morphism $X \rightarrow \text{Spec } k$ is an isomorphism in $S_b^{-1}\mathcal{S}m_k$. Therefore, X is \mathbb{A}^1 -connected (see Remark 6.2). \square

Next, we prove a comparison result between \mathbb{A}^1 -homotopy and ordinary or étale homotopy. For a complex variety X , we denote X^{an} for the associated analytic space.

Proposition 6.4. *Assume k admits a resolution of singularities. Let $X \rightarrow Y$ be a S^1 -stable \mathbb{A}^1 -0-equivalence of smooth proper k -varieties.*

- (1) *The induced map $\pi_1^{\acute{e}t}(X) \rightarrow \pi_1^{\acute{e}t}(Y)$ is an isomorphism.*
- (2) *If $k = \mathbb{C}$, the continuous map $X^{an} \rightarrow Y^{an}$ is a 1-equivalence of topological spaces.*

Proof. Since the functor of (étale) fundamental groups is birational invariant of smooth proper varieties, this is regarded as a functor on $S_b^{-1}\mathcal{S}m_k^{prop}$. On the other hand, $X \rightarrow Y$ is an isomorphism in $S_b^{-1}\mathcal{S}m_k^{prop}$ by Proposition 6.1. Thus the map of (étale) fundamental groups induced by $X \rightarrow Y$ is an isomorphism. \square

Definition 6.5. A smooth k -scheme X is called *R -rigid*, if for all $K \in \mathcal{F}_k$ the scalar extension X_K has no rational curves over K .

By [AM11, Thm. 6.2.1], a smooth proper variety X is R -rigid if and only if the canonical morphism of sheaves $X \rightarrow \pi_0^{b\mathbb{A}^1}(X)$ is an isomorphism. We see examples of R -rigid varieties.

- Example 6.6.** (1) Let C be a geometrically irreducible smooth projective curve of genus ≥ 1 . Then C is R -rigid. Indeed, for every $K \in \mathcal{F}_k$ the scalar extension C_K has genus ≥ 1 and thus C_K has no rational curves over K .
- (2) An abelian variety X is also R -rigid. Indeed, for every $K \in \mathcal{F}_k$ the scalar extension X_K is a coproduct of abelian varieties over K and thus X_K has no rational curves over K .

We give a classification of R -rigid varieties up to a S^1 -stable \mathbb{A}^1 -0-equivalences.

Proposition 6.7. *Assume k admits a resolution of singularities. Let $f : X \rightarrow Y$ be a morphism of R -rigid smooth proper k -varieties. Then the following conditions are equivalent.*

- (1) *f is an isomorphism in $\mathcal{S}m_k$.*

- (2) f is birational.
- (3) f is stable birational.
- (4) f is an \mathbb{A}^1 -weak equivalence.
- (5) f is a S^1 -stable \mathbb{A}^1 -0-equivalence.

Proof. (1) \Rightarrow (2) \Rightarrow (3) and (1) \Rightarrow (4) \Rightarrow (5) are obvious. We first prove that (3) \Rightarrow (5). By [KS15, Thm. 1.7.2], $X \rightarrow Y$ is an isomorphism in $S_b^{-1}\mathcal{S}m_k$. Thus by Proposition 6.1, $X \rightarrow Y$ is a S^1 -stable \mathbb{A}^1 -0-equivalence. Next, we prove that (5) \Rightarrow (1). Then the induced map $X(k(U))/R \rightarrow Y(k(U))/R$ is bijective for all $U \in \mathcal{S}m_k^{var}$ by Proposition 6.1. Hence, Lemma 2.2 shows that the morphism of birational sheaves

$$\pi_0^{b\mathbb{A}^1}(X) \rightarrow \pi_0^{b\mathbb{A}^1}(Y) \tag{6.1}$$

is an isomorphism. On the other hand, since X and Y are R -rigid, there exist canonical isomorphisms of sheaves $X \cong \pi_0^{b\mathbb{A}^1}(X)$ and $Y \cong \pi_0^{b\mathbb{A}^1}(Y)$. Thus by Yoneda's lemma in $\mathcal{S}m_k$, (6.1) induces an isomorphism $X \xrightarrow{\cong} Y$. \square

References

- [Aso12] A. Asok, *Birational invariants and \mathbb{A}^1 -connectedness*, J. Reine Angew. Math., **681** (2012) 39-64.
- [AH11] A. Asok and C. Haesemeyer, *Stable \mathbb{A}^1 -homotopy and R -equivalence*, J. Pure Appl. Algebra **215**(10) (2011) 2469-2472.
- [AM11] A. Asok and F. Morel, *Smooth varieties up to \mathbb{A}^1 -homotopy and algebraic h -cobordisms*, Adv. Math. **227**(5) (2011) 1990-2058.
- [CT95] J.-L. Colliot-Thélène, *Birational invariants, purity and the Gersten conjecture*, K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992), Proc. Sympos. Pure Math., **58**, Part 1, Amer. Math. Soc., Providence, RI (1995) 164.
- [Fal83] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math., **73** (1983) 349-366.
- [GZ67] P. Gabriel and M. Zisman, *Calculus of fractions and homotopy theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band **35**, Springer (1967)
- [KS07] B. Kahn and R. Sujatha, *A few localisation theorems*, Homology Homotopy Appl., **9**(2) (2007) 137-161.
- [KS15] B. Kahn and R. Sujatha, *Birational geometry and localisation of categories*, Doc. Math., Extra vol.: Alexander S. Merkurjev's sixtieth birthday (2015) 277-334.

- [KS17] B. Kahn and R. Sujatha, *Birational motives, II: Triangulated birational motives*, Int. Math. Res. Not. IMRN, **22** (2017) 67786831.
- [Kol99] J. Kollár, *Rationally connected varieties over local fields*, Ann. of Math., **150** (1999) 357-367.
- [Man86] Yu.I. Manin, *Cubic forms*, Algebra, geometry, arithmetic. Translated from the Russian by M. Hazewinkel. Second edition. North-Holland Mathematical Library, **4**. North-Holland Publishing Co., Amsterdam, 1986.
- [Mor05] F. Morel, *The stable \mathbb{A}^1 -connectivity theorems*, K-theory, **35**(1-2) (2005) 1-68.
- [Mor12] F. Morel, *\mathbb{A}^1 -algebraic topology over a field*, Lecture Notes in Math., **2052**, Springer, Heidelberg (2012).
- [MV99] F. Morel and V. Voevodsky, *\mathbb{A}^1 -homotopy theory of schemes*, Publ. Math. IHES, **90** (1999) 45-143.
- [MVW] V. Mazza, V. Voevodsky, and C. Weibel, *Lecture notes on motivic cohomology*, volume 2 of Clay Mathematics Monographs, American Mathematical Society, Providence, RI (2006).
- [SV96] A. Suslin and V. Voevodsky, *Singular homology of abstract varieties*, Invent. Math., **123**(1) (1996) 61-94.
- [Voe00] V. Voevodsky, *Triangulated categories of motives over a field*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., **143**, Princeton Univ. Press, Princeton, NJ (2000) 188-238.