

Landscapes of data sets and functoriality of persistent homology

Wojciech Chachólski^a, Alessandro De Gregorio^b, Nicola Quercioli^{c,*}, Francesca Tombari^a

^a*Mathematics Department, KTH, Lindstedtsvagen 25, Stockholm, 11428, Sweden*

^b*Mathematics Department, Politecnico di Torino, Corso Duca degli Abruzzi 24, Torino, 10129, Italy*

^c*Mathematics Department, University of Bologna, Piazza di Porta S. Donato 5, Bologna, 40126, Italy*

Abstract

The aim of this article is to describe a new perspective on functoriality of persistent homology and explain its intrinsic symmetry that is often overlooked. A data set for us is a finite collection of functions, called measurements, with a finite domain. Such a data set might contain internal symmetries which are effectively captured by the action of a set of the domain endomorphisms. Different choices of the set of endomorphisms encode different symmetries of the data set. We describe various category structures on such enriched data sets and prove some of their properties such as decompositions and morphism formations. We also describe a data structure, based on coloured directed graphs, which is convenient to encode the mentioned enrichment. We show that persistent homology preserves only some aspects of these landscapes of enriched data sets however not all. In other words persistent homology is not a functor on the entire category of enriched data sets. Nevertheless we show that persistent homology is functorial locally. We use the concept of equivariant operators to capture some of the information missed by persistent homology.

Keywords: persistent homology, topological data analysis, equivariant operators,

2020 MSC: 55N31, 62R40

1. Introduction

In this article we give an answer to the question: what is persistent homology a functor of?

*Corresponding author.

Email addresses: `wojtek@kth.se` (Wojciech Chachólski), `alessandro.degregorio@polito.it` (Alessandro De Gregorio), `nicola.quercioli2@unibo.it` (Nicola Quercioli), `tombari@kth.se` (Francesca Tombari)

We will consider data sets given by finite sets of functions on a finite set X with real values. There are several important consequences of data sets having this form. For example, they endow X with a pseudometric, enabling us to extract non-trivial homological information in form of persistent homology, one of the key invariants studied in Topological Data Analysis. A single measurement does not contain any higher non-trivial homological information. Sets of measurements however do. Thus it is essential that measurements, on a given set X , are grouped together to form various data sets. In this case persistent homology becomes a non-expansive (1-Lipschitz) function $PH_d^\Phi: \Phi \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$, assigning to each measurement in the data set Φ a tame vector space parametrized by $[0, \infty) \times \mathbf{R}$. It is important to notice that the choice of a set of measurements on X affects the pseudometric defined on it. One can use this fact to change the metric on X in order to extract more meaningful information from persistent homology. For example consider X to be a finite sample of points on a circle. If Φ consists of only one function given by the x -coordinate, then the persistent homology of this measurement is trivial in degrees greater than 0. If we enlarge the data set by adding to the x -coordinate the function given by precomposing x with rotation by 90 degrees, then the persistent homology of the function x with respect to this bigger data set gains a non-trivial homology in degree 1. This illustrates how our knowledge of an object is affected by the number and the type of measurements done on it. Furthermore in this example we gain additional information by enlarging the set of measurements through the action of some of the endomorphisms of X on the existing measurements. We can then take advantage of these actions to inject geometrical features of our choice on a given data set. For exhibiting and extracting interesting homological features of data sets, such actions are therefore important.

A data set Φ is naturally equipped with an action of the monoid of its operations $\text{End}_\Phi(X)$, which are endomorphisms of X preserving Φ . This action gives the set Φ a structure of Grothendieck graph. Persistent homology turns out to be a functor indexed by this graph, rather than simply a function. Thus not only persistent homology can be assigned to individual measurements in a data set, but operations can be used to compare persistent homologies of different measurements. That is what we call local functorial properties of persistent homology.

Persistent homology also has certain global functorial properties. There are various ways of representing data in the form of sets of measurements, we might choose different units or different parametrizations of a domain of measurements, or we might need to focus only on certain operations such as rotations. Furthermore, the same measurements might be part of different data sets. These are some of the reasons why it is essential to be able to compare data sets equipped with different structures. For that purpose we introduce the notion of incarnations of data sets to encode different actions, and SEOs to compare incarnations. An incarnation of a data set Φ is an action of a subset $M \subset \text{End}_\Phi(X)$. A SEO (set equivariant operator) between two incarnations (Φ, M) and (Ψ, N) is a pair consisting of a map $T: M \rightarrow N$ and an equivariant

(with respect to T) function $\alpha : \Phi \rightarrow \Psi$. The use of this kind of operators for the comparison of incarnations of data sets has been inspired by [1, 2], where GENEOS (group equivariant non-expansive operators) are introduced and used for applications to neural networks. If a SEO is geometric, then there is a comparison map between persistent homologies of the incarnations connected by the SEO. However if a SEO is not geometric, such as the change of units SEO, there is no direct comparison of persistent homologies of the involved incarnations. Such SEOs therefore exhibit diverse homological features of data sets enhancing the analysis. This suggests complementarity of these operators and persistent homology for a geometric analysis of a data set. Consider the change of unit as an example. In general it is the SEO obtained by composing measurements in a data set by a given real valued function defined on the real numbers. Multiplication by -1 is an example of such a SEO. It has the effect of turning the sub-level sets persistent homology of a measurement into its super-level sets persistent homology, leading in general to a completely different information about the data set. The outcome consists of two different points of view on the same object, that are not functorially comparable, but together may enhance the accuracy of the analysis of the object of interest.

2. Data sets

For us a data set, which we regard as a point in the data landscape, is given by a finite set of real valued functions on some finite set X also called measurements:

$$\Phi = \{\phi_i : X \rightarrow \mathbf{R} \mid i = 1, \dots, m\}.$$

We define $\text{dom}(\Phi)$, the **domain** of dataset Φ , to be the set X which is the domain of the functions in Φ . The most fundamental aspect of a data set Φ is that it is a set. All such data sets with different domains form a category with functions as morphisms. This is the most primitive landscape of data sets. The nature of our data sets however can be used to impose more intricate structures and more meaningful landscapes. This is reminiscent of the case of groups. The most fundamental aspect of a group is that it is a set. However the category whose morphisms are group homomorphism is a much more meaningful landscape in which to study relationships between groups. To understand relationships between topological groups, the category with continuous group homomorphisms provides an even more meaningful landscape.

In this most primitive landscape however we can already perform products and coproducts. Let $\phi : X \rightarrow \mathbf{R}$ and $\psi : Y \rightarrow \mathbf{R}$ be functions. Define $\phi + \psi : X \coprod Y \rightarrow \mathbf{R}$ to be the function that maps x in X to $\phi(x)$ and y in Y to $\psi(y)$. The **coproduct** of two data sets Φ and Ψ , denoted by $\Phi \coprod \Psi$, is defined to be the data set given by the measurements $\{\phi + 0 \mid \phi \in \Phi\} \cup \{0 + \psi \mid \psi \in \Psi\}$ on $X \coprod Y$. Their **product**, denoted by $\Phi \times \Psi$, is defined to be the data set given

by the measurements $\{\phi + \psi \mid \phi \in \Phi \text{ and } \psi \in \Psi\}$ on $X \coprod Y$. The functions:

$$\begin{array}{ccc}
 & \xrightarrow[\text{pr}_\Phi]{\phi + \psi \mapsto \phi} & \Phi \\
 \Phi \times \Psi & & \xrightarrow[\text{in}_\Phi]{\phi \mapsto \phi + 0} \Phi \coprod \Psi \\
 & \xrightarrow[\phi + \psi \mapsto \psi]{\text{pr}_\Psi} & \Psi \\
 & & \xrightarrow[\psi \mapsto 0 + \psi]{\text{in}_\Psi}
 \end{array}$$

satisfy the following universal properties, which justify the names coproduct and product:

- for any data set Π , and any two functions $\alpha: \Phi \rightarrow \Pi$ and $\beta: \Psi \rightarrow \Pi$, there is a unique function $\mu: \Phi \coprod \Psi \rightarrow \Pi$ for which $\mu \text{in}_\Phi = \alpha$ and $\mu \text{in}_\Psi = \beta$;
- for any data set Π , and any two functions $\alpha: \Pi \rightarrow \Phi$ and $\beta: \Pi \rightarrow \Psi$, there is a unique function $\mu: \Pi \rightarrow \Phi \times \Psi$ for which $\text{pr}_\Phi \mu = \alpha$ and $\text{pr}_\Psi \mu = \beta$.

Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a function. By composing with f , a data set Φ is transformed into a new data set $f\Phi := \{f\phi \mid \phi \in \Phi\}$. This operation is called **change of units** along f . The symbol $f-: \Phi \rightarrow f\Phi$ denotes the function mapping ϕ to $f\phi$. For example let $f: \mathbf{R} \rightarrow \mathbf{R}$ map $\{r \in \mathbf{R} \mid r < 0\}$ to -1 and $\{r \in \mathbf{R} \mid r \geq 0\}$ to 1 . Consider $X = \{x_1, x_2\}$, two data sets $\{1, 2\}$ and $\{-1, 1\}$ given by the constant functions $-1, 1, 2: X \rightarrow \mathbf{R}$, and a function $\alpha: \{1, 2\} \rightarrow \{-1, 1\}$ mapping 1 to -1 and 2 to 1 . Then $f\{1, 2\} = \{1\}$ and $f-: \{-1, 1\} \rightarrow f\{-1, 1\}$ is the identity. Thus there is no function $f\{1, 2\} \rightarrow f\{-1, 1\}$ making the following diagram commutative:

$$\begin{array}{ccc}
 \{1, 2\} & \xrightarrow{f-} & f\{1, 2\} = \{1\} \\
 \alpha \downarrow & & \downarrow \\
 \{-1, 1\} & \xrightarrow{f-\text{id}} & f\{-1, 1\} = \{-1, 1\}
 \end{array}$$

Consequently, for that f there is no functor F assigning to a data set Φ its change of units $f\Phi$ along f for which $f-: \Phi \rightarrow f\Phi$ is a natural transformation between F and the identity functor. If f is invertible, then $f-: \Phi \rightarrow f\Phi$ is a bijection whose inverse is given by $f^{-1}-$. The association $(\alpha: \Phi \rightarrow \Psi) \mapsto ((f-)\alpha(f^{-1}-): f\Phi \rightarrow f\Psi)$ is a functor for which $f-: \Phi \rightarrow f\Phi$ is a natural transformation between this functor and the identity functor. Changing the units along any function preserves products and coproducts i.e., $f(\Phi \coprod \Psi)$ is isomorphic to $f(\Phi) \coprod f(\Psi)$, and $f(\Phi \times \Psi)$ is isomorphic to $f(\Phi) \times f(\Psi)$. A similar reasoning is used in [7] to study brain data, in order to obtain results that are invariant under transformations given by change of units with invertible functions, and in [8] to study metric spaces that are isometric up to a rescaling of the distance functions.

Let Φ be a data set with the domain X . By composing a function $f: Y \rightarrow X$ with the measurements in Φ , we obtain a new data set $\Phi f := \{\phi f \mid \phi \in \Phi\}$ with the domain Y . This operation is called **domain change** along f . The symbol $-f: \Phi \rightarrow \Phi f$ denotes the function that maps ϕ to ϕf .

Let $f_1: Z_1 \rightarrow X$ and $f_2: Z_2 \rightarrow Y$ be functions and $f_1 \amalg f_2: Z_1 \amalg Z_2 \rightarrow X \amalg Y$ be their coproduct. For any datasets Φ and Ψ with $\text{dom}(\Phi) = X$ and $\text{dom}(\Psi) = Y$, the following equalities hold:

$$(\Phi \amalg \Psi)(f_1 \amalg f_2) = \Phi f_1 \amalg \Psi f_2, \quad (\Phi \times \Psi)(f_1 \amalg f_2) = \Phi f_1 \times \Psi f_2.$$

3. Metrics and persistent homology

We can think about a data set Φ as a subset $\Phi \subset \mathbf{R}^{|X|}$. Via this inclusion Φ inherits a metric induced by the infinity norm $\|v\|_\infty = \max\{|v_i|\}$ on $\mathbf{R}^{|X|}$. We use the symbol $\|\phi - \psi\|_\infty$ to denote the distance between ϕ and ψ in Φ . The considered data sets are not just sets anymore but metric spaces. Therefore non-expansive (1-Lipschitz) functions between data sets play a special role. For example, let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a function. If f is non-expansive, then so is the change of units along f , $f-: \Phi \rightarrow f\Phi$, that maps ϕ to $f\phi$. The domain change $-h: \Phi \rightarrow \Phi h$ is non-expansive along any h . Non-expansiveness is an important assumption to prove some stability results in [1] and it is also reasonable in applications, since it is important that these functions between data sets do not alter the information too much.

By taking all the measurements of Φ together, we can form a function $[\phi_1 \cdots \phi_m]: X \rightarrow \mathbf{R}^m$. Via this function, X inherits a pseudometric d_Φ induced by the infinity norm on \mathbf{R}^m . Explicitly $d_\Phi(x, y) := \max_{1 \leq i \leq m} |\phi_i(x) - \phi_i(y)|$. This metric plays a fundamental role as it permits us to extract persistent homologies (see [3, 6]). In this article, **persistent homology** of a data set Φ with coefficients in a field and in a given degree d assigns a vector space $PH_d^\Phi(\phi)_{r,s}$ to each measurement ϕ in Φ , for every (r, s) in $[0, \infty) \times \mathbf{R}$, and it is defined as:

$$PH_d^\Phi(\phi)_{r,s} := H_d(\text{VR}_r(\phi \leq s, d_\Phi)), \text{ where:}$$

- $\phi \leq s := \phi^{-1}(-\infty, s]$;
- $\text{VR}_r(\phi \leq s, d_\Phi)$ is the **Vietoris-Rips** complex whose simplices are given by the subsets $\sigma \subset (\phi \leq s)$ of diameter not exceeding r with respect to d_Φ ;
- H_d is the homology in degree d with coefficients in a given field.

If $s \leq s'$ and $r \leq r'$, then $(\phi \leq s) \subset (\phi \leq s')$ and therefore $\text{VR}_r(\phi \leq s) \subset \text{VR}_{r'}(\phi \leq s')$. The linear function induced on homology by this inclusion is denoted by:

$$PH_d^\Phi(\phi)_{(r,s) \leq (r',s')} : PH_d^\Phi(\phi)_{r,s} \rightarrow PH_d^\Phi(\phi)_{r',s'}.$$

These functions form a functor $PH_d^\Phi(\phi)$ indexed by the poset $[0, \infty) \times \mathbf{R}$ with values in the category of vector spaces. Since X is finite, $PH_d^\Phi(\phi)$ is **tame** (see [11]). This means that values of $PH_d^\Phi(\phi)$ are finite dimensional, and there are two finite sequences $0 = r_0 < r_1 < \cdots < r_m$ in $[0, \infty)$ and $s_0 < s_1 <$

$\dots < s_l = \infty$ in \mathbf{R} such that $PH_d^\Phi(\phi)$, restricted to subposets of the form $[r_i, r_{i+1}) \times (\infty, s_0) \subset [0, \infty) \times \mathbf{R}$ and $[r_i, r_{i+1}) \times [s_j, s_{j+1}) \subset [0, \infty) \times \mathbf{R}$, is constant. The category of such functors is denoted by $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$. Thus a data set Φ leads to a function assigning to each measurement ϕ its persistent homology in a given degree:

$$PH_d^\Phi: \Phi \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}).$$

Next, recall a definition of the **interleaving metric** in the direction of the vector $(0, 1)$ on $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ (see [9]). Let P and Q be in $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$.

- P and Q are ϵ -**interleaved** if, for all (r, s) in $[0, \infty) \times \mathbf{R}$, there are linear functions $f_{s,r}: P_{r,s} \rightarrow Q_{r,s+\epsilon}$ and $g_{s,r}: Q_{r,s} \rightarrow P_{r,s+\epsilon}$ making the following diagram commutative:

$$\begin{array}{ccccc} & & P_{r,s} & \xrightarrow{P_{(r,s) < (r,s+2\epsilon)}} & P_{r,s+2\epsilon} \\ & \nearrow g_{r,s-\epsilon} & \searrow f_{s,r} & \nearrow g_{r,s+\epsilon} & \searrow f_{r,s+2\epsilon} \\ Q_{r,s-\epsilon} & \xrightarrow{Q_{(r,s-\epsilon) < (r,s+\epsilon)}} & Q_{r,s+\epsilon} & \xrightarrow{Q_{(r,s+\epsilon) < (r,s+3\epsilon)}} & Q_{r,s+3\epsilon} \end{array}$$

- $d_{\bowtie}(P, Q) := \inf\{\epsilon \in [0, \infty) \mid P \text{ and } Q \text{ are } \epsilon\text{-interleaved}\}$.

The function $P, Q \mapsto d_{\bowtie}(P, Q)$ is an extended (∞ is allowed) metric on the set $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ called interleaving metric in the direction of the vector $(0, 1)$.

Proposition 1. *The function $PH_d^\Phi: \Phi \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ is non-expansive if the set Φ is equipped with ∞ -norm metric $\|\phi - \psi\|_\infty$ and the set $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ is equipped with the interleaving metric in the direction of the vector $(0, 1)$.*

Proof. Let $\phi, \psi: X \rightarrow \mathbf{R}$ be measurements in Φ and $\epsilon = \|\phi - \psi\|_\infty$. For every s in \mathbf{R} , the sublevel set $\phi \leq s$ is a subset of $\psi \leq s + \epsilon$, and $\psi \leq s$ is a subset of $\phi \leq s + \epsilon$. This translates into inclusions:

$$\text{VR}_r(\phi \leq s, d_\Phi) \subset \text{VR}_r(\psi \leq s + \epsilon, d_\Phi) \quad \text{VR}_r(\psi \leq s, d_\Phi) \subset \text{VR}_r(\phi \leq s + \epsilon, d_\Phi)$$

leading functions:


$$f_{s,r}: PH_d^\Phi(\phi)_{r,s} \rightarrow PH_d^\Phi(\psi)_{r,s+\epsilon} \quad g_{s,r}: PH_d^\Phi(\psi)_{r,s} \rightarrow PH_d^\Phi(\phi)_{r,s+\epsilon}.$$

These functions provide ϵ interleaving between $PH_d^\Phi(\phi)$ and $PH_d^\Phi(\psi)$, giving $\|\phi - \psi\|_\infty \geq d_{\bowtie}(PH_d^\Phi(\phi), PH_d^\Phi(\psi))$. \square

A measurement $\phi: X \rightarrow \mathbf{R}$ can be part of many data sets and its persistent homology depends on what data set this function is part of. For example, let $X = \{x_1, x_2, x_3, x_4\}$ and $\phi, \psi: X \rightarrow \mathbf{R}$ be measurements defined as follows:

$$\begin{array}{lll} \phi(x_1) = -1 & \phi(x_2) = \phi(x_3) = 0 & \phi(x_4) = 1 \\ \psi(x_3) = -1 & \psi(x_1) = \psi(x_4) = 0 & \psi(x_2) = 1 \end{array}$$

The measurement ϕ is part of two data sets $\Phi = \{\phi\}$ and $\Psi = \{\phi, \psi\}$. The induced pseudometrics d_Φ and d_Ψ on X can be depicted by the following diagrams where the continuous, dashed, and dotted lines indicate distance 0, 1 and 2 respectively:



In this case $PH_1^\Phi(\phi)_{r,s} = 0$ for all r and s , however:

$$\dim PH_1^\Psi(\phi)_{r,s} = \begin{cases} 1 & \text{if } 1 \leq s \text{ and } 1 \leq r < 2 \\ 0 & \text{otherwise} \end{cases}$$

To understand persistent homology, it is therefore paramount to understand how it changes when data sets change and here functoriality plays an essential role.

Let Φ and Ψ be data sets consisting of measurements on X and Y respectively. A function $\alpha: \Phi \rightarrow \Psi$ is called **geometric** if there is a function $f: Y \rightarrow X$, called a **realization** of α , making the following diagram commutative for every ϕ in Φ :

$$\begin{array}{ccc}
 Y & \xrightarrow{\alpha(\phi)} & \mathbf{R} \\
 f \downarrow & \searrow & \\
 X & \xrightarrow{\phi} &
 \end{array}$$

For example $-f: \Phi \rightarrow \Phi f$ is geometric, as it is realized by f .

The commutativity of the triangle above has two consequences. First, f is non-expansive with respect to the pseudometrics d_Φ on X and d_Ψ on Y . Second, for s in \mathbf{R} and ϕ in Φ , the subset $(\alpha(\phi) \leq s) \subset Y$ is mapped via f into $(\phi \leq s) \subset X$, i.e., the following diagram commutes:

$$\begin{array}{ccc}
 \alpha(\phi) \leq s & \hookrightarrow & Y \\
 f \downarrow & & \searrow \alpha(\phi) \\
 \phi \leq s & \hookrightarrow & X \xrightarrow{\phi} \mathbf{R}
 \end{array}$$

The realization f induces therefore a map of Vietoris-Rips complexes and their homologies:

$$\begin{array}{ccc}
 f_{s,r}: \text{VR}_r(\alpha(\phi) \leq s, d_\Psi) & \rightarrow & \text{VR}_r(\phi \leq s, d_\Phi); \\
 PH_d^\Psi(\alpha(\phi))_{r,s} & & PH_d^\Phi(\phi)_{r,s} \\
 \parallel & \xrightarrow{H_d(f_{r,s})} & \parallel \\
 H_d(\text{VR}_r(\alpha(\phi) \leq s, d_\Psi)) & \rightarrow & H_d(\text{VR}_r(\phi \leq s, d_\Phi)).
 \end{array}$$

If $f, f': Y \rightarrow X$ are two realizations of α , then for y in Y , $d_\Phi(f(y), f'(y)) = 0$, hence they are points of the same simplex in the Vietoris-Rips complex, implying

that $f_{r,s}$ and $f'_{r,s}$ are homotopic for all r and s . Consequently, $H_d(f_{r,s}) = H_d(f'_{r,s})$. The linear function $H_d(f_{r,s})$ depends therefore only on α and it is independent on the choice of its realization f . We denote this function by:

$$PH_d^\alpha(\phi)_{r,s}: PH_d^\Psi(\alpha(\phi))_{r,s} \rightarrow PH_d^\Phi(\phi)_{r,s}.$$

These functions are natural in r and s and induce a morphism in the category $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ between persistent homologies:

$$PH_d^\alpha(\phi): PH_d^\Psi(\alpha(\phi)) \rightarrow PH_d^\Phi(\phi).$$

If $\alpha: \Phi \rightarrow \Psi$ and $\beta: \Psi \rightarrow \Xi$ are geometric functions realized by $f: Y \rightarrow X$ and $g: Z \rightarrow Y$, then the composition $\beta\alpha: \Phi \rightarrow \Xi$ is also geometric, and realized by the composition $fg: Z \rightarrow X$. Consequently, for every measurement ϕ in Φ , $PH_d^{\beta\alpha}(\phi) = PH_d^\alpha(\phi)PH_d^\beta(\alpha(\phi))$, that assures the commutativity of the diagram:

$$\begin{array}{ccccc} PH_d^\Xi(\beta\alpha(\phi)) & \xrightarrow{PH_d^\beta(\alpha(\phi))} & PH_d^\Psi(\alpha(\phi)) & \xrightarrow{PH_d^\alpha(\phi)} & PH_d^\Phi(\phi) \\ & & \searrow & \nearrow & \\ & & PH_d^{\beta\alpha}(\phi) & & \end{array}$$

For any $\alpha: \Phi \rightarrow \Psi$, taking persistent homology leads to two functions on Φ :

$$\begin{array}{ccc} & PH_d^\Phi & \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}) \\ \Phi & \searrow & \\ & \alpha & \rightarrow \Psi \xrightarrow{PH_d^\Psi} \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}) \end{array}$$

These functions rarely coincide. However, when α is geometric, we can use the morphisms $PH_d^\alpha(\phi): PH_d^\Psi(\alpha(\phi)) \rightarrow PH_d^\Phi(\phi)$ to compare the values of these two functions on Φ . For non-geometric α , we are not equipped with such comparison morphisms and there is no reason for such a comparison to even exist. For example, consider the change of unit along the function $f: \mathbf{R} \rightarrow \mathbf{R}$, $f(x) := -x$. Then $f -: \Phi \rightarrow f\Phi$ is an isomorphism. In this case

$$PH_d^\Phi(\phi)_{r,s} := H_d(\text{VR}_r(\phi \leq s, d_\Phi)) \mid (f-)PH_d^{f\Phi}(\phi) = H_d(\text{VR}_r(\phi \geq -s, d_\Phi)).$$

Thus PH_d^Φ encodes information about sub-level sets of the measurements in Φ and $(f-)PH_d^{f\Phi}$ encodes information about super-level sets of the measurements. These persistent homologies encode therefore the same information as the so called extended persistence (see [4, 10]).

4. Actions

To describe symmetries of a data set Φ with domain X , we consider operations on X that convert measurements into measurements. By definition a

Φ -operation is a function $g: X \rightarrow X$ such that, for every measurement ϕ in Φ , the composition ϕg also belongs to Φ . If $g: X \rightarrow X$ is such an operation, then, for all ϕ and ψ in Φ :

$$\|\phi - \psi\|_\infty = \max_{x \in X} |\phi(x) - \psi(x)| \geq \max_{x \in \text{im}(g)} |\phi(x) - \psi(x)| = \|\phi g - \psi g\|_\infty.$$

Thus the function $-g: \Phi \rightarrow \Phi$ that maps ϕ to ϕg is non-expansive.

The composition of Φ -operations is again a Φ -operation, and the identity function id_X is also a Φ -operation. In this way the set of Φ -operations with the composition becomes a unitary monoid, called the **structure monoid** of Φ , and denoted by:

$$\text{End}_\Phi(X) = \{g: X \rightarrow X \mid \phi g \in \Phi \text{ for every } \phi \in \Phi\} \subset \text{End}(X).$$

A Φ -operation g is invertible if there is a Φ -operation h such that $gh = hg = \text{id}_X$. Since Φ is finite, a Φ -operation is invertible if and only if it is a bijection. Their collection is denoted by:

$$\text{Aut}_\Phi(X) = \{g: X \rightarrow X \mid g \text{ is a bijection, and } \phi g \in \Phi \text{ for every } \phi \in \Phi\}.$$

With the composition operation, $\text{Aut}_\Phi(X)$ becomes a group for which the inclusion $\text{Aut}_\Phi(X) \subset \text{End}_\Phi(X)$ is a monoid homomorphism.

A data set Φ is equipped with an associative right action:

$$\Phi \times \text{End}_\Phi(X) \rightarrow \Phi, \quad (\phi, g) \mapsto \phi g.$$

Thus Φ is not just a set, but a set with an action of the monoid $\text{End}_\Phi(X)$. To encode the symmetries of Φ induced by this action, we consider its incarnations.

An **incarnation** of Φ is a choice of a subset $M \subset \text{End}_\Phi(X)$ (in general, not necessarily a submonoid). An incarnation is denoted as a pair (Φ, M) . We think about M as an additional structure on Φ . An incarnation of the form (Φ, M) is called an M -incarnation. We also refer to an M -incarnation as an M -action. The choice of an M -action on Φ encodes certain symmetries of Φ . Different choices of M can encode different symmetries. This flexibility is important in applications. For example in data sets that represent images, we might want to focus on rotational symmetries, so we may use an appropriate action on the data set to inject the corresponding geometry. The incarnation $(\Phi, \text{End}_\Phi(X))$ is an example of a incarnation called universal.

An incarnation (Φ, M) is called a **monoid incarnation** if $M \subset \text{End}_\Phi$ is a submonoid, and our convention here is that all such submonoids contain the identity element. If (Φ, M) is an incarnation, we use the symbol $(\Phi, \langle M \rangle)$ to denote the monoid incarnation where $\langle M \rangle \subset \text{End}_\Phi(X)$ is the submonoid generated by M .

If a submonoid $M \subset \text{End}_\Phi(X)$ is a group, then (Φ, M) is called a **group incarnation**. The incarnation $(\Phi, \text{Aut}_\Phi(X))$ is an example of a group incarnation called universal.

Let (Φ, M) be an incarnation for which any element g in M is a bijection. Such incarnations are called **group-like**. For group like incarnations (Φ, M)

the finiteness implies that the monoid $\langle M \rangle$ is in fact a subgroup of $\text{Aut}_\Phi(X)$. Thus any group-like incarnation (Φ, M) leads to a group incarnation $(\Phi, \langle M \rangle)$.

Let (Φ, M) be an incarnation. For a subset $\Omega \subset \Phi$, the symbol ΩM denotes the set of all the measurements in Φ which either belong to Ω or are of the form $\omega g_1 \cdots g_k$, for some ω in Ω and some sequence of elements g_1, \dots, g_k in M . If $\Omega M = \Phi$, then Ω is said to **generate** the incarnation (Φ, M) . In the case (Φ, M) is a monoid incarnation, then any element in ΩM is of the form ωg for some ω in Ω and g in M . Note that $\Omega M = \Omega \langle M \rangle$ for every incarnation (Φ, M) .

If ψ belongs to $\phi M := \{\phi\}M$, then ψ is said to be a **deformation** of ϕ . If (Φ, M) is a group incarnation, then the relation of being a deformation is an equivalence relation. For a general incarnation however being a deformation can fail to be even a symmetric relation. Two measurements in Φ are said to be **connected** if they are related by the equivalence relation generated by the relation of being a deformation. The symbol Φ/M denotes the partition of Φ induced by this equivalence relation. We refer to Φ/M as the **quotient** of the incarnation (Φ, M) . The partitions Φ/M and $\Phi/\langle M \rangle$ coincide. If (Φ, M) is a group incarnation, then Φ/M coincide with the orbit partition of the usual group action of M on Φ .

Let (Φ, M) be an incarnation. For a measurement ψ in Φ , the symbol $[\psi]$ denotes the block in Φ/M containing ψ . Explicitly, $[\psi]$ is the subset of Φ consisting of all the measurements connected to ψ . Note that, for all g in M , if ϕ is connected to ψ , then ϕg is also connected to ψ . We thus have the following inclusions:

$$\begin{array}{ccc} M & \hookrightarrow & \text{End}_\Phi(X) \\ \downarrow & & \downarrow \\ \text{End}_{[\psi]}(X) & \hookrightarrow & \text{End}(X) \end{array}$$

The M incarnation $([\psi], M)$ of the block $[\psi]$, given by the above inclusions $M \subset \text{End}_{[\psi]}(X)$, is called a **block incarnation** of (Φ, M) . In this way we can think about $[\psi]$ and $([\psi], M)$ as a new data set.

An incarnation (Φ, M) is called **transitive** if all the elements in Φ are connected to each other. For example, let M be a finite submonoid of $\text{End}(X)$. For a given function $\phi: X \rightarrow \mathbf{R}$, define a data set $\phi M := \{\phi g \mid g \in M\}$ to consist of all functions of the form $x \mapsto \phi(g(x))$ for all g in M . Then every $g: X \rightarrow X$ in M is a ϕM -operation. The obtained incarnation $(\phi M, M)$ is transitive. Any transitive group incarnation is of such form. For all measurements ϕ in any incarnation (Φ, M) , the block incarnation $([\phi], M)$ is transitive. Any transitive incarnation is of this form.

Let (Φ, M) be an incarnation. A subset $\Omega \subset \Phi$ is called **independent** if no element in Ω is a deformation of any other element in Ω , explicitly: $\omega \notin \omega' M$ for all $\omega \neq \omega'$ in Ω .

A **basis** of (Φ, M) is an independent subset $\Omega \subset \Phi$ such that $\Omega M = \Phi$ (Ω generates (Φ, M)).

Two measurements ψ and ϕ are called **indistinguishable** if ψ is a deformation of ϕ and ϕ is a deformation of ψ . If (Φ, M) is a group incarnation, then

ψ and ϕ are indistinguishable if and only if $\psi = \phi g$ for some g in M , i.e., if ψ is a deformation of ϕ .

Proposition 2. 1. *Every incarnation has a basis.*

2. *Let $\Omega, \Omega' \subset \Phi$ be two bases of an incarnation (Φ, M) . Then there is a bijection $\sigma: \Omega \rightarrow \Omega'$ such that ω and $\sigma(\omega)$ are indistinguishable for every ω in Ω .*

Proof. (1): Let (Φ, M) be an incarnation. Choose $\Omega \subset \Phi$ to be an independent subset for which ΩM is maximal. Existence of Ω is guaranteed by finiteness of Φ . We claim that $\Omega M = \Phi$ and hence Ω is a basis. If this is not the case, let ψ be in $\Phi \setminus \Omega M$. Define $\Omega' = \{\psi\} \cup \{\omega \in \Omega \mid \omega \notin \{\psi\}M\}$. Then $\Omega'M$ contains Ω and hence ΩM . It also contains ψ . Since Ω' is independent, we would obtain a contradiction to the maximality assumption about ΩM , and thus the claim holds.

(2): Let ω be in Ω . Since $\Omega M = \Phi = \Omega'M$, there is ω' in Ω' such that $\omega \in \omega'M$. Let ω_1 in Ω be such that $\omega' \in \omega_1 M$. Then $\omega \in \omega'M \subset \omega_1 M$, and hence $\omega = \omega_1$ by the independence of Ω . The desired bijection is then given by $\omega \mapsto \omega'$. \square

According to Proposition 2, any two bases of an incarnation have the same number of elements. We define the **dimension** of an incarnation to be the cardinality of its bases. For example a transitive group incarnation has dimension 1. In fact for a transitive group incarnation any single measurement forms a basis. More generally, the dimension of a group incarnation (Φ, M) equals the cardinality of Φ/M . In this case $\Omega \subset \Phi$ is a basis if and only if, for every block Ψ in Φ/M , the intersection $\Omega \cap \Psi$ has only one element. Since being a basis depends only on the monoid $\langle M \rangle$, the dimension of a group-like incarnation (Φ, M) equals also the cardinality of Φ/M , and similarly a subset $\Omega \subset \Phi$ is a basis if and only if, for every block Ψ in the partition Φ/M , the intersection $\Omega \cap \Psi$ has only one element.

The dimension of a transitive monoid incarnation can be bigger than 1. For example, let $X = \{x_1, x_2, x_3\}$ and consider functions $\phi_1, \phi_2, \phi_3: X \rightarrow \mathbf{R}$ and $g_1, g_2, g_3: X \rightarrow X$ defined as follows:

$$\begin{array}{l|l|l|l|l|l|l} \phi_1(x_1) = 2 & \phi_2(x_1) = 2 & \phi_3(x_1) = 1 & g_1(x_1) = x_2 & g_2(x_1) = x_2 & g_3(x_1) = x_1 \\ \phi_1(x_2) = 2 & \phi_2(x_2) = 2 & \phi_3(x_2) = 2 & g_1(x_2) = x_2 & g_2(x_2) = x_2 & g_3(x_2) = x_2 \\ \phi_1(x_3) = 3 & \phi_2(x_3) = 2 & \phi_3(x_3) = 2 & g_1(x_3) = x_3 & g_2(x_3) = x_2 & g_3(x_3) = x_2 \end{array}$$

The compositions $g_i g_j$ and $\phi_i g_j$ are described by the following tables:

	g_1	g_2	g_3		g_1	g_2	g_3
g_1	g_1	g_2	g_2	ϕ_1	ϕ_1	ϕ_2	ϕ_2
g_2	g_2	g_2	g_2	ϕ_2	ϕ_2	ϕ_2	ϕ_2
g_3	g_2	g_2	g_3	ϕ_3	ϕ_2	ϕ_2	ϕ_3

Thus the functions g_1 , g_2 , and g_3 are $\Phi := \{\phi_1, \phi_2, \phi_3\}$ -operations. Furthermore the subset $M := \{\text{id}, g_1, g_2, g_3\} \subset \text{End}_\Phi(X)$ is a submonoid. The incarnation

(Φ, M) is a transitive monoid incarnation. Since the set $\{\phi_1, \phi_3\}$ is independent and generates (Φ, M) , it is a basis. Thus (Φ, M) is an example of a transitive monoid incarnation of dimension 2.

5. Nirvana

To compare incarnations of various data sets we are going to use SEOs (set equivariant operators). A **SEO** from an incarnation (Φ, M) to an incarnation (Ψ, N) , denoted as $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$, is a pair of functions $(\alpha: \Phi \rightarrow \Psi, T: M \rightarrow N)$ for which the following diagram commutes:

$$\begin{array}{ccccc} \Phi \times M & \hookrightarrow & \Phi \times \text{End}_\Phi(X) & \xrightarrow{\text{action}} & \Phi \\ \alpha \times T \downarrow & & & & \downarrow \alpha \\ \Psi \times N & \hookrightarrow & \Psi \times \text{End}_\Psi(Y) & \xrightarrow{\text{action}} & \Psi \end{array}$$

Explicitly, for ϕ in Φ and g in M , it holds $\alpha(\phi g) = \alpha(\phi)T(g)$. This implies that for ϕ in Φ and every sequence of elements g_1, \dots, g_k in M , it holds:

$$\alpha(\phi g_1 \cdots g_k) = \alpha(\phi)T(g_1) \cdots T(g_k).$$

Be however aware that in general there may not be a homomorphism $T: \langle M \rangle \rightarrow \langle N \rangle$ of monoids which extends $T: M \rightarrow N$ and makes the following diagram commutative:

$$\begin{array}{ccccccc} \Phi \times M & \hookrightarrow & \Phi \times \langle M \rangle & \hookrightarrow & \Phi \times \text{End}_\Phi(X) & \xrightarrow{\text{action}} & \Phi \\ \alpha \times T \downarrow & & \alpha \times T \downarrow & & & & \downarrow \alpha \\ \Psi \times N & \hookrightarrow & \Psi \times \langle N \rangle & \hookrightarrow & \Psi \times \text{End}_\Psi(Y) & \xrightarrow{\text{action}} & \Psi \end{array}$$

A SEO between monoid incarnations $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is called a MEO (monoid equivariant operators) if $T: M \rightarrow N$ is a monoid homomorphism. A MEO between group incarnations is also called a GEO (group equivariant operators).

Let $(\alpha_0, T_0): (\Phi_0, M_0) \rightarrow (\Phi_1, M_1)$ and $(\alpha_1, T_1): (\Phi_1, M_1) \rightarrow (\Phi_2, M_2)$ be SEOs. Then the compositions $(\alpha_1 \alpha_0, T_1 T_0)$ form a SEO. Furthermore the pair $(\text{id}_\Phi, \text{id}_M): (\Phi, M) \rightarrow (\Phi, M)$ is also a SEO. The composition of SEOs is an associative operation and defines a category structure on the collection of data set incarnations with SEOs as morphisms. This category is called **Nirvana**.

A SEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is an isomorphism if and only if both of the functions α and T are bijections. Isomorphisms preserve independence and being a basis:

Proposition 3. *If $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is an isomorphism, then a subset $\Omega \subset \Phi$ is independent or a basis if and only if its image $\alpha(\Omega) \subset \Psi$ is independent or a basis.*

Proof. Assume α and T are bijections. This assumption imply that ϕ_1 belongs to $\phi_2 M$ if and only if $\alpha(\phi_1)$ belongs to $\alpha(\phi_2)N$. It follows that two elements in Φ are (in)dependant if and only if their images via α are (in)dependent in Ψ . By the same argument, $\Omega M = \Phi$ if and only $\alpha(\Omega)T(M) = \alpha(\Phi)$. \square

According to Proposition 3 two isomorphic incarnations have the same dimension.

The universal incarnations $(\Phi, \text{End}_\Phi(X))$ and $(\Phi, \text{Aut}_\Phi(X))$ are special in the category Nirvana. For any (Φ, M) , the pair $(\text{id}, i: M \hookrightarrow \text{End}_\Phi(X))$ defines a SEO $(\Phi, M) \rightarrow (\Phi, \text{End}_\Phi(X))$ called **canonical**. If (Φ, M) is a group incarnation, then the pair $(\text{id}, i: M \hookrightarrow \text{Aut}_\phi(X))$ defines a GEO $(\Phi, M) \rightarrow (\Phi, \text{Aut}_\phi(X))$ also called canonical.

The rest of this section is devoted to present three ways of constructing SEOs.

Change of units. Choose a function $f: \mathbf{R} \rightarrow \mathbf{R}$. For any incarnation (Φ, M) , consider the data set $f\Phi$ (see Section 2). If g is a Φ -operation, then it is also a $f\Phi$ -operation. Thus there is an inclusion $\text{End}_\Phi(X) \subset \text{End}_{f\Phi}(X)$, which is an equality if f is invertible, therefore we have an incarnation $(f\Phi, M)$. If (Φ, M) is a monoid or a group incarnation, then so is $(f\Phi, M)$. The pair $(f-, \text{id}_M): (\Phi, M) \rightarrow (f\Phi, M)$ is a SEO called the change of units along f .

Assume f is invertible. If $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is a SEO, then the pair of functions $((f-)\alpha(f^{-1}-), T)$ forms a SEO between $(f\Phi, M)$ and $(f\Psi, N)$. The assignment $(\alpha, T) \mapsto ((f-)\alpha(f^{-1}-), T)$ is a self functor $C(f)$ of Nirvana also called the change of units along f . It is an equivalence of categories. Indeed,

$$\begin{aligned} C(f)C(f^{-1})((\Phi, M)) &= C(f)(f^{-1}\Phi, M) = (\Phi, M) \\ C(f)C(f^{-1})((\alpha, T)) &= C(f)((f^{-1}-)\alpha(f-), T) \\ &= ((f-)(f^{-1}-)\alpha(f-)(f^{-1}-), T) = (\alpha, T). \end{aligned}$$

The same holds for $C(f^{-1})C(f)$, hence $C(f)$ is an equivalence of categories. The SEOs $(f-, \text{id}_M): (\Phi, M) \rightarrow (f\Phi, M)$, for all incarnations (Φ, M) , form a natural transformation between the identity functor on **Nirvana** and the change of units along f functor.

Domain change. Let (Φ, M) and (Ψ, N) be incarnations of data sets consisting of measurements on X and Y respectively. A SEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is called **geometric** if there is a function $f: Y \rightarrow X$, called a realization of (α, T) , making the following diagram commutative for every ϕ in Φ and g in M :

$$\begin{array}{ccc} Y & \xrightarrow{T(g)} & Y \\ f \downarrow & & \downarrow f \\ X & \xrightarrow{g} & X \end{array} \quad \begin{array}{c} \searrow \alpha(\phi) \\ \nearrow \phi \\ \mathbf{R} \end{array}$$

For example, let (Φ, M) be an incarnation of a data set consisting of measurements on X . Then the SEO $(\text{id}_\Phi, \text{id}_M): (\Phi, M) \rightarrow (\Phi, M)$ is geometric. The identity function $\text{id}_X: X \rightarrow X$ is one of its realizations.

Let $Y \subset X$ be M -invariant: $g(y)$ belongs to Y for all y in Y and g in M . Consider the data set $\Phi|_Y$ given by the domain change along the inclusion $Y \subset X$. The restriction of g to Y is a $\Phi|_Y$ -operation for every g in M . We use the symbol $T_Y: M \rightarrow \text{End}_{\Phi|_Y}(Y)$ to denote the function that maps g in M to the restriction of g to Y . The incarnation $(\Phi|_Y, T_Y(M))$ is called the **restriction** of (Φ, M) to the invariant subset Y . The pair $(\Phi \rightarrow \Phi|_Y, T_Y)$ forms a geometric SEO. The inclusion $i_Y: Y \hookrightarrow X$ is one of its realizations.

Let $f: Y \rightarrow X$ be a bijection. Consider the data set Φf . For any g in M , the function $f^{-1}gf: Y \rightarrow Y$ is a Φf -operation. Define $T: M \rightarrow \text{End}_{\Phi f}(Y)$ to map g in M to $f^{-1}gf$. The incarnation $(\Phi f, T(M))$ is called the domain change of (Φ, M) along f . The pair $(-f: \Phi \rightarrow \Phi f, T)$ forms a geometric SEO and $f: Y \rightarrow X$ is one of its realizations.

Extending from a basis. SEOs can be effectively constructed using bases.

Proposition 4. *Let (Φ, M) and (Ψ, N) be incarnations and Ω be a basis of (Φ, M) . Then two SEOs $(\alpha, T), (\alpha', T'): (\Phi, M) \rightarrow (\Psi, N)$ are equal if and only if $T = T'$ and $\alpha(\omega) = \alpha'(\omega)$ for any ω in Ω .*

Proof. The only non trivial thing to prove in the statement of the proposition is that $\alpha = \alpha'$ when their restrictions to Ω are equal. Assume $T = T'$ and $\alpha(\omega) = \alpha'(\omega)$ for any ω in Ω . Since Ω generates (Φ, M) , any element in Φ is of the form $\phi = \omega g_1 \cdots g_k$ for some ω in Ω and a sequence of elements g_1, \dots, g_k in M . The assumption and the fact that (α, T) and (α', T) are SEOs, imply:

$$\begin{aligned} \alpha(\phi) &= \alpha(\omega g_1 \cdots g_k) = \alpha(\omega) T(g_1) \cdots T(g_k) = \\ &= \alpha'(\omega) T(g_1) \cdots T(g_k) = \alpha'(\omega g_1 \cdots g_k) = \alpha'(\phi). \end{aligned}$$

Consequently $\alpha = \alpha'$. □

According to Proposition 4, a SEO is determined by what it does on a basis of the domain. This is analogous to a linear map between vector spaces being determined by its values on a basis. However unlike for linear maps, we cannot freely map elements of a basis of an incarnation to obtain a SEO. To obtain a SEO certain relations have to be preserved. Let (Φ, M) be an incarnation. A **relation** between measurements ϕ and ψ in Φ is by definition a pair of sequences $((g_1, \dots, g_k), (h_1, \dots, h_l))$ of elements in M for which the following equality holds: $\phi g_1 \cdots g_k = \psi h_1 \cdots h_l$.

Proposition 5. *Let (Φ, M) and (Ψ, N) be incarnations, Ω be a basis of (Φ, M) , and $\bar{\alpha}: \Omega \rightarrow \Psi$ and $T: M \rightarrow N$ be functions.*

1. *Assume that for every relation $((g_1, \dots, g_k), (h_1, \dots, h_l))$ between any two elements ω, ω' in Ω , the pair $((T(g_1), \dots, T(g_k)), (T(h_1), \dots, T(h_l)))$ is a relation between $\alpha(\omega)$ and $\alpha(\omega')$ in Ψ . Under this assumption, there is a unique SEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ for which the restriction of $\alpha: \Phi \rightarrow \Psi$ to Ω is $\bar{\alpha}$.*

2. Assume (Φ, M) and (Ψ, N) , are monoid incarnations, T is a monoid homomorphism, and if $\omega g = \omega' h$ for some ω, ω' in Ω and g, h in M , then $\alpha(\omega)T(g) = \alpha(\omega')T(h)$. Under these assumptions, there is a unique MEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ for which the restriction of $\alpha: \Phi \rightarrow \Psi$ to Ω is $\bar{\alpha}$.
3. Assume (Φ, M) and (Ψ, N) are group incarnations, T is a group homomorphism, and if $\omega = \omega g$, for some ω in Ω and g in M , then $\alpha(\omega) = \alpha(\omega)T(g)$. Under these assumptions, there is a unique GEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ for which the restriction of $\alpha: \Phi \rightarrow \Psi$ to Ω is $\bar{\alpha}$.

Proof. Since the proofs are analogous, we illustrate only how to show statement (2). For every ϕ in Φ , there exist (not necessarily unique) ω in Ω and g in M such that $\phi = \omega g$. The assumption implies that the expression $\alpha(\omega)T(g)$ depends on ϕ and not on the choices of ω and g for which $\phi = \omega g$. Thus by mapping ϕ in Φ to $\alpha(\omega)T(g)$ in Ψ , we obtain a well defined function also denoted by $\alpha: \Phi \rightarrow \Psi$. The pair (α, T) is the desired MEO. The uniqueness is a consequence of Proposition 4. \square

For example assume (Φ, M) is a transitive group incarnation and (Ψ, N) is a group incarnation. Choose an element ω in Φ . Recall that any such element is a basis of (Φ, M) . Fix a group homomorphism $T: M \rightarrow N$. Then any GEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is uniquely determined by the element $\alpha(\omega)$ in Ψ . Thus by choosing a basis element ω in Φ , we can identify the collection of GEOs of the form $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ with a subset of Ψ . To describe this subset explicitly, we apply Proposition 5.2. It states that there is a GEO $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ (necessarily unique) such that $\alpha(\omega) = \psi$ if and only if the following implication holds: if $\omega = \omega g$, then $\psi = \psi T(g)$. The collection $M_\omega := \{g \in M \mid \omega = \omega g\}$ is the isotropy subgroup of ω consisting of all the elements in M that fix ω . Thus GEOs of the form $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ can be identified with the subset of all the elements in Ψ whose isotropy group contains $T(M_\omega)$.

6. Decomposition

Let (Φ, M) be an incarnation of a data set Φ . Consider its quotient Φ/M , which is a partition of Φ , and the block incarnations (Ψ, M) for every block Ψ in Φ/M (see Section 4). Let X be the domain of Φ . Recall that the domain of the data set $\coprod_{\Psi \in \Phi/M} \Psi$ is given by the disjoint union $\coprod_{\Psi \in \Phi/M} X$, and that this data set consists of functions $\coprod_{\Psi \in \Phi/M} X \rightarrow \mathbf{R}$ whose restrictions to all but one summands X in $\coprod_{\Psi \in \Phi/M} X$ is the 0 function and the restriction to the remaining summand belongs to the corresponding block of the partition Φ/M . Define:

$$M' = \left\{ \coprod_{\Psi \in \Phi/M} g: \coprod_{\Psi \in \Phi/M} X \rightarrow \coprod_{\Psi \in \Phi/M} X \quad \mid \quad g \in M \right\}.$$

Then $M' \subset \text{End}_{\coprod_{\Psi \in \Phi/M} \Psi}(\coprod_{\Psi \in \Phi/M} X)$. We call $(\coprod_{\Psi \in \Phi/M} \Psi, M')$ the diagonal incarnation. Define $T: M \rightarrow M'$ to map $g: X \rightarrow X$ in M to $\coprod_{\Psi \in \Phi/M} g$ in M' . Define $\alpha: \Phi \rightarrow \coprod_{\Psi \in \Phi/M} \Psi$ to map ϕ to the function $\coprod_{\Psi \in \Phi/M} X \rightarrow \mathbf{R}$ whose restriction to the summand X corresponding to the block $[\phi]$ is ϕ and that maps all other summands to 0. Note that both of the functions α and T are bijections. Furthermore they form a SEO between (Φ, M) and $(\coprod_{\Psi \in \Phi/M} \Psi, M')$.

Proposition 6. *The SEO $(\alpha, T): (\Phi, M) \rightarrow (\coprod_{\Psi \in \Phi/M} \Psi, M')$ is an isomorphism.*

7. Grothendieck graphs

In this section we explain a convenient data structure to encode incarnations of data sets.

A **Grothendieck graph** is a triple (V, M, E) consisting of a finite set V whose elements are called vertices, a finite set M whose elements are called colors or operations, and a subset $E \subset V \times M \times V$ whose elements are called edges, such that, for every vertex v in V , the following composition is a bijection:

$$(\{v\} \times M \times V) \cap E \hookrightarrow E \hookrightarrow V \times M \times V \xrightarrow{\text{pr}_M} M.$$

This condition assures that, for every v in V and g in M , there is a unique element in V , denoted by vg , such that (v, g, vg) is an edge in E . For example let (Φ, M) be an incarnation of a data set Φ . Define:

$$E_{\Phi, M} := \{(\phi, g, \psi) \in \Phi \times M \times \Phi \mid \phi g = \psi\}.$$

Then the triple $(\Phi, M, E_{\Phi, M})$ is a Grothendieck graph. We think about this graph as a convenient data structure representing the incarnation (Φ, M) .

Grothendieck graphs are also convenient to represent SEOs. Define a **morphism between Grothendieck graphs** (V, M, E) and (W, N, F) to be a pair of functions $\alpha: V \rightarrow W$ and $T: M \rightarrow N$ such that, if (v, g, w) belongs to E , then $(\alpha(v), T(g), \alpha(w))$ belongs to F . Such a morphism is denoted as $(\alpha, T): (V, M, E) \rightarrow (W, N, F)$. Componentwise composition defines a category structure on the collection of Grothendieck graphs and we use the symbol GGraph to denote this category. If $(\alpha, T): (\Phi, M) \rightarrow (\Psi, N)$ is a SEO, then $(\alpha, T): (\Phi, M, E_{\Phi, M}) \rightarrow (\Psi, N, E_{\Psi, N})$ is a morphism between the associated Grothendieck graphs. By assigning to a SEO (α, T) the graph morphism given by the same pair (α, T) , we obtain a fully faithful functor from the category Nirvana to GGraph .

Grothendieck graphs can also be used to encode pseudometric information on incarnations. A pseudometric on a Grothendieck graph (V, M, E) is a pseudometric d on V such that $d(v, w) \geq d(vg, wg)$ for all v and w in V , and g in M . For example, the pseudometric $\|\phi - \psi\|_\infty$ on Φ is a pseudometric on the graph $(\Phi, M, E_{\Phi, M})$.

A Grothendieck graph (V, M, E) is said to be compatible with a monoid structure on M if $(v, 1, v)$ is in E , and whenever (v_0, g_0, v_1) and (v_1, g_1, v_2) belong to E , then so does $(v_0, g_1 g_0, v_2)$. In this case the composition operation given by the association $(v_0, g_0, v_1)(v_1, g_1, v_2) \mapsto (v_0, g_1 g_0, v_2)$ defines a category structure, denoted by $\text{Gr}_M V$, with V as the set of objects and E as the set of morphisms. This category is a familiar Grothendieck construction [5, 12]. For example, the Grothendieck graph associated with a monoid incarnation (Φ, M) is compatible with the monoid structure on M . We think about $\text{Gr}_M \Phi$ as an additional structure on the data set Φ : objects are the measurements in Φ , morphisms are triples $(\phi, g, \phi g)$, where ϕ is in Φ , g is in M , and the composition of $(\phi, g, \phi g)$ and $(\phi g, h, \phi gh)$ is given by $(\phi, gh, \phi gh)$.

A **contravariant functor** indexed by a Grothendieck graph (V, M, E) with values in a category \mathcal{C} , denoted as $P: (V, M, E) \rightarrow \mathcal{C}$, is by definition a sequence of objects $\{P(v) \mid v \in V\}$ and a sequence of morphisms $\{P(v_0, g, v_1): P(v_1) \rightarrow P(v_0) \mid (v_0, g, v_1) \in E\}$ in \mathcal{C} subject to: if (v_0, g_0, v_1) , (v_1, g_1, v_2) , and (v_0, h, v_2) are edges in E , then $P(v_2, h, v_0) = P(v_2, g_1, v_1)P(v_1, g_0, v_0)$. If (V, M, E) is compatible with a monoid structure on M , then a contravariant functor indexed by (V, M, E) is simply a contravariant functor indexed by the category $\text{Gr}_M V$.

Let (Φ, M) be an incarnation of a data set Φ consisting of measurements on X , and $(\Phi, M, E_{\Phi, M})$ be the associated Grothendieck graph. For every g in M , the function $-g: \Phi \rightarrow \Phi$, mapping ϕ to ϕg , is geometric and realized by $g: X \rightarrow X$ (see Section 3). Persistent homology leads therefore to the following collections of objects and morphisms in $\text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ as explained in Section 3:

$$\begin{aligned} & \{PH_d^\Phi(\phi) \mid \phi \in \Phi\}, \\ & \{PH_d^{-g}(\phi): PH_d^\Phi(\phi g) \rightarrow PH_d^\Phi(\phi) \mid (\phi, g, \phi g) \in E_{\Phi, M}\}. \end{aligned}$$

These sequences form a functor $PH_d^\Phi: (\Phi, M, E_{\Phi, M}) \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect})$ also referred to as the persistent homology functor of the incarnation (Φ, M) .

Let $(\alpha, T): (W, N, F) \rightarrow (V, M, E)$ be a morphism and $P: (V, M, E) \rightarrow \mathcal{C}$ be a functor. The following sequences of objects and morphisms in \mathcal{C} form a contravariant functor denoted by $P(\alpha, T): (W, N, F) \rightarrow \mathcal{C}$ and called the **composition** of (α, T) with P :

$$\begin{aligned} & \{P(\alpha(v)) \mid v \in V\}, \\ & \{P(w_0, g, w_1): P(\alpha(w_1)) \rightarrow P(\alpha(w_0)) \mid (w_0, g, w_1) \in F\}. \end{aligned}$$

For example, let $(\text{id}_\Phi, i): (\Phi, M) \rightarrow (\Phi, \text{End}_\Phi(X))$ be the canonical SEO (see Section 5). Consider the induced morphism of the associated Grothendieck graphs:

$$(\text{id}_\Phi, i_M): (\Phi, M, E_{\Phi, M}) \rightarrow (\Phi, \text{End}_\Phi(X), E_{\Phi, \text{End}_\Phi(X)}).$$

Consider also the persistent homology of the universal incarnation:

$$PH_d^\Phi: (\Phi, \text{End}_\Phi(X), E_{\Phi, \text{End}_\Phi(X)}) \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}).$$

The composition of these two functors coincides with the persistent homology of the incarnation (Φ, M) :

$$PH_d^\Phi : (\Phi, M, E_{\Phi, M}) \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}).$$

In this way we obtain a commutative diagram:

$$\begin{array}{ccc} & (\Phi, \text{End}_\Phi(X), E_{\Phi, \text{End}_\Phi(X)}) & \\ \text{(id}_\Phi, i_M) \nearrow & & \searrow PH_d^\Phi \\ (\Phi, M, E_{\Phi, M}) & \xrightarrow{PH_d^\Phi} & \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}) \end{array}$$

Such a commutativity does not hold for arbitrary SEOs. Consider a SEO $(\alpha, T) : (\Phi, M) \rightarrow (\Psi, N)$. We can form two functors indexed by the graph $(\Phi, M, E_{\Phi, M})$:

$$\begin{array}{ccc} & PH_d^\Phi & \rightarrow \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}) \\ & \nearrow & \\ (\Phi, M, E_{\Phi, M}) & & \\ & \searrow \alpha & \\ & (\Psi, N, E_{\Psi, N}) & \xrightarrow{PH_d^\Psi} \text{Tame}([0, \infty) \times \mathbf{R}, \text{Vect}) \end{array}$$


These functors rarely coincide. However, in the case (α, T) is geometric, the morphisms $PH_d^\alpha(\phi) : PH_d^\Psi(\alpha(\phi)) \rightarrow PH_d^\Phi(\phi)$ (see Section 3), for all ϕ in Φ , form a natural transformation.

8. Conclusions

In the following figure we give a graphical representation of some of the concepts introduced in this article. Data sets can be equipped with three structures: a pseudometric, an incarnation describing an action, and a Grothendieck graph. We imagine Nirvana as the landscape of all possible incarnations of data sets, represented by the shaded region in the following figure. Each point in Nirvana has a lot of internal structure allowing the extraction of persistent homology. In this landscape the black arrows represent geometric SEOs and the grey ones non-geometric SEOs. Recall that geometric SEOs enable us to compare relevant persistent homology. Non-geometric SEOs contain complementary information.

Acknowledgements

The research carried out by N.Q. was partially supported by GNSAGA-INdAM (Italy). A.D. has been supported by the SmartData@PoliTO center on Big Data and Data Science and by the Italian MIUR Award “Dipartimento di Eccellenza 2018-202” - CUP: E11G18000350001. The work of W.C. and F.T. was partially supported by the Wallenberg AI, Autonomous System and Software Program (WASP) funded by Knut and Alice Wallenberg Foundation. The work of W.C. was also in part funded by VR and Göran Gustafsson foundation.



References

- [1] Mattia G. Bergomi, Patrizio Frosini, Daniela Giorgi, and Nicola Quercioli. Towards a topological-geometrical theory of group equivariant non-expansive operators for data analysis and machine learning. *CoRR*, abs/1812.11832, 2018. URL: <http://arxiv.org/abs/1812.11832>, arXiv:1812.11832.
- [2] Mattia G. Bergomi, Patrizio Frosini, Daniela Giorgi, and Nicola Quercioli. Towards a topological-geometrical theory of group equivariant non-expansive operators for data analysis and machine learning. *Nature Machine Intelligence*, 1(9):423–433, 2019. doi:10.1038/s42256-019-0087-3.
- [3] Gunnar Carlsson and Afra Zomorodian. The theory of multidimensional persistence. *Discrete Comput. Geom.*, 42(1):71–93, 2009. URL: <https://doi.org/10.1007/s00454-009-9176-0>, doi:10.1007/s00454-009-9176-0.

- [4] David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Extending persistence using Poincaré and Lefschetz duality. *Found. Comput. Math.*, 9(1):79–103, 2009. URL: <https://doi.org/10.1007/s10208-008-9027-z>, doi:10.1007/s10208-008-9027-z.
- [5] William G. Dwyer and Hans-Werner Henn. *Homotopy theoretic methods in group cohomology*. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser Verlag, Basel, 2001. URL: <https://doi.org/10.1007/978-3-0348-8356-6>, doi:10.1007/978-3-0348-8356-6.
- [6] Herbert Edelsbrunner and John Harer. Persistent homology—a survey. In *Surveys on discrete and computational geometry*, volume 453 of *Contemp. Math.*, pages 257–282. Amer. Math. Soc., Providence, RI, 2008. URL: <https://doi.org/10.1090/conm/453/08802>, doi:10.1090/conm/453/08802.
- [7] Chad Giusti, Eva Pastalkova, Carina Curto, and Vladimir Itskov. Clique topology reveals intrinsic geometric structure in neural correlations. *Proceedings of the National Academy of Sciences*, 112(44):13455–13460, 2015. URL: <https://www.pnas.org/content/112/44/13455>, arXiv:<https://www.pnas.org/content/112/44/13455.full.pdf>, doi:10.1073/pnas.1506407112.
- [8] Alessandro De Gregorio, Ulderico Fugacci, Facundo Memoli, and Francesco Vaccarino. On the notion of weak isometry for finite metric spaces, 2020. arXiv:2005.03109.
- [9] Michael Lesnick. The theory of the interleaving distance on multidimensional persistence modules. *Found. Comput. Math.*, 15(3):613–650, 2015. URL: <https://doi.org/10.1007/s10208-015-9255-y>, doi:10.1007/s10208-015-9255-y.
- [10] Steve Y. Oudot. *Persistence theory: from quiver representations to data analysis*, volume 209 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015. URL: <https://doi.org/10.1090/surv/209>, doi:10.1090/surv/209.
- [11] Martina Scalamiero, Wojciech Chachólski, Anders Lundman, Ryan Ramanujam, and Sebastian Öberg. Multidimensional persistence and noise. *Foundations of Computational Mathematics*, 17(6):1367–1406, 2017. doi:10.1007/s10208-016-9323-y.
- [12] R. W. Thomason. Homotopy colimits in the category of small categories. *Math. Proc. Cambridge Philos. Soc.*, 85(1):91–109, 1979. URL: <https://doi.org/10.1017/S0305004100055535>, doi:10.1017/S0305004100055535.