

SEMI-BIPRODUCTS IN MONOIDS

N. MARTINS-FERREIRA

ABSTRACT. It is shown that the category of semi-biproducts in monoids is equivalent to a category of pseudo-actions. A semi-biproduct in monoids is at the same time a generalization of a semi-direct product in groups and a biproduct in commutative monoids. Every Schreier extension of monoids can be seen as an instance of a semi-biproduct; namely a semi-biproduct whose associated pseudo-action has a trivial correction system. A correction system is a new ingredient that must be inserted in order to obtain a pseudo-action out of a pre-action and a factor system. In groups, every correction system is trivial. Hence, semi-biproducts there are the same as semi-direct products with a factor system, which are nothing but group extensions. An attempt to establish a general context in which to define semi-biproducts is made. As a result, a new structure of map-transformations is obtained from a category with a 2-cell structure. Examples and first basic properties are briefly explored.

February 17, 2020

1. INTRODUCTION

The purpose of this paper is to introduce the notion of a semi-biproduct and to investigate it in the category of monoids.

From a categorical point of view, a biproduct is simultaneously a product and a co-product (see [9], p.194). Furthermore, when there is a coincidence between products and co-products in a category, each hom-set carries the structure of an abelian group and parallel morphisms can be added. This means that a biproduct can be defined as a diagram of the form

$$X \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} A \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} B \quad (1)$$

such that the following conditions hold:

$$ps = 1_B \quad , \quad pk = 0_{X,B} \quad (2)$$

$$qs = 0_{B,X} \quad , \quad qk = 1_X \quad (3)$$

$$kq + sp = 1_A. \quad (4)$$

It is clear that this definition makes sense even when each hom-set is a commutative monoid. And that seems to be the most general case in which it makes sense. But we may ask the following question: how

to generalize the definition so that it can be applied in the category of groups, where semi-direct products are expected to appear?

This work answers the question and it goes beyond groups. It provides a generalization to the so called *Schreier extensions* in monoids.

A Schreier extension of monoids (see [12]) can be seen as a sequence of monoid homomorphisms

$$X \xrightarrow{k} A \xrightarrow{p} B,$$

in which $k = \ker(p)$, with the property that there exist two set-theoretical functions, $s: B \rightarrow A$ and $q: A \rightarrow X$, satisfying conditions (7)–(9), below. When condition (9) is dropped, then the remaining ones can be reorganized so that they become precisely the conditions (2)–(4), except that now q and s are not homomorphisms but simply zero-preserving maps.

Thus, the notion of semi-biproduct in monoids arrives naturally as a diagram of the form (1) where p and k are monoid homomorphisms while q and s are zero-preserving maps, satisfying conditions (2)–(4).

Let us see in more detail the long path that has given rise to this simple definition.

In **Grp**, the category of groups and group homomorphisms, every split epimorphism

$$A \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} B, \quad ps = 1_B$$

induces an exact sequence

$$X \xrightarrow{k} A \xrightarrow{p} B,$$

with $X = \text{Ker}(p)$. If fixing X in additive notation (but not necessarily abelian) and B in multiplicative notation, then, up to isomorphism, the group A is recovered as a semi-direct product of the form $X \rtimes_{\varphi} B$, with neutral element $(0, 1) \in X \times B$, and group operation

$$(x, b) + (x', b') = (x + \varphi_b(x'), bb'). \quad (5)$$

Recall that (see e.g. [10] and its references to previous work)

$$\varphi_b: X \rightarrow X$$

is defined, for every $b \in B$, as the map $\varphi_b(x) = q(s(b) + k(x))$, with $q(a) = a - sp(a) \in X$, for every $a \in A$. Moreover, when p is a surjective homomorphism, but not necessarily a split epimorphism, it is still possible to recover the group A as a semi-direct product with a factor system, $X \rtimes_{\varphi, \gamma} B$. A factor system, $\gamma: B \times B \rightarrow X$, is any map $\gamma(b, b') = q(s(b) + s(b')) = s(b) + s(b') - s(bb')$, for some chosen set theoretical section $s: B \rightarrow A$ of p , i.e., such that $ps = 1_B$. In this case, the group operation in $X \rtimes_{\varphi, \gamma} B$ becomes

$$(x, b) + (x', b') = (x + \varphi_b(x') + \gamma(b, b'), bb'). \quad (6)$$

When A is abelian, the action $\varphi_b(x) = s(b) + k(x) - s(b) \in X$ is trivial and so, the surjective homomorphisms with abelian domain are nothing but (symmetric) factor systems.

The case of monoids is different (see [1]). A surjective monoid homomorphism $p: A \rightarrow B$, in general, cannot be presented as a sequence

$$X \xrightarrow{(1,0)} X \rtimes_{\varphi,\gamma} B \xrightarrow{\pi_B} B$$

with $X = \text{Ker}(p)$ and such that the underlying set of A is in bijection with the cartesian product $X \times B$. Thus, in monoids, the transport of the structure from A to $X \times B$ is not always possible. So, there is no hope of having an isomorphism $A \cong X \rtimes_{\varphi,\gamma} B$. As we will see, the best that we can hope for is to have A (i.e. the underlying set of A) as a subset of the cartesian product. To do that we will need to introduce a new ingredient, a correction system, $\rho(x, b) = x^b$, so that we have

$$A \cong (X \rtimes_{\varphi,\rho,\gamma} B) \subseteq X \times B.$$

And yet, not every surjective monoid homomorphism exhibits that property. Take, for example, the usual addition of natural numbers,

$$\mathbb{N} \times \mathbb{N} \xrightarrow{+} \mathbb{N},$$

whose kernel is the trivial monoid.

We may ask: what kind of monoid homomorphisms has the same behaviour as in groups? The answer turns out to be the so called Schreier extensions (see [6, 12] and references therein, see also [1]).

A Schreier extension $p: A \rightarrow B$ is nothing but a monoid homomorphism with the property that there exists a set theoretical section $s: B \rightarrow A$ for which the function $X \times B \rightarrow A$, $(x, b) \mapsto k(x) + s(b)$ is a bijection of the underlying sets, with $k = \text{ker}(p)$.

Equivalently, a Schreier extension can be seen as a monoid homomorphism $p: A \rightarrow B$, together with set theoretical maps $s: B \rightarrow A$ and $q: A \rightarrow X$, with $X = \text{Ker}(p)$ and $k = \text{ker}(p): X \rightarrow A$, such that

$$ps(b) = b \quad , \quad b \in B \tag{7}$$

$$kq(a) + sp(a) = a \quad , \quad a \in A \tag{8}$$

$$q(k(x) + s(b)) = x \quad , \quad x \in X, b \in B. \tag{9}$$

The key feature of this work is the observation that, at the expense of having to insert a correction system, the condition (9) can be discarded. The correction system controls the lack of condition (9) and will be denoted by x^b . It is defined as $x^b = q(k(x) + s(b))$, for every $x \in X$ and $b \in B$. While dropping condition (9), we loose the bijection between the underlying set of A and the cartesian product of sets $X \times B$. However, if adding two simple conditions, namely $q(k(x)) = x$ and $q(s(b)) = 0$, then, A is isomorphic to a subset of the cartesian product $X \times B$. The

subset consists of those pairs in $X \times B$ that are of the form (x^b, b) for some $x \in X$ and $b \in B$ (see Theorem 1). The monoid operation is

$$(x, b) + (x', b') = ((x + b \cdot x' + (b \times b'))^{bb'}, bb'). \quad (10)$$

The correction system x^b has to satisfy some conditions (Definition 1). In the case of groups it becomes trivial, that is $x^b = x$. To see it, simply take $b' = b^{-1}$ in equation (36). The notion of a correction system is inspired by the work of Leech and Wells on extending groups by monoids ([8, 14]). The fact that the correction system $\rho(x, b) = x^b$ is invisible in groups, hides the true nature of pseudo-actions as a combination of structures φ, ρ, γ (Definition 1) satisfying one single major condition (31). Only the particular traces of (31) are familiar in groups. This is better explained in Remark 1. A Schreier extension in monoids is precisely a semi-biproduct with a trivial correction system, which explains the similar behaviour between Schreier extensions in monoids and all extensions in groups. Examples of semi-biproducts in monoids which are not Schreier extensions, that is, with a non-trivial correction system, are presented (Section 3).

From a structural point of view, we are considering sequences of monoid homomorphisms

$$X \xrightarrow{k} A \xrightarrow{p} B \quad (11)$$

together with set theoretical maps, preserving the neutral element, but not necessarily preserving the monoid operation,

$$X \xleftarrow{q} A \xleftarrow{s} B \quad (12)$$

and satisfying the conditions (2)–(4). When the maps q and s are monoid homomorphisms then this is exactly the definition of a bi-product. It is then natural to call semi-biproduct when q and s are just zero-preserving maps.

As expected, a semi-biproduct becomes a product as soon as the zero-preserving maps q and s are both morphisms. In that case it is a coproduct as well. Particular instances when just one of q or s is a morphism are worthwhile studying, but we shall no longer expect a product or a co-product to appear. In general it does not make sense to speak about the semi-biproduct of two objects, unless some kind of pseudo-action is specified. This is not so surprising since in groups, semi-biproducts are the same as semi-direct products with a factor system. Connections with homology and cohomology are expected via several notions such as abstract kernel, obstruction, etc (see for example [12] and references there), and it is clear that the notion of pseudo-action considered here (Definition 1) can be used as a tool and clarify some classical interpretations in group-cohomology.

This paper is organized as follows. First we concentrate our attention on the concrete case of monoids, introducing a new concept of pseudo-action and establishing a correspondence between pseudo-actions and semi-biproducts in monoids (Theorem 3). In Section 3 we illustrate the concept with Schreier extensions and other types of more general extensions.

At the end we address the natural question of where to define semi-biproducts. We arrive at an abstract structure of map-transformations. Its origin comes from a convenient structure of 2-cells in monoids. This appears to be a convenient setting to work in a wider context while keeping the intuition from monoids. The observation that semi-biproducts can be defined in such cases is made precise. We briefly discuss some properties of map-transformation structures, such as the property of some monomorphisms being recognizers (Theorem 4). A list of examples and general procedures on how to generate structures of map-transformations is provided.

2. MONOIDS, SEMI-BIPRODUCTS AND PSEUDO ACTIONS

In this section we define semi-biproducts in monoids, introduce a new notion of pseudo-action, and prove that pseudo-actions are equivalent to semi-biproducts. Several works have been done in this direction but the emphasis has always been on the side of (Schreier) extensions and their classification (e.g. [2, 5, 6, 12]). Here we consider a semi-biproduct as a mathematical object rather than something which arises from an extension with appropriate choices for a section and a retraction. In a certain sense this work goes in the direction of [7]. There, the Schreier condition (9) is still being considered, but it shows that it is possible to work on the wider context of unitary magmas.

2.1. Semi-biproducts in monoids. At the end of this paper we establish a general setting in which semi-biproducts can be defined. When interpreted in the case of monoids (with zero-preserving maps), it gives a diagram of the shape

$$X \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} A \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} B \quad (13)$$

such that p, k , are monoid homomorphisms, while q and s are zero-preserving maps, verifying the conditions

$$ps = 1_B \quad (14)$$

$$qk = 1_X \quad (15)$$

$$kq + sp = 1_A \quad (16)$$

$$pk = 0_{X,B} \quad (17)$$

$$qs = 0_{B,X}. \quad (18)$$

It is possible to characterize all semi-biproducts in monoids with fixed X and B . We will work with additive notation on X and multiplicative notation on B , but neither is assumed to be commutative.

Theorem 1. *Let (X, A, B, p, k, q, s) be a semi-biproduct in the category of monoids with zero-preserving maps. If we put for every $b, b' \in B$ and $x \in X$,*

$$b \cdot x = q(s(b) + k(x)) \quad (19)$$

$$x^b = q(k(x) + s(b)) \quad (20)$$

$$b \times b' = q(s(b) + s(b')) \quad (21)$$

then, for every $a, a' \in A$

$$a + a' = k(q(a) + p(a) \cdot q(a') + (p(a) \times p(a'))) + sp(a + a') \quad (22)$$

and

$$a + a' = k(u^v) + s(v) \quad (23)$$

with $u = q(a) + p(a) \cdot q(a') + (p(a) \times p(a')) \in X$ and $v = p(a + a') \in B$.

Moreover, the map $\beta: A \rightarrow X \times B$ with $\beta(a) = (q(a), p(a))$ is always injective. An element $(y, b) \in X \times B$ is in the image of β if and only if $y = x^b$ for some $x \in X$.

This means that A is in bijection with the set $\{(x^b, b) \mid (x, b) \in X \times B\}$ with the inverse map for β being $\alpha(x^b, b) = k(x^b) + s(b) = k(x) + s(b)$.

Furthermore, for every $a \in A$, $q(a) = q(a)^{p(a)}$.

Proof. We observe:

$$\begin{aligned} a + a' &= kqa + (spa + kqa') + spa' && (kq + sp = 1) \\ &= kqa + kq(spa + kqa') + spa + spa' && (ps = 1, pk = 0) \\ &= kqa + kq(spa + kqa') + kq(spa + spa') + s(pa + pa') \\ &= k(qa + q(spa + kqa') + q(spa + spa')) + sp(a + a') \\ &= k(q(a) + p(a) \cdot q(a') + (p(a) \times p(a'))) + sp(a + a') \end{aligned}$$

We observe further that for every $a \in A$, $x \in X$, $b \in B$, if $a = k(x) + s(b)$ then $a = k(x^b) + s(b)$, which proves condition (23). This follows from the fact that for every $a \in A$, we have $q(a) = q(a)^{p(a)}$. Indeed, $q(a) = q(kq(a) + sp(a))$ follows directly from $a = kq(a) + sp(a)$.

If $\beta(a) = \beta(a')$ then $q(a) = q(a')$ and $p(a) = p(a')$, hence $a = a'$. This proves β is injective.

If $(x, b) \in X \times B$ is of the form $x = q(a)$, $b = p(a)$ for some $a \in A$, then $x = x^b$ because $q(a) = q(a)^{p(a)}$ as already shown. If $(y, b) \in X \times B$ is of the form $y = x^b$ for some $x \in X$, then there exists $a = k(x) + s(b) \in A$ such that $\beta(a) = (x, b)$. □

We will now show that there is an equivalence of categories between the category of semi-biproducts in monoids and a suitable category of pseudo-actions in monoids.

2.2. Pseudo-actions in monoids. Let X and B be two monoids. For convenience, let us again use additive notation for X and multiplicative notation for B . Note that neither one is assumed to be commutative.

Definition 1. A pseudo-action of B on X consists in three different components:

(1) a pre-action, that is, a map

$$\varphi: B \rightarrow X^X$$

associating to every element $b \in B$ the map $\varphi_b: X \rightarrow X$ which is denoted as $\varphi_b(x) = b \cdot x$ and satisfies the conditions

$$1 \cdot x = x, \quad \forall x \in X \quad (24)$$

$$b \cdot 0 = 0, \quad \forall b \in B \quad (25)$$

(2) a correction system, that is, a map

$$\rho: X \times B \rightarrow X; \quad (x, b) \mapsto x^b \quad (26)$$

such that for all $x \in X$ and $b \in B$

$$x^1 = x \quad (27)$$

$$0^b = 0 \quad (28)$$

(3) a factor system, that is, a map

$$\gamma: B \times B \rightarrow X; \quad (b, b') \mapsto b \times b' \quad (29)$$

such that for every $b \in B$

$$1 \times b = 0 = b \times 1 \quad (30)$$

The three components are related via the condition (31) which must hold for every $x, x', x'' \in X$ and $b, b', b'' \in B$.

$$\begin{aligned} & (x + b \cdot ((x' + b' \cdot x'' + (b' \times b''))^{b'b''}) + (b \times b'b''))^{bb'b''} = \\ & = ((x + b \cdot x' + (b \times b'))^{bb'} + bb' \cdot x'' + (bb' \times b''))^{bb'b''} \end{aligned} \quad (31)$$

The correction system, ρ , is used to correct the fact that in general $b \cdot (x + y)$ is not equal to $b \cdot x + b \cdot y$. Instead, we have the equalities

$$(b \cdot (x + y))^b = (b \cdot x + b \cdot y)^b \quad (32)$$

$$(x + y)^b = (x + y^b)^b \quad (33)$$

$$(x^b + b \cdot y)^b = (x + (b \cdot y)^b)^b \quad (34)$$

which are obtained (see items 4, 5, 6 below) as particular cases of (31). This correction is invisible in groups (see item 2 below) which perhaps explains the reason why it has never appeared explicitly as a structure before. Nevertheless, it has implicitly been used by Leech [8] and Wells [14] (see also [5]).

Remark 1. In (31) we may observe the following particular cases of interest:

(1) If taking x, x', x'' to be zero then we get

$$(b \cdot (b' \times b'')^{bb'} + (b \times b'b''))^{bb'b''} = ((b \times b')^{bb'} + (bb' \times b''))^{bb'b''} \quad (35)$$

which, if we ignore the correction system, it becomes the usual formula for a factor system in monoids (see e.g. [6] and references therein).

(2) If taking $x = x'' = 0$, $b = 1$ then we get (by putting again x in the place of x' , b in the place of b' and b' in the place of b'' for readability)

$$(x^b + b \times b')^{bb'} = (x + b \times b')^{bb'} \quad (36)$$

which explains how different the correction system is from being an action. In particular, if X is right cancellable and B is a group then the correction system is always trivial. Indeed, if we take $b' = b^{-1}$ we obtain $x^b + (b \times b^{-1}) = x + (b \times b^{-1})$ and hence, cancelling out $b \times b^{-1}$, we get $x^b = x$.

(3) If taking $x = x' = 0$ and $b'' = 1$ then we get

$$(b \cdot (b' \cdot x'')^{b'} + (b \times b'))^{bb'} = ((b \times b')^{bb'} + bb' \cdot x'')^{bb'} \quad (37)$$

which, in groups, becomes the familiar expression

$$b \cdot (b' \cdot x'') = (b \times b') + bb' \cdot x'' - (b \times b') \quad (38)$$

stating that the factor system $b \times b'$ measures, by conjugation, the distance between a pre-action φ and an ordinary action.

(4) If taking $b = b' = 1$ and $x'' = 0$ then we get

$$(x + x')^{b''} = (x + x^{b''})^{b''} \quad (39)$$

which is exactly the same as (33)

(5) If taking $b = b'' = 1$ and $x' = 0$ then we get

$$(x^{b'} + b' \cdot x'')^{b'} = (x + (b' \cdot x'')^{b'})^{b'} \quad (40)$$

which is exactly the same as (34).

(6) If taking $b' = b'' = 1$ and $x = 0$ then we get

$$(b \cdot (x' + x''))^b = ((b \cdot x')^b + b \cdot x'')^b \quad (41)$$

which, if combined with (33) and (34), gives (32).

2.3. The equivalence. It will be clear at the end of this subsection that there is an equivalence between pseudo-actions and semi-biproducts in monoids.

Theorem 2. *Let (X, A, B, p, k, q, s) be a semi-biproduct in the category of monoids with zero-preserving maps. The system with three components*

$$b \cdot x = q(s(b) + k(x)) \quad (42)$$

$$x^b = q(k(x) + s(b)) \quad (43)$$

$$b \times b' = q(s(b) + s(b')) \quad (44)$$

is a pseudo action from B into X .

Proof. The heart of the proof relies on the decomposition

$$a + a' = k(q(a) + p(a) \cdot q(a') + (p(a) \times p(a'))) + sp(a + a')$$

which holds for every $a, a' \in A$, as proved in Theorem 1.

To obtain (31), we consider $a + (a' + a'') = (a + a') + a''$ with $x = q(a), x' = q(a'), x'' = q(a'')$ and $b = p(a), b' = p(a'), b'' = p(a'')$. To prove $b \cdot 0 = 0$, $b \times 1 = 0 = 1 \times b$ and $0^b = 0$ we use the fact that $qs(b) = 0$ for all $b \in B$ and the other conditions are easily verified. \square

This means that if starting with a semi-biproduct (X, A, B, p, k, q, s) we obtain a pseudo-action (φ, ρ, γ) with $\varphi_b(x) = b \cdot x$, $\rho(x, b) = x^b$ and $\gamma(b, b') = b \times b'$ as defined in Theorem 2. We only need to make a straightforward calculation to see that given a pseudo-action we obtain a semi-biproduct in monoids. All the necessary ingredients have been given in Theorem 1. Let us make it more precise. Given a pseudo-action (φ, ρ, γ) of B on X , we construct a synthetic (see [8]) semi-biproduct in monoids

$$X \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} A \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} B \quad (45)$$

with $A = \{(x^b, b) \mid x \in X, b \in B\}$ a monoid with neutral element $(0, 1) \in X \times B$ and whose operation is the restriction to the operation defined on $X \times B$ as

$$(x, b) + (x', b') = ((x + b \cdot x' + (b \times b'))^{bb'}, bb') \quad (46)$$

which is well defined and associative on the subset A . However, in general, it fails to be associative on the set $X \times B$. The homomorphisms p and k , as well as the zero-preserving maps q and s , are the expected ones. The monoid A is in some sense a semi-direct product with a correction system and a factor system, so, it would be appropriate to write it as $X \rtimes_{\varphi, \rho, \gamma} B$.

Theorem 3. *There is an equivalence between the category of semi-biproducts in monoids and the category of pseudo-actions.*

Proof. Start with a semi-biproduct (X, A, B, p, k, q, s) and let (φ, ρ, γ) be its associated pseudo action. The synthetic semi-biproduct constructed with the pseudo-action is isomorphic to the original semi-biproduct as illustrated

$$\begin{array}{ccc} X & \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} & A & \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} & B \\ \parallel & & \alpha \updownarrow \beta & & \parallel \\ X & \begin{array}{c} \xleftarrow{\pi_1} \\ \xrightarrow{\langle 1, 0 \rangle} \end{array} & X \rtimes_{\varphi, \rho, \gamma} B & \begin{array}{c} \xrightarrow{\pi_2} \\ \xleftarrow{\langle 0, 1 \rangle} \end{array} & B \end{array} \quad (47)$$

with the morphisms α and β defined as in Theorem 1.

If we start with a pseudo-action (φ, ρ, γ) , build its associated synthetic semi-biproduct, and extract the pseudo-action associated to it, then a new pseudo-action is derived, say $(\varphi', \rho', \gamma')$. The derived pseudo-action is obtained as $\varphi'_b(x) = (b \cdot x)^b$, $\rho'(x, b)$ is the same as $\rho(x, b) = x^b$ and $\gamma'(b, b') = (b \times b')^{bb'}$. Nevertheless these two pseudo-actions are isomorphic because $(x^b)^b = x^b$, which is a consequence of equation (33). \square

3. EXAMPLES OF SEMI-BIPRODUCTS IN MONOIDS

Every Schreier split extension ([2]) of a monoid X by a monoid B is a semi-biproduct in monoids. In this case $b \times b' = 0$ and $x^b = x$ for all $x \in X$, $b, b' \in B$. Every Schreier extension of monoids is a semi-biproduct in which $x^b = x$. In fact, a semi-biproduct of monoids (X, A, B, p, k, q, s) is a Schreier extension if and only if $x^b = x$ for all $x \in X$ and $b \in B$.

Here is a general procedure for the construction of semi-biproducts which are not necessarily Schreier extensions.

Let X (written additively) and B (written multiplicatively) be two monoids and let us suppose the existence of a subset $R \subseteq X \times B$, considered as a binary relation, so that we write xRb instead of $(x, b) \in R$, together with two maps $u: B \rightarrow X$ and $q: R \rightarrow X$ satisfying the following conditions:

$$\text{for all } x \in X, xR1 \text{ and } q(xR1) = x \quad (48)$$

$$\text{for all } b \in B, u(b)Rb \text{ and } q(u(b)Rb) = 0 \quad (49)$$

$$\text{if } xRb, yRb \text{ and } q(xRb) = q(yRb) \text{ then } x = y. \quad (50)$$

For every monoid structure on the set R , for which $0R1$ is the neutral element and the projection map $xRb \mapsto b$ is a homomorphism, we put:

$$\begin{aligned} x \oplus x' &= q(xR1 + x'R1) \\ b \times b' &= q(u(b)Rb + u(b')Rb') \\ b \cdot x &= q(u(b)Rb + xR1) \\ x^b &= q(xR1 + u(b)Rb). \end{aligned}$$

If $x \oplus x' = x + x'$ for all $x, x' \in X$ (where $+$ is the monoid operation on X) and if $q(xRb)^b = q(xRb)$, whenever xRb , then we have a semi-biproduct

$$X \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} R \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} B \quad (51)$$

with $p(xRb) = b$, $k(x) = (xR1)$, $s(b) = (u(b)Rb)$.

The monoid operation on R is given by the formula

$$(xRb) + (x'Rb') = (x + b \cdot x' + b \times b')^{bb'} Rbb'. \quad (52)$$

Indeed, k is a homomorphism because \oplus is the monoid operation defined on X , p is a homomorphism by assumption, $pk = 0$, $ps = 1$,

$qk = 1$, $qs = 0$ by construction and we have, for every xRb ,

$$xRb = q(xRb)R1 + u(b)Rb \quad (53)$$

which means that $kq + sp = 1_R$. It is not difficult to see that (53) follows from the fact that $q(xRb)^b = q(xRb)$ together with the fact that the map $\langle q, p \rangle: R \rightarrow X \times B$ is injective (condition (50) above).

In particular, when we force $q(xRb) = x$ and $u(b) = 0$ then we are reduced to the analysis of monoid structures on R . In that case, the condition $q(xRb)^b = q(xRb)$ becomes $x^b = x$ whenever xRb , and if we are looking for semi-biproducts which are not Schreier extensions then we have to consider relations which are not in bijection with the set $X \times B$. Let us work a concrete example.

Take $X = \{0, s\}$ with $s + s = s$ and $B = \{1, t\}$ with $t^2 = t$. Clearly, $R = \{(0, 1), (s, 1), (0, t)\}$ is the only possible proper subset of $X \times B$ with $xR1$ and $0Rb$ for every $x \in X$ and $b \in B$. There are two possible solutions to turn R into a monoid with $0R1$ as neutral element and so that the map $p(xRb) = b$ is a homomorphism.

One possibility is to map $0R1 \mapsto 1$, $sR1 \mapsto -1$, $0Rt \mapsto 0$ and take the usual multiplication on the set $\{-1, 0, 1\}$. However, in this case we find that $s \oplus s = 0$ while $s + s = s$. So, it does not give a semi-biproduct.

Another possibility is to consider the chain semilattice structure on R as follows

$$\begin{array}{c|ccc} + & 0R1 & sR1 & 0Rt \\ \hline 0R1 & 0R1 & sR1 & 0Rt \\ sR1 & sR1 & sR1 & 0Rt \\ 0Rt & 0Rt & 0Rt & 0Rt \end{array}$$

In this case we have

$$\begin{array}{c|c|c|c|c|c|c|c} x & b & x' & b' & x \oplus x' & b \times b' & b \cdot x & x^b \\ \hline 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & t & s & 1 & s & 0 & 0 & 0 \\ s & 1 & 0 & t & s & 0 & s & s \\ s & t & s & t & s & 0 & 0 & 0 \end{array}$$

and since $x \oplus x' = x + x'$ and $x^b = x$ whenever xRb (note that $s^t = 0$ but $(s, t) \notin R$) we obtain a semi-biproduct of monoids

$$\{0, s\} \begin{array}{c} \xleftarrow{\pi_1} \\ \xrightarrow{\iota_1} \end{array} R \begin{array}{c} \xrightarrow{\pi_2} \\ \xleftarrow{\iota_2} \end{array} \{1, t\} \quad (54)$$

with $\pi_2(xRb) = b$, $\iota_2(b) = (0Rb)$, $\pi_1(xRb) = x$, $\iota_1(x) = (xR1)$. It cannot be a Schreier extension because R has three elements while $X \times B$ has four elements.

Let us now see a simple example which illustrates the case when we take R to be in bijection with $X \times B$. Put $X = B = \mathbb{N}$ the additive monoid of natural numbers and consider the order relation $R = \{(x, b) \in \mathbb{N}^2 \mid x \geq b\}$ together with the two maps $q(x \geq b) = x - b$ and $u(b) = b$. The usual component-wise addition on R verifies all

the conditions specified by the construction scheme outlined above and hence it gives rise to a semi-biproduct. Namely, $(\mathbb{N}, R, \mathbb{N}, p, k, q, s)$ with $p(x \geq b) = b$, $k(x) = (x \geq 0)$, $q(x \geq b) = x - b$ and $s(b) = (b \geq b)$. In this case q and s are both homomorphisms and hence $x^b = x$ for every $(x, y) \in \mathbb{N} \times \mathbb{N}$ so that we obtain an isomorphism

$$\begin{array}{ccccc} \mathbb{N} & \xleftarrow{q} & R & \xrightarrow{p} & \mathbb{N} \\ & & \downarrow k & & \downarrow s \\ \mathbb{N} & \xleftarrow{\pi_1} & \mathbb{N} \times \mathbb{N} & \xrightarrow{\pi_2} & \mathbb{N} \\ & & \langle 1, 0 \rangle & & \langle 0, 1 \rangle \end{array}$$

with $\beta(x \geq b) = (x - b, b)$ and $\alpha(x, b) = (x + b \geq b)$. See Theorem 3.

4. WHAT IS THE RIGHT CONTEXT IN WHICH TO DEFINE SEMI-BIPRODUCTS?

The question of choosing an appropriate setting in which to define semi-biproducts is a natural one. We will merely give some directions that we expect to follow in future research. It all starts with the observation that zero-preserving maps, an essential part on the definition of semi-biproduct (mainly because they are closed under component-wise addition), can be seen as a particular instance of a 2-cell structure in the category of monoids. The following subsections are an attempt to explain better what we have in mind.

4.1. The category of monoids with a 2-cell structure. It is well known that a monoid can be seen as a one object category. This means that two parallel monoid homomorphisms, $f, g: A \rightarrow B$, can be seen as two functors. Thus, a natural transformation, $\tau: f \Longrightarrow g$ becomes nothing but an element $t \in B$ for which $t + f(x) = g(x) + t$ for all $x \in A$. In this case we write $\tau = (f, t, g)$. Pushing it a little bit further we may say that a *transformation* from f to g is a zero-preserving map $t: A \rightarrow B$ such that $t(x) + f(x) = g(x) + t(x)$, for all $x \in A$. This gives rise to an example of a 2-cell structure in the sense of [11]. See also [3], where they are called *wiskered categories*. A whiskered category, or a category with a 2-cell structure, is not a 2-category because the middle interchange law is not present.

A 2-cell structure over a category \mathbf{A} is a system $(H, \text{dom}, \text{cod}, 0, +)$ in which $H: \mathbf{A}^{\text{op}} \times \mathbf{A} \rightarrow \mathbf{Set}$ is a bifunctor, $\text{dom}, \text{cod}: H \rightarrow \text{hom}_{\mathbf{A}}$, $0: \text{hom}_{\mathbf{A}} \rightarrow H$ and

$$+: H \times_{\text{hom}} H \rightarrow H$$

are natural transformations satisfying certain conditions (the details can be found in [11] but will not be needed here). We only need to observe that if defining, for every two monoids A and B , the set $H(A, B)$ as consisting in all the triples of the form $\tau = (f, t, g)$, with $f, g: A \rightarrow B$ monoid homomorphisms, and $t: A \rightarrow B$ a zero-preserving

map with the property $t(x) + f(x) = g(x) + t(x)$, for all $x \in A$, then it gives rise to a 2-cell structure on the category of monoids. Indeed, $\text{dom}(\tau) = f$, $\text{cod}(\tau) = g$, $0(f) = (f, 0_{A,B}, f)$ and if $\tau' = (g, t', h)$ then $\tau' + \tau = (f, t' + t, h)$.

We point out that this is not a natural 2-cell structure and hence it does not give rise to a horizontal composition as would be expected in a 2-category. Nevertheless, we do have a reasonable formula for the horizontal composition of 2-cells, namely

$$(f, t, g) \circ (f', t', g') = (ff', tt', gg') \quad (55)$$

for homomorphisms f, f', g, g' and zero-preserving maps t and t' with appropriate domains and codomains. In a small parenthesis we may add that for the middle interchange law to hold, it would be needed that $gt' + tf'$ is equal to $tg' + ft'$, which is not the case in general.

Note that in $\tau = (f, t, g)$, when f and g are the zero homomorphisms, then t becomes a zero-preserving map with no other conditions.

We conclude this subsection by remarking that if we denote by $\text{map}(A, B)$ the collection of all zero-preserving maps from a monoid A into a monoid B , then, it is derived from the bifunctor H , as defined above, with

$$\text{map}(A, B) = \{\tau \in H(A, B) \mid \text{dom}(\tau) = 0_{A,B} = \text{cod}(\tau)\}.$$

This suggests to attempt for an abstraction of $\text{map}(A, B)$. It is a bifunctor into the category of pointed sets (as soon as the category \mathbf{A} is pointed) and it factors through the category of monoids. Moreover, there is a natural inclusion $\varepsilon: \text{hom}_{\mathbf{A}} \rightarrow \text{map}$ which, in the current example of monoids, is defined as $\varepsilon(f) = (0_{A,B}, f, 0_{A,B})$ for every homomorphism $f: A \rightarrow B$. Its compatibility with (55), when restricted from H to map , is expressed as follows:

$$(0, t, 0) \circ \varepsilon(h) = (0, th, 0) = (0, t, 0)h, \quad (56)$$

$$\varepsilon(g) \circ (0, t, 0) = (0, gt, 0) = g(0, t, 0). \quad (57)$$

The formulas on the right are the whiskering compositions between morphisms and 2-cells. Compare these with (70) and (71) below.

4.2. A (pointed) category with a map-transformation structure. In order to speak of semi-biproducts we need to introduce a setting where addition of morphisms is possible. As follows from the case in monoids, the (componentwise) addition of two parallel morphisms fails, in general, to be a morphism and it is only a zero-preserving function. This fact, together with the explanation provided in the previous subsection, suggests to introduce the notion of a (pointed) category with an abstract structure of zero-preserving maps, that we decided to call map-transformations, for convenience of notation.

Let \mathbf{A} be a pointed category and let $\text{hom}_{\mathbf{A}}$ denote its hom-functor. A *map-transformation structure* on \mathbf{A} consists in a bifunctor

$$\text{map}: \mathbf{A}^{\text{op}} \times \mathbf{A} \rightarrow \mathbf{Set}_* \quad (58)$$

that factors through the category of monoids

$$\begin{array}{ccc} \mathbf{A}^{\text{op}} \times \mathbf{A} & \xrightarrow{\text{Map}} & \mathbf{Mon} \\ & \searrow \text{map} & \downarrow F \\ & & \mathbf{Set}_* \end{array} \quad (59)$$

together with a natural inclusion (of pointed sets)

$$\varepsilon: \text{hom}_{\mathbf{A}} \rightarrow \text{map} \quad (60)$$

and, for every three objects A, B, C in \mathbf{A} , an associative composition of *map-transformations*

$$\mu_{A,B,C}: \text{map}(B, C) \times \text{map}(A, B) \rightarrow \text{map}(A, C) \quad (61)$$

such that, for every $x \in \text{map}(A, B)$, and for every two morphisms $f: A' \rightarrow A$, $g: B \rightarrow B'$,

$$\mu(\varepsilon(f), x) = \text{map}(f, 1)(x) \quad (62)$$

and

$$\mu(x, \varepsilon(g)) = \text{map}(1, g)(x). \quad (63)$$

If we put $gfx = \text{map}(f, g)(x)$ and use gx and xf instead of $gx1$ and $1xf$, then, the conditions on a map-transformation structure translate as

$$g0f = 0 \quad (64)$$

$$g(x + y)f = gfx + gyf \quad (65)$$

$$1x1 = x \quad (66)$$

$$g'(gfx)f' = (g'g)x(ff') \quad (67)$$

$$\varepsilon(0) = 0 \quad (68)$$

$$g\varepsilon(u)f = \varepsilon(guf) \quad (69)$$

$$\mu(x, \varepsilon(f)) = xf \quad (70)$$

$$\mu(\varepsilon(g), x) = gx \quad (71)$$

$$\mu(x, \mu(x', x'')) = \mu(\mu(x, x'), x'') \quad (72)$$

for every $x, y \in \text{map}(A, B)$, and $u: A \rightarrow B$, $f: A' \rightarrow A$, $f': A'' \rightarrow A'$, $g: B \rightarrow B'$, $g': B' \rightarrow B''$, and $x' \in \text{map}(A', A)$, $x'' \in \text{map}(A'', A')$.

The elements in $\text{map}(A, B)$ are called *map-transformations* from A to B and are represented with double arrows. This means that an element $x \in \text{map}(A, B)$ is displayed as

$$A \xrightarrow{x} B$$

and the element gxf is obtained as

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ f \uparrow & & \downarrow g \\ A' & \xrightarrow{gxf} & B' \end{array} \quad (73)$$

This definition and its interpretation, namely viewing an element $x \in \text{map}(A, B)$ as a super-morphism from A to B is motivated with the striking analogy between map-transformations and 2-cells in a category with a 2-cell structure (or a sesquicategory) as studied in [11]. Indeed, the list of conditions above is very similar to a list in [11], characterizing 2-cell structures in terms of a vertical composition (written additively) and left and right actions (whiskering composition).

The use of the word *transformation* seems to be appropriate in the sense that it captures the idea of something like a morphism, but different. In a category with a map-transformation structure every morphism, f , is also a (map-)transformation, $\varepsilon(f)$, but not every transformation is a morphism. Let us see some examples.

4.3. Examples of categories with map-transformations. A simple case is the category of commutative monoids where we can take the bifunctor map to be precisely the hom -functor. In this case there is no distinction between map-transformations and morphisms. We emphasize the fact that the name map-transformation is being used as a designation for the elements in a given bifunctor $\text{map}(A, B)$ which is part of a map-transformation structure, following the same analogy in which the word *morphism* is used to designate an element in $\text{hom}(A, B)$. Let us see other examples where a map-transformation is not necessarily a morphism.

Example 1. *Let \mathbf{A} be a pointed category. We can always turn \mathbf{A} into a category with a map-transformation structure. Indeed, the adjoint to the forgetful functor from monoids to pointed sets can be used to turn each $\text{hom}_{\mathbf{A}}(A, B)$, a pointed set, into a monoid. The process is standard: generate a free monoid and identify the zero morphism with the neutral element. Let us denote by $\text{map}(A, B)$ the result of that process. A typical element in $\text{map}(A, B)$ is thus a formal sum $h_1 + \cdots + h_n$ of homomorphisms $h_i: A \rightarrow B$ in \mathbf{A} . The inclusion ε is clear and the composition μ is defined as follows. If $h = h_1 + \cdots + h_n$ is an element in $\text{map}(B, C)$ and $h' = h'_1 + \cdots + h'_n$ is an element in $\text{map}(A, B)$, then $\mu(h, h') \in \text{map}(A, C)$ is the formal expression*

$$h_1 h'_1 + \cdots + h_1 h'_n + \cdots + h_n h'_1 + \cdots + h_n h'_n.$$

Observe that this expression is formed using distributivity on the right, but not on the left. This is in accordance with the fact that condition (65) (in the case of monoids) holds for f and g homomorphisms,

whereas if they are merely functions then we have $(x + y)f = xf + yf$ but $g(x + y)$ is not the same as $gx + gy$.

This example is somehow trivial, but if, for every object A in \mathbf{A} , we choose a set of formal expressions to be identified with the identity morphism, 1_A , then some interesting situations are expected in the study of semi-biproducts. Let us now see another sort of examples.

Example 2. Let \mathbf{A} be a pointed category with, say finite limits, and co-products. We can always define a map-transformation structure as follows. For each pair of objects (A, B) let $\text{map}(A, B)$ be $\text{span}(A, B)_{/\sim}$, that is, the collection of all spans (identifying equivalent ones) in \mathbf{A} from A to B . A span from A to B is an ordered pair of morphisms (u, v) as illustrated

$$A \xleftarrow{u} S \xrightarrow{v} B.$$

Two spans are said to be equivalent if there is a zig-zag of span morphisms connecting them (see e.g. [9]).

The inclusion of morphisms into spans is done by the formula $\varepsilon(h) = (1_A, h)$, for every morphism $h: A \rightarrow B$ in \mathbf{A} . The usual composition of spans can be used to define the map $\mu((u, v), (t, s)) = (u\bar{t}, s\bar{v})$, as illustrated in the diagram below (in which \bar{t} and \bar{v} are obtained by pullback)

$$\begin{array}{ccccc}
 & & \bullet & & \\
 & \swarrow \bar{t} & & \searrow \bar{v} & \\
 & \bullet & & \bullet & \\
 \swarrow u & & & & \searrow s \\
 \bullet & & \bullet & & \bullet \\
 \swarrow v & & \swarrow t & & \\
 \bullet & & \bullet & &
 \end{array}$$

This formula, in particular, can be applied to the left and right composition of morphisms with spans, as to fulfil equations (70) and (71). It remains to define a monoid structure on each $\text{map}(A, B)$. The neutral element is necessarily the insertion of the zero morphism, whereas addition is obtained as follows. Let (u, v) and (u', v') be two spans from A to B , then we define $(u, v) + (u', v') = ([u, u'], [v, v'])$ obtained via co-product as illustrated

$$A \xleftarrow{[u, u']} S + S' \xrightarrow{[v, v']} B.$$

Up to equivalence, this operation is associative, and the span $(1_A, 0_{A, B})$ is the neutral element in $\text{map}(A, B)$.

Other technical details are omitted here since we will not, at the moment, analyse this example. Moreover, it is somehow trivial. Nevertheless, if we choose a good class of morphisms in \mathbf{A} , say \mathcal{E} , containing all isomorphisms, which is closed under composition, stable under pullbacks, and if $u, u' \in \mathcal{E}$ are two morphisms with the same codomain, then the induced morphism from the coproduct of their domains, $[u, u']$,

into the common codomain, is still in \mathcal{E} . Then, we can restrict the previous example and consider only those spans (u, v) for which $u \in \mathcal{E}$. This will certainly create interesting examples. Take for instance the class of regular epimorphisms in a semi-abelian category.

The previous example is also related to the notion of imaginary morphisms in the following sense. When considering a good class of morphisms \mathcal{E} it may happen that there exists a pointed endo-functor of \mathbf{A} , say (T, η) for which $\text{map}(A, B) \cong \text{hom}_A(T(A), B)$ so that each class of spans in $\text{span}(A, B)_{/\sim}$ with the left leg in \mathcal{E} is represented by a span of the form

$$A \xleftarrow{\eta_A} T(A) \xrightarrow{f} B.$$

This means that we can *imagine* a map-transformation from A to B as a morphism from $T(A)$ to B (see for example [13] and references therein). These ideas will be developed in future work.

Let us now see a procedure to construct more concrete examples, given some concrete situations.

Example 3. Let \mathbf{A} be a pointed category and $I: \mathbf{A} \rightarrow \mathbf{Mon}$ a faithful functor. Let F denote the forgetful functor from monoids into pointed sets. Then there is a bifunctor

$$\text{map}: \mathbf{A}^{op} \times \mathbf{A} \rightarrow \mathbf{Mon}$$

with $\text{map}(A, B)$ the collection of all pointed maps from the pointed set $FI(A)$ to the pointed set $FI(B)$. The zero map $0_{A,B}: FI(A) \rightarrow FI(B)$ serves as neutral element and the usual component-wise addition of pointed maps, that is, $(f + g)(a) = f(a) + g(a) \in FI(B)$, for every $a \in FI(A)$ and $f, g \in \text{map}(A, B)$, makes $\text{map}(A, B)$ a monoid. It is clear that the element $f(a) + g(a)$ comes from the monoid $I(B)$.

The natural inclusion ε as well as the associative composition μ are not difficult to obtain in order to make \mathbf{A} a category with map-transformations. Indeed, if $h: A \rightarrow B$ is a morphism in \mathbf{A} then $\varepsilon(h)$ is simply the map $FI(h)$, while $\mu(f, g) = fg$ is nothing but the usual composition of set-theoretical pointed maps.

The previous example (as well as the following one) can be generalized to the case where \mathbf{Mon} is replaced with $\mathbf{Mon}(\mathbf{C})$, i.e. the category of internal monoids in a category \mathbf{C} with finite limits. Furthermore, it is clear that $\text{map}(A, B)$ does not need to be equal to the collection of all pointed maps from $FI(A)$ to $FI(B)$. Let us make this assertion more precise.

Example 4. For every commutative square of categories and functors

$$\begin{array}{ccc}
 & \mathbf{A} & \\
 V \swarrow & & \searrow I \\
 \mathbf{B} & & \mathbf{Mon} \\
 U \searrow & & \swarrow F \\
 & \mathbf{Set}_* &
 \end{array}$$

in which \mathbf{A} and \mathbf{B} are pointed categories and all the functors are faithful (recall that F is the forgetful functor from monoids to pointed sets) we can always find a family of submonoids

$$j_{A,B}: \text{map}(A, B) \rightarrow \text{hom}_{\mathbf{Set}_*}(FI(A), FI(B))$$

satisfying the following two conditions:

- (1) the inclusion of $\text{hom}_{\mathbf{B}}(V(A), V(B))$, for every pair of objects A, B , into $\text{hom}_{\mathbf{Set}_*}(FI(A), FI(B))$, factors through $j_{A,B}$, say, as

$$\begin{array}{ccc}
 \text{hom}_{\mathbf{B}}(V(A), V(B)) & \xrightarrow{i_{A,B}} & \text{map}(A, B) \\
 & \searrow & \downarrow j_{A,B} \\
 & & \text{hom}_{\mathbf{Set}_*}(FI(A), FI(B))
 \end{array}$$

- (2) for every three objects A, B, C , the restriction to the composition mapping of pointed maps, factors as follows

$$\begin{array}{ccc}
 \text{map}(B, C) \times \text{map}(A, B) & \xrightarrow{\mu_{A,B,C}} & \text{map}(A, C) \\
 & \searrow & \downarrow j_{A,B} \\
 & & \text{hom}_{\mathbf{Set}_*}(FI(A), FI(C))
 \end{array}$$

In this way we get a bifunctor map which is defined on morphisms, say $f: A' \rightarrow A$, and $g: B \rightarrow B'$,

$$\text{map}(f, g): \text{map}(A, B) \rightarrow \text{map}(A', B')$$

as $\text{map}(f, g)(u) = \mu(i(V(f)), \mu(u, i(V(g))))$. It is a monoid homomorphism because $I(f)$ is a monoid homomorphism and $UV(f) = FI(f)$. The inclusion of morphisms into map-transformations is done with the mapping $\varepsilon(h) = i(V(h))$.

In this paper we have considered the case when $\mathbf{A} = \mathbf{B} = \mathbf{Mon}$, $I = V$ is the identity functor, and $U = F$. It is, nevertheless, clear that several interesting other examples can be considered as well:

- (1) Take \mathbf{A} to be the category of topological groups with I its forgetful functor into monoids and V its forgetful functor into topological spaces;

- (2) Take \mathbf{A} to be the category of topological monoids with I its forgetful functor into monoids and V its forgetful functor into topological spaces;
- (3) Take \mathbf{A} to be the category of (pre)ordered groups with I its forgetful functor into monoids and V its forgetful functor into (pre)ordered sets;

Indeed, when \mathbf{A} is the category of topological groups then we have two possibilities for $\text{map}(A, B)$ (see for example [6]). It can be the set of pointed maps from $FI(A)$ to $FI(B)$, or the set of continuous pointed maps from $FI(A)$ to $FI(B)$. In the same way, the category of (pre)ordered groups (with its forgetful functor into monoids) can be enriched with a map-transformation structure in two different ways. We can take $\text{map}(A, B)$ to be the set of pointed maps from $FI(A)$ to $FI(B)$ but we can also take it to be the set of monotone pointed maps from $FI(A)$ to $FI(B)$. This idea is implicitly used in [4] where a systematic study of semi-direct products is conducted in the category of preordered groups. Other possibilities are also interesting to be considered, such is the case of those pointed maps τ which are not homomorphisms but satisfy the condition $\tau(x + y) \leq \tau(x) + \tau(y)$.

In a certain sense, all examples come from a structure of 2-cells in a pointed category with a horizontal composition. All the details that are omitted in the example below can be found in [11], see also [3].

Example 5. *Let \mathbf{A} be a pointed category equipped with a 2-cell structure $(H, \text{dom}, \text{cod}, 0, +)$. Let us suppose it comes equipped with a horizontal composition law for 2-cells, μ , not necessarily obtained from the middle interchange law. Define,*

$$\text{map}(A, B) = \{x \in H(A, B) \mid \text{dom}(x) = 0 = \text{cod}(x)\}.$$

Now, for every natural inclusion of pointed sets, $\varepsilon: \text{hom} \rightarrow \text{map}$, which satisfies conditions (70) and (71) we obtain a map-transformation structure $(\text{map}, \mu, \varepsilon)$ on \mathbf{A} .

5. SEMI-BIPRODUCTS

In this section we give a definition of semi-biproduct in a category with an abstract map-transformation structure and derive some of its immediate properties.

5.1. Semi-biproducts in categories with map-transformations.

A semi-biproduct, as a generalization of biproduct, makes sense in a category with map-transformations. Indeed, it is obtained from a semi-biproduct in monoids, such as the one displayed in diagram (13), by declaring that the zero-preserving maps q and s are map-transformations (and reformulating the conditions accordingly so that they still make sense).

Definition 2. Let $\mathbf{A} = (\mathbf{A}, \text{map}, \varepsilon, \mu)$ be a category with a structure of map-transformations. A semi-biproduct in \mathbf{A} is a diagram of the form

$$X \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} A \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} B \quad (74)$$

in which X, A, B are objects in \mathbf{A} , p, k are morphisms, q, s are map-transformations, and the following conditions hold:

$$ps = \varepsilon(1_B) \quad (75)$$

$$qk = \varepsilon(1_X) \quad (76)$$

$$kq + sp = \varepsilon(1_A) \quad (77)$$

$$pk = 0_{X,B} \quad (78)$$

$$\mu(q, s) = \varepsilon(0_{B,X}). \quad (79)$$

Recall that $ps = \text{map}(1_B, p)(s) \in \text{map}(B, B)$, $qk = \text{map}(k, 1_X)(q) \in \text{map}(X, X)$. Similarly, kq and sp are elements in $\text{map}(A, A)$ and $kq + sp \in \text{map}(A, A)$ is their sum using the monoid operation defined in the monoid $\text{Map}(A, A)$, see diagram (59).

We will see some immediate properties of this definition, further studies are postponed to future work.

5.2. Immediate properties of semi-biproducts. Let us suppose that (X, A, B, p, k, q, s) is a semi-biproduct in a category with a structure of map-transformations, such as the one displayed in diagram (74). It follows that if k has a certain desirable property, then it is a kernel for p and dually p is a cokernel for k . The desirable property is the ability to *recognize* when a map-transformation, say u , is in the image of ε or not, that is, if it is of the form $u = \varepsilon(f)$ for some morphism f or not. A morphism with this property is said to be a recognizer.

Definition 3. A monomorphism $k: X \rightarrow A$ is said to be a recognizer when for every object Y and map-transformation $u \in \text{map}(Y, X)$, if there exists $f: Y \rightarrow A$ such that $ku = \varepsilon(f)$ then $u = \varepsilon(u')$ for some $u': Y \rightarrow X$.

In other words, k recognizes morphisms if, whenever the composition of k with a map-transformation u results in a morphism ku , then, u is a morphism itself.

Dually, an epimorphism p co-recognizes morphisms if, whenever the composition of a map-transformation v with p results in a morphism vp , then, v is a morphism itself.

In groups (with zero-preserving maps) it is true that every monomorphism is a recognizer and every epimorphism is a co-recognizer. In monoids it is still true that every monomorphism is a recognizer but only surjective homomorphisms (that is, regular epimorphisms) are co-recognizers. In the category of topological groups (with zero-preserving

maps, not necessarily continuous) it is not true in general that monomorphisms are recognizers. The same phenomenon can be observed in the case of preordered groups (with zero-preserving maps, not necessarily monotone).

Theorem 4. *Let (X, A, B, p, k, q, s) be a semi-biproduct in a category with map-transformations $\mathbf{A} = (\mathbf{A}, \text{map}, \varepsilon, \mu)$, then:*

- (1) *the morphism k is a monomorphism;*
- (2) *the morphism p is an epimorphism;*
- (3) *if k recognizes morphisms then it is the kernel of p ;*
- (4) *if p co-recognizes morphisms then it is the cokernel of k .*

Proof. Let $u, v: Y \rightarrow X$ be two morphisms in \mathbf{A} with $ku = kv$. Then $qku = qkv$ and since $qk = \varepsilon(1_X)$ we have $\varepsilon(1_X)u = \varepsilon(1_X)v$. By the naturality of the inclusion of morphisms into map-transformations we have $\varepsilon(1_X u) = \varepsilon(1_X v)$ which gives $\varepsilon(u) = \varepsilon(v)$. We may conclude that $u = v$ because ε is an inclusion. This proves that k is a monomorphism. Dually we prove that p is an epimorphism.

Let $f: Y \rightarrow A$ be a morphism for which $pf = 0$. Then there exists a unique morphism $f': Y \rightarrow X$ such that $\varepsilon(f') = qf$. Indeed, the map-transformation qf is such that $kqf = f$ (because $f = kqf + spf$ and $pf = 0$). The existence of f' comes from the fact that k recognizes morphisms. This proves that k is the kernel of p . Dually we prove that p is the co-kernel of k . \square

There is an analogue to the celebrated split short five lemma for semi-biproducts.

A morphism between semi-biproducts consists of three morphisms (f_0, f_1, f_2) as illustrated

$$\begin{array}{ccccc} X & \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} & A & \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{s} \end{array} & B \\ f_0 \downarrow & & \downarrow f_1 & & \downarrow f_2 \\ X' & \begin{array}{c} \xleftarrow{q'} \\ \xrightarrow{k'} \end{array} & A' & \begin{array}{c} \xleftarrow{p'} \\ \xrightarrow{s'} \end{array} & B' \end{array}$$

such that $q'f_1 = f_0q$, $p'f_1 = f_2p$, $k'f_0 = f_1k$ and $f_1s = s'f_2$.

Theorem 5. *Let (f_0, f_1, f_2) be a morphism of semi-biproducts as displayed above. If f_0 and f_2 are isomorphisms and f_1 recognizes or co-recognizes morphisms, then f_1 is an isomorphism whose inverse f_1^{-1} is such that $\varepsilon(f_1^{-1}) = kf_0^{-1}q' + sf_2^{-1}p'$.*

Proof. Let us suppose, without loss of generality, that $X = X'$, $B = B'$, $f_0 = 1_X$, and $f_2 = 1_B$. Let us also denote f_1 simply by f . Then, it is clear that the map-transformation $g = kq' + sp'$ is such that $gf = (kq' + sp')f = kq'f + sp'f = kq + sp = \varepsilon(1_A)$. Similarly, $fg = \varepsilon(1'_A)$. Hence, f is both a monomorphism and an epimorphism. If f recognizes

or co-recognizes morphisms then it follows that g is in fact a morphism and thus $f^{-1} = g$. \square

6. CONCLUSION

We have tried to give some directions in which the study of semi-biproducts can be carried out in a general context. It seems appropriate to put it into the light of transformation structures and 2-cells. It would be interesting to explore this concept even further, moving beyond the pointed context. For any category \mathbf{A} , together with a 2-cell structure $(H, \text{dom}, \text{cod}, 0, +)$, and assuming the existence of a natural transformation $\varepsilon: \text{hom} \rightarrow H$, it would be interesting to see what is a sequence in \mathbf{A} of the form

$$X \xrightarrow{k} A \xrightarrow{p} B$$

together with 2-cells $q \in H(A, X)$ and $s \in H(B, A)$ such that the following conditions hold

$$\begin{aligned} ps = \varepsilon(1_B) & \quad , & \quad qk = \varepsilon(1_X) \\ \text{dom}(kq) = \text{cod}(sp) & \quad , & \quad kq + sp = \varepsilon(1_A). \end{aligned}$$

In this case there are no zero-morphisms and the horizontal composition μ does not need to be present. This could be applied, for example, in the category of rings.

ACKNOWLEDGEMENTS

This work is supported by the Fundação para a Ciência e a Tecnologia (FCT) and Centro2020 through the Project references: UID/Multi/04044/2019; PAMI - ROTEIRO/0328/2013 (Nº 022158); Next.parts (17963); and also by CDRSP and ESTG from the Polytechnic Institute of Leiria.

REFERENCES

- [1] D. Bourn, N. Martins-Ferreira, A. Montoli and M. Sobral, *Schreier split epimorphisms in monoids and in semirings*, *Textos de Matemática Série B* **45** (Departamento de Matemática, Universidade de Coimbra (ISBN 978-972-8564-49-0)) (2013) vi+116pp. (Cited on page 3.)
- [2] D. Bourn, N. Martins-Ferreira, A. Montoli, M. Sobral, *Schreier split epimorphisms between monoids*, *Semigroup Forum* **88** (2014) 739–752. (Cited on pages 5 and 10.)
- [3] R. Brown, *Possible connections between whiskered categories and groupoids, Leibniz algebras, automorphism structures and local-to-global questions*, *Journal of Homotopy and Related Structures* **1**(1) (2010) 1–13. (Cited on pages 12 and 19.)
- [4] M. M. Clementino, N. Martins-ferreira and A. Montoli, *On the categorical behaviour of preordered groups*, *J. Pure Appl. Algebra* **223** (2019) 4226–4245. (Cited on page 19.)
- [5] I. Fleischer, *Monoid extension theory*, *Journal of Pure and Applied Algebra* **21** (1981) 151–159. (Cited on pages 5 and 7.)

- [6] J. J. Ganci, *Schreier extensions of topological semigroups*, PhD, Louisiana State University (1975). (Cited on pages 3, 5, 8, and 19.)
- [7] M. Gran, G. Janelidze and M. Sobral, *Split extensions and semidirect products of unitary magmas*, arxiv: 1906.02310v2 (math.CT). (Cited on page 5.)
- [8] J. Leech, *Extending groups by monoids*, Journal of Algebra **74** (1982) 1–19. (Cited on pages 4, 7, and 9.)
- [9] S. Mac Lane, *Categories for the Working Mathematician*, 2ed, Graduate Texts in Mathematics 5, Springer, 1998. (Cited on pages 1 and 16.)
- [10] S. Mac Lane, *Cohomology Theory in Abstract Groups. III* Annals of Mathematics, Second Series, **50**(3) (1949) 736–761. (Cited on page 2.)
- [11] N. Martins-Ferreira, *On the notion of pseudocategory internal to a category with a 2-cell structure*, Tbil. Math. J. **8** (1) (2015) 107–141. (Cited on pages 12, 15, and 19.)
- [12] N. Martins-Ferreira, A. Montoli, M. Sobral and A. Patchkoria, *On the classification of Schreier extensions of monoids with non-abelian kernel*, to appear, Forum Mathematicum. (Cited on pages 2, 3, 4, and 5.)
- [13] A. Montoli, D. Rodelo and T. Van der Linden, *Intrinsic Schreier split extensions*, Appl. Categ. Structures (2019). (Cited on page 17.)
- [14] C. Wells, *Extension theories for monoids*, Semigroup Forum **16** (1978) 13–35. (Cited on pages 4 and 7.)

(Nelson Martins-Ferreira) INSTITUTO POLITÉCNICO DE LEIRIA, LEIRIA, PORTUGAL

E-mail address: martins.ferreira@ipleiria.pt