

Abstract rewriting internalized

Maxime LUCAS 

Inria and LS2N, Université de Nantes, France

<http://lucas-webpage.gforge.inria.fr/>

Maxime.Lucas@inria.fr

Abstract

In traditional rewriting theory, one studies a set of terms up to a set of rewriting relations. In algebraic rewriting, one instead studies a vector space of terms, up to a vector space of relations. Strikingly, although both theories are very similar, most results (such as Newman's Lemma) require different proofs in these two settings.

In this paper, we develop rewriting theory internally to a category \mathcal{C} satisfying some mild properties. In this general setting, we define the notions of termination, local confluence and confluence using the notion of reduction strategy, and prove an analogue of Newman's Lemma. In the case of $\mathcal{C} = \text{Set}$ or $\mathcal{C} = \text{Vect}$ we recover classical results of abstract and algebraic rewriting in a slightly more general form, closer to von Oostrom's notion of decreasing diagrams.

2012 ACM Subject Classification Theory of computation \rightarrow Rewrite systems; Theory of computation \rightarrow Equational logic and rewriting

Keywords and phrases Rewriting theory, Internal object, Kleene algebras, Termination

Acknowledgements I thank Mathieu Anel, Cameron Calk, and Damien Pous for fruitful conversations during the preparation of this article, Marie Kerjean for helpful advice to make this article more accessible, and Cyrille Chenavier for both.

1 Introduction

The goal of this work is to bridge the gap between two major branches of rewriting theory: abstract rewriting, stemming from the work of Newman, and algebraic rewriting. In algebraic rewriting, convergent presentations are called Gröbner basis and were introduced by Buchberger to compute basis of algebras. In particular, it allows one to solve the ideal membership problem, which is the linear equivalent of the word problem. Today Gröbner basis are in particular used in control theory.

The two theories are very similar: both define a notion of terminating, confluent or locally confluent relation, and prove a diamond Lemma showing that locally confluent terminating system are confluent (in the case of algebraic rewriting, this was explicitly done by Bergman [2]). Nevertheless, we are not aware of any treatment unifying those two theories.

The main difference between algebraic and abstract rewriting is that algebraic rewriting is not stable by contextual closure. For example, suppose given an rewriting relation \rightarrow on a vector space. If $u \rightarrow v$ is a valid rewriting step, then $u + w \rightarrow v + w$ is not necessarily valid. This is necessary to avoid non-terminating behaviours: otherwise the rewriting step $u - u - v \rightarrow v - u - v$ would be valid, but this is just $-v \rightarrow -u$, which implies that $v \rightarrow u$: in other words, \rightarrow is always symmetric and can never be terminating. The failure to take this phenomenon into account plagued many early papers studying the algebraic λ -calculus, such as [4], and was not recognized before [11].

While the existence of two rewriting theories is not an issue *per se*, the multiplication of applications of higher dimensional rewriting to various algebraic structures calls for a unified framework. Higher dimensional rewriting seeks to apply rewriting techniques to study homotopical and homological properties of algebraic objects. Already existing examples include monoids [9], algebras [5], string diagrams (encoded as Pros and ProPs) [6], and term

rewriting systems [8]. Future cases of interest include e.g. (non-symmetric, symmetric or shuffle, linear or set-theoretic) operads or linear Pros.

All these structures can be represented as monoids inside a category \mathcal{C} : taking $\mathcal{C} = \text{Set}$ we obtain (regular) monoids, monoid objects in $\mathcal{C} = \text{Vect}_k$ are k -algebras, and the different flavors of operads are all monoids in various categories of collections. This paper constitutes a first step towards a unified treatment of higher dimensional rewriting for these different objects, by developing a general theory of rewriting inside a category \mathcal{C} , omitting with the monoid structure for now. In the $\mathcal{C} = \text{Set}$ and $\mathcal{C} = \text{Vect}_k$ we recover respectively abstract and algebraic rewriting, in a slightly more general form than the one usually presented. Those two cases have already presented in earlier works, namely [7] for the case $\mathcal{C} = \text{Set}$, and [3] for the case $\mathcal{C} = \text{Vect}_k$.

There are two ways to model a relation in abstract rewriting. The first one is to see a relation \rightarrow on an object E as a subset of $E \times E$. The second one, which is the one suited for higher dimensional rewriting and that we will generalize here, is to see a relation as a set R equipped with maps $\sigma_R, \tau_R : R \rightarrow E$, associating to any $f \in R$ its source $\sigma_R f$ and target $\tau_R f$, and making (E, R, σ_R, τ_R) a directed graph¹. We therefore study the rewriting properties of graphs internal to an arbitrary category \mathcal{C} satisfying some mild properties. We now describe some of the content of the paper.

While the notions of reflexivity, symmetry and transitivity can be defined straightforwardly in this context, the main obstacle lies in defining an appropriate notion of termination for a graph R on an object E . While in abstract rewriting a relation R is *intrinsically* terminating, the situation is more subtle in algebraic rewriting. There, a relation on a vector space $E = kX$ is said to terminate *with respect to a terminating order on X* . In practice, kX is often the vector space underlying a polynomial algebra, and the terminating order is a monomial order.

In an arbitrary category \mathcal{C} , we proceed similarly, by supposing that E is endowed with a filtration $E_0 \subset E_1 \subset E_2 \subset \dots \subset E$, which encodes the terminating ordering on E . Formally, such a filtration is given by a terminating order (I, \leq) and a functor $I \rightarrow \mathcal{C}$ whose colimit is E . From this we can define the *object of normal forms* of E , denoted E_{min} , as the union of the E_i where i is a minimal element of I .

A graph R on E is then terminating if it is compatible with this filtration in a suitable way, expressed through the existence of a local strategy $h : E \rightarrow R$. When E is a set, a local strategy maps any $x \in E$ to a rewriting step $h(x)$ of source x and whose target is smaller than x . Any local strategy induces a (global) strategy H , mapping any x to a path $H(x)$ to (one of) its normal form, denoted $H^\tau(x)$. The end diagram is the following:

$$\begin{array}{ccccc} & \overset{H}{\curvearrowright} & & \overset{\iota}{\curvearrowright} & \\ R & \xrightarrow[\tau_R]{\sigma_R} & E & \xrightarrow{H^\tau} & E_{min} \end{array}$$

and it satisfies the relations:

$$\sigma_R \circ H = id_E \quad \tau_R \circ H = \iota \circ H^\tau \quad H^\tau \circ \iota = Id_{E_{min}}$$

This almost makes the previous diagram into a split coequaliser, the only equation missing being $H^\tau \circ \sigma_R = H^\tau \circ \tau_R$. We define this as our notion of confluence, which implies

¹ This can also be seen as a *labelled* rewriting relation.

immediately that if R is a terminating and confluent relation, then the quotient E/R is isomorphic to the object of normal forms E_{min} . When \mathcal{C} is a locally finitely presented category, We also define a notion of local confluence and show that, together with termination, it implies confluence, recovering Newman's Lemma in this general setting.

Organisation

In Section 2, we start by recalling some classical definitions and results of abstract and algebraic rewriting. Those will be useful throughout the paper in order to compare them to the methods developed in the subsequent sections.

In Section 3, we investigate elementary properties of graphs internal to a category \mathcal{C} , such as reflexivity, symmetry and transitivity. We also define the notions of reflexive, symmetric and transitive closures of a relation.

Section 4 is devoted to the notion of termination of a graph. We define termination of a relation as the existence of a local strategy compatible with this filtration. We finish this section by examples, showing in particular that for any terminating (abstract or algebraic) relation induces a terminating graph.

In Section 5, we define a notion of global strategy for a graph R on a filtered object E , and define when such a strategy is confluent. We show that whenever there exists a confluent strategy, then the quotient of E by R is isomorphic to the object of normal forms. We show that any local strategy induces a global strategy, although not necessarily a confluent one.

Finally in Section 6 we restrict ourselves to the case when \mathcal{C} is a locally finitely presented category. In this case, we are able to give a criterion for a local strategy to induce a confluent strategy. Interpreting this criterion as a form of local confluence, we obtain a general proof of Newman's and Bergman's diamond Lemmas.

2 Abstract and algebraic rewriting

In this section, we recall some of the standard results of abstract and algebraic rewriting. This will be useful in subsequent sections for comparison with our general theory. Since in Section 5 confluence will only be defined on terminating graphs, we do not dwell on the case of confluent but non-terminating relations. None of the results presented here are new, although perhaps algebraic rewriting is only rarely presented at this level of generality.

Contrary to the rest of this article, where we will model relations by graphs, here we stick to the more usual presentation of subsets of $E \times E$. While this choice is somewhat arbitrary, it will be useful later on in order to distinguish between termination in the sense of relations, and termination in the sense of graphs.

► **Definition 1.** *Let E be a set. By a (set-theoretic) relation, we mean a subset of $E \times E$. We say that such a relation \rightarrow is terminating if there exists no infinite sequence $a_0, a_1, a_2, \dots \in E$ such that $a_0 \rightarrow a_1 \rightarrow a_2 \rightarrow \dots$*

An element $e \in E$ is said to a normal form for \rightarrow if there exists no $y \in E$ such that $e \rightarrow y$. We denote by $NF(\rightarrow)$ the set of normal forms for \rightarrow .

We denote by $\xrightarrow{=}$, $\xrightarrow{+}$, $\xrightarrow{}$ and $\xleftrightarrow{*}$ respectively the reflexive, transitive, reflexive-transitive and symmetric-reflexive-transitive closure of \rightarrow .*

We say that \rightarrow is confluent if for any u, v, w such that $u \xrightarrow{} v$ and $u \xrightarrow{*} w$, there exists z such that $v \xrightarrow{*} z$ and $w \xrightarrow{*} z$.*

We say that \rightarrow is locally confluent if for any u, v, w such that $u \rightarrow v$ and $u \rightarrow w$, there exists z such that $v \xrightarrow{} z$ and $w \xrightarrow{*} z$.*

Finally, we denote by E / \leftrightarrow^* the quotient of E by the equivalence relation generated by \rightarrow .

► **Proposition 2.** Let \rightarrow be a terminating relation on a set E . The following are equivalent:

SC1 The relation \rightarrow is locally confluent.

SC2 The relation \rightarrow is confluent.

SC3 For any $u, v \in E$ such that $u \leftrightarrow^* v$, there exists w such that $u \xrightarrow{*} w$ and $v \xrightarrow{*} w$.

SC4 The canonical map $NF(\rightarrow) \rightarrow (E / \leftrightarrow^*)$, sending any normal form to its equivalence class modulo \leftrightarrow^* , is a bijection.

► **Remark 3.** Property **SC3** above is known as the Church-Rosser property, and is equivalent to confluence even without the hypothesis that \rightarrow is terminating.

Throughout this article, we fix k a characteristic 0 field. We now give a quick presentation of algebraic rewriting.

► **Definition 4.** Let X be a set and let kX denote the vector space spanned by X . Any $u \in kX$ can be written in a unique way as $\sum_{i=1}^n \lambda_i x_i$, where $\lambda_i \neq 0$ and $x_i \in X$. The set $\{x_1, \dots, x_n\}$ is called the support of u and is denoted by $\text{supp}(u)$.

An algebraic relation \rightarrow on kX is a subset of $X \times kX$ such that whenever $x \rightarrow u$, then $x \notin \text{supp}(u)$. Such a relation induces two (set-theoretic) relations on kX , denoted $\xrightarrow{\text{alg}}$ and $\xrightarrow{\text{wf}}$. The first is defined by $\lambda x + v \xrightarrow{\text{alg}} \lambda u + v$ for all $\lambda \in k$, $u, v \in kX$ and $x \in X$ such that $x \rightarrow u$.

The second one is the restriction of $\xrightarrow{\text{alg}}$ to the case when $\lambda \neq 0$ and $x \notin \text{supp}(v)$.

Let \leq be a terminating order on X . We say that \rightarrow is terminating with respect to \leq if whenever $x \rightarrow u$, then any y appearing in the support of u is strictly smaller than x . This implies in particular that $\xrightarrow{\text{wf}}$ is terminating, as a set-theoretic relation.

The set of normal forms for $\xrightarrow{\text{wf}}$ forms a sub-vector space of kX , a basis of which is given by elements $x \in X$ such that there exists no $u \in kX$ such that $x \rightarrow u$. The vector space of normal forms is denoted $NF(\rightarrow)$.

Finally, we denote by \leftrightarrow^* the congruence generated by \rightarrow (which is also the equivalence relation generated by $\xrightarrow{\text{alg}}$).

► **Note 5.** The use of a terminating order on X to define the termination of an algebraic relation above is fairly standard in the literature (and is the one used by Bergman in [2]), but is actually not really necessary: one could instead request that $\xrightarrow{\text{wf}}$ is terminating, as a set-theoretic relation. This is done for example in [5, Section 3.2].

Although this allows one to deduce many results on algebraic rewriting from abstract rewriting, we argue that it is still not satisfying: the definition of the relation $\xrightarrow{\text{wf}}$ is very specific to the category at hand (here $\mathcal{C} = \text{Vect}_k$) and cannot be generalised to other categories, such as the category of groups.

The following lemma clarifies the relationship between the relations \rightarrow , $\xrightarrow{\text{wf}}$ and $\xrightarrow{\text{alg}}$.

► **Lemma 6.** Let X be a set and \rightarrow be an algebraic relation on kX .

■ For any $x \in X$, the relations $x \rightarrow u$ and $x \xrightarrow{\text{wf}} u$ are equivalent.

■ If $u \xrightarrow{\text{alg}} v$, there exists $w \in kX$ such that $u \xrightarrow{\text{wf}} w \xleftarrow{\text{wf}} v$

Proof. While the first point is an easy verification, the second is more subtle. A proof can be found in the proof of [5, Theorem 4.2.1]. ◀

► **Proposition 7.** Let X be a set equipped with a terminating order \leq , and \rightarrow be an algebraic relation on kX which is terminating with respect to \leq . The following are equivalent:

AC1 For any $x \in X$, if $x \rightarrow u$ and $x \rightarrow v$ then there exists w such that $u \xrightarrow[wf]{*} w$ and

$$v \xrightarrow[wf]{*} w.$$

AC2 The relation $\xrightarrow[wf]{*}$ satisfies the equivalent properties of Proposition 2.

AC3 The canonical map $NF(\rightarrow) \rightarrow (kX / \xleftrightarrow{*})$ sending any normal form to its equivalence class modulo $\xleftrightarrow{*}$, is an isomorphism.

► **Example 8.** Take $kX = k[x]$: in other words, $X = \{1, \dots, x^n, \dots\}$. We can equip X with the terminating order induced by $x^i < x^{i+1}$. Given a unitary polynomial $P \in k[x]$ we can write $P = x^n + \sum_{i=0}^{n-1} x^i$. This induces a (confluent and terminating) algebraic rewriting system $x^n \rightarrow -\sum_{i=0}^{n-1} x^i$.


The quotient $kX / \xleftrightarrow{*}$ is none other than $k[X]/(P)$, while $NF(\rightarrow)$ is the subspace of polynomials of degree at most $(n-1)$. Taking $k = \mathbb{R}$ and $P = x^2 + 1$, the isomorphism of property **AC3** is none other than the canonical isomorphism between $\mathbb{R} + x\mathbb{R}$ and $\mathbb{R}[x]/(x^2 + 1) = \mathbb{C}$.

3 Graphs as relations

We fix a category \mathcal{C} , and suppose that \mathcal{C} is finitely complete has all countable (including finitary) coproducts.

In this section, we study define some elementary properties of graphs, seen as generalized relations.

► **Definition 9.** Let \mathcal{C} be a category, and E an object of \mathcal{C} . We denote by Gph_E the category of graphs over E . Objects are triples (R, σ_R, τ_R) , where R is an object of \mathcal{C} and $\sigma_R : R \rightarrow E$ and $\tau_R : R \rightarrow E$ are maps in \mathcal{C} .




A morphism of graphs from (R, σ_R, τ_R) to (S, σ_S, τ_S) is a map $R \rightarrow S$ commuting with σ and τ .

We will often denote a graph (R, σ_R, τ_R) simply by R .

► **Definition 10.** Let E be an object of \mathcal{C} , and let R, S be two graphs over E .

- E canonically inherits the structure of a graph using $\sigma_E = \text{id}_E$ and $\tau_E = \text{id}_E$.
- The product of R and S , denoted RS , is the graph defined by the following pullback, with $\sigma_{RS} = \sigma_R \circ \pi_1$ and $\tau_{RS} = \tau_R \circ \pi_2$:



- The sum of R and S , denoted $R + S$, is the graph $(R \coprod S, \sigma_R \coprod \sigma_S, \sigma_R \coprod \sigma_S)$, where \coprod denotes the coproduct in \mathcal{C} .
- The transitive closure of R , denoted R^+ , is the graph over E with underlying object the countable coproduct $\coprod_{i=1}^{\infty} R^{\times E^i}$, with σ_{R^+} (resp. τ_{R^+}) defined on the component $R^{\times E^i}$ as the composite $\sigma_R \circ \pi_1$ (resp. $\tau_R \circ \pi_i$) for any $i \geq 1$.
- The transitive reflexive closure of R , denoted R^* , is the graph $R^+ + E$.
- The opposite of R , denoted R° , is the graph obtained by reversing the source and target of R : $\sigma_{R^\circ} = \tau_R$ and $\tau_{R^\circ} = \sigma_R$.
- The reflexive symmetric transitive closure of R , denoted R^S , is the graph $(R + R^\circ)^*$.

► **Definition 11.** Let R be a graph over an object E in a category \mathcal{C} . We say that R is:

- reflexive if there exists a map (of graphs) $u_R : E \rightarrow R$,
- transitive if there exists a map $m_R : RR \rightarrow R$,
- symmetric if there exists a map $s_R : R^\circ \rightarrow R$.

The closure terminology of Definition 10 is justified by the following lemma:

► **Lemma 12.** Let R be a graph over an object E in a category \mathcal{C} . Then $R + E$ is a universal among the reflexive graphs under R , in the sense that $R + E$ is reflexive, and for any map $f : R \rightarrow S$ in \mathbf{Gph}_E , where S is a reflexive graph, then there exists a unique map $\tilde{f} : R + E \rightarrow S$ preserving the reflexive structure such that the composite with the inclusion $R \rightarrow R + E$ yields f .

Similarly:

- The graph $R + R^\circ$ is universal among symmetric graphs under R .
- The functor R^+ is universal among transitive graphs under R .
- The functor R^* is universal among reflexive transitive graphs under R .
- The functor R^S is universal among reflexive transitive symmetric graphs under R .

► **Example 13.** In the case $\mathcal{C} = \mathbf{Set}$, a graph over a set E is just a graph with set of objects E , and reflexivity and symmetry coincide with the usual terminology.

If \mathcal{C} is a Mal'cev variety, that is \mathcal{C} is the category of models of an algebraic theory containing a ternary operation t such that $t(x, x, y) = y$ and $t(x, y, y) = x$, then reflexivity implies symmetry and transitivity. Mal'cev varieties include $\mathcal{C} = \mathbf{Gp}$, the category of groups, whose where the ternary operation is given by $t(x, y, z) = xy^{-1}z$. Other examples also include the category of abelian groups, vector spaces, or the one of heaps.

In the case of $\mathcal{C} = \mathbf{Vect}_k$, the category of k -vector spaces (or indeed any abelian category), then a truncation of the Dold-Kan correspondence shows that the category of reflexive graphs is equivalent to the category of chain complexes concentrated in degrees 0 and 1.

Moving from a graph R to one of its closure does not change the equivalence relation presented by R . This is made formal by the following lemma.

► **Lemma 14.** If R is a graph on an object E of \mathcal{C} , let us denote by E/R the coequaliser of the diagram $\sigma_R, \tau_R : R \rightarrow E$.

Then for any graph R over E , there are canonical isomorphisms $E/R = E/R^* = E/R^S$.

4 Internalizing termination

We fix a category \mathcal{C} with all pullbacks and countable coproducts.

In this section, we define what it means for a graph to be terminating. This property is expressed in term of a rewriting strategy. When $\mathcal{C} = \mathbf{Set}$, such a strategy associates to any

x which is not a normal form a rewriting step of source x . We show how the examples of terminating rewriting relations of Section 2 fit into the general definition, but also that some non-terminating relations can be terminating as graphs.

► **Note 15.** If I is partially ordered set, we silently see I as a category, with an arrow $x \rightarrow y$ whenever $x \leq y$ in I .


► **Definition 16.** A directed set is a non-empty partially ordered set I such that for all $x, y \in I$, there exists z such that $x, y \leq z$.

Let E be an object of \mathcal{C} . A directed structure on E is the data of a directed set I together with a functor $E_\bullet : I \rightarrow \mathcal{C}$ and a natural transformation $\iota_\bullet : E_\bullet \rightarrow E$ exhibiting E as the colimit of E_\bullet . If I is a directed set, the data of E_\bullet and ι_\bullet is called an I -filtration of E , and we slightly abuse notations by saying that E is an I -filtered object if it comes equipped with an i -filtration $(E_\bullet, \iota_\bullet)$.

A morphism of I -filtered object from E to F is the data of a natural transformation $\eta : E_\bullet \Rightarrow F_\bullet$. Note that this induces a map from E to F .

We denote by $I\text{-Filt}_{\mathcal{C}}$ the category of I -filtered objects in \mathcal{C} .

► **Note 17.** Identifying an object E of \mathcal{C} with a functor from the terminal category $\top \xrightarrow{E} \mathcal{C}$, the data of a filtered object in \mathcal{C} fits into the following diagram:



► **Definition 18.** Let I be a directed set. For any $i \in I$ we denote by $I_{<i}$ the subset of elements of I smaller than i , and by I_{\min} the set of minimal elements of I .

If E is an I -filtered object of a category \mathcal{C} , then for any $i \in I$, we denote by $E_{<i}$ (resp. E_{\min}) the colimit of the restriction of the functor E_\bullet to $I_{<i}$ (resp. I_{\min}). For $i \in I_{\min}$, we define $E_{<i}$ as E_i . This defines a functor $E_{<\bullet} : I \setminus I_{\min} \rightarrow \mathcal{C}$. The universal property of the colimit induces a natural transformation $\iota_{<\bullet} : E_{<\bullet} \rightarrow E$.

If $f : E \rightarrow F$ is a map in \mathcal{C} , it induces a natural transformation $f_\bullet : E_\bullet \Rightarrow F$. For any $i \in I \setminus I_{\min}$, restricting f_\bullet to $I_{<i}$ and passing to the colimit induces a map $f_{<i} : E_{<i} \rightarrow F$, which we extend for $i \in I_{\min}$ by setting $f_{<i} := f \circ \iota_i$. This defines a natural transformation $f_{<\bullet} : E_{<\bullet} \Rightarrow F$.

► **Note 19.** By definition, I_{\min} is a discrete partially ordered set (i.e. for all $i, j \in I_{\min}$, if $i \leq j$ then $i = j$). As a result E_{\min} is actually given by the coproduct $E_{\min} = \coprod_{i \in I_{\min}} E_i$.

► **Definition 20.** A terminating graph on an I -filtered object E of a category \mathcal{C} is the data of a graph $R \in \text{Gph}_E$, together with a map $h : E \rightarrow R + E$ and a natural transformation $h^\tau : E_\bullet \Rightarrow E_{<\bullet}$ such that


TG1 I is a terminating order.

TG2 For all $i \in I$, $\sigma_{R+E} \circ h = \text{id}_E$.

TG3 For all $i \in I_{\min}$, $h \circ \iota_i = u_{R+E} \circ \iota_i$.


TG4 For all $i \in I$, $\tau_{R+E} \circ h \circ \iota_i = \iota_{<i} \circ h_i^\tau$.

The last two axioms can be represented by the following diagram:



► **Note 21.** Note also the last two axioms imply that for all $i \in I_{min}$, $h_i^\tau = id_{E_i}$.

► **Example 22.** While in terminating relations *every* rewriting step is supposed to be decreasing, the definition of terminating graph defined above only requires that there exists a decreasing rewriting step of source x whenever x is not a normal form. For example, the following graph is terminating:




Formally, we take $I = \{0, 1\}$, $E_0 = \{c, d\}$ and $E_1 = E = \{a, b, c, d\}$. As a result, the set of normal forms is $E_{min} = \{c, d\}$.

Defining a strategy (h, h^τ) on $R = \{f_1, f_2, f_3, f_4\}$ amounts to choosing for any $x \in E$ that is not a normal form a rewriting step $h(x) : x \rightarrow h^\tau(x)$. The fact that h^τ lands in $E_{<i}$ rather than E witnesses the fact that $h^\tau(x)$ is in some sense “smaller” than x . On normal forms, axiom **TG3** forces $h(c) = 1_c$ and $h(d) = 1_d$, and by axiom **TG4**, we necessarily have $h_0^\tau(c) = c$ and $h_0^\tau(d) = d$.

For $i = 1$ by naturality of h_\bullet^τ we still have $h_1^\tau(c) = c$ and $h_1^\tau(d) = d$. Finally we set $h(a) = f_3$ and $h(b) = f_4$, which forces $h^\tau(a) = c$ and $h^\tau(b) = d$.

We represent the situation by the diagram on the left hand side of the following picture. The horizontal bar denotes the filtration of E induced by I , while the thick red arrows denote the arrows selected by the local strategy.



The fact that h_i^τ lands in $E_{<i}$ is represented by the fact that the chosen arrows each go one step lower in the filtration.

The diagram on the right hand side of the figure above pictures another filtration of E (given by $I = \{0 < 1 < 2\}$, with $E_0 = \{c, d\}$, $E_1 = \{b, c, d\}$ and $E_2 = \{a, b, c, d\}$) and another local strategy, given by $h(a) = f_1$ and $h(b) = f_4$.

The following example explains how any terminating relation on a set E induces both a filtration of E , and a terminating graph on E .

► **Example 23.** Let (E, \rightarrow) be a set equipped with a terminating relation. Then E is naturally equipped with a structure of an \mathbb{N} -filtered set as follows:

$$E_0 = NF(\rightarrow) \quad E_{i+1} = \{x \in E \mid \exists y \in E_i, x \xrightarrow{=} y\}$$

Notice that for all $i \in \mathbb{N}$, $\mathbb{N}_{<i+1}$ has a terminal element (namely i), and so $E_{<i+1}$ is simply E_i .

In addition, \rightarrow induces a terminating graph (R, h, h^τ) on E , where:

- $R = \{(x, y) \mid x \rightarrow y\}$ with operations σ_R and τ_R given by the first and second projections respectively.
- For $x \in E_0$, i.e. x is a normal form, we set $h_0^\tau(x) = x$.
- For any $i \in \mathbb{N}$, take $x \in E_{i+}$. If $x \in E_i$, then we pose $h_{i+}^\tau(x) = h_i^\tau(x)$. Otherwise, by definition there exists $y \in E_i$ such that $x \rightarrow y$, and we pose $h_{i+1}^\tau(x) = y$.
- Finally, $h(x)$ is given by the pair $(x, h_i^\tau(x))$, which does not depend on i by definition of h_\bullet^τ .

The next lemma will allow us to easily build filtered objects in categories other than sets.

► **Lemma 24.** Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a cocontinuous functor. Then for any directed set I , F induces a functor

$$I\text{-Filt}_F : I\text{-Filt}_{\mathcal{C}} \rightarrow I\text{-Filt}_{\mathcal{D}}.$$

In addition, for any I -filtered object E , $F(E)_i = F(E_i)$, $F(E)_{\min} = F(E_{\min})$ and $F(E)_{<i} = F(E_{<i})$.

Proof. Let E be an I -filtered object in \mathcal{C} . Since F preserves colimits, $(F(E), I, F \circ E_\bullet, F \circ \iota_\bullet)$ is a filtered object in \mathcal{D} . The image of maps by F is as straightforward. ◀

► **Example 25.** Let X be a set equipped with a terminating order \leq , and let \rightarrow be an algebraic relation on kX which is terminating with respect to \leq .

To exhibit a filtration on kX , we first define an map $\text{ht} : X \rightarrow \mathbb{N}$, by induction on $x \in X$, using the order \leq . Without loss of generality, we can suppose that $NF(\rightarrow)$ is equal to the subspace of kX spanned by the $x \in X$ which are minimal for \leq (otherwise, replace \leq by the relation $x \prec y$ defined by $x < y$ and $y \notin NF(\rightarrow)$).

- If x is minimal for \leq , i.e. x is a normal form, we define $\text{ht}(x) = 0$.
- Otherwise, we define:

$$\text{ht}(x) := \min_{\substack{u \in kX \\ x \rightarrow u}} \left\{ \max_{y \in \text{supp}(u)} \text{ht}(y) \right\} + 1$$

We finally define a \mathbb{N} -filtration on X by setting $X_i = \{x \in X \mid \text{ht}(x) \leq i\}$. By the previous lemma, this filtration can be transported to kX . More precisely, a linear combination $u \in kX$ lies in kX_i if and only if $\text{supp}(u) \subseteq X_i$.

Then \rightarrow induces a reflexive graph R , which is the subspace of $kX \times kX$ generated by the pairs (x, u) such that $x \xrightarrow{\equiv} u$. To show that this graph is terminating, we define h^τ and h as follows:

- Let $x \in X_0$, that is x is a normal form. We then define $h^\tau(x) = x$, and extend it to kX_0 by linearity.
- For any $i \in I$, let $x \in X_{i+1}$. If $x \in X_i$, then we simply define $h_{i+1}^\tau(x) = h_i^\tau(x)$. Otherwise, by definition there exists $u \in kX_i$ such that $x \rightarrow u$, and we define $h_{i+1}^\tau(x) = u$. We finally extend h_{i+1}^τ to kX_{i+1} by linearity.
- For any $x \in X$, we finally define $h(x) = (x, h^\tau(x))$, and extend it to kX by linearity.

► **Remark 26.** Let $(R, h_\bullet, h_\bullet^\tau)$ be a terminating graph on an a filtered object E , and S be an other graph over E . For any arrow $f : R \rightarrow S$, $(S, f \circ h_\bullet, h_\bullet^\tau)$ is a terminating graph over E . In particular, the various closures of R defined in Section 3 all inherit a canonical structure of terminating graph over E , which we still denote h_\bullet and h_\bullet^τ .

5 Confluence and strategies

In this section, we define the notion of confluence and local confluence of a graph. Our criterion for confluence is that a for a confluent and terminating graph, the quotient by the graph should be isomorphic to the object of normal forms. We show this property, and prove that local confluence together with termination imply confluence: this is Newman's Lemma.

► **Definition 27.** Let \mathcal{C} be any category. A split coequalizer in \mathcal{C} is a diagram

$$\begin{array}{ccccc} & \overset{t}{\curvearrowright} & & \overset{s}{\curvearrowright} & \\ A & \xrightarrow{f} & B & \xrightarrow{e} & C \\ & \underset{g}{\curvearrowright} & & & \end{array}$$

such that:

$$e \circ f = e \circ g \quad e \circ s = id_C \quad s \circ e = g \circ t \quad f \circ t = id_B$$

For any such diagram, e is necessary a coequalizer of f and g , which we denote by $B/A = C$. In addition, such a coequaliser is absolute, meaning that it is preserved by any functor.

► **Definition 28.** Let R be a graph on a filtered object E . A (global) strategy for R is the data of a pair of morphisms $H : E \rightarrow R$ and $H^\tau : E \rightarrow E_{min}$ such that the following equations hold:

$$\sigma_R \circ H = id_E \quad \tau_R \circ H = \iota_{min} \circ H^\tau \quad H^\tau \circ \iota_{min} = Id_{E_{min}}$$

A strategy (H, H^τ) is said to be confluent if $H^\tau \circ \sigma_R = H^\tau \circ \tau_R$

The following proposition immediately follows from the definition of confluent strategy.

► **Proposition 29.** Let R be a graph over a filtered object E . If there exists a confluent strategy (H, H^τ) for R then the diagram

$$\begin{array}{ccccc} & \overset{H}{\curvearrowright} & & \overset{\iota_{min}}{\curvearrowright} & \\ R & \xrightarrow[\tau_R]{\sigma_R} & E & \xrightarrow{H^\tau} & E_{min} \end{array}$$

forms a split coequalizer. In particular, $E/R = E_{min}$.

► **Theorem 30.** *Let $(R, h_\bullet, h_\bullet^\tau)$ be a terminating graph on an I -filtered object E . Then there is a strategy (H, H^τ) for R^S , which satisfy the additional equations:*

$$H = \mu_R \circ \langle h, H \circ \tau_{R+E} \circ h \rangle \quad H^\tau = H^\tau \circ \tau_{R+E} \circ h$$

Where μ_R denotes the canonical map $RR^S \rightarrow R^S$.

Proof. Since E is the colimit of the functor E_\bullet , it is enough to define natural transformations $H_\bullet : E_\bullet \Rightarrow R^S$ and $H_\bullet^\tau : E_\bullet \Rightarrow E_{\min}$ satisfying the following equations:

$$\sigma_{R^S} \circ H_\bullet = \iota_\bullet \quad \tau_{R^S} \circ H_\bullet = \iota_{\min} \circ H_\bullet^\tau \quad H_i^\tau = \iota_i, \text{ where } i \in I_{\min}.$$

$$H_\bullet = \mu_R \circ \langle h \circ \iota_\bullet, H \circ \tau_{R+E} \circ h \circ \iota_\bullet \rangle \quad H_\bullet^\tau = H^\tau \circ \tau_{R+E} \circ h \circ \iota_\bullet$$

Note that using the equations relating h to h^τ , the last two relations can be rewritten as:

$$H_\bullet = \mu_R \circ \langle h \circ \iota_\bullet, H_{<\bullet} \circ h_\bullet^\tau \rangle \quad H_\bullet^\tau = H_{<\bullet}^\tau \circ h_\bullet^\tau$$

We proceed by induction on $i \in I$ to build such natural transformations. If i is minimal, then the third equation entirely determines H_i^τ , and we take $H_i := u_{R^S} \circ \iota_i$. Then the first equation holds because $\sigma_{R^S} \circ H_i = \sigma_{R^S} \circ u_{R^S} \circ \iota_i = \iota_i$, while for the second we have $\tau_{R^S} \circ H_i = \iota_i = \iota_{\min} \circ \iota_i$, where the second equality holds because i is minimal.

Suppose now that H_\bullet and H_\bullet^τ are defined for all $j < i$, naturally in j . Then they induce functors $H_{<i} : E_{<i} \rightarrow R^S$ and $H_{<i}^\tau : E_{<i} \rightarrow E_{\min}$. By induction hypothesis those satisfy the following equations:

$$\sigma_{R^S} \circ H_{<i} = \iota_{<i} \quad \tau_{R^S} \circ H_{<i} = \iota_{\min} \circ H_{<i}^\tau \quad H_{<i}^\tau \circ \iota_j = \iota_j, \text{ where } j \in I_{\min}.$$

Before defining H_i let us consider the two maps h_i and $H_{<i} \circ h_i^\tau$, from E_i to respectively R and R^S . Notice that we have $\tau_R \circ h_i = \iota_{<i} \circ h_i^\tau = \sigma_{R^S} \circ H_{<i} \circ h_i^\tau$, and so these two maps induce a map $\langle h_i, H_{<i} \circ h_i^\tau \rangle : E_i \rightarrow RR^S$. We finally define:

$$H_i := m_R \circ \langle h_i, H_{<i} \circ h_i^\tau \rangle \quad H_i^\tau := H_{<i}^\tau \circ h_i^\tau.$$

The fact that H_i and H_i^τ satisfy the required equations in all cases is then a straightforward verification. ◀

6 From local to global confluence

The goal of this section is to give a local confluence criteria in order to prove confluence. In this section, we suppose that \mathcal{C} is a locally finitely presentable category.

► **Definition 31.** *Let (R, h, h^τ) be a terminating graph over a filtered object E . A local-confluence structure (or lc-structure for short) on (R, h, h^τ) is the structure of a J -filtered object on R , where J is the category associated to a total terminating order such that $R_{\min} = \emptyset$, and a natural transformation $c_\bullet : R_\bullet \Rightarrow R_{<\bullet}^S$ satisfying the equations:*

$$\sigma_{R^S} \circ \iota_{<\bullet}^S \circ c_\bullet = \tau_{R+E} \circ h \circ \sigma_{R_\bullet} \quad \tau_{R^S} \circ \iota_{<\bullet}^S \circ c_\bullet = \tau_{R_\bullet}$$

The reason for the restriction to the case where \mathcal{C} is locally finitely presentable, and J is total is the following lemma, that we implicitly use in the proof of Theorem 33.

► **Lemma 32.** *Let \mathcal{C} be a locally finitely presented category, and let R be a graph on a filtered object E . Suppose that R is J -filtered. Then*

- *The functor $(R_\bullet)^* : J \rightarrow \mathcal{C}$ is a J -filtration of R^* .*
- *If J is total, then for any $j \in J$, $(R_{<j})^S = (R^*)_{<j}$.*

In addition, the same properties hold for $R + R^\circ$, and thus for R^S .

Proof. Since colimits commute with colimits, we just have to prove that $\text{colim}_j R_j^{\times_{E^n}} = R^{\times_{E^n}}$. For $n = 0, 1$ this is clear. Let us treat the case $n = 2$, the general case being similar. Note first that since J is directed the inclusion $J \rightarrow J \times J$ is final, and so $\text{colim}_j R_j \times_E R_j = \text{colim}_{i,j} R_i \times_E R_j$. Finally in a locally finitely presentable category pullbacks preserve directed colimits [1, Proposition 1.59], and so we get:

$$\text{colim}_j R_j \times_E R_j = \text{colim}_i (\text{colim}_j (R_i \times_E R_j)) = \text{colim}_i (R_i \times_E R) = R \times_E R$$

To compute $(R_{<j})^*$ we use the same technique, using the fact that since J is total then $I_{<j}$ is still directed.

The case of $R + R^\circ$ is clear since colimits commute with colimits, and the case of R^S follows by combining the two previous cases. ◀

► **Theorem 33 (Newman's Lemma for graphs).** *Let (R, h, h^τ) be a terminating graph over a filtered object E , equipped with an lc-structure. Then the strategy induced by (R, h, h^τ) on R^S is confluent.*

In particular, $E/R = E/R^S = E_{\min}$.

Proof. Let us prove by induction on $j \in J$ that the equality $H^\tau \circ \sigma_{R_j^S} = H^\tau \circ \tau_{R_j^S}$ holds.

Suppose first that j minimal. Since $R_{\min} = \coprod_{j \in J_{\min}} R_j = \emptyset$, then $R_j = \emptyset$. The required equation thus holds by unicity of the maps from the initial object.

Otherwise, by induction we have $H^\tau \circ \sigma_{R_{<j}^S} = H^\tau \circ \tau_{R_{<j}^S}$. Then we get $H^\tau \circ \sigma_{R_j} = H^\tau \circ \tau_{R_j}$. Indeed we have:

$$\begin{aligned} H^\tau \circ \sigma_{R_j} &= H^\tau \circ \tau_{R+E} \circ h \circ \sigma_{R_j} \\ &= H^\tau \circ \sigma_{R^S} \circ \iota_{<j}^S \circ c_j \\ &= H^\tau \circ \tau_{R_j^S} \circ \iota_{<j} \circ c_j \\ &= H^\tau \circ \tau_{R_j} \end{aligned}$$

By exchanging the roles of σ_{R_j} and τ_{R_j} , the same equality holds for R_j° . Let us denote by T the sum $R_j + R_j^\tau$ and by T^n the limit $T^{\times_{E^n}}$, and let us show that for all $n \in \mathbb{N}$, $H^\tau \circ \sigma_{T^n} = H^\tau \circ \tau_{T^n}$. For $n = 1$ this is the previous discussion. For $n = 0$ σ_{T^0} and τ_{T^0} coincide and the equality holds. Finally for $n \geq 2$ we have, using repeatedly $n = 1$:

$$H^\tau \circ \sigma_{T^n} = H^\tau \circ \sigma_T \circ \pi_1 = H^\tau \circ \tau_T \circ \pi_1 = H^\tau \circ \sigma_T \circ \pi_2 = \dots = H^\tau \circ \tau_T \circ \pi_n = H^\tau \circ \tau_{T^n}.$$

Putting all those equalities together we finally get $H^\tau \circ \sigma_{R_j^S} = H^\tau \circ \tau_{R_j^S}$, and by colimit $H^\tau \circ \sigma_{R^S} = H^\tau \circ \tau_{R^S}$. Hence H is indeed a confluent strategy on R^S . The equality $E/R^S = E_{\min}$ follows by Proposition 29, and $E/R = E/R^S$ by Lemma 14. ◀

► **Example 34.** Let (E, \rightarrow) be a set equipped with a terminating relation. Following Example 23, we can equip E with an \mathbb{N} -filtered structure, and \rightarrow induces a terminating graph

$R = \{(x, y) | x \rightarrow y\}$ on E . Then there is a canonical filtration of R given by the following pullback:

$$\begin{array}{ccc} R_j & \longrightarrow & E_j \\ \downarrow & & \downarrow \iota_j \\ R & \xrightarrow{\sigma_R} & E. \end{array}$$

Since \mathbf{Set} is locally finitely presented, pullbacks preserve directed colimits and thus $\text{colim}_j R_j = R$. More precisely, $(x, y) \in R$ lies in R_j if and only if x lies in E_j . In particular $R_0 = \emptyset$ since E_0 is the set of normal forms of \rightarrow .

Then the existence of an lc-structure c_\bullet on R is equivalent to the existence for any $r = (x, y) \in R$ of some equivalence path $c(r) : h^\tau(x) \xleftrightarrow{*} y$ of shape:

$$\begin{array}{ccc} x & \xrightarrow{r} & y \\ h(x) \downarrow & \nearrow c(r) & \\ h^\tau(x) & & \end{array}$$

such that any rewriting step occurring in $c(x)$ has source smaller than x . This is true whenever \rightarrow is locally confluent, and Theorem 33 recovers that R is confluent.

The same phenomenon applies to the case $\mathcal{C} = \mathbf{Ab}$, which is also locally finitely presented.

► **Note 35.** The idea of ordering the relations is not new in abstract rewriting, and is for example the main idea behind van Oostrom's notion of decreasing diagrams [10]. In fact when specialised to the case $\mathcal{C} = \mathbf{Set}$, Theorem 33 (which is more general than Newman's Lemma since, as noted in Example 22, termination in the sense of Definition XX is more general than the usual termination of relations) is a direct consequence of [10, Theorem 3.7]

References

- 1 Jiří Adámek and Jiří Rosický. *Locally presentable and accessible categories*, volume 189 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1994. URL: <https://doi.org/10.1017/CB09780511600579>, doi:10.1017/CB09780511600579.
- 2 George M. Bergman. The diamond lemma for ring theory. *Adv. in Math.*, 29(2):178–218, 1978. URL: [https://doi.org/10.1016/0001-8708\(78\)90010-5](https://doi.org/10.1016/0001-8708(78)90010-5), doi:10.1016/0001-8708(78)90010-5.
- 3 Cyrille Chenavier and Maxime Lucas. The diamond lemma for non-terminating rewriting systems using deterministic reduction strategies. 2019.
- 4 Thomas Ehrhard and Laurent Regnier. The differential lambda-calculus. *Theoret. Comput. Sci.*, 309(1-3):1–41, 2003. URL: [https://doi.org/10.1016/S0304-3975\(03\)00392-X](https://doi.org/10.1016/S0304-3975(03)00392-X), doi:10.1016/S0304-3975(03)00392-X.
- 5 Yves Guiraud, Eric Hoffbeck, and Philippe Malbos. Convergent presentations and polygraphic resolutions of associative algebras. *Math. Z.*, 293(1-2):113–179, 2019. URL: <https://doi.org/10.1007/s00209-018-2185-z>, doi:10.1007/s00209-018-2185-z.
- 6 Yves Guiraud and Philippe Malbos. Higher-dimensional categories with finite derivation type. *Theory Appl. Categ.*, 22:No. 18, 420–478, 2009.
- 7 Maxime Lucas. *Cubical categories for homotopy and rewriting*. Theses, Université Paris 7, Sorbonne Paris Cité, December 2017. URL: <https://hal.archives-ouvertes.fr/tel-01668359>.
- 8 Philippe Malbos and Samuel Mimram. Homological computations for term rewriting systems. In *1st International Conference on Formal Structures for Computation and Deduction*, volume 52 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. No. 27, 17. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2016.

- 9 Craig C. Squier, Friedrich Otto, and Yuji Kobayashi. A finiteness condition for rewriting systems. *Theoret. Comput. Sci.*, 131(2):271–294, 1994. URL: [https://doi.org/10.1016/0304-3975\(94\)90175-9](https://doi.org/10.1016/0304-3975(94)90175-9), doi:10.1016/0304-3975(94)90175-9.
- 10 Vincent van Oostrom. Confluence by decreasing diagrams converted. In *Rewriting techniques and applications*, volume 5117 of *Lecture Notes in Comput. Sci.*, pages 306–320. Springer, Berlin, 2008. URL: https://doi.org/10.1007/978-3-540-70590-1_21, doi:10.1007/978-3-540-70590-1_21.
- 11 Lionel Vaux. The algebraic lambda calculus. *Math. Structures Comput. Sci.*, 19(5):1029–1059, 2009. URL: <https://doi.org/10.1017/S0960129509990089>, doi:10.1017/S0960129509990089.