
A QUALITATIVE STUDY OF AN ANAEROBIC FOOD-WEB REVEALS THE IMPORTANCE OF HYDROGEN FOR MICROBIAL STABILITY

A PREPRINT

Szymon Sobieszek

Department of Mathematics & Statistics
McMaster University
Hamilton, ON Canada
sobieszs@mcmaster.ca

Gail S. K. Wolkowicz

Department of Mathematics & Statistics
McMaster University
Hamilton, ON Canada
wolkowicz@mcmaster.ca

Matthew J. Wade

School of Engineering
Newcastle University, Newcastle-upon-Tyne
United Kingdom
matthew.wade@ncl.ac.uk

February 2, 2022

ABSTRACT

The mathematical analysis of a three-tiered food-web describing anaerobic chlorophenol mineralisation has suggested the emergence of interesting dynamical behaviour through its specific ecological interactions, which include competition, syntrophy and product inhibition. Previous numerical analyses have revealed the possibility for a Hopf bifurcation occurring through the interior equilibrium and the role of extraneous substrate inputs in both mitigating the emergence of periodic solutions and expanding the desired stable positive steady-state, where full mineralisation occurs. Here we show that, for a generalised model, the inflow of multiple substrates results in greater dynamical complexity and prove the occurrence of a supercritical Hopf bifurcation resulting from variations in these operating parameters. Further, using numerical estimation, we also show that variations in the dilution rate can lead to Bogdanov-Takens and Bautin bifurcations. Finally, we are able to show apply persistence theory for a range of parameter sets to demonstrate unique persistence in the cases where chlorophenol and hydrogen are extraneously added to the system, mirroring recent applied studies highlighting the role of hydrogen in maintaining stable anaerobic microbial communities.

Keywords Chlorophenol mineralisation; Bifurcation analysis; Hopf bifurcation; Hydrogen

1 Introduction

A mathematical model of the complete anaerobic mineralisation of a generic monochlorophenol isomer (C_6H_4ClOH) by a canonical food-web has recently been described [18]. The system comprises three microbial species (chlorophenol degrader, phenol degrader, hydrogenotrophic methanogen) whose interactions are summarised as follows:

- Reductive dehalogenation of chlorophenol in the presence of hydrogen by the chlorophenol degrader producing phenol [13];
- Phenol is mineralised to acetate and hydrogen via the benzoyl-CoA pathway or a caproate intermediary [10];

- Anaerobic acetogenic (acetate producing) processes are known to be endergonic (the reaction results in a net loss of energy to the system). The production of hydrogen can lead to thermodynamic constraints, or inhibition, if its partial pressure is high enough. In other words, the reaction becomes decreasingly exergonic as more hydrogen is produced until it ceases to be thermodynamically favourable [15, 8]. Hydrogen scavengers such as the methanogen form a syntrophic partnership with the phenol degrader by maintaining the hydrogen partial pressure at concentrations low enough for the mineralisation reaction to proceed;
- Given that the chlorophenol degrader may also act as a syntrophic partner with the phenol degrader, a competitive interaction between the two hydrogen utilisers occurs. This positive and negative feedback loops reframes the ecological network from being a simple food-chain to a more complex food-web that allows for the possibility of periodicity. This additionally leads to the possibility of the system reducing to a self-sustaining two-species network in which the chlorophenol degrader acts as the syntrophic partner to the phenol degrader.

For reference, the food-web is presented diagrammatically in [18] (Figure 8) and [14] (Figure 1). The model is a simplified representation of the system at the population level, ignoring metabolic intermediates and dead-end products such as methane, which does not contribute to the process dynamics. Acetoclastic methanogenesis, the conversion of acetate to methane, is also omitted from the model.

Hydrogen, however, has been shown to play an important role in stability of anaerobic microbial communities through the effects of inhibition and competition [1, 4, 17]. Given that external hydrogen addition will maintain the methanogen population (no washout when the methanogen growth rate is greater than the combined dilution and decay/maintenance rates) under a wider operating parameter regime (chlorophenol inflow and dilution rate) [18], a global analysis of the model can provide deeper insights into the ecological role of hydrogen through its association with community stability and criticality of the Hopf bifurcation.

Here, we focus on the mathematical analysis of the model, extending the work reported in the literature. For example, an analytical approach was taken to characterise the existence and stability of the system equilibria with and without inclusion of a microbial decay term using a general representation of the species growth functions [14]. With no decay, local stability and the conditions giving rise to asymptotic coexistence of all three species have been shown analytically, where the possibility of periodic orbits are also not excluded [6]. However, numerical analysis has suggested the presence of a Hopf bifurcation emerging through the positive steady-state, with the concentration of influent chlorophenol as the bifurcating parameter [14].

In this work, we extend the analysis of the model providing a proof of the existence, uniqueness and stability of six identified equilibria with the addition of only chlorophenol, and the more general case where all three substrates as external inputs to the system. The procedure we use allows to identify the sufficient conditions for the emergence of a Hopf bifurcation in those inflow concentration parameters. We are also able to prove that the dimensionless model is uniformly persistent, a new result for the system.

2 The model revisited

We first present concisely the original chemostat model using identical scaling to that given by [18]

$$\begin{cases} x'_0 &= -\alpha x_0 + \mu_0(s_0, s_2) x_0 - k_A x_0, \\ x'_1 &= -\alpha x_1 + \mu_1(s_1, s_2) x_1 - k_B x_1, \\ x'_2 &= -\alpha x_2 + \mu_2(s_2) x_2 - k_C x_2, \\ s'_0 &= \alpha(u_f - s_0) - \mu_0(s_0, s_2) x_0, \\ s'_1 &= \alpha(u_g - s_1) + \omega_0 \mu_0(s_0, s_2) x_0 - \mu_1(s_1, s_2) x_1, \\ s'_2 &= \alpha(u_h - s_2) - \omega_2 \mu_0(s_0, s_2) x_0 + \omega_1 \mu_1(s_1, s_2) x_1 - \mu_2(s_2) x_2, \end{cases} \quad (1)$$

with

$$\alpha = \frac{D}{k_{m, \text{ch}} Y_{\text{ch}}}, \quad (2)$$

$$u_f = \frac{S_{\text{ch}, \text{in}}}{K_{S, \text{ch}}}, \quad u_g = \frac{S_{\text{ph}, \text{in}}}{K_{S, \text{ph}}}, \quad u_h = \frac{S_{\text{H}_2, \text{in}}}{K_{S, \text{H}_2}}, \quad (3)$$

$$\omega_0 = \frac{K_{S, \text{ch}}}{K_{S, \text{ph}}} \frac{224}{208} (1 - Y_{\text{ch}}), \quad \omega_1 = \frac{K_{S, \text{ph}}}{K_{S, \text{H}_2}} \frac{32}{224} (1 - Y_{\text{ph}}), \quad \omega_2 = \frac{16}{208} \frac{K_{S, \text{ch}}}{K_{S, \text{H}_2}}, \quad (4)$$

$$\phi_1 = \frac{k_{m, \text{ph}} Y_{\text{ph}}}{k_{m, \text{ch}} Y_{\text{ch}}}, \quad \phi_2 = \frac{k_{m, \text{H}_2} Y_{\text{H}_2}}{k_{m, \text{ch}} Y_{\text{ch}}}, \quad (5)$$

$$K_P = \frac{K_{S, \text{H}_2, \text{c}}}{K_{S, \text{H}_2}}, \quad K_I = \frac{K_{S, \text{H}_2}}{K_{I, \text{H}_2}}, \quad (6)$$

$$k_A = \frac{k_{\text{dec}, \text{ch}}}{k_{m, \text{ch}} Y_{\text{ch}}}, \quad k_B = \frac{k_{\text{dec}, \text{ph}}}{k_{m, \text{ch}} Y_{\text{ch}}}, \quad k_C = \frac{k_{\text{dec}, \text{H}_2}}{k_{m, \text{ch}} Y_{\text{ch}}}. \quad (7)$$

$$\mu_0(s_0, s_2) = \frac{s_0}{1 + s_0} \frac{s_2}{K_P + s_2}, \quad \mu_1(s_1, s_2) = \frac{\phi_1 s_1}{1 + s_1} \frac{1}{1 + K_I s_2}, \quad \mu_2(s_2) = \frac{\phi_2 s_2}{1 + s_2}, \quad (8)$$

where α is the dilution rate, u_f, u_g, u_h are the chlorophenol, phenol and hydrogen inflow concentrations, respectively, and k_A, k_B, k_C are the decay (or maintenance) terms. These are scaled to be dimensionless, as are the other parameters using the scaling provided by [18]. Briefly, $k_{m, \cdot}$ are the specific growth rates, $K_{S, \cdot}$ are the half-saturation coefficients, Y_{\cdot} are the substrate yield coefficients, K_{I, H_2} is the kinetic inhibition constant of hydrogen on the phenol degrader, and $k_{\text{dec}, \cdot}$ are the unscaled decay terms. The numeric values indicate the stoichiometric coefficients given in terms of units of Chemical Oxygen Demand rather than molarity, as is common for environmental engineering models. The $\mu_n(\cdot)$, $n = 0, 1, 2$ are the species growth functions described by double Monod, Monod with product inhibition, and Monod kinetics, respectively. Subscripts $_{\text{ch}, \text{ph}, \text{H}_2}$ relate to the chlorophenol degrader, phenol degrader, and methanogen, respectively.

For numerical bifurcation analysis given in Section 4.4, we consider the same parameter values as provided in the original work, as shown in Table 1.

Parameters	Value
ω_0	0.1854
ω_1	1656.69
ω_2	163.08
ϕ_1	1.8875
ϕ_2	3.8113
K_P	0.04
K_I	7.1429

Table 1: Parameter regimes for the system (1).

3 Reduction of the model

We are able to obtain many theoretical results assuming general forms of the growth functions provided we assume the death rates of the microbial populations are insignificant compared to the dilution rate. We thus consider the following

system that is identical to system (1), except that we assume $k_i = 0$, $i \in \{A, B, C\}$:

$$\begin{cases} x'_0 &= -\alpha x_0 + \mu_0(s_0, s_2) x_0, \\ x'_1 &= -\alpha x_1 + \mu_1(s_1, s_2) x_1, \\ x'_2 &= -\alpha x_2 + \mu_2(s_2) x_2, \\ s'_0 &= \alpha(u_f - s_0) - \mu_0(s_0, s_2) x_0, \\ s'_1 &= \alpha(u_g - s_1) + \omega_0 \mu_0(s_0, s_2) x_0 - \mu_1(s_1, s_2) x_1, \\ s'_2 &= \alpha(u_h - s_2) - \omega_2 \mu_0(s_0, s_2) x_0 + \omega_1 \mu_1(s_1, s_2) x_1 - \mu_2(s_2) x_2, \end{cases} \quad (9)$$

$$x_i(0) \geq 0, \quad s_i(0) \geq 0, \quad i \in \{1, 2, 3\}.$$

We assume that $\mu_0(s_0, s_2)$, $\mu_1(s_1, s_2)$, $\mu_2(s_2)$ are \mathcal{C}^1 functions that satisfy the following general conditions:

- For all $s_0 \geq 0$ and $s_2 \geq 0$, $\mu_0(0, s_2) = 0$, $\mu_0(s_0, 0) = 0$. As a consequence, $\partial_{s_0} \mu_0(s_0, 0) = 0$, $\partial_{s_2} \mu_0(0, s_2) = 0$. Thus we assume that the chlorophenol degrader cannot grow in the absence of either chlorophenol or hydrogen;
- For all $s_0 > 0$ and $s_2 > 0$, $\partial_{s_0} \mu_0(s_0, s_2) > 0$, $\partial_{s_2} \mu_0(s_0, s_2) > 0$. Thus we assume that the chlorophenol degrader grows on both chlorophenol and hydrogen;
- For all $s_2 \geq 0$ and $s_1 \geq 0$, $\mu_1(0, s_2) = 0$, $\partial_{s_2} \mu_1(0, s_2) = 0$. Thus we assume that the phenol degrader cannot grow in the absence of phenol;
- For all $s_1 > 0$ and $s_2 > 0$, $\partial_{s_1} \mu_1(s_1, s_2) > 0$, $\partial_{s_2} \mu_1(s_1, s_2) < 0$. Thus we assume that the supply of phenol results in growth of the phenol degrader, and that hydrogen inhibits its growth;
- $\mu_2(0) = 0$ and $\mu'_2(s_2) > 0$ for all $s_2 > 0$. Thus we assume that the mathanogen cannot grow without the presence of hydrogen, and that increasing the supply of hydrogen results in faster growth of the methanogen.

We use the prototypes μ_0 , μ_1 , and μ_2 defined in (8), which satisfy these conditions, when we are able to prove results in general, and when providing numerical simulations or bifurcation diagrams.

We now prove a lemma that we will use to show global well-posedness of system (9).

Lemma 3.1. *All solutions of system (9) with positive initial conditions remain positive and bounded for all positive times.*

If $x_i(0) = 0$, $i \in \{1, 2, 3\}$, then $x_i(t) = 0$ for all $t \geq 0$.

Proof. Consider any solution $\vec{\varphi}(t)$ with positive initial conditions. By existence and uniqueness theory, there cannot be a time $\bar{t} > 0$, such that $x_i(\bar{t}) = 0$ for some $i \in \{1, 2, 3\}$, since then $x_i(t) \equiv 0$ for all $t \in \mathbb{R}$, contradicting $x_i(0) > 0$. Hence $x_i(t) > 0$ for all $t \geq 0$. Also, if $x_i(0) = 0$ for some $i \in \{1, 2, 3\}$, then there is a solution of system (9) with $x_i(t) \equiv 0$ for all $t \in \mathbb{R}$. By existence and uniqueness theory, this is the only solution.

Now, consider $\vec{\varphi}(t)$, and suppose that there is some $\bar{t} > 0$, such that $s_0(t) > 0$ for $t \in [0, \bar{t})$, $s_0(\bar{t}) = 0$, and $s_1(t), s_2(t) \geq 0$ for $t \in [0, \bar{t}]$. Then $s'_0(\bar{t}) \leq 0$. However, from system (9), $s'_0(\bar{t}) = \alpha u_f$. If $u_f > 0$, then $s'_0(\bar{t}) > 0$, a contradiction. If $u_f = 0$, then there is a solution of system (9) with $s_0(t) \equiv 0$, which contradicts uniqueness of solutions. It follows that $s_0(t) > 0$ for all $t \geq 0$.

Next, consider $\vec{\varphi}(t)$, and suppose that there is some $\bar{t} > 0$, such that $s_2(t) > 0$ for $t \in [0, \bar{t})$, $s_2(\bar{t}) = 0$, and $s_1(t) \geq 0$ for $t \in [0, \bar{t}]$. Then $s'_2(\bar{t}) \leq 0$. However, from system (9), $s'_2(\bar{t}) = \alpha u_h + \omega_1 \mu_1(s_1(\bar{t}), 0) x_1$. If $u_h > 0$, or $s_1(\bar{t}) > 0$, then $s'_2(\bar{t}) > 0$, a contradiction. If both $u_h = 0$, and $s_1(\bar{t}) = 0$, then $s'_2(\bar{t}) = 0$, and there is another solution with $s_2(t) \equiv 0$, which contradicts uniqueness of solutions to initial value problems. It follows that $s_2(t) > 0$ for all $t \geq 0$.

Finally, consider $\vec{\varphi}(t)$, and suppose that there is some $\bar{t} > 0$, such that $s_1(t) > 0$ for $t \in [0, \bar{t})$, $s_1(\bar{t}) = 0$. Then $s'_1(\bar{t}) \leq 0$. However, from system (9), $s'_1(\bar{t}) = \alpha u_g + \omega_0 \mu_0(s_0(\bar{t}), s_2(\bar{t})) x_0$, so $s'_1(\bar{t}) > 0$, a contradiction. It follows that $s_1(t) > 0$ for all $t \geq 0$.

We have thus proved the positivity of solutions and move on to showing the boundedness of solutions.

By adding the first and the fourth equations of (9), we obtain

$$x'_0 + s'_0 = -\alpha(x_0 + s_0 - u_f),$$

hence

$$(x_0 + s_0 - u_f)' = -\alpha(x_0 + s_0 - u_f),$$

which implies that

$$x_0(t) + s_0(t) = u_f + (x_0(0) + s_0(0) - u_f) e^{-\alpha t}. \quad (10)$$

Similarly, we obtain

$$x_1(t) + \omega_0 s_0(t) + s_1(t) = \omega_0 u_f + u_g + (x_1(0) + \omega_0 s_0(0) + s_1(0) - \omega_0 u_f - u_g) e^{-\alpha t}, \quad (11)$$

and

$$\begin{aligned} \omega_2 x_0(t) + x_2(t) + \omega_0 \omega_1 s_0(t) + \omega_1 s_1(t) + s_2(t) &= \omega_0 \omega_1 u_f + \omega_1 u_g + u_h + \\ &+ (\omega_0 x_0(0) + x_2(0) + \omega_0 \omega_1 s_0(0) + \omega_1 s_1(0) + s_2(0) - \omega_0 \omega_1 u_f - \omega_1 u_g - u_h) e^{-\alpha t}. \end{aligned} \quad (12)$$

Since all terms of the sums in (10), (11) and (12) are positive for all positive initial conditions, the solutions of (9) are bounded. Also, taking the limit as $t \rightarrow \infty$ in equations (10), (11), and (12) we obtain that

$$\lim_{t \rightarrow \infty} (x_0(t) + s_0(t)) = u_f, \quad (13)$$

$$\lim_{t \rightarrow \infty} (x_1(t) + \omega_0 s_0(t) + s_1(t)) = \omega_0 u_f + u_g, \quad (14)$$

$$\lim_{t \rightarrow \infty} (\omega_2 x_0(t) + x_2(t) + \omega_0 \omega_1 s_0(t) + \omega_1 s_1(t) + s_2(t)) = \omega_0 \omega_1 u_f + \omega_1 u_g + u_h. \quad (15)$$

Starting with any positive initial conditions, the solutions of system (9) eventually satisfy

$$x_0 + s_0 = u_f, \quad (16)$$

$$x_1 + \omega_0 s_0 + s_1 = \omega_0 u_f + u_g, \quad (17)$$

$$\omega_2 x_0 + x_2 + \omega_0 \omega_1 s_0 + \omega_1 s_1 + s_2 = \omega_0 \omega_1 u_f + \omega_1 u_g + u_h. \quad (18)$$

□

We call relations (16)-(18) "conservation principles". In other words, system (9) admits a positively invariant attracting set $\Omega \subset \mathbb{R}^6$, such that

$$\begin{aligned} \Omega = \{ & (x_0, x_1, x_2, s_0, s_1, s_2) \in \mathbb{R}^6 : x_i, s_i \geq 0, i = 0, 1, 2; \\ & x_0 + s_0 = u_f, \\ & x_1 + \omega_0 s_0 + s_1 = \omega_0 u_f + u_g, \\ & \omega_2 x_0 + x_2 + \omega_0 \omega_1 s_0 + \omega_1 s_1 + s_2 = \omega_0 \omega_1 u_f + \omega_1 u_g + u_h \}. \end{aligned} \quad (19)$$

Using the conservation principles we can compute s_0 , s_1 , and s_2 as functions of x_0 , x_1 , x_2

$$\begin{aligned} s_0 &= -x_0 + u_f, \\ s_1 &= \omega_0 x_0 - x_1 + u_g, \\ s_2 &= -\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h. \end{aligned} \quad (20)$$

Now, we can reduce the analysis of the original system (9) to the analysis of the following equivalent three-dimensional system on the invariant set Ω

$$\begin{cases} x'_0 = -\alpha x_0 + \mu_0 (-x_0 + u_f, -\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h) x_0, \\ x'_1 = -\alpha x_1 + \mu_1 (\omega_0 x_0 - x_1 + u_g, -\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h) x_1, \\ x'_2 = -\alpha x_2 + \mu_2 (-\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h) x_2. \end{cases} \quad (21)$$

From now on, we will study the reduced system (21). We begin by analyzing all the possible equilibria.

3.1 Equilibria of the reduced system and their local stability

The equilibria are found by setting the right hand sides of equations in (21) equal to zero. Below, we list all the possibilities obtained this way. Since equations (20) give a one-to-one correspondence of the equilibria of system (21) with the equilibria of system (9), we also list the corresponding steady states $(x_0, x_1, x_2, s_0, s_1, s_2)$ of the six-dimensional system in each case.

Types of equilibria of system (21):

- Zero equilibrium ${}_{(000)}\mathcal{E} = (0, 0, 0)$. The corresponding equilibrium ${}_{(000)}\mathbb{E}$ in the six-dimensional system:

$${}_{(000)}\mathbb{E} = (0, 0, 0, u_f, u_g, u_h). \quad (22)$$

In this case, all the populations die, hence the only source for the substrates comes from the inflow rates u_f , u_g , and u_h .

- Boundary equilibria:

- $_{(100)}\mathcal{E} = \left(_{(100)}x_0, 0, 0\right)$, where $x_0 = _{(100)}x_0 > 0$ is a solution (if it exists) of

$$\mu_0(-x_0 + u_f, -\omega_2 x_0 + u_h) = \alpha. \quad (23)$$

The corresponding equilibrium $_{(100)}\mathbb{E}$ in the six-dimensional system:

$$_{(100)}\mathbb{E} = \left(_{(100)}x_0, 0, 0, -_{(100)}x_0 + u_f, \omega_0 _{(100)}x_0 + u_g, -\omega_2 _{(100)}x_0 + u_h\right). \quad (24)$$

In this case, the only microorganism surviving is the chlorophenol degrader. It consumes the chlorophenol, hence the value of s_0 is given as the balance between this consumption, and the supply inflow u_f . Since x_0 produces phenol this value is added to u_g in the total phenol amount s_1 . Since x_0 consumes hydrogen as well, the value $\omega_2 _{(100)}x_0$ is subtracted from s_2 as well. This steady state is not desirable because of the phenol build-up in the system.

- $_{(010)}\mathcal{E} = \left(0, _{(010)}x_1, 0\right)$, where $x_1 = _{(010)}x_1 > 0$ is a solution (if it exists) of

$$\mu_1(-x_1 + u_g, \omega_1 x_1 + u_h) = \alpha. \quad (25)$$

The corresponding equilibrium $_{(010)}\mathbb{E}$ in the six-dimensional system:

$$_{(010)}\mathbb{E} = \left(0, _{(010)}x_1, 0, u_f, -_{(010)}x_1 + u_g, \omega_1 _{(010)}x_1 + u_h\right). \quad (26)$$

In this case, only the phenol degrader survives, and hence the value of s_1 at the equilibrium is equal to the balance between its consumption and inflow u_g . Chlorophenol is not being consumed, hence its total amount equals the inflow concentration u_f . Hydrogen is being produced by the phenol degrader, and also its value is increased by the inflow u_h .

- $_{(001)}\mathcal{E} = \left(0, 0, _{(001)}x_2\right)$, where $x_2 = _{(001)}x_2 > 0$ is a solution (if it exists) of

$$\mu_2(-x_2 + u_h) = \alpha. \quad (27)$$

The corresponding equilibrium $_{(001)}\mathbb{E}$ in the six-dimensional system:

$$_{(001)}\mathbb{E} = \left(0, 0, _{(001)}x_2, u_f, u_g, -_{(001)}x_2 + u_h\right). \quad (28)$$

Here, only the methanogen is present, hence the values of chlorophenol and phenol are equal the inflow concentrations u_f and u_g , respectively.

- $_{(101)}\mathcal{E} = \left(_{(101)}x_0, 0, -\omega_2 _{(101)}x_0 + u_h - \mu_2^{-1}(\alpha)\right)$, where $x_0 = _{(101)}x_0 > 0$ is a solution (if it exists) of

$$\mu_0(-x_0 + u_f, \mu_2^{-1}(\alpha)) = \alpha. \quad (29)$$

The corresponding equilibrium $_{(101)}\mathbb{E}$ in the six-dimensional system:

$$_{(101)}\mathbb{E} = \left(_{(101)}x_0, 0, -\omega_2 _{(101)}x_0 + u_h - \mu_2^{-1}(\alpha), -_{(101)}x_0 + u_f, \omega_0 _{(101)}x_0 + u_g, \mu_2^{(-1)}(\alpha)\right). \quad (30)$$

In this case, both chlorophenol degrader and methanogen are present. The lack of phenol degrader results in phenol build-up. We can also observe competition for hydrogen between the phenol degrader and methanogen.

- $_{(011)}\mathcal{E} = \left(0, _{(011)}x_1, \omega_1 _{(011)}x_1 + u_h - \mu_2^{-1}(\alpha)\right)$, where $x_1 = _{(011)}x_1 > 0$ is a solution (if it exists) of

$$\mu_1(-x_1 + u_g, \mu_2^{-1}(\alpha)) = \alpha. \quad (31)$$

The corresponding equilibrium $_{(011)}\mathbb{E}$ in the six-dimensional system:

$$_{(011)}\mathbb{E} = \left(0, _{(011)}x_1, \omega_1 _{(011)}x_1 + u_h - \mu_2^{-1}(\alpha), u_f, -_{(011)}x_1 + u_g, \mu_2^{-1}(\alpha)\right). \quad (32)$$

This steady state represents a two-tiered food chain, with the phenol degrader and methanogen present. Hydrogen has an inhibiting effect on the phenol degrader.

– $_{(110)}\mathcal{E} = \left(_{(110)}x_0, _{(110)}x_1, 0\right)$, where $x_0 = _{(110)}x_0 > 0$ and $x_1 = _{(110)}x_1 > 0$ are solutions of

$$\begin{aligned}\mu_0(-x_0 + u_f, -\omega_2 x_0 + \omega_1 x_1 + u_h) &= \alpha, \\ \mu_1(\omega_0 x_0 - x_1 + u_g, -\omega_2 x_0 + \omega_1 x_1 + u_h) &= \alpha.\end{aligned}\quad (33)$$

The corresponding equilibrium $_{(110)}\mathbb{E}$ in the six-dimensional system:

$$_{(110)}\mathbb{E} = \left(_{(110)}x_0, _{(110)}x_1, 0, -_{(110)}x_0 + u_f, \omega_0 _{(110)}x_0 - _{(110)}x_1 + u_g, -\omega_2 _{(110)}x_0 + \omega_1 _{(110)}x_1 + u_h\right). \quad (34)$$

In this case, both the chlorophenol and phenol degraders are present, however the methanogen is washed out. Thus, full mineralisation to methane is not possible and, hence, the hydrogen accumulates to some theoretical maximum, balanced such that the inhibitory effect on the phenol degrader does not induce washout, whilst providing enough hydrogen for chlorophenol degrader activity.

- Positive (interior) equilibrium $_{(111)}\mathcal{E} = \left(x_0^*, x_1^*, x_2^*\right)$, where $x_0 = x_0^* > 0$, $x_1 = x_1^* > 0$, and $x_2 = x_2^* > 0$ are solutions of

$$\begin{aligned}\mu_0(-x_0 + u_f, -\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h) &= \alpha, \\ \mu_1(\omega_0 x_0 - x_1 + u_g, -\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h) &= \alpha, \\ \mu_2(-\omega_2 x_0 + \omega_1 x_1 - x_2 + u_h) &= \alpha.\end{aligned}\quad (35)$$

The corresponding equilibrium $_{(111)}\mathbb{E}$ in the six-dimensional system:

$$_{(111)}\mathbb{E} = \left(x_0^*, x_1^*, x_2^*, -x_0^* + u_f, \omega_0 x_0^* - x_1^* + u_g, \mu_2^{(-1)}(\alpha)\right). \quad (36)$$

Here, all species are present, and thus we observe full chlorophenol mineralisation. For this reason, asymptotic stability of this equilibrium is the desired operational state.

As it is not clear whether the listed equations have solutions, and if the solutions are unique, we now derive conditions on the parameters that address these questions.

3.1.1 Existence and uniqueness

Since there are many parameters in system (21), it was not possible to obtain explicit expressions for some of the equilibria. We did however simplify the computation by only looking for equilibria in the invariant set Ω . This assumption is reasonable, since the dynamics of the original system reduces to the dynamics on the set Ω .

- $_{(000)}\mathcal{E} = (0, 0, 0)$ equilibrium always exists.
- $_{(100)}\mathcal{E} = \left(_{(100)}x_0, 0, 0\right)$. As mentioned at the beginning of subsection 3.1.1, we are looking for the equilibria in the feasible set Ω , i.e., where all of the components in corresponding six-dimensional equilibria are nonnegative. Thus, we want $x_0 = _{(100)}x_0$ to satisfy $_{(100)}x_0 \in (0, u_f]$, and $_{(100)}x_0 \leq \frac{u_h}{\omega_2}$; hence we consider only $x_0 \in \left(0, \min\left(u_f, \frac{u_h}{\omega_2}\right)\right]$. For such x_0 the differentiable mapping $x_0 \mapsto \mu_0(u_f - x_0, -\omega_2 x_0 + u_h)$ is decreasing, and thus $_{(100)}\mathcal{E}$ exists in Ω if and only if $\alpha \in [0, \mu_0(u_f, u_h))$, and when it exists, it is unique.
- $_{(010)}\mathcal{E} = \left(0, _{(010)}x_1, 0\right)$. By a similar argument, we consider $x_1 \in (0, u_g]$. The differentiable mapping $x_1 \mapsto \mu_1(u_g - x_1, \omega_1 x_1 + u_h)$ is decreasing, so $_{(010)}\mathcal{E}$ exists in Ω if and only if $\alpha \in [0, \mu_1(u_g, u_h))$, and when it exists, it is unique.
- $_{(001)}\mathcal{E} = \left(0, 0, _{(001)}x_2\right)$. Once again, we consider only $x_2 \in (0, u_h]$, for which the differentiable mapping $x_2 \mapsto \mu_2(-x_2 + u_h)$ is decreasing, so $_{(001)}\mathcal{E}$ exists in Ω if and only if $\alpha \in [0, \mu_2(u_h))$, and when it exists, it is unique (notice that if $\mu_2(s_2) = \frac{\phi_2 s_2}{1+s_2}$, we have $_{(001)}x_2 = u_h - \frac{\alpha}{\phi_2 - \alpha}$).
- $_{(101)}\mathcal{E} = \left(_{(101)}x_0, 0, -\omega_2 _{(101)}x_0 + u_h - \mu_2^{-1}(\alpha)\right)$. For $_{(101)}x_2 = -\omega_2 _{(101)}x_0 + u_h - \mu_2^{-1}(\alpha)$, the restriction $_{(101)}x_2 > 0$ gives us the condition $_{(101)}x_0 < \frac{u_h - \mu_2^{-1}(\alpha)}{\omega_2}$ (notice that this already implies that $_{(101)}x_0 \leq \frac{u_h}{\omega_2}$).

is satisfied), and thus the requirement ${}_{(101)}x_0 > 0$ results in the first condition on α , i.e., $\alpha < \mu_2(u_h)$. We also require that ${}_{(101)}x_0 \leq u_f$. We therefore only consider the mapping $x_0 \mapsto \mu_0(-x_0 + u_f, \mu_2^{-1}(\alpha))$ for $x_0 \in \left(0, \min\left(u_f, \frac{u_h - \mu_2^{-1}(\alpha)}{\omega_2}\right)\right)$. This differentiable mapping is decreasing, so ${}_{(101)}\mathcal{E}$ exists in Ω if and only if

$$\alpha \in \left(\mu_0\left(u_f - \min\left(u_f, \frac{u_h - \mu_2^{-1}(\alpha)}{\omega_2}\right), \mu_2^{-1}(\alpha)\right), \mu_0(u_f, \mu_2^{-1}(\alpha))\right), \quad (37)$$

and $\alpha < \mu_2(u_h)$, and when it exists, it is unique.

By solving the equation

$$\mu_0(-x_0 + u_f, \mu_2^{-1}(\alpha)) = \alpha, \quad (38)$$

we obtain the following explicit formulas for ${}_{(101)}x_0$ and ${}_{(101)}x_2$ with our test prototypes μ_0, μ_1 , and μ_2 :

$${}_{(101)}x_0 = \frac{\alpha(1 + u_f)(K_P + \mu_2^{-1}(\alpha)) - u_f\mu_2^{-1}(\alpha)}{\alpha(K_P + \mu_2^{-1}(\alpha)) - \mu_2^{-1}(\alpha)}, \quad (39)$$

$${}_{(101)}x_2 = \omega_2 \frac{u_f\mu_2^{-1}(\alpha) - \alpha(1 + u_f)(K_P + \mu_2^{-1}(\alpha))}{\alpha(K_P + \mu_2^{-1}(\alpha)) - \mu_2^{-1}(\alpha)} + u_h - \mu_2^{-1}(\alpha). \quad (40)$$

- ${}_{(011)}\mathcal{E} = \left(0, {}_{(011)}x_1, \omega_1 {}_{(011)}x_1 + u_h - \mu_2^{-1}(\alpha)\right)$. For $x_2 = \omega_1 {}_{(011)}x_1 + u_h - \mu_2^{-1}(\alpha)$ the restriction ${}_{(011)}x_2 > 0$ gives the condition $x_1 > \frac{\mu_2^{-1}(\alpha) - u_h}{\omega_1}$, so the requirement ${}_{(011)}x_1 \leq u_g$ results in the first condition on α , i.e., $\alpha < \mu_2(\omega_1 u_g + u_h)$ (notice that this already implies that $\omega_1 {}_{(011)}x_1 + u_h \geq 0$ is satisfied).

We therefore only consider the mapping $x_1 \mapsto \mu_1(-x_1 + u_g, \mu_2^{-1}(\alpha))$ for $\left(\max\left(0, \frac{\mu_2^{-1}(\alpha) - u_h}{\omega_1}\right), u_g\right]$.

For such x_1 , this differentiable mapping is decreasing, so ${}_{(011)}\mathcal{E}$ exists in Ω if and only if $\alpha \in \left[0, \mu_1\left(u_g - \max\left(0, \frac{\mu_2^{-1}(\alpha) - u_h}{\omega_1}\right), \mu_2^{-1}(\alpha)\right)\right)$ and $\alpha < \mu_2(\omega_1 u_g + u_h)$, and when it exists, it is unique.

By solving the equation

$$\mu_1(-x_1 + u_g, \mu_2^{-1}(\alpha)) = \alpha, \quad (41)$$

for the prototypes given by (8), we obtain the following explicit formulas for ${}_{(011)}x_1$ and ${}_{(011)}x_2$

$${}_{(011)}x_1 = \frac{\alpha(1 + u_g)(1 + K_I\mu_2^{-1}(\alpha)) - u_g\phi_1}{\alpha(1 + K_I\mu_2^{-1}(\alpha)) - \phi_1}, \quad (42)$$

$${}_{(011)}x_2 = \omega_1 \frac{\alpha(1 + u_g)(1 + K_I\mu_2^{-1}(\alpha)) - u_g\phi_1}{\alpha(1 + K_I\mu_2^{-1}(\alpha)) - \phi_1} + u_h - \mu_2^{-1}(\alpha). \quad (43)$$

- ${}_{(110)}\mathcal{E} = \left({}_{(110)}x_0, {}_{(110)}x_1, 0\right)$. This case is much more complicated since we cannot explicitly compute x_0 as a function of x_1 , or x_1 as a function of x_0 in the same way as in the previous cases. In this case more than one equilibrium of the form ${}_{(110)}\mathcal{E}$ can exist. In the case of the growth functions defined in (8), it was proved in [14] that if $u_g = u_h = 0$, then there exist at most two equilibria of this form. By using the specific growth functions (8), the equilibria, given as positive solutions of the following system of equations

$$\begin{aligned} \frac{-x_0 + u_f}{1 - x_0 + u_f} \frac{-\omega_2 x_0 + \omega_1 x_1 + u_h}{K_P - \omega_2 x_0 + \omega_1 x_1 + u_h} &= \alpha, \\ \phi_1 \frac{\omega_0 x_0 - x_1 + u_g}{1 + \omega_0 x_0 - x_1 + u_g} \frac{1}{1 + K_I \omega_0 x_0 - x_1 + u_g} &= \alpha, \end{aligned} \quad (44)$$

must also satisfy $x_0 < u_f$ and $\max\left(0, \frac{\omega_2 x_0 - u_h}{\omega_1}\right) < x_1 < u_g + \omega_0 x_0$. Notice that the first equation in (44) is linear in x_1 , hence we can compute it as a function of x_0 , and substitute this expression into the second equation of (44), obtaining a fourth order polynomial in x_0 . Each zero of this polynomial, together with the corresponding value of x_1 , which satisfies the aforementioned conditions, will constitute an equilibrium ${}_{(110)}\mathcal{E}$ of system (21). Since the polynomial in x_0 is of order four, and x_1 is given as a function of x_0 , we can have at most four equilibria of the form ${}_{(110)}\mathcal{E}$.

- $_{(111)}\mathcal{E} = \left(\begin{smallmatrix} * \\ x_0, x_1, x_2 \end{smallmatrix} \right)$. For the interior equilibrium, we have to consider two cases, depending on the sign of $\omega_2 u_f - u_h$, since the bounds on the values of x_1 are different in each case. We are looking for solutions of system (35), for which x_0 , x_1 , and x_2 satisfy

$$\begin{aligned} x_0 &\in (0, u_f], \\ x_1 &\in \left(\max \left(0, \frac{\omega_2 x_0 - u_h}{\omega_1} \right), \omega_0 x_0 + u_g \right], \\ x_2 &\in (0, -\omega_2 x_0 + \omega_1 x_1 + u_h], \end{aligned} \quad (45)$$

if $\omega_2 u_f - u_h > 0$, and

$$\begin{aligned} x_0 &\in (0, u_f], \\ x_1 &\in (0, \omega_0 x_0 + u_g], \\ x_2 &\in (0, -\omega_2 x_0 + \omega_1 x_1 + u_h], \end{aligned} \quad (46)$$

if $\omega_2 u_f - u_h < 0$. In both cases, if we let $x_2 = -\omega_2 x_0 + \omega_1 x_1 + u_h - \mu_2^{-1}(\alpha)$ (which immediately gives us a necessary condition $\alpha < \sup_{s_2 \geq 0} \mu_2(\alpha)$), we obtain the following system for x_0 and x_1

$$\begin{aligned} \mu_0(-x_0 + u_f, \mu_2^{-1}(\alpha)) &= \alpha, \\ \mu_1(\omega_0 x_0 - x_1 + u_g, \mu_2^{-1}(\alpha)) &= \alpha. \end{aligned} \quad (47)$$

For $x_0 \in (0, u_f]$ the differentiable mapping $x_0 \mapsto \mu_0(-x_0 + u_f, \mu_2^{-1}(\alpha))$ is decreasing, so $x_0 = x_0^*$ exists if and only if $\alpha \in [0, \mu_0(u_f, \mu_2^{-1}(\alpha))]$ and when this value exists, it is unique. Now consider

$$\mu_1(\omega_0 x_0^* - x_1 + u_g, \mu_2^{-1}(\alpha)) = \alpha. \quad (48)$$

We have two cases

- $\omega_2 u_f - u_h > 0$. For $x_1 \in \left(\max \left(0, \frac{\omega_2 x_0^* - u_h}{\omega_1} \right), \omega_0 x_0^* + u_g \right]$ the differentiable mapping $x_1 \mapsto \mu_1(\omega_0 x_0^* - x_1 + u_g, \mu_2^{-1}(\alpha))$ is decreasing, so $x_1 = x_1^*$ exists if and only if $\alpha \in \left[0, \mu_1 \left(\omega_0 x_0^* - \max \left(0, \frac{\omega_2 x_0^* - u_h}{\omega_1} \right) + u_g, \mu_2^{-1}(\alpha) \right) \right]$ and when this value exists, it is unique.
- $\omega_2 u_f - u_h < 0$. Similarly, by considering $x_1 \in (0, \omega_0 x_0^* + u_g]$ it follows that $x_1 = x_1^*$ exists if and only if $\alpha \in [0, \mu_1(\omega_0 x_0^* + u_g, \mu_2^{-1}(\alpha))]$.

Having x_0^* and x_1^* defined, we let $x_2^* = -\omega_2 x_0^* + \omega_1 x_1^* + u_h - \mu_2^{-1}(\alpha)$. In order to have $x_2^* > 0$ we need $\alpha < \mu_2(-\omega_2 x_0^* + \omega_1 x_1^* + u_h)$; hence $_{(111)}\mathcal{E}$ exists in Ω if and only if

$$\begin{aligned} \alpha \in \left[0, \min \left(\mu_0(u_f, \mu_2^{-1}(\alpha)), \right. \right. \\ \left. \mu_1 \left(\omega_0 x_0^* - \max \left(0, \frac{\omega_2 x_0^* - u_h}{\omega_1} \right) + u_g, \mu_2^{-1}(\alpha) \right), \right. \\ \left. \left. \mu_2(-\omega_2 x_0^* + \omega_1 x_1^* + u_h) \right) \right], \end{aligned} \quad (49)$$

in the $\omega_2 u_f - u_h > 0$ case, and

$$\alpha \in \left[0, \min \left(\mu_0(u_f, \mu_2^{-1}(\alpha)), \mu_1(\omega_0 x_0^* + u_g, \mu_2^{-1}(\alpha)), \mu_2(-\omega_2 x_0^* + \omega_1 x_1^* + u_h) \right) \right], \quad (50)$$

in the $\omega_2 u_f - u_h < 0$ case. Although the conditions on α are implicit and very complicated, we now know that if $_{(111)}\mathcal{E}$ exists, it is unique. With the growth functions defined in (8), we can solve the equations (35)

explicitly and obtain the following formulas for the interior equilibrium

$$x_0^* = 1 + u_f + \frac{1}{K_P(\phi_2 - \alpha) + \alpha - 1}, \quad (51)$$

$$x_1^* = \omega_0 x_0^* + u_g + 1 + \frac{\phi_1}{\alpha \left(1 + K_I \frac{\alpha}{\phi_2 - \alpha}\right) - \phi_1}, \quad (52)$$

$$x_2^* = -\omega_2 x_0^* + \omega_1 x_1^* + u_h - \frac{\alpha}{\phi_2 - \alpha}. \quad (53)$$

3.1.2 Local stability results

We now study the local stability of the equilibria by considering the eigenvalues of the Jacobian evaluated at each equilibrium.

The Jacobian J for system (21) evaluated at (x_0, x_1, x_2) has the following form

$$J = \begin{bmatrix} \mu_0 - \alpha + x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2}\right) & \omega_1 x_0 \frac{\partial \mu_0}{\partial s_2} & -x_0 \frac{\partial \mu_0}{\partial s_2} \\ x_1 \left(\omega_0 \frac{\partial \mu_1}{\partial s_1} - \omega_2 \frac{\partial \mu_1}{\partial s_2}\right) & \mu_1 - \alpha + x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2}\right) & -x_1 \frac{\partial \mu_1}{\partial s_2} \\ -\omega_2 x_2 \mu_2' & \omega_1 x_2 \mu_2' & \mu_2 - \alpha - x_2 \mu_2' \end{bmatrix}. \quad (54)$$

- For the zero equilibrium ${}_{(000)}\mathcal{E}$, the corresponding Jacobian ${}_{(000)}J$ has the following form

$${}_{(000)}J = \begin{bmatrix} \mu_0 - \alpha & 0 & 0 \\ 0 & \mu_1 - \alpha & 0 \\ 0 & 0 & \mu_2 - \alpha \end{bmatrix}, \quad (55)$$

and its eigenvalues are $\lambda_1 = \mu_0(u_f, u_h) - \alpha$, $\lambda_2 = \mu_1(u_g, u_h) - \alpha$, $\lambda_3 = \mu_2(u_h) - \alpha$. This implies that if

- $\alpha > \max(\mu_0(u_f, u_h), \mu_1(u_g, u_h), \mu_2(u_h))$, then ${}_{(000)}\mathcal{E}$ is a stable node,
- $\min(\mu_0(u_f, u_h), \mu_1(u_g, u_h), \mu_2(u_h)) < \alpha < \max(\mu_0(u_f, u_h), \mu_1(u_g, u_h), \mu_2(u_h))$, then ${}_{(000)}\mathcal{E}$ is a saddle point,
- $\alpha < \min(\mu_0(u_f, u_h), \mu_1(u_g, u_h), \mu_2(u_h))$, then ${}_{(000)}\mathcal{E}$ is an unstable node.
- For the boundary equilibrium ${}_{(100)}\mathcal{E}$, the corresponding Jacobian ${}_{(100)}J$ has the following form

$${}_{(100)}J = \begin{bmatrix} {}_{(100)}x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2}\right) & \omega_1 {}_{(100)}x_0 \frac{\partial \mu_0}{\partial s_2} & -{}_{(100)}x_0 \frac{\partial \mu_0}{\partial s_2} \\ 0 & \mu_1 - \alpha & 0 \\ 0 & 0 & \mu_2 - \alpha \end{bmatrix}, \quad (56)$$

and its eigenvalues are $\lambda_1 = {}_{(100)}x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2}\right) < 0$,

$\lambda_2 = \mu_1(\omega_0 {}_{(100)}x_0 + u_g, -\omega_2 {}_{(100)}x_0 + u_h) - \alpha$, and $\lambda_3 = \mu_2(-\omega_2 {}_{(100)}x_0 + u_h) - \alpha$. Hence if

- $\alpha > \max(\mu_1(\omega_0 {}_{(100)}x_0 + u_g, -\omega_2 {}_{(100)}x_0 + u_h), \mu_2(-\omega_2 {}_{(100)}x_0 + u_h))$, then ${}_{(100)}\mathcal{E}$ is a stable node,
- $\alpha < \max(\mu_1(\omega_0 {}_{(100)}x_0 + u_g, -\omega_2 {}_{(100)}x_0 + u_h), \mu_2(-\omega_2 {}_{(100)}x_0 + u_h))$, then ${}_{(100)}\mathcal{E}$ is a saddle point.
- For the boundary equilibrium ${}_{(010)}\mathcal{E}$, the corresponding Jacobian ${}_{(010)}J$ has the following form

$${}_{(010)}J = \begin{bmatrix} \mu_0 - \alpha & 0 & 0 \\ {}_{(010)}x_1 \left(\omega_0 \frac{\partial \mu_1}{\partial s_1} - \omega_2 \frac{\partial \mu_1}{\partial s_2}\right) & {}_{(010)}x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2}\right) & -{}_{(010)}x_1 \frac{\partial \mu_1}{\partial s_2} \\ 0 & 0 & \mu_2 - \alpha \end{bmatrix}, \quad (57)$$

and its eigenvalues are $\lambda_1 = \mu_0(u_f, \omega_1 {}_{(010)}x_1 + u_h) - \alpha$, $\lambda_2 = {}_{(010)}x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2}\right) < 0$, and

$\lambda_3 = \mu_2(\omega_1 {}_{(010)}x_1 + u_h) - \alpha$. Hence if

- $\alpha > \max \left(\mu_0 \left(u_f, \omega_{1(010)} x_1 + u_h \right), \mu_2 \left(\omega_{1(010)} x_1 + u_h \right) \right)$, then ${}_{(010)}\mathcal{E}$ is a stable node,
- $\alpha < \max \left(\mu_0 \left(u_f, \omega_{1(010)} x_1 + u_h \right), \mu_2 \left(\omega_{1(010)} x_1 + u_h \right) \right)$, then ${}_{(010)}\mathcal{E}$ is a saddle point.
- For the boundary equilibrium ${}_{(001)}\mathcal{E}$, the corresponding Jacobian ${}_{(001)}J$ has the following form

$${}_{(001)}J = \begin{bmatrix} \mu_0 - \alpha & 0 & 0 \\ 0 & \mu_1 - \alpha & 0 \\ -\omega_{2(001)} x_2 \mu'_2 & \omega_{1(001)} x_2 \mu'_2 & \mu_2 - \alpha - {}_{(001)}x_2 \mu'_2 \end{bmatrix}, \quad (58)$$

and its eigenvalues are $\lambda_1 = \mu_0 \left(u_f, -{}_{(001)}x_2 + u_h \right) - \alpha$, $\lambda_2 = \mu_1 \left(u_g, -{}_{(001)}x_2 + u_h \right) - \alpha$, and $\lambda_3 = -{}_{(001)}x_2 \mu'_2 \left(-{}_{(001)}x_2 + u_h \right) < 0$. Hence if

- $\alpha > \max \left(\mu_0 \left(u_f, -{}_{(001)}x_2 + u_h \right), \mu_1 \left(u_g, -{}_{(001)}x_2 + u_h \right) \right)$, then ${}_{(001)}\mathcal{E}$ is a stable node,
- $\alpha < \max \left(\mu_0 \left(u_f, -{}_{(001)}x_2 + u_h \right), \mu_1 \left(u_g, -{}_{(001)}x_2 + u_h \right) \right)$, then ${}_{(001)}\mathcal{E}$ is a saddle point.
- For the boundary equilibrium ${}_{(101)}\mathcal{E}$, the corresponding Jacobian ${}_{(101)}J$ has the following form

$${}_{(101)}J = \begin{bmatrix} {}_{(101)}x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) & \omega_{1(101)} x_0 \frac{\partial \mu_0}{\partial s_2} & -{}_{(101)}x_0 \frac{\partial \mu_0}{\partial s_2} \\ 0 & \mu_1 - \alpha & 0 \\ -\omega_{2(101)} x_2 \mu'_2 & \omega_{1(101)} x_2 \mu'_2 & -{}_{(101)}x_2 \mu'_2 \end{bmatrix}, \quad (59)$$

where ${}_{(101)}x_2 = -\omega_{2(101)} x_0 + u_h - \mu_2^{-1}(\alpha)$. We immediately obtain one of the eigenvalues $\lambda_1 = \mu_1 \left(\omega_{0(101)} x_0 + u_g, \mu_2^{-1}(\alpha) \right) - \alpha$. The other two eigenvalues are given as the solutions of the following quadratic equation

$$\lambda^2 + a_1 \lambda + a_0 = 0, \quad (60)$$

where

$$a_1 = {}_{(101)}x_0 \left(\frac{\partial \mu_0}{\partial s_0} + \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) + {}_{(101)}x_2 \mu'_2 \quad (61)$$

$$a_0 = {}_{(101)}x_0 {}_{(101)}x_2 \frac{\partial \mu_0}{\partial s_0} \mu'_2. \quad (62)$$

Since both $a_1 > 0$, and $a_0 > 0$, by the Routh-Hurwitz criterion, all roots of the equation (60) have negative real parts. Hence if

- $\alpha > \mu_1 \left(\omega_{0(101)} x_0 + u_g, \mu_2^{-1}(\alpha) \right)$, then ${}_{(101)}\mathcal{E}$ is asymptotically stable,
- $\alpha < \mu_1 \left(\omega_{0(101)} x_0 + u_g, \mu_2^{-1}(\alpha) \right)$, then ${}_{(101)}\mathcal{E}$ is a saddle point.
- For the boundary equilibrium ${}_{(011)}\mathcal{E}$, the corresponding Jacobian ${}_{(011)}J$ has the following form

$${}_{(011)}J = \begin{bmatrix} \mu_0 - \alpha & 0 & 0 \\ {}_{(011)}x_1 \left(\omega_0 \frac{\partial \mu_1}{\partial s_1} - \omega_2 \frac{\partial \mu_1}{\partial s_2} \right) & {}_{(011)}x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) & -{}_{(011)}x_1 \frac{\partial \mu_1}{\partial s_2} \\ -\omega_{2(011)} x_2 \mu'_2 & \omega_{1(011)} x_2 \mu'_2 & -{}_{(011)}x_2 \mu'_2 \end{bmatrix}, \quad (63)$$

where ${}_{(011)}x_2 = \omega_{1(011)} x_1 + u_h - \mu_2^{-1}(\alpha)$. We immediately obtain one of the eigenvalues $\lambda_1 = \mu_0 \left(u_f, \mu_2^{-1}(\alpha) \right) - \alpha$. The other two eigenvalues are given as the solutions of the following quadratic equation

$$\lambda^2 + a_1 \lambda + a_0 = 0, \quad (64)$$

where

$$a_1 = {}_{(011)}x_1 \left(\frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) + {}_{(011)}x_2 \mu'_2, \quad (65)$$

$$a_0 = {}_{(011)}x_1 {}_{(011)}x_2 \frac{\partial \mu_1}{\partial s_1} \mu'_2. \quad (66)$$

Since both $a_1 > 0$, and $a_0 > 0$, by the Routh-Hurwitz criterion, all roots of the equation (64) have negative real parts. Hence if

- $\alpha > \mu_0(u_f, \mu_2^{-1}(\alpha))$, then ${}_{(011)}\mathcal{E}$ is asymptotically stable,
- $\alpha < \mu_0(u_f, \mu_2^{-1}(\alpha))$, then ${}_{(011)}\mathcal{E}$ is a saddle point.

- For the boundary equilibrium ${}_{(110)}\mathcal{E}$, the corresponding Jacobian ${}_{(110)}J$ has the following form

$${}_{(110)}J = \begin{bmatrix} {}_{(110)}x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) & \omega_1 {}_{(110)}x_0 \frac{\partial \mu_0}{\partial s_2} & - {}_{(110)}x_0 \frac{\partial \mu_0}{\partial s_2} \\ {}_{(110)}x_1 \left(\omega_0 \frac{\partial \mu_1}{\partial s_1} - \omega_2 \frac{\partial \mu_1}{\partial s_2} \right) & {}_{(110)}x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) & - {}_{(110)}x_1 \frac{\partial \mu_1}{\partial s_2} \\ 0 & 0 & \mu_2 - \alpha \end{bmatrix}. \quad (67)$$

We immediately obtain one eigenvalue $\lambda_1 = \mu_2 - \alpha$. The other two eigenvalues are solutions of the following quadratic equation

$$\lambda^2 + a_1 \lambda + a_0 = 0, \quad (68)$$

where

$$a_1 = {}_{(110)}x_0 \left(\frac{\partial \mu_0}{\partial s_0} + \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) + {}_{(110)}x_1 \left(\frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_1}{\partial s_2} \right), \quad (69)$$

$$a_0 = {}_{(110)}x_0 {}_{(110)}x_1 \left(\frac{\partial \mu_0}{\partial s_0} + \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) \left(\frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) - \omega_1 {}_{(110)}x_0 {}_{(110)}x_1 \frac{\partial \mu_0}{\partial s_2} \left(\omega_0 \frac{\partial \mu_1}{\partial s_1} - \omega_2 \frac{\partial \mu_1}{\partial s_2} \right). \quad (70)$$

We have $a_1 > 0$, and

$$a_0 > 0 \iff \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_2} + (\omega_2 - \omega_0 \omega_1) \frac{\partial \mu_0}{\partial s_2} \frac{\partial \mu_1}{\partial s_1} > 0. \quad (71)$$

Hence if

- $\alpha > \mu_2$ and $\frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_2} + (\omega_2 - \omega_0 \omega_1) \frac{\partial \mu_0}{\partial s_2} \frac{\partial \mu_1}{\partial s_1} > 0$ (where all the functions are evaluated at the steady state), then ${}_{(110)}\mathcal{E}$ is asymptotically stable,
- $\alpha < \mu_2$ or $\frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_2} + (\omega_2 - \omega_0 \omega_1) \frac{\partial \mu_0}{\partial s_2} \frac{\partial \mu_1}{\partial s_1} < 0$, then ${}_{(110)}\mathcal{E}$ is unstable.

- For the interior equilibrium ${}_{(111)}\mathcal{E}$, the corresponding Jacobian ${}_{(111)}J$ has the following form

$${}_{(111)}J = \begin{bmatrix} {}^*x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) & \omega_1 {}^*x_0 \frac{\partial \mu_0}{\partial s_2} & - {}^*x_0 \frac{\partial \mu_0}{\partial s_2} \\ {}^*x_1 \left(\omega_0 \frac{\partial \mu_1}{\partial s_1} - \omega_2 \frac{\partial \mu_1}{\partial s_2} \right) & {}^*x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) & - {}^*x_1 \frac{\partial \mu_1}{\partial s_2} \\ -\omega_2 {}^*x_2 \mu'_2 & \omega_1 {}^*x_2 \mu'_2 & - {}^*x_2 \mu'_2 \end{bmatrix}, \quad (72)$$

and its eigenvalues are solutions of the following cubic equation

$$\lambda^3 + a_2 \lambda^2 + a_1 \lambda + a_0 = 0, \quad (73)$$

where

$$a_2 = - {}^*x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) - {}^*x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) + {}^*x_2 \mu'_2, \quad (74)$$

$$a_1 = {}^*x_1 \frac{\partial \mu_1}{\partial s_1} \left({}^*x_0 \frac{\partial \mu_0}{\partial s_0} - (\omega_0 \omega_1 - \omega_2) {}^*x_0 \frac{\partial \mu_0}{\partial s_2} + {}^*x_2 \mu'_2 \right) + {}^*x_0 \frac{\partial \mu_0}{\partial s_0} \left(-\omega_1 {}^*x_1 \frac{\partial \mu_1}{\partial s_2} + {}^*x_2 \mu'_2 \right), \quad (75)$$

$$a_0 = {}^*x_0 {}^*x_1 {}^*x_2 \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} \mu'_2. \quad (76)$$

By the Routh-Hurwitz criterion, all eigenvalues have negative real parts if and only if $a_2 > 0$, $a_0 > 0$, and $a_2 a_1 > a_0$. We have $a_2 > 0$, and $a_0 > 0$ always, and

$$a_2 a_1 > a_0 \iff \left[- {}^*x_0 \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) - {}^*x_1 \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) + {}^*x_2 \mu'_2 \right] \cdot \left[{}^*x_1 \frac{\partial \mu_1}{\partial s_1} \left({}^*x_0 \frac{\partial \mu_0}{\partial s_0} - (\omega_0 \omega_1 - \omega_2) {}^*x_0 \frac{\partial \mu_0}{\partial s_2} + {}^*x_2 \mu'_2 \right) + {}^*x_0 \frac{\partial \mu_0}{\partial s_0} \left(-\omega_1 {}^*x_1 \frac{\partial \mu_1}{\partial s_2} + {}^*x_2 \mu'_2 \right) \right] - {}^*x_0 {}^*x_1 {}^*x_2 \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} \mu'_2 > 0. \quad (77)$$

Thus ${}_{(111)}\mathcal{E}$ is asymptotically stable if $a_2 a_1 > a_0$, and is unstable if $a_2 a_1 < a_0$.

We gather the results concerning the existence and local stability of the equilibria of (21) in table 2. All the functions in the "Local stability" column are evaluated at the corresponding steady states, and all the symbols are given by equations (78)-(84). The conditions for existence of the ${}_{(110)}\mathcal{E}$ equilibrium are discussed in subsection 3.1.1.

Equilibrium	Existence	Local stability
${}_{(000)}\mathcal{E}$	Always	$\alpha > \max(\mu_0, \mu_1, \mu_2)$
${}_{(100)}\mathcal{E}$	$\alpha < \mu_0(u_f, u_h)$	$\alpha > \max(\mu_1, \mu_2)$
${}_{(010)}\mathcal{E}$	$\alpha < \mu_1(u_g, u_h)$	$\alpha > \max(\mu_0, \mu_2)$
${}_{(001)}\mathcal{E}$	$\alpha < \mu_2(u_h)$	$\alpha > \max(\mu_0, \mu_1)$
${}_{(101)}\mathcal{E}$	${}_{(101)}\star$	$\alpha > \mu_1$
${}_{(011)}\mathcal{E}$	${}_{(011)}\star$	$\alpha > \mu_0$
${}_{(110)}\mathcal{E}$	-	$\alpha > \mu_2$ and ${}_{(110)}a_0 > 0$
${}_{(111)}\mathcal{E}$	${}_{(111)}\star$	${}_{(111)}a_2 {}_{(111)}a_1 - {}_{(111)}a_0 > 0$,

Table 2: Equilibria of system (21) together with their local stability.

$$_{(101)}\star = \alpha \in \left(\mu_0 \left(u_f - \min \left(u_f, \frac{u_h - \mu_2^{-1}(\alpha)}{\omega_2} \right), \mu_2^{-1}(\alpha) \right), \mu_0(u_f, \mu_2^{-1}(\alpha)) \right), \quad (78)$$

$$_{(011)}\star = \alpha \in \left[0, \mu_1 \left(u_g - \max \left(0, \frac{\mu_2^{-1}(\alpha) - u_h}{\omega_1} \right), \mu_2^{-1}(\alpha) \right) \right), \quad (79)$$

$$\text{and } \alpha < \mu_2(\omega_1 u_g + u_h),$$

$$_{(110)}a_0 = \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} - \omega_1 \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_2} + (\omega_2 - \omega_0 \omega_1) \frac{\partial \mu_0}{\partial s_2} \frac{\partial \mu_1}{\partial s_1}, \quad (80)$$

$$_{(111)}\star = \alpha \in \left[0, \min \left(\mu_0(u_f, \mu_2^{-1}(\alpha)), \mu_1 \left(\omega_0 x_0 - \max \left(0, \frac{\omega_2^* x_0 - u_h}{\omega_1} \right) + u_g, \mu_2^{-1}(\alpha) \right), \mu_2(-\omega_2^* x_0 + \omega_1^* x_1 + u_h) \right) \right), \quad (81)$$

$$_{(111)}a_2 = -x_0^* \left(-\frac{\partial \mu_0}{\partial s_0} - \omega_2 \frac{\partial \mu_0}{\partial s_2} \right) - x_1^* \left(-\frac{\partial \mu_1}{\partial s_1} + \omega_1 \frac{\partial \mu_1}{\partial s_2} \right) + x_2^* \mu_2', \quad (82)$$

$$_{(111)}a_1 = x_1^* \frac{\partial \mu_1}{\partial s_1} \left(x_0^* \frac{\partial \mu_0}{\partial s_0} - (\omega_0 \omega_1 - \omega_2) x_0^* \frac{\partial \mu_0}{\partial s_2} + x_2^* \mu_2' \right) + x_0^* \frac{\partial \mu_0}{\partial s_0} \left(-\omega_1 x_1^* \frac{\partial \mu_1}{\partial s_2} + x_2^* \mu_2' \right), \quad (83)$$

$$_{(111)}a_0 = x_0^* x_1^* x_2^* \frac{\partial \mu_0}{\partial s_0} \frac{\partial \mu_1}{\partial s_1} \mu_2'. \quad (84)$$

4 Analysis of the full system

4.1 Periodic orbits on the faces

We begin by ruling out possibility of having a periodic orbit in one of the invariant faces of Ω . This can be done by using general forms of prototypes μ_0 , μ_1 and μ_2 .

First, consider system (21) on the part of Ω with $x_0 = 0$, i.e.,

$$\begin{cases} x_1' = (-\alpha + \mu_1(-x_1 + u_g, \omega_1 x_1 - x_2 + u_h)) x_1, \\ x_2' = (-\alpha + \mu_2(\omega_1 x_1 - x_2 + u_h)) x_2. \end{cases} \quad (85)$$

The domain for system (85) is given by the following set Ω_{12} :

$$\Omega_{12} = \{(x_1, x_2) \in \mathbb{R}^2 : 0 \leq x_1 \leq u_g, 0 \leq x_2 \leq u_h + \omega_1 x_1\}. \quad (86)$$

Notice that no periodic orbit can intersect the axes $x_1 = 0$ or $x_2 = 0$ since they are invariant. Now, let us define an auxiliary function

$$\varphi_0(x_1, x_2) = \frac{1}{x_1 x_2}, \quad (x_1, x_2) \in \Omega_{12} \setminus (\{x_1 = 0\} \cup \{x_2 = 0\}). \quad (87)$$

Then

$$\nabla \cdot (\varphi_0(-\alpha + \mu_1) x_1, \varphi_0(-\alpha + \mu_2) x_2) = \frac{-\partial_{s_1} \mu_1 + \omega_1 \partial_{s_2} \mu_1}{x_2} - \frac{\mu_2'}{x_1} < 0 \quad (88)$$

for all (x_1, x_2) in the domain of φ_0 . Thus, by the Dulac's Criterion [9], there are no periodic orbits in the $x_1 x_2$ face.

Now, consider system (21) on the part of Ω with $x_1 = 0$, i.e.,

$$\begin{cases} x_0' = (-\alpha + \mu_0(-x_0 + u_f, -\omega_2 x_0 - x_2 + u_h)) x_0, \\ x_2' = (-\alpha + \mu_2(-\omega_2 x_0 - x_2 + u_h)) x_2, \end{cases} \quad (89)$$

defined on Ω_{02} given by

$$\Omega_{02} = \left\{ (x_0, x_2) \in \mathbb{R}^2 : 0 \leq x_0 \leq \min \left\{ u_f, \frac{u_h - x_2}{\omega_2} \right\}, 0 \leq x_2 \leq -\omega_2 x_0 + u_h \right\}. \quad (90)$$

For auxiliary function

$$\varphi_1(x_0, x_2) = \frac{1}{x_0 x_2}, \quad (x_0, x_2) \in \Omega_{02} \setminus (\{x_0 = 0\} \cup \{x_2 = 0\}) \quad (91)$$

we have

$$\nabla \cdot (\varphi_1(-\alpha + \mu_1)x_0, \varphi_1(-\alpha + \mu_2)x_2) = \frac{-\partial_{s_0}\mu_0 - \omega_2\partial_{s_2}\mu_0}{x_2} - \frac{\mu'_2}{x_0} < 0. \quad (92)$$

This, together with the fact that the axes $x_0 = 0$, $x_2 = 0$ are invariant, shows that there are no periodic orbits in the $x_0 x_2$ face.

Finally, consider system (21) on the part of Ω with $x_2 = 0$, i.e.,

$$\begin{cases} x'_0 = (-\alpha + \mu_0(-x_0 + u_f, -\omega_2 x_0 + \omega_1 x_1 + u_h))x_0, \\ x'_1 = (-\alpha + \mu_1(\omega_0 x_0 - x_1 + u_g, -\omega_2 x_0 + \omega_1 x_1 + u_h))x_1. \end{cases} \quad (93)$$

defined on Ω_{01} given by

$$\Omega_{01} = \left\{ (x_0, x_1) \in \mathbb{R}^2 : 0 \leq x_0 \leq u_f, \max \left\{ 0, \frac{\omega_2 x_0 - u_h}{\omega_1} \right\} \leq x_1 \leq \omega_0 x_0 + u_g \right\}. \quad (94)$$

Analogously to the previous cases, we define

$$\varphi_2(x_0, x_1) = \frac{1}{x_0 x_1}, \quad (x_0, x_1) \in \Omega_{01} \setminus (\{x_0 = 0\} \cup \{x_1 = 0\}) \quad (95)$$

and compute

$$\nabla \cdot (\varphi(-\alpha + \mu_0)x_0, \varphi(-\alpha + \mu_1)x_1) = \frac{-\partial_{s_0}\mu_0 - \omega_2\partial_{s_2}\mu_0}{x_1} + \frac{-\partial_{s_1}\mu_1 + \omega_1\partial_{s_2}\mu_1}{x_0} < 0. \quad (96)$$

Since the axes $x_0 = 0$, $x_1 = 0$ are invariant, by Dulac's Criterion, there are no periodic orbits in the $x_0 x_1$ face.

4.2 Hopf Bifurcation

In this work we are especially interested in developing a more systematic approach to studying the Hopf bifurcation of the interior equilibrium. That Hopf bifurcation that occurs in this model was previously observed numerically [14]. The occurrence of a stable periodic orbit in system (21) represents a situation in which all three populations of microorganisms oscillate indefinitely, and as a consequence, the substrate concentrations fluctuate. The characteristic polynomial of the Jacobian ${}_{(111)}J$ corresponding to the interior equilibrium is given by

$$\lambda^3 + a_2\lambda^2 + a_1\lambda + a_0 = 0, \quad (97)$$

where

$$a_2 = -x_0^* \left(-\frac{\partial\mu_0}{\partial s_0} - \omega_2 \frac{\partial\mu_0}{\partial s_2} \right) - x_1^* \left(-\frac{\partial\mu_1}{\partial s_1} + \omega_1 \frac{\partial\mu_1}{\partial s_2} \right) + x_2^* \mu'_2, \quad (98)$$

$$a_1 = x_1^* \frac{\partial\mu_1}{\partial s_1} \left(x_0^* \frac{\partial\mu_0}{\partial s_0} - (\omega_0\omega_1 - \omega_2) x_0^* \frac{\partial\mu_0}{\partial s_2} + x_2^* \mu'_2 \right) + x_0^* \frac{\partial\mu_0}{\partial s_0} \left(-\omega_1 x_1^* \frac{\partial\mu_1}{\partial s_2} + x_2^* \mu'_2 \right), \quad (99)$$

$$a_0 = x_0^* x_1^* x_2^* \frac{\partial\mu_0}{\partial s_0} \frac{\partial\mu_1}{\partial s_1} \mu'_2, \quad (100)$$

and the coefficients a_2 , a_1 , and a_0 depend on the parameters u_f , u_g , u_h , and α . The coefficients a_2 and a_0 are sign-definite (they are both positive), and a_1 might possibly change sign. Let us first notice that since the polynomial has order three, a real eigenvalue always exists. By the Routh-Hurwitz criterion, the above polynomial has a pair of purely imaginary eigenvalues if and only if

$$a_2 a_1 = a_0 \quad (\text{which implies } a_1 > 0). \quad (101)$$

In that case we also have

$$\lambda^3 + a_2 \lambda^2 + a_1 \lambda + a_0 = \lambda^3 + a_2 \lambda^2 + a_1 \lambda + a_2 a_1 = (\lambda + a_2)(\lambda^2 + a_1). \quad (102)$$

Hence the eigenvalues are $\lambda_1 = -a_2$ and $\lambda_{2,3} = \pm \sqrt{a_1} i$. Since eigenvalues are continuous functions of the parameters, we can see that if there is some $(u_f, u_g, u_h, \alpha) = (u_f^*, u_g^*, u_h^*, \alpha^*)$ such that $a_2 a_1 = a_0$, then if we denote by λ_1 the always present real eigenvalue, there is some $\delta > 0$ such that for $\|(u_f, u_g, u_h, \alpha) - (u_f^*, u_g^*, u_h^*, \alpha^*)\| < \delta$ we always have $\lambda_1 < 0$. By lemma 5.1, section 5.2 in [9], this implies the existence of a parameter-dependent, smooth, attracting, two dimensional, center manifold $W_{(u_f, u_g, u_h, \alpha)}^c$. In the following analysis, we consider only parameters that are in the δ -neighborhood of (u_f, u_g, u_h, α) in order to ensure that the real eigenvalue λ_1 is negative.

By the Routh-Hurwitz criterion (since we just shown that when ${}_{(111)}\mathcal{E}$ exists that a_0 and a_2 are always positive), a Hopf bifurcation occurs when the expression $a_2 a_1 - a_0$ changes sign as a parameter varies. This ensures that the real part of a pair of complex eigenvalues with nonzero imaginary part passes through 0 and hence changes sign. This is related to the transversality condition: the derivative of the real part of the eigenvalue with respect to the bifurcation parameter evaluated at the critical value when the real parts are zero is non-zero. We check this condition for specific forms of the functions μ_0 , μ_1 , and μ_2 . For the prototypes proposed in (8), and with the values of parameters from Table 1 fixed, the function

$$f(u_f, u_g, u_h, \alpha) = a_2 a_1 - a_0 \quad (103)$$

is an algebraic function in u_f , u_g , u_h , and α . Hence fixing all the parameters except one makes the function f a polynomial, the order of which depends on the choice of the free parameter. Specifically if we choose:

- α - free parameter $\implies f$ has order 27,
- u_f - free parameter $\implies f$ has order 3,
- u_g - free parameter $\implies f$ has order 3,
- u_h - free parameter $\implies f$ has order 2.

We construct bifurcation diagrams to explore the possibility of Hopf bifurcations as α varies in subsection 4.4, and begin our theoretical analysis by choosing u_f as the free parameter. We have

$$f_{u_f}(u_f) = a_2 a_1 - a_0 = b_3 u_f^3 + b_2 u_f^2 + b_1 u_f + b_0, \quad (104)$$

and we assume that there is a value $u_f = u_f^*$ such that $f_{u_f}(u_f^*) = 0$. We want to find conditions on the coefficients of f_{u_f} that guarantee that the derivative of $a_2 a_1 - a_0$ with respect to u_f is not equal to zero when $a_2 a_1 - a_0 = 0$, i.e., that u_f^* is not a local extremum of f_{u_f} . We have

$$f'_{u_f}(u_f) = 3b_3 u_f^2 + 2b_2 u_f + b_1. \quad (105)$$

The necessary condition for u_f^* to be a local extremum for f_{u_f} is $f'_{u_f}(u_f^*) = 0$, that is

$$u_f^* = \frac{-b_2 \pm \sqrt{b_2^2 - 3b_1 b_3}}{3b_3}. \quad (106)$$

We can derive sufficient conditions for u_f^* to be an extremum (for example by computing the second derivative of f_{u_f}), but the condition (106) is already very restrictive and will be sufficient for our work. We have thus obtained a sufficient condition for a Hopf bifurcation.

If we choose u_g as the free parameter, we have

$$f_{u_g}(u_h) = c_3 u_g^3 + c_2 u_g^2 + c_1 u_g + c_0, \quad (107)$$

and with the assumption that $f_{u_g}(u_g^*) = 0$, by a similar analysis as in the previous case, we obtain an analogous sufficient condition for a Hopf bifurcation in u_g .

Finally, if we choose u_h as the free parameter, we have

$$f_{u_h}(u_h) = d_2 u_h^2 + d_1 u_h + d_0. \quad (108)$$

Once again, we assume that $f_{u_h}(u_h^*) = 0$, that is

$$u_h^* = \frac{-d_1 \pm \sqrt{d_1^2 - 4d_2 d_0}}{2d_2}. \quad (109)$$

Here, u_h^* is the local extremum if and only if the discriminant of equation (108) is zero, i.e., if $d_1^2 - 4d_2 d_0 = 0$.

We summarize our results in the following theorem.

Theorem 1. *Consider system (21) with the prototypes given by (8) and with the values of parameters from Table 1 fixed. Assume that there exists a point $(u_f, u_g, u_h, \alpha) = (u_f^*, u_g^*, u_h^*, \alpha^*)$ such that $f(u_f^*, u_g^*, u_h^*, \alpha^*) = 0$ for f defined in (103). Then $f_{u_f}(u_f) = f(u_f, u_g^*, u_h^*, \alpha^*)$, $f_{u_g}(u_g) = f(u_f^*, u_g, u_h^*, \alpha^*)$, and $f(u_h) = f(u_f^*, u_g^*, u_h, \alpha^*)$ are given by the equations (104), (107), and (108), respectively. Also, there exists $\delta > 0$ such that if $\|(u_f, u_g, u_h, \alpha) - (u_f^*, u_g^*, u_h^*, \alpha^*)\| < \delta$, then*

I. if

$$u_f^* \neq \frac{-b_2 \pm \sqrt{b_2^2 - 3b_1 b_3}}{3b_3}, \quad (110)$$

then there is a Hopf bifurcation in u_f at $u_f = u_f^*$,

II. if

$$u_g^* \neq \frac{-c_2 \pm \sqrt{c_2^2 - 3c_1 c_3}}{3c_3}, \quad (111)$$

then there is a Hopf bifurcation in u_g at $u_g = u_g^*$,

III. if

$$d_1^2 - 4d_2 d_0 \neq 0, \quad (112)$$

then there is a Hopf bifurcation in u_h at $u_h = u_h^*$.

We now illustrate the theoretical results with the numerical simulations. To approximate values of the equilibria we used Maple software [11], rounding all the values to 6 significant digits. For the choice of parameters given in Table 1 and

$$\alpha = 0.01, \quad u_f = 0.5, \quad u_g = 0.0006, \quad (113)$$

it follows that the only zero of (108) occurs for $u_h = 0.102520$. In Figure 1 we plot the phase space for $u_h = 0.05$ (just before the Hopf bifurcation) using the ode15s solver from [12]. For this set of parameters, we have the following approximate values of the equilibria

$$\begin{aligned} {}_{(000)}\mathcal{E} &= (0, 0, 0) \quad (\text{unstable}), \\ {}_{(100)}\mathcal{E} &= (0.000299015, 0, 0) \quad (\text{stable}), \\ {}_{(001)}\mathcal{E} &= (0, 0, 0.0473693) \quad (\text{unstable}), \\ {}_{(110)}\mathcal{E}^{(1)} &= (0.0545964, 0.00534486, 0) \quad (\text{unstable}), \\ {}_{(110)}\mathcal{E}^{(2)} &= (0.489465, 0.0487183, 0) \quad (\text{unstable}), \\ {}_{(111)}\mathcal{E} &= (0.306611, 0.0520205, 36.2277) \quad (\text{unstable}), \end{aligned} \quad (114)$$




Figure 1: Phase space of system (21) with $\alpha = 0.01$, $u_f = 0.5$, $u_g = 0.0006$, and $u_h = 0.05$.

We can see that there exists a stable periodic orbit in the system, but depending on the initial conditions, the solution might also converge to the boundary equilibrium $_{(100)}\mathcal{E}$. Thus in this case we observe bistability.

We now repeat the simulations for $u_h = 0.3$ (after the predicted Hopf bifurcation), presented in Figure 2. With all the other parameters set to the same values as in the previous case, we have the following equilibria

$$\begin{aligned}
 _{(000)}\mathcal{E} &= (0, 0, 0) \quad (\text{unstable}), \\
 _{(100)}\mathcal{E} &= (0.00183202, 0, 0) \quad (\text{stable}), \\
 _{(001)}\mathcal{E} &= (0, 0, 0.297369) \quad (\text{unstable}), \\
 _{(110)}\mathcal{E}^{(1)} &= (0.0528596, 0.00502299, 0) \quad (\text{unstable}), \\
 _{(110)}\mathcal{E}^{(2)} &= (0.489467, 0.0485697, 0) \quad (\text{unstable}), \\
 _{(111)}\mathcal{E} &= (0.306611, 0.0520205, 36.4777) \quad (\text{stable}).
 \end{aligned} \tag{115}$$




Figure 2: Phase space of system (21) with $\alpha = 0.01$, $u_f = 0.5$, $u_g = 0.0006$, and $u_h = 0.3$.

We can see that the Hopf bifurcation occurs between $u_h = 0.05$ and $u_h = 0.3$. The stable periodic orbit is no longer present, and the interior equilibrium is stable. Once again, the boundary equilibrium $_{(100)}\mathcal{E}$ is stable, and thus we observe bistability in the system.

It is worth to notice, that increasing u_h had a stabilizing effect on the system (actually, this type of behaviour applies also to u_f and u_g). This result is especially important in the context of the modeled phenomenon, since the most desirable situation happens when the production of methane is not fluctuating. Variable rates of gas production can result in decreased productivity of the biogas plant.

4.3 Persistence

The notion of persistence is particularly important in modeling biological phenomena. Roughly speaking, we say that a system is persistent if all the species with positive initial populations survive. The formal definition is as follows.

Definition 4.1. The system

$$\begin{aligned} x'_i &= x_i f_i(x_1, x_2, \dots, x_n), \\ x_i(0) &= x_{i0} \geq 0, \quad i = 1, 2, \dots, n, \end{aligned} \quad (116)$$

is said to be weakly persistent if

$$\limsup_{t \rightarrow \infty} x_i(t) > 0, \quad i = 1, 2, \dots, n \quad (117)$$

for every trajectory with positive initial conditions, and is said to be persistent if

$$\liminf_{t \rightarrow \infty} x_i(t) > 0, \quad i = 1, 2, \dots, n \quad (118)$$

for every trajectory with positive initial conditions. This system is said to be uniformly persistent if there exists a positive number ϵ such that

$$\liminf_{t \rightarrow \infty} x_i(t) \geq \epsilon, \quad i = 1, 2, \dots, n \quad (119)$$

for every trajectory with positive initial conditions.

To prove that system (21) is persistent, we will use the Butler-McGehee lemma [16] repeatedly.

Lemma 4.1. *Suppose that x^* is a hyperbolic equilibrium point of the system*

$$\begin{aligned} x' &= f(x), \\ x(0) &= x_0, \end{aligned} \tag{120}$$

with $x \in \mathbb{R}^n$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, where f is continuously differentiable. Suppose also that x^ is in $\omega(x_0)$, the omega limit set of $\gamma^+(x_0)$ (the positive semi-orbit through x_0), but is not the entire omega limit set. Then $\omega(x_0)$ has nontrivial (i.e., different from x^*) intersection with the stable and unstable manifolds of x^* .*

As we already noticed in section 4.2, there are values of the parameters where one of the boundary equilibria and the interior equilibrium are both asymptotically stable, and hence system (21) is not persistent, even though an interior equilibrium point exists. We will thus focus on the cases for which no boundary equilibrium point of system (21) is stable.

Theorem 2. *Let system (21) have the following equilibria configuration (as represented schematically in Figure 3):*

<i>Equilibrium</i>	<i>Number of eigenvalues with positive real part</i>	<i>Number of eigenvalues with negative real part</i>
$(000)\mathcal{E}$	2	1
$(100)\mathcal{E}$	1	2
$(001)\mathcal{E}$	1	2
$(110)\mathcal{E}$	1	2

Then system (21) is persistent.




Figure 3: Schematic representation of the equilibria configuration occurring in the hypothesis of theorem 2. Black arrows represent stable and unstable manifolds of each of the equilibrium (marked by the orange dots). In the example for the parameters we select in model (21) to illustrate this theorem (see (121)), there is an asymptotically stable interior equilibrium (as shown). However, this is not necessary for the proof of theorem 2.

Proof. Since planes x_0x_1 , x_0x_2 and x_1x_2 are invariant, we know where the stable and unstable manifolds of the boundary equilibria lie. This is represented in a schematic way in Figure 3. Keeping this picture in mind should make the following argument much more transparent. Assume that a solution $\vec{x}(t) = (x_0(t), x_1(t), x_2(t))$ with an initial condition $\vec{x}^{(0)} = (x_0^{(0)}, x_1^{(0)}, x_2^{(0)})$, where $x_i^{(0)} > 0$, $i = 1, 2, 3$, is given. First, suppose that ${}_{(000)}\mathcal{E}$ belongs to $\omega(\gamma^+(\vec{x}^{(0)}))$, the omega limit set of $\gamma^+(\vec{x}^{(0)})$. Since ${}_{(000)}\mathcal{E}$ is a saddle point with one-dimensional stable manifold restricted to the x_1 -axis, it is not the entire omega limit set $\omega(\gamma^+(\vec{x}^{(0)}))$. Hence, by lemma 4.1, there is a point $x^* \neq {}_{(000)}\mathcal{E}$ in both $\omega(\gamma^+(\vec{x}^{(0)}))$ and $W^s({}_{(000)}\mathcal{E})$, the stable manifold of ${}_{(000)}\mathcal{E}$. The entire orbit through any point in an omega limit set is also in the omega limit set. The stable manifold of ${}_{(000)}\mathcal{E}$ is the x_1 -axis, and the x_1 -axis is unbounded. We have already proven in section 3 that all orbits of system (21) are bounded, and hence the omega limit set of any orbit of (21) is bounded. This contradicts the existence of such an x^* and thus ${}_{(000)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$.

Now, suppose that ${}_{(001)}\mathcal{E} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Since ${}_{(001)}\mathcal{E}$ is a saddle point with two-dimensional stable manifold restricted to the x_0x_1 -plane, $\{{}_{(001)}\mathcal{E}\}$ is not the entire omega limit set $\omega(\gamma^+(\vec{x}^{(0)}))$. Thus, using lemma 4.1, there is a point $x^* \in \omega(\gamma^+(\vec{x}^{(0)})) \cap W^s({}_{(001)}\mathcal{E}) \setminus \{{}_{(001)}\mathcal{E}\}$. Since the stable manifold $W^s({}_{(001)}\mathcal{E})$ lies entirely in the x_1x_2 -plane, and the entire orbit through x^* is in $\omega(\gamma^+(\vec{x}^{(0)}))$, by the analysis in subsection 4.1, this orbit becomes unbounded in backward time. This contradiction shows that ${}_{(001)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$.

Now, suppose that ${}_{(100)}\mathcal{E} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Similarly as in the previous cases, this implies that there exists a point $x^* \in \omega(\gamma^+(\vec{x}^{(0)})) \cap W^s({}_{(100)}\mathcal{E}) \setminus \{{}_{(100)}\mathcal{E}\}$. This time the stable manifold $W^s({}_{(100)}\mathcal{E})$ is two-dimensional and

lies entirely in the x_0x_2 -plane. By the analysis in subsection 4.1, the entire orbit through x^* (which belongs to $\omega(\gamma^+(\vec{x}^{(0)}))$) becomes unbounded in backward time or its closure contains ${}_{(001)}\mathcal{E}$. This contradiction proves that ${}_{(100)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$.

Now, suppose that ${}_{(110)}\mathcal{E} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Again, $\{{}_{(110)}\mathcal{E}\}$ is not the entire omega limit set $\omega(\gamma^+(\vec{x}^{(0)}))$, so there exists a point $x^* \in \omega(\gamma^+(\vec{x}^{(0)})) \cap W^s({}_{(110)}\mathcal{E}) \setminus \{{}_{(110)}\mathcal{E}\}$. This point lies in the x_0x_1 -plane, since $W^s({}_{(110)}\mathcal{E})$ is two-dimensional and is entirely contained in this plane. As in the previous cases, the entire orbit through x^* is in $\omega(\gamma^+(\vec{x}^{(0)}))$. Since there are no periodic orbits in the x_0x_1 face, and since $\{{}_{(100)}\mathcal{E}\} \notin \omega(\gamma^+(\vec{x}^{(0)}))$, the orbit becomes unbounded in backward time. This contradiction proves that ${}_{(110)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$.

Finally, consider any $\hat{x} = (\hat{x}_0, \hat{x}_1, \hat{x}_2)$, such that $\hat{x}_i = 0$ for at least one $i = 1, 2, 3$, and suppose that $\hat{x} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Then, the entire orbit through \hat{x} is in $\omega(\gamma^+(\vec{x}^{(0)}))$. But since this orbit lies entirely in either x_0x_1 , x_1x_2 , or x_0x_2 face, it converges to one of the boundary equilibria. This implies that this boundary equilibrium is in $\omega(\gamma^+(\vec{x}^{(0)}))$, and this possibility has been eliminated in the previous part of the proof.

We have therefore proven that

$$\liminf_{t \rightarrow \infty} x_i(t) > 0, \quad i = 1, 2, 3,$$

i.e., that system (21) is persistent. □

An example satisfying the assumptions of theorem 2 occurs for

$$\alpha = 0.0002, \quad u_f = 0.6, \quad u_g = 0, \quad u_h = 0.1. \quad (121)$$

Persistence can also be observed with the addition of phenol, i.e., with $u_g > 0$.

Theorem 3. *Let system (21) have the following equilibria configuration (as represented schematically in Figure 4):*

Equilibrium	Number of eigenvalues with positive real part	Number of eigenvalues with negative real part
${}_{(000)}\mathcal{E}$	2	1
${}_{(100)}\mathcal{E}$	1	2
${}_{(001)}\mathcal{E}$	1	2
${}_{(011)}\mathcal{E}$	1	2
${}_{(110)}\mathcal{E}$	1	2

Then system (21) is persistent.




Figure 4: Schematic representation of the equilibria configuration occurring in the hypothesis of theorem 3. Black arrows represent stable and unstable manifolds of each of the equilibrium (marked by the orange dots). In the example for the parameters we select in model (21) to illustrate this theorem (see (122)), there is an asymptotically stable interior equilibrium (as shown). However, this is not necessary for the proof of theorem 3.

Proof. The idea behind this proof is very similar to the method presented in the proof of theorem 2. Let $\vec{x}(t) = (x_0(t), x_1(t), x_2(t))$ be a solution of (21) with an initial condition $\vec{x}^{(0)} = (x_0^{(0)}, x_1^{(0)}, x_2^{(0)})$, where $x_i^{(0)} > 0$, $i = 1, 2, 3$. Since the stable and unstable manifolds of all of the equilibria, except $_{(001)}\mathcal{E}$ and $_{(011)}\mathcal{E}$, have the same configuration as in the hypothesis of theorem 2, the argument eliminating them from the omega limit set of $\gamma^+(\vec{x}^{(0)})$ is exactly the same and we only need to focus on $_{(001)}\mathcal{E}$ and $_{(011)}\mathcal{E}$ equilibria.

Suppose that $_{(001)}\mathcal{E} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Since $_{(001)}\mathcal{E}$ is a saddle point with one-dimensional stable manifold restricted to the x_2 -axis, we have $\omega(\gamma^+(\vec{x}^{(0)})) \setminus \{_{(001)}\mathcal{E}\} \neq \emptyset$. Hence, by lemma 4.1, there is a point $x^* \in \omega(\gamma^+(\vec{x}^{(0)})) \cap W^s(_{(001)}\mathcal{E}) \setminus \{_{(001)}\mathcal{E}\}$. The entire orbit through x^* , which also belongs to $\omega(\gamma^+(\vec{x}^{(0)}))$, either becomes unbounded in backward time, or converges to the $_{(000)}\mathcal{E}$ equilibrium. Since all orbits of system (21) are bounded, and $_{(000)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$, we obtain a contradiction. Hence $_{(001)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$.

Now, suppose that $_{(011)}\mathcal{E} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Since $_{(011)}\mathcal{E}$ is a saddle point with two-dimensional stable manifold restricted to the x_1x_2 -plane (it is repelling into the interior), we have $\omega(\gamma^+(\vec{x}^{(0)})) \setminus \{_{(011)}\mathcal{E}\} \neq \emptyset$. By using lemma 4.1, there exists a point $x^* \in \omega(\gamma^+(\vec{x}^{(0)})) \cap W^s(_{(011)}\mathcal{E}) \setminus \{_{(011)}\mathcal{E}\}$. The entire orbit through x^* , which also belongs to $\omega(\gamma^+(\vec{x}^{(0)}))$, either becomes unbounded in backward time, or converges to $_{(000)}\mathcal{E}$, or $_{(001)}\mathcal{E}$ (we have previously shown in subsection 4.1 that there are no periodic orbits in the x_1x_2 face). Since we have already

proven that ${}_{(000)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$, and ${}_{(001)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$, we obtain a contradiction, which proves that ${}_{(011)}\mathcal{E} \notin \omega(\gamma^+(\vec{x}^{(0)}))$.

Finally, consider any $\hat{x} = (\hat{x}_0, \hat{x}_1, \hat{x}_2)$, such that $\hat{x}_i = 0$ for at least one $i = 1, 2, 3$, and suppose that $\hat{x} \in \omega(\gamma^+(\vec{x}^{(0)}))$. Then, the entire orbit through \hat{x} is in $\omega(\gamma^+(\vec{x}^{(0)}))$. But since this orbit lies entirely in either x_0x_1 , x_1x_2 , or x_0x_2 face, it converges to one of the boundary equilibria. This implies that this boundary equilibrium is in $\omega(\gamma^+(\vec{x}^{(0)}))$, and this possibility has been eliminated in the previous part of the proof. \square

An example satisfying the assumptions of theorem 3 occurs for

$$\alpha = 0.0002, \quad u_f = 0.6, \quad u_g = 0.00015, \quad u_h = 0.1. \quad (122)$$

Remark 1. Interestingly enough, in many cases of models describing biological phenomena, persistence already implies uniform persistence. The rigorous results were obtained in [2]. In our context, the key theorem from [2] states that if \mathcal{F} is a dynamical system for which \mathbb{R}_+^n and $\partial\mathbb{R}_+^n$ are invariant, then \mathcal{F} is uniformly persistent provided that

1. \mathcal{F} is dissipative (meaning that $\forall x \in \mathbb{R}_+^n \omega(x) \neq \emptyset$ and $\bigcup_{x \in \mathbb{R}_+^n} \omega(x)$ has compact closure),
2. \mathcal{F} is weakly persistent,
3. $\partial\mathcal{F}$ (the restriction of \mathcal{F} to the boundary $\partial\mathbb{R}_+^n$) is "isolated",
4. $\partial\mathcal{F}$ is "acyclic".

These results can be easily modified so that we consider the flow \mathcal{F} on Ω defined in (19). Although $\partial\Omega$ is not invariant, the theorem from [2], as explained in [3], can be modified so that it applies in the case when $\partial\Omega$ is the union of two sets Ω_1 and Ω_2 , for which \mathcal{F} is invariant on Ω_1 and Ω_2 is repelling into the interior of Ω , provided that conditions 3. and 4. are satisfied for the restriction of \mathcal{F} to Ω_1 . In our case, the positively invariant set Ω , on which we analyze system (21) is bounded, hence condition 1. is satisfied. Condition 2. holds by theorem 2 (persistence implies weak persistence). In our context condition 3. is satisfied, because all the boundary equilibria are hyperbolic, and hence each one is the maximal invariant set in a neighbourhood of itself. Also, their union forms a covering of the omega limit sets of Ω_1 . Condition 4. is satisfied because the boundary equilibria are not cyclically linked, i.e., there is no cyclic chain of heteroclinic orbits joining them. Thus, we have shown not only persistence, but also uniform persistence of system (21) in the case of theorem 2 and theorem 3.

We have thus proven the following theorem:

Theorem 4. *Under the hypotheses of theorems 2 and 3, system (21) is uniformly persistent.*

We finish this subsection by extending the uniform persistence to the original six dimensional system (1). Notice that if $\vec{X}_0 = (x_0(0), x_1(0), x_2(0), s_0(0), s_1(0), s_2(0))$ with $x_i(0) \geq 0$, $s_i(0) \geq 0$, $i = 1, 2, 3$, then we necessarily must have $\omega(\vec{X}_0) \in \Omega$. Otherwise, there would exist a point $\vec{p} \in \mathbb{R}_+ \setminus \Omega$ and a sequence of times (t_n) with $t_n \rightarrow \infty$ for which the corresponding solution converges to \vec{p} . This would mean that Ω is not globally attracting, which was proven in section 3. Also, if $\omega(\vec{X}_0)$ has a point lying in a face with one of the x_i , $i = 1, 2, 3$ equal zero, then the entire orbit through that point would also be in $\omega(\vec{X}_0)$. Thus, if the assumptions of either theorem 2 or theorem 3 hold, the omega limit set $\omega(\vec{X}_0)$ is entirely contained in the interior of Ω . We have thus proven the following theorem:

Theorem 5. *Under the hypotheses of theorems 2 and 3, the six dimensional system (1) is uniformly persistent.*

4.4 Bifurcation diagrams

As previously stated in section 4.2, we now study numerically effects on the qualitative behaviour of system (21) when considering α as the bifurcation parameter. Throughout this section, we assume that parameters $\omega_0, \omega_1, \omega_2, \phi_1, \phi_2, K_P$, and K_I are fixed at the values given in Table 1.

We now fix the following parameters

$$u_f = 2, \quad u_g = 0, \quad u_h = 0, \quad (123)$$

and plot a one-parameter bifurcation diagram in α , with x_0 on the y -axis. All simulations were performed using [19].




Figure 5: One-parameter bifurcation diagram of system (21) with α as the bifurcation parameter and $u_f = 2$, $u_g = 0$, $u_h = 0$.




Figure 6: Close-up of on the one-parameter bifurcation diagram represented in Figure 5.

We can see that as α decreases, there is a saddle-node bifurcation, resulting in two equilibria ${}_{(110)}\mathcal{E}^{(1)}$ and ${}_{(110)}\mathcal{E}^{(2)}$ appearing (both unstable). Next, there is a transcritical bifurcation with the ${}_{(110)}\mathcal{E}^{(1)}$ equilibrium, which results in the positive equilibrium coming into the interior of the admissible region Ω . After that, a saddle-node of limit cycles bifurcation occurs, which gives birth to a stable and unstable periodic orbits. The ${}_{(111)}\mathcal{E}$ equilibrium (unstable), undergoes a Hopf bifurcation, and as a consequence it becomes asymptotically stable, and the stable periodic orbits disappears. Since these bifurcations occur for a narrow range of α , a close-up is presented in Figure 6. Stable periodic orbit represents a case in which all three populations oscillate indefinitely, and hence the production of methane fluctuates. As already mentioned in section 4.2, this situation is not a desirable one, because it might result it decreased productivity of the biogas plant. The unstable periodic orbits acts as a separatrix, giving the border of the basin of attraction of two asymptotically stable equilibria in the case of bistability.

Since by the conservation principles (20), $s_0 = u_f - x_0$, the bifurcation diagram in α with s_0 on the y -axis is similar to the one presented in Figure 5. The amount of chlorophenol in the system is inversely proportional to the concentration of the phenol degrader. As the dilution rate α decreases, concentration of the chlorophenol degrader in the interior equilibrium increases, and as a consequence, the amount of chlorophenol decreases. It thus suggests, that operating on lower dilution rates results in the most desirable dynamics, i.e., an asymptotically stable interior equilibrium and fast chlorophenol removal.

To extend the previous analysis, we now fix the following parameters

$$u_g = 0, \quad u_h = 0.1, \quad (124)$$

and plot a two-parameter bifurcation diagram of system (21), choosing α and u_f as the bifurcation parameters. Each region of the diagram is labeled and the corresponding dynamics are represented schematically in figures around it. Black dashed curve corresponds to saddle-node of equilibria bifurcation (LP), black solid curve represents saddle-node of limit cycles bifurcation (SNLC), black dotted curve denotes Hopf bifurcation (HB), and grey solid curves represent transcritical bifurcations (BP). We also denote the predicted heteroclinic bifurcation by a grey dashed curve which lies very close to the Hopf curve.




Figure 7: Two-parameter bifurcation diagram of system (21) with $u_g = 0$ and $u_h = 0.1$.




Figure 8: Close-up on region I of Figure 7. The SNLC curve intersects the HB curve at Bautin bifurcation. This results in the change of criticality of the Hopf bifurcation from supercritical (on the left) to subcritical (on the right).




Figure 9: Close-up on region II of Figure 7.

We can see that varying two parameters at the same time can lead to a much more complicated dynamics than in the case of one-parameter bifurcations. There is a generalized Hopf bifurcation, at the point at which the Hopf curve

intersects the saddle-node of limit cycles curve. This is the point where the criticality of the Hopf bifurcation changes from supercritical to subcritical, looking from left to right. The unstable periodic orbit disappears through a heteroclinic bifurcation. There are two heteroclinic orbits that form a cycle that joins the two equilibria in the x_0x_1 face, then passes into the interior, and then goes back to the boundary in the x_0x_1 face. The point at which the Hopf, homoclinic, and saddle-node of limits cycles curves intersect, represents the Bogdanov-Takens bifurcation.

From the biological viewpoint, the most interesting dynamics is occurs in regions 5 and *iii*. There, the interior equilibrium is asymptotically stable. In the case of region 5 we also observe bistability with the $_{(100)}\mathcal{E}$ equilibrium. In region *iii*, there is uniform persistence, and thus an interior compact attractor is present. As was previously anticipated by the analysis of the one-parameter bifurcation diagram, operating at low dilution rates is the most desirable approach. If α is small enough, it is possible to remain in region *iii*, even for high inflow rate u_f .

5 Conclusions

In this work we have generalised the approach presented in [14] by including multiple substrate inflow into the chemostat, while maintaining generality (in most cases) with respect to the exact form of the growth functions. We observed that allowing the inflow of multiple substrates resulted in much more complex dynamics of the system. For example, eight steady states are possible. Previously, the theoretical results were limited to existence and uniqueness of up to three equilibria (when chlorophenol was the only input substrate), and to numerical evidence suggesting that the model should be subjected to a more detailed analysis. We also observed that external addition of substrates can result in bistability - two equilibria can simultaneously be asymptotically stable. As well, there can be an orbitally asymptotically stable periodic orbit with all of the populations surviving and an asymptotically stable equilibrium with only chlorophenol degrader population surviving.

We have also confirmed the findings of the previous analysis in [14], where numerical evidence of the occurrence of a supercritical Hopf bifurcation was given. Theoretical conditions for the existence of a Hopf bifurcation were provided in the case of specific forms of the growth functions. Varying any one of the three parameters: chlorophenol, phenol, and hydrogen inflow rates, was shown to result in a Hopf bifurcation. Theoretical results for varying the dilution rate as the bifurcation parameter has been left for future work. However, we have observed numerically, that varying this parameter can result in a Hopf bifurcation and a saddle-node of limit cycles bifurcation. Our numerical investigations also showed that increasing the inflow rate of the substrates has a stabilizing effect on the entire system. From a biological engineering point of view, i.e. a bioreactor treating a monochlorophenol rich waste stream, instability would typically be undesirable in terms of process performance. Therefore, identification of control strategies to avoid periodic behaviour is an important output of this work.

Another result, particularly important for engineering applications, concerns the persistence of the system for a range of parameter sets. Knowing when the microbial populations survive is again crucial from a process control perspective, and it is one of the main theoretical results of this work. We have proven that in two configurations of equilibria (in both cases all the boundary equilibria are saddle points) we observe not only persistence, but also uniform persistence, a much stronger result. These situations occur when there is an inflow of all three substrates, but also when phenol addition is not considered (i.e., when $u_g = 0$).

Although we now know much more about the dynamics of the system, it is not fully understood. This follows from the numerical results provided by the two-parameter bifurcation diagrams. The analyses reveals that varying the dilution rate and the chlorophenol inflow simultaneously, can lead to a Bogdanov-Takens, or Bautin (generalized Hopf) bifurcations. Also, for the cases of bistability, where both a boundary and the interior equilibrium are asymptotically stable, it is of great importance to empiricists to have an estimation of the basins of attraction of these equilibria. This result is usually difficult to obtain theoretically, however numerical estimations are possible. Another factor that is of interest would be the inclusion of stochasticity in the model. In practice, even if the interior equilibrium is globally asymptotically stable, one of the microorganisms may become extinct. This might occur when a population is very small, and the stochastic noise effects result in the solution curve reaching one of the invariant faces of the admissible region.

There has been resistance to the idea that simplified models, of the type described here, are too remote from the systems they represent to be of worth to practitioners. Without experimental results to compare against model predictions, this case becomes stronger. However, we can look to emerging disciplines such as synthetic biology to help bridge the theoretical and the applied [7]. Recent studies have shown that synthetically derived anaerobic communities are able to confirm model predictions and provide insight into the ecology and dynamics of microbial communities that are relevant in practice [5]. We believe this work provides a basis by which experimental studies describing a chlorophenol mineralising food-web could be undertaken.

6 Acknowledgments

M.J.W. acknowledges the support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No. 702408 (DRAMATIC).

References

- [1] I. BASSANI, P. G. KOUGIAS, L. TREU, AND I. ANGELIDAKI, *Biogas upgrading via hydrogenotrophic methanogenesis in two-stage continuous stirred tank reactors at mesophilic and thermophilic conditions*, Environ. Sci. Technol., 49 (2015), pp. 12585–12593.
- [2] G. J. BUTLER, H. I. FREEDMAN, AND P. WALTMAN, *Uniformly persistent systems*, Proceedings of the American Mathematical Society, 96 (1986), pp. 425–430.
- [3] G. J. BUTLER AND G. S. K. WOLKOWICZ, *Predator — mediated coexistence in a chemostat: Coexistence and competition reversal*, Mathematical Modelling, 8 (1987), pp. 781 – 785.
- [4] J. CHEN, M. J. WADE, J. DOLFING, AND O. S. SOYER, *Increasing sulfate levels show a differential impact on synthetic communities comprising different methanogens and a sulfate reducer*, J. Royal Soc. Interface, 16 (2019), p. 20190129.
- [5] H. DELATTRE, J. CHEN, M. WADE, AND O. SOYER, *Thermodynamic modelling of synthetic communities predicts minimum free energy requirements for sulfate reduction and methanogenesis*, bioRxiv, (2019), p. 857276.
- [6] M. EL HAJJI, N. CHORFI, AND M. JLELI, *Mathematical modelling and analysis for a three-tiered microbial food web in a chemostat*, Electron. J. Differ. Eq., 255 (2017), pp. 1–13.
- [7] M. EL KAROUI, M. HOYOS-FLIGHT, AND L. FLETCHER, *Future Trends in Synthetic Biology — A Report*, Front. Bioeng. Biotechnol., 7 (2019), p. 175.
- [8] T. GROSSKOPF AND O. SOYER, *Microbial diversity arising from thermodynamic constraints*, ISME J., 10 (2016), pp. 2725–2733.
- [9] Y. KUZNESTOV, *Elements of Applied Bifurcation Theory*, vol. 112, Springer-Verlag, New York, 2004.
- [10] L. LEVÉN, K. NYBERG, AND A. SCHN URER, *Conversion of phenols during anaerobic digestion of organic solid waste – a review of important microorganisms and impact of temperature*, J. Env. Manage., 95 (2012), pp. 99–103.
- [11] MAPLE, version 18.02, Waterloo Maple Inc., Waterloo, Ontario, 2018.
- [12] MATLAB, version 9.5.0.944444 (R2018b), The MathWorks Inc., Natick, Massachusetts, 2018.
- [13] C. MAZUR, W. JONES, AND C. TEBES-STEVENSON, *H₂ consumption during the microbial reductive dehalogenation of chlorinated phenols and tetrachloroethene*, Biodegradation, 14 (2003), pp. 285–295.
- [14] T. SARI AND M. J. WADE, *Generalised approach to modelling a three-tiered microbial food-web*, Mathematical Biosciences, 291 (2017), pp. 21–37.
- [15] B. SCHINK, *Energetics of syntrophic cooperation in methanogenic degradation*, Microbiol. Mol. Biol. Rev., 61 (1997), pp. 262–280.
- [16] H. L. SMITH AND P. WALTMAN, *The Theory of the Chemostat: Dynamics of Microbial Competition*, Cambridge Studies in Mathematical Biology, Cambridge University Press, 1995.
- [17] N. W. SMITH, P. R. SHORTEN, E. H. ALTERMANN, N. C. ROY, AND W. C. MCNABB, *Hydrogen cross-feeders of the human gastrointestinal tract*, Gut Microbes, 10 (2019), pp. 270–288.
- [18] M. J. WADE, R. W. PATTINSON, N. G. PARKER, AND J. DOLFING, *Emergent behaviour in a chlorophenol-mineralising three-tiered microbial ‘food web’*, Journal of Theoretical Biology, 389 (2016), pp. 171–186.
- [19] XPPAUT, version 8.0, Dr. Bard Ermentrout, Dept of Mathematics, University of Pittsburgh, Pittsburgh PA, 2016.