

PROPERTIES OF THE RESOLUTION OF ALMOST GORENSTEIN ALGEBRAS

GIUSEPPE ZAPPALÀ

ABSTRACT. We study properties of the resolution of almost Gorenstein artinian algebras R/I , i.e. algebras defined by ideals I such that $I = J + (f)$, with J Gorenstein ideal and $f \in R$. Such algebras generalize the well known almost complete intersection artinian algebras. Then we give a new explicit description of the resolution and of the graded Betti numbers of almost complete intersection ideals of codimension 3 and we characterize the ideals whose graded Betti numbers can be achieved using artinian monomial ideals.

1. INTRODUCTION

Let $I \subset R = k[x_1, \dots, x_c]$ be a homogeneous ideal. In commutative algebra and algebraic geometry it is a very important subject to understand the structure of the graded minimal free resolution of I . The simplest case is when I is generated by a regular sequence. In this case the Koszul complex provides the answer to this problem, in any codimension. In the other cases we know satisfying answers only in few situations.

In codimension 2, when I is a perfect ideal, the structure of the resolution is given by the Hilbert-Burch theorem (see [Hi], [Bu]). This allowed, for instance, to characterize the graded Betti numbers of such ideals (see [Ga], [CGO]).

In codimension 3, when I is a Gorenstein ideal, the structure of the resolution is given by the Eisenbud-Buchsbaum theorem (see [BE]). Also in this case, due to this very important result, it is possible to characterize the graded Betti numbers (see [Di], [RZ2], [CV]).

When I is perfect of codimension greater than 2 or I is Gorenstein of codimension greater than 3 very little is known about the structure the resolution. One of the most studied cases are Gorenstein ideals in codimension 4 (see [KM],[Re]), although it is not known a description of the graded Betti numbers.

Another very important case consists of perfect ideals of codimension c generated by $c+1$ generators, i.e. almost complete intersection ideals. These ideals have the advantage that are directly linked to a Gorenstein ideal in a complete intersection, so there is the expectation that properties of the Gorenstein resolutions can affect their resolution. In codimension 3 the structure of the resolution of such ideals was studied in [BE] and in [RZ4].

2010 *Mathematics Subject Classification.* 13 H 10, 14 N 20, 13 D 40.

Key words and phrases. Resolution, Graded Betti numbers, Gorenstein ring, almost complete intersection.

In section 3 we study the resolution of the almost Gorenstein algebras, a very important generalization of the notion of almost complete intersection. Also these ideals are directly linked to the Gorenstein ideals. In particular, in Theorem 3.5, we state an important duality properties for the graded minimal free resolution.

In section 4 we focus on codimension 3 almost complete intersection algebras and we give an explicit new characterization of the graded Betti numbers. Finally we characterize the ideals, in this class, whose graded Betti numbers can be achieved using monomial ideals (Corollary 4.13 and Proposition 4.15).

2. PRELIMINARY FACTS

Throughout the paper k will be a infinite field and $R := k[x_1, \dots, x_c]$, $c \geq 3$, will be the standard graded polynomial k -algebra. We consider only homogeneous ideals.

Let $A = R/I$ be a standard graded artinian k -algebra, where I is a homogeneous ideal of R . Recall that A is said a *almost complete intersection algebra* if I is minimally generated by $c + 1$ elements (in this case I is said an almost complete intersection ideal). Every almost complete intersection ideal is directly linked to a Gorenstein ideal in a complete intersection.

It is possible to generalize this notion in the following way.

Definition 2.1. Let $A_Q = R/I_Q$ be a standard graded artinian k -algebra, where I_Q is a homogeneous ideal of R . We say that A is an *almost Gorenstein algebra* if

1. $I_Q = I_Z + (f)$, where I_Z is a Gorenstein ideal of codimension c and $f \in (x_1, \dots, x_c)$, $f \notin I_Z$.
2. A_Q is not a Gorenstein algebra.

In particular, if I_Z is a complete intersection ideal of codimension c , then A_Q is an almost complete intersection algebra.

Gorenstein ideals of codimension 3 are better understood, because of the structure theorem of Buchsbaum and Eisenbud (see [BE]). The possible graded Betti numbers for Gorenstein ideal of codimension 3 were described in [Di] and relatively to the Hilbert function in [RZ2].

More precisely let (d_1, \dots, d_{2n+1}) , $d_1 \leq \dots \leq d_{2n+1}$, be a sequence of positive integers. It is the sequence of the degrees of the minimal generators of a Gorenstein ideal of codimension 3 iff

- 1) $\vartheta = \frac{1}{n} \sum_{i=1}^{2n+1} d_i$ is an integer;
- 2) $\vartheta > d_i + d_{2n+3-i}$ for $2 \leq i \leq n$ (Gaeta conditions).

Let $\delta = (d_1, \dots, d_{2n+1})$, $d_1 \leq \dots \leq d_{2n+1}$, be the sequence of the degrees of the minimal generators of a Gorenstein ideal of codimension 3. Let $\text{Gor}(\delta)$ be the set of all Gorenstein ideals $I \subset k[x_1, x_2, x_3]$, whose sequence of the degrees of the minimal generators is δ . Moreover we define

$$\text{CI}_\delta = \{(a_1, a_2, a_3) \in (\mathbb{Z}^+)^3 \mid a_1 \leq a_2 \leq a_3 \text{ and } \exists I \in \text{Gor}(\delta) \text{ containing a regular sequence of degrees } (a_1, a_2, a_3)\}.$$

To study ideals linked to Gorenstein ideals in a complete intersection, in codimension 3, we need the following result.

Theorem 2.2. *The set CI_δ has a unique minimal element.*

Proof. See Theorem 3.6 in [RZ3], where this minimal element is explicitly computed. \square

We will denote the only minimal element of CI_δ with $\min(\delta)$.

3. ALMOST GORENSTEIN ALGEBRAS

Let $A_Q = R/I_Q$ be an almost Gorenstein algebra. Then $I_Q = I_Z + (f)$, where I_Z is a Gorenstein ideal. Let e be the socle degree of $A_Z = R/I_Z$. Since $f \in (x_1, \dots, x_c)$, $f \notin I_Z$ we get that $1 \leq \deg f \leq e$. We are interested on the graded minimal free resolutions of such ideals. We denote by ϑ_Z the degree of the only last syzygy of I_Z . Of course $\vartheta_Z = e + c$.

Proposition 3.1. *Let $A_Q = R/I_Q$ be an almost Gorenstein algebra, with $I_Q = I_Z + (f)$ as above. Then $I_G = I_Z : I_Q$ is a Gorenstein ideal of codimension c and $\vartheta_G = \vartheta_Z - \deg f$.*

Proof. Let

$$0 \rightarrow F_c \rightarrow F_{c-1} \rightarrow \dots \rightarrow F_1 \rightarrow R \rightarrow A_Q \rightarrow 0$$

and

$$0 \rightarrow K_c \rightarrow K_{c-1} \rightarrow \dots \rightarrow K_1 \rightarrow R \rightarrow A_Z \rightarrow 0$$

be graded minimal free resolutions. Since A_Z is Gorenstein $K_c \cong R(-\vartheta_Z)$. Moreover $F_1 \cong K'_1 \oplus R(-d^*)$, where K'_1 is a direct summand of K_1 and $d^* = \deg f$. Since $I_Z \subseteq I_Q$ we can use liaison theory. If we set $A_G = R/I_G$, we get the exact sequence

$$0 \rightarrow (A_Q)^\vee(-\vartheta_Z) \rightarrow A_Z \rightarrow A_G \rightarrow 0.$$

So we can compute a graded free resolution of A_G by mapping cone. Finally we can delete the summand $K_1^\vee(-\vartheta_Z)$ in the tail of resolution, so the last module in a minimal graded free resolution of A_G is isomorphic to $R(-\vartheta_Z + d^*)$. \square

Remark 3.2. Note that $\vartheta_Z - d^* = \vartheta_G$. We set $d = 2\vartheta_Z - \vartheta_G$. Note that $d = \vartheta_Z + d^* = \vartheta_G + 2d^*$, where $d^* = \deg f$.

Proposition 3.3. *A graded resolution (possibly not minimal) of A_Q is*

$$\begin{aligned} 0 \rightarrow G_{c-1}(-d^*) \rightarrow G_{c-2}(-d^*) \oplus K_{c-1} \rightarrow \dots \\ \dots \rightarrow G_1(-d^*) \oplus K_2 \rightarrow R(-d^*) \oplus K_1 \rightarrow R \rightarrow A_Q \rightarrow 0. \end{aligned}$$

Proof. In general we have the short exact sequence

$$0 \rightarrow (A_G)^\vee(-\vartheta_Z) \rightarrow A_Z \rightarrow A_Q \rightarrow 0.$$

Let G_\bullet and K_\bullet be graded minimal free resolution of A_G and A_Z respectively. By the duality properties of the Gorenstein resolutions we have that

$$G_\bullet \cong G_\bullet^\vee(-\vartheta_G).$$

Consequently, by Remark 3.2

$$G_\bullet(-d^*) \cong G_\bullet^\vee(-\vartheta_G - d^*) \cong G_\bullet^\vee(-\vartheta_Z).$$

So a resolution (which may be not minimal) of A_Q is

$$\begin{aligned} 0 \rightarrow G_{c-1}(-d^*) \rightarrow G_{c-2}(-d^*) \oplus K_{c-1} \rightarrow \dots \\ \dots \rightarrow G_1(-d^*) \oplus K_2 \rightarrow R(-d^*) \oplus K_1 \rightarrow R \rightarrow A_Q \rightarrow 0. \end{aligned}$$

□

One of the simplest cases is when $\deg f = e$.

Corollary 3.4. *Let $A_Q = R/I_Q$ be an almost Gorenstein algebra, with $I_Q = I_Z + (f)$ and $\deg f = e$, the socle degree of A_Z . Let*

$$0 \rightarrow R(-(e+c)) \rightarrow K_{c-1} \rightarrow \dots \rightarrow K_1 \rightarrow R \rightarrow A_Z \rightarrow 0$$

be a graded minimal free resolution of A_Z . Let us suppose that I_Z is generated in degrees less than e .

Then a graded minimal free resolution of A_Q is

$$\begin{aligned} 0 \rightarrow R(-(e+c-1)) \binom{c}{c-1} \rightarrow K_{c-1} \oplus R(-(e+c-2)) \binom{c}{c-2} \rightarrow \dots \\ \dots \rightarrow K_1 \oplus R(-(e-1)) \binom{c}{1} \rightarrow K_1 \oplus R(-e) \rightarrow R \rightarrow A_Q \rightarrow 0. \end{aligned}$$

Proof. If we set $I_G = I_Z : I_Q$, then $I_G = (x_1, \dots, x_c)$. Hence

$$G_i(-d^*) = G_i(-e) \cong R(-(i+e)) \binom{c}{i}.$$

Now we apply Proposition 3.3 to obtain a graded resolution of A_Q . Since I_Z is generated in degrees less than e , we have that $A_{Q,i} = A_{Z,i}$, for $i \leq e-1$ and $A_{Q,i} = 0$ for $i \geq e$, so such a resolution is minimal. □

If we set $H_i = G_i(-d^*)$, for $1 \leq i \leq c-1$ we have that

$$\begin{aligned} H_i = G_i(-d^*) \cong G_{c-i}^\vee(-\vartheta_G - d^*) \cong \\ \cong (H_{c-i}(d^*))^\vee(-\vartheta_G - d^*) \cong H_{c-i}^\vee(-\vartheta_G - 2d^*) \cong H_{c-i}^\vee(-d). \end{aligned}$$

So the resolution of Proposition 3.3 becomes

$$\begin{aligned} 0 \rightarrow H_{c-1} \rightarrow H_{c-2} \oplus K_{c-1} \rightarrow \dots \\ \dots \rightarrow H_{c-2}^\vee(-d) \oplus K_3 \rightarrow H_{c-1}^\vee(-d) \oplus K_2 \rightarrow R(-d^*) \oplus K_1 \rightarrow R \rightarrow A_Q \rightarrow 0. \end{aligned}$$

Theorem 3.5. *Let $A_Q = R/I_Q$ be an almost Gorenstein algebra with graded minimal free resolution*

$$0 \rightarrow F_c \rightarrow F_{c-1} \rightarrow \dots \rightarrow F_2 \rightarrow F_1 \rightarrow R \rightarrow A_Q \rightarrow 0.$$

Let us suppose that a minimal set of generators of I_Q is (f_1, \dots, f_n, f) with (f_1, \dots, f_n) minimal set of generators of a Gorenstein ideal I_Z . Let K_\bullet a graded minimal free resolution of I_Z and G_\bullet a graded minimal free resolution of $I_G = I_Z : I_Q$. Then

$$F_2 \cong F_c^\vee(-d) \oplus F_2' \cong G_1(-d^*) \oplus K_2'$$

for some F_2' and where K_2' is a direct summand of K_2 .

Proof. By Proposition 3.3, we have that F_c is a direct summand of $H_{c-1} = G_{c-1}(-d^*)$, namely $H_{c-1} \cong F_c \oplus H'_{c-1}$. By the hypotheses about the sets of generators $F_1 \cong R(-d^*) \oplus K_1$ and $F_2 \cong G_1(-d^*) \oplus K'_2 = H_1 \oplus K'_2$ where K'_2 is a direct summand of K_2 . Therefore

$$F_2 \cong H_1 \oplus K'_2 \cong H_{c-1}^\vee(-d) \oplus K'_2 \cong F_c^\vee(-d) \oplus H'_{c-1}(-d) \oplus K'_2.$$

□

Corollary 3.6. *Under the same hypotheses of Theorem 3.5,*

$$F_2 \cong F_c^\vee(-d) \oplus G'_1(-d^*) \oplus K'_2,$$

where G'_1 is a direct summand of G_1 and K'_2 is a direct summand of K_2 .

Proof. This follows immediately by the conclusion of the proof of Theorem 3.5. □

Corollary 3.7. *Let $A_Q = R/I_Q$ be an almost complete intersection algebra with graded minimal free resolution*

$$0 \rightarrow F_c \rightarrow F_{c-1} \rightarrow \dots \rightarrow F_2 \rightarrow F_1 \rightarrow R \rightarrow A_Q \rightarrow 0.$$

Then $F_2 \cong F_c^\vee(-d) \oplus F'_2$, where d is the sum of the degrees of a minimal set of generators of I_Q .

Proof. By the hypotheses I_Q is minimally generated by (f_1, \dots, f_c, f) , where (f_1, \dots, f_c) is a regular sequence. We set $I_Z = (f_1, \dots, f_c)$. Now, by applying Theorem 3.5, we get $F_2 \cong F_c^\vee(-d) \oplus F'_2$. By Remark 3.2, $d = \vartheta_Z + \deg f$, hence $d = \sum_{i=1}^c \deg f_i + \deg f$. □

This Corollary generalizes Lemma 2.1 in [RZ1].

4. ALMOST COMPLETE INTERSECTIONS IN CODIMENSION 3

In this section we study some properties of the resolutions of the almost complete intersection algebras in codimension 3.

An almost complete intersection algebra $A_Q = R/I_Q$ of codimension 3 has a graded minimal free resolution of the type

$$0 \rightarrow F_3 \rightarrow F_2 \rightarrow F_1 \rightarrow R \rightarrow A_Q \rightarrow 0.$$

We set $t_Q = \text{rank } F_3$. Of course $t_Q \geq 2$.

Theorem 4.1. *Let $A_Q = R/I_Q$ be an almost complete intersection algebra in codimension 3.*

If $t = t_Q$ is even, then a graded minimal free resolution of A_Q is

$$\begin{aligned} 0 \rightarrow \bigoplus_{i=1}^t R(-(d - s_i)) \rightarrow \bigoplus_{i=1}^3 R(-(d_i + d^*)) \oplus \bigoplus_{i=1}^t R(-s_i) \rightarrow \\ \rightarrow R(-d^*) \oplus \bigoplus_{i=1}^3 R(-d_i) \rightarrow R \rightarrow A_Q \rightarrow 0 \end{aligned}$$

where $d = d_1 + d_2 + d_3 + d^*$, $d_1 \leq d_2 \leq d_3$, $s_1 \leq \dots \leq s_t$.

If $t = t_Q$ is odd, then a graded minimal free resolution of A_Q is

$$\begin{aligned} 0 \rightarrow \bigoplus_{i=1}^t R(-(d - s_i)) \rightarrow \bigoplus_{1 \leq i < j \leq 3} R(-(d_i + d_j)) \oplus \bigoplus_{i=1}^t R(-s_i) \rightarrow \\ \rightarrow R(-d^*) \oplus \bigoplus_{i=1}^3 R(-d_i) \rightarrow R \rightarrow A_Q \rightarrow 0 \end{aligned}$$

where $d = d_1 + d_2 + d_3 + d^*$, $d_1 \leq d_2 \leq d_3$, $s_1 \leq \dots \leq s_t$.

Proof. Just using Theorem 3.5 and Lemma 2.1 in [RZ1]. \square

With the notation of Theorem 4.1, we set $s_Q = \sum_{i=1}^t s_i$.

Proposition 4.2. *We have that*

- 1) $s_Q = (\frac{t}{2} - 1)d^* + \frac{t}{2}(d_1 + d_2 + d_3) = d\frac{t}{2} - d^*$, if $t = t_Q$ is even;
- 2) $s_Q = \frac{t+1}{2}d^* + \frac{t-1}{2}(d_1 + d_2 + d_3) = d\frac{t-1}{2} + d^*$, if $t = t_Q$ is odd.

Proof. This follows from Theorem 4.1, since the alternating sum of the shifts of a graded minimal free resolution is equal to zero. \square

Let A_Q be an almost complete intersection algebra with minimal graded free resolution F_\bullet . If $F_2 = \bigoplus_{i=1}^{t+3} R(-\beta_i)$, we set $s_{(2)} = \sum_{i=1}^{t+3} \beta_i$. Analogously if $F_3 = \bigoplus_{i=1}^t R(-\gamma_i)$, we set $s_{(3)} = \sum_{i=1}^t \gamma_i$. Note that $s_Q = td - s_{(3)}$. We set

$$u_Q = s_{(2)} - s_Q = s_{(2)} + s_{(3)} - td.$$

We can compute the fundamental numerical invariant d^* directly from the graded minimal free resolution of A_Q .

Proposition 4.3. *Let $A_Q = R/I_Q$ be an almost complete intersection algebra with minimal graded free resolution F_\bullet . Let $u = u_Q$ and d be the sum of the degrees of the minimal generators of I_Q . Then*

$$d^* := \frac{u - d}{2}, \text{ if } t \text{ is even; } d^* = \frac{2d - u}{2}, \text{ if } t \text{ is odd.}$$

Proof. If t is even then

$$F_2 \cong \bigoplus_{i=1}^3 R(-(d_i + d^*)) \oplus \bigoplus_{i=1}^t R(-s_i),$$

hence $u = u_Q = \sum_{i=1}^3 (d_i + d^*) = d + 2d^*$.

If t is odd then

$$F_2 \cong \bigoplus_{1 \leq i < j \leq 3} R(-(d_i + d_j)) \oplus \bigoplus_{i=1}^t R(-s_i),$$

hence $u = u_Q = \sum_{1 \leq i < j \leq 3} (d_i + d_j) = 2(d_1 + d_2 + d_3) = 2(d - d^*)$. \square

Remark 4.4. By Theorem 3.5, $F_2 \cong F_3^\vee(-d) \oplus F_2'$. By Theorem 4.1

$$\begin{aligned} F_2' &\cong \bigoplus_{i=1}^3 R(-(d_i + d^*)) \text{ if } t_Q \text{ is even;} \\ F_2' &\cong \bigoplus_{1 \leq i < j \leq 3} R(-(d_i + d_j)) \text{ if } t_Q \text{ is odd.} \end{aligned}$$

Example 4.5. Let A_Q be an almost complete intersection algebra with minimal graded free resolution

$$\begin{aligned} 0 \rightarrow R(-20)^3 \oplus R(-21)^4 \oplus R(-22)^4 \rightarrow \\ \rightarrow R(-17) \oplus R(-18) \oplus R(-19)^5 \oplus R(-20)^4 \oplus R(-21)^3 \rightarrow \\ \rightarrow R(-8) \oplus R(-9) \oplus R(-10) \oplus R(-14) \rightarrow R \rightarrow A_Q \rightarrow 0. \end{aligned}$$

We have that $t = 11$, $d = 8 + 9 + 10 + 14 = 41$ and $d - 20 = 21$, $d - 21 = 20$, $d - 22 = 19$, so $u = 17 + 18 + 19 = 54$. Consequently

$$d^* = \frac{2d - u}{2} = 14 \Rightarrow d_1 = 8, d_2 = 9, d_3 = 10.$$

Moreover

$$s_1 = \dots = s_4 = 19, s_5 = \dots = s_8 = 20, s_9 = s_{10} = s_{11} = 21.$$

Now we compute d^* for the monomial ideals.

Example 4.6. Let $I_Q = (x^{a_1}, y^{a_2}, z^{a_3}, x^{b_1}y^{b_2}) \subset k[x, y, z]$, with $0 < b_1 < a_1$, $0 < b_2 < a_2$, $a_3 > 0$.

Its graded minimal free resolution is

$$\begin{aligned} 0 \rightarrow R(-(a_1 + b_2 + a_3)) \oplus R(-(b_1 + a_2 + a_3)) \rightarrow R(-(b_1 + a_2)) \oplus R(-(a_1 + b_2)) \oplus \\ \oplus R(-(a_1 + a_3)) \oplus R(-(a_2 + a_3)) \oplus R(-(b_1 + b_2 + a_3)) \rightarrow \\ \rightarrow R(-a_1) \oplus R(-a_2) \oplus R(-a_3) \oplus R(-(b_1 + b_2)) \rightarrow I_Q \rightarrow 0. \end{aligned}$$

So we have that

$$F'_2 \cong R(-(a_1 + a_3)) \oplus R(-(a_2 + a_3)) \oplus R(-(b_1 + b_2 + a_3)).$$

Consequently $d^* = a_3$.

Example 4.7. Let $I_Q = (x^{a_1}, y^{a_2}, z^{a_3}, x^{b_1}y^{b_2}z^{b_3}) \subset k[x, y, z]$, with $0 < b_i < a_i$, for $i = 1, 2, 3$.

Its graded minimal free resolution is

$$\begin{aligned} 0 \rightarrow R(-(a_1 + a_2 + b_3)) \oplus R(-(a_1 + b_2 + a_3)) \oplus R(-(b_1 + a_2 + a_3)) \rightarrow \\ \rightarrow R(-(b_1 + b_2 + a_3)) \oplus R(-(b_1 + a_2 + b_3)) \oplus R(-(a_1 + b_2 + b_3)) \oplus \\ \oplus R(-(a_1 + a_2)) \oplus R(-(a_1 + a_3)) \oplus R(-(a_2 + a_3)) \rightarrow \\ \rightarrow R(-a_1) \oplus R(-a_2) \oplus R(-a_3) \oplus R(-(b_1 + b_2 + b_3)) \rightarrow I_Q \rightarrow 0. \end{aligned}$$

So we have that

$$F'_2 \cong R(-(a_1 + a_2)) \oplus R(-(a_1 + a_3)) \oplus R(-(a_2 + a_3)).$$

Consequently $d^* = b_1 + b_2 + b_3$.

Note that there is always a regular sequence (f, f_2, f_3) in I_Q , with

$$\deg f = d^*, \deg f_2 = d_2, \deg f_3 = d_3.$$

Now we study the liaison of I_Q in a complete intersection of type d^*, d_2, d_3 .

Theorem 4.8. *Let $A_Q = R/I_Q$ be an almost complete intersection algebra of codimension 3, with $t = t_Q$ even.*

If $d^ \neq d_1$, then I_Q is linked to a Gorenstein ideal with $t + 1$ minimal generators in a complete intersection of type (d^*, d_2, d_3) .*

Proof. We set $\vartheta_Z = d^* + d_2 + d_3$. By liaison, using mapping cone, we get the following graded resolution of a Gorenstein algebra

$$\begin{aligned} 0 \rightarrow R(-(\vartheta_Z - d_1)) \rightarrow F_3(d_1) \oplus F_2^\vee(-\vartheta_Z) \rightarrow \\ \rightarrow F_3^\vee(-\vartheta_Z) \oplus R(-d^*) \oplus R(-d_2) \oplus R(-d_3) \rightarrow R. \end{aligned}$$

We have that

$$F_2^\vee(-\vartheta_Z) \cong \bigoplus_{i=1}^3 R(-(\vartheta_Z - d_i - d^*)) = R(-(d_2 + d_3 - d_1)) \oplus R(-d_3) \oplus R(-d_2).$$

Now we set $\vartheta_G = \vartheta_Z - d_1$. We have that

$$(F_3^\vee(-\vartheta_Z))^\vee(-\vartheta_G) = F_3(d_1)$$

and

$$R(-d^*)^\vee(-\vartheta_G) \cong R(d^* - \vartheta_G) = R(-(d_2 + d_3 - d_1)).$$

So, if this resolution were minimal, then

$$R(-d_2)^\vee(-\vartheta_G) \cong R(-d_3) \iff d_2 + d_3 = \vartheta_G \iff d^* = d_1.$$

Therefore the terms $R(-d_2)$ and $R(-d_3)$ are redundant and we obtain the graded minimal free resolution

$$\begin{aligned} 0 \rightarrow R(-\vartheta_G) \rightarrow F_3(d_1) \oplus R(-(d_2 + d_3 - d_1)) \rightarrow \\ \rightarrow F_3^\vee(-\vartheta_Z) \oplus R(-d^*) \rightarrow R. \end{aligned}$$

□

Example 4.9. Let us consider the ideal

$$(x_4^3 x_8, x_3^4, x_4^3 x_6^3, x_2^4 x_6^3 - x_1^6 x_8) \subset k[x_1, \dots, x_8].$$

Its artinian reduction defines an almost complete intersection algebra whose graded minimal free resolution is

$$\begin{aligned} 0 \rightarrow R(-11) \oplus R(-14) \rightarrow R(-7) \oplus R(-8) \oplus R(-10)^2 \oplus R(-11) \rightarrow \\ \rightarrow R(-4)^2 \oplus R(-6) \oplus R(-7) \rightarrow R. \end{aligned}$$

We see that $d^* = 4$. If we link in the complete intersection

$$(x_4^3 x_8, x_4^3 x_6^3 - x_3^4 x_7^2, x_3^4 x_5^3 - x_2^4 x_6^3 + x_1^6 x_8)$$

we get a Gorenstein algebra whose graded minimal free resolution is

$$\begin{aligned} 0 \rightarrow R(-13) \rightarrow R(-6) \oplus R(-7)^2 \oplus R(-9) \oplus R(-10) \rightarrow \\ \rightarrow R(-3) \oplus R(-4) \oplus R(-6)^2 \oplus R(-7) \rightarrow R, \end{aligned}$$

so, in this case, the term $R(-6) \oplus R(-7)$ cannot be deleted.

Theorem 4.10. *Let $A_Q = R/I_Q$ be an almost complete intersection algebra of codimension 3, with $t = t_Q$ even.*

If $d^ = d_1$, then I_Q has the same graded Betti numbers of an almost complete intersection ideal I'_Q , which is linked to a Gorenstein ideal with $t + 1$ minimal generators in a complete intersection of type (d^*, d_2, d_3) .*

Proof. By performing the liaison we get the following Gorenstein resolution as in the proof of Theorem 4.8

$$0 \rightarrow R(-\vartheta_G) \rightarrow F_3(d_1) \oplus R(-(d_2 + d_3 - d_1)) \oplus R(-d_2) \oplus R(-d_3) \rightarrow \\ \rightarrow F_3^\vee(-\vartheta_Z) \oplus R(-d^*) \oplus R(-d_2) \oplus R(-d_3) \rightarrow R.$$

Let G_\bullet be the graded minimal resolution obtained by the previous one, by omitting the term $R(-d_2) \oplus R(-d_3)$. By Remark 3.8 in [RZ2], there exist Gorenstein ideals whose graded minimal resolution is G_\bullet . By Theorem 3.6 in [RZ3], there exist Gorenstein ideals I_G whose graded minimal resolution is G_\bullet , containing a complete intersection ideal I_Z of type (d^*, d_2, d_3) . The ideal $I'_Q = I_Z : I_G$ is an almost Gorenstein ideal with the same graded Betti numbers as I_Q . \square

We have that

$$F_3^\vee(-\vartheta_Z) \cong \bigoplus_{i=1}^t R(d - s_i - \vartheta_Z) = \bigoplus_{i=1}^t R(-(s_i - d_1)),$$

so by Theorem 4.8 and Theorem 4.10, $(d^*, s_1 - d_1, \dots, s_t - d_1)$ is a sequence of the degrees of the minimal generators of a Gorenstein ideal of codimension 3.

If $(a_1, \dots, a_n) \in \mathbb{Z}^n$, we define

$$\text{ord}(a_1, \dots, a_n) = (a_{\sigma(1)}, \dots, a_{\sigma(n)})$$

the permutation such that $a_{\sigma(1)} \leq \dots \leq a_{\sigma(n)}$.

We set $(s'_1, \dots, s'_{t+1}) = \text{ord}(d^* + d_1, s_1, \dots, s_t)$. So

$$(s'_1 - d_1, \dots, s'_{t+1} - d_1)$$

is the ordered sequence of the degrees of the minimal generators of a Gorenstein ideal of codimension 3.

Moreover we need to compare d^* with d_2 and d_3 . So we define

$$p(d^*) = \begin{cases} 1 & \text{if } d^* \leq d_2 \\ 2 & \text{if } d_2 < d^* \leq d_3 \\ 3 & \text{if } d_3 < d^* \end{cases}$$

Theorem 4.11. *If t is even, let us consider a resolution of the type*

$$0 \rightarrow \bigoplus_{i=1}^t R(-(d - s_i)) \rightarrow \bigoplus_{i=2}^3 R(-(d_i + d^*)) \oplus \bigoplus_{i=1}^{t+1} R(-s'_i) \rightarrow \\ \rightarrow R(-d^*) \oplus \bigoplus_{i=1}^3 R(-d_i) \rightarrow R \rightarrow A_Q \rightarrow 0$$

where $d = d_1 + d_2 + d_3 + d^*$, $d_1 \leq d_2 \leq d_3$, $s_1 \leq \dots \leq s_t$, $(s'_1, \dots, s'_{t+1}) = \text{ord}(d^* + d_1, s_1, \dots, s_t)$. Furthermore let

$$(m_1, m_2, m_3) = \min(s'_1 - d_1, \dots, s'_{t+1} - d_1), \quad m_1 \leq m_2 \leq m_3,$$

$p = p(d^*)$ and $(e_1, e_2, e_3) = \text{ord}(d^*, d_2, d_3)$.

It is the graded minimal free resolution of an almost complete intersection algebra of codimension 3 if and only if

- 1) $\sum_{i=1}^t s_i = \frac{t}{2}d - d^*$;
- 2) $d > s'_i + s'_{t+3-i}$, for $2 \leq i \leq t+1$;
- 3) $m_p \leq d^*$ and $m_i < e_i$ for $1 \leq i \leq 3$, $i \neq p$.

Proof. If it is the graded minimal free resolution of an almost complete intersection algebra $A_Q = R/I_Q$ of codimension 3, then the first condition holds by Proposition 4.2. The second condition follows by the Gaeta conditions, i.e. we have that

$$\begin{aligned} \vartheta_G > s'_i - d_1 + s'_{t+3-i} - d_1 &\iff d^* + d_2 + d_3 - d_1 > s'_i - d_1 + s'_{t+3-i} - d_1 \\ &\iff d > s'_i + s'_{t+3-i} \text{ for } 2 \leq i \leq t+1. \end{aligned}$$

The third condition follows from the fact that there exists a Gorenstein ideal I_G with minimal generators of degrees $(s'_1 - d_1, \dots, s'_t - d_1)$, linked to I_Q , in a complete intersection of type (d^*, d_2, d_3) , with the generator of degree d^* minimal for I_G and the generators of degrees d_2 and d_3 not minimal for I_G .

Conversely if the three conditions hold, then we can build a Gorenstein ideal I_G whose minimal generators have degrees $(s'_1 - d_1, \dots, s'_t - d_1)$ and such that it is possible to link I_G in a complete intersection I_Z of type (d^*, d_2, d_3) , with the generator of degree d^* minimal for I_G and the generators of degrees d_2 and d_3 not minimal for I_G . The ideal $I_Q = I_Z : I_G$ is the almost complete intersection ideal whose graded minimal free resolution is the assigned one. \square

If t is odd we can proceed analogously.

Theorem 4.12. *If t is odd, let us consider a resolution of the type*

$$\begin{aligned} 0 \rightarrow \bigoplus_{i=1}^t R(-(d - s_i)) \rightarrow R(-(d_2 + d_3)) \oplus \bigoplus_{i=1}^{t+2} R(-s'_i) \rightarrow \\ \rightarrow R(-d^*) \oplus \bigoplus_{i=1}^3 R(-d_i) \rightarrow R \rightarrow A_Q \rightarrow 0 \end{aligned}$$

where $d = d_1 + d_2 + d_3 + d^*$, $d_1 \leq d_2 \leq d_3$, $s_1 \leq \dots \leq s_t$, $(s'_1, \dots, s'_{t+2}) = \text{ord}(d_1 + d_2, d_1 + d_3, s_1, \dots, s_t)$. Furthermore let

$$(m_1, m_2, m_3) = \min(s'_1 - d_1, \dots, s'_{t+2} - d_1), \quad m_1 \leq m_2 \leq m_3,$$

$p = p(d^*)$ and $(e_1, e_2, e_3) = \text{ord}(d^*, d_2, d_3)$.

It is the graded minimal free resolution of an almost complete intersection algebra of codimension 3 if and only if

- 1) $\sum_{i=1}^t s_i = \frac{t-1}{2}d + d^*$;
- 2) $d > s'_i + s'_{t+4-i}$, for $2 \leq i \leq t+2$;
- 3) $m_p < d^*$ and $m_i \leq e_i$ for $1 \leq i \leq 3$, $i \neq p$.

Proof. Analogous to the proof of Theorem 4.11. \square

Corollary 4.13. *Let $A_Q = R/I_Q$ be an almost complete intersection algebra of codimension 3, whose Cohen-Macaulay type is $t_Q = 2$.*

Let

$$\begin{aligned} 0 \rightarrow R(-(d-s_1)) \oplus R(-(d-s_2)) &\rightarrow R(-s_1) \oplus R(-s_2) \oplus \\ &\oplus R(-(d_1+d^*)) \oplus R(-(d_2+d^*)) \oplus R(-(d_3+d^*)) \rightarrow \\ &\rightarrow R(-d_1) \oplus R(-d_2) \oplus R(-d_3) \oplus R(-d^*) \rightarrow I_Q \rightarrow 0. \end{aligned}$$

be its graded minimal free resolution, where $d = d_1 + d_2 + d_3 + d^*$, $d_1 \leq d_2 \leq d_3$, $s_1 \leq s_2$.

Then I_Q has the same graded Betti numbers of the monomial ideal

$$J = (x^{d_2}, y^{d_3}, z^{d^*}, x^{s_2-d_3}y^{s_1-d_2}).$$

Proof. By condition 1 in Theorem 4.11

$$s_1 + s_2 = d_1 + d_2 + d_3 \iff (s_1 - d_2) + (s_2 - d_3) = d_1.$$

By condition 3 in Theorem 4.11

$$s_1 - d_1 < d_2 \text{ and } s_2 - d_1 < d_3$$

i.e.

$$d_2 + d_3 - s_2 < d_2 \text{ and } d_2 + d_3 - s_1 < d_3 \iff d_3 < s_2 \text{ and } d_2 < s_1.$$

So we can consider the ideal $J = (x^{d_2}, y^{d_3}, z^{d^*}, x^{s_2-d_3}y^{s_1-d_2})$. Using Example 4.6 we can verify that the graded Betti numbers of J are the assigned ones. \square

When the Cohen-Macaulay type is $t_Q = 3$, the graded Betti numbers of an almost complete intersection algebra not always can be realized by a monomial ideal.

Example 4.14. Let us consider the ideal $I_Q = (x^2, xz+y(x+y+z), ax+by, f)$, with a and b general linear forms and f general quintic form. It has the following graded minimal free resolution

$$0 \rightarrow R(-5)^2 \oplus R(-7) \rightarrow R(-4)^4 \oplus R(-6)^2 \rightarrow R(-2)^3 \oplus R(-5) \rightarrow I_Q \rightarrow 0.$$

We have that $d^* = 5$. In Example 4.7 we computed the resolution of whatever artinian monomial almost complete intersection ideal of codimension 3 with $t_Q = 3$. Using the same notation we must have

$$b_1 + b_2 + b_3 = 5 \Rightarrow b_1 = 1, b_2 = 2, b_3 = 2,$$

(up to change the names of b_i 's). So the ideal should be

$$J = (x^2, y^2, z^2, xy^2z^2) = (x^2, y^2, z^2),$$

a contradiction.

Now we characterize, in the case $t_Q = 3$, the graded Betti numbers which can be realized using a monomial ideal.

Proposition 4.15. *Let $A_Q = R/I_Q$ be any almost complete intersection algebra of codimension 3, $t = t_Q = 3$. Let*

$$\begin{aligned} 0 \rightarrow \bigoplus_{i=1}^3 R(-(d - s_i)) \rightarrow \bigoplus_{1 \leq i < j \leq 3} R(-(d_i + d_j)) \oplus \bigoplus_{i=1}^3 R(-s_i) \rightarrow \\ \rightarrow R(-d^*) \oplus \bigoplus_{i=1}^3 R(-d_i) \rightarrow R \rightarrow A_Q \rightarrow 0 \end{aligned}$$

be its graded minimal free resolution, with $d_1 \leq d_2 \leq d_3$, $s_1 \leq s_2 \leq s_3$.

Then the graded Betti numbers of I_Q can be realized by an artinian monomial ideal iff

$$d^* < s_i < d^* + d_i, \text{ for } 1 \leq i \leq 3.$$

Proof. Let us suppose that the graded Betti numbers of I_Q can be realized by a monomial ideal $J = (x^{a_1}, y^{a_2}, z^{a_3}, x^{b_1}y^{b_2}z^{b_3}) \subset k[x, y, z]$, with $0 < b_i < a_i$, for $i = 1, 2, 3$. We set

$$v_1 = a_1 + b_2 + b_3, v_2 = b_1 + a_2 + b_3, v_3 = b_1 + b_2 + a_3.$$

We have that $(s_1, s_2, s_3) = \text{ord}(v_1, v_2, v_3)$. Note that $d^* < v_i < d^* + d_i$ for $1 \leq i \leq 3$, therefore $d^* < s_i$ for $1 \leq i \leq 3$. Moreover $s_1 \leq v_1 < d^* + d_1$.

Now we consider s_2 . If $s_2 = v_i$ per $i \leq 2$ we deduce that

$$s_2 = v_i < d^* + d_i \leq d^* + d_2.$$

If $s_2 = v_3$ then $s_3 = v_i$ with $i \leq 2$, so

$$s_2 \leq s_3 = v_i < d^* + d_i \leq d^* + d_2.$$

Finally

$$s_3 = v_i < d^* + d_i \leq d^* + d_3.$$

Conversely if $d^* < s_i < d^* + d_i$, for $1 \leq i \leq 3$, we can consider the monomial ideal

$$J = (x^{d_1}, y^{d_2}, z^{d_3}, x^{d^*+d_1-s_1}y^{d^*+d_2-s_2}z^{d^*+d_3-s_3}).$$

It has 4 minimal generators since $d^* < s_i$ implies that $d^* + d_i - s_i < d_i$. Moreover the exponents are all positive since $s_i < d^* + d_i$. It is easy to check, using Example 4.7, that J has the requested graded Betti numbers. \square

REFERENCES

- [BE] David A. Buchsbaum, David Eisenbud (1977) *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3*, American Journal of Mathematics **99**(3), 447–485.
- [Bu] Lindsay Burch (1968) *On ideals of finite homological dimension in local rings*, Mathematical Proceedings of the Cambridge Philosophical Society (Vol. 64, No. 4, pp. 941–948). Cambridge University Press.
- [CGO] Ciro Ciliberto, Anthony V. Geramita, Ferruccio Orecchia (1988) *Remarks on a theorem of Hilbert-Burch* Bollettino della Unione Matematica Italiana, 2(3), 463–483.
- [CV] Aldo Conca, Giuseppe Valla (2000) *Betti numbers and lifting of Gorenstein codimension three ideals* Communications in Algebra, 28:3, 1371–1386, DOI: 10.1080/00927870008826900
- [Di] Susan Diesel (1996) *Irreducibility and dimension theorems for families of height 3 Gorenstein algebras*, Pacific Journal of Mathematics **172**(2), 365–397.

- [Ga] Federico Gaeta (1950) *Nuove ricerche sulle curve sghembe algebriche di residuale finito e sui gruppi di punti del piano* Annali di Matematica 31(4), 1-64, DOI: 10.1007/BF02428254
- [Hi] David Hilbert (1890) *Über die Theorie von algebraischen Formen*, Mathematische Annalen 36, 473–534, DOI: 10.1007/BF01208503
- [KM] Andrew R. Kustin, Matthew Miller (1983) *Constructing big Gorenstein ideals from small ones*, Journal of Algebra 85(2), 303–322, DOI: 10.1016/0021-8693(83)90096-0
- [Re] Miles Reid (2015) *Gorenstein in codimension 4: the general structure theory* Algebraic geometry in east AsiaTaipei 2011, 201–227, Mathematical Society of Japan DOI: 10.2969/aspm/06510201
- [RZ1] Alfio Ragusa, Giuseppe Zappalà (2013) *Characterization of the Graded Betti Numbers for Almost Complete Intersections*, Communications in Algebra, 41:2, 492–506, DOI: 10.1080/00927872.2011.558150
- [RZ2] Alfio Ragusa, Giuseppe Zappalà (2001) *Properties of 3-codimensional Gorenstein schemes*, Communications in Algebra, 29:1, 303–318, DOI: 10.1081/AGB-100000802
- [RZ3] Alfio Ragusa, Giuseppe Zappalà (2011) *On complete intersections contained in Cohen-Macaulay and Gorenstein ideals*, Algebra Colloquium 18 (Spec 1), 857–872
- [RZ4] Alfio Ragusa, Giuseppe Zappalà (2013) *On the structure of resolutions of almost complete intersections*, International Journal of Pure and Applied Mathematics 83(1), 13–23. DOI: 10.12732/ijpam.v83i1.3

DIPARTIMENTO DI MATEMATICA E INFORMATICA,
UNIVERSITÀ DI CATANIA,
VIALE A. DORIA 6, 95125 CATANIA, ITALY
E-mail address: zappalag@dmf.unict.it