A Divide and Conquer Algorithm of Bayesian Density Estimation

Ya Su

Department of Statistics, University of Kentucky, Lexington, KY 40536-0082, U.S.A., ya.su@uky.edu

Abstract

Data sets for statistical analysis become extremely large even with some difficulty of being stored on one single machine. Even when the data can be stored in one machine, the computational cost would still be intimidating. We propose a divide and conquer solution to density estimation using Bayesian mixture modeling including the infinite mixture case. The methodology can be generalized to other application problems where a Bayesian mixture model is adopted. The proposed prior on each machine or subsample modifies the original prior on both mixing probabilities as well as on the rest of parameters in the distributions being mixed. The ultimate estimator is obtained by taking the average of the posterior samples corresponding to the proposed prior on each subset. Despite the tremendous reduction in time thanks to data splitting, the posterior contraction rate of the proposed estimator stays the same (up to a log factor) as that of the original prior when the data is analyzed as a whole. Simulation studies also justify the competency of the proposed method compared to the established WASP estimator in the finite dimension case. In addition, one of our simulations is performed in a shape constrained deconvolution context and reveals promising results. The application to a GWAS data set reveals the advantage over a naive method that uses the original prior.

Some Key Words: Divide and conquer, Bayesian density estimation, Posterior contraction rate, Bayesian mixture model.

Short title:

1 Introduction

In an era of real big data, data sets for statistical analysis become extremely large even with some difficulty of storing on one single machine. Even when the data can be stored in one machine, the computational cost would still be intimidating. As an example, for most recent data sets in the Genome Wide Association Study (GWAS), the number of subjects amounts to several hundreds of thousands while the number of single-nucleotide polymorphism (SNP) goes up to one million for each individual.

A divide and conquer algorithm involves three steps. First, a partition of X_1, \ldots, X_n is distributed to J machines. For simplicity, we assume that the data is randomly partitioned with equal size so that the sample size on each machine is m = n/J. Second, individual analysis is performed to the subset data on each machine, usually in a paralleled fashion. The last step is to combine the estimators from all J machines. The computational cost of a divide and conquer algorithm is reduced tremendously thanks to the paralleled analyses on much smaller data sets. The reduction in time could be significant if the complexity of the statistical analysis is of first or higher order of sample size. In the Bayesian framework, different approaches arise in this context for various purposes. To name a few, Scott et al. (2016) came up with a simple procedure in terms of both assigning prior and combining posterior samples. Srivastava et al. (2018) unified all posterior distributions on each subset, leading to an overall posterior distribution that maintains the same concentration rate as if the whole data has been treated together. Sabnis et al. (2016) made a first step in subsetting the variables in a Bayesian factor model. Guhaniyogi et al. (2017) proposed distributed kriging for Gaussian process in spatial data.

It is well acknowledged in both frequentist and Bayesian perspectives that some debiasing or overfitting procedure needs to be done when analyzing the subset data, in order to obtain a combined estimator that achieves the same accuracy as that of the original estimator when the data is analyzed as a whole. Methods are distinguished by how the individual analysis is appropriately adjusted and the way that the estimators in different machines are combined. Here our attention is given to several recent Bayesian approaches. Within the context of signal-in-white-noise model, Szabó and van Zanten (2019) pointed out the necessity of carefully choosing among several strategies for successfully achieving the optimal convergence rate and posterior coverage probability. Scott et al. (2016) and Neiswanger et al. (2014) proposed a general framework for modifying the prior in the divide and conquer context and applied the method in various setups. Other approaches (Srivastava et al., 2018; Xue and Liang, 2019) concern modifying the likelihood in evaluating the posterior distribution together with combination techniques that find the "center" of the posterior distributions on each subset. However, these methods require caution to use when the parameter is of large or infinite dimension, with the combination strategy too simple to be justified or too complicated to compute. In addition, these combination techniques are also impossible to apply when the goal is to estimate a density on a non-Euclidean space, e.g. space of densities. A comparison with this estimator, named WASP, is of merit in the low dimension case.

Of the existing methods using a Bayesian procedure along with a data-splitting technique, none of them estimate an infinite dimensional parameter nor do they deal with prior distributions over an infinite dimensional space. Problems of this kind arise naturally in nonparametric density estimation and become attractive in high dimensional and nonlinear models, see Section 2.5 for an incomplete list of references. In such problems, the most popular choice of prior one can expect is the Dirichlet process mixture of standard densities, like Normal, Laplace, Gamma, etc. Following the success of Scott et al. (2016) in examples with a finite dimensional parameter or a simple conjugate prior, we are motivated to take a step into more complicated scenarios, for instance, when the prior belongs to the specific type as above.

In the divide and conquer framework, we propose a general methodology for assigning such priors with a focus on density estimation. The proposed prior generalizes the idea when the parameter is finite dimensional, that is, the prior is adjusted with the purpose of debiasing the subset density estimators by sacrificing the uncertainty therein. We provide the solutions to adjust priors having a Dirichlet process component and more. We use simple averaging for combining the individual estimators which facilitates the computation and meanwhile reduces excessive uncertainty without introducing bias. The ultimate density estimator after combining the individual estimators is constructed under much less computational and memory burden while still achieving the optimal rate. We show in simulations and real data that the proposed procedure is also applicable to contexts beyond density estimation. Included in this paper is a confirmed success in a density deconvolution problem. By its design, the method fits easily into other contexts as long as the prior itself is composed of a(n) finite/infinite mixture of standard probability distributions and others.

The following sections are organized in this way. Section 2 specifies two example models regarding Bayesian density estimation and the proposed priors in the divide and conquer context. In Section 3 we show that the proposed density estimator can achieve the optimal rate when the number of subsets is growing no faster than log rate of sample size. Two simulations are conducted in Section 4, illustrating the competency of our method with WASP in density estimation and the capability of an extension to density deconvolution which is motivated by a real application. The proposed method is implemented on a data set in genome-wide association study (GWAS) and results are presented in Section 5. The paper ends with discussions in Section 6.

Notations. IW(ν, Ψ) denotes an inverse wishart distribution with degrees of freedom ν and scale matrix Ψ . Ga(a, b)/IG(a, b) stands for a(n) gamma/inverse gamma distribution with shape a and rate/scale b. Dir(α) denotes a Dirichlet distribution of order K with parameter $\alpha = \{\alpha_1, \ldots, \alpha_K\}$. N_p(μ, Σ) stands for a multivariate normal distribution of dimension p with mean μ and covariance matrix Σ and N(μ, σ^2) in the case p = 1. DP(M, G) denotes a Dirichlet process with concentration parameter M and base probability measure G. Unif(θ_1, θ_2) is a uniform distribution supported on the interval [θ_1, θ_2]. All of the distributions above can be easily switched to a density by adding a dot argument as the first argument, e.g. N_p($\cdot; \mu, \Sigma$) denotes the density of N_p(μ, Σ). By convention, ϕ_{σ} refers to the density function of a univariate Normal distribution with mean 0 and standard deviation σ . The expression $a_n \approx b_n$ states that a_n and b_n are of the same rate asymptotically.

2 Model Specification

2.1 Background

Suppose $X_1, \ldots, X_n \in \mathbb{R}^p$ is an independent and identically distributed sample from an unknown density $f_0(\cdot), \mathbb{R}^p \to [0, \infty)$. We are going to illustrate our idea under both parametric and nonparametric model for $f_0(\cdot)$, both characterized by a mixtures of normal distributions. The key idea can be easily extended to mixtures of distributions other than normal.

We first consider $f_0(\cdot)$ is a finite dimensional mixture of normal distributions,

$$f_0(x) = \sum_{k=1}^{K} \pi_k \mathcal{N}_p(x; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k).$$
(1)

The mixing probabilities $\boldsymbol{\pi} = (\pi_1, \dots, \pi_K)$ lie in a (L-1)-simplex. The *k*th component normal distribution has mean and covariance matrix $\boldsymbol{\mu}_k$ and $\boldsymbol{\Sigma}_k$. Together, $\boldsymbol{\pi}$, $\{\boldsymbol{\mu}_k\}_{k=1}^K$ and $\{\boldsymbol{\Sigma}_k\}_{k=1}^K$ form the unknown parameters in the data generating model.

For ease of computation, a conjugate prior corresponding to (1) can be imposed (Srivastava et al., 2018):

$$\boldsymbol{\pi} \sim \operatorname{Dir}(\alpha_1, \dots, \alpha_K), \quad \boldsymbol{\mu}_k \mid \boldsymbol{\Sigma}_k \sim \operatorname{N}_p(\boldsymbol{0}, l\boldsymbol{\Sigma}_k), \quad \boldsymbol{\Sigma}_k \sim \operatorname{IW}(\nu, \boldsymbol{S}).$$
 (2)

On the other hand, if the form of the true density $f_0(\cdot)$ is unknown, in which case a nonparametric counterpart to the finite dimensional model (1) and (2) is a popular substitute. We are going to present the univariate case, a straightforward extension of the current algorithm and theory to a multivariate case exists; see Remark 1 for a brief discussion about the theory about the multivariate case. Specifically, the nonparametric model is

$$f(x) \sim \int \phi_{\sigma}(x-\mu)P(d\mu), \quad \sigma \sim \Pi_{\sigma}(\cdot), \quad P \sim \mathrm{DP}(M,G).$$
 (3)

The model (3) corresponds to the so-called Dirichlet process (location) mixtures of Normal (DPMN) prior, algorithms of which have been studied previously (Blei et al., 2006; Rasmussen, 2000). The asymptotics about DPMN have been investigated in Ghosal et al. (2007) for the univariate case and Shen et al. (2013) for the multivariate case.

Given the proven performance of these priors in producing a good density estimator while running on the complete data, the following sections will provide guidance on imposing priors when we work on small chunks of data spread across various machines. Before illustrating our approach regarding the density estimation problems above, we first present a general way which has been explored for models with finite dimensional parameters (Scott et al., 2016). Denote θ be the parameter of interest, $L(\theta; \boldsymbol{x})$ be the likelihood function based on data \boldsymbol{x} , $\Pi(\theta)$ be the prior on θ . In the distributed setting with J chunks, the likelihood function can be decomposed into J components, $L(\theta; \boldsymbol{x}) = \prod_{j=1}^{J} L(\theta; \boldsymbol{x}^{(j)})$, where $\boldsymbol{x}^{(j)}$ is the data belonging to the *j*th chunk. The posterior distribution of θ takes the form $\Pi(\theta \mid \boldsymbol{x}) =$ $\prod_{j=1}^{J} \{L(\theta; \boldsymbol{x}^{(j)}) \Pi(\theta)^{1/J}\}$ where the likelihood function and the prior are factorized similarly. This general idea paves the way of seeking for an appropriate prior on each chunk of data by assigning $\Pi(\theta)^{1/J}$.

Difficulties exist on how to justify the above idea for all cases of $\Pi(\theta)$ where θ could be infinite dimension. The focus of this paper is to address this issue for a family of models including but not limited to (1) and (3). It is seemingly hard to handle priors with its support on a probability space with the existing literature because these priors involve a a "distribution on distribution" component corresponding to the Dirichlet distribution or the Dirichlet process prior and some independent prior distributions on the rest of parameters in the component densities. In what follows we describe the modification to these priors in the divide and conquer context. The prior on the parameters in component distributions which usually takes a conjugate form against the likelihood will be imposed as the same type. On the other hand, we propose to make a simple adjustment on the parameters of the Dirichlet prior from the property of Dirichlet distribution or process.

2.2 Finite mixtures of Normal prior

As introduced in Section 2.1, the prior for a finite mixture of normal model takes the form (2). We start with a basic property of the Dirichlet distribution.

Proposition 1. Let $\boldsymbol{\pi} \sim \text{Dir}(\boldsymbol{\alpha})$, where $\boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_K)$. Denote $\alpha_s = \sum_{k=1}^K \alpha_k$. Then $E(\pi_k) = \alpha_k/\alpha_s$, $\operatorname{var}(\pi_k) = \alpha_k(\alpha_s - \alpha_k)/\alpha_s^2(\alpha_s + 1)$, $\operatorname{cov}(\pi_j, \pi_k) = -\alpha_j \alpha_k/\alpha_s^2(\alpha_s + 1)$.

The Dirichlet distribution prior on $\boldsymbol{\pi}$, $\text{Dir}(\alpha_1, \ldots, \alpha_K)$, is proportional to $\prod_{k=1}^K \pi_k^{\alpha_k-1}$. Proposition 1 (a proof can be found in Chapter 27 of Balakrishnan and Nevzorov (2004)) states that scaling $\boldsymbol{\alpha}$ by a factor γ scales up the variance by a factor $(1 + \alpha_s)/(1 + \gamma \alpha_s)$ while keeping the mean unchanged component-wise. This property is essential and provides directions to adjust the prior on $\boldsymbol{\pi}$. We propose to scale the parameters of the Dirichlet distribution prior on $\boldsymbol{\pi}$ by 1/J leading to $\text{Dir}(\alpha_1/J, \ldots, \alpha_K/J)$. Indeed, we can make an assertion that this simple adjustment sacrifices uncertainty in exchange for debiasing regarding $\boldsymbol{\pi}$ according to Proposition 1.

We take the same strategy as Concensus Monte Carlo (Scott et al., 2016) for adjusting the priors on the remaining parameters. Raising a power 1/J to the normal prior on $\boldsymbol{\mu}_k \mid \boldsymbol{\Sigma}_k$ leads to $N_p(0, l\boldsymbol{\Sigma}_k J)$. It can be shown easily that the effect of a power 1/J to the inverse wishart prior on $\boldsymbol{\Sigma}_k$, IW(ν, \boldsymbol{S}), is chracterized by an inverse wishart type IW($\nu/J-(p+1)(J-1)/J, \boldsymbol{S}/J$). The risk of its first parameter being possibly negative will be regulated by the likelihood when constructing the (conditional) posterior distribution of $\boldsymbol{\Sigma}_k$. A Gibbs sampler corresponding to the modified prior above is provided in Section S.1.1 in the Supplementary material.

2.3 Dirichlet process mixtures of Normal prior

Although a Dirichlet process has a remarkable stick-breaking representation (Sethuraman, 1994), unfortunately, it does not have a probability density as a Dirichlet distribution does. However, we can extend the idea in Section 2.2 to Dirichlet process since marginally a Dirichlet process follows a Dirichlet distribution. That is, if $P \sim DP(M, G)$, for any measurable finite partition $\{A_1, \ldots, A_k\}$ of the support of the base measure G, $(P(A_1), \ldots, P(A_k)) \sim Dir(G(A_1)M, \ldots, G(A_k)M)$.

Our idea is to modify the parameters associated with a Dirichlet process such that the relationships between its subsequent marginal distributions and those under the Dirichlet process with the original parameters are maintained to be the same as that in Section 2.2. This can achieved by adjusting the prior on P as DP(M/J, G) under which $(P(A_1), \ldots, P(A_k)) \sim Dir(G(A_1)M/J, \ldots, G(A_k)M/J)$ for the above partition. Hence regarding the nonparametric model (3), the following prior is suggested:

$$f(x) \sim \int \phi_{\sigma}(x-\mu)P(d\mu), \quad \sigma \propto \Pi_{\sigma}^{1/J}(\cdot), \quad P \sim \mathrm{DP}(M/J,G).$$
 (4)

The general form of the prior on σ , $\Pi_{\sigma}^{1/J}$, can be simplified if Π_{σ} takes a parametric form. In the case when a conjugate prior for σ , $\Pi_{\sigma} \sim \text{IG}(a,b)$, is adopted, it becomes $\Pi_{\sigma}^{1/J} \sim \text{IG}(a/J - (J-1)/J, b/J)$. Indeed it is just an inverse gamma type since the first argument is negative when a < (J-1), but similar arguments about the prior on Σ_k in Section 2.2 apply here. A Gibbs sampler corresponding to (4) is provided in Section S.1.2 in the Supplementary material.

2.4 A combined density estimator

Let \mathbf{X}_j be the subset of data distributed to the *j*th machine, $j = 1, \ldots, J$. Denote the posterior probability under the model in Section 2.2 or 2.3 in accordance to the *j*th subset as $\Pi_m(\cdot \mid \mathbf{X}_j)$, where subscript *m* indicates the distributed sample size. Our procedure proceeds as follows. For each subset *j*, we could obtain an estimator $f^{[j]}(\cdot) \in \mathcal{P}$ by taking a random sample from the posterior distribution $\Pi_m(\cdot \mid \mathbf{X}_j)$. The ultimate estimator is then formed by a simple average over all subset samples, $\bar{f}(\cdot) = (1/J) \sum_{j=1}^{J} f^{[j]}(\cdot)$.

Let $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$ denote the distribution of $\overline{f}(\cdot)$. Then it is easy to show that $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$ is a convex convolution of all subset posterior densities, $\Pi_m(\cdot \mid \mathbf{X}_j)$, $j = 1, \ldots, J$. Although we will provide asymptotics of $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$ in Section 3.2, we discuss the appropriateness of proposing $\overline{f}(\cdot)$, or equivalently $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$. The major consequence of us modifying the prior is that the center of $\Pi_m(\cdot \mid \mathbf{X}_j)$ is pulled towards the targeted posterior distribution while admitting larger variability. We construct the aforementioned combined density /posterior distribution as it keeps the center unchanged while reducing the variability in subset posterior distributions.

2.5 Other applications

Our divide and conquer algorithm goes beyond density estimation problem as long as the prior of the relevant model consists of a Dirichlet distribution/process component, which is often seen along with models characterized by a(n) finite/infinite mixture of standard probability distributions. The popularity of such prior has risen in recent years with appearances in high dimensional normal means problem (Bhattacharya et al., 2015), multivariate cat-

egorical data with dependency (Dunson and Xing, 2009), and nonlinear regression models (De Jonge et al., 2010; Naulet et al., 2018), just to name a few.

3 Theoretical results

3.1 Preliminary definitions

The set of density functions is $\mathcal{P} = \{f(\cdot) : f(\cdot) \ge 0, \int f(x)dx = 1\}$. We consider the metric on \mathcal{P} to be the Hellinger distance $h(\cdot, \cdot)$. For any $p, q \in \mathcal{P}, h(p,q) = \{\int (p^{1/2}(x) - q^{1/2}(x))^2 dx\}^{1/2}$. The Wasserstein space of order 2 is defined as $\mathscr{P}_2 = \{\mu \in \text{probability measure on } \mathcal{P} : \int_{\mathcal{P}} h^2(f, f_0) d\mu(f) < \infty\}$. For any $\mu, \nu \in \mathscr{P}_2, \ \Xi(\mu, \nu)$ is a set of all probability measures on $\mathcal{P} \times \mathcal{P}$ whose marginal measures are μ and ν . The Wasserstein distance of order 2 on \mathscr{P}_2 is defined as $W_2^2(\mu, \nu) = \inf_{\xi \in \Xi(\mu, \nu)} \int_{\mathcal{P} \times \mathcal{P}} h^2(p, q) d\xi(p, q)$.

In particular, if one of the probability measures is concentrated on a fixed element in $f_0 \in \mathcal{P}$, e.g., $\nu = \delta_{f_0}$, the Wasserstein metric becomes $W_2^2(\mu, \nu) = \int_{\mathcal{P}} h^2(f, f_0) d\mu(f)$. Thus in the context of this paper, the Wasserstein distance between any posterior distribution $\Pi_n(\cdot \mid \mathbf{X}_n)$ on \mathcal{P} and δ_{f_0} is

$$W_2^2(\Pi_n, \delta_{f_0}) = \int_{\mathcal{P}} h^2(f, f_0) d\Pi_n(f \mid \boldsymbol{X}_n).$$
 (5)

Hence

$$W_2^2(\Pi_n, \delta_{f_0}) \le \epsilon_n^2 + 2\Pi_n(\{f \in \mathcal{P} : h(f, f_0) \ge \epsilon_n\} \mid \boldsymbol{X}_n).$$

The above inequality is due to the fact that the Hellinger distance is less or equal to 2. Hence we conclude that a typical posterior contraction rate result, $\Pi_n(\{f \in \mathcal{P} : h(f, f_0) \ge \epsilon_n\} \mid \mathbf{X}_n) \to 0$, is sufficient to prove a convergence of $\Pi_n(\cdot \mid \mathbf{X}_n)$ to δ_{f_0} with rate ϵ_n in Wasserstein distance. In Section 3.2 we will present our theoretical results in terms of the latter.

Lemma 1. The Hellinger distance is a biconvex functional on $\mathcal{P} \times \mathcal{P}$, that is, for any $g_1, g_2, f \in \mathcal{P}$, and $\omega_1 \ge 0, \omega_2 \ge 0$ with $\omega_1 + \omega_2 = 1$, $h(\omega_1 g_1 + \omega_2 g_2, f) \le \omega_1 h(g_1, f) + \omega_2 h(g_2, f)$. Similarly, $h(f, \omega_1 g_1 + \omega_2 g_2) \le \omega_1 h(f, g_1) + \omega_2 h(f, g_2)$.

3.2 Main theorems

To see the asymptotic behavior of $\overline{f}(\cdot)$, equivalently, we can study the underlying distribution which yields $\overline{f}(\cdot)$, $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$. By definition of $\overline{f}(\cdot)$, $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$ is a (convex) convolution of the subset posterior distributions $\Pi_m(\cdot \mid \mathbf{X}_j), j = 1, \ldots, J$. Specifically, for any functional $L(\cdot)$ on \mathcal{P} ,

$$\int_{\mathcal{P}} L(f) d\overline{\Pi}_n(f \mid \boldsymbol{X}_n) = \int_{\mathcal{P} \times \dots \times \mathcal{P}} L(\sum_{j=1}^J f_j / J) d\Pi_m(f_1 \mid \boldsymbol{X}_1) \cdots d\Pi_m(f_J \mid \boldsymbol{X}_J).$$
(6)

It is trivial to show that $\overline{f}(\cdot)$ corresponds to a sample drawn from $\overline{\Pi}_n(\cdot \mid \boldsymbol{X}_n)$.

We are going to state the posterior contraction rate for $\overline{\Pi}_n(\cdot \mid \mathbf{X}_n)$. For illustration purposes, we are going to state the theory for the nonparametric density estimator under the Dirichlet process mixtures of normal prior (4).

Let $a_1, \ldots, a_5, b_1, \ldots, b_5$ and C_1, \ldots, C_4 be positive constants. Denote C^{β, L, τ_0} as the locally β -Hölder function class with functions that have finite partial derivatives $f^{(k)}(\cdot)$ up to order $k \leq \lfloor \beta \rfloor$ such that for all $k \leq \lfloor \beta \rfloor$, $|f^{(k)}(x+y) - f^{(k)}(x)| \leq L(x) \exp(\tau_0 |y|^2) |y|^{\beta - \lfloor \beta \rfloor}$.

The key assumptions are

(C1) $1 - G([-x, x]) \le b_1 \exp(-C_1 x^{a_1})$ for sufficiently large x > 0.

(C2) $\Pi_{\sigma}\{(0,x)\} \leq b_2 \exp(-C_2 x^{-a_2})$, for sufficiently small x > 0. $\Pi_{\sigma}\{(x,\infty)\} \leq b_3 x^{-a_3}$, for sufficiently large x > 0. For any s, t > 0 $\Pi_{\sigma}\{(s^{-1}(1+t)^{-1/2}, s^{-1})\} \geq b_4 s^{-a_4} t^{a_5} \exp(-C_3 s^{-1})$. (C3) $f_0(\cdot) \in C^{\beta,L,\tau_0}$. $P_0(|D^k f_0|/f_0)^{(2\beta+\epsilon)/k} < \infty$ for all integer $k \leq \lfloor\beta\rfloor$. $P_0(L/f_0)^{(2\beta+\epsilon)/\beta} < \infty$. $f_0(x) \leq b_5 \exp(-C_4|x|^{\tau})$ for sufficiently large |x| and some $\tau > 0$. (C4) $J \asymp \log n$, equivalently, $m \asymp n \log^{-1} n$.

The first three conditions are the same as those in Shen et al. (2013) in the univariate case. The last condition is on the growing rate of subsets in our divide and conquer setup.

<u>**Theorem</u></u> 1. If assumptions (C1)—(C4) are satisfied, for any j = 1, ..., J and any t > 3/2 + (1/\tau)/(2+1/\beta), the posterior distribution \Pi_m(\cdot | \mathbf{X}_j) converges to f_0(\cdot) in Wasserstein distance with contraction rate \epsilon_m = m^{-\beta/(2\beta+1)}(\log m)^t, that is,</u>**

$$W_2(\Pi_m(\cdot \mid \boldsymbol{X}_j), \delta_{f_0}) \lesssim \epsilon_m, \quad j = 1, \dots, J.$$

Here δ_{f_0} is the dirac measure on \mathcal{P} concentrating at $f_0(\cdot)$.

The next lemma states how the convergence under the Wasserstein metric for each subset posterior distribution $\Pi_m(\cdot \mid \boldsymbol{X}_j)$ controls that of the posterior distribution $\overline{\Pi}_n(\cdot \mid \boldsymbol{X}_n)$.

<u>Lemma</u> 2. $W_2(\overline{\Pi}_n(\cdot \mid \boldsymbol{X}_n), \delta_{f_0}) \leq J^{-1} \sum_{j=1}^J W_2(\Pi_m(\cdot \mid \boldsymbol{X}_j), \delta_{f_0}).$

Proof. First, from the expressions (6), (5) and Lemma 1, it can be seen that

$$\begin{split} W_2^2(\overline{\Pi}_n(\cdot \mid \boldsymbol{X}_n), \delta_{f_0}) &= \int_{\mathcal{P} \times \dots \times \mathcal{P}} h^2(\sum_{j=1}^J f_j / J, f_0) d\Pi_m(f_1 \mid \boldsymbol{X}_1) \cdots d\Pi_m(f_J \mid \boldsymbol{X}_J) \\ &\leq J^{-2} \sum_{j=1}^J \int_{\mathcal{P}} h^2(f, f_0) d\Pi_m(f \mid \boldsymbol{X}_j) \\ &+ J^{-2} \sum_{j \neq k} \int_{\mathcal{P} \times \mathcal{P}} h(f_j, f_0) h(f_k, f_0) d\Pi_m(f_j \mid \boldsymbol{X}_j) d\Pi_m(f_k \mid \boldsymbol{X}_k). \end{split}$$

By Cauchy-Schwartz inequality, all terms in the second summation on the right hand side is bounded by $\{\int_{\mathcal{P}} h^2(f, f_0) d\Pi_m(f \mid \mathbf{X}_j) \int_{\mathcal{P}} h^2(f, f_0) d\Pi_m(f \mid \mathbf{X}_k)\}^{1/2}$. Applying expression (5) again to all terms in the previous display yields

$$W_{2}^{2}(\overline{\Pi}_{n},\delta_{f_{0}}) \leq J^{-2} \left\{ \sum_{j=1}^{J} W_{2}^{2}(\Pi_{m}(\cdot \mid \boldsymbol{X}_{j}),\delta_{f_{0}}) + \sum_{j \neq k} W_{2}(\Pi_{m}(\cdot \mid \boldsymbol{X}_{j}),\delta_{f_{0}}) W_{2}(\Pi_{m}(\cdot \mid \boldsymbol{X}_{k}),\delta_{f_{0}}) \right\}$$

$$= \left\{ J^{-1} \sum_{j=1}^{J} W_{2}(\Pi_{m}(\cdot \mid \boldsymbol{X}_{j}),\delta_{f_{0}}) \right\}^{2}.$$

<u>Theorem</u> 2. If assumptions (C1)—(C4) are satisfied, the posterior distribution $\overline{\Pi}_n(\cdot \mid \boldsymbol{X}_n)$ converges to $f_0(\cdot)$ in Wasserstein distance with contraction rate $\epsilon_n = n^{-\beta/(2\beta+1)}(\log n)^u, u > 3/2 + (1/\tau)/(2 + 1/\beta) + \beta/(2\beta + 1)$, that is,

$$W_2(\overline{\Pi}_n(\cdot \mid \boldsymbol{X}_n), \delta_{f_0}) \lesssim \epsilon_n.$$

Proof. The desired inequality is implied by Theorem 1 and Lemma 2.

<u>**Remark</u> 1.** In Theorem 2, we state the one dimensional case for simplicity. We make a brief comment without details that a multivariate version of Theorem 2 exists provided a multivariate correspondence to prior (4) is imposed. In the standard Bayesian density estimation setting, the contraction rate for estimating a multivariate density has been studied previously</u> in Shen et al. (2013). When switched to the multivariate case with dimension d, the correspondence to Π_{σ} would be a prior on the covariance matrix $\Sigma \in \mathbb{R}^{d \times d}$, on which Shen et al. (2013) imposed conditions regarding the concentration of eigenvalues of the matrix. Meanwhile, the base measure $G(\cdot)$ on \mathbb{R}^d is required to satisfy a straightforward multi-dimensional extension of (C1). Under these conditions, the logic flow in deriving Theorem 2 guarantees that our divide and conquer density estimator can achieve the multivariate contraction rate as if the complete data has been used. Details will be omitted.

4 Simulation

4.1 Overview

As briefed in Section 2.2–2.4, we come up with a proper way to rebuild a simple density estimator for some existing parametric and nonparametric methods in the divide and conquer context.

The finite mixtures of normals is one example considered in Srivastava et al. (2018) regarding model (1) and prior (2). For comparison, our first simulation will imitate Section 4.2 of Srivastava et al. (2018) in the choices of p = 2, K = 2, $\alpha_1 = \alpha_2 = 1/2$, l = 100, $\nu = 2$ and $\mathbf{S} = 4\mathbf{I}_2$. The true parameter values are set as $\boldsymbol{\pi} = (0.3, 0.7)$, $\boldsymbol{\mu}_1 = (1, 2)^T$, $\boldsymbol{\mu}_2 = (7, 8)^T$, $\boldsymbol{\Sigma}_1 = \boldsymbol{\Sigma}_2 = \{\boldsymbol{\Sigma}_{ij}\}_{1 \leq i,j \leq 2}$, where $\boldsymbol{\Sigma}_{12} = 0.5$, $\boldsymbol{\Sigma}_{11} = 1$ and $\boldsymbol{\Sigma}_{22} = 2$. As mentioned in Section 2.2, we propose the following adjusted prior on each machine:

 $\boldsymbol{\pi} \sim \operatorname{Dir}(1/(2J), 1/(2J)), \quad \boldsymbol{\mu}_k \mid \boldsymbol{\Sigma}_k \sim \operatorname{N}_2(\boldsymbol{0}, 100\boldsymbol{\Sigma}_k J), \quad \boldsymbol{\Sigma}_k \sim \operatorname{IW}(5/J - 3, (4/J)\boldsymbol{I}_2).$ (7)

Srivastava et al. (2018) mainly illustrated the ability of their method, WASP, on estimating some nonlinear functions of the parameters, say $g(\pi, \mu_1, \mu_2, \Sigma_1, \Sigma_2)$. They compared the accuracy of WASP with other methods, among which the superiors were Consensus Monte Carlo and WASP. Given their close competition in estimating the functions of the parameters, it is worthwhile to examine whether the performance is consistent in estimating the original model parameters. With the acknowledgment of the possible similarity of our method and Consensus Monte Carlo in finite dimensional models, we implement our algorithm using (7). A disclaimer is that we find there are differences with the implementation of CMC in Srivastava et al. (2018) (specifically in updating π and Σ_k). WASP estimator is obtained using online code of Srivastava et al. (2018).

The second simulation we conduct is in a more complicated context, as designed for density deconvolution with shape constraints. We generate observed data W_i through a classical measurement error model, specifically, an independent sample of W = X + U, where the distribution of U is known and possibly heteroscedastic. The density of the true variable $X, f(\cdot)$, is of interest and in some application context (see Section 5 for one such application) should have a symmetric and unimodal shape. To ensure the shape of $f(\cdot)$, a multi-layer mixture prior is adopted. According to Feller (1971), any unimodal and symmetric density (with finite first derivative) $f(\cdot)$ can be represented by a mixture of uniform distributions. We focus on the scenario that the mixing distribution has a density $g(\cdot)$. The density $g(\cdot)$ is then built upon a Dirichlet process mixture of gamma distributions. Using latent variables, we can write out the hierarchical model

$$W_i \mid X_i \sim \mathcal{N}(X_i, \sigma_i^2); \quad X_i \mid \theta_i \sim \mathrm{Unif}(-\theta_i, \theta_i); \quad \theta_i \mid z, \mu \sim \mathrm{Ga}(z, z/\mu);$$
$$\mu \mid P_\mu \sim P_\mu; \quad P_\mu \mid m, D \sim \mathrm{DP}(m, D); \quad z \sim \Pi_z(\cdot), \tag{8}$$

where the gamma distributions are reparameterized by shape z and mean μ . The mixing is imposed on μ leading to a Dirichlet process location mixture of gamma distributions. The proposed prior works well in simulations and real data in a recent work under review in a peer-reviewed journal.

The second simulation setup involves a hierarchical prior through the introduced latent variables. We propose to impose the fraction 1/J on the bottom layer of the prior, namely,

$$W_{i} \mid X_{i} \sim \mathcal{N}(X_{i}, \sigma_{i}^{2}); \quad X_{i} \mid \theta_{i} \sim \mathrm{Unif}(-\theta_{i}, \theta_{i}); \quad \theta_{i} \mid z, \mu \sim \mathrm{Ga}(z, z/\mu);$$
$$\mu \mid P_{\mu} \sim P_{\mu}; \quad P_{\mu} \mid m, D \sim \mathrm{DP}(m/J, D); \quad z \sim \Pi_{z}^{1/J}(\cdot), \tag{9}$$

To the best of our knowledge, hierarchical priors have not been investigated in the divide and conquer context. Thus it highlights the capability of applying our method in a broad range. A Gibbs sampler corresponding to (9) is provided in Section S.1.3 in the Supplementary material. The true density is composed of a normal component, $N(0, 0.2^2)$, which targets a sharp peak at zero and a t distribution with degrees of freedom 5 which generates the large values of X. The two components are assigned with probabilities 0.8 and 0.2 respectively so that the resulting density has a sharp peak around zero and a small portion on the large values. The choice of error variances is $\sigma_i^2 = (0.75 + X_i/4)^2$ under which the variance of error depends on X and the *expected* variance of error is more than the variance of X.

We adopt posterior mean estimators unless specified otherwise. For WASP, we originally get an overall posterior distribution from the author's code, from which posterior samples are generated, this in turn leads to the posterior mean estimator. Our estimator is obtained first by taking the average of posterior samples of densities across the MCMC steps on each individual machine. Then our divide and conquer estimator is calculated by averaging over the estimators across the selected machines. In addition, an estimator based on the original analysis using the complete data is implemented for validating the above estimators.

4.2 Simulation Results

For the first simulation, the sample size of the complete data is n = 10,000 and the number of MCMC steps is 10,000 with burn-in steps 5,000 and thinning every 5th iterations. In addition, J = 10 machines are chosen for splitting the data. The simulation is repeated 10 times. All these setups agree with Srivastava et al. (2018) for comparison with the WASP estimator.

Table 1 summarizes and compares the accuracy of parameter estimation for μ_1 and μ_2 . Table 2 contains the counterparts for Σ_1 and Σ_2 . For the covariance matrices estimation, we are limited to present the performance of our method and the method that uses the complete data. The WASP estimators for Σ_1 and Σ_2 are missing due to incapability of obtaining posterior samples from the online codes provided in Srivastava et al. (2018).

The second simulation is performed on a data set of size 15,000 and repeated 50 times. We use 5,000 MCMC iterations with 1,000 burn-ins. The density estimators are constructed in three contexts: the complete data with the original prior, divided data on 20 machines with the original prior, divided data on 20 machines with the proposed prior. For the latter

Parameter	μ_1			μ_2		
Estimator	full	WASP	fPrior	full	WASP	fPrior
bias (×10 ⁻³)	(-2, -2)	(-2, -2)	(-2, -2)	(1, 3)	(0, 3)	(1, 3)
se (×10 ⁻³)	(4.7, 4.5)	(4.3, 4.6)	(4.7, 4.5)	(5.7, 7.5)	(5.8, 7.6)	(5.7, 7.5)

Table 1: The bias and standard error in estimating μ_1 and μ_2 using complete data (full), WASP and our method (fPrior). The reported values are magnified by a factor 10^3 so that less decimals are displayed.

Parameter	Σ_{11}		Σ_{12}		Σ_{22}	
Estimator	full	fPrior	full	fPrior	full	fPrior
bias (×10 ⁻³)	(2, -2)	(2, 0)	(2, -4)	(2, -3)	(6, 0)	(8, 3)
se (×10 ⁻³)	(4.2, 13.3)	(5.1, 7.0)	(3.5, 10.7)	(4.7, 9.5)	(13.3, 7.6)	(10.6, 11.8)

Table 2: The bias and standard error in estimating components of Σ_1 , Σ_2 ($\Sigma_1 = \Sigma_2$) using complete data (full) and our method (fPrior). We report two values for each parameter corresponding to estimators from the two components. The reported values are magnified by a factor 10³ so that less decimals are displayed

two, the estimated density is obtained by further averaging the posterior densities over the 20 machines. Figure 1 presents the estimated densities and the true density.

It is worthwhile to mention that the accuracy of our proposed prior is achieved with a significant computational gain. The algorithm runs on multiple computer nodes in a Linux OS cluster with single core assigned for individual analysis. In addition, less memory is assigned for analysis of split data than that used for analyzing the complete sample. Roughly, the overall time is about 1/5 over that without data splitting.



Figure 1: Deconvoluted density estimators using the complete data of size 15,000 and using the divided data (of equal size) on 20 machines. The truth (black solid line) is a mixture of t with degrees of freedom 5 with probability 0.2 and $N(0, 0.2^2)$ with probability 0.8. The posterior mean of MCMC samples of the density using the complete data and the original prior (green dotted dash line) can be treated as the "best" estimator. The averaged estimator using the original prior over 20 machines (red dashed line) and our estimator (blue dashed line) are displayed for comparison.

4.3 Conclusions

We implemented two distinct simulations in density estimation using finite mixture of normal distributions and density deconvolution with shape constraints. In the first simulation, the performance of our methods in estimating the mean parameter of normal distributions is very competitive with the established WASP while both of them are indifferentiable with the original analysis acting on the complete data. The second simulation showcases the capability of our method in a more complicated problem that involves a hierarchical nonparametric prior. Our estimator has a clear advantage in accuracy over a naive estimator that imposes the prior in the original analysis on split data.

5 A GWAS data set

The algorithm (8) was designed to analyze data from GWAS studies. We focus on one particular study data, GIANT Height. Here we provide a concise introduction that paves the way to apply the method, more thorough information can be found in Allen et al. (2010). The trait variable height is collected for N = 133,653 individuals of recent European ancestry. After an initial screening, the number of single-nucleotide polymorphisms (SNPs) that are of interest is reduced to n = 941,389, of whom the regression coefficients W_i and their associated standard error σ_i , $i = 1, \ldots, n$, in accordance to a simple linear regression are available.

Upon simple derivation the observed effect size W_i is related to the true effect size X_i through a measurement error model $W_i = X_i + U_i$, where $U_i \sim N(0, \sigma_{i\epsilon}^2/N)$ and $\sigma_{i\epsilon}^2$ equals the variance of regression error in the linear regression of height on the *i*th SNP. Because of the large number of individuals in the study, the variance of U_i is well estimated by the standard error of the regression coefficient W_i and thus is treated as known. By assuming all the true effect sizes X_i 's come from one distribution with $f(\cdot)$ as its density function, and acknowledging the fact that the observed effect sizes W_i are symmetric with a majority near zero, it is reasonable to infer that $f(\cdot)$ is unimodal and symmetric at zero. The natural question is how to estimate $f(\cdot)$. It is apparent that applying an efficient algorithm matters when one realizes that the number of SNPs is so large, which is typical for a GWAS data set.

The problem introduced has the same setup as our second simulation. We are interested in applying the divide and conquer algorithm (9) in this paper and compare the density estimators with those obtained by blindly using the original algorithm (8) under the identical splitting of data. The same strategy as used in the simulation section is adopted for combining the individual density estimators. These two estimators are referred to as fPrior and naive correspondingly. We select J = 50 or J = 200 machines with each assigned effect sizes of around 20,000 SNPs in the former case or around 5,000 in the latter. As usual, a posterior mean estimator is chosen as the density estimator on individual machine.

Figure 2 displays the estimated densities for the true effect size X under fPrior or naive. We display the peak and tail areas of these estimators separately, which are determined by a pre-specified cutoff 0.003. Table 3 summarizes the integrated absolute value of the difference (IAD) between the estimated densities under J = 50 and J = 200 for each method. IADs are also calculated separately for the peak region and the tail region.

One conclusion from Figure 2 and Table 3 is that estimating the effect size density using the proposed prior is less sensitive to the total machines being used than that using the naive prior and thus we believe the proposed prior is advantageous, especially in estimating the tails. We also observe that both priors are not very consistent in estimating the density around zero when different number of machines are selected reflecting the intrinsic difficulty in estimating the extremely small effect sizes. In conclusion, our method leads to a feasible solution in practice, especially when the focus is on detecting the larger effect sizes.

6 Discussions

We study a scalable Bayesian parametric or nonparametric density estimation method using a divide and conquer strategy, with a guaranteed optimal posterior convergence. In addition, our numerical and real data results show the applicability of the method to a density deconvolution problem. There is an interest to see how the idea can be used in an even broader context.

Density Area	x > 0.003		x < 0.003	
Estimator IAD $(50 - 200)$	fPrior	naive	fPrior	naive
	0.003	0.015	0.033	0.035

Table 3: The integrated absolute value between the effect size density estimators under the choices of J = 50 and J = 200, IAD (50 – 200), is compared separately for our method (fPrior) and for the naive prior (naive). The metric is calculated for two regions: the larger effect sizes (|x| > 0.003) and the extremely small effect sizes (|x| < 0.003)



Figure 2: True effect size density estimators for the GIANT Height data set containing the observed effect sizes (associated with height) corresponding to 941,389 SNPs separately. The peak region and tail region (on the positive values) are displayed in the left and right panel. Four density estimators are displayed: fPrior using divided data (of equal size) on 50 machines (red dashed line), fPrior using divided data on 200 machines (purple dashed line), the naive prior (original prior) using divided data on 50 machines (blue dotted dashed line), the naive prior using divided data on 200 machines (green dotted dashed line).

Our theoretical results indicate it is expected that the number of machines can not be chosen to be too large compared to the total sample size. In practice, how to select the total number of subsets is an open problem.

We are aware of a very nice theoretical result (Szabó and van Zanten, 2019) that investigates the optimal posterior convergence regardless of the choice for the number of subsets. They consider the signal-in-noise model with a conjugate normal prior which is regulated by the "decay" rate of the true signal. The authors cast a doubt about the existence of an adaptive version of the method. One conjecture based on our theoretical results and others is that a nonparametric prior is crucial in possibly obtaining an adaptive convergence rate, together with a control over the growth rate of the number of subsets.

Supplementary Material

The online supplementary material includes detailed algorithms for the selected examples that we used in the main paper, the additional lemmas that slightly modifies the existing ones in the literature for completeness and clearness.

Acknowledgments

Su is supported by a startup fund from College of Arts and Sciences, University of Kentucky. The Authors are grateful to Yan Zhang of Johns Hopkins University for directions to the online GWAS data set.

Appendix

Appendix

A.1 Proof of Theorem

Proof. This is a proof of Theorem 1. We will show that for the subset posterior distributions $\Pi_m(\cdot \mid \mathbf{X}_j)$ the same contraction rate is achieved as if the original prior has been assigned on each machine.

It is easy to argue that all $\Pi_m(\cdot | \mathbf{X}_j)$ share the same asymptotic results, it is sufficient to do for a single one. With a slight abuse of notation, $\Pi_m(\cdot | \mathbf{X}_m)$ denotes the posterior distribution on $f(\cdot)$ in accordance to the adjusted prior, where $\mathbf{X}_m = (X_1, \ldots, X_m)$ corresponds to a sample of size m. Thus,

$$\Pi_m(\cdot \mid \boldsymbol{X}_m) = \frac{\prod_{i=1}^m f(X_i) d\Pi_J(f)}{\int \prod_{i=1}^m f(X_i) d\Pi_J(f)}.$$

Here $\Pi_J(f)$ is the prior given by (4).

We can follow the procedure in Shen et al. (2013) which extends Ghosal et al. (2007) to derive the contraction rate for $\Pi_m(\cdot | \mathbf{X}_j)$. The former leads to an adaptive rate assuming the true function is in a locally Hölder class. The difference lies in the fraction prior that is used in this paper. The proof is built upon that of Theorem 1 in Shen et al. (2013). Here we aim to organize the outline of the proof and discuss the differences whenever necessary.

Given the assumptions, a sieve space corresponding to ϵ_m is constructed as \mathscr{F}_m . The procedure requires three major steps:

- 1. $\log \mathcal{N}(\epsilon_m, \mathscr{F}_m^c, h) \leq c_1 m \epsilon_m^2$ when $m \to \infty$,
- 2. $\Pi_J(\mathscr{F}_m^c) \le c_3 \exp[-(c_2+4)m\widetilde{\epsilon}_m^2],$
- 3. $\Pi_J(f: P_0 \log(f/f_0) \le \tilde{\epsilon}_m^2, P_0 \log^2(f/f_0) \le \tilde{\epsilon}_m^2) \ge c_4 \exp(-c_2 m \tilde{\epsilon}_m^2).$

We can show that all three steps above hold for $\mathscr{F}_m = \{f : f = \int \phi_\sigma(x-z)P(dz) \text{ with } P = \sum_{h=1}^{\infty} \pi_h \delta_{z_h}, z_h \in [-a_m, a_m] \text{ if } h \leq H_m; \sum_{h>H_m} \pi_h < \epsilon_m; \sigma_m^2 \leq \sigma^2 < \sigma_m^2 (1 + \epsilon_m^2)^{M_m} \},$ where $a_m^{a_1} = \sigma_m^{-2a_2} = M_m = m, H_m = \lfloor m \epsilon_m^2 / (\log m) \rfloor, \epsilon_m = m^{-\beta/(2\beta+1)} (\log m)^t$ and $\tilde{\epsilon}_m = m^{-\beta/(2\beta+1)} (\log m)^{t_0}, t-1/2 > t_0 = 1 + (1/\tau)/(2+1/\beta)$. The detailed proof follows the proof of Theorem 1 in Shen et al. (2013) where we see that the effect of the proposed prior leads to a contraction rate ϵ_m that is only a factor $(\log m)^{1/2}$ larger than if using the original prior.

Since the same sieve space as Shen et al. (2013) is adopted and a larger ϵ_m (a factor of $\log m$), the first step on the entropy of the sieve space remains the same.

The second step will be discussed with more details. From the definition of sieve space \mathscr{F}_m , it can be easily argued that under prior (4), $DP(M/J, G) \times \Pi_{\sigma}^{1/J}$,

$$(\mathrm{DP}(M/J,G) \times \Pi_{\sigma}^{1/J})(\mathscr{F}_{m}^{c}) \leq H_{m}G([-a_{m},a_{m}]^{c}) + Pr(\sum_{h>H_{m}}\pi_{h} > \epsilon_{m}) + \Pi_{\sigma}^{1/J}(\sigma^{-2} > \sigma_{m}^{-2}) + \Pi_{\sigma}^{1/J}(\sigma^{-2} \leq \sigma_{m}^{-2}(1+\epsilon_{m}^{2})^{-M_{m}})(\mathrm{A.1})$$

According to the assumptions on G, Π_{σ} and \mathscr{F}_m , it can be shown that

$$H_m G([-a_m, a_m]^c) \leq b_1 m \epsilon_m^2 (\log m)^{-1} \exp(-C_1 m),$$

$$\Pi_{\sigma}^{1/J} (\sigma^{-2} > \sigma_m^{-2}) \leq b_2^{1/J} \exp(-C_2 m/J),$$

$$\Pi_{\sigma}^{1/J} (\sigma^{-2} \leq \sigma_m^{-2} (1 + \epsilon_m^2)^{-M_m}) \leq b_3^{1/J} m^{a_3/(a_2 J)} (1 + \epsilon_m^2)^{-m/J}$$

$$\approx \exp(-C_3 m \epsilon_m^2/J),$$

in addition, the second term in (A.1), the upper bound for $Pr(\sum_{h>H_m} \pi_h > \epsilon_m)$, can be found using stick breaking representation for $\{\pi_h, h \ge 1\}$, that is, $\pi_h = V_h \prod_{i < h} (1 - V_i)$, $\{V_h, h \ge 1\}$ are independent beta-distributed random variables with parameter 1 and M/J. Then we can show

$$Pr(\sum_{h>H_m} \pi_h > \epsilon_m) = Pr\{\prod_{i=1}^{H_m} (1-V_i) > \epsilon_m\} = Pr\{-\sum_{i=1}^{H_m} \log(1-V_i) < \log(1/\epsilon_m)\} \\ \leq \frac{\{-(M/J)\log\epsilon_m\}^{H_m}}{\Gamma(H_m+1)} \leq \left(\frac{eM}{H_mJ}\log\frac{1}{\epsilon_m}\right)^{H_m} \asymp \exp\{-C_4m\epsilon_m^2\log m\}.$$

The last two inequalities follow from $-\sum_{i=1}^{H_m} \log(1-V_i)$ is a Gamma random variable with parameter H_m and M/J and Stirling's formula.

These upper bounds together with (A.1) yield $(DP(M/J, G) \times \Pi_{\sigma}^{1/J})(\mathscr{F}_{m}^{c}) \lesssim \exp(-Cm\epsilon_{m}^{2}/J)$ for some constant *C*. Since $m = n/J, J \asymp \log n$ and $\epsilon_{m}^{2} > \tilde{\epsilon}_{m}^{2} \log m$, we conclude that $(DP(M/J, G) \times \Pi_{\sigma}^{1/J})(\mathscr{F}_{m}^{c}) \leq c_{3} \exp\{-(c_{2}+4)m\tilde{\epsilon}_{m}^{2}\}$ for some constant c_{3} and any constant c_{2} , which will be chosen as the constant in step three.

The third step is termed "Prior thickness result" in Shen et al. (2013) and the lower bound therein is built for prior (3), $DP(M,G) \times \Pi_{\sigma}$, on a prior set $\mathcal{P}_{\sigma_m} \times \mathcal{S}_{\sigma_m}$ (in the one dimensional case $\mathcal{S}_{\sigma_m} = \{\sigma : \sigma^{-2} \in [\sigma_m^{-2}, \sigma_m^{-2}(1 + \sigma_m^{2\beta})]\}$) with $\sigma_m^\beta = \tilde{\epsilon}_m \{\log^{-1}(1/\tilde{\epsilon}_m)\}$, in recognition of Lemma 10 in Ghosal et al. (2007) and the condition of Π_{σ} . Thus it can be easily verified the lower bound holds for the same $\tilde{\epsilon}_m$ under prior (4), $DP(M/J,G) \times \Pi_{\sigma}^{1/J}$ on the prior set $\mathcal{P}_{\sigma_m} \times \mathcal{S}_{\sigma_m}$, as long as the same lower bound of Lemma 10 and that of Π_{σ} can be achieved. It is easy to show that $\Pi_{\sigma}^{1/J}(\mathcal{S}_{\sigma_m}) \geq C_4 \exp[-c_4 \tilde{\epsilon}^{-1/\beta} \{\log(1/\tilde{\epsilon}_m)\}^{2+1/\tau+1/\beta}]$ under Condition (C2) and (C4). We can also show that the same lower bound of Lemma 10 in Ghosal et al. (2007) holds with a slight modification on the condition of Dirichlet parameters. For readers' interest it is stated as Lemma S.1 and provided in the Supplementary material.

References

- Allen, H. L., Estrada, K., Lettre, G., Berndt, S. I., Weedon, M. N., Rivadeneira, F., Willer, C. J., Jackson, A. U., Vedantam, S., Raychaudhuri, S., et al. (2010). Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature*, 467, 832– 838.
- Balakrishnan, N. and Nevzorov, V. B. (2004). A primer on statistical distributions. John Wiley & Sons.
- Bhattacharya, A., Pati, D., Pillai, N. S., and Dunson, D. B. (2015). Dirichlet–Laplace priors for optimal shrinkage. *Journal of the American Statistical Association*, 110, 1479–1490.
- Blei, D. M., Jordan, M. I., et al. (2006). Variational inference for Dirichlet process mixtures. Bayesian analysis, 1, 121–143.
- De Jonge, R., Van Zanten, J., et al. (2010). Adaptive nonparametric Bayesian inference using location-scale mixture priors. *The Annals of Statistics*, 38, 3300–3320.
- Dunson, D. B. and Xing, C. (2009). Nonparametric Bayes modeling of multivariate categorical data. Journal of the American Statistical Association, 104, 1042–1051.
- Feller, W. (1971). An Introduction to Probability Theory and its Applications, volume 2. John Wiley & Sons.
- Ghosal, S., Van Der Vaart, A., et al. (2007). Posterior convergence rates of Dirichlet mixtures at smooth densities. *The Annals of Statistics*, 35, 697–723.
- Guhaniyogi, R., Li, C., Savitsky, T. D., and Srivastava, S. (2017). A divide-and-conquer Bayesian approach to large-scale kriging. arXiv preprint arXiv:1712.09767, .
- Ishwaran, H. and Zarepour, M. (2002). Exact and approximate sum representations for the Dirichlet process. *Canadian Journal of Statistics*, 30, 269–283.
- Naulet, Z., Barat, E., et al. (2018). Some aspects of symmetric Gamma process mixtures. Bayesian Analysis, 13, 703–720.
- Neiswanger, W., Wang, C., and Xing, E. P. (2014). Asymptotically exact, embarrassingly parallel MCMC. In *Proceedings of the Thirtieth Conference on Uncertainty in Artificial Intelligence*, page 623632, Arlington, Virginia, USA. AUAI Press.

- Rasmussen, C. E. (2000). The infinite Gaussian mixture model. In Advances in neural information processing systems, pages 554–560.
- Sabnis, G., Pati, D., Engelhardt, B., and Pillai, N. (2016). A divide and conquer strategy for high dimensional Bayesian Factor models. *arXiv preprint arXiv:1612.02875*, .
- Scott, S. L., Blocker, A. W., Bonassi, F. V., Chipman, H. A., George, E. I., and McCulloch, R. E. (2016). Bayes and big data: The consensus Monte Carlo algorithm. *International Journal of Management Science and Engineering Management*, 11, 78–88.
- Sethuraman, J. (1994). A constructive definition of Dirichlet priors. *Statistica Sinica*, 4, 639–650.
- Shen, W., Tokdar, S. T., and Ghosal, S. (2013). Adaptive Bayesian multivariate density estimation with Dirichlet mixtures. *Biometrika*, 100, 623–640.
- Srivastava, S., Li, C., and Dunson, D. B. (2018). Scalable Bayes via barycenter in Wasserstein space. The Journal of Machine Learning Research, 19, 312–346.
- Szabó, B. and van Zanten, H. (2019). An asymptotic analysis of distributed nonparametric methods. *Journal of Machine Learning Research*, 20, 1–30.
- Xue, J. and Liang, F. (2019). Double-Parallel Monte Carlo for Bayesian analysis of big data. Statistics and computing, 29, 23–32.

Supplementary Material to Divide and Conquer algorithm of Bayesian Density Estimation

Ya Su Department of Statistics, University of Kentucky, Lexington, KY 40536-0082, U.S.A., ya.su@uky.edu

S.1 Algorithms for example models

The algorithms for three example models in the main body of the paper are displayed. For ease of notation, let $\Omega_{-\zeta}$ be all variables in Ω but excluding ζ .

S.1.1 Finite mixtures of Normal prior

The Gibbs sampling algorithm is given below.

Denote X_{ji} the *i*th sample distributed to subset j, Z_{ji} the component indicator variable where X_{ji} pertains to that is, $X_{ji}|Z_{ji} = k \sim N_p(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k), n_{jk} = \sum_{i=1}^m I_{(Z_{ji}=k)}, \overline{X}_{jk} = n_{jk}^{-1} \sum_{i=1}^m X_{ji} I_{(Z_{ji}=k)}$ and $V_{jk} = \sum_{Z_{ji}=k} (X_{ji} - \overline{X}_{jk}) (X_{ji} - \overline{X}_{jk})^T$ corresponding to the number of samples, sample mean and scaled sample covariance matrix belonging to subset j and component k for $j = 1, \ldots, J, k = 1, \ldots, K$. Under the new prior for each subset, the conditional posterior distributions of the variables are

$$P(Z_{ji} = k | \boldsymbol{\Omega}_{-Z_i}) \propto \pi_k N_p(X_{ji}; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

$$\boldsymbol{\Sigma}_k | \boldsymbol{\Omega}_{-\{\boldsymbol{\Sigma}_k, \boldsymbol{\mu}_k\}} \sim IW \left(n_{jk} + \frac{\nu + 1}{J} - \frac{(p+1)(J-1)}{J}, V_{jk} + \frac{(lJ)^{-1}n_{jk}}{(lJ)^{-1} + n_{jk}} \overline{X}_{jk} \overline{X}_{jk}^T + \frac{1}{J} \boldsymbol{S} \right)$$

$$\boldsymbol{\mu}_k | \boldsymbol{\Omega}_{-\boldsymbol{\mu}_k} \sim N_p \left(\frac{n_{jk}}{(lJ)^{-1} + n_{jk}} \overline{X}_{jk}, \frac{1}{(lJ)^{-1} + n_{jk}} \boldsymbol{\Sigma}_k \right)$$

$$\boldsymbol{\pi} | \boldsymbol{\Omega}_{-\boldsymbol{\pi}} \sim Dir(n_{j1} + \alpha_1 J^{-1}, \dots, n_{jK} + \alpha_K J^{-1}).$$

S.1.2 Shape constraint density deconvolution

To ease computation, we approximate the Dirichlet process mixture prior with a finite mixture of Gamma distributions with K components, with a specific Dirichlet prior on the mixture probabilities (Ishwaran and Zarepour, 2002). Specifically, our hierarchical Bayes model for subsequent implementations is as follows. Let i denote the index for subject, and k be the index for the kth component, for all i = 1, ..., n, k = 1, ..., K. Let t > 1 denote a fixed constant. Then,

$$(W_i|X_i) \sim \operatorname{Normal}(X_i, \sigma_i^2); \quad (X_i|\theta_i) \sim \operatorname{Unif}(-\theta_i, \theta_i); \quad (\theta_i|Z_i = k, \alpha_k, \beta_k) \sim \operatorname{Ga}(\alpha_k, \beta_k); P(Z_i = k|p_1, \dots, p_K) = p_k; \quad (\alpha_k|\lambda, t) \sim \operatorname{Expon}(\lambda; t, \infty); \quad (\beta_k|\Xi_1, \Xi_2) \sim \operatorname{Ga}(\Xi_1, \Xi_2); (p_1 \dots, p_K) \sim \operatorname{Dir}(m/K, \dots, m/K),$$

where $\operatorname{Expon}(\lambda; \ell, u)$ denotes an exponential distribution with parameter λ truncated at (ℓ, u) . The truncation of α_k at some t > 1 makes the density of X be finite at zero. The set of hyperparameters is $(\lambda, t, \Xi_1, \Xi_2, K, m)$.

For k = 1, ..., K, let $r_k = \sum_i I_{(Z_i=k)}$ be the total number of individuals that fall into group k and $s_k = \sum_i \theta_i I_{(Z_i=k)}$ be the summation of the θ_i from the kth group. To sample from the posterior distribution of Ω , we use a Gibbs sampler for all parameters other than the α_k , combined with a Metropolis-Hastings within Gibbs for the α_k . The posterior full-conditional distributions are

$$\begin{aligned} & (X_i | \mathbf{\Omega}_{-X_i}) \sim \mathrm{N}(W_i, \sigma_i^2; -\theta_i, \theta_i); \\ & (\theta_i | \mathbf{\Omega}_{-\theta_i}) \sim \mathrm{Ga}(\alpha_{Z_i} - 1, \beta_{Z_i}; |X_i|, \infty); \\ & P(Z_i = k | \mathbf{\Omega}_{-Z_i}) \propto \Gamma(\alpha_k)^{-1} p_k (\beta_k \theta_i)^{\alpha_k} \exp(-\beta_k \theta_i); \\ & (p_1, \dots, p_K | \mathbf{\Omega}_{-\{p_1, \dots, p_K\}}) \sim \mathrm{Dir}(m/K + r_1, \dots, m/K + r_K); \\ & (\beta_k | \mathbf{\Omega}_{-\beta_k}) \sim \mathrm{Ga}(\Xi_1 + \alpha_k r_k, \Xi_2 + s_k); \\ & (\alpha_k | \mathbf{\Omega}_{-\alpha_k}) \propto \Gamma(\alpha_k)^{-r_k} \exp\{-\alpha_k (\lambda - r_k \log \beta_k - \sum_i \log(\theta_i) I_{(Z_i = k)})\}. \end{aligned}$$

The symbol $N(\mu, \sigma^2; \ell, u)$ denotes a Normal distribution with parameters (μ, σ^2) truncated at (ℓ, u) , while $Ga(\alpha, \beta; \ell, u)$ corresponds to a Gamma distribution with parameters (α, β) truncated at (ℓ, u) . Since the posterior distribution of α_k does not belong to a standard family, we implement a Metropolis-Hastings algorithm within the Gibbs sampler to update the α_k . We use a Gamma proposal distribution; specifically, $\tilde{\alpha}_k \sim Ga(2, 2/\alpha_k; t, \infty)$, and we accept the proposed $\tilde{\alpha}_k$ or keep the original α_k according to the general Metropolis-Hastings rule. The proposal distribution is truncated to reflect the prior assumption on α_k .

S.2 Additional Lemmas

The following lemma is a straightforward extension to Lemma 10 in Ghosal et al. (2007). It turns out that the same conclusion holds for a Dirichlet-distributed random variable when the sum of its associated parameters have limit zero.

Lemma S.1. For (p_1, \ldots, p_N) be an arbitrary point in the *N*-dimensional unit simplex and let (X_1, \ldots, X_N) be Dirichlet distributed with parameter $(\alpha_1, \ldots, \alpha_N)$ with $\alpha_j \leq 1$ and $\sum_{j=1}^{N} \alpha_j = m_N$. Suppose $\lim_{N\to\infty} m_N = 0$. Then for every $\epsilon^b < a\alpha_j$ and $\epsilon_N \leq 1$, there exists constants *c* and *C* that depend only on *a* and *b* such that

$$\Pr\left(\sum_{j=1}^{N} |X_j - p_j| \le 2\epsilon, \min_{1 \le j \le N} X_j \ge \epsilon^2/2\right) \ge C \exp(-cN \log \epsilon^{-1}).$$

The proof of the above lemma can follow exactly the lines of Lemma 10 in Ghosal et al. (2007) and thus is omitted.