arXiv:2002.07241v2 [math.CO] 6 May 2020

Codes with few weights arising from linear sets

Vito Napolitano and Ferdinando Zullo

Abstract

In this article we present a class of codes with few weights arising from special type of linear sets. We explicitly show the weights of such codes, their weight enumerator and possible choices for their generator matrices. In particular, our construction yields also to linear codes with three weights and, in some cases, to almost MDS codes. The interest for these codes relies on their applications to authentication codes and secret schemes, and their connections with further objects such as association schemes and graphs.

MSC 2010: 51E20, 05B25, 51E22
Keywords: Codes with few weights, Linear set, MRD-code.

This research was partially supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM). The authors were also supported by the project "VALERE: VAnviteLli pEr la RicErca" of the University of Campania "Luigi Vanvitelli".

1 Introduction

Let $q = p^h$, with p prime and $h \geq 1$. A *q-ary linear code* $\mathcal{C}$ is any $\mathbb{F}_q$-subspace of $\mathbb{F}_q^m$. If $\mathcal{C}$ has dimension r, we say that $\mathcal{C}$ is an *$[m, r]_q$-code*. A *generator matrix* G for a linear code $\mathcal{C}$ is a $(r \times m)$-matrix over $\mathbb{F}_q$ whose rows form a basis of $\mathcal{C}$, i.e.

$$\mathcal{C} = \{\mathbf{x}G \colon \mathbf{x} \in \mathbb{F}_q^r\}.$$

We may consider $\mathbb{F}_q^m$ as a metric space endowed with the *Hamming distance* d_H, i.e. $d_H(\mathbf{x}, \mathbf{y})$ is the number of entries in which $\mathbf{x}$ and $\mathbf{y}$ differ, with $\mathbf{x}, \mathbf{y} \in \mathbb{F}_q^m$. The *Hamming weight* $w_H(\mathbf{c})$ of a codeword $\mathbf{c} \in \mathcal{C}$ is the number of nonzero entries of $\mathbf{c}$, or equivalently $w_H(\mathbf{c}) = d_H(\mathbf{c}, \mathbf{0})$. Recall that the

minimum weight of a linear code coincides with its minimum distance. Let denote by A_i^H the number of codewords of $\mathcal{C}$ with Hamming weight i. The *(Hamming) weight enumerator* is defined as the following polynomial

$$1 + A_1^H z + \ldots + A_m^H z^m.$$

This polynomial gives a good deal of information about the code and it is an important invariant, and has been calculated for few families of codes. Also, it is used in the probability theory involved with different ways of decoding. A *ℓ-weight code* $\mathcal{C}$ is an $[m, r]_q$-code having ℓ nonzero weights $w_1 < \ldots < w_\ell$, i.e. if the sequence $(A_1^H, \ldots, A_m^H)$ have exactly ℓ nonzero entries. If $\ell \leq r$ we say that the code has *few weights*. Much of the focus on linear codes to date has been on codes with few weights, especially on two and three-weight codes, for their applications in secret sharing [13], authentication codes [18] and their connections with association schemes [6, 5] and with graphs [33].

Let n, r, h be positive integers such that $h \leq r - 1$ and $h + 1 \mid rn$, and let q be a prime power. Denote by $\theta_i = \frac{q^{i+1}-1}{q-1}$ for any $i \geq 0$. In this paper we deal with a geometric construction of linear $[\theta_{\frac{rn}{h+1}}, r]_{q^n}$-codes with at most $h + 1$ weights, determining their weight enumerator and the possible distribution of their weights, see Section 4. In particular, for $h = r - 1$ this construction yields r-weight codes connected with a special class of maximum rank distance codes, see Section 5. When $h = 2$ we prove that our construction yields three-weight codes, cf. Section 6, and examples of almost MDS codes in even characteristic, c.f. Remark 6.6. Finally, in Section 7, we present suitable choices for generator matrices when $h = 2$ and $h = r - 1$.

2 Some known linear codes with few weights

In the literature the known constructions for linear codes with few weights arise either from linear algebraic tools, based on the properties of the trace function, or from a geometric point of view.

In [16, 17], the authors define a class of linear codes as follows. Let $\mathrm{Tr}_{q/p}$ denote the trace function from $\mathbb{F}_q$ to $\mathbb{F}_p$, i.e. $\mathrm{Tr}_{q/p}(x) = \sum_{i=0}^{h-1} x^{p^i}$. Let $D = \{d_1, \ldots, d_m\} \subseteq \mathbb{F}_q$ and let

$$\mathcal{C}_D = \{(\mathrm{Tr}_{q/p}(xd_1), \ldots, \mathrm{Tr}_{q/p}(xd_m)) \colon x \in \mathbb{F}_q\}.$$

Then $\mathcal{C}_D$ is a linear code of length m over $\mathbb{F}_p$ and D is called its *defining set*. Most of the known constructions for codes with few weights come from

$\mathcal{C}_D$ by selecting appropriately the defining set D. For instance, in [13] the authors choose

$$D = \{x \in \mathbb{F}_q^* \colon \mathrm{Tr}_{q/p}(x^2) = 0\}.$$

Then

$$|D| = \begin{cases} p^{h-1} - 1 & h > 1 \text{ odd} \\ p^{h-1} - (-1)^{\left(\frac{p-1}{2}\right)^2 \frac{m}{2}}(p-1)p^{\frac{m-2}{2}} - 1 & h \geq 2 \text{ even} \end{cases}$$

and $\mathcal{C}_D$ has length $|D|$ and either two or three different weights according to the parity of h.

For the geometric type construction we need the following notion. A *projective* $[n, r, d]$*–system* is a finite (multi-)set $\mathcal{M}$ of points of $\mathrm{PG}(r-1, q)$, not all of which lie in a hyperplane, where $n = |\mathcal{M}|$, and

$$n - d = \max\{|\mathcal{M} \cap H| \ : H \subset \mathrm{PG}(r-1, q), \dim H = r-2\},$$

see [2]. Note that the cardinalities above are counted with multiplicities in the case of a multiset. By defining the $(r \times n)$-matrix G by taking as columns the coordinates of points of $\mathcal{M}$, then G is the generator matrix of a linear $[n, r]_q$-code $\mathcal{C}_{\mathcal{M}}$.

Conversely, let G be an $(r \times n)$-generator matrix of a nondegenerate $[n, r]_q$-linear code $\mathcal{C}$, that is a code without coordinates being zero for every codeword. The (multi-)set of one dimensional subspaces of $\mathbb{F}_q^r$ spanned by the columns of G, may be considered as a (multi-)set $\mathcal{M}$ of points of $\mathrm{PG}(r-1, q)$. So, there is a one to one correspondence between nondegenerate linear codes and projective systems. Furthermore, for any non-zero vector $\mathbf{v} = (v_1, v_2, \ldots, v_r)$ in $\mathbb{F}_q^r$, we have that the projective hyperplane

$$v_1x_1 + v_2x_2 + \cdots + v_rx_r = 0$$

contains $|\mathcal{M}| - w$ points of $\mathcal{M}$ if and only if the codeword $\mathbf{v}G$ has weight w. Thus, there exists a linear $[n, r]_q$–code with minimum distance d if and only if there exists a projective $[n, r, d]$-system. So, the number of distinct weights of $\mathcal{C}$ corresponds to the distinct sizes of the intersections of $\mathcal{M}$ with all the hyperplanes, that is with the intersection numbers of $\mathcal{M}$ with respect to the hyperplanes. See also [34].

In particular, we are interested in three-weight codes, i.e. those arising from (multi-)set with three intersection numbers with respect to hyperplanes.

Despite of the number of the algebraic constructions for three-weight codes, see e.g. [13, 14, 15, 33, 36], there are few known geometric constructions [19]. A general geometric construction for three-weight codes has been given in [1]. Let $a \in \mathbb{F}_{q^2}^*$, $b \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ and $\mathcal{B}(a,b)$ be the affine algebraic set of $\mathrm{AG}(r-1,q^2)$ of equation

$$x_{r-1}^q - x_{r-1} + a^q(x_1^{2q} + \ldots + x_{r-2}^{2q}) - a(x_1^2 + \ldots + x_{r-2}^2) = (b^q - b)(x_1^{q+1} + \ldots + x_{r-2}^{q+1}).$$

Let P_∞ be the point at infinity of the non-degenerate Hermitian variety of $\mathrm{AG}(r-1,q^2)$ with affine equation

$$x_{r-1}^q - x_{r-1} = (b^q - b)(x_1^{q+1} + \ldots + x_{r-2}^{q+1}).$$

Let r be a odd positive integer and let j be a positive integer greater than one. Providing some conditions on a, b and q (cf. [1, Section 5]), the multiset $\overline{\mathcal{B}}(a,b)$, consisting of the points of $\mathcal{B}(a,b)$ and j times the point P_∞, defines a linear $[q^{2r-3}+j, r]_{q^2}$-code $\mathcal{C}_{\overline{\mathcal{B}}(a,b)}$ with weights

- q^{2r-3};
- $q^{2r-3} - q^{2r-5}$;
- $q^{2r-3} - q^{2r-5} + q^{r-3} + j$;
- $q^{2r-3} - q^{2r-5} + q^{r-3} - q^{r-2} + j$.

When $j = q^{r-2} - q^{r-3}$ or $j = q^{2r-5} - q^{r-3}$, $\mathcal{C}_{\overline{\mathcal{B}}(a,b)}$ is a three-weight code. Note that when $\mathcal{C}_{\overline{\mathcal{B}}(a,b)}$ is a three-weight code, it is also q-*divisible*, i.e. all the nonzero weights are divisible by q.

3 Scattered linear sets

Let V be an r-dimensional $\mathbb{F}_{q^n}$-vector space. A point set L of $\Lambda = \mathrm{PG}(V, \mathbb{F}_{q^n}) = \mathrm{PG}(r-1, q^n)$ is said to be an $\mathbb{F}_q$-*linear set* of Λ of rank k if it is defined by the non-zero vectors of a k-dimensional $\mathbb{F}_q$-vector subspace U of V, i.e.

$$L = L_U := \{\langle \mathbf{u} \rangle_{\mathbb{F}_{q^n}} : \mathbf{u} \in U \setminus \{\mathbf{0}\}\}.$$

We will also denote its rank by $\mathrm{rk}(L_U)$.

Let $\Omega = \mathrm{PG}(W, \mathbb{F}_{q^n})$ be a subspace of Λ and let L_U be an $\mathbb{F}_q$-linear set of Λ, then $\Omega \cap L_U = L_{W \cap U}$, and if $\dim_{\mathbb{F}_q}(W \cap U) = i$, i.e. if the $\mathbb{F}_q$-linear set $\Omega \cap L_U = L_{W \cap U}$ has rank i, we say that Ω has *weight* i in L_U, and we write $w_{L_U}(\Omega) = i$. Note that if Ω has dimension s and L_U has rank k, then $0 \leq w_{L_U}(\Omega) \leq \min\{k, n(s+1)\}$. In particular, a point P belongs to an $\mathbb{F}_q$-linear set L_U if and only if $w_{L_U}(P) \geq 1$.

For an $\mathbb{F}_q$-linear set L_U of rank k in $\Lambda = \mathrm{PG}(r-1, q^n) = \mathrm{PG}(V, \mathbb{F}_{q^n})$ the bound

$$|L_U| \leq \theta_{k-1} = \frac{q^k - 1}{q - 1}, \tag{1}$$

holds true. Hence, L_U is called *scattered* if it achieves the bound (1), or equivalently if all of its points have weight one.

A scattered $\mathbb{F}_q$-linear set L_U of Λ with highest possible rank is a *maximum scattered $\mathbb{F}_q$–linear set* of Λ.

More recently, extending the definition given in [24] and in [32], the family of h-scattered linear sets has been introduced in [9]. An $\mathbb{F}_q$-linear set L_U of Λ is said *h-scattered linear set* if

- $\langle L_U \rangle = \Lambda$;
- for each $(h-1)$-subspace Ω of $\mathrm{PG}(r-1, q^n)$ we have that

$$w_{L_U}(\Omega) \leq h.$$

When $h = r - 1$ and $\dim_{\mathbb{F}_q} U = n$, we obtain the scattered linear sets with respect to hyperplanes introduced in [24] and in [32].

In this paper we will deal with h-scattered linear sets of maximum rank, showing that they provide families of point set with few intersection numbers with respect to hyperplanes, yielding codes (equipped with Hamming distance) with few weights. To this aim, let us recall some properties of h-scattered linear sets proved in [9]. In [9, Theorem 2.3], it has been proved a bound on the rank of a h-scattered $\mathbb{F}_q$-linear set. More precisely, if L_U is an h-scattered $\mathbb{F}_q$-linear set in $\Lambda = \mathrm{PG}(r-1, q^n)$ which is not a subgeometry, then

$$\mathrm{rk}(L_U) \leq \frac{rn}{h+1}, \tag{2}$$

which generalizes the bound given by Blokhuis and Lavrauw in [4] for the classical scattered linear sets. Constructions of h-scattered $\mathbb{F}_q$-linear sets

attaining the bound (2) have been presented in [9, Theorems 2.7 and 3.6] for $h > 1$ and in [28, Section 4], [24, Section 3] and in [32, Corollary 4.4] for $h = r - 1$. Therefore, h-scattered $\mathbb{F}_q$-linear sets attaining (2) are called *maximum h-scattered* $\mathbb{F}_q$-linear sets. From now on, we will consider only maximum h-scattered $\mathbb{F}_q$-linear sets. In [9, Theorem 2.8 and Section 5], the authors also show that such linear sets have few possible intersection numbers with respect to hyperplanes. Indeed, if L_U is a maximum h-scattered $\mathbb{F}_q$-linear set in $\Lambda = \mathrm{PG}(r-1, q^n)$ then for each hyperplane $\mathcal{H} = \mathrm{PG}(r-2, q^n)$ of Λ we have that

$$\frac{rn}{(h+1)} - n \leq w_{L_U}(\mathcal{H}) \leq \frac{rn}{(h+1)} - n + h.$$

Since L_U is scattered, if $\mathcal{H}$ is a hyperplane of Λ with $w_{L_U}(\mathcal{H}) = i$, then

$$|L_U \cap \mathcal{H}| = \theta_{i-1} = \frac{q^i - 1}{q - 1},$$

hence for any hyperplane $\mathcal{H}$ we have that the possible $h + 1$ intersection numbers with respect to hyperplanes are

$$\theta_{\frac{rn}{(h+1)}-n-1}, \ldots, \theta_{\frac{rn}{(h+1)}-n+h-1}. \tag{3}$$

Remark that $\theta_{-1} = 0$. For $h = 1$, in [4], the authors proved that there exist hyperplanes intersecting the linear set in $\theta_{\frac{rn}{2}-n-1}$ and $\theta_{\frac{rn}{2}-n}$ points, i.e. maximum scattered linear sets are two character sets. When $h > 1$ it is not known whether for all the intersection numbers in (3) there exist at least one hyperplane intersecting the linear set in such a number of points. We will solve this problem for the $h = r - 1$ and $h = 2$ cases, see Sections 5 and 6. Since the intersection numbers with respect to hyperplanes for a maximum h-scattered linear set are at most $h+1$, they provide examples of codes with few weights, as we will see in the next sections.

4 Constructions

In this section we exhibit codes (equipped with Hamming metric) with few weights arising from maximum h-scattered linear sets following the geometric construction presented in Section 2, and the papers [5, 6] and [22, Section 2.6].

Now, let L_U be a maximum h-scattered linear set in $\Lambda = \mathrm{PG}(r-1, q^n)$. Denote by N its number of points, i.e. $N = \frac{q^{\frac{rn}{h+1}}-1}{q-1}$, and let

$$L_U = \{\langle (g_{1i}, \ldots, g_{ri}) \rangle_{\mathbb{F}_{q^n}} : i \in \{1, \ldots, N\}\}.$$

Consider G as the $(r \times N)$-matrix with the points of L_U as columns. Since $\langle L_U \rangle = \Lambda$, the rank of G is r, so G is the generator matrix of an $\mathbb{F}_{q^n}$-linear code $\mathcal{C}_{L_U}$ of dimension r and length N.

Consider $\mathbf{c} = (c_1, \ldots, c_N)$ a codeword of $\mathcal{C}_{L_U}$, then there exists $(a_0, \ldots, a_{r-1}) \in \mathbb{F}_{q^n}^r$ such that

$$c_j = \sum_{i=1}^{r} a_{i-1} g_{ij}$$

for each $j \in \{1, \ldots, N\}$. Suppose that $c_j = 0$, then $a_0 g_{1j} + \cdots + a_{r-1} g_{rj} = 0$. This happens if and only if the point $\langle (g_{1j}, \ldots, g_{rj}) \rangle_{\mathbb{F}_{q^n}}$ belongs to the hyperplane with equation $a_0 x_0 + \cdots + a_{r-1} x_{r-1} = 0$. Therefore, denoting by $\mathcal{H}\colon a_0 x_0 + \cdots + a_{r-1} x_{r-1} = 0$, since L_U is scattered then the Hamming weight of c is $N - \theta_{i-1}$ if $w_{L_U}(\mathcal{H}) = i$. Hence, by (3), we have the following.

Theorem 4.1. *If L_U is a maximum h-scattered linear set in Λ, then $\mathcal{C}_{L_U}$ is a code with at most $h+1$ weights. In particular, the weights of $\mathcal{C}_{L_U}$ are*

$$N - \theta_{i-1},$$

where i runs over all the possible weights of a hyperplane with respect to L_U.

In the next sections, we will determine the weights and the weight enumerators for $h = 2$ and $h = r - 1$.

5 Weights for $h = r - 1$

As already remarked, to find the weights of the aforementioned codes is equivalent to finding all sizes of intersections with respect to the hyperplanes of a fixed maximum h-scattered linear set, i.e. its intersection numbers with respect to the hyperplanes. In Section 3, we mentioned that the possible intersection numbers of a maximum h-scattered linear set with respect to hyperplane are among a collection of $h+1$ values, but it is not known whether for each of these values we may find at least one hyperplane intersecting the linear set in such a number of points. Here, we shall show that for $h = r-1$, that all the aforementioned intersection numbers are nonzero and hence we get an r-weight code.

5.1 Connections with MRD-codes

Before considering the $h = r - 1$ case, i.e. scattered linear sets with respect to hyperplanes, we briefly point out some basic properties of MRD-codes and their connection with such linear sets. A *rank metric (or RM) code* $\mathcal{C}$ of $\mathbb{F}_q^{n\times m}$, $n \leq m$, can be considered as a subset of $\mathrm{Hom}_{\mathbb{F}_q}(U, V)$, where $\dim_{\mathbb{F}_q} U = m$ and $\dim_{\mathbb{F}_q} V = n$, with *rank distance* defined as $d(f, g) := \mathrm{rk}(f - g)$, i.e. $\dim_{\mathbb{F}_q} \mathrm{Im}(f - g)$. The minimum distance of $\mathcal{C}$ is $d := \min\{d(f, g)\colon f, g \in \mathcal{C}, f \neq g\}$.

Result 5.1. *[12] If $\mathcal{C}$ is a rank metric code of $\mathbb{F}_q^{n\times m}$, $n \leq m$, with minimum distance d, then*

$$|\mathcal{C}| \leq q^{m(n-d+1)}. \tag{4}$$

Rank metric codes for which (4) holds with equality are called *maximum rank distance (or MRD) codes*.

We will only consider $\mathbb{F}_q$-linear MRD-codes of $\mathbb{F}_q^{n\times n}$, i.e. those which can be identified with $\mathbb{F}_q$-subspaces of $\mathrm{End}_{\mathbb{F}_q}(\mathbb{F}_{q^n})$. Since $\mathrm{End}_{\mathbb{F}_q}(\mathbb{F}_{q^n})$ is isomorphic to the ring $\mathcal{L}_{n,q}$ of q-polynomials over $\mathbb{F}_{q^n}$ modulo $x^{q^n} - x$, i.e.

$$\mathcal{L}_{n,q} \simeq \left\{ \sum_{i=0}^{n-1} a_i x^{q^i},\ \ a_i \in \mathbb{F}_{q^n} \right\},$$

with addition and composition as operations, we will consider $\mathcal{C}$ as an $\mathbb{F}_q$-subspace of $\mathcal{L}_{n,q}$, see e.g. [35]. We will need the following property on the number A_i of matrices in $\mathcal{C}$ having rank i.

Result 5.2. *[26, Lemma 2.1] If $\mathcal{C}$ is an MRD-code of $\mathbb{F}_q^{n\times n}$ with minimum distance d, then $A_i > 0$ for each i such that $d \leq i \leq n$.*

More generally, the weight distribution of an MRD-code was precisely determined by Delsarte in [12] (and later by Gabidulin in [20]).

Result 5.3. *Let $\mathcal{C}$ be an MRD-code in $\mathbb{F}_q^{n\times n}$ with minimum distance d. Then*

$$A_{d+\ell} = \begin{bmatrix} n \\ d+\ell \end{bmatrix}_q \sum_{t=0}^{\ell} (-1)^{t-l} \begin{bmatrix} \ell+d \\ \ell-t \end{bmatrix}_q q^{\binom{\ell-t}{2}} (q^{n(t+1)} - 1), \tag{5}$$

for $\ell \in \{0, 1, \ldots, n-d\}$.

Given two $\mathbb{F}_q$-linear RM-codes, $\mathcal{C}_1$ and $\mathcal{C}_2$, they are equivalent if and only if there exist two invertible $\mathbb{F}_q$-linear map φ_1, $\varphi_2 \in \mathcal{L}_{n,q}$ and $\rho \in \mathrm{Aut}(\mathbb{F}_q)$ such that

$$\varphi_1 \circ f^\rho \circ \varphi_2 \in \mathcal{C}_2 \text{ for all } f \in \mathcal{C}_1,$$

where $\circ$ stands for the composition of maps modulo $x^{q^n} - x$ and $f^\rho(x) = \sum a_i^\rho x^{q^i}$ for $f(x) = \sum a_i x^{q^i}$. For a rank metric code $\mathcal{C}$, its left and right idealisers can be written as:

$$L(\mathcal{C}) = \{\varphi \in \mathcal{L}_{n,q} \colon \varphi \circ f \in \mathcal{C} \text{ for all } f \in \mathcal{C}\},$$

$$R(\mathcal{C}) = \{\varphi \in \mathcal{L}_{n,q} \colon f \circ \varphi \in \mathcal{C} \text{ for all } f \in \mathcal{C}\},$$

see [23] and [26].

The relation between scattered linear sets with respect to hyperplanes and MRD-codes has been pointed out in [24] and in [32], see also [9, Section 4.1]. More precisely, $\mathcal{C}$ is an $\mathbb{F}_q$-linear MRD-code of $\mathcal{L}_{n,q}$ with minimum distance $n-r+1$ and with left-idealiser isomorphic to $\mathbb{F}_{q^n}$ if and only if $\mathcal{C}$ is equivalent to

$$\langle f_1(x), \dots, f_r(x) \rangle_{\mathbb{F}_{q^n}}$$

for some $f_1, f_2, \dots, f_r \in \mathcal{L}_{n,q}$, which happens if and only if

$$L_U = \{\langle (f_1(x), \dots, f_r(x)) \rangle_{\mathbb{F}_{q^n}} \colon x \in \mathbb{F}_{q^n}^*\}$$

is a scattered $\mathbb{F}_q$-linear set with respect to hyperplanes of rank n in $\Lambda = \mathrm{PG}(\mathbb{F}_{q^n}^r, \mathbb{F}_{q^n}) = \mathrm{PG}(r-1, q^n)$.

5.2 Scattered $\mathbb{F}_q$-linear sets with respect to hyperplanes

Let L_U be a scattered linear set with respect to the hyperplanes of $\Lambda = \mathrm{PG}(r-1, q^n)$ with $\mathrm{rk}(L_U) = n$, i.e. L_U is a maximum $(r-1)$-scattered linear set. Since $\mathrm{rk}(L_U) = n$, then there exist $f_1(x), \dots, f_r(x) \in \mathcal{L}_{n,q}$ such that

$$L_U = \{\langle (f_1(x), \dots, f_r(x)) \rangle_{\mathbb{F}_{q^n}} \colon x \in \mathbb{F}_{q^n}^*\}.$$

For each hyperplane $\mathcal{H} = \mathrm{PG}(W, \mathbb{F}_{q^n}) = \mathrm{PG}(r-2, q^n)$ of $\Lambda = \mathrm{PG}(r-1, q^n)$, by (3) we get that

$$w_{L_U}(\mathcal{H}) \in \{0, \dots, r-1\},$$

i.e.

$$|\mathcal{H} \cap L_U| \in \{0, 1, q+1, \dots, q^{r-2} + \cdots + q + 1\}. \tag{6}$$

We show that for each $i \in \{0, \dots, r-1\}$ there exists at least one hyperplane $\mathcal{H}$ such that $w_{L_U}(\mathcal{H}) = i$.

Remark 5.4. *Note that if* $\mathcal{H} : a_0x_0 + \cdots + a_{r-1}x_{r-1} = 0$ *is a hyperplane of* Λ*, then*

$$w_{L_U}(\mathcal{H}) = n - \mathrm{rk}(a_0f_1(x) + \cdots + a_{r-1}f_r(x)) = \dim_{\mathbb{F}_q} \ker(a_0f_1(x) + \cdots + a_{r-1}f_r(x)),$$

c.f. [32, Proposition 4.2]

We prove that L_U is an r-character set with respect to hyperplanes, i.e. it has r distinct intersection numbers with respect to the hyperplanes.

Theorem 5.5. *If* L_U *is a scattered linear set with respect to hyperplanes in* $\Lambda = \mathrm{PG}(r-1, q^n)$*, then it is an* r*-character set with respect to hyperplanes with intersection numbers given in* (6)*.*

Proof. Consider $i \in \{0, \dots, r-1\}$, since L_U is a scattered linear set with respect to hyperplanes, then by Subsection 5.1 the rank metric code $\mathcal{D} = \langle f_1(x), \dots, f_r(x) \rangle_{\mathbb{F}_{q^n}}$ is an MRD-code. In particular, by Result 5.2, there exists in $\mathcal{D}$ at least one q-polynomial, say $f(x) = a_0x + \ldots + a_{r-1}x^{q^{r-1}}$, whose image has dimension $n - i$ over $\mathbb{F}_q$, i.e. its kernel has dimension i. Let $\mathcal{H}\colon a_0x_0 + \ldots + a_{r-1}x_{r-1} = 0$. By Remark 5.4, it follows that the hyperplane $\mathcal{H}$ has weight i with respect to L_U, i.e. it meets L_U in $\theta_{i-1} = q^{i-1} + \ldots + q + 1$ points. □

Corollary 5.6. *The number of hyperplanes in* Λ *having weight* i *w.r.t* L_U *(i.e. the number of hyperplanes meeting* L_U *in* θ_{i-1} *points) is* A_{n-i}*, where* A_j*'s are defined as in* (5) *and* $i \in \{0, \dots, r-1\}$*.*

Therefore, as a consequence Theorem 4.1 and Theorem 5.5 we have the following result.

Corollary 5.7. *If* L_U *is a maximum* $(r-1)$*-scattered in* Λ*, then* $\mathcal{C}_{L_U}$ *is an* r*-weight code with weights*

$$w_i = N - \theta_{i-1},$$

with $i \in \{0, \dots, r-1\}$*. Its weight enumerator is*

$$1 + \sum_{i=0}^{r-1} A_{n-i} z^{w_i},$$

where A_j*'s are given in* (5)*.*

6 Weights for $h = 2$

Let L_U be a maximum 2-scattered linear set in $\Lambda = \mathrm{PG}(r-1, q^n)$, in particular $\mathrm{rk}(L_U) = \frac{rn}{3}$. Since L_U is scattered we also have that $|L_U| = \theta_{\frac{rn}{3}-1}$. By (3), we have that for each hyperplane $\mathcal{H} = \mathrm{PG}(r-1, q^n)$ it follows that

$$w_{L_U}(\mathcal{H}) \in \left\{ \frac{(r-3)n}{3}, \frac{(r-3)n}{3} + 1, \frac{(r-3)n}{3} + 2 \right\},$$

i.e.

$$|\mathcal{H} \cap L_U| \in \left\{ \theta_{\frac{(r-3)n}{3}-1}, \theta_{\frac{(r-3)n}{3}}, \theta_{\frac{(r-3)n}{3}+1} \right\}.$$

We are going to show that L_U is a three-character set with respect to hyperplanes.

Remark 6.1. *In this section, $n \geq 3$, since we are considering linear sets generating the all space with rank $\frac{rn}{3}$, i.e.*

$$r \leq \mathrm{rk}(L_U) \leq \frac{rn}{3},$$

which cannot be satisfied when $n \leq 2$.

Denote by t_i the number of hyperplanes of Λ with weight $\frac{(r-3)n}{3} + i$ with respect to L_U, for $i \in \{0, 1, 2\}$. Since for each line ℓ of Λ we have that $|L_U \cap \ell| \leq q+1$ and counting the hyperplanes, point-hyperplane pairs $(A, \mathcal{H})$ with $A \in \mathcal{H} \cap L_U$ and point-point-hyperplane triples $(A, B, \mathcal{H})$ with $A \neq B$ and $A, B \in \mathcal{H} \cap L_U$, we get the following equations in the variables t_0, t_1 and t_2,

$$\begin{aligned}
&t_0 + t_1 + t_2 = \tfrac{q^{nr}-1}{q^n-1};\\
&\theta_{\frac{(r-3)n}{3}-1} t_0 + \theta_{\frac{(r-3)n}{3}} t_1 + \theta_{\frac{(r-3)n}{3}+1} t_2 = \theta_{\frac{rn}{3}-1} \tfrac{q^{n(r-1)}-1}{q^n-1};\\
&\theta_{\frac{(r-3)n}{3}-1} \left(\theta_{\frac{(r-3)n}{3}-1} - 1\right) t_0 + \theta_{\frac{(r-3)n}{3}} \left(\theta_{\frac{(r-3)n}{3}} - 1\right) t_1 + \theta_{\frac{(r-3)n}{3}+1} \left(\theta_{\frac{(r-3)n}{3}+1} - 1\right) t_2 =\\
&= \theta_{\frac{rn}{3}-1} \left(\theta_{\frac{rn}{3}-1} - 1\right) \tfrac{q^{n(r-2)}-1}{q^n-1}.
\end{aligned} \tag{7}$$

By the previous system we get that

$$t_0 = \frac{\left(q^{\frac{rn}{3}}-1\right)}{(q-1)^2(q+1)\left(q^n-1\right)} t_0^*;$$
$$t_1 = \frac{\left(q^{\frac{rn}{3}}-1\right)\left(q^{\frac{rn}{3}+2}-q^{\frac{rn}{3}+1}-q^n+q^2\right)}{(q-1)^2 q}; \tag{8}$$
$$t_2 = \frac{\left(q^n-q\right)\left(q^{\frac{rn}{3}}-1\right)}{(q-1)^2 q(q+1)},$$

where

$$t_0^* = q^{\frac{2rn}{3}+3}-q^{\frac{2rn}{3}+2}-q^{\frac{2rn}{3}+1}+q^{\frac{2rn}{3}}-q^{\frac{rn}{3}+n+2}+q^{\frac{rn}{3}+3}-q^{\frac{rn}{3}+1}+q^{\frac{1}{3}(r+3)n}+q^{2n}-q^{n+2}-q^{n+1}+q^3.$$

Remark 6.2. *For $r = 3$, maximum 2-scattered linear sets correspond to scattered with respect to lines (hyperplanes) in $\Lambda = \mathrm{PG}(2, q^n)$. Hence, by Theorem 5.5 we have that t_0, t_1 and t_2 are positive and by Corollary 5.6 it follows that*

$$t_i = A_{n-i}, \text{ for } i \in \{0, 1, 2\}.$$

Lemma 6.3. *The values t_0, t_1 and t_2 are nonzero for each $r \geq 3$, $q \geq 2$ and $n \geq 3$.*

Proof. By Remark 6.2, we may assume $r \geq 4$. Since $q \geq 2$, $n \geq 3$ and $r \geq 4$, it is clear that $t_2 > 0$. First, note that for any q

$$q^k > \frac{q^k-1}{q-1} = q^{k-1} + q^{k-2} + \ldots + q + 1. \tag{9}$$

Let $m = \frac{rn}{3}$. Then $m > n \geq 3$ and $t_1 > 0$ follows from

$$q^{m+2} - q^{m+1} - q^n + q^2 > q^{m+2} - q^{m+1} - q^m > 0,$$

by (9).

Now, we show that $t_0 > 0$. This follows from

$$t_0^* = q^{2m+3}-q^{2m+2}-q^{2m+1}-q^{m+n+2}+q^{2m}+q^{m+3}+q^{m+n}-q^{m+1}+q^{2n}-q^{n+2}-q^{n+1}+q^3 =$$
$$= q^{m+n+2}(q^{m-n+1}-q^{m-n}-q^{m-n-1}-1)+q^m(q^m+q^3+q^n-q)+q^n(q^n-q^2-q)+q^3$$
$$> q^{m+n+2}(q^{m-n+1}-q^{m-n}-q^{m-n-1}-1)+q^m(q^m+q^3+q^n-q)+q^n(q^n-q^2-q) > 0,$$

by (9) and since $q \geq 2$ and $n \geq 3$. ∎

Therefore, we have the following.

Theorem 6.4. *The linear set L_U is a three-character set with respect to the hyperplanes with intersection numbers $\theta_{\frac{(r-3)n}{3}-1}, \theta_{\frac{(r-3)n}{3}}$ and $\theta_{\frac{(r-3)n}{3}+1}$. Also, the number of hyperplanes intersecting L_U in $\theta_{\frac{(r-3)n}{3}+i-1}$ points is given by t_i, with $i \in \{0,1,2\}$.*

Whereas, using Theorems 4.1 and 6.4, we have the following.

Corollary 6.5. *If L_U is a maximum 2-scattered linear set in Λ, then $\mathcal{C}_{L_U}$ is a three-weight code with weights*

$$w_0 := N - \theta_{\frac{(r-3)n}{3}-1},$$

$$w_1 := N - \theta_{\frac{(r-3)n}{3}},$$

and

$$w_2 := N - \theta_{\frac{(r-3)n}{3}+1}.$$

Its weight enumerator is

$$1 + t_0 z^{w_0} + t_1 z^{w_1} + t_2 z^{w_2},$$

where t_i's are defined as in (8).

6.1 Maximum scattered linear sets with respect to lines of the plane

When $h = 2$ and $r = 3$, by (8), the number of lines t_i with weight i with respect to L_U are

$$t_0 = \frac{3q^{2n} - 2q^{n+1} - q^{n+2} + q^{n+3} - q^{2n+1} - 2q^{2n+2} + q^{2n+3} + q^3}{(q-1)^2(q+1)};$$

$$t_1 = \frac{(q^n - 1)\left(-q^{n+1} + q^{n+2} - q^n + q^2\right)}{(q-1)^2 q};$$

$$t_2 = \frac{(q^n - 1)(q^n - q)}{(q-1)^2 q(q+1)}.$$

By the aforementioned construction, we get a three-weight code $\mathcal{C}_{L_U}$ of dimension 3 and length N, with weights

$$w_1 = N = \frac{q^n - 1}{q - 1};$$

$$w_2 = N - 1 = q\frac{q^{n-1} - 1}{q - 1};$$

$$w_3 = N - q - 1 = q^2\frac{q^{n-2} - 1}{q - 1}.$$

Furthermore, the weight enumerator is

$$1 + t_2 z^{N-q-1} + t_1 z^{N-1} + t_0 z^N.$$

Remark 6.6. *Since the minimum distance is $N-q-1$, the Singleton bound for such a code is*

$$N - q - 1 \leq N - 2.$$

In particular, the defect in the Singleton bound is $q - 1$. For $q = 2$ we get an almost MDS code*, i.e. a code for which the minimum distance is the difference of its length and its dimension. De Boer in [11] introduced those codes for the first time and very recently in [27] they have been used in secret sharing schemes.*

6.2 Comparison with known families of three-weight codes

Most of the known three-weight codes beloging to the first family mentioned in Section 2 have the property that their lenght divides the order of the base field minus one, see [13, Section VI]. Our construction provides codes for which this property does not hold.

When $r > 3$ the codes we construct are q-divisible, whereas for $r = 3$ they are not, contrary to those presented in [1]. Furthermore, the codes introduced in [1] are defined over $\mathbb{F}_{q^2}$, whereas our construction yield codes defined over $\mathbb{F}_{q^n}$, for any $n \geq 3$.

7 Generator matrices

The aim of this section is to exhibit generator matrices of the aforementioned linear codes.

We start by considering an $\mathbb{F}_q$-linear set L_U which is scattered with respect to hyperplanes of rank n in $\mathrm{PG}(r-1,q^n)$, which, as seen before, may be written as

$$L_U = \{\langle (f_1(x),\dots,f_r(x))\rangle_{\mathbb{F}_{q^n}} : x \in \mathbb{F}_{q^n}\},$$

with $f_1(x),\dots,f_r(x) \in \mathcal{L}_{n,q}$. Therefore, a matrix whose columns correspond to the following set of vectors

$$\{(f_1(x),\dots,f_r(x)) : x \in \mathbb{F}_{q^n}^*\}$$

is a generator matrix for the code $\mathcal{C}_{L_U}$.

In the following table, we resume the known choices of the polynomials $f_1,\dots,f_r$ defining scattered $\mathbb{F}_q$-linear sets with respect to the hyperplanes in $\mathrm{PG}(r-1,q^n)$.

n	r	$(f_1(x), \ldots, f_r(x))$	conditions	references
		$(x, x^{q^s}, \ldots, x^{q^{s(r-1)}})$	$\gcd(s,n)=1$	[12, 20, 21]
		$(x^{q^s}, \ldots, x^{q^{s(r-2)}}, x+\delta x^{q^{s(r-1)}})$	$\gcd(s,n)=1$, $\mathrm{N}_{q^n/q}(\delta) \neq (-1)^{nr}$	[31, 25]
6	4	$(x^q, x^{q^2}, x^{q^4}, x-\delta^{q^5} x^{q^3})$	$q>4$ certain choices of δ	[7, 30]
6	4	$(x^q, x^{q^3}, x-x^{q^2}, x^{q^4}-\delta x)$	q odd $\delta^2+\delta=1$	[10, 29]
6	4	$(h^{q^2-1}x^q + h^{q-1}x^{q^2}, x^{q^3}, x^q - h^{q-1}x^{q^4}, x^q - h^{q-1}x^{q^5})$	q odd $h^{q^3+1}=-1$	[3, 37]
7	3	(x, x^q, x^{q^3})	q odd, $\gcd(s,7)=1$	[8]
7	4	$(x, x^{q^{2s}}, x^{q^{3s}}, x^{q^{4s}})$	q odd, $\gcd(s,n)=1$	[8]
8	3	(x, x^q, x^{q^3})	$q \equiv 1 \pmod 3$, $\gcd(s,8)=1$	[8]
8	5	$(x, x^{q^{2s}}, x^{q^{3s}}, x^{q^{4s}}, x^{5s})$	$q \equiv 1 \pmod 3$, $\gcd(s,8)=1$	[8]
8	6	$(x^q, x^{q^2}, x^{q^3}, x^{q^5}, x^{q^6}, x-\delta x^{q^4})$	q odd, $\delta^2=-1$	[7]

Table 1: Possible choices for $f_1, \ldots, f_r$

Hence, we have the following result.

Theorem 7.1. *Let q be a prime power, r, n be two positive integers with $r \geq 3$, let $(f_1(x), \ldots, f_r(x))$ be as in Table 1 and $N = \frac{q^n-1}{q-1}$. If G is a $(r \times N)$-matrix whose set of columns is*

$$\{(f_1(x), \ldots, f_r(x)) \colon x \in \mathbb{F}_{q^n}^*\},$$

then G is the generator matrix of a r-weight linear $[N, r]_{q^n}$-code.

For maximum 2-scattered $\mathbb{F}_q$-linear sets we have less examples. A general construction for h-scattered linear sets is given by the following result.

Theorem 7.2. [9, Theorem 2.5] *Let $V = V_1 \oplus \ldots \oplus V_t$, let L_{U_i} a h_i-scattered $\mathbb{F}_q$-linear set in $\mathrm{PG}(V_i, \mathbb{F}_{q^n})$, with $i \in \{1, \ldots, t\}$ and let*

$$U = U_1 \oplus \ldots \oplus U_t.$$

The linear set L_U is h-scattered in $\Lambda = \mathrm{PG}(V, \mathbb{F}_{q^n})$, with $h = \min\{h_1, \ldots, h_t\}$.

Hence, suppose that $3 \mid r$. So, $r = 3t$ and let $\Lambda = \mathrm{PG}(\mathbb{F}_{q^n}^{3t}, \mathbb{F}_{q^n})$. Let $\pi = \mathrm{PG}(W_i, \mathbb{F}_{q^n})$ be the projective subspace of Λ defined by

$$\pi_i \colon x_j = 0, \quad j \in \{1, \ldots, 3t\} \setminus \{3i-2, 3i-1, 3i\}$$

for $i \in \{1, \ldots, t\}$.

Thus we have

$$\Lambda = \langle \pi_1, \ldots, \pi_t \rangle,$$

and

$$\mathbb{F}_{q^n}^{3t} = W_1 \oplus \ldots \oplus W_t.$$

For each $i \in \{1, \ldots, t\}$, choose

$$(f_1^i(x), f_2^i(x), f_3^i(x))$$

as in Table 1 with $r = 3$. Then consider U_i the set of vectors $(a_1, \ldots, a_{3t})$

$$a_j = \begin{cases} f_1^i(x) & \text{if } j = i \\ f_2^i(x) & \text{if } j = i+1 \\ f_3^i(x) & \text{if } j = i+2 \\ 0 & \text{otherwise.} \end{cases}$$

It follows that U_i is an $\mathbb{F}_q$-subspace and L_{U_i} is a 2-scattered $\mathbb{F}_q$-linear set in π_i. Therefore, by Theorem 7.2 we have that

$$U = U_1 \oplus \ldots \oplus U_t =$$

$$\{(f_1^1(x_1), f_2^1(x_1), f_3^1(x_1), \ldots, f_1^t(x_t), f_2^t(x_t), f_3^t(x_t))\colon x_i \in \mathbb{F}_{q^n}\}$$

defines a maximum 2-scattered $\mathbb{F}_q$-linear set in Λ. Hence, a $(3\times\theta_{tn-1})$-matrix G over $\mathbb{F}_{q^n}$ whose columns correspond to the following set of vectors

$$\{(f_1^1(x_1), f_2^1(x_1), f_3^1(x_1), \ldots, f_1^t(x_t), f_2^t(x_t), f_3^t(x_t))\colon x_i \in \mathbb{F}_{q^n}\} \setminus \{\mathbf{0}\}$$

is a generator matrix for the code $\mathcal{C}_{L_U}$.

Therefore, we have the following.

Theorem 7.3. *Let q be a prime power, t be a positive integer $t \geq 2$, $N = \theta_{tn-1} = \frac{q^{nt}-1}{q-1}$ and $(f_1^j(x), f_2^j(x), f_3^j(x))$ as in Table 1 for each $j \in \{1, \ldots, t\}$. If G is a $(3\times N)$-matrix G over $\mathbb{F}_{q^n}$ whose columns correspond to the following set of vectors*

$$\{(f_1^1(x_1), f_2^1(x_1), f_3^1(x_1), \ldots, f_1^t(x_t), f_2^t(x_t), f_3^t(x_t))\colon x_i \in \mathbb{F}_{q^n}\} \setminus \{\mathbf{0}\},$$

then G is the generator matrix of a three-weight linear $[N, 3]_{q^n}$-code.

References

[1] A. Aguglia and L. Giuzzi: Intersection sets, three-character multisets and associated codes, *Des. Codes Cryptogr.* **83** (2017), 269–282.

[2] T. L. Alderson: A note on full weight spectrum codes, *Trans. on Combinatorics* **8** No. 3 (2019), 15–22.

[3] D. Bartoli, C. Zanella and F. Zullo: A new family of maximum scattered linear sets in PG$(1, q^6)$, *Ars Math. Contemp.* (forthcoming) arXiv:1910.02278.

[4] A. Blokhuis and M. Lavrauw: Scattered spaces with respect to a spread in PG(n, q), *Geom. Dedicata* **81** (2000), 231–243.

[5] A. R. Calderbank and J. M. Goethals: Three-weight codes and association schemes, *Philips J. Res.* **39**(4.5) (1984), 143-152.

[6] A. R. Calderbank and W. M. Kantor: The geometry of two-weight codes, *Bull. London Math. Soc.* **18** (1986), 97-122.

[7] B. Csajbók, G. Marino, O. Polverino and C. Zanella: A new family of MRD-codes, *Linear Algebra Appl.* **548** (2018), 203–220.

[8] B. Csajbók, G. Marino, O. Polverino and Y. Zhou: Maximum Rank-Distance codes with maximum left and right idealisers, arXiv:1807.08774.

[9] B. Csajbók, G. Marino, O. Polverino and F. Zullo: A special class of scattered subspaces, arXiv:1906.10590.

[10] B. Csajbók, G. Marino and F. Zullo: New maximum scattered linear sets of the projective line, *Finite Fields Appl.* **54** (2018), 133–150.

[11] M.A. De Boer: Almost MDS codes, *Des. Codes Cryptogr.* **9(2)** (1996), 143–155.

[12] P. Delsarte: Bilinear forms over a finite field, with applications to coding theory, *J. Combin. Theory Ser. A* **25** (1978), 226–241.

[13] K. Ding and C. Ding A Class of Two-Weight and Three-Weight Codes and Their Applications in Secret Sharing, *IEEE Trans. Inform. Theory* **61(11)** (2015), 5835–5842.

[14] K. Ding K. and C. Ding: Binary linear codes with three weights, *IEEE Commun. Lett.* **18** (2014), 1879–1882.

[15] C. Ding, C. Li, N. Li and Z. Zhou: Three-weight cyclic codes and their weight distributions, *Discret. Math.* **339** (2016), 415–427.

[16] C. Ding, J. Luo and H. Niederreiter: Two-weight codes punctured from irreducible cyclic codes, *Proc. Ist Int. Workshop Coding theory and Cryptogr.* (2008), 119–124.

[17] C. Ding and H. Niederreiter: Cyclotomic linear codes of order 3, *IEEE Trans. Inf. Theory* **53.6** (2007), 2274-2277.

[18] C. Ding and X. Wang: A coding theory construction of new systematic authentication codes, *Theoretical computer science* **330.1** (2005), 81–99.

[19] N. Durante: On Sets with Few Intersection Numbers in Finite Projective and Affine Spaces, *Electron. J. Combin.* **21 (4)** (2014), 1–18.

[20] E. Gabidulin: Theory of codes with maximum rank distance, *Problems of information transmission*, **21(3)** (1985), 3–16.

[21] A. Kshevetskiy and E. Gabidulin: The new construction of rank codes, *International Symposium on Information Theory*, 2005. ISIT 2005. Proceedings, pages 2105–2108, Sept. 2005.

[22] M. Lavrauw: Scattered spaces with respect to Spreads, and Eggs in Finite Projective Spaces, *Ph.D thesis*, Eindhoven University of Technology (2001).

[23] D. Liebhold and G. Nebe: Automorphism groups of Gabidulin-like codes, *Arch. Math.* **107(4)** (2016), 355–366.

[24] G. Lunardon: MRD-codes and linear sets, *J. Combin. Theory Ser. A* **149** (2017), 1–20.

[25] G. Lunardon, R. Trombetti and Y. Zhou: Generalized Twisted Gabidulin Codes, *J. Combin. Theory Ser. A* **159** (2018), 79–106.

[26] G. Lunardon, R. Trombetti and Y. Zhou: On kernels and nuclei of rank metric codes, *J. Algebraic Combin.* **46** (2017), 313–340.

[27] S. Mehta, V. Saraswat and S. Sen: Secret sharing using near-MDS codes, *Codes, Cryptology, and Information Security (C2SI 2019). LNCS,* vol. 11445, pp. 195–214. Springer (2019).

[28] V. Napolitano, O. Polverino, G. Zini and F. Zullo: Linear sets from projection of Desarguesian spreads, arXiv:2001.08685.

[29] G. Marino, M. Montanucci and F. Zullo: MRD-codes arising from the trinomial $x^q + x^{q^3} + cx^{q^5} \in \mathbb{F}_{q^6}[x]$, *Linear Algebra Appl.* **591** (2020), 99–114.

[30] O. Polverino and F. Zullo: On the number of roots of some linearized polynomials, arXiv:1909.00802.

[31] J. Sheekey: A new family of linear maximum rank distance codes, *Adv. Math. Commun.* **10(3)** (2016), 475–488.

[32] J. Sheekey and G. Van de Voorde: Rank-metric codes, linear sets and their duality, *Des. Codes Cryptogr.* (2019), https://doi.org/10.1007/s10623-019-00703-z.

[33] M. Shi and P.Solé: Three-weight codes, triple sum sets, and strongly walk regular graphs, *Designs, Codes and Cryptogr.* **87** (2019) 2395–2404.

[34] M.A. Tsfasman, S.G. Vlăduţ and D. Nogin: Algebraic geometric codes: basic notions, *mathematical surveys and monographs, American Mathematical society* **139** (2007).

[35] B. Wu and Z. Liu: Linearized polynomials over finite fields revisited, *Finite Fields Appl.* **22** (2013), 79–100.

[36] Y. Wu, Q. Yansheng and X. Shi: At most three-weight binary linear codes from generalized Moisios exponential sums, *Designs, Codes and Cryptogr.* **87(8)** (2019), 1927–1943.

[37] C. Zanella and F. Zullo: Vertex properties of maximum scattered linear sets of $\mathrm{PG}(1, q^n)$, *Discrete Math.* **343(5)** (2020).

Vito Napolitano and Ferdinando Zullo
Dipartimento di Matematica e Fisica,
Università degli Studi della Campania "Luigi Vanvitelli",
Viale Lincoln, 5
I– 81100 Caserta, Italy
{*vito.napolitano,ferdinando.zullo*}*@unicampania.it*