

Quasi-majority Functional Voting on Expander Graphs

Nobutaka Shimizu*

Takeharu Shiraga†

January 8, 2022

Abstract

Consider a distributed graph where each vertex holds one of two distinct opinions. In this paper, we are interested in synchronous *voting processes* where each vertex updates its opinion according to a predefined common local updating rule. For example, each vertex adopts the majority opinion among 1) itself and two randomly picked neighbors in *best-of-two* or 2) three randomly picked neighbors in *best-of-three*. Previous works intensively studied specific rules including best-of-two and best-of-three individually.

In this paper, we generalize and extend previous works of best-of-two and best-of-three on expander graphs by proposing a new model, *quasi-majority functional voting*. This new model contains best-of-two and best-of-three as special cases. We show that, on expander graphs with sufficiently large initial bias, any quasi-majority functional voting reaches consensus within $O(\log n)$ steps with high probability. Moreover, we show that, for any initial opinion configuration, any quasi-majority functional voting on expander graphs with higher expansion (e.g., Erdős-Rényi graph $G(n, p)$ with $p = \Omega(1/\sqrt{n})$) reaches consensus within $O(\log n)$ with high probability. Furthermore, we show that the consensus time is $O(\log n / \log k)$ of best-of- $(2k + 1)$ for $k = o(n / \log n)$.

Keywords: Distributed voting, consensus problem, expander graph, Markov chain

1 Introduction

Consider an undirected graph $G = (V, E)$ where each vertex $v \in V$ initially holds an opinion $\sigma \in \Sigma$ from a finite set Σ . In *synchronous voting process* (or simply, *voting process*), in each round, every vertex communicates with its neighbors and then all vertices simultaneously update their opinions according to a predefined protocol. The aim of the protocol is to reach a *consensus configuration*, i.e., a configuration where all vertices have the same opinion. Voting process has been extensively studied in several areas including biology, network analysis, physics and distributed computing [10, 33, 31, 23, 27, 2]. For example, in distributed computing, voting process plays an important role in the consensus problem [23, 27].

This paper is concerned with the *consensus time* of voting processes over *binary* opinions $\Sigma = \{0, 1\}$. Then voting processes have state space 2^V . A state of 2^V is called a *configuration*. The *consensus time* is the number of steps needed to reach a consensus configuration.

1.1 Previous works of specific updating rules

In *pull voting*, in each round, every vertex adopts the opinion of a randomly selected neighbor. This is one of the most basic voting process, which has been well explored in the past [34, 28, 14, 18, 8]. In

*The University of Tokyo, Japan. nobutaka.shimizu@mist.i.u-tokyo.ac.jp

†Chuo University, Japan. shiraga.076@g.chuo-u.ac.jp

particular, the expected consensus time of this process has been extensively studied in the literature. For example, Hassin and Peleg [28] showed that the expected consensus time is $O(n^3 \log n)$ for all non-bipartite graphs and all initial opinion configurations, where n is the number of vertices. From the result of Cooper, Elsässer, Ono, and Radzik [14], it is known that on the complete graph K_n , the expected consensus time is $O(n)$ for any initial opinion configuration.

In *best-of-two* (a.k.a. *2-Choices*), each vertex v samples two random neighbors (with replacement) and, if both hold the same opinion, v adopts the opinion. Otherwise, v keeps its own opinion. Doerr, Goldberg, Minder, Sauerwald, and Scheideler [21] showed that, on the complete graph K_n , the consensus time of best-of-two is $O(\log n)$ with high probability¹ for an arbitrary initial opinion configuration. Since best-of-two is simple and is faster than pull voting on the complete graphs, this model gathers special attention in distributed computing and related area [26, 15, 16, 17, 19, 20, 39]. There is a line of works that study best-of-two on expander graphs [15, 16, 17], which we discuss later.

In *best-of-three* (a.k.a. *3-Majority*), each vertex v randomly selects three random neighbors (with replacement). Then, v updates its opinion to match the majority among the three. It follows directly from Ghaffari and Lengler [26] that, on K_n with any initial opinion configuration, the consensus time of best-of-three is $O(\log n)$ w.h.p. Kang and Rivera [29] considered the consensus time of best-of-three on graphs with large minimum degree starting from a random initial configuration. Shimizu and Shiraga [39] showed that, for any initial configurations, best-of-two and best-of-three reach consensus in $O(\log n)$ steps w.h.p. if the graph is an Erdős-Rényi graph $G(n, p)$ ² of $p = \Omega(1)$.

Best-of- k ($k \geq 1$) is a generalization of pull voting, best-of-two and best-of-three. In each round, every vertex v randomly selects k neighbors (with replacement) and then if at least $\lfloor k/2 \rfloor + 1$ of them have the same opinion, the vertex v adopts it. Note that the best-of-1 is equivalent to pull voting. Abdullah and Draief [1] studied a variant of best-of- k ($k \geq 5$ is odd) on a specific class of sparse graphs that includes n -vertex random d -regular graphs³ $G_{n,d}$ of $d = o(\sqrt{\log n})$ with a random initial configuration. To the best of our knowledge, best-of- k has not been studied explicitly so far.

In *Majority* (a.k.a. *local majority*), each vertex v updates its opinion to match the majority opinion among the neighbors. This simple model has been extensively studied in previous works [6, 9, 25, 35, 36, 41]. For example, Majority on certain families of graphs including the Erdős-Rényi random graph [6, 41], random regular graphs [25] have been investigated. See [36] for further details.

Voting process on expander graphs. Expander graph gathers special attention in the context of Markov chains on graphs, yielding a wide range of theoretical applications. A graph G is λ -*expander* if $\max\{|\lambda_2|, |\lambda_n|\} \leq \lambda$, where $1 = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq -1$ are the eigenvalues of the transition matrix P of the simple random walk on G . For example, an Erdős-Rényi graph $G(n, p)$ of $p \geq (1 + \epsilon) \frac{\log n}{n}$ for an arbitrary constant $\epsilon > 0$ is $O(1/\sqrt{np})$ -expander w.h.p. [12]. An n -vertex random d -regular graph $G_{n,d}$ of $3 \leq d \leq n/2$ is $O(1/\sqrt{d})$ -expander w.h.p. [13, 40].

Cooper et al. [14] showed that the expected consensus time of pull voting is $O(n/(1 - \lambda))$ on λ -expander regular graphs for any initial configuration. Compared to pull voting, the study of best-of-two on general graphs seems much harder. Most of the previous works concerning best-of-two on expander graphs put some assumptions on the initial configuration. Let A denote the set of vertices of opinion 0 and $B = V \setminus A$. Cooper, Elsässer, and Radzik [15] showed that, for any regular

¹In this paper “with high probability” (w.h.p.) means probability at least $1 - n^{-c}$ for a constant $c > 0$.

²Recall that the Erdős-Rényi random graph $G(n, p)$ is a graph on n vertices where each of possible $\binom{n}{2}$ vertex pairs forms an edge with probability p independently.

³An n -vertex random d -regular graph $G_{n,d}$ is a graph selected uniformly at random from the set of all labelled n -vertex d -regular graphs.

λ -expander graph, the consensus time is $O(\log n)$ w.h.p. if $||A| - |B|| = \Omega(\lambda n)$. This result was improved by Cooper, Elsässer, Radzik, Rivera, and Shiraga [16]. Roughly speaking, they proved that, on λ -expander graphs, the consensus time is $O(\log n)$ if $|d(A) - d(B)| = \Omega(\lambda^2 d(V))$, where $d(S) = \sum_{v \in S} \deg(v)$ denotes the volume of $S \subseteq V$. To the best of our knowledge, the worst case consensus time of best-of- k on expander graphs has not been studied.

1.2 Our model

In this paper, we propose a new class *functional voting* of voting process, which contains many known voting processes as a special case. Let $A \subseteq V$ be the set of vertices of opinion 0 and A' be the set in the next round. Let $B = V \setminus A$ and $B' = V \setminus A'$. For $v \in V$ and $S \subseteq V$, let $N(v) = \{w \in V : \{v, w\} \in E\}$ and $\deg_S(v) = |N(v) \cap S|$.

Definition 1.1 (Functional voting). *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying $f([0, 1]) = [0, 1]$ and $f(0) = 0$. A functional voting with respect to f is a synchronous voting process defined as*

$$\begin{aligned} \Pr[v \in A'] &= f\left(\frac{\deg_A(v)}{\deg(v)}\right) & \text{if } v \in B, \\ \Pr[v \in B'] &= f\left(\frac{\deg_B(v)}{\deg(v)}\right) & \text{if } v \in A. \end{aligned}$$

We call the function f a betrayal function and the function

$$H_f(x) := x(1 - f(1 - x)) + (1 - x)f(x)$$

an updating function.

Since $f(0) = 0$, consensus configurations are absorbing states. Hence the consensus time is well-defined⁴. The intuition behind the updating function H_f is that, letting $\alpha = |A|/n$ and $\alpha' = |A'|/n$, on a complete graph K_n (with self-loop), the functional voting with respect to f satisfies $\mathbf{E}[\alpha'] = \frac{|A|}{n} \left(1 - f\left(\frac{|B|}{n}\right)\right) + \frac{|B|}{n} f\left(\frac{|A|}{n}\right) = H_f(\alpha)$.

Functional voting contains many existing models as special cases. For example, pull voting, best-of-two, and best-of-three are functional votings with respect to x , x^2 and $3x^2 - 2x^3$, respectively. In general, best-of- k is a functional voting with respect to

$$f_k(x) = \sum_{i=\lfloor k/2 \rfloor + 1}^k \binom{k}{i} x^i (1 - x)^{k-i}. \quad (1)$$

It is straightforward to check that $H_{f_k}(x) = f_k(x)$ if k is odd and $H_{f_k}(x) = f_{k+1}(x)$ if k is even. Majority is a functional voting with respect to

$$f(x) = \begin{cases} 0 & \text{if } x < \frac{1}{2}, \\ \frac{1}{2} & \text{if } x = \frac{1}{2}, \\ 1 & \text{if } x > \frac{1}{2} \end{cases} \quad (2)$$

if a vertex adopts the random opinion when it meets the tie.

⁴For disconnected graphs, there exist initial configurations that never reach consensus. Henceforth, we are concerned with connected graphs.




Figure 1: The update functions $H_f(x)$ of pull voting (solid line), best-of-three (dashed line) and best-of-seven (dotted line). One can easily observe that best-of-three and best-of-seven are quasi-majority functional voting. Intuitively speaking, quasi-majority functional voting processes have updating functions H_f with the property so-called “the rich get richer”, which coincides with Definition 1.2.

Quasi-majority functional voting. In this paper, we focus on functional voting with respect to f satisfying the following property.

Definition 1.2 (Quasi-majority.). *A function f is quasi-majority if f satisfies the following conditions.*

- (1) f is C^2 ,
- (2) $0 < f(1/2) < 1$,
- (3) $H_f(x) < x$ whenever $x \in (0, 1/2)$.
- (4) $H'_f(1/2) > 1$,
- (5) $H'_f(0) < 1$.

A voting process is a quasi-majority functional voting if it is a functional voting with respect to a quasi-majority function f .

Note that $H_f(x)$ is symmetric (i.e., $H_f(1-x) = 1 - H_f(x)$) and thus the condition (3) implies $H_f(x) > x$ for every $x \in (1/2, 1)$. Intuitively, the conditions (3) to (5) ensure the drift towards consensus. The conditions (1) and (2) are due to a technical reasons.

For each constant $k \geq 2$, best-of- k is quasi-majority functional voting but pull voting and Majority are not. Indeed, if H_{f_k} is the updating function of best-of- k , then $H'_{f_{2\ell}}(x) = H'_{f_{2\ell+1}}(x) = (2\ell + 1) \binom{2\ell}{\ell} x^\ell (1-x)^\ell$. It is straightforward to check that this function satisfies the conditions (3) to (5) if $\ell \neq 0$ (pull-voting). See Figure 1 for depiction of the updating functions of pull voting, best-of-three and best-of-seven.

1.3 Our result

In this paper, we study the consensus time of quasi-majority functional voting on expander graphs⁵. Let $T_{\text{cons}}(A)$ denote the consensus time starting from the initial configuration $A \subseteq V$. For a graph $G = (V, E)$, let $\pi = (\pi(v))_{v \in V}$ denote the *degree distribution* defined as

$$\pi(v) = \frac{\deg(v)}{2|E|}. \quad (3)$$

Note that $\sum_{v \in V} \pi(v) = 1$ holds. We denote by $\|x\|_p := (\sum_{v \in V} |x_v|^p)^{1/p}$ the ℓ^p norm of $x \in \mathbb{R}^V$. For $\pi \in [0, 1]^V$ and $A \subseteq V$, let $\pi(A) := \sum_{v \in A} \pi(v)$. Let

$$\delta(A) := \pi(A) - \pi(V \setminus A) = 2\pi(A) - 1$$

denote the *bias* between A and $V \setminus A$.

Theorem 1.3 (Main theorem). *Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Then, the following holds:*

- (i) *Let $C_1 > 0$ be an arbitrary constant and $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}$ be an arbitrary function satisfying $\varepsilon(n) \rightarrow 0$ as $n \rightarrow \infty$. Suppose that $\lambda \leq C_1 n^{-1/4}$, $\|\pi\|_2 \leq C_1/\sqrt{n}$ and $\|\pi\|_3 \leq \varepsilon/\sqrt{n}$. Then, for any $A \subseteq V$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.*
- (ii) *Let C_2 be a positive constant depending only on f . Suppose that $\lambda \leq C_2$ and $\|\pi\|_2 \leq C_2/\sqrt{\log n}$. Then, for any $A \subseteq V$ satisfying $|\delta(A)| \geq C_2 \max\{\lambda^2, \|\pi\|_2 \sqrt{\log n}\}$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.*

The following result which we show in Section 5 indicates that the consensus time of Theorem 1.3(i) is optimal up to a constant factor.

Theorem 1.4 (Lower bound). *Under the same assumption of Theorem 1.3(i), $T_{\text{cons}}(A) = \Omega(\log n)$ w.h.p. for some $A \subseteq V$.*

Theorem 1.5 (Fast consensus for $H'_f(0) = 0$). *Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Let $C > 0$ be a constant depending only on f . Suppose that $H'_f(0) = 0$, $\lambda \leq C$ and $\|\pi\|_2 \leq C/\sqrt{\log n}$. Then, for any $A \subseteq V$ satisfying $|\delta(A)| \geq C \max\{\lambda^2, \|\pi\|_2 \sqrt{\log n}\}$, it holds w.h.p. that*

$$T_{\text{cons}}(A) = O\left(\log \log n + \log |\delta(A)|^{-1} + \frac{\log n}{\log \lambda^{-1}} + \frac{\log n}{\log(|\pi\|_2 \sqrt{\log n})^{-1}}\right).$$

For example, for each constant $k \geq 2$, best-of- k is quasi-majority with $H'_f(0) = 0$.

Remark 1.6. *Roughly speaking, for $p \geq 2$, $\|\pi\|_p$ measures the imbalance of the degrees. For any graphs, $\|\pi\|_p \geq n^{-1+1/p}$ and the equality holds if and only if the graph is regular. For star graphs, we have $\|\pi\|_p \approx 1$.*

⁵Throughout the paper, we consider sufficiently large $n = |V|$.

Results of best-of- k . Our results above do not explore Majority since it is not quasi-majority. A plausible approach is to consider best-of- k for $k = k(n) = \omega(1)$ since each vertex is likely to choose the majority opinion if the number of neighbor sampling increases. Also, note that the betrayal function f_k of best-of- k given in (1) converges to that of Majority (i.e., $f_k(x) \rightarrow f(x)$ as $k \rightarrow \infty$ for each $x \in [0, 1]$, where f is the betrayal function (2) of Majority). On the other hand, if $k = O(1)$, there is a tremendous gap between best-of- k and Majority: For any functional voting on the complete graph K_n , $T_{\text{cons}}(A) = \Omega(\log n)$ for some $A \subseteq V$ from Theorem 1.4. Majority on K_n reaches the consensus in a single step if $|A| < |V \setminus A| - 1$. This motivates us to consider best-of- k for $k = k(n) \rightarrow \infty$ as $n \rightarrow \infty$. For simplicity, we focus on best-of- $(2k + 1)$ and prove the following result in Section 6.

Theorem 1.7. *Let $k = k(n)$ be such that $k = \omega(1)$ and $k = o(n/\log n)$. Let C be an arbitrary positive constant. Consider best-of- $(2k+1)$ on an n -vertex λ -expander graph with degree distribution π such that $\lambda \leq Ck^{-1/2}n^{-1/4}$, $\|\pi\|_2 \leq Cn^{-1/2}$ and $\|\pi\|_3 \leq Ck^{-1/6}n^{-1/2}$. Then, $T_{\text{cons}}(A) = O\left(\frac{\log n}{\log k}\right)$ holds w.h.p. for any $A \subseteq V$.*

1.4 Application

Here, we apply our main theorem to specific graphs and derive some useful results.

For any $p \geq (1 + \epsilon)\frac{\log n}{n}$ for an arbitrary constant $\epsilon > 0$, $G(n, p)$ is connected and $O(1/\sqrt{np})$ -expander w.h.p [12, 24].

Corollary 1.8. *Consider a best-of- k on an Erdős-Rényi graph $G(n, p)$ for an arbitrary constant $k \geq 2$. Then, $G(n, p)$ w.h.p. satisfies the following:*

(i) *Suppose that $p = \Omega(n^{-1/2})$. Then*

(a) *for any $A \subseteq V$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.*

(b) *for some $A \subseteq V$, $T_{\text{cons}}(A) = \Omega(\log n)$ w.h.p.*

(ii) *Suppose that $p \geq (1 + \epsilon)\frac{\log n}{n}$ for an arbitrary constant $\epsilon > 0$. Then, for any $A \subseteq V$ satisfying $|\delta(A)| \geq C \max\left\{\frac{1}{np}, \sqrt{\frac{\log n}{n}}\right\}$, $T_{\text{cons}}(A) = O\left(\log \log n + \log |\delta(A)|^{-1} + \frac{\log n}{\log(np)}\right)$ w.h.p., where $C > 0$ is a constant depending only on f .*

In Corollary 1.8(i), we stress that the worst-case consensus time on $G(n, p)$ was known for $p = \Omega(1)$ [39]. If $\frac{\log n}{\log(np)} = O(\log \log n)$ (or equivalently, $np = n^{\Omega(1/\log \log n)}$), Corollary 1.8(ii) implies $T_{\text{cons}}(A) = O(\log \log n + \log |\delta(A)|^{-1})$ w.h.p.

Corollary 1.9. *Let $k = k(n)$ be such that $k = \omega(1)$ and $k = O(\sqrt{n})$. Consider best-of- $(2k + 1)$ on $G(n, p)$ for $p = \Omega(k/\sqrt{n})$. Then, for any $A \subseteq V$, $T_{\text{cons}}(A) = O\left(\frac{\log n}{\log k}\right)$ holds w.h.p.*

From Corollary 1.9, best-of- n^ϵ on $G(n, n^{-1/2+\epsilon})$ for any constant $\epsilon \in (0, 1/2)$ reaches consensus in $O(1)$ steps. It is known that Majority on $G(n, Cn^{-1/2})$ satisfies $T_{\text{cons}}(A) \leq 4$ for large constant C and random $A \subseteq V$ with constant probability [6].

For $3 \leq d \leq n/2$, n -vertex random d -regular graph $G_{n,d}$ is connected and $O(1/\sqrt{d})$ -expander w.h.p. [13, 40].

Corollary 1.10. *Consider a best-of- k on an n -vertex random d -regular graph $G_{n,d}$ for an arbitrary constant $k \geq 2$. Then, $G_{n,d}$ w.h.p. satisfies the following:*

(i) Suppose that $d = \Omega(n^{1/2})$ and $d \leq n/2$. Then,

(a) for any $A \subseteq V$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.

(b) for some $A \subseteq V$, $T_{\text{cons}}(A) = \Omega(\log n)$ w.h.p.

(ii) Suppose that $d \geq C$ and $d \leq n/2$ for a constant $C > 0$ depending only on f . Then, for any $A \subseteq V$ satisfying $|\delta(A)| \geq C \max\left\{\frac{1}{d}, \sqrt{\frac{\log n}{n}}\right\}$, it holds w.h.p. that $T_{\text{cons}}(A) = O\left(\log \log n + \log |\delta(A)|^{-1} + \frac{\log n}{\log d}\right)$.

Corollary 1.11. Let $k = k(n)$ be such that $k = \omega(1)$ and $k = O(\sqrt{n})$. Consider best-of- $(2k+1)$ on an n -vertex random d -regular graph $G_{n,d}$ such that $d = \Omega(k\sqrt{n})$ and $d \leq n/2$. Then, for any $A \subseteq V$, $T_{\text{cons}}(A) = O\left(\frac{\log n}{\log k}\right)$ holds w.h.p.

We can apply Theorems 1.3 and 1.5 if the ratio of the maximum and average degree is constant as follows.

Corollary 1.12. Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Suppose that $d_{\max} \leq C_1 d_{\text{ave}}$ for an arbitrary constant $C_1 > 0$, where d_{\max} and d_{ave} denote the maximum and average degree, respectively. Then, the following holds:

(i) Suppose that $\lambda \leq C_1 n^{-1/4}$. Then

(a) for any $A \subseteq V$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.

(b) for some $A \subseteq V$, $T_{\text{cons}}(A) = \Omega(\log n)$ w.h.p.

(ii) Suppose that $\lambda \leq C_2$ for some constant $C_2 > 0$ depending only on f . Then, for any $A \subseteq V$ satisfying $|\delta(A)| \geq C_2 \max\left\{\lambda^2, \sqrt{\frac{\log n}{n}}\right\}$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.

(iii) In addition to the same assumption as (ii), suppose that $H'_f(0) = 0$. Then, it holds w.h.p. that $T_{\text{cons}}(A) = O\left(\log \log n + \log |\delta(A)|^{-1} + \frac{\log n}{\log \lambda^{-1}}\right)$.

Corollary 1.13. Let $k = k(n)$ be such that $k = \omega(1)$ and $k = o(n/\log n)$. Let C be an arbitrary constant. Consider best-of- $(2k+1)$ on an n -vertex λ -expander graph with degree distribution π such that $\lambda \leq Ck^{-1/2}n^{-1/4}$, and $d_{\max} \leq Cd_{\text{ave}}$, where d_{\max} and d_{ave} denote the maximum and average degree, respectively. Then, $T_{\text{cons}}(A) = O\left(\frac{\log n}{\log k}\right)$ holds w.h.p. for any $A \subseteq V$.

Corollaries 1.12 and 1.13 immediately follow from Theorems 1.3 to 1.5 and 1.7 since $\|\pi\|_2 = O(n^{-1/2})$. Note that if the ratio of the maximum degree d_{\max} and average degree d_{ave} is constant, $\|\pi\|_p = \Theta(1/n^{1-1/p})$ since $\pi(v) = O(1/n)$ for all $v \in V$. We obtain Corollaries 1.8 to 1.11 from Corollaries 1.12 and 1.13.

Other quasi-majority functional voting. We can consider the ρ -lazy variant of a voting process, i.e., every vertex v individually tosses its private coin and operates the voting process with probability ρ , while v does nothing with probability $1 - \rho$. Berenbrink, Giakkoupis, Kermarrec, and Mallmann-Trenn [8] studies 1/2-lazy pull voting. If the original voting process is a quasi-majority functional voting with respect to f , then the corresponding ρ -lazy variant is quasi-majority functional voting with respect to $\rho f : x \mapsto \rho f(x)$. Indeed, $H_{\rho f}(x) = (1 - \rho)x + \rho H_f(x)$.

Corollary 1.14. *Consider a ρ -lazy quasi-majority functional voting on $G(n, p)$ for an arbitrary constant $\rho \in (0, 1]$. Suppose that $p = \Omega(1/\sqrt{n})$. Then, for any $A \subseteq V$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.*

This implies the following interesting observation. In voting processes, the number of neighbor sampling queries per each vertex at each step affects the performance. In pull voting, each vertex communicates with one neighbor but it has a drawback on the slow consensus time. In best-of-two, each vertex communicates with two random neighbors and its consensus time is much faster than that of pull voting. In ρ -lazy best-of-two, each vertex queries 2ρ vertices at each round in expectation, that is less queries than pull voting if $\rho < 1/2$. On the other hand, the consensus time is much faster than pull voting.

Additionally, we can deal with *k-careful voting*. In this model, each vertex v selects k random neighbors (with replacement), and if these sampled k opinions are the same one, v adopts it. Note that one-careful voting and two-careful voting are equivalent to pull voting and best-of-two, respectively. One can check easily that, for any constant $k \geq 2$, this model is a quasi-majority functional voting with respect to $f(x) = x^k$. Note that $H'_f(0) = 0$ and $H'_f(1/2) = 1 + \frac{k-1}{2^{k-1}}$.

Corollary 1.15. *Consider a k -careful voting on $G(n, p)$ for an arbitrary constant $k \geq 2$. Suppose that $p = \Omega(1/\sqrt{n})$. Then, for any $A \subseteq V$, $T_{\text{cons}}(A) = O(\log n)$ w.h.p.*

1.5 Related work

In asynchronous voting process, in each round, a vertex is selected uniformly at random and only the selected vertex updates its opinion. Cooper and Rivera [18] introduced *linear voting model*. In this model, an opinion configuration is represented as a vector $v \in \Sigma^V$ and the vector v updates according to the rule $v \leftarrow Mv$, where M is a random matrix sampled from some probability space. This model captures a wide variety model including asynchronous push/pull voting and synchronous pull voting. Note that best-of-two and best-of-three are not included in linear voting model. Schoenebeck and Yu [37] proposed an asynchronous variant of our functional voting. The authors of [37] proved that, if the function f is symmetric (i.e., $f(1-x) = 1-f(x)$), smooth and has “majority-like” property (i.e., $f(x) > x$ whenever $1/2 < x < 1$), then the expected consensus time is $O(n \log n)$ w.h.p. on $G(n, p)$ with $p = \Omega(1)$. This perspective has also been investigated in physics (see, e.g., [10]).

Several researchers have studied best-of-two and best-of-three on complete graphs initially involving $k \geq 2$ opinions [5, 4, 7, 26]. For example, the consensus time of best-of-three is $O(k \log n)$ if $k = O(n^{1/3}/\sqrt{\log n})$ [26]. Cooper, Radzik, Rivera, and Shiraga [17] considered best-of-two and best-of-three on regular expander graphs that hold more than two opinions.

Recently, Cruciani, Natale, and Scornavacca [20] studied best-of-two with a random initial configuration on a clustered regular graph. Shimizu and Shiraga [39] obtained phase-transition results of best-of-two and best-of-three on stochastic block models.

2 Preliminary and technical result

2.1 Formal definition

Let $G = (V, E)$ be an undirected and connected graph. Let $P \in [0, 1]^{V \times V}$ be the matrix defined as

$$P(u, v) := \frac{\mathbb{1}_{\{u, v\} \in E}}{\deg(u)} \quad \forall (u, v) \in V \times V \quad (4)$$

where $\mathbb{1}_Z$ denotes the indicator of an event Z . For $v \in V$ and $S \subseteq V$, we write $P(v, S) = \sum_{s \in S} P(v, s)$.

Now, let us describe the formal definition of functional voting. For a given $A \subseteq V$, let $(X_v)_{v \in V}$ be independent binary random variables defined as

$$\begin{aligned} \Pr[X_v = 1] &= f(P(v, A)) \quad \text{if } v \in B, \\ \Pr[X_v = 0] &= f(P(v, B)) \quad \text{if } v \in A, \end{aligned} \tag{5}$$

where $B = V \setminus A$. For $A \subseteq V$ and (X_v) above, define $A' = \{v \in V : X_v = 1\}$. Note that this definition coincides with Definition 1.1 since $P(v, A) = \frac{\deg_A(v)}{\deg(v)}$. Then, a functional voting is a Markov chain A_0, A_1, \dots where $A_{t+1} = (A_t)'$.

For $A \subseteq V$, let $T_{\text{cons}}(A)$ denote the consensus time of the functional voting starting from the initial configuration A . Formally, $T_{\text{cons}}(A)$ is the stopping time defined as

$$T_{\text{cons}}(A) := \min \{t \geq 0 : A_t \in \{\emptyset, V\}, A_0 = A\}.$$

2.2 Technical background

Consider best-of-two on a complete graph K_n (with self loop on each vertex) with a current configuration $A \subseteq V$. Let $\alpha = |A|/n$. We have $P(v, A) = \alpha$ for any $v \in V$ and $A \subseteq V$. Then, for any $A \subseteq V$, $\mathbf{E}[\alpha'] = H_f(\alpha) = 3\alpha^2 - 2\alpha^3$. Thus, in each round, $\alpha' = 3\alpha^2 - 2\alpha^3 \pm O(\sqrt{\log n/n})$ holds w.h.p. from the Hoeffding bound. Therefore, the behavior of α can be written as the iteration of applying H_f .

The most technical part is the symmetry breaking at $\alpha = 1/2$. Note that $H_f(1/2) = 1/2$ and thus, the argument above does not work in the case of $|\alpha - 1/2| = o(\sqrt{\log n/n})$. To analyze this case, the authors of [21, 11] proved the following technical lemma asserting that α w.h.p. escapes from the area in $O(\log n)$ rounds.

Lemma 2.1 (Lemma 4.5 of [11] (informal)). *For any constant C , it holds w.h.p. that $|\alpha - 1/2| \geq C\sqrt{\log n/n}$ in $O(\log n)$ rounds (the hidden constant factor depends on C) if*

- (i) *For any constant h , there is a constant $C_0 > 0$ such that, if $|\alpha - 1/2| = O(\sqrt{\log n/n})$ then $\Pr[|\alpha' - 1/2| > h/\sqrt{n}] > C_0$.*
- (ii) *If $|\alpha - 1/2| = O(\sqrt{\log n/n})$ and $|\alpha - 1/2| = \Omega(1/\sqrt{n})$, $\Pr[|\alpha' - 1/2| \leq (1 + \epsilon)|\alpha - 1/2|] \leq \exp(-\Theta((\alpha - 1/2)^2 n))$ for some constant $\epsilon > 0$.*

Intuitively speaking, the condition (ii) means that the bias $|\alpha' - 1/2|$ is likely to be at least $(1 + \epsilon)|\alpha - 1/2|$ for some constant $\epsilon > 0$. The condition (ii) is easy to check using the Hoeffding bound. The condition (i) means that α' has a fluctuation of size $\Omega(1/\sqrt{n})$ with a constant probability. We can check condition (i) using the Central Limit Theorem (the Berry-Esseen bound, see Lemma A.5). The Central Limit Theorem implies that the normalized random variable $(\alpha' - \mathbf{E}[\alpha'])/\sqrt{\mathbf{Var}[\alpha']}$ converges to the standard normal distribution as $n \rightarrow \infty$. In other words, α' has a fluctuation of size $\Theta(\sqrt{\mathbf{Var}[\alpha']})$ with constant probability. Now, to verify the condition (i), we evaluate $\mathbf{Var}[\alpha']$. On K_n , it is easy to show that $\mathbf{Var}[\alpha'] = \Theta(1/n)$, which implies the condition (i).

The authors of [16, 17] considered best-of-two on expander graphs. They focused on the behavior of $\pi(A)$ instead of α . Roughly speaking, they proved that $\mathbf{E}[\pi(A')] - 1/2 \geq (1 + \epsilon)(\pi(A) - 1/2) - O(\lambda^2)$. At the heart of the proof, they showed the following result.

Lemma 2.2 (Special case of Lemma 3 of [17]). *Consider a λ -expander graph with degree distribution π . Then, for any $S \subseteq V$,*

$$\left| \sum_{v \in V} \pi(v) P(v, S)^2 - \pi(S)^2 \right| \leq \lambda^2 \pi(S) (1 - \pi(S)).$$

Then, from the Hoeffding bound, we have $\mathbf{E}[\pi(A') - 1/2] \geq (1 + \epsilon)(\pi(A) - 1/2) - O(\lambda^2 + \|\pi\|_2 \sqrt{\log n})$. Thus, if the initial bias $|\pi(A) - 1/2|$ is $\Omega(\max\{\lambda^2, \sqrt{\log n/n}\})$, we can show that the consensus time is $O(\log n)$.

Unfortunately, we can not apply the same technique to estimate $\mathbf{Var}[\pi(A')]$ on expander graphs, and due to this reason, it seems difficult to estimate the worst-case consensus time on expander graphs. Actually, any previous works put assumptions on the initial bias due to the same reason. It should be noted that Lemma 2.1 is well-known in the literature. For example, Cruciani et al. [20] used Lemma 2.1 from random initial configurations.

The technique of estimating $\mathbf{E}[\pi(A')]$ by Cooper et al. [16, 17] is specialized in best-of-two. Thus, it is not straightforward to prove the estimation of $\mathbf{E}[\pi(A')]$ for voting processes other than best-of-two.

2.3 Our technical contribution

For simplicity, in this part, we focus on a quasi-majority functional voting with respect to a *symmetric* function f (i.e., $f(1 - x) = 1 - f(x)$ for every $x \in [0, 1]$) on a λ -expander graph with degree distribution π . For example, $f(x) = 3x^2 - 2x^3$ of best-of-three is a symmetric function. Note that $f = H_f$ if f is symmetric. Similar results mentioned in this subsection holds for non-symmetric f (see Section 3.3). For a C^2 function $h : \mathbb{R} \rightarrow \mathbb{R}$, let

$$K_1(h) := \max_{x \in [0, 1]} |h'(x)|, \quad K_2(h) := \max_{x \in [0, 1]} |h''(x)|$$

be some constants⁶ depending only on h . The following technical result enables us to estimate $\mathbf{E}[\pi(A')]$ and $\mathbf{Var}[\pi(A')]$ of functional voting.

Lemma 2.3. *Consider a functional voting with respect to a symmetric C^2 function f on a λ -expander graph with degree distribution π . Let $g(x) := f(x)(1 - f(x))$. Then, for all $A \subseteq V$,*

$$\begin{aligned} |\mathbf{E}[\pi(A')] - H_f(\pi(A))| &\leq \frac{K_2(f)}{2} \lambda^2 \pi(A) (1 - \pi(A)), \\ |\mathbf{Var}[\pi(A')] - \|\pi\|_2^2 g(\pi(A))| &\leq K_1(g) \lambda \sqrt{\pi(A) (1 - \pi(A))} \|\pi\|_3^{3/2}. \end{aligned}$$

Note that, if f is symmetric, the corresponding functional voting satisfies that $\mathbf{Pr}[v \in A'] = f(P(v, A))$ for any $v \in V$. Thus we have

$$\mathbf{E}[\pi(A')] = \sum_{v \in V} \pi(v) f(P(v, A)), \quad \mathbf{Var}[\pi(A')] = \sum_{v \in V} \pi(v)^2 g(P(v, A)).$$

To evaluate $\mathbf{E}[\pi(A')]$ and $\mathbf{Var}[\pi(A')]$ above, we prove the following key lemma that is a generalization of Lemma 2.2 and implies Lemma 2.3.

⁶For example, for $f(x) = 3x^2 - 2x^3$ of best-of-three, $f''(x) = 6 - 12x$ and $K_2(f) = 6$. It should be noted that we deal with f not depending on G except for best-of- k with $k = \omega(1)$ in Section 6.

Lemma 2.4 (Special case of Lemmas 3.2 and 3.3). *Consider a λ -expander graph with degree distribution π . Then, for any $S \subseteq V$ and any C^2 function $h : \mathbb{R} \rightarrow \mathbb{R}$,*

$$\left| \sum_{v \in V} \pi(v) h(P(v, S)) - h(\pi(S)) \right| \leq \frac{K_2(h)}{2} \lambda^2 \pi(S) (1 - \pi(S)),$$

$$\left| \sum_{v \in V} \pi(v)^2 h(P(v, S)) - \|\pi\|_2^2 h(\pi(S)) \right| \leq K_1(h) \lambda \sqrt{\pi(S) (1 - \pi(S))} \|\pi\|_3^{3/2}.$$

2.4 Proof sketch of Theorem 1.3

We present proof sketch of Theorem 1.3(i). From the assumption of Theorem 1.3(i) and Lemma 2.3, if $|\pi(A) - 1/2| = o(1)$, we have $\mathbf{Var}[\pi(A')] = \Theta(\|\pi\|_2^2 g(\pi(A))) = \Theta(\|\pi\|_2^2 g(1/2 + o(1))) = \Theta(1/n)$. Moreover, $\mathbf{E}[\pi(A')] = H_f(\pi(A)) \pm O(\pi(A)/\sqrt{n})$ holds for any $A \subseteq V$. Hence, from the Hoeffding bound, $\pi(A') = H_f(\pi(A)) + O(\sqrt{\log n/n})$ holds w.h.p. for any $A \subseteq V$.

- If $|\pi(A) - 1/2| = O(\sqrt{\log n/n})$, we use Lemma 2.1 to obtain an $O(\log n)$ round symmetry breaking. In this phase, since $|\pi(A) - 1/2| = o(1)$, $\mathbf{Var}[\pi(A') - 1/2] = \Theta(1/n)$. Then, from the Berry-Esseen theorem (Lemma A.5), we can check the condition (i). To check the condition (ii), we invoke the condition $H'_f(1/2) > 1$ of the quasi-majority function. From Taylor's theorem and the assumption of Lemma 2.1(ii) ($\pi(A) - 1/2 = \Omega(1/\sqrt{n})$), $\mathbf{E}[\pi(A') - 1/2] = H_f(\pi(A)) - H_f(1/2) - O(1/\sqrt{n}) \approx (1 + \epsilon_1)(\pi(A) - 1/2)$ for some positive constant $\epsilon_1 > 0$. Note that $H_f(1/2) = 1/2$.
- If $C_1 \sqrt{\log n/n} \leq |\pi(A) - 1/2| \leq C_2$ for sufficiently large constant C_1 and some constant $C_2 > 0$, we use the Hoeffding bound and then obtain $\pi(A') - 1/2 \approx (1 + \epsilon_1)(\pi(A) - 1/2) - O(\sqrt{\log n/n}) \geq (1 + (\epsilon_1/2))(\pi(A) - 1/2)$ w.h.p. Hence, $O(\log n)$ rounds suffice to yield a constant bias. (Note that this argument holds when $|\pi(A) - 1/2| \leq C_2$ due to the remainder term of Taylor's theorem.)
- If $C_3 \leq \pi(A) < 1/2$, it is straightforward to see that $\pi(A') = H_f(\pi(A)) + O(\sqrt{\log n/n}) \leq \pi(A) - \epsilon_2$ w.h.p. for some constant $\epsilon_2 > 0$. Note that we invoke the property that $H_f(x) < x$ whenever $0 < x < 1/2$.
- If $\pi(A) \leq C_3$ for sufficiently small constant C_3 , we use the Markov inequality to show $\pi(A_t) = O(n^{-3})$ w.h.p. for some $t = O(\log n)$. Since $\pi(A) \geq 1/n^2$ whenever $A \neq \emptyset$, this implies that the consensus time is $O(\log n)$ w.h.p. Note that, since $H'_f(0) < 1$, we have $\mathbf{E}[\pi(A')] \leq H_f(\pi(A)) + O(\pi(A)/\sqrt{n}) \approx H'_f(0)\pi(A) + O(\pi(A)/\sqrt{n}) \leq (1 - \epsilon_3)\pi(A)$ for some constant $\epsilon_3 > 0$.

In the proof of Theorem 1.7, we modify Lemma 2.1 and apply the same argument.

3 Estimation of $\mathbf{E}[\pi(A')]$ and $\mathbf{Var}[\pi(A')]$

In this section, we prove Lemma 2.4 by showing Lemmas 3.2 and 3.3, which are generalizations of Lemma 2.4 in terms of *reversible Markov chain*. This enables us to evaluate $\mathbf{E}[\pi(A')]$ and $\mathbf{Var}[\pi(A')]$ for functional voting with respect to a C^2 function f (see Section 3.3 for functional voting with respect to non-symmetric f).

3.1 Technical tools for reversible Markov chains

To begin with, we briefly summarize the notation of Markov chain, which we will use in this section⁷. Let V be a set of size n . A *transition matrix* P over V is a matrix $P \in [0, 1]^{V \times V}$ satisfying $\sum_{v \in V} P(u, v) = 1$ for any $u \in V$. Let $\pi \in [0, 1]^V$ denote the *stationary distribution* of P , i.e., a probability distribution satisfying $\pi P = \pi$. A transition matrix P is *reversible* if $\pi(u)P(u, v) = \pi(v)P(v, u)$ for any $u, v \in V$. It is easy to check that the matrix (4) is a reversible transition matrix and its stationary distribution is (3). Let $\lambda_1 \geq \dots \geq \lambda_n$ denote the eigenvalues of P . If P is reversible, it is known that $\lambda_i \in \mathbb{R}$ for all i . Let $\lambda = \max\{|\lambda_2|, |\lambda_n|\}$ be the second largest eigenvalue in absolute value⁸.

For a function $h : \mathbb{R} \rightarrow \mathbb{R}$ and subsets $S, T \subseteq V$, consider the quantity $Q_h(S, T)$ defined as

$$Q_h(S, T) := \sum_{v \in S} \pi(v) h(P(v, T)). \quad (6)$$

The special case of $h(x) = x$, that is, $Q(S, T) := \sum_{v \in S} \pi(v) P(v, T)$, is well known as *edge measure* [30] or *ergodic flow* [3, 32]. Note that, for any reversible P and subsets $S, T \subseteq V$, $Q(S, T) = Q(T, S)$ holds. The following result is well known as a version of the *expander mixing lemma*.

Lemma 3.1 (See, e.g., p.163 of [30]). *Suppose P is reversible. Then, for any $S, T \subseteq V$,*

$$|Q(S, T) - \pi(S)\pi(T)| \leq \lambda \sqrt{\pi(S)\pi(T)(1 - \pi(S))(1 - \pi(T))}.$$

We show the following lemma which gives a useful estimation of $Q_h(S, T)$.

Lemma 3.2. *Suppose P is reversible. Then, for any $S, T \subseteq V$ and any C^2 function $h : \mathbb{R} \rightarrow \mathbb{R}$,*

$$\left| Q_h(S, T) - \pi(S)h(\pi(T)) - h'(\pi(T))(Q(S, T) - \pi(S)\pi(T)) \right| \leq \frac{K_2(h)}{2} \lambda^2 \pi(T)(1 - \pi(T)).$$

Proof of Lemma 3.2. From Taylor's theorem, it holds for any $x, y \in [0, 1]$ that

$$|h(x) - h(y) - h'(y)(x - y)| \leq \frac{K_2(h)}{2} (x - y)^2.$$

Hence

$$\begin{aligned} & \left| Q_h(S, T) - \pi(S)h(\pi(T)) - h'(\pi(T))(Q(S, T) - \pi(S)\pi(T)) \right| \\ &= \left| \sum_{v \in S} \pi(v) \left(h(P(v, T)) - h(\pi(T)) - h'(\pi(T))(P(v, T) - \pi(T)) \right) \right| \\ &\leq \sum_{v \in S} \pi(v) \left| h(P(v, T)) - h(\pi(T)) - h'(\pi(T))(P(v, T) - \pi(T)) \right| \\ &\leq \sum_{v \in S} \pi(v) \frac{K_2(h)}{2} (P(v, T) - \pi(T))^2 \leq \frac{K_2(h)}{2} \sum_{v \in V} \pi(v) (P(v, T) - \pi(T))^2 \\ &\leq \frac{K_2(h)}{2} \lambda^2 \pi(T)(1 - \pi(T)). \end{aligned}$$

Note that the last inequality follows from Corollary A.2. □

⁷For further detailed arguments about reversible Markov chains, see e.g., [30].

⁸If P is *ergodic*, i.e., for any $u, v \in V$, there exists a $t > 0$ such that $P^t(u, v) > 0$ and $\text{GCD}\{t > 0 : P^t(x, x) > 0\} = 1$, $1 > \lambda_2$ and $\lambda_n > -1$. For example, the transition matrix of the simple random walk on a connected and non-bipartite graph is ergodic.

Next, consider

$$R_h(S, T) := \sum_{v \in S} \pi(v)^2 h(P(v, T)) \quad (7)$$

for a function $h : \mathbb{R} \rightarrow \mathbb{R}$ and $S, T \subseteq V$. For notational convenience, for $S \subseteq V$, let $\pi_2(S) := \sum_{v \in S} \pi(v)^2$. We show the following lemma that evaluates $R_h(S, T)$.

Lemma 3.3. *Suppose that P is reversible. Then, for any $S, T \subseteq V$ and any C^2 function $h : \mathbb{R} \rightarrow \mathbb{R}$,*

$$|R_h(S, T) - \pi_2(S)h(\pi(T))| \leq K_1(h) \|\pi\|_3^{3/2} \lambda \sqrt{\pi(T)(1 - \pi(T))}.$$

Proof. We first observe that

$$|h(x) - h(y)| \leq K_1(h)|x - y| \quad (8)$$

holds for any $x, y \in [0, 1]$ from Taylor's theorem. Hence,

$$\begin{aligned} & \left| R_h(S, T) - \pi_2(S)h(\pi(T)) \right| \\ &= \left| \sum_{v \in S} \pi(v)^2 (h(P(v, T)) - h(\pi(T))) \right| \leq \sum_{v \in S} \pi(v)^2 |h(P(v, T)) - h(\pi(T))| \\ &\leq \sum_{v \in S} \pi(v)^2 K_1(h) |P(v, T) - \pi(T)| \leq K_1(h) \sum_{v \in V} \pi(v)^2 |P(v, T) - \pi(T)|. \end{aligned}$$

Then, applying the Cauchy-Schwarz inequality and Corollary A.2,

$$\begin{aligned} \sum_{v \in V} \pi(v)^2 |P(v, T) - \pi(T)| &\leq \sqrt{\left(\sum_{v \in V} \pi(v)^3 \right) \left(\sum_{v \in V} \pi(v) (P(v, T) - \pi(T))^2 \right)} \\ &\leq \|\pi\|_3^{3/2} \lambda \sqrt{\pi(T)(1 - \pi(T))} \end{aligned}$$

and we obtain the claim. \square

Remark 3.4. *The results of this paper can be extended to voting processes where the sampling probability is determined by a reversible transition matrix P . This includes voting processes on edge-weighted graphs $G = (V, E, w)$, where $w : E \rightarrow \mathbb{R}$ denotes an edge weight function. Consider the transition matrix P defined as follows: $P(u, v) = w(\{u, v\}) / \sum_{x: \{u, x\} \in E} w(\{u, x\})$ for $\{u, v\} \in E$ and $P(u, v) = 0$ for $\{u, v\} \notin E$. A weighted functional voting with respect to f is determined by $\mathbf{Pr}[v \in A' | v \in B] = f(P(v, B))$ and $\mathbf{Pr}[v \in B' | v \in A] = f(P(v, A))$. For simplicity, in this paper, we do not explore the weighted variant and focus on the usual setting where P is the matrix (4) and its stationary distribution π is (3).*

3.2 Proof of Lemma 2.4

For the first inequality, by substituting V to S of Lemma 3.2, we obtain

$$\left| Q_h(V, T) - h(\pi(T)) \right| \leq \frac{K_2(h)}{2} \lambda^2 \pi(T)(1 - \pi(T)).$$

Note that $Q(V, T) = Q(T, V) = \pi(T)$ from the reversibility of P . Similarly, we obtain the second inequality by substituting V to S of Lemma 3.3. \square

3.3 Non-symmetric functions

This section is devoted to evaluate $\mathbf{E}[\pi(A')]$ and $\mathbf{Var}[\pi(A')]$ for non-symmetric f . To be more specifically, we prove the following.

Lemma 3.5. *Consider a functional voting with respect to a C^2 function f on a λ -expander graph. Then, for all $A \subseteq V$,*

$$|\mathbf{E}[\pi(A')] - H_f(\pi(A))| \leq K_2(f)\lambda(|2\pi(A) - 1| + \lambda)\pi(A)(1 - \pi(A)).$$

Lemma 3.6. *Consider a functional voting with respect to a C^2 function f on a λ -expander graph. Let $g(x) := f(x)(1 - f(x))$. Then, for all $A \subseteq V$,*

$$\left| \mathbf{Var}[\pi(A')] - \|\pi\|_2^2 g\left(\frac{1}{2}\right) \right| \leq K_1(g) \left(\frac{1}{2} \|\pi\|_2^2 |2\pi(A) - 1| + 2\|\pi\|_3^{3/2} \lambda \sqrt{\pi(A)(1 - \pi(A))} \right).$$

Recall that we use $B = V \setminus A$ for $A \subseteq V$. Then, it is clear that

$$\mathbf{E}[\pi(A')] = \pi(A) - \sum_{v \in A} \pi(v)f(P(v, B)) + \sum_{v \in B} \pi(v)f(P(v, A)), \quad (9)$$

$$\mathbf{Var}[\pi(A')] = \sum_{v \in A} \pi(v)^2 g(P(v, B)) + \sum_{v \in B} \pi(v)g(P(v, A)). \quad (10)$$

Proof of Lemma 3.5. From Definition 1.1, (6) and (9), we have

$$\mathbf{E}[\pi(A')] = \pi(A) - Q_f(A, B) + Q_f(B, A), \quad (11)$$

$$H_f(\pi(A)) = \pi(A) - \pi(A)f(\pi(B)) + \pi(B)f(\pi(A)). \quad (12)$$

For notational convenience, for $S, T \subseteq V$, let

$$\begin{aligned} \Delta_f(S, T) &:= Q_f(S, T) - \pi(S)f(\pi(T)) - f'(\pi(T))(Q(S, T) - \pi(S)\pi(T)) \\ &= Q_h(S, T) - \pi(S)f(\pi(T)) - f'(\pi(T))(Q(T, S) - \pi(T)\pi(S)). \end{aligned}$$

The equality follows from the reversibility of P (see Section 3). From Lemma 3.2, we have

$$|\Delta_f(S, T)| \leq \frac{K_2(f)}{2} \lambda^2 \pi(T)(1 - \pi(T)).$$

Then, combining (11) and (12), we have

$$\begin{aligned} & \left| \mathbf{E}[\pi(A')] - H_f(\pi(A)) \right| \\ &= \left| Q_f(B, A) - \pi(B)f(\pi(A)) - Q_f(A, B) + \pi(A)f(\pi(B)) \right| \\ &= \left| \Delta_f(B, A) + f'(\pi(A))(Q(A, B) - \pi(A)\pi(B)) \right. \\ & \quad \left. - \Delta_f(A, B) - f'(\pi(B))(Q(A, B) - \pi(A)\pi(B)) \right| \\ &\leq |\Delta_f(B, A)| + |\Delta_f(A, B)| + \left| f'(\pi(A)) - f'(\pi(B)) \right| |Q(A, B) - \pi(A)\pi(B)| \\ &\leq K_2(f)\lambda^2 \pi(A)\pi(B) + K_2(f)|\pi(A) - \pi(B)|\lambda\pi(A)\pi(B). \end{aligned}$$

and we obtain the claim. Note that the last inequality follows from Taylor's theorem (8) and Lemma 3.1. \square

Proof of Lemma 3.6. From (7) and (10),

$$\mathbf{Var}[\pi(A')] = R_g(A, B) + R_g(B, A).$$

Thus, applying Lemma 3.3 yields

$$\left| \mathbf{Var}[\pi(A')] - \left(\pi_2(A)g(\pi(B)) + \pi_2(B)g(\pi(A)) \right) \right| \leq 2K_1(g)\|\pi\|_3^{3/2}\lambda\sqrt{\pi(A)\pi(B)}. \quad (13)$$

Next, using Taylor's theorem (8),

$$\begin{aligned} & \left| \pi_2(A)g(\pi(B)) + \pi_2(B)g(\pi(A)) - \|\pi\|_2^2 g\left(\frac{1}{2}\right) \right| \\ &= \left| \pi_2(A) \left(g(\pi(B)) - g\left(\frac{1}{2}\right) \right) + \pi_2(B) \left(g(\pi(A)) - g\left(\frac{1}{2}\right) \right) \right| \\ &\leq K_1(g)\pi_2(A) \left| \pi(B) - \frac{1}{2} \right| + K_1(g)\pi_2(B) \left| \pi(A) - \frac{1}{2} \right| = K_1(g)\|\pi\|_2^2 \left| \pi(A) - \frac{1}{2} \right|. \end{aligned} \quad (14)$$

The last equality follows since $|\pi(A) - 1/2| = |\pi(B) - 1/2|$. Combining (13) and (14), we obtain the claim. \square

4 Proofs of Theorems 1.3 and 1.5

Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Let A_0, A_1, \dots , be the sequence given by the functional voting with initial configuration $A_0 \subseteq V$. Theorems 1.3 and 1.5 follow from the following lemma.

Lemma 4.1. *Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Let $\epsilon_h(f) := H'_f(1/2) - 1$, $\epsilon_c(f) := 1 - H'_f(0)$ and $K(f) := \max\{K_2(f), K_2(H_f)\}$ be three positive constants depending only on f . Then, the following holds:*

- (I) *Let $C_1 > 0$ be an arbitrary constant and $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}$ be an arbitrary function satisfying $\varepsilon(n) \rightarrow 0$ as $n \rightarrow \infty$. Suppose that $\lambda \leq C_1 n^{-1/4}$, $\|\pi\|_2 \leq C_1/\sqrt{n}$ and $\|\pi\|_3 \leq \varepsilon/\sqrt{n}$. Then, for any $A_0 \subseteq V$ such that $|\delta(A_0)| \leq c_1 \log n/\sqrt{n}$ for an arbitrary constant $c_1 > 0$, $|\delta(A_t)| \geq c_1 \log n/\sqrt{n}$ within $t = O(\log n)$ steps w.h.p.*
- (II) *Suppose that $\lambda \leq \frac{\epsilon_h(f)}{2K(f)}$. Then, for any $A_0 \subseteq V$ s.t. $\frac{2\max\{K(f), 8\}}{\epsilon_h(f)} \max\{\lambda^2, \|\pi\|_2\sqrt{\log n}\} \leq |\delta(A_0)| \leq \frac{\epsilon_h(f)}{K(f)}$, $|\delta(A_t)| \geq \frac{\epsilon_h(f)}{K(f)}$ within $t = O(\log |\delta(A_0)|^{-1})$ steps w.h.p.*
- (III) *Let c_2, c_3 be two arbitrary constants satisfying $0 < c_2 < c_3 < 1/2$ and $\epsilon(f) := \min_{x \in [c_2, c_3]} (x - H_f(x))$ be a positive constant depending f, c_2, c_3 . Suppose that $\lambda \leq \frac{\epsilon(f)}{2K(f)}$ and $\|\pi\|_2 \leq \frac{\epsilon(f)}{4\sqrt{\log n}}$. Then, for any $A_0 \subseteq V$ satisfying $c_2 \leq \pi(A_0) \leq c_3$, $\pi(A_t) \leq c_2$ within constant steps w.h.p.*
- (IV) *Suppose that $\lambda \leq \frac{\epsilon_c(f)}{2K(f)}$ and $\|\pi\|_2 \leq \frac{\epsilon_c(f)^2}{32K(f)\sqrt{\log n}}$. Then, for any $A_0 \subseteq V$ satisfying $\pi(A_0) \leq \frac{\epsilon_c(f)}{8K(f)}$, $\pi(A_t) = 0$ within $t = O(\log n)$ steps w.h.p.*
- (V) *Suppose that $H'_f(0) = 0$, $\lambda \leq \frac{1}{10K(f)}$ and $\|\pi\|_2 \leq \frac{1}{64K(f)\sqrt{\log n}}$. Then, for any $A_0 \subseteq V$ satisfying $\pi(A_0) \leq \frac{1}{7K(f)}$, it holds w.h.p. that $\pi(A_t) = 0$ within*

$$t = O\left(\log \log n + \frac{\log n}{\log \lambda^{-1}} + \frac{\log n}{\log(\|\pi\|_2\sqrt{\log n})^{-1}}\right) \text{ steps.}$$

Proof of Theorem 1.3(ii). Since $\|\pi\|_2 \geq 1/\sqrt{n}$, we have $|\delta(A_0)| = \Omega(\sqrt{\log n/n})$. This implies that Phase (II) takes at most $O(\log n)$. Thus, we obtain the claim since we can merge Phases (II) to (IV) by taking appropriate constants c_2, c_3 in Phase (III). \square

Proof of Theorem 1.3(i). Under the assumption of Theorem 1.3(i), for any positive constant C , a positive constant C' exists such that $C(\lambda^2 + \|\pi\|_2 \sqrt{\log n}) \leq C' \frac{\log n}{\sqrt{n}}$. Thus, we can combine Phase (I) and Theorem 1.3(ii), and we obtain the claim. \square

Proof of Theorem 1.5. Combining Phases (II), (III) and (V), we obtain the claim. \square

4.1 Proof of Lemma 4.1

For notational convenience, let

$$\begin{aligned}\alpha &:= \pi(A), \alpha' := \pi(A'), \alpha_t := \pi(A_t), \\ \delta &:= \delta(A) = 2\alpha - 1, \delta' := \delta(A'), \delta_t := \delta(A_t).\end{aligned}$$

4.2 Phase (I): $0 \leq |\delta| \leq c_1 \log n / \sqrt{n}$

We use the following lemma to show Lemma 4.1(I).

Lemma 4.2 (Lemma 4.5 of [11]). *Consider a Markov chain $(X_t)_{t=1}^\infty$ with finite state space Ω and a function $\Psi : \Omega \rightarrow \{0, \dots, n\}$. Let C_3 be arbitrary constant and $m = C_3 \sqrt{n} \log n$. Suppose that Ω, Ψ and m satisfies the following conditions:*

(i) *For any positive constant h , there exists a positive constant $C_1 < 1$ such that*

$$\mathbf{Pr} [\Psi(X_{t+1}) < h\sqrt{n} \mid \Psi(X_t) \leq m] < C_1.$$

(ii) *Three positive constants γ, C_2 and h exist such that, for any $x \in \Omega$ satisfying $h\sqrt{n} \leq \Psi(x) < m$,*

$$\mathbf{Pr} [\Psi(X_{t+1}) < (1 + \gamma)\Psi(X_t) \mid X_t = x] < \exp \left(-C_2 \frac{\Psi(x)^2}{n} \right).$$

Then, $\Psi(X_t) \geq m$ holds w.h.p. for some $t = O(\log n)$.

Let us first prove the following lemma concerning the growth rate of $|\delta|$, which we will use in the proofs of (I) and (II) of Lemma 4.1.

Lemma 4.3. *Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Let $\epsilon_h(f) := H'_f(1/2) - 1$ and $K(f) := \max\{K_2(f), K_2(H_f)\}$ be positive constants depending only on f . Suppose that $\lambda \leq \frac{\epsilon_h(f)}{2K(f)}$. Then, for any $A \subseteq V$ satisfying $\frac{2K(f)}{\epsilon_h(f)} \lambda^2 \leq |\delta| \leq \frac{\epsilon_h(f)}{K(f)}$,*

$$\mathbf{Pr} \left[|\delta'| \leq \left(1 + \frac{\epsilon_h(f)}{8} \right) |\delta| \right] \leq 2 \exp \left(-\frac{\epsilon_h(f)^2 \delta^2}{128 \|\pi\|_2^2} \right).$$

Proof. Combining Lemma 3.5 and Taylor's theorem, we have

$$\begin{aligned}
\left| \mathbf{E}[\delta'] - H_f' \left(\frac{1}{2} \right) \delta \right| &= 2 \left| \mathbf{E}[\alpha'] - \frac{1}{2} - H_f' \left(\frac{1}{2} \right) \left(\alpha - \frac{1}{2} \right) \right| \\
&= 2 \left| \mathbf{E}[\alpha'] - H_f(\alpha) + H_f(\alpha) - H_f \left(\frac{1}{2} \right) - H_f' \left(\frac{1}{2} \right) \left(\alpha - \frac{1}{2} \right) \right| \\
&\leq 2K_2(f)\lambda(|\delta| + \lambda)\alpha(1 - \alpha) + K_2(H_f) \left(\alpha - \frac{1}{2} \right)^2 \\
&\leq \left(\frac{K(f)}{2}\lambda + \frac{K(f)}{4}|\delta| \right) |\delta| + \frac{K(f)}{2}\lambda^2
\end{aligned} \tag{15}$$

Note that $H_f(1/2) = 1/2$ from the definition. From assumptions of $\lambda \leq \frac{\epsilon_h(f)}{2K(f)}$, $|\delta| \leq \frac{\epsilon_h(f)}{K(f)}$ and $\lambda^2 \leq \frac{\epsilon_h(f)}{2K(f)}|\delta|$, we have

$$\left| H_f' \left(\frac{1}{2} \right) \delta - |\mathbf{E}[\delta']| \right| \leq \left| H_f' \left(\frac{1}{2} \right) \delta - \mathbf{E}[\delta'] \right| \leq \frac{3}{4}\epsilon_h(f)|\delta|.$$

Hence, it holds that

$$|\mathbf{E}[\delta']| \geq \left| H_f' \left(\frac{1}{2} \right) \delta - \frac{3}{4}\epsilon_h(f)|\delta| \right| = (1 + \epsilon_h(f))|\delta| - \frac{3}{4}\epsilon_h(f)|\delta| = \left(1 + \frac{\epsilon_h(f)}{4} \right) |\delta|.$$

We observe that, for any $\kappa > 0$,

$$\mathbf{Pr} \left[|\delta'| \leq |\mathbf{E}[\delta']| - \kappa \right] \leq 2 \exp \left(-\frac{\kappa^2}{2\|\pi\|_2^2} \right) \tag{16}$$

from Corollary A.4. Note that $\delta' = \sum_{v \in V} \pi(v)(2X_v - 1)$ for independent indicator random variables $(X_v)_{v \in V}$ (see (5) for the definition of X_v). Thus,

$$\begin{aligned}
\mathbf{Pr} \left[|\delta'| \leq \left(1 + \frac{\epsilon_h(f)}{8} \right) |\delta| \right] &= \mathbf{Pr} \left[|\delta'| \leq \left(1 + \frac{\epsilon_h(f)}{4} \right) |\delta| - \frac{\epsilon_h(f)}{8} |\delta| \right] \\
&\leq \mathbf{Pr} \left[|\delta'| \leq |\mathbf{E}[\delta']| - \frac{\epsilon_h(f)}{8} |\delta| \right] \leq 2 \exp \left(-\frac{\epsilon_h(f)^2 \delta^2}{128\|\pi\|_2^2} \right)
\end{aligned}$$

and we obtain the claim. \square

Proof of Lemma 4.1(I). We check the conditions (i) and (ii) of Lemma 4.2 with letting $\Psi(A) = \lfloor n|\delta(A)| \rfloor$ and $m = c_1 \sqrt{n} \log n$.

Condition (i). First, we show the following claim that evaluates $\mathbf{Var}[\delta']$.

Claim 4.4. *Under the same assumption as Lemma 4.1(I),*

$$\frac{\epsilon_{\text{var}}(f)}{n} \leq \mathbf{Var}[\delta'] \leq \frac{5C_1^2}{n}$$

where $\epsilon_{\text{var}}(f) := f(1/2)(1 - f(1/2))$ is a positive constant depending only on f .

Proof of the claim. From Lemma 3.6 and assumptions, we have

$$\begin{aligned} \left| \frac{\mathbf{Var}[\delta']}{4} - \|\pi\|_2^2 g \left(\frac{1}{2} \right) \right| &= \left| \mathbf{Var}[\alpha'] - \|\pi\|_2^2 g \left(\frac{1}{2} \right) \right| \leq K_1(g) \left(\|\pi\|_2^2 \frac{|\delta|}{2} + \|\pi\|_3^{3/2} \lambda \right) \\ &\leq \frac{K_1(g)}{n} \left(C_1^2 c_1 \frac{\log n}{\sqrt{n}} + C_1 \epsilon^{3/2} \right) = \frac{1}{n} \cdot o(1). \end{aligned}$$

Note that $\mathbf{Var}[\delta'] = \mathbf{Var}[2\alpha' - 1] = 4 \mathbf{Var}[\alpha']$. Since $\|\pi\|_2^2 \geq 1/n$, we have

$$\frac{\epsilon_{\text{var}}(f)}{n} \leq \frac{4\epsilon_{\text{var}}(f) - o(1)}{n} \leq \mathbf{Var}[\delta'] \leq \frac{4C_1^2 + o(1)}{n} \leq \frac{5C_1^2}{n}.$$

□

From Corollary A.6 with letting $Y_v = \pi(v)(2X_v - 1)$, we have

$$\begin{aligned} \mathbf{Pr} \left[|\delta'| \leq x \sqrt{\frac{\epsilon_{\text{var}}(f)}{n}} \right] &\leq \mathbf{Pr} \left[|\delta'| \leq x \sqrt{\mathbf{Var}[\delta']} \right] \leq \Phi(x) + \frac{5.6 \|\pi\|_3^3}{\mathbf{Var}[\delta']^{3/2}} \\ &\leq \Phi(x) + 5.6 \frac{\epsilon^3}{n^{3/2}} \cdot \frac{n^{3/2}}{\epsilon_{\text{var}}(f)^{3/2}} = \Phi(x) + o(1) \end{aligned} \quad (17)$$

for any $x \in \mathbb{R}$, where $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} dy$. Thus, for any constant $h > 0$, there exists some constant $C > 0$ such that

$$\mathbf{Pr}[\Psi(A') < h\sqrt{n} \mid \Psi(A) \leq m] < C,$$

which verifies the condition (i).

Condition (ii). Set $h = \frac{2K(f)}{\epsilon_h(f)} C_1^2$ and assume $h\sqrt{n} \leq \Psi(A) < m$. Then

$$\frac{2K(f)}{\epsilon_h(f)} \lambda^2 n \leq \frac{2K(f)}{\epsilon_h(f)} C_1^2 \sqrt{n} = h\sqrt{n} \leq \Psi(A) \leq |\delta|n = o(n).$$

Thus, we can apply Lemma 4.3 and positive constants γ, C exist such that, for any $h\sqrt{n} \leq \Psi(A) \leq c_1 \sqrt{n} \log n$,

$$\mathbf{Pr}[\Psi(A') < (1 + \gamma)\Psi(A)] < \exp \left(-C \frac{\Psi(A)^2}{n} \right).$$

Note that $\|\pi\|_2^2 = \Theta(1/n)$ from the assumption. This verifies the condition (ii).

Thus, we can apply Lemma 4.2 and we obtain the claim. □

4.3 Phase (II): $\frac{2 \max\{K(f), 8\}}{\epsilon_h(f)} \max\{\lambda^2, \|\pi\|_2 \sqrt{\log n}\} \leq |\delta| \leq \frac{\epsilon_h(f)}{K(f)}$

Proof of Lemma 4.1(II). Since $|\delta| \geq \frac{16}{\epsilon_h(f)} \|\pi\|_2 \sqrt{\log n}$ from assumptions, applying Lemma 4.3 yields

$$\mathbf{Pr} \left[|\delta'| \leq \left(1 + \frac{\epsilon_h(f)}{8} \right) |\delta| \right] \leq \frac{2}{n^2}.$$

Thus, it holds with probability larger than $(1 - 2/n^2)^t$ that $|\delta_t| \geq \left(1 + \frac{\epsilon_h(f)}{8} \right)^t |\delta_0|$ and we obtain the claim by substituting $t = O(\log |\delta_0|^{-1})$. □

4.4 Phase (III): $0 < c_2 \leq \alpha \leq c_3 < 1/2$

Proof of Lemma 4.1(III). We first observe that, for any $\kappa > 0$,

$$\Pr \left[|\alpha' - \mathbf{E}[\alpha']| \geq \kappa \|\pi\|_2 \sqrt{\log n} \right] \leq 2n^{-2\kappa} \quad (18)$$

from Lemma A.3. Note that $\alpha' = \sum_{v \in V} \pi(v) X_v$ for independent indicator random variables $(X_v)_{v \in V}$. Hence, applying Lemma 3.5 yields

$$|\alpha' - H_f(\alpha)| \leq |\alpha' - \mathbf{E}[\alpha']| + |\mathbf{E}[\alpha'] - H_f(\alpha)| \leq \|\pi\|_2 \sqrt{\log n} + \frac{K_2(f)}{4} (|\delta| + \lambda) \lambda \quad (19)$$

with probability larger than $1 - 2/n^2$. Then, for any $\alpha \in [c_2, c_3]$, it holds with probability larger than $1 - 2/n^2$ that

$$\alpha' \leq H_f(\alpha) + \frac{K(f)}{2} \lambda + \|\pi\|_2 \sqrt{\log n} \leq \alpha - \epsilon(f) + \frac{\epsilon(f)}{4} + \frac{\epsilon(f)}{4} \leq \alpha - \frac{\epsilon(f)}{2}.$$

Thus, for $\alpha_0 \in [c_2, c_3]$, $\alpha_t \leq c_2$ within $t = 2(c_3 - c_2)/\epsilon(f) = O(1)$ steps w.h.p. \square

4.5 Phase (IV): $0 \leq \alpha \leq \frac{\epsilon_c(f)}{8K(f)}$

We show the following lemma which is useful for proving (IV) and (V) of Lemma 4.1.

Lemma 4.5. *Let $\epsilon \in (0, 1]$ be an arbitrary constant. Consider functional voting on an n -vertex connected graph with degree distribution π . Suppose that, for some $\alpha_* \in [0, 1]$ and $K \in [0, 1 - \epsilon]$,*

$$\mathbf{E}[\alpha'] \leq K\alpha$$

for any $A \subseteq V$ satisfying $\alpha \leq \alpha_$ and $\|\pi\|_2 \leq \frac{\epsilon\alpha_*}{2\sqrt{\log n}}$. Then, for any $A_0 \subseteq V$ satisfying $\alpha_0 \leq \alpha_*$, $\alpha_t = 0$ w.h.p. within $O\left(\frac{\log n}{\log K^{-1}}\right)$ steps.*

Proof. For any $\alpha \leq \alpha_*$, from (18) and assumptions of $\mathbf{E}[\alpha'] \leq K\alpha$ and $\|\pi\|_2 \leq \frac{\epsilon\alpha_*}{2\sqrt{\log n}}$, it holds with probability larger than $1 - 2/n^4$ that

$$\alpha' \leq \mathbf{E}[\alpha'] + 2\|\pi\|_2 \sqrt{\log n} \leq K\alpha + \epsilon\alpha_* \leq (1 - \epsilon)\alpha_* + \epsilon\alpha_* = \alpha_*.$$

Thus, for any $\alpha_0 \leq \alpha_*$, we have

$$\begin{aligned} \mathbf{E}[\alpha_t] &= \sum_{x \leq a_*} \mathbf{E}[\alpha_t | \alpha_{t-1} = x] \Pr[\alpha_{t-1} = x] + \sum_{x > a_*} \mathbf{E}[\alpha_t | \alpha_{t-1} = x] \Pr[\alpha_{t-1} = x] \\ &\leq \sum_{x \leq a_*} Kx \Pr[\alpha_{t-1} = x] + \Pr[\alpha_{t-1} > a_*] \leq K \mathbf{E}[\alpha_{t-1}] + \frac{2t}{n^4} \\ &\leq \dots \leq K^t \alpha_0 + \frac{2t^2}{n^4} \leq K^t + \frac{2t^2}{n^4}. \end{aligned}$$

This implies that, $\mathbf{E}[\alpha_t] = O(n^{-3})$ within $t = O\left(\frac{\log n}{\log K^{-1}}\right)$ steps. Let $\pi_{\min} := \min_{v \in V} \pi(v) \geq 1/(2|E|) \geq 1/n^2$. Markov inequality yields

$$\Pr[\alpha_t = 0] = 1 - \Pr[\alpha_t \geq \pi_{\min}] \geq 1 - \frac{\mathbf{E}[\alpha_t]}{\pi_{\min}} = 1 - O(1/n)$$

and we obtain the claim. \square

Proof of Lemma 4.1 of (IV). Combining Lemma 3.5 and Taylor's theorem,

$$\begin{aligned}
|\mathbf{E}[\alpha'] - H'_f(0)\alpha| &= |\mathbf{E}[\alpha'] - H_f(\alpha) + H_f(\alpha) - H_f(0) - H'_f(0)(\alpha - 0)| \\
&\leq K_2(f)\lambda(|\delta| + \lambda)\alpha(1 - \alpha) + \frac{K_2(H_f)}{2}\alpha^2 \\
&\leq 2K(f)\lambda\alpha + \frac{K(f)}{2}\alpha^2.
\end{aligned} \tag{20}$$

Hence, for any $\alpha \leq \frac{\epsilon_c(f)}{8K(f)}$, we have

$$\begin{aligned}
\mathbf{E}[\alpha'] &\leq \left(H'_f(0) + 2K(f)\lambda + \frac{K(f)}{2}\alpha \right) \alpha \\
&\leq \left(1 - \epsilon_c(f) + \frac{\epsilon_c(f)}{4} + \frac{\epsilon_c(f)}{4} \right) \alpha = \left(1 - \frac{\epsilon_c(f)}{2} \right) \alpha.
\end{aligned}$$

Letting $\epsilon = \epsilon_c(f)/2$, $K = 1 - \epsilon_c(f)/2$ and $\alpha_* = \frac{\epsilon_c(f)}{8K(f)}$, from the assumption, $\|\pi\|_2 \leq \frac{\epsilon_c(f)^2}{32K(f)\sqrt{\log n}} = \frac{\epsilon\alpha_*}{2\sqrt{\log n}}$. Thus, we can apply Lemma 4.5 and we obtain the claim. \square

4.6 Phase (V): $H'_f(0) = 0$ and $0 \leq \alpha \leq \frac{1}{7K(f)}$

Proof of Lemma 4.1(V). In this case, from (20),

$$\mathbf{E}[\alpha'] \leq 2K(f)\lambda\alpha + \frac{K(f)}{2}\alpha^2. \tag{21}$$

We consider the following two cases.

Case 1. $\max \left\{ \lambda, \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \right\} \leq \alpha \leq \frac{1}{7K(f)}$: In this case, combining (18) and (21), it holds with probability larger than $1 - 2/n^2$ that

$$\alpha' \leq \left(\frac{2K(f)\lambda}{\alpha} + \frac{K(f)}{2} + \frac{\|\pi\|_2\sqrt{\log n}}{\alpha^2} \right) \alpha^2 \leq \frac{7K(f)}{2}\alpha^2.$$

Applying this inequality iteratively, for any $\alpha_0 \leq 7K(f)^{-1}$,

$$\alpha_t \leq \frac{7K(f)}{2}\alpha_{t-1}^2 \leq \dots \leq \frac{2}{7K(f)} \left(\frac{7K(f)}{2}\alpha_0 \right)^{2^t} \leq \frac{2}{7K(f)2^{2^t}}.$$

holds with probability larger than $(1 - 2/n^2)^t$. This implies that, within $t = O(\log \log n)$ steps,

$\alpha_t \leq \max \left\{ \lambda, \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \right\}$ w.h.p. Note that $\max \left\{ \lambda, \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \right\} \geq \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \geq \sqrt{\frac{\sqrt{\log n/n}}{K(f)}}$ since $\|\pi\|_2^2 \geq 1/n$.

Case 2. $\alpha \leq \max \left\{ \lambda, \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \right\}$: Set $\alpha_* = \max \left\{ \lambda, \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \right\} \geq \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}}$, $K = \frac{5K(f)}{2}\lambda + \frac{1}{2}\sqrt{K(f)\|\pi\|_2\sqrt{\log n}}$ and $\epsilon = 1/4$. Then, from $\lambda \leq \frac{1}{10K(f)}$ and $\|\pi\|_2 \leq \frac{1}{64K(f)\sqrt{\log n}}$,

$$\|\pi\|_2 = (\sqrt{\|\pi\|_2})^2 \leq \frac{\sqrt{\|\pi\|_2}}{8\sqrt{K(f)\sqrt{\log n}}} = \sqrt{\frac{\|\pi\|_2\sqrt{\log n}}{K(f)}} \frac{\epsilon}{2\sqrt{\log n}} \leq \frac{\epsilon\alpha_*}{2\sqrt{\log n}},$$

$$K \leq \frac{1}{2} + \frac{1}{16} \leq 1 - \epsilon,$$

$$\mathbf{E}[\alpha'] \leq \left(2K(f)\lambda + \frac{K(f)}{2}\alpha \right) \alpha \leq \left(2K(f)\lambda + \frac{K(f)}{2}\lambda + \frac{1}{2}\sqrt{K(f)\|\pi\|_2\sqrt{\log n}} \right) \alpha = K\alpha.$$

$K \leq 1/4 + 1/2 = 3/4$. Thus, applying Lemma 4.5, we obtain the claim. \square

5 Proof of Theorem 1.4

This section is devoted to prove Theorem 1.4. In particular, we show the following theorem.

Theorem 5.1. *Let $C > 0$ be an arbitrary constant. Consider a quasi-majority functional voting with respect to f on an n -vertex λ -expander graph with degree distribution π . Suppose that $\max\{\lambda, \|\pi\|_2\} \leq n^{-C}$. Then, for any $A \subseteq V$ satisfying $|\delta(A)| \leq n^{-C}$, $T_{\text{cons}}(A) = \Omega(\log n)$ w.h.p.*

Proof of Theorem 5.1. From (15),

$$\begin{aligned} |\mathbf{E}[\delta']| &\leq H'_f \left(\frac{1}{2} \right) |\delta| + \left(\frac{K(f)}{2} \lambda + \frac{K(f)}{4} |\delta| \right) |\delta| + \frac{K(f)}{2} \lambda^2 \\ &\leq \left(1 + \epsilon_h(f) + \frac{3K(f)}{4} \right) |\delta| + K(f) \lambda^2. \end{aligned}$$

Recall that $\delta' = \sum_{v \in V} (2\pi_v - 1)$ for independent indicator random variables $(X_v)_{v \in V}$ (5). Thus, for any $\kappa > 0$,

$$\Pr[|\delta'| \geq |\mathbf{E}[\delta']| + \kappa] \leq \exp\left(-\frac{\kappa^2}{2\|\pi\|_2^2}\right)$$

from Corollary A.4. Hence, it holds with probability larger than $1 - 2/n^2$ that

$$|\delta'| \leq c|\delta| + K(f)\lambda^2 + 2\|\pi\|_2\sqrt{\log n},$$

where we put $c := 1 + \epsilon_h(f) + \frac{3K(f)}{4} > 1$. Then, applying this inequality iteratively with $t = (C/2) \log_c n$ steps,

$$\begin{aligned} |\delta_t| &\leq c|\delta_{t-1}| + K(f)\lambda^2 + 2\|\pi\|_2\sqrt{\log n} \\ &\leq \dots \leq c^t|\delta_0| + tc^t \left(K(f)\lambda^2 + 2\|\pi\|_2\sqrt{\log n} \right) \\ &\leq \frac{n^{C/2}}{n^C} + n^{C/2} \log_c n^{C/2} \left(\frac{K(f)}{n^{2C}} + \frac{2\sqrt{\log n}}{n^C} \right) = o(1) \end{aligned}$$

w.h.p., and we obtain the claim. Note that we use our assumptions of $|\delta_0|, \max \lambda, \|\pi\|_2 \leq n^{-C}$ in the last inequality. \square

6 Proof of Theorem 1.7

We show Theorem 1.7. The proof is almost same as the one given in Section 4 but we need some special care. We assume $k = \omega(1)$ and thus k is sufficiently large. Consider best-of- $(2k+1)$ on an n -vertex λ -expander graph with degree distribution π . Suppose that the graph satisfies the conditions of Theorem 1.7. Let A_0, A_1, \dots , be the sequence given by the best-of- $(2k+1)$ with initial configuration $A_0 \subseteq V$. For notational convenience, let

$$\begin{aligned} \alpha &:= \pi(A), \alpha' := \pi(A'), \alpha_t := \pi(A_t), \\ \delta &:= \delta(A) = 2\alpha - 1, \delta' := \delta(A'), \delta_t := \delta(A_t). \end{aligned}$$

The dynamics of best-of- $(2k+1)$ are divided into four phases. More specifically, we prove the following key result that corresponds to Lemma 4.1.

Lemma 6.1. Consider best-of- $(2k+1)$ on an n -vertex λ -expander graph with degree distribution π . Suppose that the graph satisfies the conditions of Theorem 1.7. Then, the following holds:

- (I) For any $A_0 \subseteq V$ satisfying $|\delta_0| \leq 300C \log n / \sqrt{n}$, $|\delta_t| \geq 300C \log n / \sqrt{n}$ within $t = O(\log n / \log k)$ steps w.h.p.
- (II) For any $A_0 \subseteq V$ satisfying $|\delta_0|$ satisfying $300C \log n / \sqrt{n} \leq |\delta_0| \leq \frac{1.25}{\sqrt{k}}$, $|\delta_t| > \frac{1.25}{\sqrt{k}}$ within $t = O(\log n / \log k)$ steps w.h.p.
- (III) For any $A_0 \subseteq V$ satisfying $\frac{1.25}{\sqrt{k}} \leq |\delta_0| \leq 0.9$, $|\delta_1| > 0.9$ w.h.p.
- (IV) For any $A_0 \subseteq V$ satisfying $0.9 \leq |\delta_0| < 1$, $|\delta_t| = 1$ (or equivalently, the voting process reaches consensus) within $t = O(\log n / \log k)$ steps w.h.p.

Proof of Theorem 1.7 using Lemma 6.1. Theorem 1.7 is straightforward from Lemma 6.1. For any initial configuration $A_0 \subseteq V$, A_0 satisfies one of (I) to (IV). If A_0 satisfies (IV), the consensus time is $O(\log n / \log k)$. Otherwise, from Lemma 6.1, for some $t = O(\log n / \log k)$, A_t satisfies $|\delta(A_t)| > 0.9$ and then apply Lemma 6.1(IV). \square

The rest of this section is devoted to prove Lemma 6.1. We begin with preparing useful facts concerning with best-of- $(2k+1)$. Let f_{2k+1} be the betrayal function of best-of- $(2k+1)$. Then, we have

$$\begin{aligned} \left| f'_{2k+1} \left(\frac{1}{2} \right) \right| &= (2k+1) \binom{2k}{k} 4^{-k} \geq 1.05\sqrt{k}, \\ |f'_{2k+1}(x)| &\leq \left| f'_{2k+1} \left(\frac{1}{2} \right) \right| \leq \frac{3}{\sqrt{\pi}} \sqrt{k} \leq 2\sqrt{k}, \\ |f''_{2k+1}(x)| &\leq \left| f''_{2k+1} \left(\frac{1}{2} + \frac{1}{2\sqrt{2k-1}} \right) \right| < 1.6k \end{aligned}$$

for sufficiently large k . Here, we used $\frac{4^k}{\sqrt{\pi k}} \left(1 - \frac{1}{8k}\right) \leq \binom{2k}{k} \leq \frac{4^k}{\sqrt{\pi k}}$.

From Lemma 2.3 and Lemma 3.6 (note that $f_{2k+1}(x)$ satisfies $f_{2k+1}(x) + f_{2k+1}(1-x) = 1$), it holds for all $A \subseteq V$ that

$$|\mathbf{E}[\alpha'] - f_{2k+1}(\alpha)| \leq 0.4k\lambda(\delta + \lambda)\alpha(1-\alpha) \quad (22)$$

$$|\mathbf{Var}[\alpha'] - g_{2k+1}(1/2)\|\pi\|_2^2| \leq 2\sqrt{k} \left(\frac{\|\pi\|_2^2}{2} |\delta| + \lambda \|\pi\|_3^{3/2} \right), \quad (23)$$

where $g_{2k+1}(x) = f(x)(1-f(x))$. Note that $g'_{2k+1}(x) = f'_{2k+1}(x)(1-2f_{2k+1}(x))$ satisfies $|g'_{2k+1}(x)| \leq |f'_{2k+1}(x)| \leq 2\sqrt{k}$. Thus, from the Hoeffding bound (Lemma A.3), it holds w.h.p. that

$$\begin{aligned} |\alpha' - f_{2k+1}(\alpha)| &\leq 0.4k\lambda^2\alpha(1-\alpha) + \|\pi\|_2\sqrt{\log n} \\ &\leq 0.4k\lambda^2\alpha(1-\alpha) + \sqrt{\frac{C \log n}{n}}. \end{aligned} \quad (24)$$

On the other hand, it is routine to check the following facts.

$$\lambda\sqrt{k\|\pi\|_3^3} = o(n^{-1}), \quad (25)$$

$$\frac{1}{n} \leq \|\pi\|_2^2 \leq \frac{C}{n}, \quad (26)$$

$$k\lambda^2 = O(1/\sqrt{n}). \quad (27)$$

We begin with proving the following result that corresponds to Lemma 4.3.

Lemma 6.2. *There exists constants $h, c > 0$ such that, for any $A \subseteq V$ satisfying $h/\sqrt{nk} \leq |\delta| \leq 1.25/\sqrt{k}$,*

$$\Pr[|\delta'| < 0.025\sqrt{k}|\delta|] \leq \exp(-ckn\delta^2).$$

Proof. Let h be a sufficiently large constant and let $A \subseteq V$ be a configuration satisfying $\frac{h}{\sqrt{kn}} \leq |\delta| \leq \frac{1.25}{\sqrt{k}}$. From (22), (27) and Taylor's theorem, we have

$$\begin{aligned} |\mathbf{E}[\delta']| &\geq \left| 2f_{2k+1}\left(\frac{1}{2} + \frac{\delta}{2}\right) - 1 \right| - 0.4k\lambda^2\alpha(1-\alpha) \\ &\geq f'_{2k+1}\left(\frac{1}{2}\right)|\delta| - \max_{0 \leq z \leq 1} |f''_{2k+1}(z)| \frac{\delta^2}{2} - 0.1k\lambda^2 \\ &\geq 0.05\sqrt{k}|\delta| + (\sqrt{k}|\delta| - 0.8k\delta^2) - 0.1k\lambda^2 \\ &\geq 0.05\sqrt{k}|\delta| + \frac{0.01h}{\sqrt{n}} - 0.1k\lambda^2 \\ &\geq 0.05\sqrt{k}|\delta|. \end{aligned}$$

In the fourth inequality, note that $\sqrt{k}|\delta| \geq 0.8k\delta^2$ holds if $|\delta| \leq 1.25/\sqrt{k}$. In the last inequality, we used $\lambda = O(k^{-0.5}n^{-0.25})$ and thus $k\lambda^2 = O(1/\sqrt{n}) \leq 0.01h/\sqrt{n}$ for sufficiently large constant h . Then, from Corollary A.4, we have

$$\begin{aligned} \Pr[|\delta'| < 0.025\sqrt{k}|\delta|] &\leq \Pr[|\delta'| < 0.5|\mathbf{E}[\delta']|] \\ &\leq 2 \exp\left(-\frac{0.5|\mathbf{E}[\delta']|^2}{\|\pi\|_2^2}\right) \\ &\leq \exp(-ckn\delta^2) \end{aligned}$$

for some suitable constant $c > 0$. In the last inequality, we used (26). \square

6.1 Phase (I): $0 \leq |\delta| \leq 300C \log n / \sqrt{n}$

In this part, we show Lemma 6.1(I). The proof is almost same as that of Lemma 4.1(I) that is presented in Section 4.2. The difference is that we use the following result, which is a slight modification of Lemma 4.2.

Lemma 6.3 (Modification of Lemma 4.2). *Consider a Markov chain $(X_t)_{t=1}^\infty$ with finite state space Ω and a function $\Psi : \Omega \rightarrow [0, n]$. Let C_1 be an arbitrary constant and $m = C_1\sqrt{n} \log n$. Let $k = k(n)$ be a function such that $k(n) \rightarrow \infty$ as $n \rightarrow \infty$. Suppose that Ω, Ψ and m satisfies the following conditions:*

(i) *For any positive constant h , there exists a positive constant $C_2 < 1$ such that*

$$\Pr\left[\Psi(X_{t+1}) < h\sqrt{\frac{n}{k}} \mid \Psi(X_t) \leq m\right] < \frac{C_2}{\sqrt{k}}.$$

(ii) *Three positive constants C_3, C_4 and h exist such that, for any $x \in \Omega$ satisfying $h\sqrt{n/k} \leq \Psi(x) < m$,*

$$\Pr\left[\Psi(X_{t+1}) < C_3\sqrt{k}\Psi(X_t) \mid X_t = x\right] < \exp\left(-C_4 \frac{k\Psi(x)^2}{n}\right).$$

Then, $\Psi(X_t) \geq m$ holds w.h.p. for some $t = O(\log n / \log k)$.

We prove Lemma 6.3 in Section 6.5. Lemma 6.1(I) is immediate from Lemma 6.3 with letting $\Psi(A) = n|\delta|$ and $C_1 = 300C$. Hence, it suffices to verify the conditions (i) and (ii).

Condition (i). First we evaluate the variance $\mathbf{Var}[\delta']$.

Claim 6.4. *Under the same assumption as Lemma 6.1(I),*

$$\mathbf{Var}[\delta'] \geq \frac{0.99}{n}.$$

Proof of the claim. Note that $\mathbf{Var}[\delta'] = 4 \mathbf{Var}[\alpha']$. From (23), we can evaluate the variance $\mathbf{Var}[\alpha']$ as follows:

$$\begin{aligned} \mathbf{Var}[\alpha'] &\geq g_{2k+1}(1/2) \|\pi\|_2^2 - \sqrt{k}|\delta| \|\pi\|_2^2 - 2\sqrt{k}\lambda \|\pi\|_3^{3/2} \\ &\geq \frac{1}{4n} - 3\sqrt{k} \left(300C^2 \sqrt{\frac{\log n}{n^3}} + \lambda \|\pi\|_3^{3/2} \right) \quad (\text{from (26)}) \\ &= \frac{1 - o(1)}{4n} \quad (\text{since } k = o(\log n/n) \text{ and (25)}) \\ &\geq \frac{0.99}{4n}. \end{aligned}$$

□

From Corollary A.6, for any positive real x , we have

$$\Pr \left[|\delta'| \leq x \sqrt{\frac{0.99}{n}} \right] \leq \Phi(x) + \frac{5.6 \|\pi\|_3^3}{\mathbf{Var}[\delta']^{3/2}} = \Phi(x) + o(1),$$

where $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} dy$ (see (17)). This yields the condition (i).

Condition (ii). This condition directly follows Lemma 6.2 by substituting $|\delta| = \frac{\Psi(A)}{n}$.

6.2 Phase (II): $300C \log n / \sqrt{n} \leq |\delta| \leq 1.25/\sqrt{k}$

Since $|\delta| \geq 300C \log n / \sqrt{n}$, from Lemma 6.2, we have

$$\Pr[|\delta'| < 0.025\sqrt{k}|\delta|] \leq \exp \left(-\frac{ckn(\log n)^2}{n} \right) \leq n^{-2}$$

if k is sufficiently large. Thus, we have $|\delta_t| \geq (0.025\sqrt{k})^t \cdot 300C \log n / \sqrt{n}$ with probability $(1 - n^{-2})^t$. Therefore, for some $t = O(\log n / \log k)$, $|\delta_t| \geq 1.25/\sqrt{k}$ with probability $1 - n^{-1.9}$.

6.3 Phase (III): $1.25/\sqrt{k} < |\delta| \leq 0.9$

We may assume that $\delta \geq 0$ without loss of generality (otherwise, consider A^c). From (24), we have

$$\begin{aligned}\delta' &\geq 2f_{2k+1}\left(\frac{1}{2} + \frac{\delta}{2}\right) - 1 - 0.4k\lambda^2 - 2\sqrt{\frac{K \log n}{n}} \\ &\geq 2f_{2k+1}\left(\frac{1}{2} + \frac{\delta}{2}\right) - 1 - o(1).\end{aligned}$$

We claim that $2f_{2k+1}\left(\frac{1}{2} + \frac{\delta}{2}\right) - 1 > 0.9$ during this phase (for sufficiently large n and k). Let $\text{Bin}(N, p)$ denote the random variable of binomial distribution with N trials and probability p . Then, from the definition of f_{2k+1} , it holds that

$$\begin{aligned}f_{2k+1}\left(\frac{1}{2} + \delta\right) &= \mathbf{Pr}\left[\text{Bin}\left(2k+1, \frac{1}{2} + \delta\right) \geq k+1\right] \\ &= 1 - \mathbf{Pr}\left[\text{Bin}\left(2k+1, \frac{1}{2} + \delta\right) \leq k\right].\end{aligned}\tag{28}$$

Let $\mu = (2k+1)(1/2 + \delta)$ be the expectation of $\text{Bin}(2k+1, 1/2 + \delta)$. Then, since $\mu - k \geq 2k\delta$, we have

$$\begin{aligned}\mathbf{Pr}\left[\text{Bin}\left(2k+1, \frac{1}{2} + \delta\right) \leq k\right] &\leq \mathbf{Pr}\left[\text{Bin}\left(2k+1, \frac{1}{2} + \delta\right) \leq \mu - (\mu - k)\right] \\ &\leq \mathbf{Pr}\left[\text{Bin}\left(2k+1, \frac{1}{2} + \delta\right) \leq \mu - 2k\delta\right] \\ &\leq \exp(-2k\delta^2).\end{aligned}\tag{29}$$

In the third inequality, we applied the Hoeffding bound (Lemma A.3). If $\delta \geq \frac{1.25}{\sqrt{k}}$, by combining (28) and (29), we obtain

$$\begin{aligned}f_{2k+1}\left(\frac{1}{2} + \delta\right) &\geq 1 - \exp(-2k\delta^2) \\ &\geq 1 - e^{-3.125} \\ &> 0.92.\end{aligned}$$

Thus, from (24), $\delta' \geq 0.92 - o(1) > 0.9$ holds w.h.p.

6.4 Phase (IV): $0.9 < |\delta| \leq 1$

We may assume $\pi(A_0) \leq 0.1$ without loss of generality. We claim that $\pi(A_t) < \frac{1}{n^2}$ for some $t = O(\log n / \log k)$, which implies $A_t = \emptyset$ (since $\pi(S) \geq \frac{1}{2m} \geq \frac{1}{n^2}$ whenever $S \neq \emptyset$).

Observe that

$$\begin{aligned}f_{2k+1}(x) &= \sum_{i=k+1}^{2k+1} \binom{2k+1}{i} x^i (1-x)^{2k+1-i} \\ &\leq (4x)^k \\ &\leq \frac{x}{4k}\end{aligned}$$

whenever $x \leq 0.1 \leq 4^{-1}(16k)^{1/(k-1)}$ with $k \geq 2$. Therefore, from (22), we have

$$\mathbf{E}[\alpha'] \leq \left(\frac{1}{4k} + 0.4k\lambda^2 \right) \alpha.$$

From (24) and the upper bound of $\mathbf{E}[\alpha']$ above, it holds with probability $1 - O(n^{-3})$ that $\alpha' \leq 0.9$ conditioned on $\alpha \leq 0.1$. Thus,

$$\mathbf{E}[\pi(A_t)] \leq \left(\frac{1}{4k} + 0.4k\lambda^2 \right)^t + n^{-3+o(1)} \leq n^{-3+o(1)}$$

for some $t = O(\log n / \log k + \log n / \log \lambda^{-1}) = O(\log n / \log k)$ (note that $\lambda^{-1} = \Omega(n^{1/4})$ from (27)). For this t , we have $\mathbf{Pr}[A_t \neq \emptyset] \leq \mathbf{Pr}[\pi(A_t) \geq n^{-2}] \leq n^2 \mathbf{E}[\pi(A_t)] = O(n^{-1})$. This completes the proof of Lemma 6.1 as well as Theorem 1.7.

6.5 Proof of Lemma 6.3

The proof is essentially given in [11]. By inspecting the proof of [11] with evaluating constant terms carefully, we obtain Lemma 6.3. For completeness, let us present it here.

Let $m = C_1 \sqrt{n} \log n$. Let $\tau = \inf\{t \in \mathbb{N} : \Psi(X_t) \geq m\}$ and $\{\tau(i)\}_{i \in \mathbb{N}}$ be the hitting times defined as

$$\begin{cases} \tau(0) = 0, \\ \tau(i) = \inf_{t \in \mathbb{N}} \{t : \tau(i-1) < t < \tau, f(X_t) \geq h\sqrt{n/k}\}. \end{cases}$$

Let R_1, R_2, \dots be the sequence of random variables defined as $R_i = X_{\tau(i)}$. It is shown in [11] that

- The sequence $(R_i)_{i \in \mathbb{N}}$ is a Markov chain.
- The sequence $(R_i)_{i \in \mathbb{N}}$ satisfies

$$\mathbf{Pr}[\Psi(R_{i+1}) < C_3 \sqrt{k} \Psi(R_i) | R_i = x] < \exp \left(-C_4 \frac{k \Psi(x)^2}{n} \right)$$

for any $x \in \Omega$ that $h\sqrt{n/k} \leq \Phi(x) < m$.

We claim that $\Psi(R_i) \geq m$ for some $i = O(\log n / \log k)$. To prove this, we use the Markov inequality. Fix a state $x \in \Omega$ such that $h\sqrt{n/k} \leq \Psi(x) < m$ for a sufficiently large constant h . Let $Y_i = \exp(-\frac{\Psi(R_i)}{\sqrt{n}})$ for each i . Let $y = \exp(-\frac{\Psi(x)}{\sqrt{n}})$ and $z = z(x) = \frac{\sqrt{k}\Psi(x)}{\sqrt{n}} \geq h$ for $x \in \Omega$. Note that

$e^z = y^{-\sqrt{k}}$. Then, we have

$$\begin{aligned}
& \mathbf{E}[Y_{i+1}|R_i = x] \\
& \leq \mathbf{Pr}[\Psi(R_{i+1}) < C_3\sqrt{k}\Psi(x)] + \mathbf{Pr}[\Psi(R_{i+1}) \geq C_3\sqrt{k}\Psi(x)] \cdot \exp\left(-C_3\sqrt{k}\frac{\Psi(x)}{\sqrt{n}}\right) \\
& \leq \exp\left(-C_4\frac{k\Psi(x)^2}{n}\right) + \exp\left(-C_3\frac{\sqrt{k}\Psi(x)}{\sqrt{n}}\right) \\
& = \exp(-C_4z^2) + \exp(-C_3z) \\
& = y^{-\frac{C_3}{2}\sqrt{k}} \left(\exp\left(\frac{C_3}{2}z - C_4z^2\right) + \exp\left(-\frac{C_3}{2}z\right) \right) \\
& \leq \frac{1}{2}y^{\frac{C_3}{2}\sqrt{k}} \quad (\text{since } z \geq h \text{ is sufficiently large and } C_2 > 1) \\
& \leq \begin{cases} \frac{1}{2} & \text{if } \frac{1}{2} < y_i \leq 1, \\ \frac{y}{C_3\sqrt{k}} & \text{if } y_i \leq \frac{1}{2}. \end{cases}
\end{aligned}$$

In the second part of the last inequality, we assume that $k \geq 2$; hence, it holds that $r^a \leq \frac{r}{a}$ for $0 \leq r \leq \frac{1}{2}$ if $a \geq 2$. Note that for each $i \geq 1$, the random variable $\Psi(R_i) = \Psi(X_{\tau(i)})$ satisfies $h\sqrt{n/k} \leq \Psi(R_i) < m$. Then, we have

$$\mathbf{E}[Y_i] \leq \frac{1}{2} \left(\frac{1}{C_3\sqrt{k}} \right)^{i-2}$$

and thus, by the Markov inequality,

$$\begin{aligned}
\mathbf{Pr}[\Psi(R_i) < m] &= \mathbf{Pr}\left[Y_i > \exp\left(-\frac{m}{\sqrt{n}}\right)\right] \\
&\leq \exp\left(\frac{m}{\sqrt{n}}\right) \frac{1}{2} \left(\frac{1}{C_3\sqrt{k}} \right)^{i-2} \\
&= \frac{n^{C_1}}{2(C_3\sqrt{k})^{i-2}} \\
&\leq n^{-1}
\end{aligned}$$

for $i = \lfloor C_5 \log n / \log k \rfloor$ for some constant C_5 that depends on C_1 and C_3 .

Finally, we consider $\tau(\lfloor C_5 \log n / \log k \rfloor)$. Let W_0, W_1, \dots be binary random variables defined as

$$W_t = \begin{cases} 1 & \text{if } \Psi(X_t) \geq h\sqrt{\frac{n}{k}}, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $\mathbf{Pr}[\tau(T_1) \geq T_2] = \mathbf{Pr}[\sum_{t=1}^{T_2} W_t \leq T_1]$. Let $\hat{W}_0, \hat{W}_1, \dots$ be i.i.d. binary random variables such that $\mathbf{E}[\hat{W}_t] = 1 - \frac{C_1}{\sqrt{k}}$. From the condition (i), for every T , the sum $\sum_{t=1}^T \hat{W}_t$ has stochastic

dominance over $\sum_{t=1}^T W_t$. Therefore, setting $T_1 = \lfloor \frac{C_4 \log n}{\log k} \rfloor$ and $T_2 = \lceil \frac{2C_4 \log n}{\log k} \rceil$, we obtain

$$\begin{aligned}
\Pr \left[\tau \left(\left\lfloor \frac{C_5 \log n}{\log k} \right\rfloor \right) \geq T_2 \right] &= \Pr \left[\sum_{t=1}^{T_2} W_t \leq \left\lfloor \frac{C_5 \log n}{\log k} \right\rfloor \right] \\
&\leq \Pr \left[\sum_{t=1}^{T_2} W_t \leq \frac{C_5 \log n}{\log k} \right] \\
&\leq \Pr \left[\sum_{t=1}^{T_2} \hat{W}_t \leq \frac{C_5 \log n}{\log k} \right] \\
&\leq \Pr \left[\sum_{t=1}^{T_2} (1 - \hat{W}_t) \geq T_2 - \frac{C_5 \log n}{\log k} \right] \\
&\leq \Pr \left[\sum_{t=1}^{T_2} (1 - \hat{W}_t) \geq \frac{C_5 \log n}{\log k} \right] \\
&\leq 2^{T_2} \left(\frac{C_1}{\sqrt{k}} \right)^{\frac{C_5 \log n}{\log k}} \\
&\leq n^{O(1/\log k) - \frac{C_5}{2}}.
\end{aligned}$$

In the fifth inequality, we used the union bound over the choice for \hat{W}_t . Note that $1 - \hat{W}_t = 1$ with probability $\frac{C_1}{\sqrt{k}}$. \square

7 Conclusion

In this paper we propose functional voting as a generalization of several known voting processes. We show that the consensus time is $O(\log n)$ for any quasi-majority functional voting on $O(n^{-1/2})$ -expander graphs with balanced degree distributions. This result extends previous works concerning voting processes on expander graphs. Possible future direction of this work includes

1. Does $O(\log n)$ worst-case consensus time holds for quasi-majority functional voting on graphs with less expansion (i.e., $\lambda = \omega(n^{-1/2})$)?
2. Is there some relationship between best-of- k and Majority?

Acknowledgements

This work is supported by JSPS KAKENHI Grant Number 19J12876 and 19K20214, Japan.

References

- [1] M. A. Abdullah and M. Draief. Global majority consensus by local majority polling on graphs of a given degree sequence. *Discrete Applied Mathematics*, 1(10):1–10, 2015.
- [2] Y. Afek, N. Alon, O. Barad, E. Hornstein, N. Barkai, and Z. Bar-Joseph. A biological solution to a fundamental distributed computing problem. *Science*, 331(6014):183–185, 2011.

- [3] D. Aldous and J. Fill. Reversible Markov chains and random walks on graphs. <http://statwww.berkeley.edu/pub/users/aldous/RWG/book.html>.
- [4] L. Becchetti, A. Clementi, E. Natale, F. Pasquale, R. Silvestri, and L. Trevisan. Simple dynamics for plurality consensus. *Distributed Computing*, 30(4):293–306, 2017.
- [5] L. Becchetti, A. Clementi, E. Natale, F. Pasquale, and L. Trevisan. Stabilizing consensus with many opinions. In *Proceedings of the 27th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 620–635, 2016.
- [6] I. Benjamini, S.-O. Chan, R. O’Donnell, O. Tamuz, and L.-Y. Tardos. Convergence, unanimity and disagreement in majority dynamics on unimodular graphs and random graphs. *Stochastic Processes and their Applications*, 126(9):2719–2733, 2016.
- [7] P. Berenbrink, A. Clementi, R. Elsässer, P. Kling, F. Mallmann-Trenn, and E. Natale. Ignore or comply? On breaking symmetry in consensus. In *Proceedings of the ACM Symposium on Principles of Distributed Computing (PODC)*, pages 335–344, 2017.
- [8] P. Berenbrink, G. Giakkoupis, A.-M. Kermarrec, and F. Mallmann-Trenn. Bounds on the voter model in dynamic networks. In *Proceedings of the 43rd International Colloquium on Automata, Languages, and Programming (ICALP)*, 2016.
- [9] E. Berger. Dynamic monopolies of constant size. *Journal of Combinatorial Theory Series B*, 83(2):191–200, 2001.
- [10] R. P.-S. C. Castellano, M. A. Muñoz. The non-linear q -voter model. *Physical Review E*, 80, 2009.
- [11] A. Clementi, M. Ghaffari, L. Gualà, E. Natale, F. Pasquale, and G. Scornavacca. A tight analysis of the parallel undecided-state dynamics with two colors. In *Proceedings of the 43rd International Symposium on Mathematical Foundations of Computer Science (MFCS)*, 117(28):1–15, 2018.
- [12] A. Coja-Oghlan. On the laplacian eigenvalues of $G_{n,p}$. *Combinatorics, Probability and Computing*, 16(6):923–946, 2007.
- [13] N. Cook, L. Goldstein, and T. Johnson. Size biased couplings and the spectral gap for random regular graphs. *The Annals of Probability*, 46(1):72–125, 2018.
- [14] C. Cooper, R. Elsässer, H. Ono, and T. Radzik. Coalescing random walks and voting on connected graphs. *SIAM Journal on Discrete Mathematics*, 27(4):1748–1758, 2013.
- [15] C. Cooper, R. Elsässer, and T. Radzik. The power of two choices in distributed voting. In *Proceedings of the 41st International Colloquium on Automata, Languages, and Programming (ICALP)*, 2:435–446, 2014.
- [16] C. Cooper, R. Elsässer, T. Radzik, N. Rivera, and T. Shiraga. Fast consensus for voting on general expander graphs. In *Proceedings of the 29th International Symposium on Distributed Computing (DISC)*, pages 248–262, 2015.
- [17] C. Cooper, T. Radzik, N. Rivera, and T. Shiraga. Fast plurality consensus in regular expanders. In *Proceedings of the 31st International Symposium on Distributed Computing (DISC)*, 91(13):1–16, 2017.

- [18] C. Cooper and N. Rivera. The linear voting model. *In Proceedings of the 43rd International Colloquium on Automata, Languages, and Programming (ICALP)*, 55(144):1–12, 2016.
- [19] E. Cruciani, E. Natale, A. Nusser, and G. Scornavacca. Phase transition of the 2-choices dynamics on core-periphery networks. *In Proceedings of the 17th International Conference on Autonomous Agents and Multiagent Systems (AAMAS)*, pages 777–785, 2018.
- [20] E. Cruciani, E. Natale, and G. Scornavacca. Distributed community detection via metastability of the 2-choices dynamics. *In Proceedings of the 33rd AAAI Conference on Artificial Intelligence (AAAI)*, pages 6046–6053, 2019.
- [21] B. Doerr, L. A. Goldberg, L. Minder, T. Sauerwald, and C. Scheideler. Stabilizing consensus with the power of two choices. *In Proceedings of the 23rd Annual ACM Symposium on Parallelism in Algorithms and Architectures (SPAA)*, pages 149–158, 2011.
- [22] B. Doerr and F. Neumann. *Theory of evolutionary computation: Recent developments in discrete optimization*. Springer International Publishing, 2020.
- [23] M. Fischer, N. Lynch, and M. Merritt. Easy impossibility proofs for distributed consensus problems. *Distributed Computing*, 1(1):26–39, 1986.
- [24] A. Frieze and M. Karoński. *Introduction to random graphs*. Cambridge University Press, 2016.
- [25] B. Gärtner and A. N. Zehmakan. Majority model on random regular graphs. *In Proceedings of the 13th Latin American Symposium on Theoretical Informatics (LATIN)*, pages 572–583, 2018.
- [26] M. Ghaffari and J. Lengler. Nearly-tight analysis for 2-choice and 3-majority consensus dynamics. *In Proceedings of the ACM Symposium on Principles of Distributed Computing (PODC)*, pages 305–313, 2018.
- [27] S. Gilbert and D. Kowalski. Distributed agreement with optimal communication complexity. *In Proceedings of the 21st Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 965–977, 2010.
- [28] Y. Hassin and D. Peleg. Distributed probabilistic polling and applications to proportionate agreement. *Information and Computation*, 171(2):248–268, 2001.
- [29] N. Kang and R. Rivera. Best-of-three voting on dense graphs. *In Proceedings of the 31st ACM Symposium on Parallelism in Algorithms and Architectures (SPAA)*, pages 115–121, 2019.
- [30] D. A. Levin and Y. Peres. *Markov chain and mixing times: second edition*. The American Mathematical Society, 2017.
- [31] T. M. Liggett. *Interacting particle systems*. Springer-Verlag, 1985.
- [32] R. Montenegro and P. Tetali. *Mathematical aspects of mixing times in Markov chains*. NOW Publishers, 2006.
- [33] E. Mossel, J. Neeman, and O. Tamuz. Majority dynamics and aggregation of information in social networks. *Autonomous Agents and Multiagent Systems*, 28(3):408–429, 2014.

- [34] T. Nakata, H. Imahayashi, and M. Yamashita. Probabilistic local majority voting for the agreement problem on finite graph. *In Proceedings of the 5th Annual International Computing and Combinatorics Conference (COCOON)*, pages 330–338, 1999.
- [35] D. Peleg. Size bounds for dynamic monopolies. *Discrete Applied Mathematics*, 86(2–3):263–273, 1998.
- [36] D. Peleg. Local majorities, coalitions and monopolies in graphs: a review. *Theoretical Computer Science*, 282(2):231–257, 2002.
- [37] G. Schoenebeck and F. Yu. Consensus of interacting particle systems on Erdős-Rényi graphs. *In Proceedings of the 29th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1945–1964, 2018.
- [38] I. G. Shevtsova. An improvement of convergence rate estimates in the Lyapunov theorem. *Doklady Mathematics*, 82(3):862–864, 2010.
- [39] N. Shimizu and T. Shiraga. Phase transitions of best-of-two and best-of-three on stochastic block models. *In Proceedings of the 33rd International Symposium on Distributed Computing (DISC)*, pages 32:1–32:17, 2019.
- [40] K. Tikhomirov and P. Youssef. The spectral gap of dense random regular graphs. *The Annals of Probability*, 47(1):362–419, 2019.
- [41] A. N. Zehmakan. Opinion forming in Erdős-Rényi random graph and expanders. *In Proceedings of the 29th International Symposium on Algorithms and Computation (ISAAC)*, pages 4:1–4:13, 2018.

A Tools

Lemma A.1 (Lemma 3 of [17]). *Suppose that P is reversible. Then, for any $S \subseteq V$,*

$$\left| \sum_{v \in V} \pi(v) P(v, S)^2 - \pi(S)^2 \right| \leq \lambda^2 \pi(S) (1 - \pi(S)).$$

Corollary A.2. *Suppose that P is reversible. Then, for any $S \subseteq V$,*

$$\sum_{v \in V} \pi(v) (P(v, S) - \pi(S))^2 \leq \lambda^2 \pi(S) (1 - \pi(S)).$$

Proof. Since $Q(V, S) = Q(S, V) = \pi(S)$ for any reversible P and $S \subseteq V$, we have

$$\begin{aligned} \sum_{v \in V} \pi(v) (P(v, S) - \pi(S))^2 &= \sum_{v \in V} \pi(v) P(v, S)^2 + \pi(S)^2 - 2\pi(S) Q(V, S) \\ &= \sum_{v \in V} \pi(v) P(v, S)^2 - \pi(S)^2 \leq \lambda^2 \pi(S) (1 - \pi(S)). \end{aligned}$$

Here, we invoked Lemma A.1 in the last inequality. □

Lemma A.3 (The Hoeffding bound (see, e.g., Theorem 10.9 of [22])). *Let Y_1, Y_2, \dots, Y_n be independent random variables. Assume that each Y_i takes values in a real interval $[a_i, b_i]$ of length $c_i := b_i - a_i$. Let $Y = \sum_{i=1}^n Y_i$. Then, for any $\kappa > 0$,*

$$\begin{aligned}\Pr[Y \geq \mathbf{E}[Y] + \kappa] &\leq \exp\left(-\frac{2\kappa^2}{\sum_{i=1}^n c_i^2}\right), \\ \Pr[Y \leq \mathbf{E}[Y] - \kappa] &\leq \exp\left(-\frac{2\kappa^2}{\sum_{i=1}^n c_i^2}\right).\end{aligned}$$

Corollary A.4. *Let Y_1, Y_2, \dots, Y_n be independent random variables. Assume that each Y_i takes values in a real interval $[a_i, b_i]$ of length $c_i := b_i - a_i$. Let $Y = \sum_{i=1}^n Y_i$. Then, for any $\kappa > 0$,*

$$\begin{aligned}\Pr[|Y| \geq |\mathbf{E}[Y]| + \kappa] &\leq 2 \exp\left(-\frac{2\kappa^2}{\sum_{i=1}^n c_i^2}\right), \\ \Pr[|Y| \leq |\mathbf{E}[Y]| - \kappa] &\leq 2 \exp\left(-\frac{2\kappa^2}{\sum_{i=1}^n c_i^2}\right).\end{aligned}$$

Proof. For the first inequality, it is straightforward to see that

$$\begin{aligned}\Pr[|Y| \geq |\mathbf{E}[Y]| + \kappa] &= \Pr[|Y| - |\mathbf{E}[Y]| \geq \kappa] \leq \Pr[|Y - \mathbf{E}[Y]| \geq \kappa] \\ &\leq 2 \exp\left(-\frac{2\kappa^2}{\sum_{i=1}^n c_i^2}\right).\end{aligned}$$

Note that $|x| - |y| \leq |x - y|$ for any $x, y \in \mathbb{R}$. Similarly, it holds that

$$\begin{aligned}\Pr[|Y| \leq |\mathbf{E}[Y]| - \kappa] &= \Pr[|\mathbf{E}[Y]| - |Y| \geq \kappa] \leq \Pr[|\mathbf{E}[Y] - Y| \geq \kappa] \\ &\leq 2 \exp\left(-\frac{2\kappa^2}{\sum_{i=1}^n c_i^2}\right),\end{aligned}$$

and we obtain the claim. \square

Lemma A.5 (Berry-Esseen theorem (see, e.g., [38])). *Let Y_1, Y_2, \dots, Y_n be independent random variables such that $\mathbf{E}[Y_i] = 0$, $\mathbf{E}[Y_i^2] > 0$, $\mathbf{E}[|Y_i|^3] < \infty$ for all $i \in [n]$, and $\sum_{i=1}^n \mathbf{E}[Y_i^2] = 1$. Let $Y = \sum_{i=1}^n Y_i$ and $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} dy$ (the cumulative distribution function of the standard normal distribution). Then*

$$\sup_{x \in \mathbb{R}} |\Pr[Y \leq x] - \Phi(x)| \leq 5.6 \sum_{i=1}^n \mathbf{E}[|Y_i|^3].$$

Corollary A.6. *Let Y_1, Y_2, \dots, Y_n be independent random variables, $c = (c_1, \dots, c_n) \in \mathbb{R}^n$ be a vector, and $Y = \sum_{i=1}^n Y_i$. Suppose that, for all $i \in [n]$, $|Y_i - \mathbf{E}[Y_i]| \leq c_i < \infty$ and $\mathbf{Var}[Y] > 0$. Let $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} dy$. Then, for any positive $x \in \mathbb{R}$,*

$$\Pr[|Y| \leq x \sqrt{\mathbf{Var}[Y]}] \leq \Phi(x) + \frac{5.6 \|c\|_3^3}{\mathbf{Var}[Y]^{3/2}}.$$

Proof. For each $i \in [n]$, let

$$Z_i := \frac{Y_i - \mathbf{E}[Y_i]}{\sqrt{\mathbf{Var}[Y]}}, \quad Z := \sum_{i \in [n]: \mathbf{E}[Z_i^2] > 0} Z_i = \sum_{i \in [n]} Z_i.$$

Note that $\mathbf{E}[Z_i^2] = 0 \iff \sum_z z^2 \mathbf{Pr}[Z_i = z] = 0 \iff \mathbf{Pr}[Z_i = 0] = 1$. For all $i \in \{j \in [n] : \mathbf{E}[Z_j^2] > 0\}$, it is easy to check that $\mathbf{E}[Z_i] = 0$, $\mathbf{E}[Z_i^2] > 0$, and $\mathbf{E}[|Z_i|^3] \leq \frac{c_i^3}{\mathbf{Var}[Y]^{3/2}} < \infty$. Furthermore,

$$\sum_{i \in [n]: \mathbf{E}[Z_i^2] > 0} \mathbf{E}[Z_i^2] = \sum_{i \in [n]} \mathbf{E}[Z_i^2] = \frac{\sum_{i \in [n]} \mathbf{E}[(Y_i - \mathbf{E}[Y_i])^2]}{\mathbf{Var}[Y]} = 1.$$

Thus, we can apply Lemma A.5 to Z and it holds that

$$\begin{aligned} \left| \mathbf{Pr} \left[\frac{Y - \mathbf{E}[Y]}{\sqrt{\mathbf{Var}[Y]}} \leq x \right] - \Phi(x) \right| &= \left| \mathbf{Pr} \left[\sum_{i=1}^n Z_i \leq x \right] - \Phi(x) \right| = |\mathbf{Pr}[Z \leq x] - \Phi(x)| \\ &\leq 5.6 \sum_{i \in [n]: \mathbf{E}[Z_i^2] > 0} \mathbf{E}[|Z_i|^3] \\ &\leq 5.6 \sum_{i=1}^n \frac{c_i^3}{\mathbf{Var}[Y]^{3/2}} = \frac{5.6 \|c\|_3^3}{\mathbf{Var}[Y]^{3/2}}. \end{aligned} \quad (30)$$

Next we observe that

$$\mathbf{Pr} \left[|Y| \geq x \sqrt{\mathbf{Var}[Y]} \right] = \mathbf{Pr} \left[Y \geq x \sqrt{\mathbf{Var}[Y]} \right] + \mathbf{Pr} \left[Y \leq -x \sqrt{\mathbf{Var}[Y]} \right] \quad (31)$$

holds. If $\mathbf{E}[Y] \geq 0$, we have

$$\begin{aligned} \mathbf{Pr} \left[|Y| \geq x \sqrt{\mathbf{Var}[Y]} \right] &\geq \mathbf{Pr} \left[Y \geq x \sqrt{\mathbf{Var}[Y]} + \mathbf{E}[Y] \right] \\ &\geq 1 - \mathbf{Pr} \left[Y - \mathbf{E}[Y] \leq x \sqrt{\mathbf{Var}[Y]} \right] \geq 1 - \Phi(x) - \frac{5.6 \|c\|_3^3}{\mathbf{Var}[Y]^{3/2}} \end{aligned}$$

from (30). Similarly, if $\mathbf{E}[Y] \leq 0$, (30) yields

$$\begin{aligned} \mathbf{Pr} \left[|Y| \geq x \sqrt{\mathbf{Var}[Y]} \right] &\geq \mathbf{Pr} \left[Y \leq -x \sqrt{\mathbf{Var}[Y]} + \mathbf{E}[Y] \right] \\ &= \mathbf{Pr} \left[Y - \mathbf{E}[Y] \leq -x \sqrt{\mathbf{Var}[Y]} \right] \geq \Phi(-x) - \frac{5.6 \|c\|_3^3}{\mathbf{Var}[Y]^{3/2}}. \end{aligned}$$

Thus, the claim holds for both cases. Note that $\Phi(-x) = 1 - \Phi(x)$ holds. \square