

Weighted Hurwitz numbers, τ -functions and matrix integrals^{†*}

J. Harnad^{1,2†}

¹*Department of Mathematics and Statistics, Concordia University
1455 de Maisonneuve Blvd. W. Montreal, QC H3G 1M8 Canada*

²*Centre de recherches mathématiques, Université de Montréal,
C. P. 6128, succ. centre ville, Montréal, QC H3C 3J7 Canada*

Abstract

The basis elements spanning the Sato Grassmannian element corresponding to the KP τ -function that serves as generating function for rationally weighted Hurwitz numbers are shown to be Meijer G -functions. Using their Mellin-Barnes integral representation the τ -function, evaluated at the trace invariants of an externally coupled matrix, is expressed as a matrix integral. Using the Mellin-Barnes integral transform of an infinite product of Γ functions, a similar matrix integral representation is given for the KP τ -function that serves as generating function for quantum weighted Hurwitz numbers.

1 Hurwitz numbers: classical and weighted

The fact that KP and 2D-Toda τ -functions of hypergeometric type serve as generating functions for *weighted Hurwitz numbers* was shown in [2–5], generalizing the case of *simple* (single and double) Hurwitz numbers [8, 9]. Sections 1.1 and 1.2 below, and Section 2 give a brief review of this theory, together with two illustrative examples: rational and quantum weighted Hurwitz numbers. In Section 3, it is shown how evaluation of such τ -functions at the trace invariants of a finite matrix may be expressed either as a Wronskian determinant or as a matrix integral. The content of subsections 3.2–3.4 are largely drawn from [6, 7], in which further details and proofs of the main results may be found.

*Text of invited presentation at: Quantum Theory and Symmetries, XIth International symposium, Centre de recherches mathématiques, Montréal, July 1-5, 2019.

†e-mail: harnad@crm.umontreal.ca

1.1 Geometric meaning of classical Hurwitz numbers

The Hurwitz number $H(\mu^{(1)}, \dots, \mu^{(k)})$ is the number of inequivalent branched N -sheeted covers $\Gamma \rightarrow \mathbf{P}^1$ of the Riemann sphere, with k branch points (Q_1, \dots, Q_k) , whose *ramification profiles* are given by k *partitions* $(\mu^{(1)}, \dots, \mu^{(k)})$ of N , normalized by dividing by the order $|\text{aut}(\Gamma)|$ of its automorphism group. The *Euler characteristic* χ and *genus* g of the covering curve is given by the *Riemann-Hurwitz formula*:

$$\chi = 2 - 2g = 2N - d, \quad d := \sum_{i=1}^l \ell^*(\mu^{(i)}), \quad (1.1)$$

where $\ell^*(\mu) := |\mu| - \ell(\mu) = N - \ell(\mu)$ is the *colength* of the partition.

The *Frobenius-Schur* formula gives $H(\mu^{(1)}, \dots, \mu^{(k)})$ in terms of S_N characters:

$$H(\mu^{(1)}, \dots, \mu^{(k)}) = \sum_{\lambda, |\lambda|=N} h^{k-2}(\lambda) \prod_{j=1}^k \frac{\chi_\lambda(\mu^{(j)})}{z_{\mu^{(j)}}}, \quad |\mu^{(i)}| = N, \quad (1.2)$$

where $h(\lambda) = \left(\det \frac{1}{(\lambda_i - i + j)!} \right)^{-1}$ is the product of the hook lengths of the partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_{\ell(\lambda)} > 0)$, $\chi_\lambda(\mu^{(j)})$ is the *irreducible character* of representation λ evaluated on the conjugacy class $\mu^{(j)}$, and

$$z_{\mu^{(j)}} := \prod_i i^{m_i(\mu^{(j)})} (m_i(\mu^{(j)}))! \quad (1.3)$$

is the order of the stabilizer of any element of $\text{cyc}(\mu^{(j)})$ (and $m_i(\mu^{(j)}) = \#$ parts of partition $\mu^{(j)}$ equal to i)

1.2 Weighted Hurwitz numbers [2–5]

Define the *weight generating function* $G(z)$, or its dual $\tilde{G}(z)$, as an infinite (or finite) product or sum (formal or convergent).

$$\begin{aligned} G(z) &= \prod_{i=1}^{\infty} (1 + z c_i) = 1 + \sum_{j=1}^{\infty} g_j z^j \\ \tilde{G}(z) &= \prod_{i=1}^{\infty} (1 - z c_i)^{-1} = 1 + \sum_{j=1}^{\infty} \tilde{g}_j z^j. \end{aligned} \quad (1.4)$$

The weight for a branched covering with ramification profiles $(\mu^{(1)}, \dots, \mu^{(k)})$ is defined to be:

$$\mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) := \frac{1}{k!} \sum_{\sigma \in S_k} \sum_{1 \leq i_1 < \dots < i_k} c_{i_{\sigma(1)}}^{\ell^*(\mu^{(1)})} \dots c_{i_{\sigma(k)}}^{\ell^*(\mu^{(k)})}$$

$$\widetilde{\mathcal{W}}_{\widetilde{G}}(\mu^{(1)}, \dots, \mu^{(k)}) := \frac{(-1)^{\sum_{i=1}^k \ell^*(\mu^{(i)}) + k}}{k!} \sum_{\sigma \in S_k} \sum_{1 \leq i_1 \leq \dots \leq i_k} c_{i_{\sigma(1)}}^{\ell^*(\mu^{(1)})} \dots c_{i_{\sigma(k)}}^{\ell^*(\mu^{(k)})} \quad (1.5)$$

Weighted double Hurwitz numbers $H_G^d(\mu, \nu)$, $H_{\widetilde{G}}^d(\mu, \nu)$ for n -sheeted branched coverings of the Riemann sphere having a pair of unweighted branch points (Q_0, Q_∞) , with ramification profiles of type (μ, ν) , and k additional weighted branch points (Q_1, \dots, Q_k) with ramification profiles $(\mu^{(1)}, \dots, \mu^{(k)})$ are defined as:

$$H_G^d(\mu, \nu) := \sum_{k=1}^d \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} \mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu),$$

$$H_{\widetilde{G}}^d(\mu, \nu) := \sum_{k=1}^d \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} \widetilde{\mathcal{W}}_{\widetilde{G}}(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu),$$

where \sum' denotes the sum over all partitions other than the cycle type of the identity element $(1)^n$. If Q_∞ is not a branch point; i.e. $\nu = (1)^n$, we have a weighted *single* Hurwitz number

$$H_G^d(\mu) := H_G^d(\mu, (1)^n). \quad (1.6)$$

Two cases of particular interest are: *rational weight generating functions*:

$$G_{\mathbf{c}, \mathbf{d}}(z) := \frac{\prod_{l=1}^L (1 + c_l z)}{\prod_{m=1}^M (1 - d_m z)}. \quad (1.7)$$

and *quantum weight generating function (quantum exponential)*:

$$\widetilde{G}(z) = H_q(z) := \prod_{i=0}^{\infty} (1 - q^i z)^{-1} = \sum_{n=0}^{\infty} \frac{z^n}{(q; q)_n}, \quad (1.8)$$

where

$$(q; q)_n := (1 - q)(1 - q^2) \dots (1 - q^n) \quad (1.9)$$

for some parameter q , with $|q| < 1$.

The corresponding *rationaly weighted* (single) Hurwitz numbers are:

$$H_{G_{\mathbf{c}, \mathbf{d}}}^d(\mu, \nu) := \sum_{\substack{1 \leq k, l \\ k+l \leq d}} \sum_{\substack{\mu^{(1)}, \dots, \mu^{(k)}, \nu^{(1)}, \dots, \nu^{(l)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) + \sum_{j=1}^l \ell^*(\nu^{(j)}) = d \\ |\mu^{(i)}| = |\nu^{(j)}| = N}} \mathcal{W}_{G_{\mathbf{c}, \mathbf{d}}}(\mu^{(1)}, \dots, \mu^{(k)}; \nu^{(1)}, \dots, \nu^{(l)}) \\ \times H(\mu^{(1)}, \dots, \mu^{(k)}, \nu^{(1)}, \dots, \nu^{(l)}, \mu),$$

where the rational weight factor is:

$$\begin{aligned} & \mathcal{W}_{G_{c,d}}(\mu^{(1)}, \dots, \mu^{(k)}; \nu^{(1)}, \dots, \nu^{(l)}) \\ &:= \frac{(-1)^{\sum_{j=1}^l \ell^*(\nu^{(j)}) - l}}{k!l!} \sum_{\substack{\sigma \in S_k \\ \sigma' \in S_l}} \sum_{\substack{1 \leq a_1 < \dots < a_k \leq M \\ 1 \leq b_1 \dots \leq b_k \leq L}} c_{a_{\sigma(1)}}^{\ell^*(\mu^{(1)})} \dots c_{a_{\sigma(k)}}^{\ell^*(\mu^{(k)})} d_{b_{\sigma'(1)}}^{\ell^*(\nu^{(1)})} \dots d_{b_{\sigma'(l)}}^{\ell^*(\nu^{(l)})}. \end{aligned}$$

The *quantum weighted* (single) Hurwitz numbers are

$$H_{H_q}^d(\mu) := \sum_{k=1}^d \sum_{\substack{\mu^{(1)}, \dots, \mu^{(k)}, |\mu^{(i)}| = N \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} \widetilde{\mathcal{W}}_{H_q}(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu), \quad (1.10)$$

where the *quantum weight factor* is

$$\widetilde{\mathcal{W}}_{H_q}(\mu^{(1)}, \dots, \mu^{(k)}) := \frac{(-1)^{d-k}}{k!} \sum_{\sigma \in S_k} \prod_{j=1}^k \frac{1}{(1 - q^{\sum_{i=1}^j \ell^*(\mu^{(\sigma(i)))})}.$$

2 Hypergeometric τ -functions as generating functions for weighted Hurwitz numbers [2–5]

To construct a KP τ -function of hypergeometric type that serves as generating function for weighted Hurwitz numbers for a given weight generating function G , choose a small parameter β and define coefficients $r_\lambda^{(G,\beta)}$ that are of *content product form*:

$$r_\lambda^{(G,\beta)} := \prod_{(ij) \in \lambda} r_{j-i}^{(G,\beta)} = \prod_{(ij) \in \lambda} G((j-i)\beta), \quad (2.1)$$

where

$$r_j^{(G,\beta)} := G(j\beta) = \frac{\rho_j^{(G,\beta)}}{\beta \rho_{j-1}^{(G,\beta)}}, \quad (2.2)$$

with

$$\begin{aligned} \rho_j^{(G,\beta)} &:= \beta^j \prod_{i=1}^j G(i\beta) =: e^{T_j^G(\beta)}, \quad \rho_0 = 1, = \frac{\rho_j^{(G,\beta)}}{\beta \rho_{j-1}^{(G,\beta)}}, \\ \rho_{-j}^{(G,\beta)} &:= \beta^{-j} \prod_{i=1}^{j-1} \frac{1}{G(-i\beta)} =: e^{T_{-j}^G(\beta)}, \quad j = 1, 2, \dots \end{aligned} \quad (2.3)$$

We then have [3, 5]:

Theorem 2.1 (Hypergeometric Toda τ -functions associated to weight generating function $G(z)$). *The double Schur function series*

$$\tau^{(G,\beta)}(\mathbf{t}, \mathbf{s}) := \sum_{\lambda} \beta^{|\lambda|} r_{\lambda}^{(G,\beta)} s_{\lambda}(\mathbf{t}) s_{\lambda}(\mathbf{s}) \quad (2.4)$$

defines a 2D-Toda τ -function (at lattice value $n = 0$) .

We now use the *Frobenius character formula*

$$s_{\lambda}(\mathbf{t}) = \sum_{\mu, |\mu|=|\lambda|} \frac{\chi_{\lambda}(\mu) p_{\mu}(\mathbf{t})}{z_{\mu}}, \quad s_{\lambda}(\mathbf{s}) = \sum_{\nu, |\nu|=|\lambda|} \frac{\chi_{\lambda}(\nu) p_{\nu}(\mathbf{s})}{z_{\nu}} \quad (2.5)$$

to change the basis of Schur functions to *power sum symmetric functions*

$$p_{\mu}(\mathbf{t}) := \prod_{i=1}^{\ell(\mu)} p_{\mu_i}(\mathbf{t}), \quad p_j(\mathbf{t}) = jt_j, \quad p_{\nu}(\mathbf{s}) := \prod_{i=1}^{\ell(\nu)} p_{\nu_i}(\mathbf{s}), \quad p_j(\mathbf{s}) = js_j. \quad (2.6)$$

Theorem 2.2 (Hypergeometric Toda τ -functions as generating function for weighted double Hurwitz numbers [3, 5]). *The τ -function $\tau^{(G,\beta)}(\mathbf{t}, \mathbf{s})$ can equivalently be expressed as a double infinite series in the bases of power sum symmetric functions as follows*

$$\tau^{(G,\beta)}(\mathbf{t}, \mathbf{s}) = \sum_{d=0}^{\infty} \sum_{\substack{\mu, \nu, \\ |\mu|=|\nu|=d}} \beta^{|\mu|+d} H_G^d(\mu, \nu) p_{\mu}(\mathbf{t}) p_{\nu}(\mathbf{s}). \quad (2.7)$$

It is thus a generating function for the numbers $H_G^d(\mu, \nu)$ of weighted n -fold branched coverings of the sphere, with a pair of specified branch points having ramification profiles (μ, ν) and genus given by the Riemann-Hurwitz formula

$$2 - 2g = \ell(\mu) + \ell(\nu) - d, \quad d = \sum_{i=1}^k \ell^*(\mu^{(i)}). \quad (2.8)$$

Corollary 2.3 (Hypergeometric KP τ -functions as generating functions for weighted single Hurwitz numbers). *Set: $\mathbf{s} = \beta^{-1} \mathbf{t}_0 := (\beta^{-1}, 0, 0, \dots)$.*

Then the series

$$\begin{aligned} \tau^{(G,\beta)}(\mathbf{t}, \beta^{-1} \mathbf{t}_0) &:= \tau^{(G,\beta)}(\mathbf{t}) = \sum_{\lambda} (h(\lambda))^{-1} r_{\lambda}^{(G,\beta)} s_{\lambda}(\mathbf{t}) \\ &= \sum_{d=0}^{\infty} \sum_{\mu} \beta^d H_G^d(\mu) p_{\mu}(\mathbf{t}) \end{aligned}$$

is a KP τ -function which is a generating function for weighted single numbers $H_G^d(\mu)$ for $|\mu|$ -fold branched coverings of the sphere, with a branch point having ramification profile (μ) at Q_0 and genus given by the Riemann-Hurwitz formula.

$$2 - 2g = |\mu| + \ell(\mu) - d. \quad (2.9)$$

3 Wronskian and matrix integral representation of $\tau^{(G,\beta)}([X])$

In [6, 7] new matrix integral representations were derived for the τ -functions that serve as generating functions for rationally and quantum weighted Hurwitz numbers. The main result is that, using Laurent series and Mellin-Barnes integral representations of the adapted bases for the respective elements of the infinite Grassmannian corresponding to these cases, the τ -functions may be expressed as Wronskian determinants or as matrix integrals.

3.1 Adapted basis, recursion operators, quantum spectral curve

Henceforth, we always set:

$$\mathbf{s} = \beta^{-1}\mathbf{t}_0 := (\beta^{-1}, 0, 0, \dots) \quad (3.1)$$

and

$$\tau^{(G,\beta)}(\mathbf{t}) := \tau^{(G,\beta)}(\mathbf{t}, \beta^{-1}\mathbf{t}_0) \quad (3.2)$$

is a KP τ -function of hypergeometric type.

For $k \in \mathbf{Z}$, define:

$$\begin{aligned} \phi_k(x) &:= \frac{\beta}{2\pi i x^{k-1}} \oint_{|\zeta|=\epsilon} \rho^{(G,\beta)}(\zeta) e^{\beta^{-1}x\zeta} \frac{d\zeta}{\zeta^k}, \\ &= \beta x^{1-k} \sum_{j=0}^{\infty} \frac{\rho_{j-k}^{(G,\beta)}}{j!} \left(\frac{x}{\beta}\right)^j, \end{aligned} \quad (3.3)$$

where

$$\rho^{(G,\beta)}(\zeta) := \sum_{i=-\infty}^{k-1} \rho_{-i-1}^{(G,\beta)} \zeta^i. \quad (3.4)$$

Then $\{\phi_k(1/z)\}_{k \in \mathbf{N}^+}$ is a basis for the element $w^{(G,\beta)}$ of the Sato Grassmannian that determines the KP τ -function $\tau^{(G,\beta)}(\mathbf{t})$ [1].

3.2 Quantum and classical spectral curve

Theorem 3.1 (Quantum spectral curve and eigenvalue equations [1]). *The functions $\phi_k(x)$ satisfy*

$$\mathcal{L}\phi_k(x) := (xG(\beta\mathcal{D}) - \mathcal{D})\phi_k(x) = (k-1)\phi_k(x) \quad (3.5)$$

where $\mathcal{D} := x \frac{d}{dx}$ is the Euler operator.

The classical spectral curve is

$$y = G(\beta xy). \quad (3.6)$$

3.2.1 Rational weighting case

For $G(z) = G_{\mathbf{c}, \mathbf{d}}(z)$, denote $\phi_k(x) =: \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$. Then

$$\zeta \prod_{l=1}^L \left(\mathcal{D} + \frac{1}{\beta c_l} \right) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} + (\mathcal{D} + k - 1) \prod_{m=1}^M \left(\mathcal{D} - 1 - \frac{1}{\beta d_m} \right) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} = 0, \quad (3.7)$$

where

$$\zeta := -\kappa_{\mathbf{c}, \mathbf{d}} x, \quad \kappa_{\mathbf{c}, \mathbf{d}} := (-1)^M \frac{\prod_{l=1}^L \beta c_l}{\prod_{m=1}^M \beta d_m}. \quad (3.8)$$

3.2.2 Mellin-Barnes integral representation: Meijer G -functions [6, 7]

It may be shown that $\phi_k^{(\mathbf{c}, \mathbf{d}, \beta)}$ has the Mellin-Barnes integral representation:

$$\begin{aligned} \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} &= C_k^{(\mathbf{c}, \mathbf{d}, \beta)} G_{L, M+1}^{1, L} \left(\begin{matrix} 1 - \frac{1}{\beta c_1}, \dots, 1 - \frac{1}{\beta c_L} \\ 1 - k, 1 + \frac{1}{\beta d_1}, \dots, 1 + \frac{1}{\beta d_M} \end{matrix} \middle| -\kappa_{\mathbf{c}, \mathbf{d}} x \right) \\ &= \frac{C_k^{(\mathbf{c}, \mathbf{d}, \beta)}}{2\pi i} \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta c_\ell}\right) (-\kappa_{\mathbf{c}, \mathbf{d}} x)^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta d_m}\right)} ds. \\ &\sim \frac{\beta \rho_{-k}(\mathbf{c}, \mathbf{d})}{(\kappa x)^{k-1}} {}_L F_M \left(\begin{matrix} 1 - k + \frac{1}{\beta c_1}, \dots, 1 - k + \frac{1}{\beta c_L} \\ 1 - k - \frac{1}{\beta d_1}, \dots, 1 - k - \frac{1}{\beta d_M} \end{matrix} \middle| \kappa_{\mathbf{c}, \mathbf{d}} x \right) \end{aligned} \quad (3.9)$$

where

$$C_k^{(\mathbf{c}, \mathbf{d}, \beta)} := \frac{\prod_{j=1}^M \Gamma\left(-\frac{1}{\beta d_j}\right)}{(-\beta)^{k-1} \prod_{\ell=1}^L \Gamma\left(\frac{1}{\beta c_\ell}\right)}, \quad (3.10)$$

The contour \mathcal{C}_k is chosen so that the poles at $1 - k, 2 - k, \dots$ are to the right and the poles at $\{-i - \frac{1}{\beta c_j}\}_{j=1, \dots, L, i \in \mathbf{N}^+}$ to the left. (See Figure 1.)




Figure 1: The contours of integration for the function $\phi_k^{(\mathbf{c}, \mathbf{d}, \beta)}$ in the case $L > M + 1$.

3.2.3 Quantum case expressed as Mellin-Barnes integrals [7]

The following is an integral representation of $\phi_k^{(H_q, \beta)}(x)$, valid for all $x \in \mathbf{C}$,

$$\phi_k^{(H_q, \beta)} = \frac{1}{2\pi i} \int_{\mathcal{C}_k} A_{H_q, k}(s) x^s ds, \quad (3.11)$$

where

$$A_{H_q, k}(z) := (-\beta)^{1-k} \Gamma(1-k-z) \prod_{m=0}^{\infty} \left((-\beta q^m)^{-z} \frac{\Gamma(-\beta^{-1} q^{-m})}{\Gamma(z - \beta^{-1} q^{-m})} \right). \quad (3.12)$$

The contour \mathcal{C}_k is defined as starting at $+\infty$ immediately above the real axis, proceeding to the left above the axis, winding around the poles at the integers $s = -k, -k+1, \dots$ in a counterclockwise sense and continuing below the axis back to $+\infty$.

3.3 Determinantal representation of $\tau^{(G, \beta)}(\mathbf{t})$

If $\tau^{(G, \beta)}(\mathbf{t})$ is evaluated at the trace invariants of diagonal $X \in \text{Mat}^{n \times n}$

$$\begin{aligned} \mathbf{t} &= [X], \quad t_i = \frac{1}{i} \text{tr} X^i, \\ X &:= \text{diag}(x_1, \dots, x_n), \end{aligned} \quad (3.13)$$

it is expressible as the ratio of $n \times n$ determinants

$$\tau^{(G, \beta)}([X]) = \frac{\prod_{i=1}^n x_i^{n-1} \det(\phi_i(x_j))_{1 \leq i, j \leq n}}{\prod_{i=1}^n \rho_{-i} \Delta(x)}, \quad (3.14)$$

where

$$\Delta(x) = \prod_{1 \leq i < j \leq n} (x_i - x_j) = \det(x_i^{n-j})_{1 \leq i, j \leq n} \quad (3.15)$$

is the Vandermonde determinant.

3.3.1 Eulerian Wronskian representation

It follows from the recursion relations

$$\beta(\mathcal{D} + k - 1)\phi_k = \phi_{k-1}, \quad k \in \mathbf{Z}, \quad (3.16)$$

that

$$\tau^{(G, \beta)}([X]) = \gamma_n \left(\prod_{i=1}^n x_i^{n-1} \right) \frac{\det(\mathcal{D}^{i-1} \phi_n(x_j))_{1 \leq i, j \leq n}}{\Delta(x)}, \quad (3.17)$$

where

$$\gamma_n := \frac{\beta^{\frac{1}{2}n(n-1)}}{\prod_{i=1}^n \rho_{-i}}. \quad (3.18)$$

3.4 Matrix integral representation of $\tau^{(G, \beta)}([X])$ [6, 7]

3.4.1 Wronskian representation: rational case

For rational weight generating functions $G = G_{\mathbf{c}, \mathbf{d}}$, and any $n \in \mathbf{N}^+$, let

$$\begin{aligned} \phi_n^{(\mathbf{c}, \mathbf{d}, \beta)}(e^y) &= \int_{\mathcal{C}_n} A_n^{(\mathbf{c}, \mathbf{d}, \beta)}(s) e^{ys} ds, \\ A_n^{(\mathbf{c}, \mathbf{d}, \beta)}(s) &:= \frac{C_n^{(\mathbf{c}, \mathbf{d}, \beta)} \Gamma(1 - n - s) \prod_{l=1}^L \Gamma\left(s + \frac{1}{\beta c_l}\right) (-\kappa_{\mathbf{c}, \mathbf{d}})^s}{2\pi i \prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta d_m}\right)}. \end{aligned}$$

Define the diagonal matrix $Y = \text{diag}(y_1, \dots, y_n)$

$$X = e^Y, \quad Y = \ln(X), \quad x_i = e^{y_i}, \quad i = 1, \dots, n. \quad (3.19)$$

Then $\tau^{(G_{\mathbf{c}, \mathbf{d}}, \beta)}([X])$ becomes a ratio of Wronskian determinants

$$\tau^{(G_{\mathbf{c}, \mathbf{d}}, \beta)}([X]) = \gamma_n \left(\prod_{i=1}^n x_i^{n-1} \right) \frac{\det\left(\left(\phi_n^{(\mathbf{c}, \mathbf{d}, \beta)}\right)^{(i-1)}(e^{y_j})\right)_{1 \leq i, j \leq n}}{\Delta(e^y)}. \quad (3.20)$$

3.4.2 Matrix integral representation of $\tau^{(G,\beta)}([X])$: rational case

It follows [6] that

$$\tau^{(G_{\mathbf{c},\mathbf{d},\beta})}([X]) = \frac{\beta^{\frac{1}{2}n(n-1)}(\prod_{i=1}^n x_i^{n-1})\Delta(\ln(x))}{(\prod_{i=1}^n i!)\Delta(x)} \mathbf{Z}_{d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}}(X), \quad (3.21)$$

where

$$\mathbf{Z}_{d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}}(X) = \int_{M \in \text{Nor}_{\mathcal{C}_n}^{n \times n}} d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}(M) e^{\text{tr} Y M} \quad (3.22)$$

and

$$d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}(M) := (\Delta(\zeta)^2 \det(A_n^{(\mathbf{c},\mathbf{d},\beta)}(M)) d\mu_0(U) \prod_{j=1}^n d\zeta_j$$

is a conjugation invariant measure on the space of normal matrices

$$M = U Z U^\dagger \in \text{Nor}_{\mathcal{C}_n}^{n \times n}, \quad U \in U(n), \quad Z = \text{diag}(\zeta_1, \dots, \zeta_n) \quad (3.23)$$

with eigenvalues $\zeta_i \in \mathbf{C}$ supported on the contour \mathcal{C}_n .

3.4.3 Wronskian representation: quantum case

For quantum weight generating functions $G = H_q$, and any $n \in \mathbf{N}^+$, let

$$\begin{aligned} \phi_n^{(H_q,\beta)}(e^y) &= \int_{\mathcal{C}_n} A_n^{(\mathbf{c},\mathbf{d},\beta)}(s) e^{ys} ds, \\ A_{H_q,n}(z) &:= (-\beta)^{1-n} \Gamma(1-n-z) \prod_{m=0}^{\infty} \left((-\beta q^m)^{-z} \frac{\Gamma(-\beta^{-1} q^{-m})}{\Gamma(z - \beta^{-1} q^{-m})} \right). \end{aligned}$$

Define the diagonal matrix $Y = \text{diag}(y_1, \dots, y_n)$

$$X = e^Y, \quad Y = \ln(X), \quad x_i = e^{y_i}, \quad i = 1, \dots, n, \quad (3.24)$$

Then $\tau^{(H_q,\beta)}([X])$ becomes a ratio of Wronskian determinants

$$\tau^{(H_q,\beta)}([X]) = \gamma_n \left(\prod_{i=1}^n x_i^{n-1} \right) \frac{\det \left((\phi_n^{(\mathbf{c},\mathbf{d},\beta)})^{(i-1)}(e^{y_j}) \right)_{1 \leq i,j \leq n}}{\Delta(e^y)}. \quad (3.25)$$

3.4.4 Matrix integral representation of $\tau^{(G,\beta)}([X])$: quantum case

It similarly follows [7] that

$$\tau^{(H_q,\beta)}([X]) = \frac{\beta^{\frac{1}{2}n(n-1)}(\prod_{i=1}^n x_i^{n-1})\Delta(\ln(x))}{(\prod_{i=1}^n i!)\Delta(x)} \mathbf{Z}_{d\mu_q}(\ln(X)), \quad (3.26)$$

$$\text{where } \mathbf{Z}_{d\mu_{(q,n)}}(X) = \int_{M \in \text{Nor}_{\mathcal{C}_n}^{n \times n}} d\mu_{(q,n)}(M) e^{\text{tr} Y M},$$

$$\text{and } d\mu_{(q,n)}(M) := (\Delta(\zeta)^2 \det(A_{H_q,n}(M)))$$

is a conjugation invariant measure on the space of normal matrices

$$M = U Z U^\dagger \in \text{Nor}_{\mathcal{C}_n}^{n \times n}, \quad U \in U(n), \quad Z = \text{diag}(\zeta_1, \dots, \zeta_n) \quad (3.27)$$

with eigenvalues $\zeta_i \in \mathbf{C}$ supported on the contour \mathcal{C}_n .

Acknowledgements. This work was partially supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Fonds de recherche du Québec, Nature et technologies (FRQNT).

References

- [1] A. Alexandrov, G. Chapuy, B. Eynard and J. Harnad, “Weighted Hurwitz numbers and topological recursion: an overview”, *J. Math. Phys.* **59**, 081102: 1-20 (2018).
- [2] M. Guay-Paquet and J. Harnad, “2D Toda τ -functions as combinatorial generating functions”, *Lett. Math. Phys.* **105**, 827-852 (2015).
- [3] J. Harnad and A. Yu. Orlov, “Hypergeometric τ -functions, Hurwitz numbers and enumeration of paths”, *Commun. Math. Phys.* **338**, 267-284 (2015).
- [4] J. Harnad, “Weighted Hurwitz numbers and hypergeometric τ -functions: an overview”, *AMS Proceedings of Symposia in Pure Mathematics* **93**, 289-333 (2016).
- [5] M. Guay-Paquet and J. Harnad, “Generating functions for weighted Hurwitz numbers”, *J. Math. Phys.* **58**, 083503 (2017).
- [6] M. Bertola and J. Harnad, “Rationally weighted Hurwitz numbers, Meijer G -functions and matrix integrals”, *J. Math. Phys.* **60**, 103504 (2019).
- [7] J. Harnad and B. Runov, “Matrix model generating function for quantum weighted Hurwitz numbers”, *J. Phys. A* **53**, 065201 (2020).
- [8] A. Okounkov, “Toda equations for Hurwitz numbers”, *Math. Res. Lett.* **7**, 447-453 (2000).
- [9] A. Okounkov and R. Pandharipande, “Gromov-Witten theory, Hurwitz theory and completed cycles,” *Ann. Math.* **163**, 517-560 (2006).