

PARTITIONS OF FLAT ONE -VARIATE FUNCTIONS AND A FOURIER RESTRICTION THEOREM FOR RELATED PERTURBATIONS OF THE HYPERBOLIC PARABOLOID

STEFAN BUSCHENHENKE, DETLEF MÜLLER, AND ANA VARGAS

ABSTRACT. We continue our research on Fourier restriction for hyperbolic surfaces, by studying local perturbations of the hyperbolic paraboloid $z = xy$ which are of the form $z = xy + h(y)$, where $h(y)$ is a smooth function which is flat at the origin. The case of perturbations of finite type had already been handled before, but the flat case imposes several new obstacles. By means of a decomposition into intervals on which $|h'''|$ is of a fixed size λ , we can apply methods devised in preceding papers, but since we loose control on higher order derivatives of h we are forced to rework the bilinear method for wave packets that are only slowly decaying. Another problem lies in the passage from bilinear estimates to linear estimates, for which we need to require some monotonicity of h''' .

CONTENTS

1. Introduction	2
2. A level-size decomposition	4
3. Fourier restriction over rectangles on which $h''' \sim \lambda$	8
3.1. Admissible pairs of sets U_1, U_2 on which transversalities are of a fixed size	9
3.2. Definition of admissible pairs within $Q \times Q$	10
3.3. A prototypical admissible pair in the curved box case and the crucial scaling transformation	14
4. The bilinear method with slowly decaying wave packets for the re-scaled prototypical case	16
4.1. Slowly decaying wave packets	17
4.2. Proof of Theorem 4.1 - Setup of the argument	22
4.3. Dyadic decompositions and reduction	25
4.4. The geometric argument	27
4.5. Completion of the proof of Theorem 4.1	30
5. Passage to linear restriction estimates and proof of Theorem 1.1	33
References	38

2010 *Mathematical Subject Classification.* 42B25.

Key words and phrases. hyperbolic hypersurface, Fourier restriction.

The first author was partially supported by the ERC grant 307617.

The first two authors were partially supported by the DFG grants MU 761/ 11-1 and MU 761/ 11-2.

The third author was partially supported by grants MTM2013-40945 (MINECO) and MTM2016-76566-P (Ministerio de Ciencia, Innovación y Universidades), Spain.

1. INTRODUCTION

In this article we continue our study of Fourier restriction to hyperbolic hypersurfaces that we had begun in [BMV17, BMV19].

In contrast to hyperbolic surfaces, Fourier restriction to hypersurfaces with non-negative principal curvatures has been studied intensively by many authors (see, e.g., [Bo91], [Bo95a], [Bo95b], [MVV96], [MVV99], [TVI00], [TVII00], [W01], [T01], [LV10], [BoG11], [BMV16], [Gu16], [Gu17], [Sto17a]). For the case of hypersurfaces of non-vanishing Gaussian curvature but principal curvatures of different signs, besides Tomas-Stein type Fourier restriction estimates (see, e.g., [To75], [Str77], [Gr81], [St86], [IKM10], [IM11], [IM15]), until recently the only case which had been studied successfully was the case of the hyperbolic paraboloid (or "saddle") in \mathbb{R}^3 : in 2015, independently S. Lee [L05] and A. Vargas [V05] established results analogous to Tao's theorem [T03] on elliptic surfaces (such as the 2-sphere), with the exception of the end-point, by means of the bilinear method. Recently, B. Stovall [Sto17b] was able to include also the end-point case. Moreover, C. H. Cho and J. Lee [ChL17], and J. Kim [K17], improved the range by adapting ideas by Guth [Gu16], [Gu17] which are based on the polynomial partitioning method. For further information on the history of the restriction problem, we refer the interested reader to our previous paper [BMV17].

We shall here study surfaces S which are local perturbations of the hyperbolic paraboloid $z = xy$ that are given as the graph of a function $\phi(x, y) := xy + h(y)$, where the function h is smooth and satisfies

$$(1.1) \quad h(0) = h'(0) = h''(0) = 0,$$

i.e.,

$$(1.2) \quad S := \{(x, y, xy + h(y)) : (x, y) \in \Omega\},$$

where Ω is a sufficiently small neighborhood of the origin. The Fourier restriction problem, introduced by E. M. Stein in the seventies (for general submanifolds), asks for the range of exponents \tilde{p} and \tilde{q} for which an a priori estimate of the form

$$\left(\int_S |\hat{f}|^{\tilde{q}} d\sigma \right)^{1/\tilde{q}} \leq C \|f\|_{L^{\tilde{p}}(\mathbb{R}^n)}$$

holds true for every Schwartz function $f \in \mathcal{S}(\mathbb{R}^3)$, with a constant C independent of f . Here, $d\sigma$ denotes the surface measure on S .

As usual, it will be more convenient to use duality and work in the adjoint setting. If \mathcal{R} denotes the Fourier restriction operator $g \mapsto \mathcal{R}g := \hat{g}|_S$ to the surface S , its adjoint operator \mathcal{R}^* is given by $\mathcal{R}^*f(\xi) = \mathcal{E}f(-\xi)$, where \mathcal{E} denotes the "Fourier extension" operator given by

$$\mathcal{E}f(\xi) := \widehat{f d\sigma}(\xi) = \int_S f(x) e^{-i\xi \cdot x} d\sigma(x),$$

with $f \in L^q(S, \sigma)$. The restriction problem is therefore equivalent to the question of finding the appropriate range of exponents for which the estimate

$$\|\mathcal{E}f\|_{L^r(\mathbb{R}^3)} \leq C \|f\|_{L^q(S, d\sigma)}$$

holds true with a constant C independent of the function $f \in L^q(S, d\sigma)$.

By identifying a point $(x, y) \in \Omega$ with the corresponding point $(x, y, \phi(x, y))$ on S , we may regard our Fourier extension operator \mathcal{E} as well as an operator mapping functions on Ω to

functions on \mathbb{R}^3 , which in terms of our phase function $\phi(x, y) = xy + h(y)$ can be expressed more explicitly in the form

$$\mathcal{E}f(\xi) = \int_{\Omega} f(x, y) e^{-i(\xi_1 x + \xi_2 y + \xi_3 \phi(x, y))} \eta(x, y) dx dy,$$

if $\xi = (\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3$, with a suitable smooth density η .

We remark that it is not really necessary to assume condition (1.1). Indeed, one can easily show by means of a suitable affine-linear change of coordinates that we can always remove the Taylor polynomial of degree 2 of h associated to $y = 0$ from h to reduce the restriction estimates to the cases where the vanishing condition (1.1) holds true, so let us assume this condition henceforth.

Note that if h is of finite type at the origin, then this assumption implies that there is some $m \geq 1$ such that $h^{(m+2)}(0) \neq 0$, i.e., $h(y) = y^{m+2}a(y)$, with $a(0, 0) \neq 0$. This case had already been treated in [BMV19], so what remained open is the case where h is flat at 0, i.e., where $h^{(k)}(0) = 0$ for every $k \in \mathbb{N}$.

Our main result, which generalizes Theorem 1.1 in [BMV19], allows to treat also the latter case under a monotonicity assumption on h''' :

Theorem 1.1. *Assume that $r > 10/3$ and $1/q' > 2/r$, and let \mathcal{E} denote the Fourier extension operator associated to the graph S in (1.2) of the above phase function $\phi(x, y) := xy + h(y)$, where the function h is smooth and satisfies (1.1). Assume further that either the function h is of finite type at the origin, or flat and such that h''' is monotonic. Then, if Ω is a sufficiently small neighborhood of the origin,*

$$\|\mathcal{E}f\|_{L^r(\mathbb{R}^3)} \leq C_{r,q} \|f\|_{L^q(\Omega)}$$

for all $f \in L^q(\Omega)$.

As already mentioned, the novelty of this result lies in the case where h is flat at the origin. Assuming this, we may reduce ourselves to the region where $y > 0$, and the monotonicity assumption then allows us to assume henceforth without loss of generality that $h'''(y) > 0$ on that region.

Our approach does not allow to include also functions h where h''' admits too many oscillations, such as the function φ in Example 2.1.

A treatment of more general smooth perturbations $h(y)$ that are flat at the origin imposes indeed various serious problems, since one seems to lose control over possible bilinear estimates when we have many disjoint intervals on which $h'''(y)$ is of a certain size λ , but in between these intervals $h'''(y)$ is much bigger than λ . It is even possible that the zero set of h''' is a totally disconnected Cantor type set C of positive Lebesgue measure $|C| > 0$, and in this case it is not even clear if any kind of meaningful wave packet decomposition over this set C is still possible. Note that wave packet decompositions play a fundamental role also in other approaches to Fourier restriction, based e.g. on multilinear methods, or polynomial partitioning techniques.

For the proof of Theorem 1.1, we shall build on the methods developed in [BMV19] and [BMV17]. However, serious new obstacles do arise compared to our discussion of the finite type case in [BMV19].

One major difficulty stems from the fact that our interval decomposition result in the subsequent Theorem 2.2 will in fact allow us to decompose the y -region into intervals I_λ on which the third derivative of h is of a fixed dyadic size $\lambda > 0$, but the price that we shall pay is that we loose any reasonable control on higher order derivatives of h compared to the level λ on these subintervals. For the usual wave packet decompositions of $\mathcal{E}f$ this means that our wave packets will possibly no longer be rapidly decaying away from their central tubes - indeed only a rather slow decay can still be guaranteed. For this reason, we shall have to rework the usual bilinear analysis, making use not only of the classical tubes associated to our wave packets, but also of further “hollow tubes” that take into account the contributions by regions far away from the central tube.

A second major problem is that, unlike in the finite type perturbation case, in the case of a flat perturbation function h we shall in general no longer be able to simply sum the contributions given by the subintervals I_λ . To resolve this issue, we shall apply a suitable bootstrap argument in Section 5, which will again make use of bilinear estimates.

CONVENTION: Unless stated otherwise, $C > 0$ will stand for an absolute constant whose value may vary from occurrence to occurrence. We will use the notation $A \sim_C B$ to express that $\frac{1}{C}A \leq B \leq CA$. In some contexts where the size of C is irrelevant we shall drop the index C and simply write $A \sim B$. Similarly, $A \lesssim B$ will express the fact that there is a constant C (which does not depend on the relevant quantities in the estimate) such that $A \leq CB$, and we write $A \ll B$, if the constant C is sufficiently small.

2. A LEVEL-SIZE DECOMPOSITION

Assume that $\varphi : I \rightarrow \mathbb{R}$ is a sufficiently smooth real valued function on some compact interval $I \subset \mathbb{R}$. Our goal in this section will be to show that we can decompose the subset of I on which φ does not vanish into countably many subintervals on each which φ will be comparable in size to some fixed dyadic number λ (depending on the subinterval), with a good control on the number of these subintervals.

If φ is of finite type near some given point, say, the origin, that is, if there is some $m \in \mathbb{N}$ such that $\varphi(0) = \dots = \varphi^{(m-1)}(0) = 0 \neq \varphi^{(m)}(0)$, such a decomposition is quite easily established (at least on a sufficiently small neighborhood of the origin). Indeed, a Taylor expansion shows that $\varphi(x) \sim ax^m$ near the origin, with $a \neq 0$. Even more, we can also control the derivatives of φ :

$$\varphi^{(k)}(x) \sim a_m x^{m-k}$$

for all $k = 0, \dots, m-1$ and all x near 0. Thus, we can simply perform a dyadic decomposition with respect to the origin, and on each of its dyadic subintervals φ will be comparable in size to some fixed dyadic number, and these numbers will essentially be different for different dyadic intervals.

But, if φ is flat at the origin, that is, if $\varphi^{(k)}(0) = 0$ for all $k \in \mathbb{N}$, we have no such easy control, since φ and its derivatives may highly oscillate near the origin. Let us give an example:

Example 2.1. Let $\varphi(x) = \exp(-1/x^2) \sin(1/x)$ and $\lambda > 0$. Let $\epsilon_0 := \log(1/\lambda)^{-1/2}$ (so that $\exp(-1/\epsilon_0^2) = \lambda$.) We are interested in the number of connected component intervals into which the level band set $U_\lambda := \{x \in [0, 1] : \lambda < \varphi(x) < 4\lambda\}$ decomposes. To this end, we consider $x \in U_\lambda \cap [\epsilon, 2\epsilon]$, $\epsilon > \epsilon_0$.

Then $\frac{1}{2\epsilon} \leq \frac{1}{x} \leq \frac{1}{\epsilon}$, that means $\sin(1/x)$ may oscillate $\sim \frac{1}{\epsilon}$ times, hence $U_\lambda \cap [\epsilon, 2\epsilon]$ consists of $\sim \frac{1}{\epsilon}$ intervals. Dyadic summation shows that U_λ consists of $\sim \frac{1}{\epsilon_0} = \sqrt{\log(1/\lambda)}$ intervals.

The example shows that the level band sets U_λ of a flat function may consist of a large number (typically growing with the size of $1/\lambda$) of intervals, though we may hope for some control on this number of intervals. The following theorem does indeed provide such a control.

Theorem 2.2. *Let $I = [a, b]$ be a compact interval and $\varphi \in C^r(I, \mathbb{R})$, $r \geq 1$, and put $C_r := \|\varphi^{(r)}\|_{L^\infty(I)}$. Then there exists a decomposition of $\{\varphi \neq 0\}$ into pairwise disjoint intervals $I_{\lambda, \iota}$, where λ ranges over the set of all positive dyadic numbers $\lambda \leq \|\varphi\|_\infty$, and where for any given λ , the index ι is from some index set \mathcal{I}_λ , such that the following hold true:*

- (i) $|\mathcal{I}_\lambda| \leq 10r(1 + |I|C_r^{1/r}\lambda^{-1/r}) \lesssim 1 + \lambda^{-1/r}$.
- (ii) For any λ dyadic, $\iota \in \mathcal{I}_\lambda$ and any $x \in I_{\lambda, \iota}$ we have $\frac{1}{2}\lambda < |\varphi(x)| < 4\lambda$.

An immediate consequence is the following related result from [SJ 74]:

Corollary 2.3 (Sjölin). *Let $I \subset \mathbb{R}$ be a compact interval and assume that $\varphi \in C^\infty(I, \mathbb{R})$. Denote by E the set of zeros of φ , and by $\{I_k\}_k$ the component intervals of $I \setminus E$. Then for every $\varepsilon > 0$ the series $\sum_k (\sup_{I_k} |\varphi|)^\varepsilon$ is convergent.*

The proof of the theorem will be based on the following auxiliary lemma:

Lemma 2.4. *Let $a > 0$, $b \in \mathbb{R}$, let J be a compact interval and $\varphi \in C^r(J)$, $r \geq 1$. If there exist points $t_0 < \dots < t_r$ in J such that for every $m = 0, \dots, r$*

$$(2.1) \quad (-1)^m (\varphi(t_m) - b) \geq a,$$

then

$$(2.2) \quad |J| > 2 \left(\frac{a}{\|\varphi^{(r)}\|_\infty} \right)^{1/r}.$$

The same conclusion also holds under the assumption $(-1)^{m+1}(\varphi(t_m) - b) \geq a$.

Proof. By the mean value theorem, we find $t_m^1 \in (t_m, t_{m+1})$, $m = 0, \dots, r-1$, with

$$\varphi'(t_m^1)(t_{m+1} - t_m) = \varphi(t_{m+1}) - \varphi(t_m).$$

Then we have

$$(2.3) \quad (-1)^{m+1} \varphi'(t_m^1) = \frac{(-1)^{m+1}(\varphi(t_{m+1}) - b) + (-1)^m(\varphi(t_m) - b)}{t_{m+1} - t_m} \geq \frac{2}{|J|} a.$$

Note that this condition on φ' is similar to our assumption (2.1) on φ , but now with $b = 0$ and a replaced by $2a/|J|$. We can thus iterate this procedure. Formally, the following can be verified by induction:

$$(2.4) \quad (-1)^{m+k} \left(\varphi^{(k)}(t_m^k) - b^{(k)} \right) \geq \left(\frac{2}{|J|} \right)^k a$$

for certain $t_0^k, \dots, t_{r-k}^k \in J$, and where $b^{(0)} = b$, $b^{(k)} = 0$ for $k > 0$. We can keep iterating as long as we can differentiate. Thus, finally we arrive for $k = r$ at the estimate

$$(2.5) \quad \|\varphi^{(r)}\|_\infty \geq (-1)^{m+r} \varphi^{(r)}(t_0^r) \geq \left(\frac{2}{|J|} \right)^r a,$$

which gives (2.2). \square

Proof of Theorem 2.2. We shall show how to decompose the set $\{\varphi > 0\}$ (assuming this set is non-empty); the set $\{\varphi < 0\}$ can then be treated in a similar way. Denote by J_l , $l \in L$, the connected component intervals of $\{\varphi > 0\}$ (L is at most countable). If J_l does not intersect the boundary of I , then we may write $J_l = (a_l, b_l)$, with $a_l < b_l$. Otherwise, if $a \in J_l$, then $\varphi(a) > 0$, and we may write $J_l = [a_l, b_l] = [a, b_l]$. Similarly, if $b \in J_l$, then $\varphi(b) > 0$, and we may write $J_l = (a_l, b_l] = (a_l, b]$.

In a second step, given $l \in L$, we shall further decompose the interval J_l . To this end, we choose a point $c_0^l \in J_l$ so that $\varphi(c_0^l)$ is a dyadic number. This is always possible, unless J_l intersects the boundary and there is some dyadic number λ such that $\lambda < \varphi(x) < 2\lambda$ for every $x \in J_l$. But then J_l does already have the desired property, and since there are at most two intervals J_l of that type, we may ignore these cases in the sequel. Our aim is to construct subintervals of J_l which are as large as possible so that φ will change by a dyadic number on them.

Accordingly, we choose $c_1^l > c_0^l$ maximal in $(a_l, b_l]$ such that $\varphi(c_1^l) \in [\frac{1}{2}\varphi(c_0^l), 2\varphi(c_0^l)]$. Then, recursively, if we have already constructed c_0^l, \dots, c_k^l , we choose $c_{k+1}^l := \max\{x \in (c_k^l, b_l] : \frac{1}{2}\varphi(c_k^l) \leq \varphi(x) \leq 2\varphi(c_k^l)\}$, so that $\varphi(c_{k+1}^l) \in \{\frac{1}{2}\varphi(c_k^l), 2\varphi(c_k^l)\}$, or $c_{k+1}^l = b$. The latter case can only happen if $J_l = (a_l, b]$ and $\varphi(b) > 0$, in which case $\varphi(b)$ may or may not be a dyadic number. If $c_{k+1}^l = b$, we stop the construction; otherwise, we get a countable sequence of points $c_k^l, k \in \mathbb{N}$. We claim that in the latter case, $\lim_{k \rightarrow \infty} c_k^l = b_l$:

Obviously $c_k^l, k \rightarrow \infty$, is a convergent sequence, on which φ assumes only dyadic values, and by construction we have

$$(2.6) \quad \max\{\varphi(c_k^l), \varphi(c_{k+1}^l)\} \leq 2|\varphi(c_k^l) - \varphi(c_{k+1}^l)| \leq 2\|\varphi'\|_\infty |c_k^l - c_{k+1}^l|.$$

Therefore $\lim_{k \rightarrow \infty} \varphi(c_k^l) = 0$, which implies the claim.

We construct points c_k^l with indices $k < 0$ and $c_0^l > c_{-1}^l > c_{-2}^l > \dots$ in an analogous manner by "moving to the left", and obtain that $\lim_{k \rightarrow -\infty} c_k^l = a_l$ in case where we do not reach $a_l = a$ in finitely many steps. In this way we obtain a measure-theoretic disjoint decomposition

$$J_l = \bigcup_{k \in \mathbb{Z}^l} [c_k^l, c_{k+1}^l],$$

where $\mathbb{Z}^l \subset \mathbb{Z}$ is of the form $\mathbb{Z}^l = \{k \in \mathbb{Z} : K_-^l < k < K_+^l\}$, $-\infty \leq K_-^l < K_+^l \leq +\infty$.

Recall that then $\varphi(c_k^l)$ and $\varphi(c_{k+1}^l)$ are dyadic numbers which differ exactly by a factor 2, with the possible exception of the cases where $c_k^l = a$, or $c_{k+1}^l = b$.

We sort the points c_k^l according to which dyadic value $\varphi(c_k^l)$ takes. For bounding the number of component intervals $[c_k^l, c_{k+1}^l]$, we may ignore the intervals at the boundary of I containing a , respectively b , since there are at most two of them. For any dyadic parameter $\lambda > 0$, let

$$(2.7) \quad A(\lambda) := \{c_k^l : \{\varphi(c_k^l), \varphi(c_{k+1}^l)\} = \{\lambda, 2\lambda\}\}.$$

Note that by construction, for any $c_k^l \in A(\lambda)$ and $c_k^l < x < c_{k+1}^l$, we have

$$(2.8) \quad \lambda/2 < \varphi(x) < 4\lambda.$$

Therefore the sets $I_{\lambda,\iota} := [c_k^l, c_{k+1}^l)$, indexed by $\iota = c_k^l \in A(\lambda)$, together with a similar decomposition of $\{\varphi < 0\}$ and possibly two boundary intervals, give the desired decomposition.

For the bound in (i), it is then enough to show that

$$(2.9) \quad |A(\lambda)| \leq 4r(1 + |I|C_r^{1/r}\lambda^{-1/r}).$$

Using (2.6), we see that for $c_k^l \in A(\lambda)$ we have

$$\lambda \leq 2\|\varphi'\|_\infty |c_k^l - c_{k+1}^l|.$$

This implies that $A(\lambda)$ is finite (and in particular, that the number of connected components J_l of I is indeed countable). More precisely, we obtain the bound

$$|A(\lambda)| \leq 2\|\varphi'\|_\infty |I|\lambda^{-1},$$

which already gives the bound in (i) when $r = 1$.

When $r \geq 2$, we have to make use of estimates of the number of oscillations of certain sizes of the function φ and its derivatives and apply Lemma 2.4. To this end, we fix now a dyadic parameter λ , and order the elements of $A(\lambda)$ by $y_1 < \dots < y_M$, where $M := |A(\lambda)|$.

We claim that we have

$$(2.10) \quad \{\varphi(y_m), \varphi(y_{m+1})\} = \{\lambda, 2\lambda\} \quad \text{for all } m = 1, \dots, M-1.$$

We know by (2.7) that the $\varphi(y_m)$ can only take the values λ , or 2λ . Given $y_m = c_k^l$ for some k and l , we assume without loss of generality that $\varphi(y_m) = 2\lambda$, and hence $\varphi(c_{k+1}^l) = \lambda$ (the case where $\varphi(y_m) = \lambda$ can be treated in a similar way).

If we had $\varphi(y_{m+1}) = 2\lambda$, then we would choose $c_{k'}^{l'} > c_{k+1}^l$ minimal such that $\varphi(c_{k'}^{l'}) \geq 2\lambda$. Then necessarily $\varphi(c_{k'-1}^{l'}) < 2\lambda$, and since this is a dyadic number, even $\varphi(c_{k'-1}^{l'}) \leq \lambda$. But then we would have

$$2\lambda \leq \varphi(c_{k'}^{l'}) \leq 2\varphi(c_{k'-1}^{l'}) \leq 2\lambda,$$

so that all the above inequalities must be equalities and hence $c_{k'-1}^{l'} \in A(\lambda)$ while $y_m < c_{k'-1}^{l'} < c_{k'}^{l'} \leq y_{m+1}$, where the three points $y_m, c_{k'-1}^{l'}$ and y_{m+1} are in $A(\lambda)$. This would lead to a contradiction. Therefore $\varphi(y_{m+1}) = \lambda$, which verifies (2.10).

Let us assume for the sake of concreteness that, say, $\varphi(y_1) = \lambda$ (the proof will be quite similar when $\varphi(y_1) = 2\lambda$). Then it is easy to see that (2.10) implies that for every m

$$(-1)^m \left(\varphi(y_m) - \frac{3}{2}\lambda \right) = \frac{\lambda}{2}.$$

In the next step, we organize our sequence $\{y_m\}$ into groups of $r+1$ points: We may assume without loss of generality that $M = |A(\lambda)| \geq 4r$. Let

$$I_j := [y_{j+1}, y_{j+r+1}], \quad j \in (r+1)\mathbb{N}, \quad \frac{j}{r+1} \leq \frac{M}{r+1} - 1.$$

This means we have at least $\frac{M}{r+1} - 1 \geq \frac{M}{4r}$ pairwise disjoint intervals I_j such that in each I_j we have points y_{j+1+m} , $m = 0, \dots, r$, with

$$(-1)^{m+j+1} \left(\varphi(y_{j+1+m}) - \frac{3}{2}\lambda \right) = \frac{\lambda}{2}.$$

We can thus apply Lemma 2.4 and find that

$$|I_j| \geq 2 \left(\frac{\lambda}{2C_r} \right)^{1/r} \geq \left(\frac{\lambda}{C_r} \right)^{1/r}.$$

We conclude that

$$|I| \geq \sum_j |I_j| \geq \frac{M}{4r} \left(\frac{\lambda}{C_r} \right)^{1/r},$$

and hence

$$M \leq 4r |I| C_r^{1/r} \lambda^{-1/r}.$$

□

3. FOURIER RESTRICTION OVER RECTANGLES ON WHICH $h''' \sim \lambda$

In the sequel, we shall always assume that h is flat at the origin. By applying Theorem 2.2 to $\varphi := h'''$, we are led to first restricting ourselves to intervals $I_{\lambda,\epsilon}$ on which we may assume that $h'''(x) \sim \lambda$, where $\lambda > 0$ is a fixed dyadic number. Note also that by choosing Ω sufficiently small in Theorem 1.1, we may assume that

$$(3.1) \quad 0 < \lambda < \lambda_1, \quad \text{where } \lambda_1 \ll 1 \quad \text{is sufficiently small.}$$

Let us therefore now assume that on some interval $I = (c_I - d_I, c_I + d_I)$ of positive length $2d_I > 0$ contained in $(-1, 1)$ we know that

$$h'''(y) \sim \lambda.$$

Then a standard change of coordinates argument (cf. [BMV19]) allows to pass essentially to a situation where $c_I = 0, d_I = 1$, and where (1.1) holds true again. Indeed, write $y = c_I + d_I y'$, and consider $\phi(x, y) = xy + h(y)$. Changing coordinates $y = c_I + d_I y'$, a Taylor expansion of h then leads to

$$\phi(x, y) = x(c_I + d_I y') + h(c_I) + h'(c_I)d_I y' + \frac{h''(c_I)}{2} d_I^2 (y')^2 + d_I H(y'),$$

where $H^{(k)}(0) = 0$ for $k = 0, 1, 2$, and $H'''(y') \sim d_I^2 \lambda =: \epsilon$. Thus

$$\begin{aligned} \phi(x, y)/d_I &= y'(x + \frac{h''(c_I)}{2} d_I y') + H(y') + \text{affine linear terms} \\ &= x' y' + H(y') + \text{affine linear terms}, \end{aligned}$$

if we put $x' := x + \frac{h''(c_I)}{2} d_I y'$. Let us therefore define $\tilde{\phi}(x', y') := x' y' + H(y')$. Then $H'''(y') \sim \epsilon$, where $0 < \epsilon \ll 1$. Thus, if we put $\Omega_I := \Omega \cap \{y \in I\}$ and denote by

$$\mathcal{E}_I f(\xi) = \int_{\Omega_I} f(x, y) e^{-i(\xi_1 x + \xi_2 y + \xi_3 \phi(x, y))} \eta(x, y) dx dy$$

the contribution of the y -Interval I to $\mathcal{E}f$, and define correspondingly

$$\tilde{\mathcal{E}}_I f(\xi) = \int_{(-1, 1)^2} f(x', y') e^{-i(\xi_1 x' + \xi_2 y' + \xi_3 \tilde{\phi}(x', y'))} \tilde{\eta}(x', y') dx' dy',$$

with a suitable cutoff function $\tilde{\eta}$, then an easy scaling argument shows that the following estimates for \mathcal{E}_I and $\tilde{\mathcal{E}}_I$ are equivalent:

$$(3.2) \quad \|\tilde{\mathcal{E}}_I f\|_{L^r} \leq C \|f\|_{L^q};$$

$$(3.3) \quad \|\mathcal{E}_I g\|_{L^r} \leq C d_I^{1-2/r-1/q} \|g\|_{L^q}$$

for all g with $\text{supp } g \subset I$ (and support in x sufficiently small).

It will therefore suffice to prove estimate (3.2), and to this end recall that $H(0) = H'(0) = H''(0) = 0$ and $H'''(y) \sim \epsilon \ll 1$. Taylor expansions then show that $\|H\|_{C^2} \lesssim \epsilon$. We may thus re-write $H = \epsilon h$, i.e.,

$$(3.4) \quad \phi(x, y) = xy + \epsilon h(y), \quad \epsilon \ll 1 \quad ((x, y) \in Q := [-1, 1]^2),$$

with a new function h satisfying the following conditions:

$$(3.5) \quad h(0) = h'(0) = h''(0) = 0 \quad \text{and} \quad C_3/4 \leq h'''(y) \leq C_3 \quad \text{for all } |y| \leq 1,$$

where the constant C_3 will be assumed to be fixed constant $C_3 > 0$. Note that (3.5) implies that

$$\|h\|_{C^2} \lesssim C_3.$$

We like to stress the point that we cannot assume any reasonable control on derivatives of order 4 or higher of h . We therefore call such a function h of y a *perturbation function of coarse cubic type* of the phase xy , in contrast to the notion of a perturbation of cubic type defined by (3.1) in [BMV19], where we had in addition also required a suitable control on higher order derivatives.

A major goal of this paper will be to prove the following uniform restriction estimate for perturbations of coarse cubic type of the parabolic hyperboloid:

Theorem 3.1. *Assume that ϕ is given by (3.4) on the cube Q , where $h(y)$ is a perturbation of coarse cubic type, let S be given as the graph of ϕ . By \mathcal{E} we denote again the Fourier extension operator associated to S . Assume further that $r > 10/3$ and $1/q' > 2/r$. Then*

$$\|\mathcal{E}f\|_{L^r(\mathbb{R}^3)} \leq C_{r,q} \|f\|_{L^q(Q)} \quad \text{for all } f \in L^q(Q),$$

where the constant $C_{r,q}$ may depend on the constant C_3 in (3.5), but not on any further property of h , and not on ϵ .

The proof will be based on a modification of the bilinear method, taking into account the slow decay of wave packets that we are available only in this context. We shall therefore largely follow our approach from [BMV19] only mostly concentrate on those points of the arguments which will require new ideas.

3.1. Admissible pairs of sets U_1, U_2 on which transversalities are of a fixed size.

Recall that the bilinear approach is based on bilinear estimates of the form

$$(3.6) \quad \|\mathcal{E}_{U_1}(f_1) \mathcal{E}_{U_2}(f_2)\|_p \leq C(U_1, U_2) \|f_1\|_2 \|f_2\|_2.$$

Here, \mathcal{E}_{U_1} and \mathcal{E}_{U_2} are the Fourier extension operators associated to patches of sub-surfaces $S_i := \text{graph } \phi|_{U_i} \subset S$, $i = 1, 2$, with $U_i \subset \Omega$. What is crucial for obtaining useful bilinear estimates is that the two patches of surface S_1 and S_2 satisfy certain *transversality conditions*, which are stronger than just assuming that S_1 and S_2 are transversal as hypersurfaces (i.e., that all normals to S_1 are transversal to all normals to S_2). Indeed, what is needed in addition is the following (cf. [BMV17], [L05], [V05], [LV10], or [Be16]):

Denoting by $H\phi$ the Hessian of ϕ , we consider the following quantity

$$(3.7) \quad \tilde{\Gamma}_z^\phi(z_1, z_2, z'_1, z'_2) := \langle (H\phi)^{-1}(z)(\nabla\phi(z_2) - \nabla\phi(z_1)), \nabla\phi(z'_2) - \nabla\phi(z'_1) \rangle.$$

If its modulus is bounded from below by a constant $c > 0$ for all $z_i = (x_i, y_i)$, $z'_i = (x'_i, y'_i) \in U_i$, $i = 1, 2$, $z = (x, y) \in U_1 \cup U_2$, then we have (3.6) for $p > 5/3$, with a constant $C(U_1, U_2)$ that depends only on this constant c and on upper bounds for the derivatives of ϕ . If U_1 and U_2 are sufficiently small (with sizes depending on upper bounds of the first and second order derivatives of ϕ and a lower bound for the determinant of $H\phi$) this condition reduces to the estimate

$$(3.8) \quad |\Gamma_z^\phi(z_1, z_2)| \geq c,$$

for $z_i = (x_i, y_i) \in U_i$, $i = 1, 2$, $z = (x, y) \in U_1 \cup U_2$, where

$$(3.9) \quad \Gamma_z^\phi(z_1, z_2) := \langle (H\phi)^{-1}(z)(\nabla\phi(z_2) - \nabla\phi(z_1)), \nabla\phi(z_2) - \nabla\phi(z_1) \rangle.$$

It is easy to check that for $\phi(x, y) = xy + \epsilon h(y)$, we have

$$(3.10) \quad \Gamma_z^\phi(z_1, z_2) =: 2(y_2 - y_1) \tau_z(z_1, z_2),$$

where

$$(3.11) \quad \tau_z(z_1, z_2) := x_2 - x_1 + \epsilon[h'(y_2) - h'(y_1) - \frac{1}{2}h''(y)(y_2 - y_1)].$$

As in [BMV19], it will be particularly important to look at the expression (3.11) when $z = z_1 \in U_1$, and $z = z_2 \in U_2$, so that the two “transversalities”

$$(3.12) \quad \tau_{z_1}(z_1, z_2) = x_2 - x_1 + \epsilon[h'(y_2) - h'(y_1) - \frac{1}{2}h''(y_1)(y_2 - y_1)]$$

$$(3.13) \quad \tau_{z_2}(z_1, z_2) = x_2 - x_1 + \epsilon[h'(y_2) - h'(y_1) - \frac{1}{2}h''(y_2)(y_2 - y_1)]$$

become relevant. Note the following relation between these quantities:

$$(3.14) \quad \begin{aligned} |\tau_{z_1}(z_1, z_2) - \tau_{z_2}(z_1, z_2)| &= \frac{\epsilon}{2} |h''(y_2) - h''(y_1)| |y_2 - y_1| \sim \epsilon |h'''(\eta)| (y_2 - y_1)^2 \\ &\sim \epsilon (y_2 - y_1)^2, \end{aligned}$$

where η is some intermediate point.

Following Section 3 in [BMV19], we shall try to devise neighborhoods U_1 and U_2 of two given points $z_1^0 = (x_1^0, y_1^0)$ and $z_2^0 = (x_2^0, y_2^0)$ on which these quantities are roughly constant for $z_i = (x_i, y_i) \in U_i$, $i = 1, 2$, and which are also essentially chosen as large as possible. The corresponding pair (U_1, U_2) of neighborhoods of z_1^0 respectively z_2^0 will be called an *admissible pair*. For a motivation of the precise definition of admissible pairs that we shall give in the next subsection, we refer to [BMV19].

3.2. Definition of admissible pairs within $Q \times Q$. To begin with, we fix a large dyadic constant $C_0 \gg 1$.

In a first step, we perform a classical **dyadic decomposition in the y -variable**:

For a given dyadic number $0 < \rho \lesssim 1$, we denote for $j \in \mathbb{Z}$ such that $|j|\rho \leq 1$ by $I_{j,\rho}$ the dyadic interval $I_{j,\rho} := [j\rho, j\rho + \rho)$ of length ρ , and by $V_{j,\rho}$ the corresponding horizontal “strip” $V_{j,\rho} := [-1, 1] \times I_{j,\rho}$ within Q . Given two dyadic intervals J, J' of the same size, we say that they are *related* if their parents are adjacent but they are not adjacent. We divide each dyadic interval J in a disjoint union of dyadic subintervals $\{I_J^k\}_{1 \leq k \leq C_0/8}$, of length $8|J|/C_0$. Then, we define (I, I') to be an *admissible pair of dyadic intervals* if and only if there are J and J' related dyadic intervals and $1 \leq k, j \leq C_0/8$ such that $I = I_J^k$ and $I' = I_{J'}^j$.

We say that a pair of strips $(V_{j_1, \rho}, V_{j_2, \rho})$ is *admissible* and write $V_{j_1, \rho} \smile V_{j_2, \rho}$, if $(I_{j_1, \rho}, I_{j_2, \rho})$ is a pair of admissible dyadic intervals. Notice that in this case,

$$(3.15) \quad C_0/8 < |j_2 - j_1| < C_0/2.$$

One can easily see that this leads to the following disjoint decomposition of $Q \times Q$:

$$(3.16) \quad Q \times Q = \bigcup_{\rho} \left(\bigcup_{V_{j_1, \rho} \smile V_{j_2, \rho}} V_{j_1, \rho} \times V_{j_2, \rho} \right),$$

where the first union is meant to be over all such dyadic ρ 's.

In a second step, we perform a non-standard **Whitney type decomposition of any given admissible pair of strips**, to obtain subregions in which the transversalities are roughly constant.

To simplify notation, we fix ρ and an admissible pair $(V_{j_1, \rho}, V_{j_2, \rho})$, and simply write $I_i := I_{j_i, \rho}$, $V_i := V_{j_i, \rho}$, $i = 1, 2$, so that I_i is an interval of length ρ with left endpoint $j_i \rho$, and

$$(3.17) \quad V_1 = [-1, 1] \times I_1, \quad V_2 = [-1, 1] \times I_2,$$

are rectangles of dimension $2 \times \rho$, which are vertically separated at scale $C_0 \rho$. More precisely, for $z_1 = (x_1, y_1) \in V_1$ and $z_2 = (x_2, y_2) \in V_2$ we have $|y_2 - y_1| \in |j_2 \rho - j_1 \rho| + [-\rho, \rho]$, i.e.,

$$(3.18) \quad C_0 \rho / 2 \leq |y_2 - y_1| \leq C_0 \rho.$$

Let $0 < \delta \lesssim \epsilon^{-1} \rho^{-2}$ be a dyadic number (note that δ could be big, depending on ρ), and let \mathcal{J} be the set of points which partition the interval $[-1, 1]$ into (dyadic) intervals of the same length $\epsilon \rho^2 \delta$.

Similarly, for $i = 1, 2$, we choose a finite equidistant partition \mathcal{I}_i of width $\rho(1 \wedge \delta)$ of the interval I_i by points $y_i^0 \in \mathcal{I}_i$. Note: if $\delta > 1$, then $\rho(1 \wedge \delta) = \rho$, and we can choose for \mathcal{I}_i just the singleton $\mathcal{I}_i = \{y_i^0\}$, where y_i^0 is the left endpoint of I_i .

Definition 3.2. For any parameters $x_1^0, t_2^0 \in \mathcal{J}$, $y_1^0 \in \mathcal{I}_1$ defined in the previous lines and y_2^0 the left endpoint of I_2 , we define the sets

$$(3.19) \quad U_1^{x_1^0, y_1^0, \delta} := \{(x_1, y_1) : 0 \leq y_1 - y_1^0 < \rho(1 \wedge \delta), 0 \leq x_1 - x_1^0 + \epsilon \frac{h''(y_1^0)}{2} (y_1 - y_1^0) < \epsilon \rho^2 \delta\},$$

$$U_2^{t_2^0, y_1^0, y_2^0, \delta} := \{(x_2, y_2) : 0 \leq y_2 - y_2^0 < \rho, \\ 0 \leq x_2 - t_2^0 + \epsilon [h'(y_2) - h'(y_1^0) - \frac{h''(y_1^0)}{2} (y_2 - y_1^0)] < \epsilon \rho^2 \delta\},$$

and the points

$$(3.20) \quad z_1^0 = (x_1^0, y_1^0), \quad z_2^0 = (x_2^0, y_2^0)$$

where

$$x_2^0 := t_2^0 - \epsilon [h'(y_2^0) - h'(y_1^0) - \frac{h''(y_1^0)}{2} (y_2^0 - y_1^0)].$$

Observe that then

$$z_1^0 \in U_1^{x_1^0, y_1^0, \delta} \subset V_1 \quad \text{and} \quad z_2^0 \in U_2^{t_2^0, y_1^0, y_2^0, \delta} \subset V_2.$$

Indeed, z_i^0 is in some sense the “lower left” vertex of U_i , and the horizontal projection of $U_2^{t_2^0, y_1^0, y_2^0, \delta}$ equals I_2 . Note that if we define a^0 by

$$(3.21) \quad a^0 := \tau_{z_1^0}(z_1^0, z_2^0),$$

then $x_1^0 + a^0 = t_2^0$. Notice also that we may re-write

$$(3.22) \quad U_2^{t_2^0, y_1^0, y_2^0, \delta} = \{z_2 = (x_2, y_2) : 0 \leq \tau_{z_1^0}(z_1^0, z_2) - a^0 < \epsilon \rho^2 \delta, 0 \leq y_2 - y_2^0 < \rho\}.$$

In particular, $U_1^{x_1^0, y_1^0, \delta}$ is essentially a parallelepiped of side lengths $\sim \epsilon \rho^2 \delta \times \rho(1 \wedge \delta)$, containing the point (x_1^0, y_1^0) , whose longer side has slope y_1^0 with respect to the y -axis.

Moreover, if $\delta \ll 1$, then $U_2^{t_2^0, y_1^0, y_2^0, \delta}$ is a thin curved box of width $\sim \epsilon \rho^2 \delta$ and length $\sim \rho$, contained in a rectangle of dimension $\sim \rho^2 \times \rho$ whose axes are parallel to the coordinate axes (namely the part of a $\rho^2 \delta$ -neighborhood of a curve of curvature $\sim \epsilon$ containing the point (x_1^0, y_1^0) which lies within the horizontal strip V_2). The case $\delta \ll 1$ will therefore be called the **curved box case**.

If $\delta \gtrsim 1$, then $U_2^{t_2^0, y_1^0, y_2^0, \delta}$ is essentially a rectangular box of dimension $\sim \epsilon \rho^2 \delta \times \rho$ lying in the same horizontal strip. The case $\delta > 1$ will therefore be called the **straight box case**.

Note also that we have chosen to use the parameter t_2^0 in place of using x_2^0 here, since with this choice by (3.11) the identity

$$(3.23) \quad \tau_{z_1^0}(z_1^0, z_2^0) = t_2^0 - x_1^0$$

holds true, which will become quite useful in the sequel. We next have to relate the parameters $x_1^0, t_2^0, y_1^0, y_2^0$ in order to give a precise definition of an admissible pair.

Here, and in the sequel, we shall always assume that the points z_1^0, z_2^0 associated to these parameters are given by (3.20).

Definition 3.3. *Let us call a pair $(U_1^{x_1^0, y_1^0, \delta}, U_2^{t_2^0, y_1^0, y_2^0, \delta})$ an admissible pair of type 1 (at scales δ, ρ and contained in $V_1 \times V_2$), if the following two conditions hold true:*

$$(3.24) \quad \frac{C_0^2}{4} \epsilon \rho^2 \delta \leq |\tau_{z_1^0}(z_1^0, z_2^0)| = |t_2^0 - x_1^0| < 4 C_0^2 \epsilon \rho^2 \delta,$$

$$(3.25) \quad \frac{C_0^2}{512} \epsilon \rho^2 (1 \vee \delta) \leq |\tau_{z_2^0}(z_1^0, z_2^0)| < 5 C_0^2 \epsilon \rho^2 (1 \vee \delta).$$

By \mathcal{P}^δ we shall denote the set of all admissible pairs of type 1 at scale δ (and ρ , contained in $V_1 \times V_2$), and by \mathcal{P} the corresponding union over all dyadic scales δ .

Observe that, by (3.14), we have $\tau_{z_2^0}(z_1^0, z_2^0) - \tau_{z_1^0}(z_1^0, z_2^0) \sim \epsilon(y_2^0 - y_1^0)^2$. In view of (3.24) and (3.18) this shows that condition (3.25) is automatically satisfied, unless $\delta \sim 1$.

We remark that it would indeed be more appropriate to denote the sets \mathcal{P}^δ by $\mathcal{P}_{V_1 \times V_2}^\delta$, but we want to simplify the notation. In all instances in the rest of the paper \mathcal{P}^δ will be associated to a fixed admissible pair of strips (V_1, V_2) , so that our imprecision will not cause any ambiguity. The next lemma can be proved by closely following the arguments in the proof of the corresponding Lemma 2.1 in [BMV17] and just using (3.5).

Lemma 3.4. *If $(U_1^{x_1^0, y_1^0, \delta}, U_2^{t_2^0, y_1^0, y_2^0, \delta})$ is an admissible pair of type 1, then for all $(z_1, z_2) \in (U_1^{x_1^0, y_1^0, \delta}, U_2^{t_2^0, y_1^0, y_2^0, \delta})$,*

$$|\tau_{z_1}(z_1, z_2)| \sim_8 C_0^2 \epsilon \rho^2 \delta \text{ and } |\tau_{z_2}(z_1, z_2)| \sim_{1000} C_0^2 \epsilon \rho^2 (1 \vee \delta).$$

Up to now we focused on the case $|\tau_{z_1}(z_1, z_2)| \lesssim |\tau_{z_2}(z_1, z_2)|$. For the symmetric case, corresponding to the situation where $|\tau_{z_1}(z_1, z_2)| \gtrsim |\tau_{z_2}(z_1, z_2)|$, by interchanging the roles of z_1 and z_2 we define accordingly for any $t_1^0, x_2^0 \in \mathcal{J}$, y_1^0 the left endpoint of I_1 and $y_2^0 \in \mathcal{I}_2$ the sets $\tilde{U}_1^{t_1^0, y_1^0, y_2^0, \delta}$ and $\tilde{U}_2^{x_2^0, y_2^0, \delta}$ in analogy with our discussion in [BMV17], and denote the corresponding admissible pairs $(\tilde{U}_1^{t_1^0, y_1^0, y_2^0, \delta}, \tilde{U}_2^{x_2^0, y_2^0, \delta})$ as *admissible pairs of type 2*. We shall skip the details.

By $\tilde{\mathcal{P}}^\delta$, we shall denote the set of all admissible pairs of type 2 at scale δ (and ρ , contained in $V_1 \times V_2$), and by $\tilde{\mathcal{P}}$ the corresponding unions over all dyadic scales δ .

In analogy with Lemma 3.4, we have

Lemma 3.5. *If $(\tilde{U}_1, \tilde{U}_2) = (\tilde{U}_1^{t_1^0, y_1^0, y_2^0, \delta}, \tilde{U}_2^{x_2^0, y_2^0, \delta}) \in \tilde{\mathcal{P}}^\delta$ is an admissible pair of type 2, then for all $(z_1, z_2) \in (\tilde{U}_1, \tilde{U}_2)$ we have*

$$|\tau_{z_1}(z_1, z_2)| \sim_{1000} C_0^2 \epsilon \rho^2 (1 \vee \delta) \text{ and } |\tau_{z_2}(z_1, z_2)| \sim_8 C_0^2 \epsilon \rho^2 \delta.$$

The crucial bilinear estimates that we shall establish for admissible pairs are given in the following theorem, which extends Theorem 4.3 in [BMV19] to the case of coarse cubic perturbations $h(y)$:

Theorem 3.6. *Let $p > 5/3$, $q \geq 2$. Then, for every admissible pair $(U_1, U_2) \in \mathcal{P}^\delta$ at scale δ , the following bilinear estimates hold true: If $\delta > 1$ and $\epsilon \delta \rho^2 \leq 1$, then*

$$\|\mathcal{E}_{U_1}(f) \mathcal{E}_{U_2}(g)\|_p \leq C_{p,q} (\epsilon \delta \rho^3)^{2(1 - \frac{1}{p} - \frac{1}{q})} \|f\|_q \|g\|_q.$$

If $\delta \leq 1$, then

$$\|\mathcal{E}_{U_1}(f) \mathcal{E}_{U_2}(g)\|_p \leq C_{p,q} (\epsilon \rho^3)^{2(1 - \frac{1}{p} - \frac{1}{q})} \delta^{5 - \frac{3}{q} - \frac{6}{p}} \|f\|_q \|g\|_q.$$

The constants in these estimates are independent of the given admissible pair, of ϵ, ρ and of δ . The same estimates are valid for admissible pairs $(\tilde{U}_1, \tilde{U}_2) \in \tilde{\mathcal{P}}^\delta$ of type 2.

The case $\delta \leq 1$ of this theorem can easily be reduced by means of suitable coordinate transformations to the main result in the next subsection on “prototypical admissible pairs” (compare [BMV19].) Its proof will require new techniques.

The case where $\delta > 1$, which is in some sense more classical and easier to deal with, can be handled by combining the new techniques that we shall explain for the case $\delta \leq 1$ with more classical methods as described in Subsection 4.2 of [BMV19]. We shall leave the details of the treatment of this case to the interested reader.

Given the bilinear estimates of Theorem 3.6, we can then exactly follow our arguments from Section 5 in [BMV19] in order to pass from these bilinear estimates to the linear Fourier extension estimate in Theorem 3.1 and in this way complete the proof of Theorem 3.1. Indeed, one easily checks that all arguments in Section 5 of [BMV19] effectively only require a control on the first three derivatives of F , but not on higher order derivatives, as had already been observed in Section 5 of [BMV19], so that these arguments work as well for perturbations of coarse cubic type h in place of cubic type perturbations.

3.3. A prototypical admissible pair in the curved box case and the crucial scaling transformation. In this section we shall present a “prototypical” case where U_1 and U_2 will form an admissible pair of type 1 centered at $z_1^0 = 0 \in U_1$ and $z_2^0 \in U_2$, with $\epsilon \sim 1, \rho \sim 1$ and $\delta \ll 1$, i.e., $|y_1^0 - y_2^0| \sim 1$, and $|\tau_{z_2^0}(z_1^0, z_2^0)| \sim 1$ but $|\tau_{z_1^0}(z_1^0, z_2^0)| \sim \delta \ll 1$. This means that we shall be in the curved box case.

Fix a small number $0 < c_0 \ll 1$ ($c_0 = 10^{-10}$ will, for instance, work). Assume that $0 < \delta \leq 1/10$, and put

$$(3.26) \quad U_1 := [0, c_0^2 \delta] \times [0, c_0 \delta]$$

$$(3.27) \quad U_2 := \{(x_2, y_2) : 0 \leq y_2 - b < c_0, 0 \leq x_2 + F'(y_2) - a < c_0^2 \delta\},$$

where $|b| \sim_2 1$, $|a| \sim_4 \delta$ and F is a function of *coarse cubic type* in the sense of (3.5), i.e.,

$$(3.28) \quad F(0) = F'(0) = F''(0) = 0 \quad \text{and} \quad C_3/4 \leq F'''(y) \leq C_3 \quad \text{for all } y.$$

Remark. Note that in the case $\epsilon = 1$, if we set $C_0 = 1/c_0$, $\rho = c_0$, then any admissible pair $(U_1, U_2) = (U_1^{0,0,\delta}, U_2^{a,0,b,\delta})$, as in (3.19), would satisfy (3.26) and (3.27) with the above conditions on a and b and suitable F .

Our bilinear result in this prototypical case is as follows:

Theorem 3.7 (prototypical case). *Let $p > 5/3$, and let U_1, U_2 be as in (3.26), (3.27). Assume further that $\phi(x, y) = xy + F(y)$, where F is a real-valued smooth perturbation function of cubic type, i.e., satisfying estimates (3.28), and denote by*

$$\mathcal{E}_{U_i} f(\xi) = \int_{U_i} f(x, y) e^{-i(\xi_1 x + \xi_2 y + \xi_3 \phi(x, y))} \eta(x, y) dx dy, \quad i = 1, 2,$$

the corresponding Fourier extension operators. Then, if the constants c_0 and $\delta \ll 1$ in (3.26), (3.27) are sufficiently small,

$$(3.29) \quad \|\mathcal{E}_{U_1}(f_1), \mathcal{E}_{U_2}(f_2)\|_p \leq C_p \delta^{\frac{7}{2} - \frac{6}{p}} \|f_1\|_2 \|f_2\|_2$$

for every $f_1 \in L^2(U_1)$ and every $f_2 \in L^2(U_2)$, where the constant C_p will only depend on p and the constants C_l in (3.28).

Since according to Lemma 3.4 the transversalities $\tau_{z_1}(z_1, z_2)$ and $\tau_{z_2}(z_1, z_2)$ are of quite different sizes when $\delta \ll 1$, as in [BMV19] we first apply a scaling transformation in x by a factor δ in order to adjust these transversalities to the same level:

We introduce new coordinates (\bar{x}, \bar{y}) by writing $x = \delta \bar{x}$, $y = \bar{y}$, and then re-scale the phase function ϕ by putting

$$\phi^s(\bar{x}, \bar{y}) := \frac{1}{\delta} \phi(\delta \bar{x}, \bar{y}) = \bar{x} \bar{y} + \frac{F(\bar{y})}{\delta}.$$

Denote by U_i^s the corresponding re-scaled domains, i.e.,

$$\begin{aligned} U_1^s &= \{(\bar{x}_1, \bar{y}_1) : 0 \leq \bar{x}_1 < c_0^2, 0 \leq \bar{y}_1 < c_0 \delta\}, \\ U_2^s &= \{(\bar{x}_2, \bar{y}_2) : 0 \leq \bar{x}_2 + \frac{F'(\bar{y}_2)}{\delta} - \bar{a} < c_0^2, 0 \leq \bar{y}_2 - \bar{b} < c_0\}, \end{aligned}$$

where c_0 is small and $|\bar{a}| = |a/\delta| \sim 1$ and $\bar{b} = b \sim 1$. By $S_i^s, i = 1, 2$, we denote the corresponding scaled surface patches

$$S_i^s := \{(\bar{x}, \bar{y}, \phi^s(\bar{x}, \bar{y})) : (\bar{x}, \bar{y}) \in U_i^s\}.$$

Observe that

$$\nabla\phi^s(\bar{x}, \bar{y}) = (\bar{y}, \bar{x} + F'(\bar{y})/\delta),$$

and

$$(3.30) \quad H\phi^s(\bar{x}, \bar{y}) = \begin{pmatrix} 0 & 1 \\ 1 & F''(\bar{y})/\delta \end{pmatrix},$$

so that in particular

$$(3.31) \quad |\nabla\phi^s(\bar{z})| \lesssim 1$$

for all $\bar{z} \in U_1^s \cup U_2^s$.

Assume next that $\bar{z}_1 \in U_1^s$ and $\bar{z}_2 \in U_2^s$. Since $|\bar{y}_1| \leq c_0\delta$, $|\bar{y}_2| \sim 1$, we see that

$$(3.32) \quad \begin{cases} |\frac{F'(\bar{y}_1)}{\delta}| \sim \frac{|F'''(\eta_1)\bar{y}_1^2|}{\delta} \lesssim C_3 c_0^2 \frac{\delta^2}{\delta} = c_0^2 C_3 \delta, & \frac{|F''(\bar{y}_1)|}{\delta} \sim \frac{|F'''(\tilde{\eta}_1)\bar{y}_1|}{\delta} \lesssim c_0 C_3, \\ |\frac{F'(\bar{y}_2)}{\delta}| \sim \frac{|F'''(\eta_2)\bar{y}_2^2|}{\delta} \sim \frac{C_3}{\delta}, & \frac{|F''(\bar{y}_2)|}{\delta} \sim \frac{|F'''(\tilde{\eta}_2)\bar{y}_2|}{\delta} \sim \frac{C_3}{\delta} \end{cases}$$

(for suitable choices of intermediate points $\eta_i, \tilde{\eta}_i$). Moreover, we then also see that

$$(3.33) \quad \nabla\phi^s(\bar{z}_2) - \nabla\phi^s(\bar{z}_1) = (\bar{y}_2 - \bar{y}_1, \bar{x}_2 + \frac{F'(\bar{y}_2)}{\delta} - (\bar{x}_1 + \frac{F'(\bar{y}_1)}{\delta})) = (\bar{b}, \bar{a}) + \mathcal{O}(c_0).$$

Following further on the proof of Lemma 2.3 in [BMV17], assume that we translate the two patches of surface S_1^s and S_2^s in such a way that the two points \bar{z}_1 and \bar{z}_2 coincide after translation, and assume that the vector $\omega = (\omega_1, \omega_2)$ is tangent to the corresponding intersection curve $\gamma(t)$ at this point. Then (3.33) shows that we may assume without loss of generality that

$$(3.34) \quad \omega = (-\bar{a}, \bar{b}) + \mathcal{O}(c_0).$$

In combination with (3.32) this implies that

$$H\phi^s(\bar{z}_i) \cdot {}^t\omega = \begin{pmatrix} 0 & 1 \\ 1 & F''(\bar{y}_i)/\delta \end{pmatrix} \begin{pmatrix} -\bar{a} + \mathcal{O}(c_0) \\ \bar{b} + \mathcal{O}(c_0) \end{pmatrix}.$$

Thus, if $i = 1$, then by (3.32),

$$(3.35) \quad H\phi^s(\bar{z}_1) \cdot {}^t\omega = \begin{pmatrix} \bar{b} + \mathcal{O}(c_0) \\ -\bar{a} + \mathcal{O}(c_0) \end{pmatrix} \quad \text{and} \quad |H\phi^s(\bar{z}_1) \cdot {}^t\omega| \sim 1,$$

and if $i = 2$, then

$$(3.36) \quad H\phi^s(\bar{z}_2) \cdot {}^t\omega = \begin{pmatrix} \bar{b} + \mathcal{O}(c_0) \\ -\bar{a} + \bar{b}F''(\bar{y}_2)/\delta + \mathcal{O}(c_0)/\delta \end{pmatrix} \quad \text{and} \quad |H\phi^s(\bar{z}_1) \cdot {}^t\omega| \sim 1/\delta,$$

if $\delta \ll 1$ is sufficiently small.

Following [BMV17], the refined transversalities that we need to control in order to apply the bilinear method in the next section are given by

$$(3.37) \quad |TV_i^s(\bar{z}_1, \bar{z}_2)| := \left| \frac{\det({}^t(\nabla\phi^s(\bar{z}_1) - \nabla\phi^s(\bar{z}_2)), H\phi^s(\bar{z}_i) \cdot {}^t\omega)}{\sqrt{1 + |\nabla\phi^s(\bar{z}_1)|^2} \sqrt{1 + |\nabla\phi^s(\bar{z}_2)|^2} |H\phi^s(\bar{z}_i) \cdot {}^t\omega|} \right|, \quad i = 1, 2.$$

But, if $i = 1$, then by (3.33), (3.36), (3.31) and (3.32) we see that

$$|\det({}^t(\nabla\phi^s(\bar{z}_1) - \nabla\phi^s(\bar{z}_2)), H\phi^s(\bar{z}_1) \cdot {}^t\omega)| = \left| \det \begin{pmatrix} \bar{b} + \mathcal{O}(c_0) & \bar{b} + \mathcal{O}(c_0) \\ \bar{a} + \mathcal{O}(c_0) & -\bar{a} + \mathcal{O}(c_0) \end{pmatrix} \right| \sim 1,$$

hence $|TV_1^s(\bar{z}_1, \bar{z}_2)| \sim 1$.

And, if $i = 2$, then by (3.33), (3.35), (3.31) and (3.32) we have

$$|\det({}^t(\nabla\phi^s(\bar{z}_1) - \nabla\phi^s(\bar{z}_2)), H\phi^s(\bar{z}_2) \cdot {}^t\omega)| = \left| \det \begin{pmatrix} \bar{b} + \mathcal{O}(c_0) & \bar{b} + \mathcal{O}(c_0) \\ \bar{a} + \mathcal{O}(c_0) & -\bar{a} + \bar{b}F''(\bar{y}_2)/\delta + \mathcal{O}(c_0)/\delta \end{pmatrix} \right|,$$

hence also $|TV_2^s(\bar{z}_1, \bar{z}_2)| \sim (1/\delta)/(1/\delta) \sim 1$, provided δ and c_0 are sufficiently small.

We have thus proved the following lemma:

Lemma 3.8. *The transversalities for the scaled patches of surface S_i^s , $i = 1, 2$, satisfy*

$$|TV_i^s(\bar{z}_1, \bar{z}_2)| \sim 1, \quad i = 1, 2.$$

In the next section, we shall establish the crucial bilinear Fourier extension estimates for the pair of scaled patches of surface S_1^s and S_2^s .

4. THE BILINEAR METHOD WITH SLOWLY DECAYING WAVE PACKETS FOR THE RE-SCALED PROTOTYPICAL CASE

First, recall from the previous Subsection 3.3 that we had passed from our original coordinates (x, y) to the coordinates (\bar{x}, \bar{y}) by means of the scaling transformation $(x, y) = A(\bar{x}, \bar{y}) = (\delta\bar{x}, \bar{y})$, and had put $\phi^s(\bar{z}) := \phi(A\bar{z})/\mathfrak{a}$, with $\mathfrak{a} := \det A = \delta$.

By means of simple scaling argument (compare Subsection 3.1 of [BMV17]), estimate (3.29) is equivalent the following bilinear Fourier extension estimate for the scaled patches of surface S_i^s which were defined as the graphs of ϕ^s over the sets $U_i^s, i = 1, 2$:

$$(4.1) \quad \|\mathcal{E}_{U_1^s}(f_1) \mathcal{E}_{U_2^s}(f_2)\|_p \leq C_p \delta^{\frac{5}{2} - \frac{4}{p}} \|f_1\|_2 \|f_2\|_2,$$

for every $f_1 \in L^2(U_1^s)$ and every $f_2 \in L^2(U_2^s)$.

In order to defray our notation, let us in the sequel drop the superscripts s , and write simply U_i and S_i in place of U_i^s and S_i^s , ϕ in place of ϕ^s , and denote the coordinates (\bar{x}, \bar{y}) again by (x, y) , etc.. I.e., we assume that

$$\phi(x, y) = xy + \frac{F(y)}{\delta},$$

where F is a perturbation function of coarse cubic type in the sense of (3.28), i.e.,

$$F(0) = F'(0) = F''(0) = 0 \quad \text{and} \quad C_3/4 \leq F'''(y) \leq C_3 \quad \text{for all } y,$$

and that

$$(4.2) \quad \begin{aligned} U_1 &= \{(x_1, y_1) : 0 \leq x_1 < c_0^2, 0 \leq y_1 < c_0\delta\}, \\ U_2 &= \{(x_2, y_2) : 0 \leq x_2 + \frac{F'(y_2)}{\delta} - a < c_0^2, 0 \leq y_2 - b < c_0\}, \end{aligned}$$

where c_0 is assumed to be sufficiently small and $|a| \sim 1$ and $b \sim 1$. The corresponding transversalities will then be denoted by $TV_i(z_1, z_2)$, where $z_i = (x_i, y_i)$, $i = 1, 2$.

Note that for $z_1 \in U_1$ and $z_2 \in U_2$, and $k = 0, 1, 2, 3$, we have $|F^{(k)}(y_1)| \sim |y_1|^{3-k} \lesssim \delta^{3-k}$, whereas $|F''(y_2)| \sim 1$.

In view of (3.30), (3.31) and Lemma 3.8, the following properties of $\nabla\phi$, the operator norm of the Hessian matrix $H\phi$, and of the transversalities of ϕ are immediate or easy to verify:

For all $z_1 \in U_1$, $z_2 \in U_2$, we have

$$(4.3) \quad |\nabla \phi(z_i)| \lesssim 1 \quad i = 1, 2;$$

$$(4.4) \quad |TV_i(z_1, z_2)| \sim 1, \quad i = 1, 2;$$

$$(4.5) \quad \|H(\phi)(z_1)\| \sim 1, \quad \|H(\phi)(z_2)\| \sim 1/\delta;$$

$$(4.6) \quad |\partial_1^j \partial_2^k \phi(z_1)| \lesssim \delta^{2-j-k}, \quad |\partial_1^j \partial_2^k \phi(z_2)| \lesssim \delta^{-1}, \quad \text{if } k \leq 3, j+k \geq 2.$$

Note that the last estimates can in some cases be improved, but they will be sufficient for our purposes. With this modified notation, our major goal in this section will be to prove the following

Theorem 4.1. *Under the preceding assumptions on ϕ and U_1, U_2 , the following bilinear estimate holds true uniformly for every $f_1 \in L^2(U_1)$ and every $f_2 \in L^2(U_2)$:*

$$\|\mathcal{E}_{U_1}(f_1) \mathcal{E}_{U_2}(f_2)\|_p \leq C_p \delta^{\frac{5}{2} - \frac{4}{p}} \|f_1\|_2 \|f_2\|_2.$$

Our basic approach to this theorem will follow the lines in our preceding article [BMV16], which in turn is following the structure in Lee's article [L05]. We shall explain in more detail those parts where new ideas are needed, and shall be brief about those arguments which are essentially the same as in the classical bilinear method. For further details on the latter parts, we refer the reader to [L05], [V05], and also [T03].

4.1. Slowly decaying wave packets. As a first step, we need to introduce a modified version of the so-called wave packet decomposition. Wave packet decompositions have become by now a standard tool for the study of problems in harmonic analysis related to geometric properties of submanifolds. In those applications, hitherto one always had a good control over all derivatives (up to a fixed, but arbitrary order) of the phase functions arising in the given contexts.

In our present setting, however, we have to deal with phase functions $\phi(x, y) = xy + F(y)/\delta$, where F only satisfies the conditions in (3.28). In particular, we cannot assume any control over derivatives of F of order 4 or higher! Consequently, our “wave packets” will lack a rapid decay away from the axes of the tubes arising in the wave packet decomposition. Let us briefly discuss how much we can salvage from the usual wave packet decomposition, and where we need modifications.

For a point $(\xi, \tau) \in \mathbb{R}^2 \times \mathbb{R}$, we will write $\xi = (\xi^1, \xi^2)$, to distinguish from subscripts referring to S_1 and S_2 .

In the next lemma, we shall derive a slowly decaying wave packet decomposition for more general classes of phase functions ϕ and corresponding surfaces given as the graph of ϕ over an open and bounded subset $U \subset \mathbb{R}^2$, under suitable conditions on derivatives of ϕ . By

$$\mathcal{E}_U f(\xi, \tau) = \int_U f(x, y) e^{-i(\xi^1 x + \xi^2 y + \tau \phi(x, y))} \alpha(x, y) dx dy,$$

we denote again the corresponding Fourier extension operator, where $\alpha \in C_0^\infty$ is a suitable amplitude.

In Subsection 4.2, we shall later apply this lemma directly to S_1 , with $\kappa := 1$, which is possible in view of (4.3) – (4.6). Note here that for $R > 1/\delta$ the set U'_1 has essentially the same dimension and shape as U_1 , so that (4.3)–(4.6) can be assumed to hold true even on the

larger set U'_1 . The lemma could as well be applied to S_2 , with $\kappa := 1/\delta$, since again U'_2 has essentially the same dimension and shape as U_2 . However, as for S_2 , we shall in Subsection 4.2 first re-parametrize S_2 in order to match our wave packet decomposition with some problems arising in the induction on scales argument, and then apply the lemma in the new coordinates.

Lemma 4.2. *Let $\delta > 0$ and $\kappa \geq 1$. Let $U \subset \mathbb{R}^2$ be an open and bounded subset and let $U' := U + \mathcal{O}(\delta)$ be suitable thickening of the set U of order δ . Assume further that ϕ a smooth phase function defined on U' such that*

$$(4.7) \quad \|H\phi(z)\| \sim \kappa \geq 1 \text{ and } |\partial_1^j \partial_2^k \phi(z)| \lesssim (\delta\kappa)^{3-k-j} \delta^{-1}$$

for every $z \in U'$ and $k \leq 3$, $j + k \geq 2$. Let us also assume that $\partial_1^2 \phi \equiv 0$.

Then for every $R \geq 1/\delta$ the function $\mathcal{E}_U f$ can be decomposed into slowly decaying wave packets adapted to ϕ with central tubes of radius R and length R^2/κ . More precisely, consider the index sets $\mathcal{Y} := R\mathbb{Z}^2$ and $\mathcal{V} := R^{-1}\mathbb{Z}^2 \cap U'$, and define for $w = (\eta, v) \in \mathcal{Y} \times \mathcal{V} =: \mathcal{W}$ the tube

$$(4.8) \quad T_w = \{(\xi, \tau) \in \mathbb{R}^2 \times \mathbb{R} : |\xi - \eta + \tau \nabla \phi(v)| \leq R, |\tau| \leq R^2/\kappa\}.$$

Then there exist functions (“slowly decaying wave packets”) p_w and coefficients $c_w \in \mathbb{C}$, $w \in \mathcal{W}$, such that $\mathcal{E}_U f$ can be decomposed into

$$\mathcal{E}_U f(\xi, \tau) = \sum_{w \in \mathcal{W}} c_w p_w(\xi, \tau)$$

for every $\tau \in \mathbb{R}$ with $|\tau| \leq R^2/\kappa$, $\xi = (\xi^1, \xi^2) \in \mathbb{R}^2$, in such a way that the following hold true:

- (P1) $p_w = \mathcal{E}_U(\mathcal{F}^{-1}(p_w(\cdot, 0)))$.
- (P2) $\text{supp } \mathcal{F}p_w \subset B((v, \phi(v)), 2/R)$.
- (P3) p_w decays away from T_w as follows:

$$|p_w(\xi, \tau)| \leq CR^{-1} \left(1 + \frac{|\xi^1 - \eta^1 + \tau \partial_1 \phi(v)|}{R}\right)^{-2} \left(1 + \frac{|\xi^2 - \eta^2 + \tau \partial_2 \phi(v)|}{R}\right)^{-2}.$$

In particular, $\|p_w(\cdot, \tau)\|_2 \lesssim 1$.

- (P4) For all subsets $W \subset \mathcal{W}$, we have $\|\sum_{w \in W} p_w(\cdot, \tau)\|_2 \lesssim |W|^{\frac{1}{2}}$.

- (P5) $\|(c_w)_w\|_{\ell^2} \lesssim \|f\|_{L^2}$.

Remarks 4.3. (a) Observe that property (P3) implies the weaker estimate

$$(4.9) \quad |p_w(\xi, \tau)| \leq CR^{-1} \left(1 + \frac{\text{dist}((\xi, \tau), T_w)}{R}\right)^{-2},$$

which means that the mass of p_w is in a certain weak sense mostly localized to T_w (however, the decay outside T_w is now much slower than classically). This estimate will obviously hold true later in Subsection 4.2 for the wave packets associated to $\mathcal{E}_{U_1} f_1$, but as well for those associated to $\mathcal{E}_{U_2} f_2$, and in the last parts of the proof where interactions between wave packets of type 1 with wave packets of type 2 will become relevant we can just rely on these weaker estimates (4.9) in place of the more refined ones given by (P3).

(b) The condition $\partial_1^2 \phi \equiv 0$ will be satisfied in our applications and simplifies the integrations by parts argument in the proof a lot, but is quite surely not necessary. We shall, however, not dwell on this here.

Proof. We shall closely follow our preceding paper [BMV16] and mainly focus on those parts of the proof which will require new arguments.

The initial construction remains the usual one, as in (for instance) [L05]: Let $\psi, \hat{\eta} \in C_0^\infty(B(0, 1))$ be chosen in a such a way that for $\psi_v(z) := \psi(R(z - v))$ and $\chi_\eta(\xi) := \chi(\frac{\xi - \eta}{R})$, we have $\sum_{v \in \mathcal{V}} \psi_v = 1$ on U and $\sum_{\eta \in \mathcal{Y}} \chi_\eta = 1$. We also chose a slightly bigger function $\tilde{\psi} \in C_0^\infty(B(0, 3))$ such that $\tilde{\psi} = 1$ on $B(0, 2) \supset \text{supp } \psi + \text{supp } \hat{\eta}$, and put $\tilde{\psi}_v(z) := \tilde{\psi}(R(z - v))$, where we again assume that $z = (x, y)$. Then the functions

$$F_{(\eta, v)} := \mathcal{F}^{-1}(\widehat{\psi_v f \chi_\eta}) = (\psi_v f) * \check{\chi}_\eta, \quad \eta \in \mathcal{Y}, v \in \mathcal{V},$$

are essentially well-localized in both position and momentum/frequency space and supported in balls of radius $\sim 1/R$. Note also that f supported in U we have $\mathcal{E}_U f = \mathcal{E} f$. Define then

$$q_w := \mathcal{E}(F_w), \quad w = (\eta, v) \in \mathcal{W};$$

up to a certain factor c_w , which will be determined later, these are already the announced wave packets, i.e., $q_w = c_w p_w$.

We may also assume for simplicity that the amplitude in the definition of the extension operator \mathcal{E} is identically 1 on the sets U_i . Since $f = \sum_{w \in W} F_w$, we then have the decomposition $\mathcal{E}_U f = \sum_{w \in W} q_w$. Let us concentrate on property (P3) - the other properties are then rather easy to establish. It is easy to see that we have for every $w \in \mathcal{W}$

$$q_w(\xi, \tau) = (2\pi)^{-2} R^{-2} \int K(\xi - \zeta, \tau) \widehat{F}_w(\zeta) d\zeta,$$

with the kernel

$$K(\xi, \tau) := \int e^{i(\xi(\frac{z}{R} + v) + \tau\phi(\frac{z}{R} + v))} \tilde{\psi}(z) \alpha(\frac{z}{R} + v) dz.$$

To simplify the proof a bit, let us pretend in the sequel that $\alpha(\frac{z}{R} + v) \equiv 1$ on the support of $\tilde{\psi}$; since all derivatives of $\alpha(\frac{z}{R} + v)$ are uniformly bounded on the support of $\tilde{\psi}$, we shall see that the presence of the factor $\alpha(\frac{z}{R} + v)$ has indeed no effect on the subsequent integration by parts arguments.

We shall show that

$$(4.10) \quad |K(\xi, \tau)| \lesssim \left(1 + \frac{|\xi^1 + \tau \partial_1 \phi(v)|}{R}\right)^{-2} \left(1 + \frac{|\xi^2 + \tau \partial_2 \phi(v)|}{R}\right)^{-2}.$$

To this end, we may clearly assume that

$$(4.11) \quad |\xi^1 + \tau \partial_1 \phi(v)| \gg R,$$

or

$$(4.12) \quad |\xi^2 + \tau \partial_2 \phi(v)| \gg R.$$

Let us start with the case where both conditions hold true. We first perform integrations by parts with respect to the first component x of $z = (x, y)$ in the oscillatory integral

$$K := \int e^{i\Phi(z)} \tilde{\psi}(z) dz,$$

with phase

$$\Phi(z) := \xi\left(\frac{z}{R} + v\right) + \tau\phi\left(\frac{z}{R} + v\right).$$

Recalling that $\partial_1^2\phi \equiv 0$, hence $\partial_1^2\Phi(z) \equiv 0$, we see that

$$K = \int i e^{i\Phi(z)} \partial_1 \frac{\tilde{\psi}(z)}{\partial_1 \Phi(z)} dz = i \int e^{i\Phi(z)} \frac{\partial_1 \tilde{\psi}(z)}{\partial_1 \Phi(z)} dz.$$

Repeating this argument N times, we obtain

$$(4.13) \quad K = i^N \int e^{i\Phi(z)} \frac{\partial_1^N \tilde{\psi}(z)}{[\partial_1 \Phi(z)]^N} dz.$$

To shorten the notation in the subsequent computation, let us put $A(z) := \frac{i^N \partial_1^N \tilde{\psi}(z)}{[\partial_1 \Phi(z)]^N}$.

Integrating by parts twice now with respect to the second component y of $z = (x, y)$, we obtain

$$\begin{aligned} K &= \int i e^{i\Phi(z)} \partial_2 \frac{A(z)}{\partial_2 \Phi(z)} dz \\ &= - \int i e^{i\Phi(z)} \left(\frac{\partial_2^2 \Phi(z)}{(\partial_2 \Phi(z))^2} A(z) - \frac{\partial_2 A(z)}{\partial_2 \Phi(z)} \right) dz \\ &= \int e^{i\Phi(z)} \partial_2 \left(\frac{\partial_2^2 \Phi(z)}{(\partial_2 \Phi(z))^3} A(z) - \frac{\partial_2 A(z)}{(\partial_2 \Phi(z))^2} \right) dz \\ &= \int e^{i\Phi(z)} \left(\frac{\partial_2^3 \Phi(z)}{(\partial_2 \Phi(z))^3} A(z) - 3 \frac{(\partial_2^2 \Phi(z))^2}{(\partial_2 \Phi(z))^4} A(z) + 3 \frac{\partial_2^2 \Phi(z)}{(\partial_2 \Phi(z))^3} \partial_2 A(z) - \frac{\partial_2^2 A(z)}{(\partial_2 \Phi(z))^2} \right) dz \\ &= \int e^{i\Phi(z)} \left(\frac{\partial_2^3 \Phi(z)}{(\partial_2 \Phi(z))^3} A(z) - 3 \frac{(\partial_2^2 \Phi(z))^2}{(\partial_2 \Phi(z))^4} A(z) + 3 \frac{\partial_2^2 \Phi(z)}{(\partial_2 \Phi(z))^3} \partial_2 A(z) \right) dz \\ &\quad - \int e^{i\Phi(z)} \frac{\partial_2^2 A(z)}{(\partial_2 \Phi(z))^2} dz. \end{aligned}$$

For the first term, we stop integrating by parts in y , since we have no control of derivatives of Φ of order higher than 3. However, the second term can be integrated by parts once more.

Notice also that for all $j, k \in \mathbb{N}$ such that $k \leq 3$ and $j + k \geq 2$ we have

$$|\partial_1^j \partial_2^k \Phi(z)| \lesssim \frac{|\tau|(\delta\kappa)^{3-j-k}}{\delta R^{j+k}} \leq (\delta\kappa R)^{2-j-k} \leq 1,$$

since we assume that $|\tau|\kappa \leq R^2$, $\delta\kappa R \geq \delta R \geq 1$. Therefore, we see that for every $N \in \mathbb{N}$

$$(4.14) \quad |K| \lesssim C_N \int_{\text{supp } \tilde{\psi}} |\partial_1 \Phi(z)|^{-N} |\partial_2 \Phi(z)|^{-3} dz.$$

Observe next that for $|\tau| \leq R^2/\kappa$ and $z \in \text{supp } \tilde{\psi}$, in view of (4.7) we have for $j = 1, 2$

$$|\tau \partial_2 \phi(z/R + v) - \tau \partial_j \phi(v)| \lesssim |\tau| (|\partial_1 \partial_j \phi| + |\partial_2 \partial_j \phi|) \frac{|z|}{R} \lesssim \frac{R^2}{\kappa} \cdot \kappa \cdot \frac{1}{R} \lesssim R.$$

In view of (4.11), (4.12), these estimates imply that for $j = 1, 2$,

$$|\partial_j \Phi(z)| = |\xi^j/R + \tau \partial_j \phi(z/R + v)/R| \gtrsim |\xi^j + \tau \partial_j \phi(v)|/R,$$

so we obtain (4.10) for this case.

If only one of the conditions (4.11) or (4.12) holds, we only integrate by parts in one of the variables and arrive again at (4.10).

Following the proof in [L05], we conclude that

$$\begin{aligned}
|q_w(\xi, \tau)| &\lesssim R^{-2} \int \left| K(\xi - \zeta - \eta, \tau) \widehat{F}_w(\zeta + \eta) \right| d\zeta \\
&= R^{-2} \int \left| K(\xi - \zeta - \eta, \tau) \chi\left(\frac{\zeta}{R}\right) \widehat{\psi_v f}(\zeta + \eta) \right| d\zeta \\
&\lesssim M(\widehat{\psi_v f})(\eta) \left(1 + \frac{|\xi^1 - \eta^1 + \tau \partial_1 \phi(v)|}{R}\right)^{-2} \left(1 + \frac{|\xi^2 - \eta^2 + \tau \partial_2 \phi(v)|}{R}\right)^{-2},
\end{aligned}$$

where M denotes the Hardy-Littlewood maximal operator. Thus, by choosing $c_w = c_{(\eta, v)} := RM(\widehat{\psi_v f})(\eta)$, we obtain (P3).

Properties (P1) and (P2) follow from the definition of the wave packets as usual.

To prove (P4), as a first step note that it is easily seen by Plancherel's theorem and (P2) that

$$\left\| \sum_{w \in W} p_w(\cdot, \tau) \right\|_2 \lesssim \left(\sum_v \left\| \sum_{\eta: (\eta, v) \in W} p_{(\eta, v)}(\cdot, \tau) \right\|_2^2 \right)^{1/2}.$$

As we lack fast decay in (P3), we need to be a little more careful in establishing the orthogonality in the y -parameter in (P4). We have to show that for any $Y \subset \mathcal{Y}$

$$\left\| \sum_{\eta \in Y} p_{(\eta, v)}(\cdot, \tau) \right\|_2 \lesssim |Y|^{1/2}.$$

By a linear transformation, we may assume that $v = 0$. Then

$$\begin{aligned}
\left\| \sum_{\eta \in Y} p_{(\eta, v)}(\cdot, \tau) \right\|_2^2 &= \sum_{\eta, \eta' \in Y} \int p_{\eta, v}(\xi, \tau) \bar{p}_{\eta', v}(\xi, \tau) d\xi \\
&\leq CR^{-2} \sum_{\eta, \eta' \in Y} \prod_{i=1}^2 \int \left(1 + \frac{|\xi^i - \eta^i|}{R}\right)^{-2} \left(1 + \frac{|\xi^i - \eta'^i|}{R}\right)^{-2} d\xi^i \\
&= C \sum_{\substack{k, k' \in \mathbb{Z}^2: \\ Rk, Rk' \in Y}} \prod_{i=1}^2 \int (1 + |\xi^i - k^i|)^{-2} (1 + |\xi^i - k'^i|)^{-2} d\xi^i \\
&\leq C \sum_{\substack{k \in \mathbb{Z}^2: \\ Rk \in Y}} \prod_{i=1}^2 \int (1 + |\xi^i - k^i|)^{-2} \sum_{k'_i \in \mathbb{Z}} (1 + |\xi^i - k'^i|)^{-2} d\xi^i \\
&\lesssim |Y|.
\end{aligned}$$

For (P5), we refer to [L05]. □

4.2. Proof of Theorem 4.1 - Setup of the argument. Due to a by now standard argument, it is sufficient to prove the following local estimate: For any $p > 5/3$ and $\epsilon > 0$, there exist a constant $C_{p,\epsilon}$ such that

$$(4.15) \quad \|\mathcal{E}_{U_1}(f_1), \mathcal{E}_{U_2}(f_2)\|_{L^p(Q(R'))} \leq C_{p,\epsilon} R'^\epsilon \delta^{\frac{5}{2}-\frac{4}{p}} \|f_1\|_2 \|f_2\|_2$$

for all $f_1 \in L^2(U_1)$, $f_2 \in L^2(U_2)$ and $R' \gg 1$, where $Q(R')$ is a cuboid that we will choose below.

As already mentioned in Subsection 4.1, we can apply the wave packet decomposition from Lemma 4.2 directly to $\mathcal{E}_{U_1}f_1$, with $\kappa = 1$. The wave packets associated to $\mathcal{E}_{U_1}f_1$ and the patch of hypersurface S_1 are mainly concentrated on tubes $T_{w_1, w_1} = (\eta_1, v_1) \in \mathcal{W}_1$, which are “horizontal” translates of tubes of the form

$$T_{(v_1)} := \{(\xi, \tau) : |\xi + \tau \nabla \phi(v_1)| \leq R, |\tau| \leq R^2\},$$

but, in contrast to the classical situation, they do not decay rapidly away from those tubes. Notice the standard fact that $T_{(v_1)}$ is a tube of dimension $R \times R \times R^2$ whose long axis is pointing in the direction of the normal vector $N(v_1) := (\nabla \phi(v_1), -1)$ to S_1 at the point $(v_1, \phi(v_1))$, $v_1 \in U_1$.

Note next that the projections of tubes T_{w_1} to ξ -space are contained in a $\delta R^2 \times R^2$ -rectangle. Therefore we define $Q(R')$ to be the $\delta R^2 \times R^2 \times R^2 = \frac{R'^2}{\delta} \times \frac{R'^2}{\delta^2} \times \frac{R'^2}{\delta^2}$ cuboid centered at the origin. Then for any $(\xi, \tau) \in Q(R')$, we have $|\tau| \leq R^2$, so we can decompose

$$(4.16) \quad \mathcal{E}_{U_1}f_1(\xi, \tau) = \sum_{w_1 \in \mathcal{W}_1} c_{w_1} p_{w_1}(\xi, \tau).$$

As for the wave packets associated to $\mathcal{E}_{U_2}f_2$ and the patch of hypersurface S_2 , these should then mainly be concentrated on shorter tubes T_{w_2} , of length δR^2 in order to fit well into the cuboid $Q(R')$. Moreover, we would like to translate them not horizontally, but in directions of the ξ^2, τ coordinates so that they can fill up $Q(R')$.

For $\mathcal{E}_{U_2}f_2$, we therefore need to switch coordinates first: Since $y_2 \sim 1$ on U_2 , we may solve the equation $u = \phi(x, y)$ for x , i.e., $x = \frac{u - F(y)/\delta}{y}$, and can accordingly re-parametrize $S_2 = \{(x, y, \phi(x, y)) : (x, y) \in U_2\}$ in the form

$$\begin{aligned} S_2 &= \{(\tilde{\phi}(y, u), y, u) : (y, u) \in \tilde{U}_2\}, \text{ with} \\ \tilde{\phi}(y, u) &:= \frac{u - F(y)/\delta}{y}, \\ \tilde{U}_2 &:= \{(y, u) : 0 \leq u + \frac{F'(y) - \frac{F(y)}{y}}{\delta} - a < c_0^2, 0 \leq y - b < c_0\}. \end{aligned}$$

Since $|y| \sim 1$ for $(y, u) \in \tilde{U}_2$, it is easy to see that (compare (4.6))

$$|\partial_1^j \partial_2^k \tilde{\phi}(y, u)| \lesssim \delta^{-1} \quad \text{and} \quad \|H(\tilde{\phi})(y, u)\| \sim 1/\delta.$$

This suggests to change variables to $\tilde{\xi}^1 := \tau$, $\tilde{\tau} := \xi^1$ and $\tilde{\xi}^2 := \xi^2$, so that the new $\tilde{\tau}$ -coordinate points in the shorter ξ^1 -direction of $Q(R')$. Applying Lemma 4.2 in the new

coordinates with $\kappa := 1/\delta$, we obtain a slowly decaying wave packet decomposition w.r. to the coordinates $(\tilde{\tau}, \tilde{\xi}^2, \tilde{\xi}^1)$. For instance, (P3) becomes

$$|p_{w_2}(\xi, \tau)| \leq CR^{-1} \left(1 + \frac{|\tilde{\xi}^1 - \eta_2^1 + \tilde{\tau} \partial_1 \tilde{\phi}(\tilde{v}_2)|}{R} \right)^{-2} \left(1 + \frac{|\tilde{\xi}^2 - \eta_2^2 + \tilde{\tau} \partial_2 \tilde{\phi}(\tilde{v}_2)|}{R} \right)^{-2}.$$

The parameters \tilde{v}_2 correspond the parameters v_2 in the original coordinates through the relation

$$(v_2, \phi(v_2)) = (\tilde{\phi}(\tilde{v}_2), \tilde{v}_2).$$

In particular, the wave packets p_{w_2} associated to $\mathcal{E}_{U_2} f_2$ and the patch of hypersurface S_2 are mainly concentrated on tubes T_{w_2} whose axis point in the direction of the normal to S_2 at the point $(v_2, \phi(v_2))$ and are translates of tubes of the form

$$T_{(v_2)} := \{(\xi, \tau) : |\xi + \tau \nabla \phi(v_2)| \leq R, |\tau| \leq \delta R^2\},$$

w.r to the coordinates (ξ^2, τ) . We parametrize these tubes again by a pair of parameters $w_2 = (\eta_2, v_2) \in \mathcal{W}_2$, where η_2 represents the translation parameter.

Then for any $(\xi, \tau) = (\tilde{\tau}, \tilde{\xi}^2, \tilde{\xi}^1) \in Q(R')$, we have $|\tilde{\tau}| \leq \delta R^2$, so we can decompose

$$(4.17) \quad \mathcal{E}_{U_2} f_2(\xi, \tau) = \sum_{w_2 \in \mathcal{W}_2} c_{w_2} p_{w_2}(\xi, \tau).$$

This kind of change of coordinates and the effect on the wave packet decomposition was introduced already in our previous paper [BMV16], therefore we will omit the details.

Observe that the intersection of two transversal tubes T_{w_1} and T_{w_2} is essentially a cube q of side length R .

Another difference to the classical method occurs in the localisation to these cubes q . We want to choose functions χ_q which decay rapidly away from q (by scaling a fixed function χ in the usual way), but so that the Fourier transform $\hat{\chi}_q$ has compact support in a $1/R$ cube centered at the origin. We can do this in a manner so that still $\sum_q \chi_q = 1$ on $Q(R')$. The advantage is that then $p_{w_1} \chi_q$ essentially behaves like a wave packet again. In particular, we still have a well localised Fourier support, compare property (P2) from Lemma 4.2:

$$(4.18) \quad \text{supp } \mathcal{F}(p_w \chi_q) \subset B((v, \phi(v)), 3R^{-1}).$$

The same idea had already been applied in [BMV16].

The above wave packet decompositions based on Lemma 4.2 applies for any $R \geq 1/\delta$. Since we want to induct on the scale, we introduce $R' := \delta R$, so that we can induct over $R' \geq 1$ (compare again with [BMV16]). More precisely:

Definition 4.4. *For any $\alpha > 0$, we say that $E(\alpha)$ holds true if*

$$(4.19) \quad \|\mathcal{E}_{U_1}(f_1), \mathcal{E}_{U_2}(f_2)\|_{L^p(Q(R'))} \leq C_{p,\alpha} R'^\alpha \delta^{\frac{5}{2} - \frac{4}{p}} \|f_1\|_2 \|f_2\|_2$$

for all $R' \geq 1$ and $f_1 \in L^2(U_1)$, $f_2 \in L^2(U_2)$.

Let us fix a sufficiently small constant $\gamma > 0$ throughout this section. In our induction on scales argument, at various places harmless powers of the form $R'^{c\gamma}$ will arise, with possibly different values of the constants $c > 0$. With a slight abuse of notation, we shall nevertheless denote all of them by the same letter c .

We will then show that hypothesis $E(\alpha)$ essentially implies $E(\alpha(1-\gamma))$. In this argument, we will apply $E(\alpha)$ to $\tilde{R}' := R'^{1-\gamma} \geq 1$ in place of R' . Observe here the inequality $R \geq 1/\delta$ is not stable under taking powers of R ; this is why we cannot induct on the parameter R , but have to pass to the parameter R' .

For classical wave packet decompositions, the fast decay of the wave packets away from their “central” tubes allows, for instance, to treat the L^∞ -norm of a sum of wave packets all pointing in the same direction as if they had disjoint supports. Even though our rough wave packets are no longer rapidly decaying (we only have the weaker decay given by (P3)), the following lemma still holds true.

Lemma 4.5. *Let q be a cube as before, and let $d_1, d_2 \geq 0$.*

- (i) *Let $v_1 \in \mathcal{V}_1$ and $Y_1 \subset \mathcal{Y}_1$ such that for any $\eta_1 \in Y_1$ we have $\text{dist}(T_{(\eta_1, v_1)}, q) \gtrsim d_1 R$. Then*

$$\left\| \sum_{\eta_1 \in Y_1} p_{(\eta_1, v_1)} \chi_q \right\|_\infty \lesssim (1 + d_1)^{-1} R^{-1}.$$

- (ii) *Let $v_2 \in \mathcal{V}_2$ and $Y_2 \subset \mathcal{Y}_2$ such that for any $\eta_2 \in Y_2$ we have $\text{dist}(T_{(\eta_2, v_2)}, q) \gtrsim d_2 R$. Then*

$$\left\| \sum_{\eta_2 \in Y_2} p_{(\eta_2, v_2)} \chi_q \right\|_\infty \lesssim (1 + d_2)^{-1} R^{-1}.$$

- (iii) *Let p_{w_1}, p_{w_2} be transversal wave packets as above. Then*

$$(4.20) \quad \int_{\mathbb{R}^3} |p_{w_1} p_{w_2}| d(\xi, \tau) \lesssim R.$$

Proof. We prove (i) - the proof of (ii) is analogous. Let us also assume for simplicity that $v_1 = 0$; the general case can be treated in the same manner.

First, assume that $d_1 \geq 1$. Fix $(\xi, \tau) \in \mathbb{R}^3$. If $\text{dist}((\xi, \tau), q) \ll d_1 R$, then $\text{dist}(T_{(\eta_1, v_1)}, (\xi, \tau)) \gtrsim d_1 R$, and thus

$$\begin{aligned} \sum_{\eta_1 \in Y_1} |p_{(\eta_1, v_1)}(\xi, \tau)| &\lesssim R^{-1} \sum_{\eta_1 \in Y_1} \left(1 + \frac{|\xi^1 - \eta_1^1|}{R}\right)^{-2} \left(1 + \frac{|\xi^2 - \eta_1^2|}{R}\right)^{-2} \\ &\lesssim R^{-1} \sum_{\substack{k^1, k^2 \in \mathbb{Z}: \\ k^1 + k^2 \geq d_1}} (1 + |k^1|)^{-2} (1 + |k^2|)^{-2} \\ &\lesssim (1 + d_1)^{-1} R^{-1}. \end{aligned}$$

On the other hand, if $\text{dist}((\xi, \tau), q) \gtrsim d_1 R$, then we only obtain

$$\sum_{\eta_1 \in Y_1} |p_{(\eta_1, v_1)}(\xi, \tau)| \lesssim R^{-1} \sum_{k^1, k^2 \in \mathbb{Z}} (1 + |k^1|)^{-2} (1 + |k^2|)^{-2} \leq R^{-1},$$

but we can use the rapid decay of χ_q to estimate

$$\chi_q(\xi, \tau) \lesssim (d_1 R)^{-N} \ll d_1^{-N},$$

which combined gives an even better estimate than required for (i).

If $d_1 \leq 1$, the proof is even simpler.

For (iii), we only use the coordinate-free decay estimate (4.9), which is indeed sufficient for (4.20): Due to transversality, by an affine linear change of variables, we may assume that T_{w_1}

is parallel to, say, the τ -axis and T_{w_2} is parallel to the ξ^1 -axis, and both tubes go through the origin. Then

$$\begin{aligned}
\int_{\mathbb{R}^3} |p_{w_1} p_{w_2}| d(\xi, \tau) &\lesssim R^{-2} \int_{\mathbb{R}^3} \left(1 + \frac{|(\xi^1, \xi^2)|}{R}\right)^{-2} \left(1 + \frac{|(\xi^2, \tau)|}{R}\right)^{-2} d(\xi, \tau) \\
&= R \int_{\mathbb{R}^3} (1 + |(\xi^1, \xi^2)|)^{-2} (1 + |(\xi^2, \tau)|)^{-2} d(\xi, \tau) \\
&\leq R \int_{\mathbb{R}^3} (1 + |\xi^1|)^{-4/3} (1 + |\xi^2|)^{-2/3-2/3} (1 + |\tau|)^{-4/3} d(\xi, \tau) \\
&\lesssim R.
\end{aligned}$$

□

4.3. Dyadic decompositions and reduction. Similar to the classical bilinear approach, we have to count how many tubes (and also dilates of them) arising in the wave packet decompositions (4.16) and (4.17) can interact at a given point.

To this end, for any $W_j \subset \mathcal{W}_j$, $j = 1, 1$, and cube $q \subset Q(R')$, we define

$$W_j^1(q) := \{w_j \in W_j : R'^\gamma T_{w_j} \cap q \neq \emptyset\}.$$

Since our wave packets have only slow decay away from the tubes, we also want to have control over the tubes which lie at a certain distance to q . Equivalently, that means that a hollow tube of bigger diameter passes through q . For any dyadic ν with $1 < \nu < \nu_{\max} \sim R'^{1-2\gamma}$, we therefore define the *hollow tube* of diameter $\nu R R'^\gamma$ by

$$T_{w_j}^\nu := (\nu R'^\gamma T_{w_j}) \setminus \left(\frac{\nu}{2} R'^\gamma T_{w_j}\right),$$

where we specify that rT_{w_j} means scaling T_{w_j} around the central axis by the factor r . Then let

$$(4.21) \quad W_j^\nu(q) := \{w_j \in W_j : T_{w_j}^\nu \cap q \neq \emptyset\}.$$

This definition extends to the case $\nu = 1$ if we set $T_{w_j}^1 := R'^\gamma T_{w_j}$. Finally, we will see that the weak decay of our wave packets does not require a further decomposition beyond $\nu = \nu_{\max}$, hence we set $T_{w_j}^{\nu_{\max}} := \mathbb{R}^3 \setminus (\frac{\nu_{\max}}{2} R'^\gamma T_{w_j})$, and define again $W_j^{\nu_{\max}}(q)$ by (4.21). In the following, let \mathcal{D} denote the set of all dyadic numbers ν with $1 \leq \nu \leq \nu_{\max}$.

Decomposing dyadically as usually in the bilinear argument, we may assume $|W_j^\nu(q)|$ to be essentially constant, i.e., we only need to consider cubes q from the set

$$(4.22) \quad Q^\mu := \{q : |W_j^\nu(q)| \sim \mu_{\nu,j} \ \forall j = 1, 2, \ \nu \in \mathcal{D}\}.$$

Here, $\mu = (\mu_{\nu,j})_{\nu,j}$, $j = 1, 2$, $\nu \in \mathcal{D}$, is a collection of dyadic numbers. Further we define

$$(4.23) \quad Q^{\mu,\nu}(w_j) := \{q \in Q^\mu : T_{w_j}^\nu \cap q \neq \emptyset\},$$

and for any collection of dyadic numbers $\lambda_j = (\lambda_{\nu,j})_\nu$, $j = 1, 2, \nu \in \mathcal{D}$,

$$(4.24) \quad W_j^{\lambda_j,\mu} := \{w_j \in W_j : |Q^{\mu,\nu}(w_j)| \sim \lambda_{\nu,j} \ \forall \nu \in \mathcal{D}\}, \quad j = 1, 2.$$

Finally let

$$(4.25) \quad W_j^{\lambda_j,\mu,\nu_j}(q) := W_j^{\lambda_j,\mu} \cap W_j^{\nu_j}(q).$$

Using these dyadic decompositions, we can make the following reduction:

Lemma 4.6. *Assume that for all $W_1 \subset \mathcal{W}_1$, $W_2 \subset \mathcal{W}_2$, all dyadic $\lambda_1, \lambda_2, \mu$ and all dyadic $\nu_1, \nu_2 = 1, \dots, \nu_{\max}$, the following estimate holds true:*

$$(4.26) \quad \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(Q(R'))} \leq C_{p, \alpha} R'^\alpha \delta^{\frac{5}{2} - \frac{4}{p}} |W_1|^{1/2} |W_2|^{1/2}.$$

Then for any $\gamma > 0$, $E(\alpha + \gamma)$ holds true.

Remark 4.7. *There is actually only a logarithmic loss in place of R'^γ in Lemma 4.6, but we will not dwell on that.*

Therefore we may and shall assume from now on that $W_1 \subset \mathcal{W}_1$, $W_2 \subset \mathcal{W}_2$, and λ_1, λ_2 and μ are fixed.

Following the next step in the bilinear method, we divide $Q(R')$ into $R'^{\mathcal{O}(\gamma)}$ cubes b which are translates of $R'^{-4\gamma}Q(R') = Q(R'^{1-2\gamma})$. For a fixed tube T_{w_j} , we denote by $b(w_j)$ a cube b for which the cardinality of $\{q \in Q^{\mu, 1}(w_j) : q \cap b \neq \emptyset\}$ is maximal over all b . If there are several such b , we pick one of them.

We say that w_j and b are related and write $w_j \sim b$, if b is a neighbor of $b(w_j)$. If we split the set of integration in (4.26) into

$$\begin{aligned} & \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(Q(R))}^p \\ &= \sum_b \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(b)}^p \\ &\leq \sum_b I^{\sim b} + \sum_b I^{\not\sim b}, \end{aligned}$$

where $I^{\sim b}$ denotes the contribution to $\left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(b)}^p$ by all $w_1 \sim b$ and $w_2 \sim b$, and $I^{\not\sim b}$ the contribution by all w_1 and w_2 such that $w_1 \not\sim b$ or $w_2 \not\sim b$.

For any fixed b , as in the classical bilinear argument, the part $I^{\sim b}$ can be estimated applying the induction hypothesis $E(\alpha)$ on the cube b , which is a translate of $Q(R'^{1-2\gamma})$. Then the sum over all b 's can eventually be controlled since for any given w_j there are only $\mathcal{O}(1)$ b 's such that $b \sim w_j$, and we obtain

$$(4.27) \quad \sum_b I^{\sim b} \lesssim C_{p, \alpha} R'^{\alpha(1-2\gamma)} \delta^{\frac{5}{2} - \frac{4}{p}} |W_1|^{1/2} |W_2|^{1/2}.$$

We omit the details, since this part is still standard.

In order to estimate the remainder part $I^{\not\sim b}$, let us define

$$W_j^{\lambda_j, \mu, \not\sim b} := \{w_j \in W_j^{\lambda_j, \mu} : w_j \not\sim b\}.$$

Then it remains to prove that for some absolute constant $c > 0$ independent of $R, W_1, W_2, \delta, \gamma$ and λ, μ, ν and any b we have an estimate of the form

$$(4.28) \quad \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(b)}^p \leq C_{p, \alpha} R'^{c\gamma} \delta^{\frac{5}{2} - \frac{4}{p}} |W_1|^{1/2} |W_2|^{1/2},$$

and an analogous estimate with the roles of W_1 and W_2 interchanged.

Indeed, given these estimates, it is easy to complete the proof of Theorem 3.7. To this end, notice that summing these estimates over all b merely increases the constant c in the exponent of $R'^{c\delta}$, since there are only $R'^{\mathcal{O}(\gamma)}$ cubes b . Combining then (4.27) and (4.28), we obtain

$$(4.29) \quad \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(Q(R))} \leq C_{p, \alpha} R'^{\alpha(1-2\gamma) \vee c\gamma} \delta^{\frac{5}{2} - \frac{4}{p}} |W_1|^{1/2} |W_2|^{1/2}.$$

Hence, by Lemma 4.6, we see that $E(\alpha)$ implies $E(\alpha(1 - \gamma) \vee c\gamma)$ (with possibly yet another constant c).

We should mention here that the base case for the induction on scales is not that straight forward. Usually some crude estimate with a big loss in the power of R' will suffice, but we need the sharp dependency on δ in (4.19). However, our argument from [BMV16] and [BMV17] does apply also here (compare Remark 4.10). Thus, by induction, we see that $E(\alpha)$ holds for arbitrary small α , which proves Theorem 3.7, provided we have established (4.28).

4.4. The geometric argument. A key element of the bilinear method is a sophisticated geometric argument. It allows to estimate the number of tubes whose directions move according to normals to the surface along certain points on an intersection curve of the surfaces S_1 and S_2 . By an intersection curve, we mean the following: the patches of surface S_1 and S_2 are disjoint, but by means of suitable translations we may achieve that they do intersect along a curve.

More precisely, for any $v'_1 \in \mathcal{V}_1$ and $v'_2 \in \mathcal{V}_2$, we define the intersection curve $\Pi_{v'_1, v'_2} := (S_1 - (v'_1, \phi(v'_1))) \cap (S_2 - (v'_2, \phi(v'_2)))$. Moreover, if $W_1 \subset \mathcal{W}_1$, we put

$$[W_1]^{\Pi_{v'_1, v'_2}} := \{w_1 = (\eta_1, v_1) \in W_1 : (v_1 - v'_1, \phi(v_1) - \phi(v'_1)) \in \Pi_{v'_1, v'_2} + \mathcal{O}(R^{-1})\}.$$

Note that it is only the component v_1 of w_1 which is relevant for w_1 to be in $[W_1]^{\Pi_{v'_1, v'_2}}$. We therefore also introduce the projection of this set to all of its v_1 -components, i.e.,

$$[W_1]_{\mathcal{V}_1}^{\Pi_{v'_1, v'_2}} := \{v_1 \in \mathcal{V}_1 : \exists \eta_1 \text{ such that } (\eta_1, v_1) \in [W_1]^{\Pi_{v'_1, v'_2}}\} \subset \mathcal{V}_1.$$

There is an a priori estimate on the cardinality of $[W_1]_{\mathcal{V}_1}^{\Pi_{v'_1, v'_2}}$: As the projection of any intersection curve $(S_1 - (v'_1, \phi(v'_1))) \cap (S_2 - (v'_2, \phi(v'_2)))$ to the z -plane intersects the set $U_1 - v'_1$ diagonally (compare (3.34)), that intersection curve has length $\lesssim \delta$. Since by (4.3) the directions of the tubes are $1/R$ -separated, we have for all $W_1 \subset \mathcal{W}_1$

$$(4.30) \quad \sup_{v'_1 \in \mathcal{W}_1, v'_2 \in \mathcal{V}_2} \left| [W_1]_{\mathcal{V}_1}^{\Pi_{v'_1, v'_2}} \right| \lesssim \delta R = R'.$$




FIGURE 1. The geometry in Lemma 4.8

In a similar way, we define the sets $[W_2]^{\Pi_{v'_1, v'_2}} \subset W_2$ and $[W_2]_{\mathcal{V}_2}^{\Pi_{v'_1, v'_2}} \subset \mathcal{V}_2$ simply by interchanging the roles of S_1 and S_2 , and obtain the analogue to (4.30) for $\left|[W_2]_{\mathcal{V}_2}^{\Pi_{v'_1, v'_2}}\right|$.

If we consider the “cone” generated by the family of all tubes T_{w_1} which pass through some fixed cube q_0 and whose directions are given by the normals to S_1 at all points $(v_1, \phi(v_1))$ with $v_1 \in [W_1]_{\mathcal{V}_1}^{\Pi_{v'_1, v'_2}}$ (for some v'_1, v'_2), then the geometric meaning of our transversality assumption (4.4) is exactly that all tubes T_{w_2} of type 2 pass transversally through this cone. The analogous statement holds true for the “cone” generated by the family of all tubes T_{w_2} which pass through some fixed cube q_0 and whose directions are given by the normals to S_2 at all points $(v_2, \phi(v_2))$ with $v_2 \in [W_2]_{\mathcal{V}_2}^{\Pi_{v'_1, v'_2}}$.

These facts are crucial for the following lemma.

Lemma 4.8. *For all dyadic μ, λ , $1 \leq \nu_1, \nu_2 < \nu_{\max}$, all cubes q_0 and $v'_1 \in \mathcal{V}_1, v'_2 \in \mathcal{V}_2$ we have*

$$(4.31) \quad \lambda_{\nu_1, 1} \mu_{\nu_2, 2} \left| [W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q_0)]^{\Pi_{v'_1, v'_2}} \right| \lesssim R'^{c\gamma} \nu_1^2 \nu_2^2 |W_2|,$$

$$(4.32) \quad \lambda_{\nu_2, 2} \mu_{\nu_1, 1} \left| [W_2^{\lambda_2, \mu, \nu_2, \not\sim b}(q_0)]^{\Pi_{v'_1, v'_2}} \right| \lesssim R'^{c\gamma} \nu_1^2 \nu_2^2 |W_1|.$$

Remark 4.9. *The classical version of this argument corresponds to the case $\nu_1 = \nu_2 = 1$, cf. [L05].*

Proof. We shall only prove (4.31), the proof of (4.32) being analogous.

Let us consider the “cone” generated by the tubes $T_{w_1}^{\nu_1}$ associated to all $w_1 \in [W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q_0)]^{\Pi_{v'_1, v'_2}}$, however, without the part near q_0 where the tubes overlap. More precisely, we define

$$\Gamma_1^{\nu_1} := \bigcup \{T_{w_1}^{\nu_1} : w_1 \in [W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q_0)]^{\Pi_{v'_1, v'_2}}\} \setminus (q_0 + Q(R'^{1-2\gamma}))$$

and the corresponding set of cubes q by

$$Q_{\Gamma_1}^{\nu_1} := \{q \in Q^\mu : q \cap \Gamma_1^{\nu_1} \neq \emptyset\}.$$

Since the directions of the tubes are only allowed to move as the normals to S_1 corresponding to points on the intersection curve $\Pi_{v'_1, v'_2}$, the set $\Gamma_1^{\nu_1}$ is indeed a part of a cone thickened by $\nu_1 RR'\gamma$. We may decompose $\Gamma_1^{\nu_1}$ into disjoint layers $\Gamma_1^{\nu_1}(s)$, $s = 1, \dots, \nu_1$, which are essentially translates of Γ_1^1 , that is, parts of a cone thickened by $RR'\gamma$. Correspondingly, let

$$Q_{\Gamma_1}^{\nu_1}(s) := \{q \in Q^\mu : q \cap \Gamma_1^{\nu_1}(s) \neq \emptyset\}, \quad s = 1, \dots, \nu_1.$$

The proof will be accomplished by verifying the following three estimates:

- (i) $\lambda_{\nu_1, 1} \left| [W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q_0)]^{\Pi_{v'_1, v'_2}} \right| \lesssim R'^{c\gamma} \nu_1 |Q_{\Gamma_1}^{\nu_1}| :$
- (ii) $|Q_{\Gamma_1}^{\nu_1}| \leq \sum_{s=1}^{\nu_1} |Q_{\Gamma_1}^{\nu_1}(s)| \leq \nu_1 \sup_s |Q_{\Gamma_1}^{\nu_1}(s)|;$
- (iii) $\mu_{\nu_2, 2} \sup_s |Q_{\Gamma_1}^{\nu_1}(s)| \lesssim R'^{c\gamma} \nu_2^2 |W_2|.$

It is obvious that (i) – (iii) imply estimate (4.31). Estimate (ii) is clear too.

To verify (i), we observe that $\lambda_{\nu_1, 1} \sim |\{q \in Q^\mu : T_{w_1}^{\nu_1} \cap q \neq \emptyset\}|$ for any $w_1 \in W_1^{\lambda_1, \mu}$, hence by Fubini

$$\begin{aligned} \lambda_{\nu_1, 1} \left| [W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q_0)]^{\Pi_{v'_1, v'_2}} \right| &\lesssim \sum_{w_1 \in [W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q_0)]^{\Pi_{v'_1, v'_2}}} |\{q \in Q^\mu : T_{w_1}^{\nu_1} \cap q \neq \emptyset\}| \\ &\lesssim \sum_{q \in Q_{\Gamma_1}^{\nu_1}} |\{w_1 \in [W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q_0)]^{\Pi_{v'_1, v'_2}} : T_{w_1}^{\nu_1} \cap q \neq \emptyset\}| \\ &\lesssim R'^{c\gamma} |Q_{\Gamma_1}^{\nu_1}| \nu_1. \end{aligned}$$

The last estimate holds true because while the tubes T_{w_1} passing through q_0 with directions given by normals to the surface when moving along the intersection curve overlap at most of order $\mathcal{O}(R'^{c\gamma})$, our hollow thickened tubes $T_{w_1}^{\nu_1}$ overlap at most of order $\mathcal{O}(\nu_1 R'^{c\gamma})$.

It remains to prove (iii). To this end, let us fix an $s \in \{1, \dots, \nu_1\}$, i.e., a certain layer $\Gamma_1^{\nu_1}(s)$ of our thickened cone. Note next that every tube T_{w_2} of type 2 is transversal to the cone $\Gamma_1^{\nu_1}$, hence also to $\Gamma_1^{\nu_1}(s)$, and it essentially cuts out a “cap” of radius $\nu_2 RR'\gamma$ and thickness $RR'\gamma$ from $\Gamma_1^{\nu_1}(s)$ (cf. picture 1).

We therefore decompose $\Gamma_1^{\nu_1}(s)$ into caps P of dimensions $\nu_2 RR'\gamma \times \nu_2 RR'\gamma \times RR'\gamma$. Of course we only need to consider those caps P which interact with at least one of the q 's, and therefore put

$$\mathfrak{P}_{\Gamma_1}^\mu := \{ \text{caps } P \subset \Gamma_1^{\nu_1}(s) : \exists q \in Q^\mu : q \cap P \neq \emptyset \}.$$

Then

$$(4.33) \quad |Q_{\Gamma_1}^{\nu_1}(s)| \leq \sum_{P \in \mathfrak{P}_{\Gamma_1}^\mu} |\{q \in Q_{\Gamma_1}^{\nu_1}(s) : q \cap P \neq \emptyset\}| \leq \nu_2^2 R'^{2\gamma} |\mathfrak{P}_{\Gamma_1}^\mu|.$$

On the other hand, for any $q \in Q^\mu$ such that $q \cap P \neq \emptyset$, we have

$$\mu_{\nu_2, 2} \sim |\{w_2 \in W_2 : T_{w_2}^{\nu_2} \cap q \neq \emptyset\}| \leq |\{w_2 \in W_2 : T_{w_2}^{\nu_2} \cap 2P \neq \emptyset\}|.$$

Therefore by (4.33) and Fubini's theorem

$$\begin{aligned}
\mu_{\nu_2,2}|Q_{\Gamma_1}^{\nu_1}(s)| &\leq \nu_2^2 R'^{2\gamma} \mu_{\nu_2,2} |\mathfrak{P}_{\Gamma_1}^\mu| \\
&\lesssim \nu_2^2 R'^{2\gamma} \sum_{P \in \mathfrak{P}_{\Gamma_1}^\mu} |\{w_2 \in W_2 : T_{w_2}^{\nu_2} \cap 2P \neq \emptyset\}| \\
&\leq \nu_2^2 R'^{2\gamma} \sum_{w_2 \in W_2} |\{P \in \mathfrak{P}_{\Gamma_1}^\mu : T_{w_2}^{\nu_2} \cap 2P \neq \emptyset\}| \\
&\lesssim R'^{c\gamma} \nu_2^2 |W_2|.
\end{aligned}$$

The last inequality holds because the tubes T_{w_2} are transversal to the cone $\Gamma_1^{\nu_1}$. This verifies (iii). \square

4.5. Completion of the proof of Theorem 4.1. We were still left with proving (4.28), i.e., the estimate

$$\left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^p(b)}^p \leq C_{p,\alpha} R'^{c\gamma} \delta^{\frac{5}{2} - \frac{4}{p}} |W_1|^{1/2} |W_2|^{1/2}.$$

As usual, this is accomplished by interpolating an L^1 and an L^2 -estimate. The L^1 -estimate is the usual straight-forward application of Cauchy-Schwarz' inequality:

$$\begin{aligned}
&\left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^1(b)} \\
&\leq \left\| \sum_{w_1 \in W_1^{\lambda_1, \mu, \gamma^b}} p_{w_1} \right\|_{L^2(b)} \cdot \left\| \sum_{w_2 \in W_2^{\lambda_2, \mu}} p_{w_2} \right\|_{L^2(b)} \\
&\lesssim (R^2 \cdot \delta R^2)^{1/2} |W_1|^{1/2} |W_2|^{1/2} \\
(4.34) \quad &= \delta^{-3/2} R'^2 |W_1|^{1/2} |W_2|^{1/2}.
\end{aligned}$$

For the L^2 -estimate, in order to defray the notation, let us abbreviate $W_1^0(q) := W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q)$, $W_2^0(q) := W_2^{\lambda_2, \mu, \nu_2}(q)$.

Recall also that χ_q is not the characteristic function of q , but a smooth version of it which decays rapidly away from q . Nevertheless, Schur's test (cf. the similar argument on p. 856 in [BMV16]) implies that

$$\left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^0(q)} \sum_{w_2 \in W_2^0(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)}^2 \lesssim \sum_{q \in Q^\mu} \left\| \sum_{w_1 \in W_1^0(q)} \sum_{w_2 \in W_2^0(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)}^2.$$

We proceed in the usual way by applying Plancherel's theorem in order to exploit information on the Fourier support of the wave packets. It is here where we need (4.18) to keep track of the localization by χ_q :

$$\begin{aligned}
& \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^0(q)} \sum_{w_2 \in W_2^0(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)}^2 \\
& \lesssim \sum_{q \in Q^\mu} \sum_{w_1, w'_1 \in W_1^0(q)} \sum_{w_2, w'_2 \in W_2^0(q)} \int p_{w_1} p_{w'_2} \bar{p}_{w'_1} \bar{p}_{w_2} \chi_q \\
& \lesssim \sum_{q \in Q^\mu} \sum_{\substack{w_1 \in W_1^0(q) \\ w_2 \in W_2^0(q)}} \sum_{v'_1 \in [W_1^0(q)]^{\Pi_{v_1, v_2}}} |q| \left\| p_{w_1} p_{w_2} \sum_{\substack{\eta'_1: \\ (\eta'_1, v'_1) \in W_1^0(q)}} \sum_{\substack{w'_2 \in W_2^0(q): \\ v_1 + v'_2 = v'_1 + v_2}} p_{w'_1} p_{w'_2} \chi_q \right\|_{L^\infty}.
\end{aligned}$$

Note also that

$$\begin{aligned}
(4.35) \quad & \left\| p_{w_1} p_{w_2} \sum_{\substack{\eta'_1: \\ (\eta'_1, v_1) \in W_1^0(q)}} \sum_{\substack{w'_2 \in W_2^0(q): \\ v_1 + v'_2 = v'_1 + v_2}} p_{w'_1} p_{w'_2} \chi_q \right\|_{L^\infty} \\
& \leq \left\| p_{w_1} |\chi_q|^{1/4} \right\|_{L^\infty} \left\| p_{w_2} |\chi_q|^{1/4} \right\|_{L^\infty} \left\| \sum_{\substack{\eta'_1: \\ w'_1 = (\eta'_1, v'_1) \in W_1^0(q)}} p_{w'_1} |\chi_q|^{1/4} \right\|_{L^\infty} \left\| \sum_{\substack{w'_2 \in W_2^0(q): \\ v'_2 = v'_1 + v_2 - v_1}} p_{w'_2} |\chi_q|^{1/4} \right\|_{L^\infty}.
\end{aligned}$$

Moreover, if $w'_1 \in W_1^0(q)$, then $T_{w'_1}^{\nu_1} \cap q \neq \emptyset$, and thus $\text{dist}(T_{w'_1}, q) \gtrsim \nu_1 R R'^\gamma \geq \nu_1 R$ provided $\nu_1 > 1$, because then $T_{w'_1}^{\nu_1}$ is a hollow tube. Lemma 4.5 (i) with $d_1 := \nu_1$ (actually with $|\chi_q|^{1/4}$ in place of χ_q) then implies that

$$\left\| \sum_{\substack{\eta'_1: \\ w'_1 = (\eta'_1, v'_1) \in W_1^0(q)}} p_{w'_1} |\chi_q|^{1/4} \right\|_{L^\infty} \lesssim \nu_1^{-1} R^{-1}.$$

If $\nu_1 = 1$, we can still apply Lemma 4.5 (i), however now with $d_1 := 0$, and arrive at the same estimate. In a similar way, Lemma 4.5 (ii) allows to estimate the last factor in (4.35), and altogether we find that

$$\left\| p_{w_1} p_{w_2} \sum_{\substack{\eta'_1: \\ (\eta'_1, v_1) \in W_1^0(q)}} \sum_{\substack{w'_2 \in W_2^0(q): \\ v_1 + v'_2 = v'_1 + v_2}} p_{w'_1} p_{w'_2} \chi_q \right\|_{L^\infty} \lesssim \nu_1^{-2} \nu_2^{-2} R^{-4}.$$

We thus conclude that

$$\begin{aligned}
(4.36) \quad & \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^0(q)} \sum_{w_2 \in W_2^0(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)}^2 \\
& \lesssim R^{-1} \nu_1^{-2} \nu_2^{-2} \sum_{q \in Q^\mu} |W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q)| |W_2^{\lambda_2, \mu, \nu_2}(q)| \sup_{w_1, w_2} \left| [W_1^{\lambda_1, \mu, \nu_1, \gamma^b}(q)]^{\Pi_{v_1, v_2}} \right|.
\end{aligned}$$

In the case that $\nu_1, \nu_2 < \nu_{\max}$, we can further apply the geometric Lemma 4.8:

$$\begin{aligned}
& \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)}^2 \\
& \lesssim R'^{c\gamma} R^{-1} \sum_{q \in Q^\mu} |W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q)| \frac{|W_2^{\lambda_2, \mu, \nu_2}(q)|}{\mu_{\nu_2, 2}} \frac{|W_2|}{\lambda_{\nu_1, 1}} \\
& \lesssim R'^{c\gamma} R^{-1} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1, \not\sim b}} \frac{|Q^{\mu, \nu_1}(w_1)|}{\lambda_{\nu_1, 1}} |W_2| \\
& \lesssim R'^{c\gamma} R^{-1} |W_1| |W_2|,
\end{aligned}$$

since $|W_2^{\lambda_2, \mu, \nu_2}(q)|/\mu_{\nu_2, 2} \sim 1 \sim |Q^{\mu, \nu_1}(w_1)|/\lambda_{\nu_1, 1}$. In the second last inequality, we have again applied Fubini's theorem.

In the case that $\nu_1 = \nu_{\max}$ or $\nu_2 = \nu_{\max}$, we can use simpler estimates. Note that $\nu_1 \vee \nu_2 = \nu_{\max} \sim R'^{1-2\gamma}$, whereas by (4.30) we have $\left| [W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q)]_{\nu_1}^{\Pi_{\nu_1, \nu_2}} \right| \lesssim R'$. Hence (4.36) and Fubini's theorem give

$$\begin{aligned}
& \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^0(q)} \sum_{w_2 \in W_2^0(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)}^2 \\
& \lesssim R^{-1} \nu_1^{-2} \nu_2^{-2} \sum_{w_1 \in W_1} \sum_{w_2 \in W_2} |\{q \in Q^\mu : q \cap T_{w_1}^{\nu_1} \neq \emptyset \neq q \cap T_{w_2}^{\nu_2}\}| R' \\
& \lesssim R^{-1} |W_1| |W_2| \nu_1^{-1} \nu_2^{-1} (\nu_1 \wedge \nu_2) R' \\
& \lesssim R'^{c\gamma} R^{-1} |W_1| |W_2|.
\end{aligned}$$

For the second last inequality, we used that $|T_{w_1}^{\nu_1} \cap T_{w_2}^{\nu_2}| \sim \nu_1^2 \nu_2 |q|$ if, say, $\nu_1 \leq \nu_2$. Altogether, in both cases we obtain the estimate

$$\begin{aligned}
& \left\| \sum_{q \in Q^\mu} \sum_{w_1 \in W_1^{\lambda_1, \mu, \nu_1, \not\sim b}(q)} \sum_{w_2 \in W_2^{\lambda_2, \mu, \nu_2}(q)} p_{w_1} p_{w_2} \chi_q \right\|_{L^2(b)} \\
& \lesssim R'^{c\gamma} R^{-1/2} |W_1|^{1/2} |W_2|^{1/2} \\
& = \delta^{1/2} R'^{-1/2} R'^{c\gamma} |W_1|^{1/2} |W_2|^{1/2}.
\end{aligned}$$

Interpolating this with the L^1 estimate (4.34), we obtain (4.28), provided $p > 5/3$.

Remark 4.10. For the base case of our induction, i.e., (4.19) with any big power α of R' , it is again enough to estimate the corresponding wave packets

$$\left\| \sum_{w_1 \in W_1} \sum_{w_2 \in W_2} p_{w_1} p_{w_2} \right\|_p \lesssim R'^\alpha |W_1|^{1/2} |W_2|^{1/2}.$$

The L^1 estimate is similar as before, and for the L^2 -estimate we can begin similarly as above, using Lemma 4.5 (i), (ii) and now also (iii) to see that

$$\begin{aligned}
& \left\| \sum_{w_1 \in W_1} \sum_{w_2 \in W_2} p_{w_1} p_{w_2} \right\|_2^2 \\
& \lesssim \sum_{\substack{w_1 \in W_1 \\ w_2 \in W_2}} \sum_{v'_1 \in [W_1]^{\Pi_{v_1, v_2}}} \sum_{\substack{v'_2: \\ v_1 + v'_2 = v'_1 + v_2}} \int |p_{w_1} p_{w_2}| d(\xi, \tau) \left\| \sum_{\eta'_1} p_{w'_1} \right\|_{L^\infty} \left\| \sum_{\eta'_2} p_{w'_2} \right\|_{L^\infty} \\
& \lesssim R^{-1} |W_1| |W_2| \sup_{v_1, v_2} \left| [W_1]_{\mathcal{V}_1}^{\Pi_{v_1, v_2}} \right|.
\end{aligned}$$

Applying (4.30) now and interpolating with the L^1 estimate gives the correct power of δ .

5. PASSAGE TO LINEAR RESTRICTION ESTIMATES AND PROOF OF THEOREM 1.1

To prove Theorem 1.1, assume that $r > 10/3$ and $1/q' > 2/r$ and put $p := r/2$, so that $p > 5/3$, $1/q' > 1/p$. By interpolation with the trivial estimate for $r = \infty, q = 1$, it is enough to prove the result for r close to $10/3$ and q close to $5/2$, i.e., p close to $5/3$ and q close to $5/2$. Hence, we may assume that $p < 2$, $p < q < 2p = r$. Recall also that we already assume that $h(y)$ is flat at the origin, that $h'''(y) > 0$ for $y > 0$, and that $\text{supp } f \subset \{(x, y) \in \Omega : y \geq 0\}$. Let us therefore in the sequel assume that Ω is a rectangle of the form $\Omega = [-1, 1] \times [0, \eta_1]$, where $\eta_1 > 0$ is a sufficiently small dyadic number. Corresponding to η_1 , we also choose a small dyadic number $\lambda_1 > 0$ so that $|h'''(y)| \leq \lambda_1$ on Ω . Note that we can choose λ_1 sufficiently small by choosing η_1 small enough.

The problem in passing from our restriction estimates in Theorem 3.1 to the result in Theorem 1.1 lies in the fact that, unlike in the case of finite type perturbations studied in [BMV17], we can here usually no longer simply sum the estimates given by Theorem 3.1 for the contributions by the horizontal strips on which $h''' \sim \lambda$ over all dyadic values $0 < \lambda \leq \lambda_1$, since we have insufficient control on the lengths of the intervals $I_{\lambda, \epsilon}$ given by Theorem 2.2. To overcome this problem, we shall apply once more the bilinear method, making use of bilinear estimates for pairs of surface patches on which h''' is sufficiently small, which, however, will rather easily be established, following basically the approach from [V05] (and [L05]) devised for the unperturbed parabolic hyperboloid. In a final step, we shall then fuse these bilinear estimates with the linear estimates from Theorem 3.1 by means of a kind of bootstrap argument. It will be in this part where we shall have to make the stronger assumption that h''' is monotonic (say, increasing).

A key step in [V05], [L05] consists in devising a suitable Whitney-type decomposition of $\Omega \times \Omega$ into direct products of rectangles; more details will be given later. This leads us to considering bilinear estimates over pairs of “close” rectangles $W_1 = J_1 \times I_1, W_2 = J_2 \times I_2$ contained in Ω , where J_1, J_2 are intervals of dyadic length $d \leq 1$, and I_1, I_2 are intervals of dyadic length $r \leq \eta_1$, and which are separated in the x -coordinate of size d and in the y -coordinates of size r .

Consider such a pair W_1, W_2 , and set for $i = 1, 2$,

$$\begin{aligned}
W_i^> &:= \{(x, y) \in W_i : h'''(y) > dr/100\}, \\
W_i^< &:= \{(x, y) \in W_i : h'''(y) \leq dr/100\}.
\end{aligned}$$

By applying Theorem 2.2 to the function $\varphi := h'''$ let us decompose the interval $[0, \eta_1]$ into intervals $I_{\lambda, \iota}$ on which $\lambda/2 < h'''(y) \leq 4\lambda$ ($\lambda > 0$ dyadic). Note that if h''' is monotonic, for each λ there will be at most one corresponding ι , but for the time being this will not yet be relevant and we could as well work with non-monotonic h''' . Correspondingly, for any subset W of Ω we put $W^{\lambda, \iota} := \{(x, y) \in W : y \in I_{\lambda, \iota}\}$.

Choose next \tilde{q} so that $1/q' > 1/\tilde{q}' > 2/r$. Applying the argument from the beginning of Section 3 and making use of the equivalence of (3.2) and (3.3) in combination with Theorem 3.1 we then find that

$$\|\mathcal{E}_{W_i^{\lambda, \iota}} f\|_{L^r} \leq C_{r, \tilde{q}} \|f|_{W_i^{\lambda, \iota}}\|_{L^{\tilde{q}}},$$

since $d_{I_{\lambda, \iota}}^{1-2/r-1/\tilde{q}} \leq 1$, and thus in combination with Hölder's estimate we obtain

$$\|\mathcal{E}_{W_i^{\lambda, \iota}} f\|_{L^r} \leq C_{r, \tilde{q}} (dr)^{\frac{1}{\tilde{q}} - \frac{1}{q}} \|f|_{W_i}\|_{L^q}.$$

Write $\varepsilon := (1/\tilde{q} - 1/q)/3$. Then $\varepsilon > 0$, and if $\lambda \geq dr/400$, then $(dr)^{\frac{1}{\tilde{q}} - \frac{1}{q}} \lesssim \lambda^{2\varepsilon} (dr)^\varepsilon$. Decomposing $W_1^>$ into such sets $W_i^{\lambda, \iota}$, we then see that by choosing $r \in \mathbb{N}$ in Theorem 2.2 sufficiently large we can sum the preceding estimates over the corresponding dyadic λ 's with $\lambda \leq \lambda_1$ and ι 's and arrive at the following uniform linear estimates:

$$(5.1) \quad \|\mathcal{E}_{W_i^>} f\|_{L^r} \leq C_{r, q} \lambda_1^\varepsilon |W_i|^\varepsilon \|f|_{W_i^>}\|_{L^q}, \quad i = 1, 2,$$

where we have used the notation $f_A := f\chi_A$ for any subset $A \subset \Omega$. Recall that this part of the argument does not require any monotonicity assumption on h''' yet.

Our next step, namely the proof of the following bilinear estimate for the operators $\mathcal{E}_{W_i^<}$, will, in contrast, make use of this monotonicity assumption:

$$(5.2) \quad \|\mathcal{E}_{W_1^<} f \mathcal{E}_{W_2^<} g\|_{L^p} \leq C_{r, q} |W_1|^{\frac{1}{q'} - \frac{1}{p}} |W_2|^{\frac{1}{q'} - \frac{1}{p}} \|f|_{W_1^<}\|_{L^q} \|g|_{W_2^<}\|_{L^q}.$$

Proof. By the monotonicity of h''' , the set $J^< := \{y \in [0, \lambda_1] : h'''(y) \leq dr/100\}$ is an interval of the form $J^< = [0, \eta]$, with $0 \leq \eta \leq \eta_1$. Looking at the Taylor expansion of h around 0, we see that this implies that

$$|h^{(k)}(y)| \leq dr/100 \quad \text{for all } y \in J^<, k = 0, 1, 2, 3.$$

Let us next assume w.l.o.g. that the rectangle W_1 is located “below” W_2 , and that $W_2^< \neq \emptyset$. Then

$$W_1^< = W_1 \quad \text{and} \quad W_2^< = W_2 \cap \{(x, y) : y \in [0, \eta]\}.$$

Denote by $z_1^0 = (x_1^0, y_1^0)$ the lower left vertex of W_1 , and translate the coordinates (x, y) so that z_1^0 becomes the origin. Then, in the new coordinates, the function ϕ assumes the form

$$\phi(x, y) = xy + H(y),$$

except for affine-linear terms which have no effect on the bilinear estimates, and W_1 assumes the form $W_1 = [0, d] \times [0, r]$, and $W_2^<$ the form $W_2^< = [2d, 3d] \times [2r, \tilde{\eta}]$, with $2r \leq \tilde{\eta} \leq 3r$, or $W_2^< = [-2d, -d] \times [2r, \tilde{\eta}]$. Moreover, H will also satisfy the estimates

$$(5.3) \quad |H^{(k)}(y)| \leq dr/100 \quad \text{for all } y \in [0, \tilde{\eta}], k = 0, 1, 2, 3.$$

We next pass to the re-scaled coordinates (x', y') defined by $x = dx'$, $y = ry'$, and put

$$\phi^s(x', y') := \frac{1}{dr} \phi(dx', ry') = x'y' + H^s(y'),$$

where $H^s(y') := H(ry')/dr$. The rectangles $W_1^<$ and $W_2^<$ correspond to $(W_1^<)^s := [0, 1] \times [0, 1]$ and $W_2^< = [2, 3] \times [2, \eta^s]$, or $(W_2^<)^s = [-2, -1] \times [2, \eta^s]$, in these coordinates, with $2 \leq \eta^s \leq 3$. Finally note that by (5.3), we have

$$(5.4) \quad |(H^s)^{(k)}(y')| \leq 1/100 \quad \text{for all } y' \in [0, \eta^s], k = 0, 1, 2, 3,$$

so that the function $\phi^s(x', y')$ is a very small perturbation of $x'y'$, whereas $(W_1^<)^s$ and $(W_2^<)^s$ are contained in squares of side length 1 which are 1-separated in each coordinate. One can then easily check that both transversality functions $|TV_1^s|$ and $|TV_2^s|$ associated to ϕ^s (cf. (3.37)) are of size ~ 1 on $(W_1^<)^s \times (W_2^<)^s$. Thus, if we had also a good control on higher order derivatives of H^s of the form $|(H^s)^{(k)}(y')| \leq C_k$ for all $y' \in [0, \eta^s]$, $k \geq 4$, we could immediately argue as in [V05], or even apply directly Theorem 1.1 in [L05], to prove the following bilinear estimate

$$(5.5) \quad \|\mathcal{E}_{(W_1^<)^s}^s f \mathcal{E}_{(W_2^<)^s}^s g\|_{L^p} \leq C_{r,q} \|f_{(W_1^<)^s}\|_{L^q} \|g_{(W_2^<)^s}\|_{L^q}$$

for the scaled surface given as the graph of ϕ^s , from which (5.2) follows immediately by scaling back to our original coordinates.

However, this control of the higher order derivatives is here no longer available, only (5.4), but we had already seen in Section 4 how to establish the required bilinear estimates even under such weaker assumptions, working with slowly decaying wave packets, and the same reasoning can be applied to our present situation, and thus we can still verify the required estimates in (5.5). \square

Given the estimates (5.1) and (5.2), we can finally complete the proof of Theorem 1.1. We first decompose the interval $[-1, 1]$ for any dyadic number 2^{j_1} into dyadic subintervals $J_k^{j_1}$ of length $d = 2^{-j_1}$, $j_1 \geq 1$, in the usual way and say that two such dyadic subintervals $J_k^{j_1}$ and $J_{k'}^{j_1}$ are *related* and write $J_k^{j_1} \approx J_{k'}^{j_1}$ if they are not adjacent but have adjacent dyadic parent intervals of length 2^{1-j_1} . In a similar way, we decompose the interval $[0, \eta_1]$ into dyadic intervals $I_l^{j_2}$ of length $r = 2^{-j_2}$ and define when two such intervals $I_l^{j_2}$ and $I_{l'}^{j_2}$ are related and write $I_l^{j_2} \approx I_{l'}^{j_2}$ in the same way as before. Finally, we put $W_{k,l}^{j_1,j_2} := J_k^{j_1} \times I_l^{j_2}$, and say that two such rectangles $W_{k,l}^{j_1,j_2}$ and $W_{k',l'}^{j_1,j_2}$ of dimension $d \times r$ are *related* and write $W_{k,l}^{j_1,j_2} \approx W_{k',l'}^{j_1,j_2}$ if $J_k^{j_1} \approx J_{k'}^{j_1}$ and $I_l^{j_2} \approx I_{l'}^{j_2}$. This leads to a kind of Whitney decomposition of $\Omega \times \Omega$ away from its diagonal D into rectangular boxes, i.e.,

$$(5.6) \quad \Omega \times \Omega \setminus D = \bigcup_{j_1 \geq 1, j_2 \geq \log_2(1/\lambda_1)} \bigcup_{W_{k,l}^{j_1,j_2} \approx W_{k',l'}^{j_1,j_2}} W_{k,l}^{j_1,j_2} \times W_{k',l'}^{j_1,j_2}.$$

Since $\|\mathcal{E}f\|_{L^r}^2 = \|\mathcal{E}f\mathcal{E}g\|_{L^p}$, if we choose $g := f$, it will suffice to estimate $\|\mathcal{E}f\mathcal{E}g\|_{L^p}$, and writing

$$f_{k,l}^{j_1,j_2} := f \chi_{W_{k,l}^{j_1,j_2}}, \quad g_{k',l'}^{j_1,j_2} := g \chi_{W_{k',l'}^{j_1,j_2}},$$

by (5.6) we may decompose

$$\mathcal{E}f\mathcal{E}g = \sum_{j_1 \geq 1, j_2 \geq \log_2(1/\lambda_1)} \sum_{W_{k,l}^{j_1,j_2} \approx W_{k',l'}^{j_1,j_2}} (\mathcal{E}f_{k,l}^{j_1,j_2}) (\mathcal{E}g_{k',l'}^{j_1,j_2}),$$

so that

$$(5.7) \quad \|\mathcal{E}f\mathcal{E}g\|_{L^p} \leq \sum_{j_1 \geq 1, j_2 \geq \log_2(1/\lambda_1)} \left\| \sum_{W_{k,l}^{j_1,j_2} \approx W_{k',l'}^{j_1,j_2}} (\mathcal{E}f_{k,l}^{j_1,j_2}) (\mathcal{E}g_{k',l'}^{j_1,j_2}) \right\|_{L^p}.$$

We shall estimate each summand separately. To this end, let us fix j_1, j_2 , and defray the notation by writing $d := 2^{-j_1}, r := 2^{-j_2}$, and $W_{1;k,l} := W_{k,l}^{j_1,j_2}$, $W_{2;k',l'} := W_{k',l'}^{j_1,j_2}$, and $f_{k,l} := f_{k,l}^{j_1,j_2}$, $g_{k',l'} := g_{k',l'}^{j_1,j_2}$. We also shortly write $k \approx k', l \approx l'$ in place of $W_{1;k,l} \approx W_{2;k',l'}$. Note that $|k - k'| \leq 2$ and $|l - l'| \leq 2$, if $k \approx k', l \approx l'$. Then, with the preceding notation, since d and r are given, we may decompose

$$\begin{aligned} \mathcal{E}f_{k,l} &= \mathcal{E}_{W_{1;k,l}^>} f_{k,l} + \mathcal{E}_{W_{1;k,l}^<} f_{k,l}, \\ \mathcal{E}g_{k',l'} &= \mathcal{E}_{W_{2;k',l'}^>} g_{k',l'} + \mathcal{E}_{W_{2;k',l'}^<} g_{k',l'}, \end{aligned}$$

which leads to a decomposition of $(\mathcal{E}f_{k,l}^{j_1,j_2})(\mathcal{E}g_{k',l'}^{j_1,j_2})$ into four terms, whose contributions we shall compute separately.

We begin by estimating $I := \|\sum_{k \approx k', l \approx l'} \mathcal{E}_{W_{1;k,l}^>} f_{k,l} \mathcal{E}_{W_{2;k',l'}^>} g_{k',l'}\|_p$. To this end, we first apply a standard "orthogonality" argument (compare for the proof of Lemma 6.1 in [TVV98]), followed by an application of the Cauchy-Schwarz inequality, to see that

$$I^p \lesssim \sum_{k \approx k', l \approx l'} \|\mathcal{E}_{W_{1;k,l}^>} f_{k,l} \mathcal{E}_{W_{2;k',l'}^>} g_{k',l'}\|_p^p \leq \sum_{k \approx k', l \approx l'} \|\mathcal{E}_{W_{1;k,l}^>} f_{k,l}\|_r^p \|\mathcal{E}_{W_{2;k',l'}^>} g_{k',l'}\|_r^p.$$

Next, since $|k - k'| \leq 2$ and $|l - l'| \leq 2$, applying again Cauchy-Schwarz, we may then essentially estimate I by

$$I \lesssim \left(\sum_{k,l} \|\mathcal{E}_{W_{1;k,l}^>} f_{k,l}\|_r^r \right)^{1/r} \left(\sum_{k,l} \|\mathcal{E}_{W_{1;k,l}^>} g_{k,l}\|_r^r \right)^{1/r}$$

(plus at most 8 terms of similar kind). By (5.1), this can eventually be estimated by

$$(5.8) \quad I \lesssim \lambda_1^\varepsilon (cd)^\varepsilon \left(\sum_{k,l} \|f_{k,l}\|_q^r \right)^{1/r} \left(\sum_{k,l} \|g_{k,l}\|_q^r \right)^{1/r} \leq (cd)^\varepsilon \|f\|_q \|g\|_q,$$

since $r > q$, where $\varepsilon > 0$.

We next turn to the term $II := \|\sum_{k \approx k', l \approx l'} \mathcal{E}_{W_{1;k,l}^<} f_{k,l} \mathcal{E}_{W_{2;k',l'}^<} g_{k',l'}\|_p$. Making again use of the afore-mentioned "orthogonality" argument in combination with the bilinear estimate (5.2), we find that

$$II^p \lesssim (cd)^{\varepsilon p} \sum_{k \approx k', l \approx l'} \|f_{k,l}\|_q^p \|g_{k',l'}\|_q^p,$$

if we assume that $1/q' - 1/p > \varepsilon$. From here on we can argue in a similar way as before to see that also

$$(5.9) \quad II \lesssim (cd)^\varepsilon \left(\sum_{k,l} \|f_{k,l}\|_q^r \right)^{1/r} \left(\sum_{k,l} \|g_{k,l}\|_q^r \right)^{1/r} \leq (cd)^\varepsilon \|f\|_q \|g\|_q.$$

As for the two “mixed term” sums III and IV , which can be handled in analogous ways, let us just look at one of them, say $III := \|\sum_{k \approx k', l \approx l'} \mathcal{E}_{W_{1;k,l}^>} f_{k,l} \mathcal{E}_{W_{2;k',l'}^<} g_{k',l'}\|_p$. We can first argue as for I to see that

$$III^p \leq \sum_{k \approx k', l \approx l'} \|\mathcal{E}_{W_{1;k,l}^>} f_{k,l}\|_r^p \|\mathcal{E}_{W_{2;k',l'}^<} g_{k',l'}\|_r^p \leq \sum_{k \approx k', l \approx l'} \|\mathcal{E}_{W_{1;k,l}^>} f_{k,l}\|_r^p \|\mathcal{E}_{W_{2;k',l'}^<} g_{k',l'}\|_r^p.$$

Here, by (5.1), we can again estimate

$$\|\mathcal{E}_{W_{1;k,l}^>} f_{k,l}\|_r^p \lesssim (\lambda_1^\varepsilon (cd)^\varepsilon \|f_{k,l}\|_q)^p,$$

but we do not know a priori that the operator $\mathcal{E}_{W_{2;k',l'}^<}$ is bounded.

Therefore we first replace the operator \mathcal{E} in the preceding estimates by the truncated operator $\mathcal{E}^{\lambda_0} := \mathcal{E}_{A_{\lambda_0}}$, for any positive dyadic number $0 < \lambda_0 < \lambda_1$, where $A_{\lambda_0} := \{(x, y) \in \Omega : h'''(y) > \lambda_0\}$. Then, by (5.1),

$$(5.10) \quad \|\mathcal{E}^{\lambda_0} f\|_{L^r} \leq C_{r,q}(\lambda_0) \|f\|_q, \quad i = 1, 2,$$

and we choose for C_{r,q,λ_0} the smallest possible constant for this estimate. Then we can also estimate $\|\mathcal{E}_{W_{2;k',l'}^<}^{\lambda_0} g_{k',l'}\|_r \leq C_{r,q}(\lambda_0) \|g_{k',l'}\|_q$, and if we denote III_{λ_0} the same expression as III , only with \mathcal{E} replaced by \mathcal{E}^{λ_0} , then we find again by the Cauchy-Schwarz' inequality and a similar reasoning as before that

$$(5.11) \quad III_{\lambda_0} \leq C \lambda_1^\varepsilon (cd)^\varepsilon C_{r,q}(\lambda_0) \|f\|_q \|g\|_q.$$

The same kind of estimate also holds for the second “mixed term” IV_{λ_0} . Combining the latter estimate and (5.8)–(5.11), and choosing $f = g$, we then find that

$$\left\| \sum_{W_{k,l}^{j_1,j_2} \approx W_{k',l'}^{j_1,j_2}} (\mathcal{E} f_{k,l}^{j_1,j_2}) (\mathcal{E} g_{k',l'}^{j_1,j_2}) \right\|_p \leq C (dr)^\varepsilon [1 + \lambda_1^\varepsilon C_{r,q}(\lambda_0)] \|f\|_q \|g\|_q.$$

Summing finally over all dyadic $d = 2^{-j_1}$ and $r = 2^{-j_2}$, by (5.7) we find that

$$\|\mathcal{E}_{\lambda_0} f\|_r \leq C [1 + \lambda_1^\varepsilon C_{r,q}(\lambda_0)]^{\frac{1}{2}} \|f\|_q.$$

Thus

$$C_{r,q}(\lambda_0) \leq C [1 + \lambda_1^\varepsilon C_{r,q}(\lambda_0)]^{\frac{1}{2}},$$

and choosing λ_1 sufficiently small, we see that $C_{r,q}(\lambda_0)$ is uniformly bounded in λ_0 , i.e., there is a constant $C_{r,q}$ such that

$$(5.12) \quad \|\mathcal{E}_{\lambda_0} f\|_r \leq C \|f\|_q \quad \text{for all } 0 < \lambda_0 \leq \lambda_1.$$

Since $\mathcal{E}_{\lambda_0} f \rightarrow \mathcal{E} f$ uniformly on compact sets as $\lambda_0 \rightarrow 0$, this implies that also $\|\mathcal{E} f\|_r \leq C \|f\|_q$, so that the proof of Theorem 1.1 is complete.

REFERENCES

- [Be16] Bejenaru, I., Optimal bilinear restriction estimates for general hypersurfaces and the role of the shape operator. *Int. Math. Res. Not. IMRN* (2017), no. 23, 7109–7147.
- [Bo91] Bourgain, J., Besicovitch-type maximal operators and applications to Fourier analysis. *Geom. Funct. Anal.* 22 (1991), 147–187.
- [Bo95a] Bourgain, J., Some new estimates on oscillatory integrals. *Essays in Fourier Analysis in honor of E. M. Stein*. Princeton Math. Ser. 42, Princeton University Press, Princeton, NJ 1995, 83–112.
- [Bo95b] Bourgain, J., Estimates for cone multipliers. *Oper. Theory Adv. Appl.* 77 (1995), 1–16.
- [BoG11] Bourgain, J., Guth, L., Bounds on oscillatory integral operators based on multilinear estimates. *Geom. Funct. Anal.*, Vol.21 (2011) 1239–1295.
- [BMV16] Buschenhenke, S., Müller, D., Vargas, A., A Fourier restriction theorem for a two-dimensional surface of finite type. *Anal. PDE* 10-4 (2017), 817–891.
- [BMV17] Buschenhenke, S., Müller, D., Vargas, A., A Fourier restriction theorem for a perturbed hyperbolic paraboloid. *Proc. London Math. Soc.* (3) 120 (2020), no. 1, 124–154.
- [BMV19] Buschenhenke, S., Müller, D., Vargas, A., On Fourier restriction for finite-type perturbations of the hyperbolic paraboloid, preprint 2019, arXiv:1902.05442v2.
- [ChL17] Cho, C.-H., Lee, J., Improved restriction estimate for hyperbolic surfaces in \mathbb{R}^3 . *J. Funct. Anal.* 273 (2017), no. 3, 917–945.
- [Gr81] Greenleaf, A., *Principal Curvature and Harmonic Analysis*. Indiana Univ. Math. J. Vol. 30, No. 4 (1981).
- [Gu16] Guth, L. A restriction estimate using polynomial partitioning. *J. Amer. Math. Soc.* 29 (2016), no. 2, 371–413.
- [Gu17] Guth, L., Restriction estimates using polynomial partitioning II. *Acta Math.* Vol. 221, No. 1 (2016), 81–142.
- [IKM10] Ikromov, I. A., Kempe, M., Müller, D., Estimates for maximal functions associated with hypersurfaces in \mathbb{R}^3 and related problems in harmonic analysis. *Acta Math.* 204 (2010), 151–271.
- [IM11] Ikromov, I. A., Müller, D., Uniform estimates for the Fourier transform of surface carried measures in \mathbb{R}^3 and an application to Fourier restriction. *J. Fourier Anal. Appl.*, 17 (2011), no. 6, 1292–1332.
- [IM15] Ikromov, I. A., Müller, D., *Fourier restriction for hypersurfaces in three dimensions and Newton polyhedra*. Annals of Mathematics Studies, 194. Princeton University Press, Princeton, NJ, 2016.
- [K17] Kim, J., Some remarks on Fourier restriction estimates, preprint 2017. arXiv:1702.01231
- [L05] Lee, S., Bilinear restriction estimates for surfaces with curvatures of different signs, *Transactions of the American Mathematical Society*, Vol. 358, No. 8, 2511–2533, 2005.
- [LV10] Lee, S., Vargas, A., Restriction estimates for some surfaces with vanishing curvatures. *J. Funct. Anal.* 258 (2010), no. 9, 2884–2909.
- [MVV96] Moyua, A., Vargas, A., Vega, L., Schrödinger maximal function and restriction properties of the Fourier transform. *Internat. Math. Res. Notices* 16 (1996), 793–815.
- [MVV99] Moyua, A., Vargas, A., Vega, L., Restriction theorems and maximal operators related to oscillatory integrals in \mathbb{R}^3 . *Duke Math. J.*, 96 (3), (1999), 547–574.
- [SJ 74] Sjölin, Per, Fourier multipliers and estimates of the Fourier transform of measures carried by smooth curves in \mathbb{R}^2 , *Studia Math.* 51 (1974), 169–182.
- [St86] Stein, E.M., *Oscillatory Integrals in Fourier Analysis*. Beijing Lectures in Harmonic Analysis. Princeton Univ. Press 1986.
- [Sto17a] Stovall, B., Linear and bilinear restriction to certain rotationally symmetric hypersurfaces. *Trans. Amer. Math. Soc.* 369 (2017), no. 6, 4093–4117.
- [Sto17b] Stovall, B., Scale invariant Fourier restriction to a hyperbolic surface. *Anal. PDE* 12 (2019), no. 5, 1215–1224.
- [Str77] Strichartz, R. S., Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations. *Duke Math. J.* 44 (1977), no. 3, 705–714.
- [T01] Tao, T., Endpoint bilinear restriction theorems for the cone, and some sharp null-form estimates. *Math. Z.* 238 (2001), 215–268.
- [T03] Tao, T., A Sharp bilinear restriction estimate for paraboloids. *Geom. Funct. Anal.* 13, 1359–1384, 2003.

- [TVV98] Tao, T., Vargas, A., Vega, L., A bilinear approach to the restriction and Keakey conjectures. J. Amer. Math. Soc. 11 (1998) no. 4 , 967–1000.
- [TVI00] Tao, T., Vargas, A., A bilinear approach to cone multipliers I. Restriction estimates. Geom. Funct. Anal. 10, 185–215, 2000.
- [TVII00] Tao, T., Vargas, A., A bilinear approach to cone multipliers II. Applications. Geom. Funct. Anal. 10, 216–258, 2000.
- [To75] Tomas, P. A., A restriction theorem for the Fourier transform. Bull. Amer. Math. Soc. 81 (1975), 477–478.
- [V05] Vargas, A., Restriction theorems for a surface with negative curvature, Math. Z. 249, 97–111 (2005).
- [W01] Wolff, T., A Sharp Bilinear Cone Restriction Estimate. Ann. of Math., Second Series, Vol. 153, No. 3, 661–698, 2001.

S. BUSCHENHENKE: MATHEMATISCHES SEMINAR, C.A.-UNIVERSITÄT KIEL, LUDEWIG-MEYN-STRASSE 4, D-24118 KIEL, GERMANY

E-mail address: buschenhenke@math.uni-kiel.de

URL: <http://www.math.uni-kiel.de/analysis/de/buschenhenke>

D. MÜLLER: MATHEMATISCHES SEMINAR, C.A.-UNIVERSITÄT KIEL, LUDEWIG-MEYN-STRASSE 4, D-24118 KIEL, GERMANY

E-mail address: mueller@math.uni-kiel.de

URL: <http://www.math.uni-kiel.de/analysis/de/mueller>

A. VARGAS: DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN

E-mail address: ana.vargas@uam.es

URL: <http://matematicas.uam.es/~AFA/>