

# THE GREATEST COMMON VALUATION OF $\phi_n$ AND $\psi_n^2$ AT POINTS ON ELLIPTIC CURVES

PAUL VOUTIER AND MINORU YABUTA

**ABSTRACT.** Given a minimal model of an elliptic curve,  $E/K$ , over a finite extension,  $K$ , of  $\mathbb{Q}_p$  for any rational prime,  $p$ , and any point  $P \in E(K)$  of infinite order, we determine precisely  $\min(v(\phi_n(P)), v(\psi_n^2(P)))$ , where  $v$  is a normalised valuation on  $K$  and  $\phi_n(P)$  and  $\psi_n(P)$  are polynomials arising from multiplication by  $n$  on this model of the curve.

## 1. INTRODUCTION

Let  $E/K$  be an elliptic curve given by a Weierstrass equation and let  $\{\psi_n(X, Y)\}_{n \geq 1}$  be the sequence of associated division polynomials. We can also define a related family of polynomials,  $\phi_n(X, Y)$ , such that

$$x([n]P) = \frac{\phi_n(x(P), y(P))}{\psi_n^2(x(P), y(P))},$$

for any point,  $P = (x(P), y(P)) \in E(K)$ , satisfying the Weierstrass equation.

For many problems related to integral (and more generally,  $S$ -integral) points on elliptic curves, it is important to know, or at least bound,  $\gcd(\phi_n(P), \psi_n^2(P))$ . For example, Ayad [1] used such information to find all  $S$ -integral points on some elliptic curves of rank 1 (see his Sections 6–8 for three specific examples demonstrating his technique). More recently, Stange [11], building on work of Ingram [4], showed that if  $E/\mathbb{Q}$  is an elliptic curve in minimal Weierstrass form and  $P \in E(\mathbb{Q})$  is a non-torsion point, then there is at most one value of  $n$  larger than a bound that can be made effective such that  $[n]P$  is integral. This leads to results (see Corollary 1.2 and the discussion that follows in [11]) related to the Hall-Lang Conjecture [5].

Good knowledge of  $\gcd(\phi_n(P), \psi_n^2(P))$  is also required for problems involving elliptic divisibility sequences – this is not unrelated to the fact that the above two applications also involve rank 1 subgroups of  $E(\mathbb{Q})$ . An example of this is Lemma 3.5 (and Remark 3.6) in [13]. A weaker version of that Lemma 3.5 would have led to a weaker version of the main result there. In fact, trying to understand and generalise that

---

*Key words and phrases.* Elliptic curves, Elliptic division polynomials, valuations.

Lemma 3.5 is how our interest in this subject arose. In this paper, we determine this gcd precisely. This is a special case of the more general result we prove here.

Our results are significantly better than previous results. Previous results typically depend on the resultant of  $\phi_n(P)$  and  $\psi_n^2(P)$ , resulting in exponents that are  $O(n^4)$  (e.g., Lemmas 4 and 5, along with the claims used in their proofs, in [3]), whereas here our results are precise and we show that the actual growth of the exponents is  $O(n^2)$ .

Throughout this paper, we let  $p$  be a rational prime,  $K$  a finite extension of  $\mathbb{Q}_p$ ,  $R$  the ring of integers of  $K$ , with maximal ideal  $\mathcal{M}$ ,  $\pi$  a uniformiser for  $R$  (i.e.,  $\mathcal{M} = \pi R$ ), residue field  $k = R/\mathcal{M}$  and  $v$  a valuation for  $K$  normalised so that  $v(\pi) = 1$ .

We let  $\lambda_v : E(K) \setminus \{O\} \rightarrow \mathbb{R}$  be the local height function for  $E$  at  $v$  as defined by Silverman in [10]. We use this local height function to be consistent with the one we used in our earlier papers. See [2, Section 4] for helpful notes about the various normalisations of local height functions.

**Theorem 1.1.** *Assume we have a minimal Weierstrass model of an elliptic curve  $E/K$  and that  $P \in E(K)$  is of infinite order. Let  $n$  be a positive integer and put*

$$k_{v,n}(P) = \min(v(\phi_n(P)), v(\psi_n^2(P))).$$

*If  $P$  modulo  $\pi$  is non-singular, then  $k_{v,n}(P) = \min(0, n^2 v(x(P)))$ .*

*If  $P$  modulo  $\pi$  is singular, then*

$$k_{v,n}(P) = -\frac{2[K : \mathbb{Q}_p]}{\log |k|} \lambda_v(P) n^2 + \epsilon_{v,n}(P),$$

*where  $\epsilon_{v,n}(P)$  is as in Table 1.1.*

**Remark 1.2.** (i)  $c_v$  is the size of the component group of  $E$  at  $v$ , which we define at the end of Subsection 2.1.

(ii)  $m_P$  is the smallest positive integer such that  $[m_P]P$  modulo  $\pi$  is non-singular.

(iii) If the Kodaira symbol is  $I_0$ ,  $I_1$ ,  $II$  or  $II^*$ , then we cannot have singular reduction. Hence these symbols do not appear in Table 1.1.

(iv) In the entry for  $I_{2m}^*$  with  $c_v = 4$  in Table 1.1,  $v(\phi_2(P))$  can take only two values, either 4 or  $2m + 4$  – see Lemma 2.10.

(v)  $a_{P,v}$  in the entry for  $I_m$  in Table 1.1 is the component of the Néron model special fibre containing  $P$ . See Lemma 5.1 and the surrounding text in [10] for more information. We use  $m$  instead of  $m_P$  for the modulus here as it results in a simpler expression for  $\epsilon_{v,n}$ . We let  $n'$  be the smallest non-negative representative of the congruence class  $a_{P,v}n \bmod m$ , so  $0 \leq n' < m$ .

Kodaira symbol	$m_P$	$-2[K : \mathbb{Q}_p]\lambda_v(P)/\log k $	$\epsilon_{v,n}(P)$
$III^*$	2	$3/2$	0 if $n \equiv 0 \pmod{m_P}$ $-3/2$ if $n \not\equiv 0 \pmod{m_P}$
$IV^*$	3	$4/3$	0 if $n \equiv 0 \pmod{m_P}$ $-4/3$ if $n \not\equiv 0 \pmod{m_P}$
$III$	2	$1/2$	0 if $n \equiv 0 \pmod{m_P}$ $-1/2$ if $n \not\equiv 0 \pmod{m_P}$
$IV$	3	$2/3$	0 if $n \equiv 0 \pmod{m_P}$ $-2/3$ if $n \not\equiv 0 \pmod{m_P}$
$I_m^*, c_v = 2$	2	1	0 if $n \equiv 0 \pmod{m_P}$ $-1$ if $n \not\equiv 0 \pmod{m_P}$
$I_m^*, m$ odd, $c_v = 4$ [2] $P$ non-singular mod $\pi$	2	1	0 if $n \equiv 0 \pmod{m_P}$ $-1$ if $n \not\equiv 0 \pmod{m_P}$
$I_m^*, m$ odd, $c_v = 4$ , [2] $P$ singular mod $\pi$	4	$(m+4)/4$	0 if $n \equiv 0 \pmod{m_P}$ $-(m+4)/4$ if $n \equiv 1, 3 \pmod{m_P}$ $-1$ if $n \equiv 2 \pmod{m_P}$
$I_{2m}^*, c_v = 4$	2	$v(\phi_2(P))/4$	0 if $n \equiv 0 \pmod{m_P}$ $-v(\phi_2(P))/4$ if $n \not\equiv 0 \pmod{m_P}$
$I_m$	$\frac{m}{\gcd(a_{P,v}, m)}$	$\frac{a_{P,v}(m - a_{P,v})}{m}$	$-\frac{n'(m - n')}{m}$ if $a_{P,v}n \equiv n' \pmod{m}$

TABLE 1.1.  $\epsilon_{v,n}(P)$  values

**Remark 1.3.** The referee has kindly shared with us other ways that the quantities in our theorem arise. For example, when calculating height pairings for  $P, Q \in E(K)$ , our  $-\epsilon_{v,n}(P)$  is the correction term,  $\text{contr}_v(P) = \text{contr}_v(P, P)$ , that arises in Section 11.8 of [7]. In the notation of that section, we have  $i = j$  here, so the values can be read from the first row in Table 4 on the top of page 111 of [7].

## 2. PRELIMINARIES

**2.1. Notation.** Let  $E/K$  be an elliptic curve given by the Weierstrass equation

$$E/K : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6,$$

with  $a_1, a_2, a_3, a_4, a_6 \in R$ .

We will also require the following quantities

$$\begin{aligned}
b_2 &= a_1^2 + 4a_2, \\
b_4 &= 2a_4 + a_1a_3, \\
b_6 &= a_3^2 + 4a_6, \\
b_8 &= a_1^2a_6 + 4a_2a_6 - a_1a_3a_4 + a_2a_3^2 - a_4^2, \\
c_4 &= b_2^2 - 24b_4, \\
c_6 &= -b_2^3 + 36b_2b_4 - 216b_6, \\
\Delta &= -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6, \\
j &= c_4^3/\Delta,
\end{aligned}$$

where  $\Delta$  is the *discriminant* of the Weierstrass equation. Note that  $4b_8 = b_2b_6 - b_4^2$  and  $1728\Delta = c_4^3 - c_6^2$ . If  $\text{char}(\overline{K}) \neq 2$ , then  $E/K$  is also given by  $y^2 = 4x^3 + b_2x^2 + 2b_4x + b_6$ .

For positive integers  $n$ , we define the *division polynomials*  $\psi_n, \phi_n \in \mathbb{Z}[a_1, a_2, a_3, a_4, a_6][x, y]$  by

$$\begin{aligned}
\psi_1 &= 1, \\
\psi_2 &= 2y + a_1x + a_3, \\
\psi_3 &= 3x^4 + b_2x^3 + 3b_4x^2 + 3b_6x + b_8, \\
\psi_4 &= \psi_2(2x^6 + b_2x^5 + 5b_4x^4 + 10b_6x^3 + 10b_8x^2 + (b_2b_8 - b_4b_6)x + b_4b_8 - b_6^2),
\end{aligned}$$

and then inductively by the formulas

$$\begin{aligned}
\psi_{2m+1} &= \psi_{m+2}\psi_m^3 - \psi_{m-1}\psi_{m+1}^3 && \text{for } m \geq 2, \\
\psi_2\psi_{2m} &= \psi_m(\psi_{m+2}\psi_{m-1}^2 - \psi_{m-2}\psi_{m+1}^2) && \text{for } m \geq 3.
\end{aligned}$$

We also have

$$\begin{aligned}
\phi_1 &= x, \\
(2.1) \quad \phi_n &= x\psi_n^2 - \psi_{n-1}\psi_{n+1} && \text{for } n \geq 2.
\end{aligned}$$

We will sometimes need expressions for these polynomials that depend only on  $x$ :

$$(2.2) \quad \phi_2(x) = x^4 - b_4x^2 - 2b_6x - b_8,$$

$$(2.3) \quad \psi_2^2(x) = 4x^3 + b_2x^2 + 2b_4x + b_6.$$

In what follows, we will often use  $P$  as the argument of these polynomials, rather than  $x(P)$  and  $y(P)$ .

For a finite extension,  $L$ , of  $K$ , we will use  $R_L$ ,  $\mathcal{M}_L$  and  $v_L$  for the ring of integers of  $L$ , its maximal ideal and the associated valuation, respectively.

We put  $c_v = |E(K)/E_0(K)|$ , where  $E_0(K) = \left\{ P \in E(K) : \tilde{P} \in \tilde{E}(k)_{\text{ns}} \right\}$ , the set of points of  $E(K)$  with non-singular reduction modulo  $\pi$ . This quotient group is known as the component group of  $E$  at  $v$ , so  $c_v$  is the order of the component group.

Tate's Algorithm [12] to compute the special fibre of a Néron model will play a crucial role in many parts of our work. We will use Silverman's presentation of it in [9, Chapter IV, Section 9].

## 2.2. Simplifying $R_n(a, \ell)$ .

**Definition 2.1.** To any pair  $(a, \ell)$  of integers satisfying  $\ell > 0$ , we associate an integer sequence,  $\{R_n(a, \ell)\}_{n \geq 0}$ , defined by

$$R_n(a, \ell) = \frac{n^2 \widehat{a}(\ell - \widehat{a}) - \widehat{na}(\ell - \widehat{na})}{2\ell},$$

where  $\widehat{x}$  denotes the least non-negative residue of  $x$  modulo  $\ell$ .

This sequence is identical to the sequence  $R_n(a, \ell)$  defined in Definition 8.1 of [11], but the expression here is simpler. We prove that now.

**Lemma 2.2.** *Let  $a, \ell, n$  be non-negative integers with  $\ell \geq 1$  and let  $\widehat{x}$  denote the least non-negative residue of  $x$  modulo  $\ell$ . Then*

$$n^2 \widehat{a}(\ell - \widehat{a}) \equiv \widehat{na}(\ell - \widehat{na}) \pmod{2\ell}.$$

*As a consequence,  $R_n(a, \ell)$  here is identical to  $R_n(a, \ell)$  in Definition 8.1 of [11].*

*Proof.* Write  $a = a_1\ell + a_2$  where  $0 \leq a_2 < \ell$  (i.e.,  $\widehat{a} = a_2$ ). Then we can write  $na = (na_1 + a_3)\ell + na_2 - a_3\ell$  with  $0 \leq na_2 - a_3\ell < \ell$  for some integer  $a_3$ .

Thus

$$\begin{aligned} n^2 \widehat{a}(\ell - \widehat{a}) &= n^2 a_2(\ell - a_2) = n^2 a_2 \ell - n^2 a_2^2, \\ \widehat{na}(\ell - \widehat{na}) &= (na_2 - a_3\ell)(\ell - na_2 + a_3\ell) = a_2 n \ell - a_2^2 n^2 + 2a_2 a_3 n \ell - a_3^2 \ell^2 - a_3 \ell^2. \end{aligned}$$

Subtracting these two expressions, we obtain

$$n^2 \widehat{a}(\ell - \widehat{a}) - \widehat{na}(\ell - \widehat{na}) = (n^2 - n) a_2 \ell - 2a_2 a_3 n \ell + (a_3^2 + a_3) \ell^2.$$

Since  $x^2 \pm x$  is even for any integer  $x$ , the congruence in the lemma holds.

The simpler expression for  $R_n(a, \ell)$  is immediate.  $\square$

**2.3. Stange's results for  $v(\psi_n(P))$ .** We next state some theorems from Stange's paper [11]. Following Definition 5.3 of [11], we let

$$(2.4) \quad b \in p\mathbb{Z}^{>0} \cup \{1\}, \quad e \in \mathbb{Z}^{>0}, \quad h \in \mathbb{Z}^{\geq 0}, \quad j \in \mathbb{Z}^{\geq 0}, \quad s \in \mathbb{Z}^{>0} \cup \{\infty\}, \quad w \in \mathbb{Z}^{\geq 0} \cup \{\infty\},$$

and for  $n \in \mathbb{Z}$ , put

$$(2.5) \quad S_n(p, b, e, h, s, w) = \begin{cases} b^j s + \frac{b^j - 1}{b - 1} h + e(v_p(n) - j) + w & v_p(n) > j, \\ b^{v_p(n)} s + \frac{b^{v_p(n)} - 1}{b - 1} h & v_p(n) \leq j, \end{cases}$$

where  $v_p$  is the valuation on  $\mathbb{Q}$  associated to the rational prime,  $p$  (recall our notation in Section 1).

We shall use the following values for these quantities here.

–  $b$  will be the smallest exponent of  $T$  with the valuation  $v$  of its coefficient less than  $v(p)$  in the expansion of multiplication-by- $p$ ,  $[p]T$ , in the formal group of the elliptic curve  $E$ , or else  $b = 1$ , if no such coefficient exists.

–  $e = v(p)$ .

–  $h$  will be the valuation of the coefficient of  $T^b$ , or else  $h = 0$ , if  $b = 1$ .

–  $n_P$  will be the smallest positive integer such that  $\widetilde{[n_P]P} = \widetilde{\mathcal{O}}$ , where  $\widetilde{\cdot}$  is the reduction map from  $E(K)$  to  $\widetilde{E}(k)$  and  $k$  is the residue field.

–  $s = s_P = v(x([n_P]P)/y([n_P]P))$ .

–  $j = j_P = 0$  if  $b = 1$ ; otherwise  $j$  will be the smallest non-negative integer such that

$$e \leq b^j((b-1)s + h).$$

–  $w = w_P = 0$  unless  $b > 1$  and we have equality in the definition of  $j$  above. In this case (i.e.,  $b > 1$  and we have equality in the definition of  $j$ ), put

$$(2.6) \quad w = v\left(\frac{x([p^{j+1}n_P]P)}{y([p^{j+1}n_P]P)}\right) - bv\left(\frac{x([p^j n_P]P)}{y([p^j n_P]P)}\right) - h,$$

which may be equal to  $+\infty$ .

To simplify our notation in what follows, we will often write  $S_n(P)$  instead of  $S_n(p, b, v(p), h, s_P, w_P)$ .

**Remark 2.3.** (i) In the expression for  $w$  in (2.6), we have  $b$  as the coefficient of the second term. This corrects an error in the expression for  $w$  in Lemma 5.1(iii) of [11], where the exponent  $p$  should be  $b$ .

(ii) We have  $S_n \in \mathbb{Z}^{>0} \cup \{\infty\}$ .

(iii) In keeping with the conventional notation of  $e$  for the ramification index, we use  $e$  here, where Stange has used  $d$ .

(iv) When  $b = 1$ , we have  $h = 0$  and  $j = 0$ . In this case, we use the convention  $(b^0 - 1)/(b - 1)h = 0$  to avoid the indeterminate form  $(b^0 - 1)/(b - 1) = 0/0$  in the above expressions for  $S_n(P)$ .

**Lemma 2.4.** ([11], Theorem 6.1) *Assume that  $E$  is in minimal Weierstrass form and  $P$  has non-singular reduction. Then*

$$v(\psi_n(P)) = \min \left\{ 0, \frac{v(x(P))}{2} \right\} n^2 + \begin{cases} S_{n/n_P}(P) & \text{if } n_P \mid n, \\ 0 & \text{if } n_P \nmid n. \end{cases}$$

Furthermore,  $v(x(P)) < 0$  if and only if  $n_P = 1$ .

**Remark 2.5.** From the definition of  $n_P$ ,  $v(x([n]P)) < 0$  if and only if  $n_P \mid n$ . Therefore if  $n_P \nmid n$ , then  $v(\phi_n(P)) \geq v(\psi_n^2(P))$ .

**Lemma 2.6.** ([11], Theorem 9.3) *Suppose that  $E$  is in minimal Weierstrass form with multiplicative reduction,  $P$  has singular reduction, and let  $n_P$  be as above. Then*

$$v(\psi_n(P)) = R_n(a_P, \ell_P) + \begin{cases} S_{n/n_P}(p, p, v(p), 0, s_P, w_P) & n_P \mid n, \\ 0 & n_P \nmid n, \end{cases}$$

where  $\ell_P = -v(j(E))$  and  $a_P$  is the component of the Néron model special fibre ( $\cong \mathbb{Z}/\ell_P\mathbb{Z}$ ) containing  $P$ .

**Lemma 2.7.** (i) *Suppose that  $E$  is in minimal Weierstrass form,  $P$  has non-singular reduction, and  $n_P$  is as above. If  $n_P \mid n$  or  $v(x([n]P)) = 0$ , then*

$$v(\phi_n(P)) = \min \{0, v(x(P))\} n^2.$$

(ii) *Suppose that  $E$  is in minimal Weierstrass form with multiplicative reduction and  $P$  has singular reduction. If  $n_P \nmid n$ , then*

$$v(\phi_n(P)) = 2R_n(a_P, \ell_P),$$

where  $\ell_P = -v(j(E))$  and  $a_P$  is the component of the Néron model special fibre ( $\cong \mathbb{Z}/\ell_P\mathbb{Z}$ ) containing  $P$ .

*Proof.* (i) Suppose that  $n_P | n$ . From Lemma 2.4,

$$v(\psi_n(P)) = \min \left\{ 0, \frac{v(x(P))}{2} \right\} n^2 + S_{n/n_P}(P).$$

We also have

$$v(\phi_n(P)) = v(\psi_n^2(P)) + v(x([n]P)).$$

Since  $n_P | n$ , from the proof of Theorem 6.1 of [11], we have  $v(-x([n]P)/y([n]P)) = S_{n/n_P}(P)$ . From the minimal Weierstrass equation for  $E$ , we have  $3v(x([n]P)) = 2v(y([n]P))$ , since  $v(x([n]P)) < 0$ . Hence  $v(x([n]P)) = -2v(-x([n]P)/y([n]P))$  and so

$$v(\phi_n(P)) = v(\psi_n^2(P)) - 2S_{n/n_P}(P).$$

It follows from our expression above for  $v(\psi_n(P))$  that  $v(\phi_n(P)) = \min \{0, v(x(P))\} n^2$ .

If  $v(x([n]P)) = 0$ , then

$$v(\phi_n(P)) = v(\psi_n^2(P)) + v(x([n]P))v(\phi_n(P)) = v(\psi_n^2(P)).$$

So if  $n_P \nmid n$ , then (i) follows immediately from Lemma 2.4.

(ii) Suppose that  $n_P | n$ . From the proof of Lemma 11.4 in [11],

$$v(\psi_n(P)) = R_n(a_P, \ell_P) + v(x([n]P)/y([n]P)).$$

Therefore,

$$\begin{aligned} v(\phi_n(P)) &= v(\psi_n^2(P)) + v(x([n]P)) \\ &= 2R_n(a_P, \ell_P) + 2v(x([n]P)/y([n]P)) + v(x([n]P)). \end{aligned}$$

Since  $3v(x([n]P)) = 2v(y([n]P))$ , we have

$$v(x([n]P)/y([n]P)) = v(x([n]P)) - v(y([n]P)) = -v(x([n]P))/2.$$

Hence  $v(\phi_n(P)) = 2R_n(a_P, \ell_P)$ . □

**2.4. Non-integral  $x(P)$ .** By Proposition 2.2(iii) of [11], if  $E$  is given by a  $v$ -integral Weierstrass equation, where  $v$  is a nonarchimedean valuation, and  $v(x(P)), v(y(P)) < 0$ , then  $v(\phi_n(P)) = n^2 v(x(P))$ . From the minimal Weierstrass equation for  $E$ , we have  $3v(x(P)) = 2v(y(P))$ , so it suffices that  $v(x(P)) < 0$ .

The degree of  $\psi_n^2(P)$  as a polynomial in  $x(P)$  is  $n^2 - 1$  (see Proposition 2.2(ii) of [11], for example). So if  $E$  is given by a  $v$ -integral Weierstrass equation, where  $v$  is a nonarchimedean valuation, and  $v(x(P)) < 0$ , then  $v(\psi_n^2(P)) \geq (n^2 - 1)v(x(P))$ .

Thus  $k_{v,n}(P) = n^2 v(x(P))$  in this case. Also notice that if  $v(x(P)) < 0$ , then  $\tilde{P} = \tilde{\mathcal{O}}$  and so  $P$  is non-singular modulo  $\pi$ . This establishes Theorem 1.1 in this case.

**2.5. Points with singular reduction.** The following lemma will allow us to use the Weierstrass equations obtained in the course of Tate's Algorithm to simplify our work. Throughout this section, we will let  $P' = (x', y')$  be the image of  $P = (x, y)$  under a change of variables of the form  $x = u^2x' + r$  and  $y = u^3y' + u^2sx' + t$  with  $r, s, t, u \in R$ .

**Lemma 2.8.** (i) *Let  $P = (x, y) \in K^2$  satisfy the equation*

$$f(x, y) = y^2 + a_1xy + a_3y - x^3 - a_2x^2 - a_4x - a_6 = 0,$$

*with  $a_1, a_2, a_3, a_4, a_6 \in R$  and  $v(a_3) > 0$ ,  $v(a_4) > 0$  and  $v(a_6) > 0$ .*

*$P$  has singular reduction if and only if  $v(x) > 0$  and  $v(y) > 0$ .*

(ii) *Let  $n_P$  be as defined in Subsection 2.3. If  $v(u) = 0$ , then  $n_{P'} = n_P$ .*

(iii) *Suppose that  $E$  has multiplicative reduction and  $P \in E(K)$  has singular reduction. Let  $a_P$  be the component of the cyclic group  $E(K)/E_0(K)$  that contains  $P$ . If  $v(u) = 0$ , then  $a_{P'} = a_P$ .*

(iv) *If  $v(u) = 0$ , then  $k_{v,n}(P) = k_{v,n}(P')$ . As a consequence, if  $E$  is in minimal Weierstrass form, then the changes of variable in Tate's Algorithm leave  $k_{v,n}(P)$  unchanged.*

*Proof.* (i)

$$(2.7) \quad \begin{aligned} P \text{ has singular reduction} &\iff v\left(\frac{\partial f}{\partial x}(P)\right) > 0 \text{ and } v\left(\frac{\partial f}{\partial y}(P)\right) > 0 \\ &\iff \begin{cases} v(a_1y - 3x^2 - 2a_2x) > 0 \text{ and} \\ v(2y + a_1x) > 0. \end{cases} \end{aligned}$$

The last logical equivalence here holds using the expressions for  $\partial f/\partial x(P)$  and  $\partial f/\partial y(P)$  since we assume that  $v(a_3) > 0$  and  $v(a_4) > 0$ .

If  $v(x) > 0$  and  $v(y) > 0$ , then it is immediate that  $P$  has singular reduction. So we may assume that  $P$  has singular reduction.

Subtracting 2 times the first expression in (2.7) from  $a_1$  times the second one, we can eliminate  $y$  and obtain

$$v(x(6x + a_1^2 + 4a_2)) > 0.$$

Similarly, using the second expression in (2.7) to eliminate  $y$  from  $f(x, y)$ , we have

$$v(x^2(4x + a_1^2 + 4a_2)) > 0,$$

since  $v(a_3) > 0$  and  $v(a_4) > 0$ .

If  $v(x) > 0$ , then from  $f(x, y) = 0$ ,  $v(a_3) > 0$  and  $v(a_6) > 0$ , we have  $v(y) > 0$ .

Assume that  $v(x) \leq 0$ . Then

$$v(4x + a_1^2 + 4a_2) > 0 \quad \text{and} \quad v(6x + a_1^2 + 4a_2) > 0.$$

Subtracting these two expressions, we obtain  $v(2x) > 0$ , which contradicts  $v(x) \leq 0$  when  $v(2) = 0$ .

If  $v(2) > 0$ , then from (2.7), we have  $v(a_1y - 3x^2) > 0$  and  $v(a_1x) > 0$ . Subtracting  $y$  times the second expression from  $x$  times the first expression, we obtain  $v(3x^2) > 0$ , so  $v(x) > 0$ . As above,  $v(x) > 0$  also implies that  $v(y) > 0$ .

(ii) We know that  $\tilde{P} = \tilde{\mathcal{O}}$  if and only if  $v(x) < 0$ .

Suppose that  $v(x') < 0$ . We have  $v(x) = v(u^2x' + r) \geq \min\{v(x'), v(r)\}$  with equality if  $v(x') \neq v(r)$ . Since  $r \in R$ , we have  $v(r) \geq 0$ . Therefore  $v(x) = v(x') < 0$  and  $n_P | n_{P'}$ . The same argument also shows that  $n_{P'} | n_P$ , so  $n_P = n_{P'}$ .

(iii) This follows from the expression for  $a_P$  in Lemma 5.1 of [10] (denoted there as  $n$ ), using the facts that  $\Delta = u^{12}\Delta'$ ,  $2y + a_1x + a_3 = u^3(2y' + a'_1x' + a'_3)$  and  $v(u) = 0$ .

(iv) By Proposition 2.2(iv) of [11], our change of variables gives

$$(2.8) \quad \psi_n(P) = u^{n^2-1}\psi_n(P').$$

From this and (2.1), we obtain

$$(2.9) \quad \begin{aligned} \phi_n(P) &= x\psi_n^2(P) - \psi_{n-1}(P)\psi_{n+1}(P) \\ &= (u^2x(P') + r)u^{2n^2-2}\psi_n^2(P') - u^{2n^2}\psi_{n-1}(P')\psi_{n+1}(P') \\ &= u^{2n^2-2}(u^2\phi_n(P') + r\psi_n^2(P')). \end{aligned}$$

From (2.8) and  $v(u) = 0$ , we have  $v(\psi_n^2(P')) = v(\psi_n^2(P))$ .

If  $v(\phi_n(P)) < v(\psi_n^2(P)) = v(\psi_n^2(P'))$ , then  $v(\phi_n(P')) = v(\phi_n(P))$  from (2.9). So  $k_{v,n}(P') = v(\phi_n(P')) = k_{v,n}(P)$ .

Suppose that  $v(\phi_n(P)) \geq v(\psi_n^2(P)) = v(\psi_n^2(P'))$ . If  $v(\phi_n(P')) \geq v(r\psi_n^2(P'))$ , then  $k_{v,n}(P') = v(\psi_n^2(P'))$ , as desired. If  $v(\phi_n(P')) < (r\psi_n^2(P'))$ , then  $v(\phi_n(P)) = v(\phi_n(P'))$ , from (2.9). So  $v(\phi_n(P')) \geq v(\psi_n^2(P'))$ , as required.

The statement regarding the changes of variables in Tate's Algorithm now follows because it is only in Step 11 (i.e., when we do not start with a minimal model) that we perform a change of variables with  $v(u) \neq 0$ .  $\square$

**2.6. Additive reduction.** In order to use Stange's results above, we will often need to work in a finite extension of  $K$ . Here we present results on how the valuations behave when we work in such extensions. In addition, in Lemmas 2.10 and 2.11 we provide

some results we will need for the valuation of  $\phi_n(P)$  and  $\psi_n(P)$  in order to finish the proof of Theorem 1.1.

**Lemma 2.9.** *Let  $E/K$  be an elliptic curve having additive reduction and  $P \in E(K)$  having singular reduction.*

*Let  $x = u^2x' + r$  and  $y = u^3y' + u^2sx' + t$  be a change of variables from  $E$  to an elliptic curve  $E'$  in minimal Weierstrass form with  $r, s, t, u \in R_L$ , where  $L$  is a finite extension of  $K$ . Write  $P'$  for the image of  $P$  under this change of variables. Put*

$$T(E) = \begin{cases} v_L(\Delta_E)/6 & \text{if } E' \text{ has good reduction,} \\ v_L(c_4(E))/2 & \text{if } E' \text{ has multiplicative reduction.} \end{cases}$$

(i) *We have  $v_L(u) = T(E)/2$  and*

$$(2.10) \quad v_L(\psi_n(P)) = (n^2 - 1)T(E)/2 + v_L(\psi_n(P')),$$

$$(2.11) \quad v_L(\phi_n(P)) = (n^2 - 1)T(E) + v_L(u^2\phi_n(P') + r\psi_n^2(P')).$$

(ii) *If  $[n]P$  has non-singular reduction,  $v(a_3) > 0$ ,  $v(a_4) > 0$  and  $v(a_6) > 0$ , then*

$$(2.12) \quad v_L(\phi_n(P)) = (n^2 - 1)T(E) + v_L(u^2\phi_n(P')).$$

*Proof.* (i) Equation (2.10) follows from [11]. We use Theorem 7.1 when  $E$  has potential good reduction and Theorem 9.3 when  $E$  has potential multiplicative reduction.

For Equation (2.11), we use (2.9) in the proof of Lemma 2.8(iv):

$$\phi_n(P) = u^{2n^2-2} (u^2\phi_n(P') + r\psi_n^2(P')).$$

Suppose that  $E'$  has good reduction. Since  $\Delta_{E'} = u^{-12}\Delta_E$ , we have  $v_L(\Delta_{E'}) = v_L(\Delta_E) - 12v_L(u)$ . It must equal 0, since  $E'$  has good reduction. Therefore,  $v_L(u) = v_L(\Delta_E)/12$ .

Now suppose that  $E'$  has multiplicative reduction. Since  $c_4(E') = u^{-4}c_4(E)$ , we have  $v_L(c_4(E')) = v_L(c_4(E)) - 4v_L(u)$ . By [8, Proposition VII.5.1(b)],  $v_L(c_4(E')) = 0$ , since  $E'$  has multiplicative reduction. Therefore,  $v_L(u) = v_L(c_4(E))/4$ .

(ii) Putting  $V_{P',n} = v_L(u^2\phi_n(P') + r\psi_n^2(P'))$ , we will show that  $V_{P',n} = v_L(u^2\phi_n(P'))$ .

Since  $[n]P$  has non-singular reduction, by our assumptions on the  $a_i$ 's and Lemma 2.8(i), we have  $v_L(\phi_n(P)) \leq v_L(\psi_n^2(P))$ .

From (2.11), we have  $V_{P',n} = v_L(\phi_n(P)) - (n^2 - 1)T(E)$ . Applying  $v_L(\phi_n(P)) \leq v_L(\psi_n^2(P))$  and then (2.10), we obtain

$$(2.13) \quad V_{P',n} \leq v_L(\psi_n^2(P')).$$

We now consider the case when  $v_L(r) > 0$ . Combining  $V_{P',n} \geq \min(v_L(u^2\phi_n(P')), v_L(r\psi_n^2(P')))$  with (2.13), since  $v_L(r) > 0$ , it follows that  $V_{P',n} = v_L(u^2\phi_n(P'))$ .

Next we consider the case when  $v_L(r) = 0$ . Since  $P$  has singular reduction, by Lemma 2.8(i), we have  $v_L(x(P)) > 0$ . Since  $v_L(r) = 0$ , we have  $v_L(x(P) - r) = 0$ . Therefore

$$v_L(x(P')) = v_L(u^{-2}(x(P) - r)) = -v_L(u^2) < 0,$$

the last inequality holding by [8, Proposition VII.5.1(c)], our definition of  $T(E)$  and the relationship between  $T(E)$  and  $u$ .

Hence  $v_L(x([n]P')) \leq v_L(x(P')) < 0$ . The first inequality here follows from the our arguments in Subsection 2.4, which show that  $v(\phi_n(P')) = n^2 v(x(P'))$  and  $v(\psi_n^2(P')) \geq (n^2 - 1)v(x(P'))$ . Hence  $v_L(\phi_n(P')) - v_L(\psi_n^2(P')) \leq -v_L(u^2)$ , which implies that  $v_L(u^2 \phi_n(P')) \leq v_L(\psi_n^2(P')) = v_L(r \psi_n^2(P'))$ . If  $v_L(u^2 \phi_n(P')) < v_L(r \psi_n^2(P'))$  then  $V_{P',n} = v_L(u^2 \phi_n(P'))$ . If  $v_L(u^2 \phi_n(P')) = v_L(r \psi_n^2(P'))$ , then  $V_{P',n} \geq v_L(r \psi_n^2(P'))$ . Combining this with  $V_{P',n} \leq v_L(r \psi_n^2(P'))$  (from (2.13) and  $v_L(r) = 0$ ), we obtain  $V_{P',n} = v_L(r \psi_n^2(P')) = v_L(u^2 \phi_n(P'))$ .  $\square$

**Lemma 2.10.** *Let  $E/K$  be an elliptic curve having Kodaira type  $I_m^*$  with  $m \geq 1$ , and  $P \in E(K)$  having singular reduction. By Tate's algorithm, we may assume that the Weierstrass equation for  $E$  has  $v(a_1) \geq 1$ ,  $v(a_2) = 1$ ,  $v(a_3) \geq \lfloor m/2 \rfloor + 2$ ,  $v(a_4) \geq \lfloor (m-1)/2 \rfloor + 3$  and  $v(a_6) \geq m+3$ .*

(i) *Let  $m$  be odd. If  $v(x(P)) = 1$ , then  $[2]P$  has non-singular reduction. Here  $v(\phi_2(P)) = v(\psi_3(P)) = 4$  and  $v(\psi_2^2(P)) \geq 4$ .  $1 < v(x(P)) < (m+3)/2$  is not possible. If  $v(x(P)) \geq (m+3)/2$ , then  $[2]P$  has singular reduction. Here  $v(\phi_2(P)) = v(\psi_3(P)) = m+4$  and  $v(\psi_2^2(P)) = m+3$ .*

(ii) *If  $m$  is even, then  $[2]P$  has non-singular reduction. If  $v(x(P)) = 1$ , then  $v(\phi_2(P)) = v(\psi_3(P)) = 4$  and  $v(\psi_2^2(P)) \geq 4$ .  $1 < v(x(P)) < (m+2)/2$  is not possible. If  $v(x(P)) \geq (m+2)/2$ , then  $v(\phi_2(P)) = v(\psi_3(P)) = m+4$  and  $v(\psi_2^2(P)) = m+4$ .*

*Proof.* We write  $x$  in place of  $x(P)$  for convenience in what follows. Since  $v(a_3) > 0$ ,  $v(a_4) > 0$ ,  $v(a_6) > 0$  and  $P$  has singular reduction, from Lemma 2.8(i), we have  $v(x) > 0$ .

(i) Since  $m$  is odd, write  $m = 2k-1$  for some positive integer  $k$ . From the inequalities in the lemma, we have  $v(a_1) \geq 1$ ,  $v(a_2) = 1$ ,  $v(a_3) \geq k+1$ ,  $v(a_4) \geq k+2$  and  $v(a_6) \geq 2k+2$ . In addition, from Proposition 1(a) in Section III of [6] (note that  $n$  there equals our  $k+1$ ), we have  $v(b_6) = 2k+2$  and  $v(b_8) = 2k+3$ . As a consequence, for  $p \geq 3$ , we also have  $v(b_2) = 1$  and  $v(b_4) \geq k+2$ . For  $p = 2$ , we have  $v(b_2) \geq 2$  and  $v(b_4) \geq k+2$ . Furthermore, from the proof of Proposition 1(a) in Section III of [6], we find that  $v(a_3) = k+1$  when  $m$  is odd and  $p = 2$ .

(i-a) Let  $p \geq 3$ . For use with the expression for  $\phi_2(P)$  in (2.2), we have  $v(x^4) = 4v(x)$ ,  $v(b_4x^2) \geq 2v(x) + k + 2$ ,  $v(2b_6x) = v(x) + 2k + 2$  and  $v(b_8) = 2k + 3$ . Similarly, for use in the expression for  $\psi_2^2(P)$  in (2.3), we have  $v(4x^3) = 3v(x)$ ,  $v(b_2x^2) = 2v(x) + 1$ ,  $v(2b_4x) \geq v(x) + k + 2$  and  $v(b_6) = 2k + 2$ .

If  $v(x) = 1$ , then these inequalities imply that  $v(\phi_2(P)) = v(x^4) = 4$  and  $v(\psi_2^2(P)) \geq 4$  (the expression for  $\psi_2^2(P)$  in (2.3) shows that  $v(\psi_2^2(P)) \geq v(4x^3) = 3$ , but  $v(\psi_2^2(P))$  is even), so  $[2]P$  has non-singular reduction.

If  $2 \leq v(x) \leq k$ , then from the inequalities above, we have  $v(2b_4x) \geq v(x) + k + 2 \geq 2v(x) + 2$  and  $v(b_6) = 2k + 2 \geq 2v(x) + 2$ . Hence  $v(\psi_2^2(P)) = v(b_2x^2) = 2v(x) + 1$ , which is not possible.

If  $v(x) \geq k + 1 = (m + 3)/2$ , then  $v(\phi_2(P)) = v(b_8) = 2k + 3$  and  $v(\psi_2^2(P)) = v(b_6) = 2k + 2$ . Thus  $v(x([2]P)) = 1$ , so  $[2]P$  is singular.

(i-b) Let  $p = 2$ . For use with the expression for  $\phi_2(P)$  in (2.2), we have  $v(x^4) = 4v(x)$ ,  $v(b_4x^2) \geq 2v(x) + k + 2$ ,  $v(2b_6x) \geq v(x) + 2k + 3$  and  $v(b_8) = 2k + 3$ . For the expression for  $\psi_2^2(P)$  in (2.3), we have  $v(4x^3) \geq 3v(x) + 2$ ,  $v(b_2x^2) \geq 2v(x) + 2$ ,  $v(2b_4x) \geq v(x) + k + 3$  and  $v(b_6) = 2k + 2$ .

We consider separately three cases according to the value of  $v(x)$ .

If  $v(x) = 1$ , then we have  $v(\phi_2(P)) = v(x^4) = 4$  and  $v(\psi_2^2(P)) \geq v(b_2x^2) \geq 4$ , so  $[2]P$  has non-singular reduction.

Next assume that  $2 \leq v(x) \leq k$ . Using our inequalities for the  $v(a_i)$ 's, we see that  $v(x^3 + a_2x^2 + a_4x + a_6) = v(a_2x^2) = 2v(x) + 1$ , since  $v(a_4x) \geq k + 2 + v(x) \geq 2v(x) + 2$  and  $v(a_6) \geq 2k + 2 \geq 2v(x) + 2$ . We also have  $v(y^2) = 2v(y)$ ,  $v(a_1xy) \geq 1 + v(x) + v(y)$ ,  $v(a_3y) = k + 1 + v(y) \geq v(a_1xy)$ .

If  $v(y) \leq v(x)$ , then  $v(y^2 + a_1xy + a_3y) = v(y^2) \leq 2v(x)$ . If  $v(y) > v(x)$ , then  $v(y^2 + a_1xy + a_3y) > 2v(x) + 1$ . In both cases, the valuation of the left-hand side of the Weierstrass equation differs from the valuation of the right-hand side, so  $2 \leq v(x) \leq k$  is not possible.

Lastly, assume that  $v(x) \geq k + 1$  with  $k \geq 1$ . Then  $v(\phi_2(P)) = v(b_8) = 2k + 3$  and  $v(\psi_2^2(P)) = v(b_6) = 2k + 2$ , so  $v(x([2]P)) = 1$ . Therefore  $[2]P$  has singular reduction.

(ii) Since  $m$  is even, write  $m = 2k$  for some positive integer  $k$ . From the inequalities in the lemma, we have  $v(a_1) \geq 1$ ,  $v(a_2) = 1$ ,  $v(a_3) \geq k + 2$ ,  $v(a_4) \geq k + 2$  and  $v(a_6) \geq 2k + 3$ . In addition, from Proposition 1(b) in Section III of [6], we have  $v(b_8) = 2k + 4$ . So, for  $p \geq 3$ , we have  $v(b_2) = 1$ ,  $v(b_4) \geq k + 2$  and  $v(b_6) \geq 2k + 3$ . For  $p = 2$ , we have  $v(b_2) \geq 2$ ,  $v(b_4) \geq k + 3$  and  $v(b_6) \geq 2k + 4$ . Furthermore, from

the proof of Proposition 1(b) in Section III of [6], we find that  $v(a_4) = k + 2$  when  $m$  is even and  $p = 2$ .

(ii-a) Let  $p \geq 3$ . For use with the expression for  $\phi_2(P)$  in (2.2), we have  $v(x^4) = 4v(x)$ ,  $v(b_4x^2) \geq 2v(x) + k + 2$ ,  $v(2b_6x) \geq v(x) + 2k + 3$  and  $v(b_8) = 2k + 4$ . For use with the expression for  $\psi_2^2(P)$  in (2.3), we have  $v(4x^3) = 3v(x)$ ,  $v(b_2x^2) = 2v(x) + 1$ ,  $v(2b_4x) \geq v(x) + k + 2$ , and  $v(b_6) \geq 2k + 3$ .

If  $v(x) = 1$ , then  $v(\phi_2(P)) = v(x^4) = 4$  and  $v(\psi_2^2(P)) \geq 4$ , so  $[2]P$  is non-singular. If  $2 \leq v(x) \leq k$ , then  $v(\psi_2^2(P)) = v(b_2x^2) = 2v(x) + 1$ , which is not possible. If  $v(x) \geq k + 1$ , then  $v(\psi_2^2(P)) \geq 2k + 4$  and  $v(\phi_2(P)) = v(b_8) = 2k + 4$ . Therefore  $[2]P$  has non-singular reduction.

(ii-b) Let  $p = 2$ . From the expression for  $\phi_2(P)$  in (2.2), we have  $v(x^4) = 4v(x)$ ,  $v(b_4x^2) \geq 2v(x) + k + 3$ ,  $v(2b_6x) \geq v(x) + 2k + 5$  and  $v(b_8) = 2k + 4$ . From the expression for  $\psi_2^2(P)$  in (2.3), we have  $v(4x^3) \geq 3v(x) + 2$ ,  $v(b_2x^2) \geq 2v(x) + 2$ ,  $v(2b_4x) \geq v(x) + k + 4$  and  $v(b_6) \geq 2k + 4$ .

If  $v(x) = 1$ , then  $v(\phi_2(P)) = v(x^4) = 4$  and  $v(\psi_2^2(P)) \geq 4$ , so  $[2]P$  has non-singular reduction.

Assume that  $2 \leq v(x) \leq k$ . We proceed in the same way as for such  $v(x)$  in case (i-b).

Assume that  $v(x) \geq k + 1$ . We have  $v(4x^3) \geq 3k + 5$ ,  $v(b_2x^2) \geq 2k + 4$ ,  $v(2b_4x) \geq 2k + 5$  and  $v(b_6) \geq 2k + 4$ . Therefore  $v(\psi_2^2(P)) \geq 2k + 4$ . We also have  $v(x^4) \geq 4k + 4$ ,  $v(b_4x^2) \geq 3k + 5$ ,  $v(2b_6x) \geq 3k + 6$  and  $v(b_8) = 2k + 4$ . Therefore  $v(\phi_2(P)) = 2k + 4$  and hence  $[2]P$  has non-singular reduction.  $\square$

**Lemma 2.11.** *Let  $E/K$  be an elliptic curve. Suppose  $P \in E(K)$  has singular reduction. Upon applying Tate's algorithm, we have the following relationships.*

- (i) *If  $m_P = 2$ , then  $v(\phi_2(P)) = v(\psi_3(P))$ .*
- (ii) *If  $m_P = 3$  and  $E/K$  has additive reduction, then  $v(\phi_3(P)) = 3v(\psi_2^2(P))$ .*

*Proof.* All assertions below for the values of the  $v(a_i)$ 's and  $v(b_i)$ 's can be found in Silverman's presentation of Tate's Algorithm in [9, Chapter IV, Section 9]. We will also use, sometimes implicitly, the fact that  $v(x(P)) > 0$ . This follows from Lemma 2.8(i).

(i) Since  $m_P = 2$ , it follows that  $v(\phi_2(P)) \leq v(\psi_2^2(P))$ . From Lemma 2.8(i), we have  $v(x(P)) > 0$ . So from the formula  $\phi_2 = x\psi_2^2 - \psi_1\psi_3$  (which follows from (2.1) with  $n = 2$ ) we obtain  $v(\phi_2(P)) = v(\psi_1(P)\psi_3(P)) = v(\psi_3(P))$ .

(ii) We again use (2.1), here with  $n = 3$ , so we have  $\phi_3 = x\psi_3^2 - \psi_2\psi_4$ . Since  $m_P = 3$ , it follows that  $v(\phi_3(P)) \leq v(\psi_3^2(P))$  by Lemma 2.8(i). Thus  $v(\phi_3(P)) = v(\psi_2(P)\psi_4(P))$ , so the result would follow if we showed that  $v(\psi_4(P)) = 5v(\psi_2(P))$ .

Since  $m_P = 3$ , it follows that  $3|c_v$ , so since  $E/K$  has additive reduction, either  $E/K$  has Kodaira type  $IV$  or Kodaira type  $IV^*$ . We consider these two cases separately.

If  $E/K$  has Kodaira type  $IV$ , then we are in Step 5 of Tate's Algorithm. We have  $v(b_2) \geq 1$ ,  $v(b_4) \geq 2$ ,  $v(b_6) = 2$  and  $v(b_8) \geq 3$ . By using the expression for  $\psi_2^2(P)$  in (2.3), we obtain  $v(\psi_2^2(P)) = v(b_6) = 2$ . From the expression for  $\psi_4(P)$  in Subsection 2.1, we have  $v(\psi_4(P)) = v(\psi_2(P)) + v(b_6^2) = 5 = 5v(\psi_2(P))$ , as desired.

If  $E/K$  has Kodaira type  $IV^*$ , then we are in Step 8 of Tate's Algorithm, where  $v(a_1) \geq 1$ ,  $v(a_2) \geq 2$ ,  $v(a_3) \geq 2$ ,  $v(a_4) \geq 3$  and  $v(a_6) \geq 4$ . In addition,  $v(a_3^2/\pi^4 + 4a_6/\pi^4) = v(a_3^2 + 4a_6) - 4 = 0$ . If  $p = 2$ , then  $v(b_2) \geq 2$ ,  $v(b_4) \geq 3$ ,  $v(b_6) = 4$  (from  $v(a_3^2 + 4a_6) - 4 = 0$  above) and  $v(b_8) \geq 6$ . If  $p \geq 3$ , then  $v(b_2) \geq 2$ ,  $v(b_4) \geq 3$ ,  $v(b_6) = 4$  and  $v(b_8) \geq 4$ . Proceeding in the same way as for Kodaira type  $IV$ , we have  $v(\psi_2^2(P)) = v(b_6) = 4$  and  $v(\psi_4(P)) = v(\psi_2(P)) + v(b_6^2) = 10 = 5v(\psi_2(P))$ .  $\square$

### 3. PROOF OF THEOREM 1.1

**3.1. Multiplicative reduction.** Suppose that  $E$  has multiplicative reduction and that  $P$  is singular modulo  $\pi$ .

As in the proof of Lemma 11.3 of [11], we have

$$k_{v,n}(P) = 2v(\psi_n(P)) - 2v(D_n) = 2R_n(a_{P,v}, m),$$

where  $a_{P,v}$  is the component of the Néron model special fibre ( $\cong \mathbb{Z}/m\mathbb{Z}$ ) containing  $P$ .

For non-split multiplicative reduction, we still have  $k_{v,n}(P) = 2R_n(a_{P,v}, m)$  with  $m$  even and  $a_{P,v} = m/2$ .

From Proposition 5(c) in [2], we see that the coefficient of  $n^2$  in  $2R_n(a_{P,v}, m)$  is as stated in our theorem and in the entry for  $I_m$  in Table 1.1. The remaining term matches  $\epsilon_{v,n}(P)$  in the entry for  $I_m$  in Table 1.1 by a simple elementary manipulation.

**3.2. Additive reduction.** Throughout this section, by Tate's algorithm, we may assume that the Weierstrass equation for  $E$  has  $v(a_3) > 0$ ,  $v(a_4) > 0$  and  $v(a_6) > 0$ .

By Proposition 5.5 in Chapter VII of [8],  $E$  has potential good reduction if and only if its  $j$ -invariant is integral. By Tableaux  $I$ ,  $II$  and  $IV$  of [6], we compute the values of  $v(j_E) = v(c_4^3(E)/\Delta_E)$ , and find that if the Kodaira type of  $E$  is  $III$ ,  $IV$ ,  $III^*$ ,  $IV^*$  or  $I_0^*$ , then  $v(j_E) \geq 0$ . For Kodaira type  $I_m^*$  with  $m \geq 1$ , if  $p \geq 3$ , then  $v(\Delta_E) = 6 + m$  and

$v(c_4(E)) = 2$ , so  $v(j_E) = -m < 0$ . Therefore such  $E$  have potential good reduction if and only if  $p = 2$  and  $v(j_E) \geq 0$ .

We write  $x(P) = x$  and  $x(P') = x'$  for convenience.

**Lemma 3.1.** *Suppose that  $E/K$  is an elliptic curve having additive reduction and  $P \in E(K)$  having singular reduction.*

(i) *There exists a finite extension,  $L$ , of  $K$  such that  $E$  has either good or multiplicative reduction over  $K$ . So there exists a change of variables,  $x = u^2x' + r$  and  $y = u^3y' + u^2sx' + t$ , from  $E$  to an elliptic curve  $E'$  in minimal Weierstrass form with  $r, s, t, u \in R_L$ . Let  $P'$  be the image of  $P$  under this change of variables.*

(ii) *Let  $P'$  be as in part (i). If  $P'$  has singular reduction, then  $\ell_{P'} = 2a_{P'} = v_L(\Delta_{E'})$ , where  $\ell_{P'} = -v(j(E'))$  and  $a_{P'}$  is the component of the Néron model special fibre ( $\cong \mathbb{Z}/\ell_{P'}\mathbb{Z}$ ) containing  $P'$ .*

*Proof.* (i) This is the semi-stable reduction theorem. See Proposition VII.5.4 of [8].

(ii) Since  $P'$  has singular reduction,  $E'$  has multiplicative reduction by part (i) and  $E$  has Kodaira type  $I_m^*$  with  $m \geq 1$  (see, for example, the passage below Théorème 2 on page 121 of [6]) and  $v(j_E) = v(c_4^3(E)/\Delta_E) < 0$ . Here  $c_v = 2$  or  $4$ .

From Tableaux I–III and Théorème 3 on page 121 in [6], we find that

$$(3.1) \quad v(\Delta_E) = m + 4 + v(c_4(E)).$$

From Lemma 2.9, we know that  $v_L(u) = T(E)/2 = v_L(c_4(E))/4$ . Combining this with the expression for  $v_L(\Delta_{E'})$  in Table 3.1 on page 45 of [8], it follows that

$$(3.2) \quad v_L(\Delta_{E'}) = v_L(\Delta_E) - 3v_L(c_4(E)) = (m + 4)e_L - 2v_L(c_4(E)).$$

By Proposition 2.2(iv) of [11] and  $v_L(u) = v_L(c_4(E))/4$ ,

$$v_L(\psi_2(P')) = v_L(\psi_2(P)) - 3v_L(u) = v_L(\psi_2(P)) - \frac{3}{4}v_L(c_4(E)).$$

By Lemma 5.1 of [10],

$$a_{P'} = \min \{v_L(\psi_2(P')), v_L(\Delta_{E'})/2\}.$$

Applying (3.1) and then (3.2), we have

$$\ell_{P'} = -v_L(j_E) = -v_L(c_4^3(E)/\Delta_E) = (m + 4)e_L - 2v_L(c_4(E)) = v_L(\Delta_{E'}).$$

We consider two cases in order to determine  $a_{P'}$ .

(a) First, assume that  $[2]P$  has non-singular reduction. We establish some information about  $n_{P'}$ .

If  $v_L(x(P')) < 0$ , then  $n_{P'} = 1$ .

Next we consider  $v_L(x(P')) \geq 0$ . We saw above that  $v_L(u^2) = v_L(c_4(E))/2$ . So  $v_L(u^2) > 0$ , by Proposition VII.5.1(c) of [8]. Since  $v_L(x(P')) \geq 0$  and  $v_L(u^2) > 0$ , it follows that  $v_L(x(P) - r) \geq v_L(u^2) > 0$ . Thus, since  $v_L(x(P)) > 0$  by Lemma 2.8(i), we must also have  $v_L(r) > 0$ . Since  $[2]P$  is non-singular, by Lemma 2.8(i) we have  $v_L(x([2]P)) \leq 0$ . As a result,

$$v_L(x([2]P')) = v_L(u^{-2}(x([2]P) - r)) \leq -2v_L(u) < 0,$$

and hence  $n_{P'}|2$ .

With this information about  $n_{P'}$ , we proceed to determine  $a_{P'}$ .

Suppose that  $v_L(\psi_2(P')) < v_L(\Delta_{E'})/2$ . Then  $a_{P'} = v_L(\psi_2(P'))$ . From  $\ell_{P'} = v_L(\Delta_{E'})$ , we have  $\ell_{P'} > 2a_{P'}$ . Then  $\widehat{a_{P'}} = a_{P'}$  and  $\widehat{2a_{P'}} = 2a_{P'}$ , where  $\widehat{x}$  denotes the least non-negative residue of  $x$  modulo  $\ell_{P'}$ . Hence

$$R_2(a_{P'}, \ell_{P'}) = \frac{4a_{P'}(\ell_{P'} - a_{P'})}{2\ell_{P'}} - \frac{2a_{P'}(\ell_{P'} - 2a_{P'})}{2\ell_{P'}} = a_{P'},$$

where  $R_n$  is as defined in Definition 2.1.

From Lemma 2.7(ii), we have  $v_L(\phi_2(P')) = 2R_2(a_{P'}, \ell_{P'})$ . So  $v_L(\phi_2(P')) = 2a_{P'} = v_L(\psi_2^2(P'))$ . Therefore  $v_L(x([2]P')) = 0$ , which contradicts  $n_{P'} = 1$  or 2. So  $v_L(\psi_2(P')) \geq v_L(\Delta_{E'})/2$ . Hence  $a_{P'} = v_L(\Delta_{E'})/2$  and it follows that  $\ell_{P'} = 2a_{P'}$ .

(b) Next, assume that  $[2]P$  has singular reduction. From Lemma 2.10, this only happens when  $m$  is odd and  $v(\psi_2^2(P)) = m + 3$ . Using this, Proposition 2.2(iv) of [11] (again noting that our change of variables is defined using Silverman's convention, which differs from Stange's in [11]) and  $v_L(u) = v_L(c_4(E))/4$ , we find that

$$v_L(\psi_2(P')) = v_L(\psi_2(P)) - 3v_L(u) = \frac{(m+3)e_L}{2} - \frac{3v_L(c_4(E))}{4}.$$

We find that  $v(c_4(E)) \geq 2$  from Tableaux I–V of [6]. Applying this and (3.2) to the preceding expression for  $v_L(\psi_2(P'))$ , we obtain

$$v_L(\psi_2(P')) - \frac{v_L(\Delta_{E'})}{2} = \frac{v_L(c_4(E)) - 2e_L}{4} \geq 0.$$

So  $v_L(\psi_2(P')) \geq v_L(\Delta_{E'})/2$ . Hence  $a_{P'} = v_L(\Delta_{E'})/2$  and it follows that  $\ell_{P'} = 2a_{P'}$ .  $\square$

**Lemma 3.2.** *Suppose that  $E/K$  is an elliptic curve having additive reduction and  $P \in E(K)$  having singular reduction. Let  $E'$ ,  $L$ ,  $P'$  and  $r$  be as in Lemma 3.1(i).*

(i) *If  $n \equiv 0 \pmod{m_P}$  and  $P'$  has non-singular reduction, then*

$$(3.3) \quad k_{v,n}(P) = v(\phi_n(P)) = \begin{cases} v(x(P) - r)n^2 & \text{if } v_L(x(P')) \leq 0, \\ v(u^2)n^2 & \text{if } v_L(x(P')) > 0. \end{cases}$$

(ii) If  $n \equiv 0 \pmod{m_P}$  and  $P'$  has singular reduction, then

$$(3.4) \quad k_{v,n}(P) = v(\phi_n(P)) = \frac{1}{4}(m+4)n^2.$$

(iii) If  $n \equiv \pm 1 \pmod{m_P}$  and  $P'$  has non-singular reduction, then

$$(3.5) \quad k_{v,n}(P) = v(\psi_n^2(P)) = v(x(P) - r)(n^2 - 1).$$

(iv) If  $n \equiv \pm 1 \pmod{m_P}$  and  $P'$  has singular reduction, then

$$(3.6) \quad k_{v,n}(P) = v(\psi_n^2(P)) = \frac{1}{4}(m+4)(n^2 - 1).$$

*Proof.* We will first show that when  $n \equiv 0 \pmod{m_P}$ , we have  $n_{P'} | n$ .

If  $v_L(x(P')) < 0$ , then  $n_{P'} = 1$ , and so  $n_{P'} | n$ . If  $v_L(x(P')) \geq 0$ , then using the same argument as in case (a) of the proof of Lemma 3.1(ii), we find that  $n_{P'} | n$ .

(i) Since  $v(x([n]P)) \leq 0$ , we have  $v(\phi_n(P)) \leq v(\psi_n^2(P))$  by Lemma 2.8(i). Therefore  $k_{v,n}(P) = v(\phi_n(P))$ .

We first assume that  $v_L(x(P')) \leq 0$ . We can apply Lemma 2.7(i), and obtain  $v_L(\phi_n(P')) = v_L(x(P'))n^2$ . So by (2.12) in Lemma 2.9(ii) with  $T(E) = v_L(u^2)$ ,

$$\begin{aligned} v_L(\phi_n(P)) &= (n^2 - 1)v_L(u^2) + v_L(u^2\phi_n(P')) = (v_L(u^2) + v_L(x(P'))n^2) \\ &= v_L(x(P) - r)n^2. \end{aligned}$$

Now assume that  $v_L(x(P')) > 0$ . Applying Lemma 2.7(i), we obtain  $v_L(\phi_n(P')) = 0$ . As above, we have

$$v_L(\phi_n(P)) = (n^2 - 1)v_L(u^2) + v_L(u^2\phi_n(P')) = v_L(u^2)n^2.$$

(ii) Since  $P'$  has singular reduction, it follows that  $E'$  has multiplicative reduction. From Lemma 3.1(ii), we know that  $\ell_{P'} = 2a_{P'} = v_L(\Delta_{E'})$ .

Since  $m_P$  is even for  $I_m^*$ , from our assumption that  $n \equiv 0 \pmod{m_P}$ , we see that  $n$  is even. So  $na_{P'} \equiv 0 \pmod{\ell_{P'}}$ . Therefore

$$(3.7) \quad R_n(a_{P'}, \ell_{P'}) = \frac{n^2 a_{P'} (2a_{P'} - a_{P'})}{4a_{P'}} = \frac{a_{P'} n^2}{4}.$$

Since  $n \equiv 0 \pmod{m_P}$ , we have  $k_{v,n}(P) = v(\phi_n(P))$ . Since  $E'$  has multiplicative reduction and  $n_{P'} | n$ , from Lemma 2.7(ii) we have  $v_L(\phi_n(P')) = 2R_n(a_{P'}, \ell_{P'})$ . From (2.12) in Lemma 2.9, followed by (3.2) and (3.7), and noting that  $v_L(u) = v_L(c_4(E))/4$ ,

we obtain

$$\begin{aligned}
v_L(\phi_n(P)) &= v_L(u^2)(n^2 - 1) + v_L(u^2\phi_n(P')) \\
&= v_L(u^2)n^2 + 2R_n(a_{P'}, \ell_{P'}) \\
&= v_L(u^2)n^2 + \frac{a_{P'}n^2}{2} \\
&= v_L(u^2)n^2 + \frac{v_L(\Delta_{E'})}{4} \\
&= \frac{1}{2}v_L(c_4(E))n^2 + \frac{n^2}{2} \left( \frac{(m+4)e_L}{2} - v_L(c_4(E)) \right) \\
&= \frac{1}{4}(m+4)n^2e_L.
\end{aligned}$$

Hence

$$k_{v,P}(P) = v(\phi_n(P)) = \frac{1}{4}(m+4)n^2.$$

If  $n \equiv \pm 1 \pmod{m_P}$ , then  $[n]P$  has singular reduction and  $v(x([n]P)) > 0$ , by Lemma 2.8(i). Hence  $v(\phi_n(P)) > v(\psi_n^2(P))$  and so  $k_{v,n}(P) = v(\psi_n^2(P))$ .

We can write  $n = k \pm 1$  for some positive integer  $k$  with  $k \equiv 0 \pmod{m_P}$ .

(iii) We break the proof into two parts, depending on  $v_L(x(P'))$ .

Assume first that  $v_L(x(P')) \geq 0$ . By the same argument as at the start of the proof, we have  $n_{P'} \mid k$  and since  $n_{P'} > 1$  (by our assumption that  $v_L(x(P')) \geq 0$ ), we have  $n_{P'} \nmid n$ . So by Lemma 2.4 we have  $v_L(\psi_n(P')) = 0$ . By using (2.10) in Lemma 2.9 with  $T(E) = v_L(u^2)$ , we obtain

$$v_L(\psi_n^2(P)) = v_L(u^2)(n^2 - 1) = v_L(x(P) - r)(n^2 - 1).$$

Now assume that  $v_L(x(P')) < 0$ . Then  $n_{P'} = 1$  and  $n_{P'} \mid n$ . From Lemma 2.4,

$$v_L(\psi_n^2(P')) = v_L(x(P'))n^2 + 2S_n(P').$$

From the definition of  $S_n(P')$  in (2.5) along with the conditions in (2.4), we have  $S_n(P') \geq s_{P'} = v_L(x(P')/y(P'))$ . Noting that  $3v_L(x(P')) = 2v_L(y(P'))$ , we obtain  $s_{P'} = -v_L(x(P'))/2$ . Hence

$$v_L(\psi_n^2(P')) \geq v_L(x(P'))(n^2 - 1)$$

and so, from (2.10) in Lemma 2.9(i),

$$v_L(\psi_n^2(P)) \geq v_L(u^2)(n^2 - 1) + v_L(x(P'))(n^2 - 1) = v_L(x(P) - r)(n^2 - 1).$$

We can write

$$(3.8) \quad v_L(\psi_n^2(P)) = v_L(x(P) - r)(n^2 - 1) + \alpha_n,$$

where  $\alpha_n \geq 0$ . Since  $n_{P'} = 1$ , this argument works for any value of  $n$ , not just  $n \equiv \pm 1 \pmod{m_p}$ . Hence (3.8) holds for all  $n$  too.

We will now prove that, in fact,  $v_L(\psi_n^2(P)) = v_L(x(P) - r)(n^2 - 1)$ . Since  $v_L(x([k]P)) \leq 0$  and  $v_L(x(P)) > 0$ , we have  $v_L(\phi_k(P)) \leq v_L(\psi_k^2(P)) < v_L(x(P)\psi_k^2(P))$ , so from the formula  $\phi_k = x\psi_k^2 - \psi_{k-1}\psi_{k+1}$  and applying (3.8) with  $k-1$  and  $k+1$  instead of  $n$ , we obtain

$$(3.9) \quad v_L(\phi_k(P)) = v_L(\psi_{k-1}(P)\psi_{k+1}(P)) = v_L(x(P) - r)k^2 + \alpha_{k-1}/2 + \alpha_{k+1}/2.$$

On the other hand, since  $k \equiv 0 \pmod{m_P}$ , from part (i) of this lemma, we have  $v_L(\phi_k(P)) = v_L(x(P) - r)k^2$ . Therefore  $\alpha_{k-1} = \alpha_{k+1} = 0$ .

(iv) Here  $E'$  has multiplicative reduction. So by Lemma 3.1(ii), we know that  $a_{P'} = v_L(\Delta_{E'})/2$  and  $\ell_{P'} = 2a_{P'}$ .

Since  $m_P = 2$  or  $4$ , we see that  $n$  is odd. So  $na_{P'} \equiv a_{P'} \pmod{2a_{P'}}$ . From  $\ell_{P'} = 2a_{P'}$ , we have  $\widehat{a_{P'}} = a_{P'}$  and  $\widehat{na_{P'}} = a_{P'}$ , where  $\widehat{x}$  denotes the least non-negative residue of  $x$  modulo  $\ell_{P'}$ . Hence

$$(3.10) \quad v_L(\psi_n(P')) = R_n(a_{P'}, \ell_{P'}) = \frac{a_{P'}(n^2 - 1)}{4}.$$

By (2.10) in Lemma 2.9, followed by (3.2) and (3.10), and noting that  $v_L(u) = v_L(c_4(E))/4$ , we obtain

$$\begin{aligned} v_L(\psi_n^2(P)) &= \frac{1}{2}v_L(c_4(E))(n^2 - 1) + \frac{1}{2}(n^2 - 1)\left(\frac{(m+4)e_L}{2} - v_L(c_4(E))\right) \\ &= \frac{1}{4}(m+4)e_L(n^2 - 1). \end{aligned}$$

Hence

$$k_{v,n}(P) = v(\psi_n^2(P)) = \frac{1}{4}(m+4)(n^2 - 1).$$

□

**Lemma 3.3.** *Suppose that  $E/K$  is an elliptic curve having additive reduction and  $P \in E(K)$  having singular reduction.*

(i) *If  $c_v = 2$ , then*

$$k_{v,n}(P) = \begin{cases} v(\psi_3(P))n^2/4 & \text{if } n \equiv 0 \pmod{2}, \\ v(\psi_3(P))(n^2 - 1)/4 & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

(ii) *If  $c_v = 3$ , then*

$$k_{v,n}(P) = \begin{cases} v(\psi_2^2(P))n^2/3 & \text{if } n \equiv 0 \pmod{3}, \\ v(\psi_2^2(P))(n^2 - 1)/3 & \text{if } n \equiv 1, 2 \pmod{3}. \end{cases}$$

(iii) *Assume that  $c_v = 4$ .*

If  $[2]P$  has non-singular reduction, then

$$k_{v,n}(P) = \begin{cases} v(\psi_3(P)) n^2/4 & \text{if } n \equiv 0 \pmod{2}, \\ v(\psi_3(P)) (n^2 - 1)/4 & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

If  $[2]P$  has singular reduction, then

$$k_{v,n}(P) = \begin{cases} v(\psi_3(P)) n^2/4 & \text{if } n \equiv 0 \pmod{4}, \\ v(\psi_3(P)) (n^2 - 1)/4 & \text{if } n \equiv 1, 3 \pmod{4}, \\ v(\psi_3(P)) n^2/4 - 1 & \text{if } n \equiv 2 \pmod{4}. \end{cases}$$

*Proof.* Let  $E'$ ,  $L$ ,  $P'$  and the change of variables be as in Lemma 3.1(i).

(i) Since  $c_v = 2$ , it follows that  $E$  has Kodaira type  $III$ ,  $III^*$  or  $I_m^*$  and that  $m_P = 2$ , since  $P$  has singular reduction.

(i-a) We first assume that  $P'$  has non-singular reduction and  $v_L(x(P')) > 0$ . From (3.3) with  $n = 2$ , we have  $v(\phi_2(P)) = 4v(u^2)$ . From (3.5) with  $n = 3$ , we have  $v(\psi_3(P)) = 4v(x(P) - r)$ . From Lemma 2.11(i) along with Lemma 2.10, we have  $v(\phi_2(P)) = v(\psi_3(P))$ , so  $v(\psi_3(P)) = v(u^2)$  if  $n \equiv 0 \pmod{2}$  and  $v(\psi_3(P)) = v(x(P) - r)$  if  $n \equiv 1 \pmod{2}$ .

Therefore, by parts (i) and (iii) of Lemma 3.2,  $k_{v,n}(P) = v(u^2) n^2 = v(\psi_3(P)) n^2/4$  if  $n \equiv 0 \pmod{2}$  and  $k_{v,n}(P) = v(x(P) - r) (n^2 - 1) = v(\psi_3(P)) (n^2 - 1)/4$  if  $n \equiv 1 \pmod{2}$ .

(i-b) Next assume that  $P'$  has non-singular reduction and  $v_L(x(P')) \leq 0$ . From (3.5) with  $n = 3$ , we have  $v(\psi_3(P)) = 4v(x(P) - r)$ .

Therefore, by parts (i) and (iii) of Lemma 3.2,  $k_{v,n}(P) = v(x(P) - r) n^2 = v(\psi_3(P)) n^2/4$  if  $n \equiv 0 \pmod{2}$  and  $k_{v,n}(P) = v(x(P) - r) (n^2 - 1) = v(\psi_3(P)) (n^2 - 1)/4$  if  $n \equiv 1 \pmod{2}$ .

(i-c) Next assume that  $P'$  has singular reduction. This can only happen if  $E$  has Kodaira type  $I_m^*$ , so by Lemma 2.10, we have again  $v(\psi_3(P)) = m + 4$ . Therefore, by parts (ii) and (iv) of Lemma 3.2, we obtain the desired result here too.

(ii) Since  $c_v = 3$ ,  $E$  has Kodaira type  $IV$  or  $IV^*$  and  $m_P = 3$ , since  $P$  has singular reduction. This case happens when  $E$  has potential good reduction. So  $P'$  has non-singular reduction.

From (3.3) with  $n = 3$ , we have either  $v(\phi_3(P)) = 9v(u^2)$  or  $v(\phi_3(P)) = 9v(x(P) - r)$ . In either case, we find by (3.3) that  $v(\phi_n(P)) = v(\phi_3(P)) n^2/9$  holds for  $n \equiv 0 \pmod{3}$ . By part (ii) of Lemma 2.11, we have  $v(\phi_3(P)) = 3v(\psi_2^2(P))$ , so  $v(\phi_n(P)) = v(\psi_2^2(P)) n^2/3$ .

From (3.5) with  $n = 2$ , we have  $v(\psi_2^2(P)) = 3v(x(P) - r)$ . So if  $n \equiv \pm 1 \pmod 3$ , then  $v(\psi_n(P)) = v(\psi_2^2(P))(n^2 - 1)/3$ , using (3.5) again.

(iii) Lastly, we consider  $c_v = 4$ . This case happens only when the Kodaira type is  $I_m^*$ .

(iii-a) Assume that  $[2]P$  has non-singular reduction. Thus,  $m_P = 2$  and  $v(\phi_2(P)) \leq v(\psi_2^2(P))$  by Lemma 2.8(i). Since  $P$  has singular reduction, we have  $v(x(P)) \geq 1$  (again, by Lemma 2.8(i)).

If  $n \equiv 0 \pmod 2$ , then (3.3) or (3.4) holds. As shown in the proof of part (i),  $k_{v,n} = v(\psi_3(P))n^2/4$ .

Similarly, if  $n \equiv 1 \pmod 2$ , then we can apply (3.5) and (3.6) as in the proof of part (i) to show that  $k_{v,n} = v(\psi_3(P))(n^2 - 1)/4$ .

(iii-b) Assume that  $[2]P$  has singular reduction. From Lemma 2.10(ii), this case happens only when  $m$  is odd. In this case,  $m_P = 4$ .

If  $n \equiv \pm 1 \pmod 4$ , then  $[n]P$  is singular, so  $k_{v,n} = v(\psi_n^2(P))$ . By Lemma 2.10(i),  $v(\psi_3(P)) = m + 4$ . If  $P'$  has singular reduction, then by (3.6), we obtain  $k_{v,n} = v(\psi_3(P))(n^2 - 1)/4$ .

If  $P'$  has non-singular reduction, then by (3.5), we obtain  $k_{v,n} = v(x(P) - r)(n^2 - 1)$ . Applying (3.5), but with  $n = 3$ , we find that  $8v(x(P) - r) = 2v(\psi_3(P))$ , so the desired result follows.

Assume that  $n \equiv 0 \pmod 4$ . If  $P'$  has singular reduction, then (3.4) applies. So, from Lemma 2.10(i) we have  $v(\psi_3(P)) = m + 4$ . Therefore  $k_{v,n} = v(\psi_3(P))n^2/4$ .

If  $P'$  has non-singular reduction, then by (3.3), we obtain  $k_{v,n} = v(x(P) - r)n^2$  if  $v_L(x(P')) \leq 0$  and  $k_{v,n} = v(u^2)n^2$ , if  $v_L(x(P')) > 0$ .

We saw above that  $8v(x(P) - r) = 2v(\psi_3(P))$ , so  $k_{v,n} = v(\psi_3(P))n^2/4$ , if  $v_L(x(P')) \leq 0$ .

Applying (3.3) with  $n = 4$ , we find that  $16v(u^2) = v(\phi_4(P))$  if  $v_L(x(P')) > 0$ . So  $k_{v,n} = v(\phi_4(P))n^2/16$ , if  $v_L(x(P')) > 0$ .

Using (3.9) with  $k = 4$  (note that it is applicable since  $[4]P$  is non-singular and  $P$  is singular), we have  $v(\phi_4(P)) = v(\psi_3(P)\psi_5(P))$ . By (3.5) and part (iii) of this lemma with  $n = 5$  (note that we have already proven part (iii) for such  $n$ ), we have  $v(\psi_5^2(P)) = 6v(\psi_3(P))$ , so  $v(\psi_5(P)) = 3v(\psi_3(P))$ . Hence  $v(\phi_4(P)) = 4v(\psi_3(P))$ , and our desired result follows.

Assume that  $n \equiv 2 \pmod 4$ . Since  $[n]P$  has singular reduction and  $[2n]P$  has non-singular reduction, from Lemma 2.10(i), it must be that  $v(x([n]P)) = 1$ . Therefore  $v(\psi_n^2(P)) = v(\phi_n(P)) - 1$ . Lemma 2.10(i) tells us that  $v(x(P)) \geq 2$ , so  $v(\phi_n(P)) <$

$v(x(P)\psi_n^2(P))$ . From the formula  $\phi_n = x\psi_n^2 - \psi_{n-1}\psi_{n+1}$  along with the expressions just found when  $n \equiv \pm 1 \pmod{4}$ , we obtain

$$\begin{aligned} v(\phi_n(P)) &= v(\psi_{n-1}(P)\psi_{n+1}(P)) \\ &= \frac{1}{4}v(\psi_3(P))((n-1)^2 - 1) + \frac{1}{4}v(\psi_3(P))((n+1)^2 - 1) \\ &= \frac{1}{4}v(\psi_3(P))n^2. \end{aligned}$$

□

We now show how Theorem 1.1 follows when  $E$  has additive reduction.

If  $E$  has Kodaira type  $IV$  or  $IV^*$ , then

$$(3.11) \quad \frac{1}{3}v(\psi_2^2(P)) = -2\frac{[K:\mathbb{Q}_p]}{\log|k|}\lambda_v(P),$$

by Proposition 5(d) in [2] (noting that our definition of  $\lambda_v$ , following [10], is one-half that in [2]).

If  $E$  has Kodaira type  $III$ ,  $III^*$ ,  $I_0^*$  or  $I_m^*$ , then

$$(3.12) \quad \frac{1}{4}v(\psi_3(P)) = -2\frac{[K:\mathbb{Q}_p]}{\log|k|}\lambda_v(P),$$

by Proposition 5(e) in [2].

Applying (3.11) and (3.12) with Table 2 of [2] to Lemma 3.3, we complete the proof of Theorem 1.1. For  $I_{2m}^*$  and  $c_v = 4$ , we use  $v(\phi_2(P)) = v(\psi_3(P))$  from Lemma 2.10 to get an expression in terms of  $v(\phi_2(P))$  rather than  $v(\psi_3(P))$ .

#### ACKNOWLEDGEMENTS

The authors greatly appreciate the suggestion from Federico Pellarin that we investigate possible connections between the quantity,  $k_{v,n}$ , and local heights. This led us to the current formulation of our Theorem 1.1. We also thank the referee for their careful reading of our manuscript, helpful suggestions for improving it and interesting remarks that broadened our understanding.

#### REFERENCES

1. M. Ayad, *Points  $S$ -entiers des courbes elliptiques*, Manuscripta Math. 76 (1992), 305–324.
2. J. E. Cremona, M. Prickett, S. Siksek, *Height difference bounds for elliptic curves over number fields*, J. Number Theory **116** (2006) 42–68.
3. P. Ingram, *Elliptic divisibility sequences over certain curves*, J. Number Theory 123 (2007), 473–486.
4. P. Ingram, *Multiples of integral points on elliptic curves*, J. Number Theory 129 (2009), 182–208.

5. S. Lang, *Conjectured Diophantine estimates on elliptic curves*, In “Arithmetic and geometry, I”, Progr. Math. 35, 155–171, Birkhäuser, Boston, 1983.
6. I. Papadopoulos, *Sur la classification de Neron des courbes elliptiques en caractéristique résiduelle 2 et 3*, J. Number Theory 44 (1993), 119–152.
7. M. Schütt, T. Shioda, *Elliptic Surfaces*, In “Algebraic Geometry in East Asia — Seoul 2008”, Advanced Studies in Pure Mathematics 60 (2010), 51–160, World Scientific Publishing, Singapore.
8. J. H. Silverman, *The Arithmetic of Elliptic Curves, second edition*, Graduate Texts in Math., vol. 106, Springer-Verlag, New York, 2009.
9. J. H. Silverman, *Advanced Topics in Arithmetic of Elliptic Curves*, Graduate Texts in Math., vol. 151, Springer-Verlag, New York, 1994.
10. J. H. Silverman, *Computing heights on elliptic curves*, Math. Comp. 51 (1988), 339–358.
11. K. E. Stange, *Integral points on elliptic curves and explicit valuations of division polynomials*, Canad. J. Math. 68 (2016), 1120–1158.
12. J. Tate, *Algorithm for determining the type of a singular fiber in an elliptic pencil*, in “Modular Functions of One Variable IV”, Lecture Notes in Math. 476, Springer-Verlag, Berlin, 1975
13. P. Voutier and M. Yabuta, *Primitive divisors of certain elliptic divisibility sequences*, Acta Arith. 151 (2012) 165–190.

*Current address:* London, UK

*Email address:* paul.voutier@gmail.com

*Current address:* Senri High School, 17-1, 2 chome, Takanodai, Suita, Osaka, 565-0861, Japan

*Email address:* yabutam@senri.osaka-c.ed.jp, rinri216@msf.biglobe.ne.jp