

NAIVE \mathbb{A}^1 -HOMOTOPIES ON RULED SURFACES

CHETAN BALWE AND ANAND SAWANT

ABSTRACT. We explicitly describe the \mathbb{A}^1 -chain homotopy classes of morphisms from a smooth henselian local scheme into a smooth projective surface, which is birationally ruled over a curve of genus > 0 . We consequently determine the sheaf of naive \mathbb{A}^1 -connected components of such a surface and show that it does not agree with the sheaf of its genuine \mathbb{A}^1 -connected components when the surface is not a minimal model. However, the sections of the sheaves of both naive and genuine \mathbb{A}^1 -connected components over schemes of dimension ≤ 1 agree. As a consequence, we show that the Morel-Voevodsky singular construction on a smooth projective surface, which is birationally ruled over a curve of genus > 0 , is not \mathbb{A}^1 -local if the surface is not a minimal model.

1. INTRODUCTION

A basic question in \mathbb{A}^1 -homotopy theory is to determine the set of *genuine \mathbb{A}^1 -homotopy classes* of morphisms from one scheme to another and study how they compare with the corresponding *naive \mathbb{A}^1 -homotopy classes*. Since the \mathbb{A}^1 -homotopy category $\mathcal{H}(k)$ over a field k is built out of the category of simplicial sheaves on the big Nisnevich site of smooth schemes over k by taking a suitable localization, one expects naive and genuine \mathbb{A}^1 -homotopy classes of morphisms from a smooth henselian local scheme U into a smooth scheme X to be closely related to one another.

We will focus on smooth projective schemes X over a base field k . The genuine \mathbb{A}^1 -homotopy classes of morphisms from a smooth henselian local scheme U into X are just the sections $\pi_0^{\mathbb{A}^1}(X)(U) = \text{Hom}_{\mathcal{H}(k)}(U, X)$ of the sheaf of \mathbb{A}^1 -connected components of X on U . On the other hand, the naive \mathbb{A}^1 -homotopy classes $\mathcal{S}(X)(U)$ of morphisms from a smooth henselian scheme U into X are the quotient of the set of morphisms of schemes $U \rightarrow X$ by the equivalence relation generated by naive \mathbb{A}^1 -homotopies, that is, morphisms $U \times \mathbb{A}^1 \rightarrow X$. If the Morel-Voevodsky singular construction $\text{Sing}_*^{\mathbb{A}^1} X$ is \mathbb{A}^1 -local (see Section 2 for precise definitions), the sheaves $\mathcal{S}(X)$ and $\pi_0^{\mathbb{A}^1}(X)$ agree. However, $\text{Sing}_*^{\mathbb{A}^1} X$ is \mathbb{A}^1 -local in very special cases (see [1, Theorem 2.3.5, Remark 3.3.8] for a general result about \mathbb{A}^1 -locality of $\text{Sing}_*^{\mathbb{A}^1}$) and is not \mathbb{A}^1 -local in general (see [3], [4] for examples).

It is easy to see that $\mathcal{S}(X)(U)$ and $\pi_0^{\mathbb{A}^1}(X)(U)$ always agree when X has dimension ≤ 1 , and the same in case of a non-uniruled surface X can be obtained from the dimension ≤ 1 case (see [3, Theorem 3.14, Corollary 3.15]). A result of Asok and Morel ([2, Theorem 2.4.3]; see also [3, Theorem 3.9, Corollary 3.10]) states that for a proper scheme X over a field k ,

Chetan Balwe was supported by SERB-DST MATRICES Grant: MTR/2017/000690.

Anand Sawant acknowledges support of the Department of Atomic Energy, Government of India, under project no. 12-R&D-TFR-5.01-0500. He was supported by DFG Individual Research Grant SA3350/1-1 while he was affiliated to Ludwig-Maximilians-Universität, München, where a part of this work was carried out.

one has a bijection

$$(1.1) \quad \mathcal{S}(X)(\mathrm{Spec} F) \xrightarrow{\sim} \pi_0^{\mathbb{A}^1}(X)(\mathrm{Spec} F),$$

for every finitely generated, separable field extension F of k . However, the corresponding result for a general U cannot hold, as shown by the counterexample constructed in [3, Section 4] (for U the spectrum of a henselian discrete valuation ring and X a particular threefold).

When the base field k is algebraically closed of characteristic 0, the set $\pi_0^{\mathbb{A}^1}(X)(U)$ has been explicitly determined in [5], where X is a smooth projective surface, birationally ruled over a curve of genus > 0 . In this companion paper to [5], we determine the sheaf of naive \mathbb{A}^1 -connected components of a (birationally) ruled surface over an algebraically closed field of characteristic 0. The main result of the paper provides an extension of (1.1) to sections over schemes of dimension ≤ 1 , when X is a smooth projective surface.

Theorem 1. *Let k be an algebraically closed field of characteristic 0 and let X be a smooth projective surface over k , which is birationally ruled over a curve of genus > 0 . Then the natural map*

$$\mathcal{S}(X)(U) \rightarrow \pi_0^{\mathbb{A}^1}(X)(U)$$

is a bijection, for every henselian local scheme U over k of dimension ≤ 1 .

Theorem 1 is known to hold in the case of non-uniruled surfaces by [3, Corollary 3.15] and the case of rational surfaces is easy to handle since they are covered by affine spaces [2, Lemma 2.2.11]. Thus, Theorem 1 in fact holds for all smooth projective surfaces over k . It has been proved in [5] that the sheaf $\pi_0^{\mathbb{A}^1}(X)$ for such an X agrees with the second iteration of the functor \mathcal{S} on X . Theorem 1 will be proved by directly obtaining a precise description of $\mathcal{S}(X)(U)$ and comparing it with the description of $\mathcal{S}^2(X)(U)$ obtained in [5].

We emphasize here that our method of proof of Theorem 1 gives an explicit description of $\mathcal{S}(X)(U)$ for any smooth henselian local k -scheme U in terms of an algebraic condition, which can be easily verified. The key result pertaining to this is Proposition 4.10, which gives this description in the case of a single point blowup. The general case reduces to this case, as outlined in Section 5 and is of independent interest.

As an application of the method of our proof, we get the non- \mathbb{A}^1 -locality of the Morel-Voevodsky singular construction applied to any smooth projective surface birationally ruled over a curve of genus > 0 , which is not a minimal model.

Theorem 2. *Let k be an algebraically closed field of characteristic 0 and let X be a smooth projective surface over k , which is birationally ruled over a curve of genus > 0 . If X is not a minimal model, then the natural epimorphism*

$$\mathcal{S}(X) \rightarrow \mathcal{S}^2(X) = \pi_0^{\mathbb{A}^1}(X)$$

is not an isomorphism of Nisnevich sheaves. In particular, $\mathrm{Sing}_^{\mathbb{A}^1} X$ is not \mathbb{A}^1 -local.*

The methodology used in the proofs of Theorems 1 and 2 is to classify the naive \mathbb{A}^1 -homotopy classes of morphisms from a smooth henselian local k -scheme U into X birationally ruled over a curve C of genus > 0 by fixing a morphism $\gamma: U \rightarrow C$ and classifying homotopy classes of morphisms $U \rightarrow X$ over γ by tracking where they map the closed point of U into X . Preliminaries about \mathbb{A}^1 -connectedness are given in Section 2, whereas results about local geometry of ruled surfaces are recalled in Section 3. We first handle the case of the blowup of one point on the projectivization of a rank 2 vector bundle on C in Section 4. In

Section 5, the main results are obtained for a special class of blowups called *nodal blowups* by using an appropriate modification of the arguments given in Section 4. One then reduces the general case to these blowups by employing the same method used in [5], which analyzes local structure of naive \mathbb{A}^1 -homotopy classes of blowups of such surfaces at infinitely near points. We point out that although the strategy of the proofs of the main results is similar to the one in [5], the key geometric input in this paper (Proposition 4.10 and its proof) is very different in spirit and of independent interest to commutative algebraists and algebraic geometers.

2. NAIVE AND GENUINE \mathbb{A}^1 -CONNECTED COMPONENTS

We fix a base field k and denote by Sm/k the category of smooth, finite-type, separated schemes over $\mathrm{Spec} k$. We will denote by $\mathcal{H}(k)$ the unstable \mathbb{A}^1 -homotopy category constructed by Morel and Voevodsky [6]. One obtains $\mathcal{H}(k)$ from the category of simplicial Nisnevich sheaves of sets on Sm/k (also called *spaces* over k for brevity) by localizing with respect to Nisnevich local weak equivalences and further localizing with respect to the class of projection maps $\mathcal{X} \times \mathbb{A}^1 \rightarrow \mathcal{X}$, for every simplicial Nisnevich sheaf of sets \mathcal{X} . All presheaves on Sm/k will be extended to essentially smooth schemes (that is, schemes which are filtered inverse limits of diagrams of smooth schemes in which the transition maps are étale, affine morphisms) by defining $\mathcal{F}(\varinjlim U_\alpha) = \varinjlim \mathcal{F}(U_\alpha)$. By a *smooth henselian local scheme* over k , we will mean the henselization of the local ring at a smooth point of a scheme over k .

We will follow the conventions and notation of [3] and [5] regarding \mathbb{A}^1 -connected components. We briefly recall the notions relevant to this paper below.

Definition 2.1. Let \mathcal{X} be a space over k . Define $\mathrm{Sing}_*^{\mathbb{A}^1} \mathcal{X}$ to be the simplicial sheaf given by

$$(\mathrm{Sing}_*^{\mathbb{A}^1} \mathcal{X})_n = \underline{\mathrm{Hom}}(\Delta^n, \mathcal{X}_n),$$

where Δ^\bullet denotes the cosimplicial sheaf

$$\Delta^n = \mathrm{Spec} \frac{k[x_0, \dots, x_n]}{\langle \sum_i x_i = 1 \rangle}$$

with natural face and degeneracy maps analogous to the ones on topological simplices.

Given a simplicial sheaf of sets \mathcal{X} on Sm/k , we will denote by $\pi_0(\mathcal{X})$ the presheaf on Sm/k that associates with $U \in Sm/k$ the coequalizer of the diagram $\mathcal{X}_1(U) \rightrightarrows \mathcal{X}_0(U)$, where the maps are the face maps coming from the simplicial data of \mathcal{X} . We will denote the Nisnevich sheafification functor by a_{Nis} .

Definition 2.2. The sheaf of \mathbb{A}^1 -chain connected components of a space \mathcal{X} is defined to be

$$\mathcal{S}(\mathcal{X}) := a_{\mathrm{Nis}} \left(\pi_0 \mathrm{Sing}_*^{\mathbb{A}^1} \mathcal{X} \right).$$

Thus, when \mathcal{X} is the space given by a scheme X over k , for every smooth henselian local k -scheme U , the set $\mathcal{S}(X)(U)$ is the quotient of $X(U)$ by the equivalence relation generated by $\sigma_0^*(h) = \sigma_1^*(h)$, where $h \in X(U \times \mathbb{A}^1)$ is a naive \mathbb{A}^1 -homotopy of U into X and $\sigma_i: U \rightarrow U \times \mathbb{A}^1$ denotes the i -section.

Definition 2.3. Let $X \in Sm/k$ and let U be an essentially smooth scheme over k .

- (1) An \mathbb{A}^1 -homotopy of U in X is a morphism $h: U \times \mathbb{A}_k^1 \rightarrow X$. We say that $f, g \in X(U)$ are \mathbb{A}^1 -homotopic if there exists an \mathbb{A}^1 -homotopy $h: U \times \mathbb{A}^1 \rightarrow X$ such that $\sigma_0^*(h) = f$ and $\sigma_1^*(h) = g$.

- (2) An \mathbb{A}^1 -chain homotopy of U in X is a finite sequence $\mathcal{H} = (h_1, \dots, h_r)$ where each h_i is an \mathbb{A}^1 -homotopy of U in X such that $\sigma_1^*(h_i) = \sigma_0^*(h_{i+1})$ for $1 \leq i \leq r-1$. We say that $f, g \in X(U)$ are \mathbb{A}^1 -chain homotopic if there exists an \mathbb{A}^1 -chain homotopy $h = (h_1, \dots, h_r)$ such that $\sigma_0^*(h_1) = f$ and $\sigma_1^*(h_r) = g$.
- (3) The total space $\mathrm{Sp}(h)$ of an \mathbb{A}^1 -homotopy $h: U \times \mathbb{A}_k^1 \rightarrow X$ of U in X is just $U \times \mathbb{A}_k^1$. The total space $\mathrm{Sp}(\mathcal{H})$ of an \mathbb{A}^1 -chain homotopy $\mathcal{H} = (h_1, \dots, h_r)$ is the disjoint union of the total spaces of h_1, \dots, h_r . We write $f_{\mathcal{H}}: \mathrm{Sp}(\mathcal{H}) \rightarrow U$ for the canonical projection on U . The morphism $h_{\mathcal{H}}: \mathrm{Sp}(\mathcal{H}) \rightarrow X$ is just the morphism defining the chain homotopy.

Definition 2.4. The sheaf of \mathbb{A}^1 -connected components of a space \mathcal{X} is defined to be

$$\pi_0^{\mathbb{A}^1}(\mathcal{X}) = a_{\mathrm{Nis}} \mathrm{Hom}_{\mathcal{H}(k)}(-, \mathcal{X}).$$

Equivalently, $\pi_0^{\mathbb{A}^1}(\mathcal{X}) = a_{\mathrm{Nis}}(\pi_0 L_{\mathbb{A}^1} \mathcal{X})$, where $L_{\mathbb{A}^1}$ denotes an \mathbb{A}^1 -fibrant replacement functor for the injective Nisnevich local model structure.

Given schemes U and X over k , two elements of $X(U)$ are said to be *naively* \mathbb{A}^1 -homotopic if they are \mathbb{A}^1 -chain homotopic. Two elements of $X(U)$ are said to be *genuinely* \mathbb{A}^1 -homotopic if they map to the same element in $\pi_0^{\mathbb{A}^1}(X)(U)$.

Remark 2.5. If $\mathrm{Sing}_*^{\mathbb{A}^1} \mathcal{X}$ is \mathbb{A}^1 -local in the sense of [2, Definition A.2.1], then the natural epimorphism $\mathcal{S}(\mathcal{X}) \rightarrow \pi_0^{\mathbb{A}^1}(\mathcal{X})$ is an isomorphism (see [2, Remark 2.2.9]).

3. PRELIMINARIES ON RULED SURFACES

Theorem 1 and Theorem 2 will be proved by using the classification of ruled surfaces and an induction on the *complexity* of the blowup involved. In this section, we will briefly outline the formalism developed in [5] to handle blowups of ruled surfaces and state some results about local geometry of ruled surfaces proved in [5, Section 4]. We follow the conventions of [5, Section 4], where we also refer the reader to for more details. We will focus on smooth projective surfaces, which are obtained from $\mathbb{P}^1 \times C$ by a finite number (possibly zero) of successive blowups at smooth, closed points, where C is the henselization of the local ring of a closed point on a smooth projective curve of genus ≥ 1 . The results for general ruled surfaces will follow from the ones in this special case by standard arguments (see the proofs of the main results in Section 5).

Throughout the rest of the article, we will assume that k is an algebraically closed field of characteristic 0, although this assumption is not always necessary for some of the results proved in the rest of the article. We fix the following setting for this section.

Notation 3.1.

- (1) Let x be a variable and let A be the henselization of the polynomial ring $k[x]$ at the maximal ideal $\langle x \rangle$. Let $C = \mathrm{Spec} A$ and let c_0 denote the closed point of C . Every 1-dimensional regular henselian local ring containing k is isomorphic to A .
- (2) Let Y and Z be variables which denote the homogeneous coordinates on \mathbb{P}_C^1 . Thus, $\mathbb{P}_C^1 = \mathrm{Proj} A[Y, Z]$.
- (3) Let y denote the rational function Y/Z on \mathbb{P}_C^1 which is defined on the open subscheme of \mathbb{P}_C^1 defined by the condition $Z \neq 0$. This open subscheme is simply $\mathrm{Spec} A[y] \simeq \mathbb{A}_C^1$ and the point $(c_0, [0 : 1])$ is defined by the ideal $\langle x, y \rangle$ of $A[y]$.

- (4) Let ℓ_∞ denote the closed subscheme $C \times \{[0 : 1]\}$ of \mathbb{P}_C^1 , which is the divisor of zeros of y . The closed subscheme $C \times \{[1 : 0]\}$ is the divisor of poles of y and will be denoted by $\ell_{-\infty}$.

Definition 3.2. Let X be any scheme that is obtained from \mathbb{P}_C^1 by a finite number (possibly zero) of successive blowups at smooth, closed points.

- (1) The preimage of c_0 under the morphism $X \rightarrow C$ is a connected scheme, the irreducible components of which are isomorphic to \mathbb{P}_k^1 . We refer to these as *lines* on X .
- (2) We define a *pseudo-line* to be any curve that is either a line on X or the proper transform of ℓ_∞ .
- (3) A *node* on X is defined to be the intersection point of two pseudo-lines. It is easy to see that any node is the point of intersection of exactly two pseudo-lines. Thus, for instance, the point $(c_0, [0 : 1])$ is the only node on the scheme \mathbb{P}_C^1 .
- (4) We will denote by \mathcal{N} the collection of schemes X admitting a morphism $X \rightarrow \mathbb{P}_C^1$, which factors as

$$X = X_r \xrightarrow{\pi_r} X_{r-1} \xrightarrow{\pi_{r-1}} \cdots \xrightarrow{\pi_1} X_0 = \mathbb{P}_C^1$$

with $r \geq 0$, where for all $i \geq 1$, $\pi_i : X_i \rightarrow X_{i-1}$ is the blowup of X_{i-1} at some *node* of X_{i-1} . Elements of \mathcal{N} will be referred to as *nodal blowups*.

Since all the nodes in all the $X \in \mathcal{N}$ lie over the point $(c_0, [0 : 1])$, we see that the ideals defining these nodes are generated by rational functions in x and y . We record the following lemma from [5, Lemma 4.12].

Lemma 3.3. *Let $X \in \mathcal{N}$. Then:*

- (a) *If a line l on X has a parameter of the form x^a/y^b , then a and b are coprime non-negative integers.*
- (b) *If C_1 and C_2 are pseudo-lines on X with parameters x^a/y^b and x^c/y^d which intersect at a node defined by the ideal $\langle x^a/y^b, y^d/x^c \rangle$, then $ad - bc = 1$ (and so, in particular, $a/b > c/d$).*

This shows that for pair of coprime integers (a, b) with $a < b$, there is at most one line on X with parameter x^a/y^b . It can be shown that if such lines exist on two schemes X_1, X_2 in \mathcal{N} , then the canonical birational maps from X_1 to X_2 will take the lines into each other. If $X \in \mathcal{N}$ has n lines, we can associate to it the sequence of rational numbers $r_0 = -\infty < r_1 = 0 < \cdots < r_n = 1 < r_{n+1} = \infty$ such that:

- For $1 \leq i \leq n$, if r_i is written in reduced fractional form as a_i/b_i for some non-negative integers a_i, b_i , then X contains a line parameterized by x^{a_i}/y^{b_i} . We will label this line as ℓ_{r_i} . For $i = 0$ and $i = n + 1$, the symbol ℓ_{r_i} will denote the pseudo-lines $\ell_{-\infty}$ and ℓ_∞ respectively.
- For $1 \leq i \leq n$, the line ℓ_{r_i} only meets the pseudo-lines $\ell_{r_{i-1}}$ and $\ell_{r_{i+1}}$. If r_i and r_{i+1} are written in reduced fractional form as $r_i = a_i/b_i$ and $r_{i+1} = a_{i+1}/b_{i+1}$ then they meet in a node defined by the maximal ideal $\langle x^{a_{i+1}}/y^{b_{i+1}}, y^{b_i}/x^{a_i} \rangle$. (Here, if $r_{i+1} = \infty$, we choose $a_{i+1} = 1$ and $b_{i+1} = 0$.)

4. THE CASE OF THE BLOW-UP OF A MINIMAL MODEL AT A SINGLE CLOSED POINT

The aim of this section is to prove the following result, which proves both Theorem 1 and 2 stated in the introduction in a special case of the blowup of a single closed point on $\mathbb{P}^1 \times C$, where C is a curve of genus > 0 .

Theorem 4.1. *Let E be a \mathbb{P}^1 -bundle over a smooth projective curve C of genus > 0 . Let X be the blowup of E at a single closed point. Then*

- (a) $\mathcal{S}(X)(U) = \mathcal{S}^2(X)(U)$, where $U = \text{Spec } R$, for a henselian discrete valuation ring R over k ;
- (b) $\mathcal{S}(X) \neq \mathcal{S}^2(X)$.

Along the course of proving Theorem 4.1, we will give a complete description of \mathbb{A}^1 -chain homotopy classes of a smooth henselian local scheme U into X .

Fix a smooth henselian local ring (R, \mathfrak{m}) , $U = \text{Spec } R$. We fix a morphism $\gamma: U \rightarrow C$ and consider pullbacks of X and E with respect to γ which we denote by X_γ and E_γ . Thus X_γ is obtained by blowing up a closed subscheme of E_γ . But since a \mathbb{P}^1 -bundle is étale locally trivial, we see that $E_\gamma \simeq \mathbb{P}_U^1$. Thus we may assume that $E \rightarrow C$ is the trivial \mathbb{P}^1 -bundle \mathbb{P}_C^1 .

Thus, let us assume that $E = \mathbb{P}_C^1 = \mathbb{P}_k^1 \times C$ and X has been obtained from \mathbb{P}_C^1 by blowing up the point $p := (c_0, (0 : 1))$ where c_0 is a closed point in C . Suppose the image of $\gamma: U \rightarrow C$ does not contain c_0 . Then X_γ is isomorphic to \mathbb{P}_U^1 . On the other hand, if γ maps the whole of U into the point c_0 , then $X_\gamma = U \times T$ where T consists of two copies of \mathbb{P}_k^1 intersecting transversely in a single point. In both these cases, any two sections of $X_\gamma \rightarrow U$ are \mathbb{A}^1 -chain homotopic and thus the required result follows immediately. Thus we may now focus on the case in which γ maps the closed point of U to c_0 but does not map the whole of U into c_0 .

Let ω be a uniformizing parameter in the ring \mathcal{O}_{C, c_0} . Then X_γ is obtained from $\mathbb{P}_U^1 = \text{Proj } R[Y, Z]$ by blowing up the closed subscheme corresponding to the homogeneous ideal $\langle \gamma^*(\omega), Y \rangle$. The remainder of this section will be devoted to analyzing the \mathbb{A}^1 chain homotopy classes in $X_\gamma(U)$, where we view X_γ as a U -scheme (thus, the projections of all the \mathbb{A}^1 -chain homotopies to U must be constant). We now set up some notation to analyze the scheme X_γ .

Notation 4.2. The following notation will stay in effect for the remainder of this section:

- (a) Let R denote the henselization of a ring of the form $\mathcal{O}_{V, v}$ where V is a smooth scheme over k and v is a point of V . Let \mathfrak{m} be the maximal ideal of R and let $\kappa = R/\mathfrak{m}$. We will denote $\text{Spec } R$ by U and the closed point of U by u . For $r \in R$, $U(r)$ will denote the closed subscheme of r corresponding to the ideal $\langle r \rangle$.
- (b) Y, Z will denote the homogeneous coordinates on $\mathbb{P}_R^1 = \text{Proj } R[Y, Z]$.
- (c) For any $r_0 \in \mathfrak{m} \setminus \{0\}$, we define

$$I(r_0) := \text{ideal sheaf on } \mathbb{P}_U^1 \text{ corresponding to the} \\ \text{homogeneous ideal } \langle r_0, Y \rangle.$$

Let $T(r_0)$ denote the closed subscheme of \mathbb{P}_U^1 corresponding to this ideal sheaf. Let $\theta_{r_0}: X(r_0) \rightarrow \mathbb{P}_U^1$ denote the blowup of \mathbb{P}_U^1 at $T(r_0)$.

We wish to examine the sections of morphisms $X(r_0) \rightarrow U$. The scheme \mathbb{P}_U^1 is covered by affine patches isomorphic to \mathbb{A}_U^1 given by the conditions $Y \neq 0$ and $Z \neq 0$. We will denote these by $\text{Spec } R[Y/Z]$ and $\text{Spec } R[Z/Y]$ respectively. Using the fact that U is a local scheme it is easy to see that any section of \mathbb{P}_U^1 factors through $\text{Spec } R[Y/Z]$ or $\text{Spec } R[Z/Y]$. The

sections of $\text{Spec } R[Y/Z] \rightarrow U$ are given by R -algebra homomorphisms $R[Y/Z] \rightarrow R$. Such a homomorphism is determined by a choice for the image of Y/Z .

Notation 4.3. We denote by $\beta_r: U \rightarrow \text{Spec } R[Y/Z] \rightarrow \mathbb{P}_U^1$ the section of $\mathbb{P}_U^1 \rightarrow U$ determined by $Y/Z \mapsto r$. Similarly, we denote by $\alpha_r: U \rightarrow \text{Spec } R[Z/Y] \rightarrow \mathbb{P}_U^1$ the section of $\mathbb{P}_U^1 \rightarrow U$ determined by $Z/Y \mapsto r$. Thus we see that if r is a unit in R , then $\alpha_r = \beta_{1/r}$.

A section of $X(r_0) \rightarrow U$ can be composed with $X(r_0) \rightarrow \mathbb{P}_U^1$ to give a section of $\mathbb{P}_U^1 \rightarrow U$. Of course, not every section of $\mathbb{P}_U^1 \rightarrow U$ can be lifted to give a section of $X(r_0) \rightarrow U$. Since $r_0 \neq 0$, any section of $\mathbb{P}_U^1 \rightarrow U$ maps the generic point of U outside the scheme $T(r_0)$. As $X(r_0) \rightarrow \mathbb{P}_U^1$ is an isomorphism outside $T(r_0)$, it follows that if a lift of a section of $\mathbb{P}_U^1 \rightarrow U$ to $X(r_0)$ exists, then it must be unique.

Notation 4.4. Let $r_0 \in \mathfrak{m} \setminus \{0\}$. For $r \in R$, if a lift of $\alpha_r: U \rightarrow \mathbb{P}_U^1$ to $X(r_0)$ exists, it will be denoted by $\alpha_r^{r_0}$. Similarly, we define $\beta_r^{r_0}$ to be the lift of β_r to $X(r_0)$, if it exists.

A section $\alpha: U \rightarrow \mathbb{P}_U^1$ of $\mathbb{P}_U^1 \rightarrow U$, lifts to $X(r_0)$ if and only if the ideal sheaf $\alpha^*(I(r_0))$ is principal. Since U is local, this ideal sheaf corresponds to an ideal of R . Unless α is of the form β_r where $r \in \mathfrak{m}$, we have $\alpha^*(I(r_0)) = \mathcal{O}_U$. When $\alpha = \beta_r$ for $r \in \mathfrak{m}$, the ideal sheaf $\alpha^*(I(r_0))$ corresponds to the ideal $\langle r_0, r \rangle$ in R .

Lemma 4.5. Let $r_0 \in \mathfrak{m} \setminus \{0\}$. Let $\alpha: U \rightarrow \mathbb{P}_U^1$ be a section of $\mathbb{P}_U^1 \rightarrow U$ which admits a lift to $X(r_0)$. Then one of the following must hold:

- (a) $\alpha^*(I(r_0)) = \langle 1 \rangle$. In this case, α is of the form α_r where $r \in R$.
- (b) $\alpha^*(I(r_0)) = \langle r_0 \rangle$. In this case, α is of the form β_r where $r_0 | r$.
- (c) $\alpha^*(I(r_0)) = \langle r \rangle$ for some $r \in R$ such that $r | r_0$ but $r_0 \nmid r$. In this case, α is of the form $\beta_{r'}$ where r'/r is a unit in R .

Proof. We have $\alpha^*(I(r_0)) = \langle 1 \rangle$ if and only if α factors through the complement of the scheme $T(r_0)$ of \mathbb{P}_U^1 . This is so if and only if α is of the form α_r where $r \in R$. This proves (a).

We now have to consider the case that α is of the form $\beta_r: U \rightarrow \mathbb{P}_U^1$ for some $r \in \mathfrak{m}$. In this case, $\alpha^*(I(r_0)) = \langle r_0, r \rangle$. Since R is a local domain, this ideal is equal to $\langle r_0 \rangle$ if $r_0 | r$ or is equal to $\langle r \rangle$ if $r | r_0$. This proves (b) and (c). \square

Notation 4.6. Let $r_0 \in \mathfrak{m} \setminus \{0\}$. Then for $r \in R$, $\mathcal{A}_r^{r_0}$ denotes the set of sections $\alpha: U \rightarrow \mathbb{P}_U^1$ such that the ideal sheaf $\alpha^*(I(r_0))$ corresponds to the ideal $\langle r \rangle$ in R . By Lemma 4.5, $\mathcal{A}_r^{r_0}$ is non-empty if and only if $r | r_0$.

Remark 4.7. We note a geometric interpretation of the families $\mathcal{A}_r^{r_0}$. Observe that the preimage of c_0 under the morphism $X \rightarrow C$ is the union of two lines L_1 and L_2 . Here L_1 is the proper transform of the line $\{c_0\} \times \mathbb{P}_k^1 \subset \mathbb{P}_C^1$ and L_2 is the exceptional divisor. A section $\alpha: U \rightarrow \mathbb{P}_U^1$ maps the closed point u of U into L_2 if and only if r is not a unit. It maps u into L_1 if and only if $r | r_0$ and $r_0 \nmid r$.

Thus, the elements of $\mathcal{A}_1^{r_0}$ correspond to those lifts of γ which map u into $L_1 \setminus L_2$. The elements of $\mathcal{A}_{r_0}^{r_0}$ are those sections which map u into $L_2 \setminus L_1$. The set of sections which map u to the point $L_1 \cap L_2$ is precisely the union of the sets $\mathcal{A}_r^{r_0}$ where r is not a unit, $r | r_0$ and $r_0 \nmid r$.

This indicates a certain symmetry between the sets $\mathcal{A}_1^{r_0}$ and $\mathcal{A}_{r_0}^{r_0}$. Indeed, these two sets can be transformed into each other by means of an *elementary transformation*.

We will use the following special case of [5, Proposition 3.7].

Lemma 4.8. *Let $r_0 \in \mathfrak{m} \setminus \{0\}$. Let α and α' be sections of $\mathbb{P}_U^1 \rightarrow U$ which are connected by an \mathbb{A}^1 -chain homotopy \mathcal{H} . Then there exists $r \in R$ such that $r|r_0$ and $\alpha, \alpha' \in \mathcal{A}_r^{r_0}$. Moreover, in this case $h_{\mathcal{H}}^*(I(r_0))$ is generated by $f_{\mathcal{H}}^*(r)$.*

Lemma 4.8 gives us a necessary condition for two elements β_{r_1} and β_{r_2} to be \mathbb{A}^1 -homotopic. However, we will see now that it is not strong enough. The following two results give a complete description of \mathbb{A}^1 -homotopy classes of the sections of $X(r_0) \rightarrow U$.

Lemma 4.9. *Let $r_0 \in \mathfrak{m} \setminus \{0\}$. Any two elements of $\mathcal{A}_r^{r_0}$ are \mathbb{A}^1 -homotopic via homotopies that lift to $X(r_0)$ if $r = 1$ or r_0 .*

Proof. First consider the case $r = 1$. Suppose α, α' are in $\mathcal{A}_1^{r_0}$. Then it is easy to see that $\alpha = \alpha_{r_1}$ and $\alpha' = \alpha_{r_2}$ for some units $r_1, r_2 \in R^\times$. Then these two elements are homotopic in \mathbb{P}_U^1 via the homotopy given by the R -algebra homomorphism

$$R[Z/Y] \rightarrow R[T], \quad Z/Y \mapsto r_1^{-1}(1 - T) + r_2^{-1}T.$$

Note that we are using the parameter Z/Y instead of Y/Z , so $\alpha_{r_i}^*(Z/Y) = r_i^{-1}$ for $i = 1, 2$. This defines a morphism $\mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1$, which factors through the open immersion $\mathbb{P}_U^1 \setminus T(r_0) \hookrightarrow \mathbb{P}_U^1$ and hence, lifts to $X(r_0)$.

The case $r = r_0$ can be deduced from the case $r = 1$ by means of an elementary transformation (as mentioned in Remark 4.7). However, we will simply write out an explicit homotopy. If α, α' are in $\mathcal{A}_{r_0}^{r_0}$, they are of the form β_{r_1} and β_{r_2} where $r_0|r_1$ and $r_0|r_2$. Then consider the homotopy $h: \mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1$ given by the R -algebra homomorphism $R[Y/Z] \rightarrow R[T]$, $Y/Z \mapsto r_1(1 - T) + r_2T$. Then $h^*(I(r_0))$ is the sheaf corresponding to the ideal $\langle r_0, r_1(1 - T) + r_2T \rangle = \langle r_0 \rangle$ which is principal. Thus h lifts to $X(r_0)$ as claimed. \square

Proposition 4.10. *Let $r_0 \in \mathfrak{m} \setminus \{0\}$. Suppose $r_1, r_2 \in R$ are non-units such that $r_i|r_0$ but $r_0 \nmid r_i$ for $i = 1, 2$. Let $r' = r_0/r_1$. Then $\beta_{r_i}^{r_0}$ for $i = 1, 2$ are \mathbb{A}^1 -chain homotopic in $X(r_0)$ if and only if the following two conditions hold:*

- (a) r_1/r_2 is a unit.
- (b) $r_1/r_2 - 1 \in \sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle} \subset \sqrt{\langle r_1, r' \rangle}$.

Proof. We first prove the necessity of the conditions. Clearly, it suffices to treat the case where $\beta_{r_1}^{r_0}$ and $\beta_{r_2}^{r_0}$ are \mathbb{A}^1 -homotopic in $X(r_0)$.

The fact that r_1/r_2 is a unit is already known to us by Lemma 4.8. An \mathbb{A}^1 -homotopy $h: \mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1$ is given by the choice of an invertible sheaf on \mathbb{A}_U^1 along with the choice of two generating sections. Since U is essentially smooth, we have $\text{Pic}(U) \simeq \text{Cl}(U)$ and $\text{Pic}(\mathbb{A}_U^1) \simeq \text{Cl}(\mathbb{A}_U^1)$, where Cl denotes the group of Weil divisors. The projection $\mathbb{A}_U^1 \rightarrow U$ induces an isomorphism $\text{Cl}(U) \simeq \text{Cl}(\mathbb{A}_U^1)$. Since U is local, this implies that any invertible sheaf on \mathbb{A}_U^1 is isomorphic to $\mathcal{O}_{\mathbb{A}_U^1}$. Thus we see that the homotopy h can be given by two generating global sections of $\mathcal{O}_{\mathbb{A}_U^1}$, that is, two polynomials $p(T), q(T) \in R[T]$ such that $\langle p(T), q(T) \rangle$ is the unit ideal. If $D(q)$ denotes the open subscheme of \mathbb{A}_U^1 where q is a unit, then h maps $D(q)$ into the open subscheme $\mathbb{A}_U^1 = \text{Spec } R[Y/Z] \subset \mathbb{P}_U^1$ and the morphism $h|_{D(q)}$ is given by the R -algebra homomorphism $Y/Z \mapsto p(T)/q(T)$. Notice that $p(0)/q(0) = r_1$ and $p(1)/q(1) = r_2$. By the first equality, we see that $r_1|p(0)$ and thus, in particular, $p(0)$ is a non-unit. However, $\langle p(0), q(0) \rangle$ is the unit ideal in R (since $\langle p, q \rangle$ is the unit ideal in $R[T]$).

Thus $q(0)$ is a unit in R . Thus $q(T)$ is a primitive polynomial, that is, its content $\text{cont}(q)$, is a unit.

Now suppose that the ideal sheaf $h^*(I(r_0))$ is locally principal. Since $R[T, q^{-1}]$ is a unique factorization domain, the ideal $h^*(I(r_0))(D(q))$ is principal. By the formula defining h , this ideal is equal to $\langle r_0, p/q \rangle$. Since h lifts to $X(r_0)$, we know that this ideal is actually equal to $r_1 R[T, q^{-1}]$. Therefore,

$$p \in r_1 R[T, q^{-1}] \cap R[T] = \bigcup_{n \geq 0} (r_1 : q^n).$$

We claim that $(r_1 : q^n) = \langle r_1 \rangle$ for every $n \geq 0$. Indeed, suppose $f \in (r_1 : q^n)$. If $n = 0$, there is nothing to prove. So, we assume $n > 0$. There exists an element $g(T) \in R[T]$ and an integer n such that $r_1 g = f q^n$ in $R[T]$. But then, $q^n | r_1 g$ in $R[T]$. As q is a primitive polynomial, $q^n | (r_1 g / \text{cont}(r_1 g)) = g / \text{cont}(g)$. In particular, $q^n | g$ and thus $r_1 | f$ in $R[T]$, as claimed. This shows that $r_1 | p$ in $R[T]$ and we write $p = r_1 p'$ where $p' \in R[T]$.

The ideal $\langle r_1 p', q \rangle$ is a unit ideal. Thus, modulo r_1 , we see that the polynomial $q(T)$ must be a unit. Hence, if $q(T) = q_0 + q_1 T + \dots$, we see that q_0 is a unit (as we have already seen before) and q_i is nilpotent modulo r_1 for $i \geq 1$.

In $R[T, q^{-1}]$, we have $\langle r_0, r_1 p' / q \rangle = \langle r_1 \rangle$, which implies that $\langle r', p' / q \rangle = \langle 1 \rangle$. Hence, there exists an integer $n \geq 0$ such that q^n lies in the ideal $\langle r', p' \rangle$ in $R[T]$. Since $\langle p, q \rangle = 1$, we have $\langle p^n, q^n \rangle = 1$, which implies that $\langle (p')^n, q^n \rangle = 1$. Thus, $\langle r', p' \rangle$, which contains $\langle (p')^n, q^n \rangle$ is equal to the unit ideal. Thus if $p'(T) = p'_0 + p'_1 T + \dots$, we see that p'_0 is a unit and p'_i is nilpotent modulo r' for $i \geq 1$.

Using $p(1)/q(1) = r_2$, we see that

$$r_2 = r_1 \left(\frac{u_1 + t_1}{u_2 + t_2} \right),$$

where u_i are units, t_1 is nilpotent modulo r' and t_2 is nilpotent modulo r_1 . Using $p(0)/q(0) = r_1$, we see that $u_1/u_2 = 1$. Thus, we may assume that $u_1 = u_2 = 1$. Let

$$t_3 = -\frac{t_2}{1 + t_2}.$$

Then we see that t_3 is nilpotent modulo r_1 and $r_2 = r_1(1 + t_1)(1 + t_3)$. Thus

$$\frac{r_2}{r_1} - 1 = t_1 + t_3 + t_1 t_3$$

and it is obvious that this element lies in $\sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle}$ as claimed. This proves the necessity of the conditions.

We now prove the sufficiency. Let s_1 and s' be the square-free parts of r_1 and $r' := r_0/r_1$. Then $\sqrt{\langle r_1 \rangle} = \langle s_1 \rangle$ and $\sqrt{\langle r' \rangle} = \langle s' \rangle$. By (b), we have $r_2/r_1 - 1 = s_1 \delta + s' \delta'$, for some $\delta, \delta' \in R$. We define $r_3 = r_1(1 + s_1 \delta_1)$.

Consider the rational function

$$f_1(T) = \frac{r_3}{1 + s_1 \delta_1 T}.$$

The ideal $\langle r_3, 1 + s_1 \delta_1 T \rangle$ is the unit ideal since $s_1 \delta_1$ is nilpotent modulo r_1 and consequently, nilpotent modulo r_3 . Thus f_1 defines a homotopy

$$h_1 : \mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1; Y/Z \mapsto \frac{r_3}{1 + s_1 \delta_1 T}.$$

Since $f_1(0) = r_3$ and $f_1(1) = r_1$, this homotopy connects β_{r_3} to β_{r_1} . Now, let p be a point of \mathbb{A}_U^1 and let $z \in U$ denote its image under the projection $\mathbb{A}_U^1 \rightarrow U$. We see that the ideal

$$h_1^*(I(r_0))_p = \left\langle r_0, \frac{r_3}{1 + s_1 \delta_1 T} \right\rangle_p \subset R[T]_p$$

is principal if $1 + s_1 \delta_1 T$ is a unit at p , since $r_3 | r_0$. If $1 + s_1 \delta_1 T$ is not a unit, then s_1, δ_1 and T are all units at p . Thus, r_1 is also a unit at p . Since r_3 is a unit multiple of r_1 , we see that r_3 is a unit at p and hence, $r_3/(1 + s_1 \delta_1 T) = \infty$ modulo z . Thus, $h(p) = (z, (1 : 0))$, which is not in the closed subscheme defined by $I(r_0)$. Therefore, $h_1^*(I(r_0))_p$ is the unit ideal in this case. Consequently, $h_1^*(I(r_0))$ is a locally principal ideal, which allows us to conclude that h_1 lifts to $X(r_0)$.

Consider

$$f_2(T) = r_3 + r_1 s' \delta' T = r_3 \left(1 + \frac{s' \delta' T}{1 + s_1 \delta_1} \right).$$

This defines a homotopy

$$h_2 : \mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1; Y/Z \mapsto r_3 \left(1 + \frac{s' \delta' T}{1 + s_1 \delta_1} \right).$$

This homotopy lifts to $X(r_0)$ as the ideal $\langle r_0, r_3(1 + s' \delta' T(1 + s_1 \delta_1)^{-1}) \rangle$ is principal (since $s' \delta'$ is nilpotent modulo $r_0/r_3 = r'(1 + s_1 \delta_1)$). We observe that $f_2(0) = r_3$ and $f_2(1) = r_2$ and thus, h_2 connects β_{r_3} to β_{r_2} . This completes the proof. \square

Proof of Theorem 4.1. Let ω be a uniformizing parameter in the ring \mathcal{O}_{C, c_0} . Then X_γ is obtained from $\mathbb{P}_U^1 = \text{Proj } R[Y, Z]$ by blowing up the closed subscheme corresponding to the homogeneous ideal $\langle \gamma^*(\omega), Y \rangle$. Write $r_0 := \gamma^*(\omega)$. Lemma 4.9 and Proposition 4.10 give a classification of the \mathbb{A}^1 -homotopy classes of sections of $X_\gamma \rightarrow U$.

By Lemma 4.9, we are reduced to considering \mathbb{A}^1 -chain homotopy classes of sections of the form $\beta_r^{r_0}$, where $r \in \mathfrak{m}$ and $r | r_0$ but $r_0 \nmid r$. Set $r' = r_0/r$. If $U = \text{Spec } R$ is one-dimensional, then R is a discrete valuation ring, and it can be immediately seen that

$$\sqrt{\langle r \rangle} + \sqrt{\langle r' \rangle} = \sqrt{\langle r, r' \rangle}.$$

By [5, Proof of Theorem 5.1, Claims 1 and 2], it follows that $\mathcal{S}(X)(U) = \mathcal{S}^2(X)(U)$, for every smooth henselian local scheme U of dimension 1 over k , as desired.

The fact that $\mathcal{S}(X) \neq \mathcal{S}^2(X)$ is shown by the following concrete example. Let $\tilde{R} = k[x, y]_{\langle x, y \rangle}^h$. Let $r_0 = x(y^2 + x)$, $r_1 = x$. Thus $r' = y^2 + x$. We have

$$\sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle} = \langle x, y^2 + x \rangle = \langle x, y^2 \rangle,$$

while $\sqrt{\langle r_1, r' \rangle} = \langle x, y \rangle$. We take $r_2 = x(1 + y) = x + xy$. Thus

$$r_2/r_1 - 1 = y \notin \sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle}.$$

By Proposition 4.10, $\beta_{r_1}^{r_0}$ and $\beta_{r_2}^{r_0}$ are not \mathbb{A}^1 -chain homotopic. But by [5, Proposition 5.3, Claim 1 and Claim 2], they map to the same element in $\mathcal{S}^n(X)(\text{Spec } \tilde{R})$, for any $n \geq 2$. Therefore, $\mathcal{S}(X)(\text{Spec } \tilde{R}) \neq \mathcal{S}^2(X)(\text{Spec } \tilde{R})$. This completes the proof of Theorem 4.1.

5. PROOF OF THE MAIN RESULTS

We begin by proving Theorem 1 and Theorem 2 for nodal blowups.

Theorem 5.1. *Let $X \in \mathcal{N}$ be a nodal blowup in the sense of Definition 3.2.*

(a) *The natural map*

$$\mathcal{S}(X)(U) \rightarrow \mathcal{S}^2(X)(U) = \pi_0^{\mathbb{A}^1}(X)(U)$$

is a bijection, for every henselian local scheme U over k of dimension ≤ 1 .

(b) $\mathcal{S}(X) \neq \mathcal{S}^2(X)$.

Proof. There is a sequence of morphisms

$$(5.1) \quad X = X_p \rightarrow \cdots X_1 \rightarrow X_0 = \mathbb{P}_C^1,$$

where for all $i \geq 1$, $X_i \rightarrow X_{i-1}$ is a blowup at some node. There exist ordered pairs of non-negative integers $(m_0, n_0), (m_1, n_1), \dots, (m_p, n_p), (m_{p+1}, n_{p+1})$ with the following properties:

- m_i and n_i are coprime or each i ;
- $(m_0, n_0) = (0, 1)$, $(m_1, n_1) = (1, 0)$;
- For any i , there exist integers j and k such that $1 \leq j < k < i$ such that $m_k n_j - m_j n_k = 1$ and $m_i = m_j + m_k$, $n_i = n_j + n_k$;
- $X_i = X_{i-1}[x^{m_i}/y^{n_i}]$ (in the notation of [5, Proposition 4.2]). In other words, X_i is obtained from X_{i-1} by resolving indeterminacies of the rational function x^{m_i}/y^{n_i} .

Note that $n_i \neq 0$ for any $i \neq 1$. We relabel these ordered pairs as $(a_0, b_0), (a_1, b_1), \dots, (a_{p+1}, b_{p+1})$, in such a way that $a_0/b_0 = 0 < a_1/b_1 < \dots < a_p/b_p$ and $(a_{p+1}, b_{p+1}) = (1, 0)$. Set $s_i = a_i/b_i$ for $0 \leq i \leq p$ and $s_{p+1} = \infty$. Note that $s_p = a_p/b_p$ is always an integer. There are exactly $p + 2$ pseudo-lines on X , namely $\ell_0 = \ell_{s_0}, \ell_{s_1}, \dots, \ell_{s_p}$ and $\ell_{s_{p+1}} = \ell_\infty$. For $i \neq j$, the pseudo-lines ℓ_{s_i} and ℓ_{s_j} meet if and only if $j = i + 1$. The node in which ℓ_{s_i} and $\ell_{s_{i+1}}$ meet is locally given by the ideal $\langle x^{a_{i+1}}/y^{b_{i+1}}, y^{b_i}/x^{a_i} \rangle$. From this description, it is clear that X can be obtained from \mathbb{P}_C^1 by blowing up the ideal $\mathcal{I} = \prod_{i=1}^p \langle x^{a_i}, y^{b_i} \rangle$. We accordingly relabel the indices in (5.1) in such a way that for every i , X_{i+1} is obtained from X_i by blowing up the ideal $\langle x^{a_i}, y^{b_i} \rangle$.

We may assume by using elementary transformations that every pair (a_i, b_i) satisfies $a_i/b_i \leq 1$ and that $(a_p, b_p) = (1, 1)$. This follows by the analogue of [5, Proposition 5.3]; we thus omit the details.

If $\gamma: U \rightarrow C$ maps the closed point of U to the generic point of C , then α factors through the open subscheme $\mathbb{P}_{C \setminus \{c_0\}}^1$ of X . Hence any two γ -morphisms $\alpha_1, \alpha_2: U \rightarrow X$ are \mathbb{A}^1 -chain homotopic. On the other hand, if γ maps the generic point of U to the closed point of C , then any γ -morphism $\alpha: U \rightarrow X$ factors through the closed fibre of the morphism $X \rightarrow C$. The closed fibre of $X \rightarrow C$ is a connected scheme, the components of which are copies of \mathbb{P}_k^1 . Thus, again, in this case too, any two morphisms $\alpha_1, \alpha_2: U \rightarrow C$ which lift γ are \mathbb{A}^1 -chain homotopic. Hence, from this point onward, we will restrict our attention to the case in which γ maps the closed point of U to the closed point of C and the generic point of U to the generic point of C .

Let X_γ denote the pullback $X \times_{C, \gamma} U$. Then X_γ is the blowup of \mathbb{P}_U^1 at the pullback I_γ of the ideal \mathcal{I} with respect to the morphism $\mathbb{P}_U^1 \rightarrow \mathbb{P}_C^1$. We continue to use the variables Y and Z as homogeneous coordinates on $\mathbb{P}_U^1 = U \times_{C, \gamma} \mathbb{P}_C^1$ and we will also denote the pullback of the rational function $y = Y/Z$ on \mathbb{P}_C^1 by y . Set $r_0 = \gamma^*(x)$. Then $\mathcal{I}_\gamma = \prod_{i=1}^p I_i$

where $I_i = \langle r_0^{a_i}, y^{b_i} \rangle$. Applying [5, Proposition 3.7] to the blowup $X_\gamma \rightarrow \mathbb{P}_U^1$, we see that if $\alpha_1, \alpha_2: U \rightarrow \mathbb{P}_C^1$ are γ -morphisms which are connected by an \mathbb{A}^1 -chain homotopy \mathcal{H} which lifts to X , then there exists $r \in R$ such that the ideals $\alpha_1^*(\mathcal{I})$, $\alpha_2^*(\mathcal{I})$ and $h_{\mathcal{H}}^*(\mathcal{I})$ are all generated by r . The support of the ideal \mathcal{I} is the point $(c_0, [0 : 1])$. Hence, α_1 maps the closed point of U to $(c_0, [0 : 1])$ if and only if α_2 does the same. Any two morphisms $\alpha_1, \alpha_2: U \rightarrow \mathbb{P}_C^1$ which map the closed point of U to $\ell_0 \setminus \{(c_0, [0 : 1])\}$ are clearly \mathbb{A}^1 -chain homotopic via a homotopy which factors through the open subscheme $\mathbb{P}_C^1 \setminus \{(c_0, [0 : 1])\}$. Such a homotopy clearly lifts to X . Hence, we see that any two γ -morphisms of this type are \mathbb{A}^1 -chain homotopic. Hence, we will now focus on γ -morphisms $U \rightarrow \mathbb{P}_C^1$ which map the closed point of U to $(c_0, [0 : 1])$.

For any $r \in R$, let $\beta_r: U \rightarrow \mathbb{P}_U^1$ be the morphism induced by the R -algebra homomorphism $R[y] \rightarrow R$, $y \mapsto r$. Note that any section $\beta: U \rightarrow \mathbb{P}_U^1$ such that $\beta(u) = (u, [0 : 1])$ is of the form β_r for some unique $r \in \mathfrak{m}$. Such a section lifts to X_γ if and only if the ideal $\langle r_0^{a_i}, r^{b_i} \rangle$ is principal for every i , $1 \leq i \leq p$.

By [5, Proposition 3.7], we see that the the ideals (or ideal sheaves) $\beta_{r_1}^*(I_1)$, $\beta_{r_2}^*(I_1)$, $h_{\mathcal{H}}^*(I_1)$ are all generated by a fixed element of R , which we denote by r . we now have two cases:

Case A: $r_0 | r$. This happens when the ideals $\alpha_1^*(I_1)$, $\alpha_2^*(I_1)$ are generated by r_0 . Thus $r_0 | r_1$ and $r_0 | r_2$.

Case B: $r | r_0$ but $r_0 \nmid r$. In this case, for $i = 1, 2$, the ideal $\alpha_i^*(I_1) = \langle r_0, r_i \rangle$ is generated by r_i . As $\langle r_0, r_1 \rangle = \langle r_0, r_2 \rangle$, we see that r_1 and r_2 must be unit multiples of r .

At least one of the cases A or B must hold for the lifts of both β_{r_1} and β_{r_2} to X_γ to be \mathbb{A}^1 -chain homotopic.

If we are in Case A, consider the homotopy

$$h: \text{Spec } R[T] = \mathbb{A}_U^1 \rightarrow \mathbb{A}_U^1 = \text{Spec } R[y]$$

defined by $y \mapsto r_1 T + r_2(1 - T)$. Then as $a_i \leq b_i$ for all i , and as r_0 divides r_1 and r_2 , we get

$$h^*(\langle r_0^{a_i}, y^{b_i} \rangle) = \langle r_0^{a_i}, (r_1 T + r_2(1 - T))^{b_i} \rangle = \langle r_0^{a_i} \rangle$$

which is a principal ideal. Thus, the homotopy h lifts to X_γ and the lifts of β_{r_1} and β_{r_2} are \mathbb{A}^1 -chain homotopic.

Now, we consider Case B. Thus, r is an element of R such that $r | r_0$, $r_0 \nmid r$ and r_1 and r_2 unit elements of \mathfrak{m} such that r_i/r is a unit for $i = 1, 2$. Let $r_1, r_2 \in \mathfrak{m}$ be such that the sections β_{r_1} and β_{r_2} lift to X_γ and are connected by an \mathbb{A}^1 -chain homotopy \mathcal{H} which also lifts to X_γ . As $(a_p, b_p) = (1, 1)$, the sections β_{r_1} , β_{r_2} and the \mathbb{A}^1 -chain homotopy \mathcal{H} lift to $X_1 \times_{U, \gamma} \mathbb{P}_U^1$ which is obtained from \mathbb{P}_U^1 by blowing up the ideal $I_1 = \langle r_0, y \rangle$. By Proposition 4.10, r_1/r_2 must be a unit and we must have

$$(5.2) \quad r_1/r_2 - 1 \in \sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle} \subset \sqrt{\langle r_1, r' \rangle},$$

where $r' = r_0/r_1$. Now suppose that the condition (5.2) holds. We will show that the lifts of β_{r_1} and β_{r_2} are \mathbb{A}^1 -chain homotopic by an argument analogous to the one used in the proof of Proposition 4.10. Let s_1 and s' be the square-free parts of r_1 and $r' := r_0/r_1$ and write $r_2/r_1 - 1 = s_1 \delta_1 + s' \delta'$, for some $\delta_1, \delta' \in R$. We define $r_3 = r_1(1 + s_1 \delta_1)$. Consider the rational function

$$f_1(T) = \frac{r_3}{1 + s_1 \delta_1 T}.$$

The ideal $\langle r_3, 1 + s_1\delta_1 T \rangle$ is the unit ideal since $s_1\delta_1$ is nilpotent modulo r_1 and consequently, nilpotent modulo r_3 . Thus f_1 defines a homotopy

$$h_1: \mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1; Y/Z \mapsto \frac{r_3}{1 + s_1\delta_1 T}.$$

Since $f_1(0) = r_3$ and $f_1(1) = r_1$, this homotopy connects β_{r_3} to β_{r_1} . Now, let p be a point of \mathbb{A}_U^1 which projects to $z \in U$ under the projection $\mathbb{A}_U^1 \rightarrow U$. We see that for every i ,

$$h_1^*(I_i)_p = \langle r_0^{a_i}, \left(\frac{r_3}{1 + s_1\delta_1 T} \right)^{b_i} \rangle$$

is principal if $1 + s_1\delta_1 T$ is a unit at p . If $1 + s_1\delta_1 T$ is not a unit, then r_1 is also a unit at p and it follows that $h_1^*(I_i)_p$ is the unit ideal since $h_1(p) \notin \text{Supp}(I_i)$. Consequently, $h_1^*(\mathcal{I}_\gamma)$ is a locally principal ideal, which allows us to conclude that h_1 lifts to X_γ .

Consider

$$f_2(T) = r_3 + r_1 s' \delta' T = r_3 \left(1 + \frac{s' \delta' T}{1 + s_1 \delta_1} \right).$$

This defines a homotopy

$$h_2: \mathbb{A}_U^1 \rightarrow \mathbb{P}_U^1; Y/Z \mapsto r_3 \left(1 + \frac{s' \delta' T}{1 + s_1 \delta_1} \right).$$

If $r_0^{a_i} | r_3^{b_i}$, then there is nothing to prove. Assume that $r_3^{b_i} | r_0^{a_i}$ and $r_0^{a_i} \nmid r_3^{b_i}$ at $p \in \mathbb{A}_U^1$. If $\mathfrak{p} \subset R[T]$ is the prime ideal corresponding to p , then $r_0^{a_i} / r_3^{b_i} \in \mathfrak{p}$ implies that $s' \in \mathfrak{p}$. Thus,

$$h_2^*(I_i)_p = \langle r_3^{b_i} \times r_0^{a_i} / r_3^{b_i}, \left(1 + \frac{s' \delta' T}{1 + s_1 \delta_1} \right) \rangle_p$$

is principal. If $s' \notin \mathfrak{p}$, then r_0/r_3 is a unit at p . Since

$$\frac{r_0^{a_i}}{r_3^{b_i}} = \left(\frac{r_0}{r_3} \right)^{a_i} \left(\frac{1}{r_3} \right)^{b_i - a_i}$$

is regular at p and r_0/r_3 is a unit, $1/r_3$ is regular at p . Therefore, r_3 and hence, r_0 is a unit at p . Consequently, $\frac{r_0^{a_i}}{r_3^{b_i}}$ is a unit at p , showing that $h_2^*(\mathcal{I}_\gamma)$ is a locally principal ideal, which allows us to conclude that h_2 lifts to X_γ . Thus, we have proved that in Case B, the lifts of β_{r_1} and β_{r_2} are \mathbb{A}^1 -chain homotopic if and only if the condition (5.2) holds.

The proof of part (a) can be now completed exactly as in the proof of Theorem 4.1 by observing that when R is a discrete valuation ring, one has $\sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle} = \sqrt{\langle r_1, r' \rangle}$.

We now prove part (b) by constructing a counterexample similar to the one given in the proof of Theorem 4.1. For X as above, consider the 2-dimensional henselian local ring $\tilde{R} = k[x, y]_{\langle x, y \rangle}^h$ with $U = \text{Spec } \tilde{R}$ and consider the morphism $\gamma: U \rightarrow C$ defined by $r_0 = x^{b_1}(y^2 + x)$. Consider $\beta_{r_1}, \beta_{r_2}: U \rightarrow \mathbb{P}_U^1$ determined by $r_1 = x$ and $r_2 = x(y + 1)$. Note that β_{r_1} and β_{r_2} lift to X_γ , since for every $i > 1$ we have $a_i/b_i > a_1/b_1$. Their lifts both map the closed point of U to the point of X_γ defined by the ideal $(y, x^{a_1}/y^{b_1})$, where in fact, $a_1 = 1$. Now, $r' = r_0/r_1 = y^2 + x$ and we have

$$\sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle} = \langle x, y^2 + x \rangle = \langle x, y^2 \rangle,$$

while $\sqrt{\langle r_1, r' \rangle} = \langle x, y \rangle$. Thus

$$r_2/r_1 - 1 = y \notin \sqrt{\langle r_1 \rangle} + \sqrt{\langle r' \rangle}.$$

By the criterion (5.2), we conclude that the lifts of β_{r_1} and β_{r_2} are not \mathbb{A}^1 -chain homotopic, but are 1-ghost homotopic by [5, Proof of Theorem 5.1, Claim 2]. Consequently, $\mathcal{S}(X)(U) \neq \mathcal{S}^2(X)(U)$. By Remark 2.5, it follows that $\text{Sing}_*^{\mathbb{A}^1} X$ is not \mathbb{A}^1 -local. \square

We are now set to prove Theorem 1 and Theorem 2 for a general smooth projective ruled surface. Since the method used here is exactly parallel to the one used in [5, Section 6], we will only outline the proof and refer the reader to [5] for details.

Proofs of Theorem 1 and Theorem 2. Let E be the projectivization of a rank 2 vector bundle over a smooth projective curve C of genus $g > 0$ over an algebraically closed field k of characteristic 0. Let X be a smooth projective surface with minimal model E . Let U be a smooth henselian local scheme over k . Since C is \mathbb{A}^1 -rigid, by [5, Lemma 2.12], if two morphisms $U \rightarrow X$ are \mathbb{A}^1 -chain homotopic, then their compositions with the map $X \rightarrow C$ have to be the same. Thus, we fix a morphism $\gamma: U \rightarrow C$ and characterize \mathbb{A}^1 -chain homotopy classes of morphisms $U \rightarrow X$ over γ . Note that all the \mathbb{A}^1 -chain homotopies of γ -morphisms $U \rightarrow X$ factor through the pullback of X by the morphism $\text{Spec } \mathcal{O}_{C, c_0}^h \rightarrow C$. Note that the pullback of E by the morphism $\text{Spec } \mathcal{O}_{C, c_0}^h \rightarrow C$ is just $\mathbb{P}^1 \times \text{Spec } \mathcal{O}_{C, c_0}^h$. We can therefore replace C with $\text{Spec } \mathcal{O}_{C, c_0}^h$ and assume that C is as in Notation 3.1. Since the case $X = E$ is clear from [5, Proposition 2.13], we will henceforth assume that X is obtained by blowing up \mathbb{P}_C^1 at nonzero, finitely many successive blowups at smooth, closed points $\{Q_1, \dots, Q_m\}$ and by blowing up the ideal \mathcal{J} . For any $Q \in \{Q_1, \dots, Q_m\}$, points in the exceptional divisor over Q are said to be *infinitely near* to Q . Let N_Q denote the number of points in $\{Q_1, \dots, Q_m\}$, which are infinitely near to Q . Define

$$N_X := \max\{N_{Q_1}, \dots, N_{Q_m}\}.$$

Theorem 1 is proved by induction on N_X , which should be viewed as the *complexity* of X as a blowup of \mathbb{P}_C^1 .

Fix a smooth 1-dimensional henselian local ring (R, \mathfrak{m}) , $U = \text{Spec } R$ and a morphism $\gamma: U \rightarrow C$. Denote the pullback of X with respect to γ by X_γ . Let $\alpha, \alpha': U \rightarrow \mathbb{P}_C^1$ be morphisms over γ that lift to X . In [5, Proof of Theorem 1.5], it is shown that we can find a smooth, projective ruled surface X' such that

- X' is obtained by taking successive blowups of finitely many smooth, closed points of \mathbb{P}_C^1 ;
- \tilde{X} is an open subscheme of X' ;
- α, α' lift to X' and their lifts in X' are n -ghost homotopic if and only if the lifts of α, α' in \tilde{X} are n -ghost homotopic; and
- $N_{X'} < N_X$.

We observe that \mathbb{A}^1 -chain homotopies are nothing but 0-ghost homotopies. One can now finish the proof of Theorem 1 using Theorem 5.1 and the induction hypothesis.

For the proof of Theorem 2, observe that one can factor $X \rightarrow \mathbb{P}_C^1$ as $X \rightarrow Y \rightarrow \mathbb{P}_C^1$ such that Y is a nodal blowup. One can now take the counterexample given in the proof of Theorem 4.1 and observe that it lifts to Y_γ and X_γ to obtain $\mathcal{S}(X) \neq \mathcal{S}^2(X)$. By Remark 2.5, it follows that $\text{Sing}_*^{\mathbb{A}^1} X$ is not \mathbb{A}^1 -local, when X is not a minimal model. \square

REFERENCES

- [1] A. Asok, M. Hoyois and M. Wendt: *Affine representability results in \mathbb{A}^1 -homotopy theory II: principal bundles and homogeneous spaces*, Geom. Topol. 22 (2018), no. 2, 1181–1225. [1](#)
- [2] A. Asok and F. Morel: *Smooth varieties up to \mathbb{A}^1 -homotopy and algebraic h-cobordisms*. Adv. Math. 227(2011), no. 5, 1990–2058. [1](#), [2](#), [4](#)
- [3] C. Balwe, A. Hogadi and A. Sawant: *\mathbb{A}^1 -connected components of schemes*, Adv. Math. 282 (2015), 335–361. [1](#), [2](#), [3](#)
- [4] C. Balwe and A. Sawant: *R-equivalence and \mathbb{A}^1 -connectedness in anisotropic groups*, Int. Math. Res. Not. IMRN 2015, No. 22, 11816–11827. [1](#)
- [5] C. Balwe and A. Sawant: *\mathbb{A}^1 -connected components of ruled surfaces*, to appear Geom. Topol., Preprint - arXiv:1911.05549 [math.AG]. [2](#), [3](#), [4](#), [5](#), [8](#), [10](#), [11](#), [12](#), [14](#)
- [6] F. Morel and V. Voevodsky: *\mathbb{A}^1 -homotopy theory of schemes*, Inst. Hautes Études Sci. Publ. Math. 90(1999) 45–143. [3](#)

DEPARTMENT OF MATHEMATICAL SCIENCES, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH
MOHALI, KNOWLEDGE CITY, SECTOR-81, MOHALI 140306, INDIA.

Email address: `cbalwe@iisermohali.ac.in`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, CO-
LABA, MUMBAI 400005, INDIA.

Email address: `asawant@math.tifr.res.in`