

The effective potential and universality in GUT inspired gauge-Higgs unification

Shuichiro Funatsu¹, Hisaki Hatanaka², Yutaka Hosotani³,
Yuta Orikasa⁴ and Naoki Yamatsu⁵

¹*Institute of Particle Physics and Key Laboratory of Quark and Lepton Physics (MOE),
Central China Normal University, Wuhan, Hubei 430079, China*

²*Osaka, Osaka 536-0014, Japan*

³*Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan*

⁴*Institute of Experimental and Applied Physics, Czech Technical University in Prague,
Husova 240/5, 110 00 Prague 1, Czech Republic*

⁵*Department of Physics, Kyushu University, Fukuoka 819-0395, Japan*

Abstract

The effective potential for the Aharonov-Bohm phase θ_H in the fifth dimension in GUT inspired $SO(5) \times U(1) \times SU(3)$ gauge-Higgs unification is evaluated to show that dynamical electroweak symmetry breaking takes place with $\theta_H \neq 0$, the 4D Higgs boson mass 125 GeV being generated at the quantum level. The cubic and quartic self-couplings (λ_3, λ_4) of the Higgs boson are found to satisfy universal relations, i.e. they are determined, to high accuracy, solely by θ_H , irrespective of values of other parameters in the model. For $\theta_H = 0.1$ (0.15), λ_3 and λ_4 are smaller than those in the standard model by 7.7% (8.1%) and 30% (32%), respectively.

1 Introduction

The Higgs boson is responsible for the electroweak symmetry breaking in the standard model (SM). The Higgs potential is arranged such that the Higgs field spontaneously develops a nonvanishing vacuum expectation value. Its couplings to quarks and leptons (Yukawa couplings) are determined such that the observed quark-lepton mass spectrum is reproduced. Although the SM seems consistent with almost all experimental data so far obtained, it is yet to be seen whether or not the Higgs boson is exactly what is postulated in the SM. Unlike the gauge sector in the SM, the Higgs sector lacks a principle, which leaves arbitrariness in the theory. The Higgs boson mass acquires large quantum corrections which must be cancelled by fine-tuning of parameters in the model.

One approach to overcome these difficulties is gauge-Higgs unification in which the 4D Higgs boson is identified with the 4D fluctuation mode of an Aharonov-Bohm (AB) phase in the fifth dimension. The 4D Higgs field is contained in the extra-dimensional component of gauge potentials. As an AB phase the Higgs boson is massless at the tree level, but acquires a finite mass at the quantum level, independent of a cutoff scale and regularization method. The gauge hierarchy problem is naturally solved.[1]-[6]

Recently substantial advances have been made in gauge-Higgs unification. Realistic models have been constructed which yield nearly the same phenomenology as the SM at low energies and give many predictions to be explored at LHC and ILC. Most of gauge-Higgs unification models are constructed on orbifolds such as $M^4 \times (S^1/Z_2)$ and the Randall-Sundrum (RS) warped space. Chiral fermions naturally emerge on orbifolds.[7] The $SU(2)_L$ doublet Higgs field must appear as a zero mode of the fifth-dimensional component of gauge fields. This condition leads to gauge groups such as $SU(3) \times U(1)_X \times SU(3)_C$ or $SO(5) \times U(1)_X \times SU(3)_C$, among which the latter accommodates the custodial symmetry in the Higgs sector. Quark-lepton multiplets are introduced such that with orbifold conditions specified zero modes appear precisely for quarks and leptons, but not for exotic light fermions. They must have observed couplings to W and Z bosons, and their masses must be reproduced. Further the effective potential for the AB phase θ_H must have a global minimum at $\theta_H \neq 0$ so that the electroweak gauge symmetry is dynamically broken to $U(1)_{\text{EM}}$. As a model satisfying these conditions $SO(5) \times U(1)_X \times SU(3)_C$ gauge-Higgs unification is formulated in the RS space.[8]-[20]

In the RS space, which is an AdS spacetime sandwiched by UV and IR branes, wave functions of dominant components of W and Z bosons are almost constant in the bulk region so that gauge couplings of quarks and leptons turn out nearly the same as those

in the SM. The hierarchy between the Kaluza-Klein (KK) mass scale (~ 10 TeV) and the weak scale (~ 100 GeV) naturally emerges. Two typical ways of introducing fermions have been investigated. In one type of the models (the A model) quarks and leptons are introduced in the vector representation of $SO(5)$. The model predicts large parity violation in the Z' couplings of quarks and leptons, which can be checked in the early stage of the ILC experiments with polarized electron and positron beams.[14, 17, 21, 22]

It has been noticed, however, that there arises a difficulty in promoting the A model to grand unification.[23]-[27] The natural extension of the $SO(5) \times U(1)_X \times SU(3)_C$ model is $SO(11)$ gauge-Higgs grand unification.[24] Up-type quarks are contained in the spinor representation of $SO(11)$, but not in the vector representation so that up-type quarks in the A model do not appear from the $SO(11)$ gauge-Higgs unification. A new way of introducing fermion multiplets has been found which can be embedded into the $SO(11)$ gauge-Higgs grand unification.[16] In this GUT inspired model, or the B model, quarks and leptons are introduced in the spinor and singlet representations of $SO(5)$. It has been shown that quarks and leptons have correct gauge couplings. Furthermore the flavor mixing is nicely incorporated with gauge-invariant brane interactions in the B model. The CKM matrix is obtained, and remarkably flavor changing neutral current (FCNC) interactions are naturally suppressed.[20]

In this paper we evaluate the effective potential $V_{\text{eff}}(\theta_H)$ in GUT inspired $SO(5) \times U(1)_X \times SU(3)_C$ gauge-Higgs unification. It will be shown that with appropriate choice of parameters $V_{\text{eff}}(\theta_H)$ has global minimum at $\theta_H \neq 0$ and the Higgs boson mass $m_H = 125$ GeV is obtained. The cubic and quartic self-couplings of the Higgs boson are determined from $V_{\text{eff}}(\theta_H)$. We shall show that those cubic and quartic self-couplings are, to high accuracy, determined as functions of θ_H only. They do not depend on other parameters of the theory. It will be explained how this universality results in the model.

The effective potential $V_{\text{eff}}(\theta_H)$ is important in discussing phase transitions at finite temperature as well. Recently a possibility of having first-order phase transitions in gauge-Higgs unification has been argued.[28] At the moment the nature of phase transitions at finite temperature in $SO(5) \times U(1) \times SU(3)$ gauge-Higgs unification remains unclear.

The Higgs boson as an AB phase in gauge-Higgs unification has similarity to that in composite Higgs models in which the Higgs boson appears as a pseudo-Nambu-Goldstone boson.[9, 29, 30] In both scenarios the Higgs boson field has a character of a phase, but has a quite different mechanism for acquiring its mass. In gauge-Higgs unification the Higgs boson mass is generated by gauge-invariant dynamics of the AB phase, whereas it results from ungauged part of global symmetry in composite Higgs models. Further in

gauge-Higgs unification left-handed and right-handed components of quarks and leptons are normally localized in opposite branes; if left-handed components are localized near UV (IR) brane, then right-handed components are localized near IR (UV) brane. In typical composite Higgs models all light quarks and leptons are assumed to be localized near UV brane. This leads to big difference in phenomenology associated with Z' or techni-rho bosons. In gauge-Higgs unification in RS space there appears large parity violation in Z' couplings of quarks and leptons,[14, 31] whereas such asymmetry is absent in composite Higgs models. Gauge-Higgs unification is strictly regulated by gauge principle.

The paper is organized as follows. In Section 2 the model is introduced. In Section 3 the effective potential $V_{\text{eff}}(\theta_H)$ is evaluated. We show that dynamical EW symmetry breaking takes place. The cubic and quartic self-couplings, λ_3 and λ_4 , of the Higgs boson are evaluated from $V_{\text{eff}}(\theta_H)$. It is observed there that λ_3 and λ_4 are determined to high accuracy as functions of θ_H , irrespective of other parameters in the model. The origin of the θ_H universality in the RS space is clarified in Section 4. In Section 5 the spectrum of dark fermions is evaluated. Section 6 is devoted to summary. Mass spectra of all fields in the model are summarized in Appendix A. Functions used for the evaluation of $V_{\text{eff}}(\theta_H)$ are summarized in Appendix B.

2 Model

The GUT inspired $SO(5) \times U(1)_X \times SU(3)_C$ gauge-Higgs unification has been introduced in refs. [16, 20]. It is defined in the RS warped space with metric given by[32]

$$ds^2 = g_{MN}dx^M dx^N = e^{-2\sigma(y)}\eta_{\mu\nu}dx^\mu dx^\nu + dy^2, \quad (2.1)$$

where $M, N = 0, 1, 2, 3, 5$, $\mu, \nu = 0, 1, 2, 3$, $y = x^5$, $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$, $\sigma(y) = \sigma(y + 2L) = \sigma(-y)$, and $\sigma(y) = ky$ for $0 \leq y \leq L$. In terms of the conformal coordinate $z = e^{ky}$ ($1 \leq z \leq z_L = e^{kL}$) in the region $0 \leq y \leq L$

$$ds^2 = \frac{1}{z^2} \left(\eta_{\mu\nu} dx^\mu dx^\nu + \frac{dz^2}{k^2} \right). \quad (2.2)$$

The bulk region $0 < y < L$ ($1 < z < z_L$) is anti-de Sitter (AdS) spacetime with a cosmological constant $\Lambda = -6k^2$, which is sandwiched by the UV brane at $y = 0$ ($z = 1$) and the IR brane at $y = L$ ($z = z_L$). The KK mass scale is $m_{\text{KK}} = \pi k / (z_L - 1) \simeq \pi k z_L^{-1}$ for $z_L \gg 1$.

In addition to gauge fields $A_M^{SU(3)_C}$, $A_M^{SO(5)}$ and $A_M^{U(1)_X}$ of $SU(3)_C$, $SO(5)$, and $U(1)_X$, we introduce matter fields listed in Table 1. Fields defined in the bulk satisfy orbifold boundary conditions. Each gauge field satisfies

$$\begin{pmatrix} A_\mu \\ A_y \end{pmatrix} (x, y_j - y) = P_j \begin{pmatrix} A_\mu \\ -A_y \end{pmatrix} (x, y_j + y) P_j^{-1} \quad (2.3)$$

where $(y_0, y_1) = (0, L)$. $P_0 = P_1 = I_3$ for $A_M^{SU(3)_C}$ and $P_0 = P_1 = 1$ for $A_M^{U(1)_X}$. $P_0 = P_1 = P_5^{SO(5)} = \text{diag}(I_4, -1)$ for $A_M^{SO(5)}$ in the vector representation and $P_0 = P_1 = P_4^{SO(5)} = \text{diag}(I_2, -I_2)$ in the spinor representation, respectively. Quark and lepton multiplets satisfy

$$\begin{aligned} \Psi_{(\mathbf{3},\mathbf{4})}^\alpha(x, y_j - y) &= -P_4^{SO(5)} \gamma^5 \Psi_{(\mathbf{3},\mathbf{4})}^\alpha(x, y_j + y) , \\ \Psi_{(\mathbf{3},\mathbf{1})}^{\pm\alpha}(x, y_j - y) &= \mp \gamma^5 \Psi_{(\mathbf{3},\mathbf{1})}^{\pm\alpha}(x, y_j + y) , \\ \Psi_{(\mathbf{1},\mathbf{4})}^\alpha(x, y_j - y) &= -P_4^{SO(5)} \gamma^5 \Psi_{(\mathbf{1},\mathbf{4})}^\alpha(x, y_j + y) , \end{aligned} \quad (2.4)$$

where $\alpha = 1 \sim 3$. Dark fermion multiplets satisfy

$$\begin{aligned} \Psi_F^\beta(x, y_j - y) &= (-1)^j P_4^{SO(5)} \gamma^5 \Psi_F^\beta(x, y_j + y) , \\ \Psi_{(\mathbf{1},\mathbf{5})}^{\pm\gamma}(x, y_j - y) &= \pm P_5^{SO(5)} \gamma^5 \Psi_{(\mathbf{1},\mathbf{5})}^{\pm\gamma}(x, y_j + y) , \end{aligned} \quad (2.5)$$

where $\beta = 1 \sim N_F$ and $\gamma = 1 \sim N_V$.

Table 1: $SU(3)_C \times SO(5) \times U(1)_X$ content of matter fields is shown in the GUT inspired B model and previous A model. In the A model only $SU(3)_C \times SO(4) \times U(1)_X$ symmetry is preserved on the UV brane so that the $SU(2)_L \times SU(2)_R$ content is shown for brane fields. The B model is analyzed in the present paper.

	B model	A model
quark	$\Psi_{(\mathbf{3},\mathbf{4})}^\alpha : (\mathbf{3}, \mathbf{4})_{\frac{1}{6}}, \Psi_{(\mathbf{3},\mathbf{1})}^{\pm\alpha} : (\mathbf{3}, \mathbf{1})_{-\frac{1}{3}}^{\pm}$	$\Psi_1^\alpha : (\mathbf{3}, \mathbf{5})_{\frac{2}{3}}, \Psi_2^\alpha : (\mathbf{3}, \mathbf{5})_{-\frac{1}{3}}$
lepton	$\Psi_{(\mathbf{1},\mathbf{4})}^\alpha : (\mathbf{1}, \mathbf{4})_{-\frac{1}{2}}$	$\Psi_3^\alpha : (\mathbf{1}, \mathbf{5})_{-1}, \Psi_4^\alpha : (\mathbf{1}, \mathbf{5})_0$
dark fermion	$\Psi_F^\beta : (\mathbf{3}, \mathbf{4})_{\frac{1}{6}}, \Psi_{(\mathbf{1},\mathbf{5})}^{\pm\gamma} : (\mathbf{1}, \mathbf{5})_0^{\pm}$	$\Psi_F^\delta : (\mathbf{1}, \mathbf{4})_{\frac{1}{2}}$
brane fermion	$\chi^\alpha : (\mathbf{1}, \mathbf{1})_0$	$\hat{\chi}_{1,2,3R}^q : (\mathbf{3}, [\mathbf{2}, \mathbf{1}])_{\frac{7}{6}, \frac{1}{6}, -\frac{5}{6}}$ $\hat{\chi}_{1,2,3R}^l : (\mathbf{1}, [\mathbf{2}, \mathbf{1}])_{-\frac{3}{2}, \frac{1}{2}, -\frac{1}{2}}$
brane scalar	$\Phi_{(\mathbf{1},\mathbf{4})} : (\mathbf{1}, \mathbf{4})_{\frac{1}{2}}$	$\hat{\Phi} : (\mathbf{1}, [\mathbf{1}, \mathbf{2}])_{\frac{1}{2}}$
symmetry of brane interactions	$SU(3)_C \times SO(5) \times U(1)_X$	$SU(3)_C \times SO(4) \times U(1)_X$

The bulk action of each gauge field, $A_M^{SU(3)_C}$, $A_M^{SO(5)}$, or $A_M^{U(1)_X}$, is given by

$$S_{\text{bulk}}^{\text{gauge}} = \int d^5x \sqrt{-\det G} \left[-\text{tr} \left(\frac{1}{4} F^{MN} F_{MN} + \frac{1}{2\xi} (f_{\text{gf}})^2 + \mathcal{L}_{\text{gh}} \right) \right], \quad (2.6)$$

where $\sqrt{-\det G} = 1/kz^5$ and $F_{MN} = \partial_M A_N - \partial_N A_M - ig[A_M, A_N]$ with each 5D gauge coupling constant g . The gauge fixing f_{gf} and ghost terms \mathcal{L}_{gh} have been specified in ref. [16]. Each fermion multiplet $\Psi(x, y)$ in the bulk has its own bulk-mass parameter c . [33] The covariant derivative is given by

$$\begin{aligned}\mathcal{D}(c) &= \gamma^A e_A{}^M \left(D_M + \frac{1}{8} \omega_{MBC} [\gamma^B, \gamma^C] \right) - c\sigma'(y) , \\ D_M &= \partial_M - ig_S A_M^{SU(3)} - ig_A A_M^{SO(5)} - ig_B Q_X A_M^{U(1)} .\end{aligned}\quad (2.7)$$

Here $\sigma' = d\sigma(y)/dy$ and $\sigma'(y) = k$ for $0 < y < L$. g_S, g_A, g_B are $SU(3)_C, SO(5), U(1)_X$ gauge coupling constants. The bulk part of the action for the fermion multiplets are given, with $\bar{\Psi} = i\Psi^\dagger \gamma^0$, by

$$\begin{aligned}S_{\text{bulk}}^{\text{fermion}} &= \int d^5x \sqrt{-\det G} \left\{ \sum_J \bar{\Psi}^J \mathcal{D}(c_J) \Psi^J \right. \\ &\quad \left. - \sum_\alpha \left(m_{D_\alpha} \bar{\Psi}_{(\mathbf{3},1)}^{+\alpha} \Psi_{(\mathbf{3},1)}^{-\alpha} + \text{H.c.} \right) - \sum_\gamma \left(m_{V_\gamma} \bar{\Psi}_{(\mathbf{1},5)}^{+\gamma} \Psi_{(\mathbf{1},5)}^{-\gamma} + \text{H.c.} \right) \right\},\end{aligned}\quad (2.8)$$

where the sum \sum_J extends over $\Psi^J = \Psi_{(\mathbf{3},4)}^\alpha, \Psi_{(\mathbf{1},4)}^\alpha, \Psi_{(\mathbf{3},1)}^{\pm\alpha}, \Psi_F^\beta$ and $\Psi_{(\mathbf{1},5)}^{\pm\gamma}$.

The action for the brane scalar field $\Phi_{(\mathbf{1},4)}(x)$ is given by

$$\begin{aligned}S_{\text{brane}}^\Phi &= \int d^5x \sqrt{-\det G} \delta(y) \\ &\quad \times \left\{ - (D_\mu \Phi_{(\mathbf{1},4)})^\dagger D^\mu \Phi_{(\mathbf{1},4)} - \lambda_{\Phi_{(\mathbf{1},4)}} (\Phi_{(\mathbf{1},4)}^\dagger \Phi_{(\mathbf{1},4)} - |w|^2)^2 \right\},\end{aligned}\quad (2.9)$$

where $D_\mu = \partial_\mu - ig_A A_\mu^{SO(5)} - i\frac{1}{2}g_B A_\mu^{U(1)}$. The action for the gauge-singlet brane fermion $\chi^\alpha(x)$ is

$$S_{\text{brane}}^\chi = \int d^5x \sqrt{-\det G} \delta(y) \left\{ \frac{1}{2} \bar{\chi}^\alpha \gamma^\mu \partial_\mu \chi^\alpha - \frac{1}{2} M^{\alpha\beta} \bar{\chi}^\alpha \chi^\beta \right\}. \quad (2.10)$$

$\chi^\alpha(x)$ satisfies the Majorana condition $\chi^c = \chi$;

$$\chi = \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad \chi^c = \begin{pmatrix} +\eta^c \\ -\xi^c \end{pmatrix} = e^{i\delta_C} \begin{pmatrix} +\sigma^2 \eta^* \\ -\sigma^2 \xi^* \end{pmatrix}. \quad (2.11)$$

On the UV brane there are $SU(3)_C \times SO(5) \times U(1)_X$ -invariant brane interactions among the bulk fermion, brane fermion, and brane scalar fields. Relevant parts of the brane interactions are given by

$$S_{\text{brane}}^{\text{int}} = - \int d^5x \sqrt{-\det G} \delta(y) \times$$

$$\left\{ \kappa^{\alpha\beta} \bar{\Psi}_{(3,4)}^\alpha \Phi_{(1,4)} \cdot \Psi_{(3,1)}^{+\beta} + \tilde{\kappa}^{\alpha\beta} \bar{\chi}^\beta \tilde{\Phi}_{(1,4)}^\dagger \Psi_{(1,4)}^\alpha + \text{H.c.} \right\} \quad (2.12)$$

where κ 's and $\tilde{\kappa}$'s are coupling constants and

$$\Phi_{(1,4)} = \begin{pmatrix} \Phi_{[2,1]} \\ \Phi_{[1,2]} \end{pmatrix}, \quad \tilde{\Phi}_{(1,4)} = \begin{pmatrix} i\sigma^2 \Phi_{[2,1]}^* \\ -i\sigma^2 \Phi_{[1,2]}^* \end{pmatrix}. \quad (2.13)$$

When $\langle \Phi_{(1,4)} \rangle = (0, 0, 0, w)^t$, (2.12) generates additional mass terms

$$\int d^5x \sqrt{-\det G} \delta(y) \left\{ 2\mu^{\alpha\beta} \bar{d}_R'^\alpha D_L^{+\beta} + \text{H.c.} \right\}, \quad \mu^{\alpha\beta} = \frac{\kappa^{\alpha\beta} w}{\sqrt{2}}, \quad (2.14)$$

in the down-type quark sector and

$$- \int d^5x \sqrt{-\det G} \delta(y) \frac{m_B^{\alpha\beta}}{\sqrt{k}} (\bar{\chi}^\beta \nu_R'^\alpha + \bar{\nu}_R'^\alpha \chi^\beta), \quad m_B^{\alpha\beta} = \tilde{\kappa}^{\alpha\beta} w \sqrt{k} \quad (2.15)$$

in the neutrino sector. With the Majorana masses in (2.10), the mass term (2.15) induces inverse seesaw mechanism in the neutrino sector.[26] Further $\langle \Phi_{(1,4)} \rangle \neq 0$ breaks $SO(4) \times U(1)_X$ down to $SU(2)_L \times U(1)_Y$. We assume that $w \gg m_{\text{KK}}$. The 4D $SU(2)_L$ gauge coupling is given by $g_w = g_A/\sqrt{L}$. The 5D gauge coupling $g_Y^{5\text{D}}$ of $U(1)_Y$ and the 4D bare Weinberg angle at the tree level, θ_W^0 , are given by

$$g_Y^{5\text{D}} = \frac{g_A g_B}{\sqrt{g_A^2 + g_B^2}}, \quad \sin \theta_W^0 = \frac{s_\phi}{\sqrt{1 + s_\phi^2}}, \quad s_\phi = \frac{g_B}{\sqrt{g_A^2 + g_B^2}}. \quad (2.16)$$

The bare Weinberg angle θ_W^0 with a given θ_H is determined to fit the LEP1 data for $e^+e^- \rightarrow \mu^+\mu^-$ at $\sqrt{s} = m_Z$. [34] Approximately $\sin^2 \theta_W^0 \simeq 0.1140 + 0.1186 \cos \theta_H - 0.0014 \cos 2\theta_H$. Evaluated gauge couplings turn out very close to those in the SM with $\sin^2 \theta_W = 0.2312$. [20]

The 4D Higgs boson $\Phi_H(x)$ is contained in the $SO(5)/SO(4)$ part of $A_y^{SO(5)}$. In the z coordinate $A_z = (kz)^{-1} A_y$ ($1 \leq z \leq z_L$), and

$$\begin{aligned} A_z^{(j5)}(x, z) &= \frac{1}{\sqrt{k}} \phi_j(x) u_H(z) + \dots, \\ u_H(z) &= \sqrt{\frac{2}{z_L^2 - 1}} z, \\ \Phi_H(x) &= \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_2 + i\phi_1 \\ \phi_4 - i\phi_3 \end{pmatrix}. \end{aligned} \quad (2.17)$$

At the quantum level Φ_H develops a nonvanishing expectation value. Without loss of generality we assume $\langle\phi_1\rangle, \langle\phi_2\rangle, \langle\phi_3\rangle = 0$ and $\langle\phi_4\rangle \neq 0$, which is related to the Aharonov-Bohm (AB) phase θ_H in the fifth dimension. Eigenvalues of

$$\hat{W} = P \exp \left\{ i g_A \int_{-L}^L dy A_y \right\} \cdot P_1 P_0 \quad (2.18)$$

are gauge invariant. For $A_y = (2k)^{-1/2} \phi_4(x) v_H(y) T^{(45)}$, where $v_H(y) = k e^{ky} u_H(z)$ for $0 \leq y \leq L$ and $v_H(-y) = v_H(y) = v_H(y + 2L)$, one finds

$$\begin{aligned} \hat{W} &= \exp \left\{ i f_H^{-1} \phi_4(x) \cdot 2T^{(45)} \right\} , \\ f_H &= \frac{2}{g_A} \sqrt{\frac{k}{z_L^2 - 1}} = \frac{2}{g_w} \sqrt{\frac{k}{L(z_L^2 - 1)}} , \\ \theta_H &= \frac{\langle\phi_4\rangle}{f_H} . \end{aligned} \quad (2.19)$$

Note

$$A_z^{(45)}(x, z) = \frac{1}{\sqrt{k}} \{ \theta_H f_H + H(x) \} u_H(z) + \cdots \quad (2.20)$$

where $H(x)$ is the neutral Higgs boson field. There is a large gauge transformation which shifts θ_H by 2π , preserving the boundary conditions. Physics is invariant under $\theta_H \rightarrow \theta_H + 2\pi$. We shall evaluate the effective potential $V_{\text{eff}}(\theta_H)$ in the next section.

3 Effective potential

The effective potential $V_{\text{eff}}(\theta_H)$ at the one-loop level is evaluated from the mass spectra of all fields which depend on θ_H . After the Wick rotation into the Euclidean signature it is expressed as

$$V_{\text{eff}}(\theta_H) = \sum \pm \frac{1}{2} \int \frac{d^4 p_E}{(2\pi)^4} \sum_n \ln \{ p_E^2 + m_n(\theta_H)^2 \}, \quad (3.1)$$

where the sign $+$ ($-$) corresponds to bosons (fermions). When the Kaluza-Klein (KK) spectrum $\{m_n(\theta_H)\}$ is determined by zeros of a function $\rho(z; \theta_H)$, namely by

$$\rho(m_n; \theta_H) = 0 \quad (n = 1, 2, 3, \cdots), \quad (3.2)$$

then $V_{\text{eff}}(\theta_H)$ is given [35] by

$$V_{\text{eff}}(\theta_H) = \sum \pm \frac{1}{(4\pi)^2} \int_0^\infty dy y^3 \ln \rho(iy; \theta_H) . \quad (3.3)$$

The θ_H -dependent part of $V_{\text{eff}}^{1\text{loop}}(\theta_H)$ is finite, independent of a cutoff and regularization method employed.

The spectrum-determining functions $\rho(z; \theta_H)$ for all fields in the model have been given in ref. [16]. They are summarized in Appendix A for convenience. Relevant contributions come from W and Z gauge fields, top-bottom quark multiplets, and dark fermions in the spinor and vector representations. Contributions from light quarks and leptons are negligible. To avoid unnecessary confusion in the following argument, we denote the effective potential as $V_{\text{eff}}(\theta)$. Physical value θ_H corresponds to the global minimum of $V_{\text{eff}}(\theta)$, namely $dV_{\text{eff}}/d\theta|_{\theta=\theta_H} = 0$. One finds

$$\begin{aligned} V_{\text{eff}}(\theta) = & 2(3 - \xi^2)A_W(\theta) + (3 - \xi^2)A_Z(\theta) + 3\xi^2 A_S(\theta) \\ & - 12A_{\text{top}}(\theta) - 12A_{\text{bottom}}(\theta) - 12n_F A_F(\theta) - 8n_V A_V(\theta) , \\ A_p(\theta) \equiv & \frac{(kz_L^{-1})^4}{(4\pi)^2} \int_0^\infty dq q^3 \ln \left\{ 1 + \sum_{n=1}^2 Q_p^{(n)}(q) \cos(n\theta) \right\} , \end{aligned} \quad (3.4)$$

where n_F and n_V are the number of Ψ_F and $\Psi_{(1,5)}^\pm$, and ξ is a gauge parameter in the generalized R_ξ gauge. The integration variable has been changed from y in (3.3) to $q = k^{-1}z_L y$. In the following we take $z_L = e^{kL} = 10^{10}$ and $\xi = 0$. The contributions from W , Z towers and Goldstone boson tower are given by

$$\begin{aligned} Q_W^{(1)}(q) &= Q_Z^{(1)}(q) = Q_S^{(1)}(q) = 0 , \\ Q_W^{(2)}(q) &= -\frac{1}{-4iz_L q^{-1} \hat{C}'(q) \hat{S}(q) + 1} , \\ Q_Z^{(2)}(q) &= -\frac{1 + s_\phi^2}{-4iz_L q^{-1} \hat{C}'(q) \hat{S}(q) + 1 + s_\phi^2} , \\ Q_S^{(2)}(q) &= -\frac{1}{-2iz_L q^{-1} \hat{C}(q) \hat{S}(q) + 1} . \end{aligned} \quad (3.5)$$

$\hat{C}(q)$, $\hat{S}(q)$ etc. in the expressions above and $\hat{C}_L(q, c)$, $\hat{S}_L(q, c)$ etc. in the expressions below are given in Appendix B. They are expressed in terms of modified Bessel functions.

Top and bottom quark contributions are given by

$$\begin{aligned} Q_{\text{top}}^{(2)}(q) &= Q_{\text{bottom}}^{(2)}(q) = 0 , \\ Q_{\text{top}}^{(1)}(q) &= -\frac{1}{2\hat{S}_L(q; c_t) \hat{S}_R(q; c_t) + 1} , \\ Q_{\text{bottom}}^{(1)}(q) &= -\frac{\widetilde{S_L S_R}}{2\hat{S}_L(q; c_t) \hat{S}_R(q; c_t) \widetilde{S_L S_R} + 2|\mu|^2 \hat{C}_R(q; c_t) \hat{S}_R(q; c_t) \widetilde{C_L S_L} + 1} , \end{aligned}$$

$$\begin{aligned}
\widetilde{S_L S_R} &= \hat{S}_L(q; c_{D_b} + \tilde{m}_{D_b}) \hat{S}_R(q; c_{D_b} - \tilde{m}_{D_b}) + \hat{S}_L(q; c_{D_b} - \tilde{m}_{D_b}) \hat{S}_R(q; c_{D_b} + \tilde{m}_{D_b}) \\
&\quad + \hat{C}_L(q; c_{D_b} + \tilde{m}_{D_b}) \hat{C}_R(q; c_{D_b} - \tilde{m}_{D_b}) + \hat{C}_L(q; c_{D_b} - \tilde{m}_{D_b}) \hat{C}_R(q; c_{D_b} + \tilde{m}_{D_b}) - 2 , \\
\widetilde{C_L S_L} &= \hat{C}_L(q; c_{D_b} + \tilde{m}_{D_b}) \hat{S}_L(q; c_{D_b} - \tilde{m}_{D_b}) + \hat{C}_L(q; c_{D_b} - \tilde{m}_{D_b}) \hat{S}_L(q; c_{D_b} + \tilde{m}_{D_b}) . \quad (3.6)
\end{aligned}$$

In the above expressions we have assumed that the brane interaction term (2.12) is diagonal in generation space. c_{D_α} is the bulk mass parameter of $\Psi_{(3,1)}^{\pm\alpha}$ and $\tilde{m}_{D_\alpha} = m_{D_\alpha}/k$. Numerically the contribution of bottom quark is very small and may be ignored. There are two kinds of dark fermions (Ψ_F^β and $\Psi_{(1,5)}^{\pm\gamma}$). Their contributions are given by

$$\begin{aligned}
Q_F^{(1)}(q) &= \frac{1}{2\hat{S}_L(q; c_F)\hat{S}_R(q; c_F) + 1} , \\
Q_F^{(2)}(q) &= 0 , \\
Q_V^{(1)}(q) &= 0 , \\
Q_V^{(2)}(q) &= -\frac{2}{\hat{B}_0(q; c_V, \tilde{m}_V)} , \\
\hat{B}_0(q; c_V, \tilde{m}_V) &= \\
&\quad \hat{C}_L(q; c_V + \tilde{m}_V) \hat{C}_R(q; c_V - \tilde{m}_V) + \hat{C}_L(q; c_V - \tilde{m}_V) \hat{C}_R(q; c_V + \tilde{m}_V) \\
&\quad + \hat{S}_L(q; c_V + \tilde{m}_V) \hat{S}_R(q; c_V - \tilde{m}_V) + \hat{S}_L(q; c_V - \tilde{m}_V) \hat{S}_R(q; c_V + \tilde{m}_V) . \quad (3.7)
\end{aligned}$$

For the sake of simplicity we set degenerate bulk mass parameters c_F for Ψ_F^β , and degenerate masses $m_V = k\tilde{m}_V$ and bulk mass parameters c_V for $\Psi_{(1,5)}^{\pm\gamma}$. We note that contributions from gauge bosons and $\Psi_{(1,5)}^{\pm\gamma}$ fields to $V_{\text{eff}}(\theta)$ are periodic in θ with a period π , whereas those from top-bottom quarks and Ψ_F^β fields are periodic with a period 2π .

The parameters of the model are determined in the following steps. (i) We pick the value of θ_H . In other words we are going to adjust the parameters of the model such that $V_{\text{eff}}(\theta)$ has a global minimum at $\theta = \theta_H$. (ii) We take $z_L = 10^{10}$. Then k is determined for m_Z to be reproduced, and the KK mass scale $m_{\text{KK}} = \pi k(z_L - 1)^{-1}$ is fixed. (iii) The bulk mass parameters of $\Psi_{(3,4)}^\alpha$ and $\Psi_{(1,4)}^\alpha$ are fixed from the masses of up-type quarks and charged leptons. In particular, c_t is determined by m_t . (iv) The bulk mass parameters c_{D_α} of $\Psi_{(3,1)}^{\pm\alpha}$ and brane interaction coefficients $\mu^{\alpha\beta}$ are determined so as to reproduce the masses of down-type quarks and CKM matrix. Similarly the Majorana mass terms $M^{\alpha\beta}$ and brane interactions $\tilde{\kappa}^{\alpha\beta}$ are determined so as to reproduce neutrino masses and PMNS matrix. As remarked above, these parameters are numerically irrelevant for $V_{\text{eff}}(\theta)$. (v) At

this stage there remain five parameters to be determined; (n_F, c_F) of Ψ_F^β and (n_V, c_V, \tilde{m}_V) of $\Psi_{(1,5)}^{\pm\gamma}$. There are two conditions to be satisfied;

$$\begin{aligned} (a) : \quad & \left. \frac{dV_{\text{eff}}}{d\theta} \right|_{\theta=\theta_H} = 0 , \\ (b) : \quad & m_H^2 = \left. \frac{1}{f_H^2} \frac{d^2 V_{\text{eff}}}{d\theta^2} \right|_{\theta=\theta_H} , \end{aligned} \quad (3.8)$$

where $m_H = 125.1 \text{ GeV}$. The second condition for the Higgs boson mass m_H follows from the fact that the effective potential for the 4D Higgs field $H(x)$ is given by $V_{\text{eff}}(\theta_H + f_H^{-1}H)$ as inferred from (2.20). The conditions (3.8) give two constraints to be satisfied among the five parameters $(n_F, c_F, n_V, c_V, \tilde{m}_V)$. We first fix, for instance, (n_F, n_V, c_V) and determine (c_F, \tilde{m}_V) by (3.8).

One may wonder whether the arbitrary choice of the parameters in the last step diminishes prediction power of the model. Quite surprisingly many of the physical quantities do not depend on such details in the parameter choice, being determined solely by θ_H . There appears the θ_H -universality which will be explained in the next section.

We give some examples. The parameters fixed in the steps (i) to (iv) above are tabulated in Table 2. In Fig. 1 the effective potential for $\theta_H = 0.1$, $n_F = n_V = 2$ and $c_V = 0$ is displayed. $c_F = 0.319$ and $\tilde{m}_V = 0.0806$ are chosen to satisfy (3.8). One observes that the electroweak symmetry is dynamically broken. In Fig. 2 contributions of relevant fields to the effective potential $V_{\text{eff}}(\theta)$ are displayed. There is a lower bound for θ_H in order to reproduce the top quark mass. $\theta_H \geq \theta_{c1}$ where $\theta_{c1} \sim 0.015$ for $z_L = 10^{10}$. Similarly there is a constraint for the warp factor. For $\theta_H = 0.1$, $n_F = n_V = 2$, the top quark mass is reproduced only if $z_L \geq z_{L1} \sim 10^{8.1}$ and dynamical electroweak symmetry breaking is achieved only if $z_L \leq z_{L2} \sim 10^{15.5}$.

Table 2: Parameters determined for $z_L = 10^{10}$ and $\theta_H = 0.05, 0.1, 0.15$. We have set $\mu^{\alpha\beta} = \mu_\alpha \delta_{\alpha\beta}$, and have taken $c_{D_b} = 1.04$. μ_b is determined so as to reproduce m_b .

θ_H	$k [10^{13} \text{ GeV}]$	$m_{\text{KK}} [\text{TeV}]$	c_t	c_{D_b}	μ_b
0.05	7.68	24.1	-0.226	1.04	0.106
0.10	3.84	12.1	-0.227	1.04	0.104
0.15	2.57	8.07	-0.230	1.04	0.0990

The effective potential $V_{\text{eff}}(\theta)$ has more information. By expanding $V_{\text{eff}}(\theta_H + H/f_H)$, one finds Higgs self-couplings $\lambda_n H^n$. The n -th self-coupling λ_n is given by

$$\lambda_n \equiv \left. \frac{1}{n! f_H^n} \frac{d^n V_{\text{eff}}}{d\theta^n} \right|_{\theta=\theta_H} . \quad (3.9)$$




Figure 1: The effective potential for $\theta_H = 0.1$, $n_F = 2$, $n_V = 2$ and $c_V = 0$. The global minimum is located at $\theta = \theta_H$.




Figure 2: Contributions of relevant fields to the effective potential for $\theta_H = 0.1$, $n_F = 2$, $n_V = 2$ and $c_V = 0$ are displayed.

The couplings λ_3 and λ_4 are plotted in Fig. 3 and Fig. 4 as functions of θ_H for $c_V = 0.2$, $n_F = n_V = 2$. The fitting curves are given by

$$\begin{aligned} \text{B model : } \lambda_3/\text{GeV} &= 39.6 \cos \theta_H - 5.21(1 + \cos 2\theta_H) - 0.00911 \cos 3\theta_H , \\ \lambda_4 &= -0.0695 + 0.0852 \cos \theta_H + 0.00725 \cos 2\theta_H . \end{aligned} \quad (3.10)$$

In Figs. 3 and 4, λ_3 and λ_4 in the A model are also plotted, for which the fitting curves are given by

$$\begin{aligned} \text{A model : } \lambda_3/\text{GeV} &= 32.4 \cos \theta_H - 2.26(1 + \cos 2\theta_H) - 1.1 \cos 3\theta_H , \\ \lambda_4 &= -0.00264 - 0.0129 \cos \theta_H + 0.0363 \cos 2\theta_H . \end{aligned} \quad (3.11)$$

Note that λ_3 vanishes at $\theta_H = \frac{1}{2}\pi$ as a consequence of the H parity in GHU models.[36] For $\theta_H \gtrsim 0.6$, λ_4 becomes negative, which, however, does not mean the instability. The θ -dependent part of $V_{\text{eff}}(\theta)$ is finite, bounded from below. In gauge-Higgs unification there does not arise the vacuum instability problem which afflicts most of 4D field theories.

From the experimental constraints from the LEP1, LEP2 data and from the LHC data for the nonobservation of Z' events it is inferred that $\theta_H \lesssim 0.11$. For $\theta_H \sim 0.1$ (0.15), λ_3 and λ_4 are smaller than those in the SM by 7.7% (8.1%) and 30% (32%), respectively. As explained above, there is a lower bound for θ_H in GHU, namely $\theta_H > \theta_{c1}$, and there does not exist $\theta_H \rightarrow 0$ limit. It is not surprising that λ_3 and λ_4 deviate from the values in the SM even for small θ_H . The couplings λ_n ($n \geq 5$) are generated at the one loop level both in GHU and in the SM, which turn out finite. The Higgs couplings to quarks, leptons, and W , Z bosons in GHU for small θ_H are very close to those in the SM. There are additional contributions coming from KK modes in GHU. It may be interesting as well to measure these couplings λ_n ($n \geq 5$) in future experiments to test predictions from GHU and the SM.




Figure 3: The cubic coupling λ_3 of the Higgs boson. The fitting curves are given by $\lambda_3/\text{GeV} = 39.6 \cos \theta_H - 5.21(1 + \cos 2\theta_H) - 0.00911 \cos 3\theta_H$ for the B-model and $\lambda_3/\text{GeV} = 32.4 \cos \theta_H - 2.26(1 + \cos 2\theta_H) - 1.1 \cos 3\theta_H$ for the A-model. The SM value is $\lambda_{3,SM} = 31.5 \text{ GeV}$.




Figure 4: The quartic coupling λ_4 of the Higgs boson. The fitting curves are given by $\lambda_4 = -0.0695 + 0.0852 \cos \theta_H + 0.00725 \cos 2\theta_H$ for the B-model and $\lambda_4 = -0.00264 - 0.0129 \cos \theta_H + 0.0363 \cos 2\theta_H$ for the A-model. The SM value is $\lambda_{4,SM} = 0.0320$.

Table 3: θ_H universality in λ_3 and λ_4 for $\theta_H = 0.1$ and $z_L = 10^{10}$. With given (n_F, n_V, c_V) , c_F and \tilde{m}_V are determined to satisfy the condition (3.8), and λ_3 and λ_4 are evaluated by (3.9).

n_F	n_V	c_V	c_F	\tilde{m}_V	$\lambda_3(\text{GeV})$	λ_4
2	2	0.	0.319	0.0806	29.03	0.02083
2	2	0.2	0.319	0.0777	29.03	0.02083
2	2	0.5	0.322	-0.0371	29.02	0.02078
4	2	0.	0.425	0.0794	29.02	0.02082
4	2	0.2	0.425	0.0765	29.02	0.02082
4	2	0.5	0.426	-0.0350	29.01	0.02076
2	4	0.	0.318	0.0964	29.03	0.02084
2	4	0.2	0.318	0.0937	29.03	0.02084
2	4	0.5	0.319	0.0615	29.03	0.02083

4 θ_H universality

As remarked in the previous section there remains the arbitrariness in the choice of the parameters in the model. Among the five parameters $(n_F, c_F, n_V, c_V, \tilde{m}_V)$ there are only two conditions in (3.8) to be obeyed. In the examples given in the previous section we first fixed (n_F, n_V, c_V) and determined (c_F, \tilde{m}_V) by (3.8). The Higgs cubic and quartic couplings, λ_3 and λ_4 , are evaluated with this choice. One might wonder how λ_3 and λ_4 depend on the choice of the parameters (n_F, n_V, c_V) .

In this section we shall show that λ_3 and λ_4 are determined, to high accuracy, as functions of θ_H only, but do not depend on the details of the parameter choice. It has been known in GHU that the 3-point Higgs couplings to W , Z , quarks and leptons also have the same property.[37] These physical quantities are determined by θ_H to high accuracy. It may be called as the θ_H universality. The θ_H universality leads to profound power for predictions. Once the value of θ_H is determined by one of the physical quantities, then the values of other physical quantities are predicted.

In Table 3 evaluated values of (λ_3, λ_4) for $\theta_H = 0.1$ are shown with various choices of (n_F, n_V, c_V) . Although the values of determined c_F and \tilde{m}_V depend on the choice of (n_F, n_V, c_V) , the evaluated values of λ_3 and λ_4 are universal to high accuracy. λ_3 and λ_4 are determined as functions of θ_H only.

There is a reason for the θ_H universality. We first examine global behavior of $V_{\text{eff}}(\theta)$ with a given θ_H . Notice that the function $A_p(\theta)$ in (3.4) is expanded as

$$A_p(\theta) = \frac{(kz_L^{-1})^4}{(4\pi)^2} \sum_{\ell=1}^{\infty} \sum_{n=1}^2 \alpha_p^{(n,\ell)} \cos^\ell n\theta ,$$

$$\alpha_p^{(n,\ell)} \equiv \frac{(-1)^{\ell+1}}{\ell} \int_0^\infty dq q^3 (Q_p^{(n)}(q))^\ell . \quad (4.1)$$

As either $Q_p^{(1)}(q)$ or $Q_p^{(2)}(q)$ with given p vanishes in (3.4), $\alpha_p^{(1,\ell)}$ or $\alpha_p^{(2,\ell)} = 0$ for each p in (4.1).

To understand qualitative behavior of $V_{\text{eff}}(\theta)$, let us approximate $A_p(\theta)$ in (3.4) by

$$\frac{(4\pi)^2}{(kz_L^{-1})^4} A_p(\theta) = \begin{cases} \alpha_p^{(2,1)} \cos 2\theta & \text{for } p = W, Z, S, V, \\ \alpha_p^{(1,1)} \cos \theta + \alpha_p^{(1,2)} \cos^2 \theta & \text{for } p = \text{top}, F. \end{cases} \quad (4.2)$$

Note that $|\alpha_p^{(n,2)}/\alpha_p^{(n,1)}| < 0.05$. As the contributions from $p = \text{top}, F$ are one order of magnitude larger than those from $p = W, Z, S, V$, the $\cos^2 \theta$ terms have been retained for top and F . $V_{\text{eff}}(\theta)$ in this approximation, denoted as $V_{\text{app}}(\theta)$, is given by

$$\begin{aligned} V_{\text{app}}(\theta) &= \frac{(kz_L^{-1})^4}{(4\pi)^2} \left(-B_1 \cos \theta + B_2 \cos 2\theta \right) , \\ B_1 &= 12\alpha_{\text{top}}^{(1,1)} + 12n_F \alpha_F^{(1,1)} , \\ B_2 &= \alpha_{\text{gauge}} - 8n_V \alpha_V^{(2,1)} - 6\alpha_{\text{top}}^{(1,2)} - 6n_F \alpha_F^{(1,2)} , \\ \alpha_{\text{gauge}} &= 2(3 - \xi^2) \alpha_W^{(2,1)} + (3 - \xi^2) \alpha_Z^{(2,1)} + 3\xi^2 \alpha_S^{(2,1)} . \end{aligned} \quad (4.3)$$

The condition (a) in (3.8) leads to $B_1 = 4B_2 \cos \theta_H$. Then the condition (b) in (3.8) implies that

$$B_2 \sim \frac{16\pi^2}{g_w^2 (kL)^2} \left(\frac{m_H}{m_W} \right)^2 \quad (4.4)$$

where the relations $m_{\text{KK}} \sim \pi k z_L^{-1}$, $m_W \sim (k/L)^{1/2} z_L^{-1} \sin \theta_H$, $f_H \sim 2m_W/g_w \sin \theta_H$ have been made use of. It follows that

$$\begin{aligned} V_{\text{app}}(\theta) &= V_0 u(\theta) , \\ u(\theta) &= -4 \cos \theta_H \cos \theta + \cos 2\theta , \\ V_0 &= \frac{m_W^2 m_H^2}{g_w^2 \sin^4 \theta_H} . \end{aligned} \quad (4.5)$$

The cubic and quartic Higgs self-couplings are given by

$$\begin{aligned} \lambda_3^{\text{app}} &\sim \frac{g_w m_H^2}{4m_W} \cos \theta_H , \\ \lambda_4^{\text{app}} &\sim \frac{g_w^2 m_H^2}{96m_W^2} (7 \cos^2 \theta_H - 4) . \end{aligned} \quad (4.6)$$

The approximate formulas (4.5) and (4.6) represent qualitative behavior of the effective potential $V_{\text{eff}}(\theta)$, but exhibit slight deviation from the values in Table 3 and the fitting curves (3.10) and (3.11). We first note that the form of $V_{\text{app}}(\theta)$ is fixed, once one makes an Ansatz that $V_{\text{eff}}(\theta)$ is expressed in terms of two functions $\cos \theta$ and $\cos 2\theta$. The relevant quantities are B_1 and B_2 , but not detailed values of the parameters in the models considered. In the A- or B-model the same universality relations (4.6) result in this approximation. It is easy to confirm that the formulas (4.6) reproduce the SM values at $\theta_H = 0$;

$$\lambda_3^{\text{app}}|_{\theta_H=0} = \lambda_{3,SM} \ , \quad \lambda_4^{\text{app}}|_{\theta_H=0} = \lambda_{4,SM} \ . \quad (4.7)$$

We also note that $u(\pi) - u(0) = 8 \cos \theta_H$ and $u(\theta_H) - u(0) = -2(1 - \cos \theta_H)^2$. For small θ_H , $u(\pi) - u(0) \sim 8$ and $u(\theta_H) - u(0) \sim -\frac{1}{2}\theta_H^4$, which explains the behavior of $V_{\text{eff}}(\theta)$ for $\theta_H = 0.1$ seen in fig. 1.

To understand the θ_H universality demonstrated in the previous section, refinement of the arguments is necessary. The universality was first found in the A-model of $SO(5) \times U(1)$ gauge-Higgs unification.[13] The mechanism for yielding the θ_H universality has been explained in ref. [38]. We generalize the argument for the current B-model. The important observation is that λ_3 and λ_4 are determined by the local behavior of the effective potential $V_{\text{eff}}(\theta_H)$ in the vicinity of the global minimum at $\theta = \theta_H$, and the universality reflects the local, but not global behavior of $V_{\text{eff}}(\theta_H)$.

The effective potential $V_{\text{eff}}(\theta)$, (3.4), is decomposed into three parts;

$$V_{\text{eff}}(\theta) = \frac{(kz_L^{-1})^4}{(4\pi)^2} \left\{ h_0(\theta) + n_F h_F(\theta; c_F, z_L) + n_V h_V(\theta; c_V, \tilde{m}_V, z_L) \right\} \quad (4.8)$$

where $h_0(\theta)$ represents the contributions from gauge and top quark fields. With θ_H , z_L and ξ specified, k is determined from m_Z and c_t is subsequently determined by m_t so that $h_0(\theta)$ is fixed. All other parameters associated with quarks and leptons are irrelevant for $V_{\text{eff}}(\theta)$. There remain five parameters $(n_F, c_F, n_V, c_V, \tilde{m}_V)$ to be specified in (4.8). They must be adjusted such that the two conditions in (3.8) are satisfied. The important feature in the RS space with $z_L \gg 1$ is that the θ dependence of $h_F(\theta; c_F, z_L)$ and $h_V(\theta; c_V, \tilde{m}_V, z_L)$ factorizes near $\theta = \theta_H$

$$\begin{aligned} h_F(\theta; c_F, z_L) &\simeq \alpha_F(c_F, z_L) \tilde{h}_F(\theta) \ , \\ h_V(\theta; c_V, \tilde{m}_V, z_L) &\simeq \alpha_V(c_V, \tilde{m}_V, z_L) \tilde{h}_V(\theta) \end{aligned} \quad (4.9)$$

to very high accuracy. This can be confirmed numerically from the formula for $A_p(\theta)$ in (3.4). The relation (4.9) implies, for instance, that the ratio $h_F(\theta; c_F^{(1)}, z_L)/h_F(\theta; c_F^{(2)}, z_L)$

is θ -independent near θ_H . For $\theta_H = 0.1$, $z_L = 10^{10}$, and $(c_F^{(1)}, c_F^{(2)}) = (0.3, 0.4)$, the ratio varies from 1.7916 to 1.7896 in the range $0.09 \leq \theta \leq 0.11$. The variation is only 0.1%. We stress that this factorization formulas are valid only locally, namely near $\theta = \theta_H$, and $\tilde{h}_F(\theta)$ and $\tilde{h}_V(\theta)$ depend on θ_H . The z_L dependence of $h_F(\theta; c_F, z_L)$ and $h_V(\theta; c_V, \tilde{m}_V, z_L)$ is also tiny in the range $10^8 \lesssim z_L \lesssim 10^{15}$.

Let us pick a set of values (n_F, n_V, c_V) and determine (c_F, \tilde{m}_V) by (3.8). Making use of (4.9), one finds

$$\left[\frac{dh_0}{d\theta} + n_F \alpha_F(c_F, z_L) \frac{d\tilde{h}_F}{d\theta} + n_V \alpha_V(c_V, \tilde{m}_V, z_L) \frac{d\tilde{h}_V}{d\theta} \right]_{\theta=\theta_H} = 0 ,$$

$$\left[\frac{d^2 h_0}{d\theta^2} + n_F \alpha_F(c_F, z_L) \frac{d^2 \tilde{h}_F}{d\theta^2} + n_V \alpha_V(c_V, \tilde{m}_V, z_L) \frac{d^2 \tilde{h}_V}{d\theta^2} \right]_{\theta=\theta_H} = \frac{(4\pi)^2 m_H^2 f_H^2}{(kz_L^{-1})^4} . \quad (4.10)$$

We do this procedure for two sets; $(n_F, n_V, c_V) = (n_F^{(1)}, n_V^{(1)}, c_V^{(1)})$ and $(n_F^{(2)}, n_V^{(2)}, c_V^{(2)})$. Then (4.10) implies that

$$n_F^{(1)} \alpha_F(c_F^{(1)}, z_L) = n_F^{(2)} \alpha_F(c_F^{(2)}, z_L) \equiv \beta_F ,$$

$$n_V^{(1)} \alpha_V(c_V^{(1)}, \tilde{m}_V^{(1)}, z_L) = n_V^{(2)} \alpha_V(c_V^{(2)}, \tilde{m}_V^{(2)}, z_L) \equiv \beta_V . \quad (4.11)$$

Although values of (c_F, \tilde{m}_V) depend on the choice of (n_F, n_V, c_V) , $\beta_F = n_F \alpha_F(c_F, z_L)$ and $\beta_V = n_V \alpha_V(c_V, \tilde{m}_V, z_L)$ are universal, provided solutions exist. Consequently one obtains

$$V_{\text{eff}}(\theta) \simeq \frac{(kz_L^{-1})^4}{(4\pi)^2} \bar{h}(\theta) , \quad \bar{h}(\theta) = h_0(\theta) + \beta_F \tilde{h}_F(\theta) + \beta_V \tilde{h}_V(\theta) . \quad (4.12)$$

It immediately follows that

$$\lambda_3(\theta_H) = \frac{g_w m_H^2 \sin \theta_H}{12 m_W} \frac{\bar{h}^{(3)}(\theta_H)}{\bar{h}^{(2)}(\theta_H)} ,$$

$$\lambda_4(\theta_H) = \frac{g_w^2 m_H^2 \sin^2 \theta_H}{96 m_W^2} \frac{\bar{h}^{(4)}(\theta_H)}{\bar{h}^{(2)}(\theta_H)} , \quad (4.13)$$

which explains the θ_H universality observed in the previous section. The relevant quantities for λ_3 and λ_4 are $\beta_F(\theta_H)$ and $\beta_V(\theta_H)$, but not (n_F, n_V, c_V) . As mentioned above, the z_L -dependence of $h_F(\theta; c_F, z_L)$ and $h_V(\theta; c_V, \tilde{m}_V, z_L)$ is weak. The θ_H universality stays valid to good approximation even for varying z_L . For instance, for $\theta_H = 0.1$ and $(n_F, n_V, c_V) = (2, 2, 0)$, the resultant (λ_3, λ_4) is $(28.93 \text{ GeV}, 0.02042)$ for $z_L = 1.237 \times 10^8$, which should be compared to $(29.03 \text{ GeV}, 0.02083)$ for $z_L = 10^{10}$.

The θ_H universality is observed in other physical quantities. The Higgs boson couplings g_{WWH} and g_{ZZH} to W , Z , and Yukawa couplings y_f to quarks and leptons are given, to good approximation, by [20, 37]

$$\begin{aligned} g_{WWH} &= g_w m_W \cos \theta_H , \\ g_{ZZH} &= \frac{g_w m_Z}{\cos \theta_W^0} \cos \theta_H , \\ y_f &= \begin{cases} \frac{m_f}{v_{\text{SM}}} \cos \theta_H & \text{in the A model} \\ \frac{m_f}{v_{\text{SM}}} \cos^2 \frac{1}{2} \theta_H & \text{in the B model} \end{cases} \end{aligned} \quad (4.14)$$

where $v_{\text{SM}} = f_H \sin \theta_H = 2m_W/g_w$. For small θ_H , deviation in the Higgs couplings in (4.14) is small, whereas deviation in λ_3 and λ_4 becomes substantial. We remark that the relations in (4.14) have been derived in the composite Higgs model where the parameter $\sqrt{\xi} = v/f$ corresponds to θ_H in GHU.[29, 30]

5 Dark Fermions

Although the θ_H universality holds for various couplings associated with the Higgs boson, masses of dark fermions Ψ_F and $\Psi_{(1,5)}^\pm$, for instance, sensitively depend on the choice of the parameters (n_F, n_V, c_V) . They are determined by (A.12) for Ψ_F , and by (A.13) and (A.14) for charged and neutral components of $\Psi_{(1,5)}^\pm$. In Table 4 their masses are tabulated for various θ_H with $n_F = n_V = 2$. Dark fermions have relatively small masses compared with the KK mass scale m_{KK} . The lightest neutral component of the dark fermions can be a candidate for dark matter.

Table 4: KK mass and dark fermion masses are shown in the unit of TeV for various θ_H with $z_L = 10^{10}$, $n_F = n_V = 2$ and $c_V = 0.2$. Charged and neutral components of $\Psi_{(1,5)}^\pm$ have nearly the same masses.

θ_H	m_{KK}	Ψ_F	$\Psi_{(1,5)}^\pm$
0.05	24.1	6.30	5.60
0.10	12.1	3.42	2.84
0.15	8.07	2.38	1.91
0.20	6.08	1.82	1.45

In Fig. 5 the mass of Ψ_F is plotted as a function of θ_H for several n_F . The mass decreases as n_F increases. Similar behavior is obtained for $\Psi_{(1,5)}^\pm$ as n_V is varied with c_V fixed.




Figure 5: θ_H and n_F dependence of the mass of dark fermion Ψ_F . $z_L = 10^{10}$, $n_V = 2$, $c_V = 0.2$.

6 Summary

In this paper we have examined the effective potential $V_{\text{eff}}(\theta_H)$ in GUT inspired $SO(5) \times U(1) \times SU(3)$ gauge-Higgs unification to confirm that electroweak symmetry breaking is dynamically induced by the Hosotani mechanism. From $V_{\text{eff}}(\theta_H)$ the cubic and quartic self-couplings, λ_3 and λ_4 , of the Higgs boson are determined. We have shown the θ_H universality of these couplings, i.e. they are determined as functions of θ_H to high accuracy, irrespective of the details of other parameters in the theory. For $\theta_H = 0.1$ (0.15), λ_3 and λ_4 are smaller than those in the standard model by 7.7% (8.1%) and 30% (32%), respectively. The θ_H universality in λ_3 and λ_4 is understood as a result of the factorization property of each component in the contributions to the effective potential, which is valid to high accuracy in the Randall-Sundrum warped space with $z_L \gg 1$.

The θ_H universality gives the model great prediction power. Once the value of θ_H is determined by one of the experimental data, then many other physical quantities such as masses and couplings of various particles are predicted. It has been known that gauge-Higgs unification models in the RS space predict large parity violation in the couplings of quarks and leptons to Z' particles (KK modes of γ , Z and Z_R). Its effect can be clearly seen in electron-positron collision experiments with polarized electron/positron beams in which θ_H is the most important parameter. Z' particles can be directly produced at LHC, and parity-violating couplings would manifest in the rapidity distribution in $t\bar{t}$ production. CKM mixing with natural FCNC suppression is also incorporated in the GUT inspired gauge-Higgs unification. It is curious to pin down the behavior of the model at finite temperature and implications to cosmology. $SO(5) \times U(1) \times SU(3)$ gauge-Higgs

unification is one of the most promising scenarios beyond the standard model. We shall come back to these issues in future.

Acknowledgements

This work was supported in part by European Regional Development Fund-Project Engineering Applications of Microworld Physics (No. CZ.02.1.01/0.0/0.0/16_019/0000766) (Y.O.), by the National Natural Science Foundation of China (Grant Nos. 11775092, 11675061, 11521064, 11435003 and 11947213) (S.F.), by the International Postdoctoral Exchange Fellowship Program (IPEFP) (S.F.), and by Japan Society for the Promotion of Science, Grants-in-Aid for Scientific Research, No. 19K03873 (Y.H.) and Nos. 18H05543 and 19K23440 (N.Y.).

A Mass spectrum

In evaluating the effective potential $V_{\text{eff}}(\theta_H)$ in Section 3, one needs to know the mass spectrum of each KK tower of the fields in the model. It is sufficient to know the form of functions whose zeros determine the mass spectrum. These functions have been given in ref. [16]. We summarize them in this appendix for the convenience.

We first introduce

$$F_{\alpha,\beta}(u, v) \equiv J_\alpha(u)Y_\beta(v) - Y_\alpha(u)J_\beta(v) , \quad (\text{A.1})$$

where $J_\alpha(u)$ and $Y_\alpha(u)$ are the first and second kind Bessel functions. For gauge fields we define

$$\begin{aligned} C(z; \lambda) &= \frac{\pi}{2} \lambda z z_L F_{1,0}(\lambda z, \lambda z_L), \\ S(z; \lambda) &= -\frac{\pi}{2} \lambda z F_{1,1}(\lambda z, \lambda z_L), \\ C'(z; \lambda) &= \frac{\pi}{2} \lambda^2 z z_L F_{0,0}(\lambda z, \lambda z_L), \\ S'(z; \lambda) &= -\frac{\pi}{2} \lambda^2 z F_{1,1}(\lambda z, \lambda z_L). \end{aligned} \quad (\text{A.2})$$

For fermion fields with a bulk mass parameter c , we define

$$\begin{aligned} \begin{pmatrix} C_L \\ S_L \end{pmatrix} (z; \lambda, c) &= \pm \frac{\pi}{2} \lambda \sqrt{z z_L} F_{c+\frac{1}{2}, c \mp \frac{1}{2}}(\lambda z, \lambda z_L) , \\ \begin{pmatrix} C_R \\ S_R \end{pmatrix} (z; \lambda, c) &= \mp \frac{\pi}{2} \lambda \sqrt{z z_L} F_{c-\frac{1}{2}, c \pm \frac{1}{2}}(\lambda z, \lambda z_L) , \end{aligned} \quad (\text{A.3})$$

and

$$\begin{aligned}
\mathcal{C}_{R1}(z; \lambda, c, \tilde{m}) &= C_R(z; \lambda, c + \tilde{m}) + C_R(z; \lambda, c - \tilde{m}), \\
\mathcal{C}_{R2}(z; \lambda, c, \tilde{m}) &= S_R(z; \lambda, c + \tilde{m}) - S_R(z; \lambda, c - \tilde{m}), \\
\mathcal{S}_{L1}(z; \lambda, c, \tilde{m}) &= S_L(z; \lambda, c + \tilde{m}) + S_L(z; \lambda, c - \tilde{m}), \\
\mathcal{S}_{L2}(z; \lambda, c, \tilde{m}) &= C_L(z; \lambda, c + \tilde{m}) - C_L(z; \lambda, c - \tilde{m}), \\
\mathcal{C}_{L1}(z; \lambda, c, \tilde{m}) &= C_L(z; \lambda, c + \tilde{m}) + C_L(z; \lambda, c - \tilde{m}), \\
\mathcal{C}_{L2}(z; \lambda, c, \tilde{m}) &= S_L(z; \lambda, c + \tilde{m}) - S_L(z; \lambda, c - \tilde{m}), \\
\mathcal{S}_{R1}(z; \lambda, c, \tilde{m}) &= S_R(z; \lambda, c + \tilde{m}) + S_R(z; \lambda, c - \tilde{m}), \\
\mathcal{S}_{R2}(z; \lambda, c, \tilde{m}) &= C_R(z; \lambda, c + \tilde{m}) - C_R(z; \lambda, c - \tilde{m}).
\end{aligned} \tag{A.4}$$

A.1 Gauge bosons

The mass spectrum $\{m_n = k\lambda_n\}$ of W and W_R towers is determined by

$$\begin{aligned}
W \text{ tower} &: 2S(1; \lambda)C'(1; \lambda) + \lambda \sin^2 \theta_H = 0, \\
W_R \text{ tower} &: C(1; \lambda) = 0.
\end{aligned} \tag{A.5}$$

The spectrum of γ , Z , Z_R and A_z towers is determined by

$$\begin{aligned}
\gamma \text{ tower} &: C'(1; \lambda) = 0, \\
Z \text{ tower} &: 2S(1; \lambda)C'(1; \lambda) + (1 + s_\phi^2)\lambda \sin^2 \theta_H = 0, \\
Z_R \text{ tower} &: C(1; \lambda) = 0, \\
A_z \text{ tower} &: S(1; \lambda)C'(1; \lambda) + \lambda \sin^2 \theta_H = 0.
\end{aligned} \tag{A.6}$$

A.2 Fermions

With given up-type quark masses $m_Q = (m_u, m_c, m_t)$ the bulk mass parameter $c_Q = (c_u, c_c, c_t)$ of up-type quark multiplets is fixed by

$$S_L(1; \lambda, c_Q)S_R(1; \lambda, c_Q) + \sin^2 \frac{\theta_H}{2} = 0, \tag{A.7}$$

where $\lambda = \lambda_Q = m_Q/k$. Then the spectrum of up-type quark towers is determined by (A.7). In the down-type quark sector there are brane interactions which mix d'^α and $D^{+\beta}$ through (2.14). When brane interactions are diagonal in the generation space, $\mu^{\alpha\beta} = \delta^{\alpha\beta} \mu_\alpha$, the spectrum of down-type quark tower is determined by

$$\left(S_L^Q S_R^Q + \sin^2 \frac{\theta_H}{2} \right) (\mathcal{S}_{L1}^D \mathcal{S}_{R1}^D - \mathcal{S}_{L2}^D \mathcal{S}_{R2}^D)$$

$$+ |\mu_1|^2 C_R^Q S_R^Q (\mathcal{S}_{L1}^D \mathcal{C}_{L1}^D - \mathcal{S}_{L2}^D \mathcal{C}_{L2}^D) = 0 ,$$

$$S_L^Q = S_L(1; \lambda, c_Q), \quad \mathcal{S}_{L1}^D = \mathcal{S}_{L1}(1; \lambda, c_D, \tilde{m}_D), \quad \text{etc.} \quad (\text{A.8})$$

where $\mu_1 = (\mu_d, \mu_s, \mu_b)$, $c_D = (c_{D_d}, c_{D_s}, c_{D_b})$ and $\tilde{m}_D = (\tilde{m}_{D_d}, \tilde{m}_{D_s}, \tilde{m}_{D_b})$. The parameters μ_1, c_D, \tilde{m}_D are determined such that $\lambda = (\lambda_d, \lambda_s, \lambda_b) = k^{-1}(m_d, m_s, m_b)$ solves (A.8) in each generation. For the third generation, for instance, we take $(\mu_b, c_{D_b}, \tilde{m}_{D_d}) = (0.1, 1.044, 1.0)$. Only top quark multiplet among quark multiplets gives a relevant contribution to $V_{\text{eff}}(\theta_H)$. By considering general $\mu^{\alpha\beta}$ the CKM mixing is incorporated with natural FCNC suppression.[20]

With given charged lepton masses $m_L = (m_e, m_\mu, m_\tau)$ the bulk mass parameter $c_L = (c_e, c_\mu, c_\tau)$ of charged lepton multiplets is fixed by

$$S_L(1; \lambda, c_L) S_R(1; \lambda, c_L) + \sin^2 \frac{\theta_H}{2} = 0, \quad (\text{A.9})$$

where $\lambda = \lambda_L = m_L/k$. Then the spectrum of charged lepton towers is determined by (A.9). In the neutrino sector brane interactions mix $\nu^\alpha, \nu'^\alpha, \chi^\alpha$. When both $M^{\alpha\beta} = \delta^{\alpha\beta} M_\alpha$ in (2.10) and $m_B^{\alpha\beta} = \delta^{\alpha\beta} m_B^\alpha$ in (2.15) are diagonal, the spectrum of neutrino tower is determined by

$$(k\lambda - M) \left(S_L^L S_R^L + \sin^2 \frac{\theta_H}{2} \right) + \frac{m_B^2}{k} S_R^L C_R^L = 0 ,$$

$$S_R^L = S_R(1; \lambda, c_L), \quad \text{etc.} \quad (\text{A.10})$$

where $M = (M_1, M_2, M_3)$ and $m_B = (m_B^1, m_B^2, m_B^3)$. With $c_L < -\frac{1}{2}$ the light neutrino mass is given by

$$m_\nu \sim \frac{m_L^2 M}{(2|c_L| - 1) m_B^2} \quad (\text{A.11})$$

in each generation. Contributions from lepton multiplets to $V_{\text{eff}}(\theta_H)$ are negligible.

The spectrum of dark fermion Ψ_F tower is determined by

$$S_L(1; \lambda, c_F) S_R(1; \lambda, c_F) + \cos^2 \frac{\theta_H}{2} = 0 . \quad (\text{A.12})$$

The spectrum of charged components of dark fermions $\Psi_{(1,5)}^\pm$ tower is determined by

$$\mathcal{S}_{L1}(1; \lambda, c_V) \mathcal{S}_{R1}(1; \lambda, c_V) - \mathcal{S}_{L2}(1; \lambda, c_V) \mathcal{S}_{R2}(1; \lambda, c_V) = 0, \quad (\text{A.13})$$

whereas the spectrum of neutral component tower is determined by

$$\{B_0(\lambda, c_V, \tilde{m}_V) - 2 \cos 2\theta_H\}^2 = 0 ,$$

$$\begin{aligned}
B_0(\lambda, c, \tilde{m}) &= C_L(1; \lambda, c + \tilde{m})C_R(1; \lambda, c - \tilde{m}) + C_L(1; \lambda, c - \tilde{m})C_R(1; \lambda, c + \tilde{m}) \\
&+ S_L(1; \lambda, c + \tilde{m})S_R(1; \lambda, c - \tilde{m}) + S_L(1; \lambda, c - \tilde{m})S_R(1; \lambda, c + \tilde{m}).
\end{aligned} \tag{A.14}$$

There are two degenerate towers.¹

B Useful functions

As shown in the formula (3.3), a mass-determining function $\rho(m; \theta_H)$ is analytically continued to $\rho(iy; \theta_H)$. We summarize functions used in the evaluation of $V_{\text{eff}}(\theta_H)$ in Section 3. We introduce

$$\hat{F}_{\alpha, \beta}(u, v) \equiv I_\alpha(u)K_\beta(v) - e^{-i(\alpha - \beta)\pi} K_\alpha(u)I_\beta(v), \tag{B.1}$$

where $I_\alpha(u)$ and $K_\alpha(u)$ are first and second kind modified Bessel functions. In terms of $\hat{F}_{\alpha, \beta}(u, v)$ we define

$$\begin{aligned}
\hat{C}(q) &= q\hat{F}_{1,0}(qz_L^{-1}, q), \\
\hat{S}(q) &= iqz_L^{-1}\hat{F}_{1,1}(qz_L^{-1}, q), \\
\hat{C}'(q) &= q^2z_L^{-1}\hat{F}_{0,0}(qz_L^{-1}, q), \\
\hat{S}'(q) &= -iq^2z_L^{-2}\hat{F}_{1,1}(qz_L^{-1}, q)
\end{aligned} \tag{B.2}$$

for gauge fields. For fermion fields with $c > 0$ we define

$$\begin{aligned}
\hat{C}_L(q; c) &= qz_L^{-1/2}\hat{F}_{c+\frac{1}{2}, c-\frac{1}{2}}(qz_L^{-1}, q), \\
\hat{S}_L(q; c) &= iqz_L^{-1/2}\hat{F}_{c+\frac{1}{2}, c+\frac{1}{2}}(qz_L^{-1}, q), \\
\hat{C}_R(q; c) &= qz_L^{-1/2}\hat{F}_{c-\frac{1}{2}, c+\frac{1}{2}}(qz_L^{-1}, q), \\
\hat{S}_R(q; c) &= -iqz_L^{-1/2}\hat{F}_{c-\frac{1}{2}, c-\frac{1}{2}}(qz_L^{-1}, q).
\end{aligned} \tag{B.3}$$

For $c < 0$, we use the the relations

$$\hat{C}_L(q; -c) = \hat{C}_R(q; c), \quad \hat{S}_L(q; -c) = -\hat{S}_R(q; c). \tag{B.4}$$

References

¹There was a typo in (D.16) of ref. [16]. The last term in the second line, $s_H^2 c_H^2 (\mathcal{C}_{R1}^V \mathcal{C}_{L2}^V - \mathcal{C}_{R2}^V \mathcal{S}_{L1}^V)^2$, should be $s_H^2 c_H^2 (\mathcal{C}_{R1}^V \mathcal{S}_{L2}^V - \mathcal{C}_{R2}^V \mathcal{S}_{L1}^V)^2$. With this correction (D.16) of ref. [16] coincides with (A.14) in the current paper.

- [1] Y. Hosotani, “*Dynamical mass generation by compact extra dimensions*”, *Phys. Lett. B***126**, 309 (1983).
- [2] Y. Hosotani, “*Dynamics of nonintegrable phases and gauge symmetry breaking*”, *Ann. Phys. (N.Y.)* **190**, 233 (1989).
- [3] A. T. Davies and A. McLachlan, “*Gauge group breaking by Wilson loops*”, *Phys. Lett. B***200**, 305 (1988); “*Congruency class effects in the Hosotani model*”, *Nucl. Phys. B***317**, 237 (1989).
- [4] H. Hatanaka, T. Inami, C.S. Lim, “*The gauge hierarchy problem and higher dimensional gauge theories*”, *Mod. Phys. Lett. A***13**, 2601 (1998).
- [5] H. Hatanaka, “*Matter representations and gauge symmetry breaking via compactified space*”, *Prog. Theoret. Phys.* **102**, 407 (1999).
- [6] M. Kubo, C.S. Lim and H. Yamashita, “*The Hosotani mechanism in bulk gauge theories with an orbifold extra space S^1/Z_2* ”, *Mod. Phys. Lett. A***17**, 2249 (2002).
- [7] A. Pomarol and M. Quiros, “*The Standard Model from extra dimensions*”, *Phys. Lett. B***438**, 255 (1998).
- [8] C.A. Scrucca, M. Serone, L. Silvestrini, “*Electroweak symmetry breaking and fermion masses from extra dimensions*”, *Nucl. Phys. B***669**, 128 (2003).
- [9] K. Agashe, R. Contino and A. Pomarol, “*The minimal composite Higgs model*”, *Nucl. Phys. B***719**, 165 (2005).
- [10] G. Cacciapaglia, C. Csaki, S.C. Park, “*Fully radiative electroweak symmetry breaking*”, *JHEP* **0603**, 099 (2006).
- [11] A. D. Medina, N. R. Shah and C. E. M. Wagner, “*Gauge-Higgs unification and radiative electroweak symmetry breaking in warped extra dimensions*”, *Phys. Rev. D***76**, 095010 (2007).
- [12] Y. Hosotani, K. Oda, T. Ohnuma and Y. Sakamura, “*Dynamical electroweak symmetry breaking in $SO(5) \times U(1)$ gauge-Higgs unification with top and bottom quarks*”, *Phys. Rev. D***78**, 096002 (2008); *Erratum-ibid.* **D79**, 079902 (2009).
- [13] S. Funatsu, H. Hatanaka, Y. Hosotani, Y. Orikasa and T. Shimotani, “*Novel universality and Higgs decay $H \rightarrow \gamma\gamma, gg$ in the $SO(5) \times U(1)$ gauge-Higgs unification*”, *Phys. Lett. B***722**, 94 (2013).
- [14] S. Funatsu, H. Hatanaka, Y. Hosotani and Y. Orikasa, “*Distinct signals of the gauge-Higgs unification in e^+e^- collider experiments*”, *Phys. Lett. B***775**, 297 (2017).
- [15] Y. Adachi and N. Maru, “*Revisiting electroweak symmetry breaking and the Higgs boson mass in gauge-Higgs unification*”, *Phys. Rev. D***98**, 015022 (2018).

- [16] S. Funatsu, H. Hatanaka, Y. Hosotani, Y. Orikasa and N. Yamatsu, “*GUT inspired $SO(5) \times U(1) \times SU(3)$ gauge-Higgs unification*”, *Phys. Rev. D* **99**, 095010 (2019).
- [17] J. Yoon and M.E. Peskin, “*Fermion pair production in $SO(5) \times U(1)$ gauge-Higgs unification models*”, arXiv:1811.07877 [hep-ph].
- [18] J. Yoon and M.E. Peskin, “*Dissection of an $SO(5) \times U(1)$ gauge-Higgs unification model*”, *Phys. Rev. D* **100**, 015001 (2019).
- [19] S. Funatsu, “*Forward-backward asymmetry in the gauge-Higgs unification at the International Linear Collider*”, *Eur. Phys. J. C* **79**, 854 (2019).
- [20] S. Funatsu, H. Hatanaka, Y. Hosotani, Y. Orikasa and N. Yamatsu, “*CKM matrix and FCNC suppression in $SO(5) \times U(1) \times SU(3)$ gauge-Higgs unification*”, arXiv:1909.00190 [hep-ph], to appear in *Phys. Rev. D*.
- [21] S. Bilokin, R. Pöschl and F. Richard, “*Measurement of b quark EW couplings at ILC*”, arXiv:1709.04289 [hep-ex]; F. Richard, “*Bhabha scattering at ILC250*”, arXiv:1804.02846 [hep-ex]; A. Irles, R. Pöschl, F. Richard and H. Yamamoto, “*Complementarity between ILC250 and ILC-GigaZ*”, arXiv:1905.00220 [hep-ex]; A. Irles, R. Pöschl and F. Richard, “*Production and measurement of $e^+e^- \rightarrow c\bar{c}$ signatures at the 250 GeV ILC*”, arXiv:2002.05805 [hep-ex].
- [22] K. Fujii et al., “*Physics case for the 250 GeV stage of the International Linear Collider*”, arXiv:1710.07621 [hep-ex]; ILC Collaboration (H. Aihara et al.), “*The International Linear Collider. A Global Project*”, arXiv:1901.09829 [hep-ex]; Philip Bambade et al., “*The International Linear Collider. A Global Project*”, arXiv:1903.01629 [hep-ex];
- [23] G. Burdman and Y. Nomura, “*Unification of Higgs and gauge fields in five dimensions*”, *Nucl. Phys. B* **656**, 3 (2003); N. Haba, Masatomi Harada, Y. Hosotani and Y. Kawamura, “*Dynamical rearrangement of gauge symmetry on the orbifold S^1/Z_2* ”, *Nucl. Phys. B* **657**, 169 (2003); *Erratum-ibid.* **669**, 381 (2003); N. Haba, Y. Hosotani, Y. Kawamura and T. Yamashita, “*Dynamical symmetry breaking in gauge Higgs unification on orbifold*”, *Phys. Rev. D* **70**, 015010 (2004); C.S. Lim and N. Maru, “*Towards a realistic grand gauge-Higgs unification*”, *Phys. Lett. B* **653**, 320 (2007); K. Kojima, K. Takenaga and T. Yamashita, “*Grand gauge-Higgs unification*”, *Phys. Rev. D* **84**, 051701(R) (2011); “*Gauge symmetry breaking patterns in an $SU(5)$ grand gauge-Higgs unification*”, *Phys. Rev. D* **95**, 015021 (2017); M. Frigerio, J. Serra and A. Varagnolo, “*Composite GUTs: Models and expectations at the LHC*”, *JHEP* **1106**, 029 (2011); M. Kakizaki, S. Kanemura, H. Taniguchi and T. Yamashita, “*Higgs sector as a probe of supersymmetric grand unification with*

- the Hosotani mechanism”, *Phys. Rev. D* **89**, 075013 (2014); N. Maru and Y. Yatagai, “Fermion mass hierarchy in grand gauge-Higgs unification”, arXiv:1903.08359 [hep-ph];
- [24] Y. Hosotani and N. Yamatsu, “Gauge-Higgs grand unification”, *Prog. Theoret. Exp. Phys.* **2015**, 111B01 (2015).
 - [25] A. Furui, Y. Hosotani, and N. Yamatsu, “Toward realistic gauge-Higgs grand unification”, *Prog. Theoret. Exp. Phys.* **2016**, 093B01 (2016); Y. Hosotani and N. Yamatsu, “Electroweak symmetry breaking and mass spectra in six-dimensional gauge-Higgs grand unification”, *Prog. Theoret. Exp. Phys.* **2018**, 023B05 (2018).
 - [26] Y. Hosotani and N. Yamatsu, “Gauge-Higgs seesaw mechanism in 6-dimensional grand unification”, *Prog. Theoret. Exp. Phys.* **2017**, 091B01 (2017).
 - [27] C. Englert, D.J. Miller and D.D. Smaranda, “Phenomenology of GUT-inspired gauge-Higgs unification”, *Phys. Lett. B* **802**, 135261 (2020).
 - [28] Y. Adachi and N. Maru, “Strong first order electroweak phase transition in gauge-Higgs unification at finite temperature”, arXiv:1911.05963 [hep-ph].
 - [29] G. F. Giudicea, C. Grojean, A. Pomarol, and R. Rattazzi, “The strongly-interacting light Higgs”, *JHEP* **0706**, 045 (2007); R. Contino, C. Grojean, M. Moretti, F. Piccinini, and R. Rattazzi, “Strong double Higgs production at the LHC”, *JHEP* **1005**, 089 (2010); R. Contino, M. Ghezzi, M. Moretti, G. Panico, F. Piccinini, and A. Wulzer, “Anomalous couplings in double Higgs production”, *JHEP* **1208**, 154 (2012); R. Contino, M. Ghezzi, C. Grojean, M. Muhlleitner, and M. Spira, “Effective Lagrangian for a light Higgs-like scalar”, *JHEP* **1307**, 035 (2013).
 - [30] G. Cacciapaglia, C. Pica and F. Sannino, “Fundamental Composite Dynamics: A Review”, arXiv:2002.04914 [hep-ph]; R. Contino, “The Higgs as a Composite Nambu-Goldstone Boson”, TASI 2009, 235; arXiv:1005.4269 [hep-ph]; B. Bellazzini, C. Csaki, and J. Serra, “Composite Higgses”, *Eur. Phys. J. C* **74**, 2766 (2014).
 - [31] S. Funatsu, H. Hatanaka, Y. Hosotani, Y. Orikasa, and T. Shimotani, “LHC signals of the $SO(5) \times U(1)$ gauge-Higgs unification”, *Phys. Rev. D* **89**, 095019 (2014).; S. Funatsu, H. Hatanaka, Y. Hosotani and Y. Orikasa, “Collider signals of W' and Z' bosons in the gauge-Higgs unification”, *Phys. Rev. D* **95**, 035032 (2017).
 - [32] L. Randall and R. Sundrum, “A large mass hierarchy from a small extra dimension”, *Phys. Rev. Lett.* **83**, 3370 (1999).
 - [33] T. Gherghetta and A. Pomarol, “Bulk fields and supersymmetry in a slice of AdS”, *Nucl. Phys. B* **586**, 141 (2000).

- [34] M. Tanabashi et al. (Particle Data Group), “*Review of Particle Physics*”, *Phys. Rev. D* **98**, 030001 (2018), Chap. 10. “*Electroweak Model and Constraints on New Physics*”.
- [35] A. Falkowski, “*Holographic pseudo-Goldstone boson*”, *Phys. Rev. D* **75**, 025017 (2007).
- [36] Y. Hosotani, M. Tanaka and N. Uekusa, “*H parity and the stable Higgs boson in the $SO(5) \times U(1)$ gauge-Higgs unification*”, *Phys. Rev. D* **82**, 115024 (2010).
- [37] Y. Hosotani and Y. Kobayashi, “*Yukawa couplings and effective interactions in gauge-Higgs unification*”, *Phys. Lett. B* **674**, 192 (2009).
- [38] Y. Hosotani, “*Gauge-Higgs Unification – Beyond the Standard Model*” (in Japanese), in SGC Library -143, SAIENSU-SHA Co., Ltd, Tokyo (2018), Section 8.1.