

Global Convergence and Variance-Reduced Optimization for a Class of Nonconvex-Nonconcave Minimax Problems

Junchi Yang *

Negar Kiyavash †

Niao He *

September 9, 2021

Abstract

Nonconvex minimax problems appear frequently in emerging machine learning applications, such as generative adversarial networks and adversarial learning. Simple algorithms such as the gradient descent ascent (GDA) are the common practice for solving these nonconvex games and receive lots of empirical success. Yet, it is known that these vanilla GDA algorithms with constant step size can potentially diverge even in the convex setting. In this work, we show that for a subclass of nonconvex-nonconcave objectives satisfying a so-called two-sided Polyak-Lojasiewicz inequality, the alternating gradient descent ascent (AGDA) algorithm converges globally at a linear rate and the stochastic AGDA achieves a sublinear rate. We further develop a variance reduced algorithm that attains a provably faster rate than AGDA when the problem has the finite-sum structure.

1 Introduction

We consider minimax optimization problems of the forms

$$\min_{x \in \mathbb{R}^{d_1}} \max_{y \in \mathbb{R}^{d_2}} f(x, y) \triangleq \mathbb{E}[F(x, y; \xi)], \quad (1)$$

and

$$\min_{x \in \mathbb{R}^{d_1}} \max_{y \in \mathbb{R}^{d_2}} f(x, y) \triangleq \frac{1}{n} \sum_{i=1}^n f_i(x, y), \quad (2)$$

where ξ is a random vector with support Ξ , and $f(x, y)$ is a possibly nonconvex-nonconcave function. Minimax problems have been widely studied in game theory and operations research. Recent emerging applications in machine learning have further stimulated a surge of interest in these problems. For example, generative adversarial networks (GANs) [Goodfellow et al., 2016] can be viewed as a two-player game between a generator that produces synthetic data and a discriminator that differentiates between true data and synthetic data. In reinforcement learning, solving Bellman equations can also be reformulated as minimax optimization problems [Chen and Wang, 2016, Dai et al., 2017, 2018]. Other applications include robust optimization [Namkoong and Duchi, 2016, 2017], adversarial machine learning [Sinha et al., 2017, Madry et al., 2017], unsupervised learning [Xu et al., 2005], and so on.

*Department of Industrial and Enterprise Systems Engineering, University of Illinois at Urbana-Champaign, IL 61801, USA (junchiy2@illinois.edu, niaoh@illinois.edu).

†School of Management of Technology, École Polytechnique Fédérale de Lausanne, Switzerland (negar.kiyavash@epfl.ch).

The most natural and frequently used methods for solving minimax problems (1) and (2) are the gradient descent ascent (GDA) algorithms (or their stochastic variants), with either simultaneous or alternating updates of the primal-dual variables, referred to as SGDA and AGDA, respectively, throughout the paper. While these algorithms have received much empirical success especially in adversarial training, it is known that these GDA algorithms with constant stepsizes could fail to converge for general smooth function [Mescheder et al., 2018], even for the bilinear games [Gidel et al., 2019]; even when they do converge, the stable limit point may not be a local Nash equilibrium [Daskalakis et al., 2018, Mazumdar and Ratliff, 2018]. On the other hand, GDA algorithms can converge linearly to the saddle point for strongly-convex-strongly-concave functions [Facchinei and Pang, 2007]. Moreover, for many simple nonconvex-nonconcave objective functions, such as, $f(x, y) = x^2 + 3 \sin^2 x \sin^2 y - 4y^2 - 10 \sin^2 y$, we also observe that GDA algorithms with constant stepsizes indeed converge to the global Nash equilibrium (or saddle point), at a linear rate (see Figure 1). This also holds true for their stochastic variants, albeit at a sublinear rate. These facts naturally raise a question: *Is there a general condition under which GDA algorithms converge to the global optima?*




Figure 1: (a) Surface plot of the nonconvex-nonconcave function $f(x, y) = x^2 + 3 \sin^2 x \sin^2 y - 4y^2 - 10 \sin^2 y$; (b) Convergence of SGDA and AGDA; (c) Convergence of stochastic SGDA and stochastic AGDA; (d) Trajectories of four algorithms

Furthermore, the use of variance reduction techniques has played a prominent role in improving the convergence over stochastic or batch algorithms for both convex and nonconvex minimization problems, which have been extensively studied in the past few years; see, e.g., [Johnson and Zhang, 2013, Reddi et al., 2016a,b, Xiao and Zhang, 2014], just to name a few. However, when it comes to the minimax problems, there are limited results, except under convex-concave setting [Palaniappan and Bach, 2016, Du and Hu, 2019]. This leads to another open question: *Can we improve GDA algorithms for nonconvex-nonconcave minimax problems?*

1.1 Our contributions

In this paper, we address these two questions and specifically focus on the alternating gradient descent ascent, namely AGDA. This is due to several considerations. First of all, it has been recently shown that alternating updates of GDA are more stable than simultaneous updates [Gidel et al., 2019, Bailey et al., 2019]. Note that for a convex-concave matrix game, SGDA may diverge while AGDA is proven to always have bounded iterates [Gidel et al., 2019]. See Figure 2 for a simple illustration. Secondly, in general, it is much more challenging to analyze AGDA than SGDA. There is a lack of discussion on the convergence of AGDA for general minimax problems in the literature. Our contributions are summarized as follows.

First, we identify a general condition that relaxes the convex-concavity requirement of the objective function while still guaranteeing global convergence of AGDA and stochastic AGDA (Stoc-AGDA). We call this the two-sided PL condition, which requires that both players' utility functions satisfy Polyak-Łojasiewicz




Figure 2: Consider the objective $f(x, y) = \log(1 + e^x) + 3xy - \log(1 + e^y)$: (a) Convergence of AGDA and SGDA under the stepsize $\tau = 0.01$; (b) Trajectories of two algorithms under the stepsize $\tau = 0.01$; (c) Convergence of AGDA and SGDA under stepsize $\tau = 0.025$; (d) Trajectories of two algorithms with stepsize $\tau = 0.025$;

(PL) inequality [Polyak, 1963]. Such conditions indeed hold true for several applications, including robust least square, generative adversarial imitation learning for linear quadratic regulator (LQR) dynamics [Cai et al., 2019], and potentially many others in adversarial learning [Du et al., 2019], robust phase retrieval [Sun et al., 2018, Zhou et al., 2016], robust control [Fazel et al., 2018], and etc. We show that under the two-sided PL condition, AGDA with proper constant stepsizes converges globally to a saddle point at a linear rate of $\mathcal{O}(1 - \kappa^{-3})^t$, while Stoc-AGDA with proper diminishing stepsizes converges to a saddle point at a sublinear rate of $\mathcal{O}(\kappa^5/t)$, where κ is the underlying condition number.

Second, for minimax problems with the finite sum structure, we introduce a variance-reduced AGDA algorithm (VR-AGDA) that leverages the idea of stochastic variance reduced gradient (SVRG) [Johnson and Zhang, 2013, Reddi et al., 2016a] with the alternating updates. We prove that VR-AGDA achieves the complexity of $\mathcal{O}((n^{2/3}\kappa^3 \log(1/\epsilon)))$ in the region $n \leq \kappa^9$ and $\mathcal{O}(n + \kappa^9) \log(1/\epsilon)$ in the region $n \geq \kappa^9$, where n is the number of component functions. This greatly improves over the $\mathcal{O}(n\kappa^3 \log \frac{1}{\epsilon})$ complexity of AGDA when applied to the finite sum minimax problems. We summarize the results of these algorithms in Table 1. Our numerical experiments further demonstrate that VR-AGDA performs significantly better than AGDA and Stoc-AGDA, especially for problems with large condition numbers. To our best knowledge, this is the first work to provide a variance reduced algorithm and theoretical guarantees in the nonconvex-nonconcave regime of minimax optimization.

Algorithms	AGDA	Stoc-AGDA	VR-AGDA
Complexity	$\mathcal{O}(n\kappa^3 \log \frac{1}{\epsilon})$	$\mathcal{O}\left(\frac{\kappa^5}{\mu_2 \epsilon}\right)$	$\mathcal{O}\left(n^{\frac{2}{3}}\kappa^3 \log \frac{1}{\epsilon}\right), n \leq \kappa^9$ $\mathcal{O}((n + \kappa^9) \log \frac{1}{\epsilon})$

Table 1: Complexities of three algorithms for the finite-sum problem (2), where $\kappa \triangleq l/\mu_1$ is condition number, l is Lipschitz gradient constant, μ_1 and μ_2 are the two-side PL constants with $\mu_1 \leq \mu_2$. See Section 4 for more details.

1.2 Related work

Nonconvex minimax problems. There has been a recent surge in research on solving minimax optimization beyond the convex-concave regime [Sinha et al., 2017, Chen et al., 2017, Qian et al., 2019, Thekumparampil et al., 2019, Lin et al., 2018, Nouiehed et al., 2019, Abernethy et al., 2019], but they differ from our work from various perspectives. For example, Chen et al. [2017], Sinha et al. [2017], Lin et al. [2019], Thekumparampil et al. [2019] considered the minimax problem when the objective function is nonconvex in x

but concave in y and focused on achieving convergence to stationary points. Their algorithms require solving the inner maximization or some sub-problems with high accuracy at every iteration, which are different from AGDA. Lin et al. [2018] considered a general class of weakly-convex weakly-concave minimax problems and proposed an inexact proximal point method to find an ϵ -stationary point. Their convergence result relies on assuming the existence of a solution to the corresponding Minty variational inequality, which is often hard to verify. Abernethy et al. [2019] recently showed the linear convergence of a second-order iterative algorithm, called Hamiltonian gradient descent (HGD), for a subclass of “sufficiently bilinear” functions. Compared with their work, the PL condition we consider in this paper is easier to verify and GDA algorithms are much simpler.

PL condition. Recently, Nouiehed et al. [2019] studied a class of minimax problems where the objective only satisfies a one-sided PL condition and introduced the GDmax algorithm, which takes multiple ascent steps at every iteration. Our work differs from [Nouiehed et al., 2019] in two aspects: (i) we consider the two-sided PL condition which guarantees global convergence¹; (ii) we consider AGDA which takes one ascent step at every iteration. Another closely related work is Cai et al. [2019]. The authors considered a specific application in generative adversarial imitation learning with linear quadratic regulator dynamics. This is a special example that falls under the two-sided PL condition.

Variance-reduced minimax optimization. There exists a few works that apply variance reduction techniques to minimax optimization. Palaniappan and Bach [2016], Luo et al. [2019] provided linear-convergent algorithms for strongly-convex-strongly-concave objectives, based on simultaneous updates. Du and Hu [2019] extended the result to convex-strongly-concave objectives with full-rank coupling bilinear term. In contrast, we are dealing with a much broader class of objectives that are possibly nonconvex-nonconcave. We point out that Luo et al. [2020] recently introduced a variance-reduced algorithm for finding the stationary point of nonconvex-strongly-concave problems, which is again different from our setting.

The rest of this paper is organized as follows. In Section 2, we introduce the two-sided PL condition and show the equivalence of three min-max optimality criteria under this condition. In Section 3, we describe deterministic and stochastic AGDA algorithms, and provide convergence analyses of those algorithms under the two-sided PL condition. In Section 4, we introduce the variance-reduced AGDA algorithm and establish its convergence results. In Section 5, we provide numerical performance of these algorithms for robust least square and imitation learning for LQR.

2 Global optima and two-sided PL condition

Throughout this paper, we assume that the function $f(x, y)$ in (1) is continuously differentiable and has Lipschitz gradient. We state it as a basic assumption. Here $\|\cdot\|$ is used to denote the Euclidean norm.

Assumption 1 (Lipschitz gradient). *There exists a positive constant $l > 0$ such that*

$$\begin{aligned}\|\nabla_x f(x_1, y_1) - \nabla_x f(x_2, y_2)\| &\leq l[\|x_1 - x_2\| + \|y_1 - y_2\|], \\ \|\nabla_y f(x_1, y_1) - \nabla_y f(x_2, y_2)\| &\leq l[\|x_1 - x_2\| + \|y_1 - y_2\|],\end{aligned}$$

holds for all $x_1, x_2 \in \mathbb{R}^{d_1}, y_1, y_2 \in \mathbb{R}^{d_2}$.

¹We also show that AGDA can find ϵ -stationary point for minimax problems under the one-sided PL condition within $\mathcal{O}(1/\epsilon^2)$ iterations in Appendix D.

We now define three notions of optimality for minimax problems. The most direct notion of optimality is global minimax point, at which x^* is an optimal solution to the function $g(x) := \max_y f(x, y)$ and y^* is an optimal solution to $\max_y f(x^*, y)$. In the two-player zero-sum game, the notion of saddle point is also widely used [Von Neumann et al., 2007, Nash, 1953]. For a saddle point (x^*, y^*) , x^* is an optimal solution to $\min_x f(x, y^*)$ and y^* is an optimal solution to $\max_y f(x^*, y)$.

Definition 1 (Global optima).

1. (x^*, y^*) is a global minimax point, if for any (x, y) :

$$f(x^*, y) \leq f(x^*, y^*) \leq \max_{y'} f(x, y'). \quad (3)$$

2. (x^*, y^*) is a saddle point, if for any (x, y) :

$$f(x^*, y) \leq f(x^*, y^*) \leq f(x, y^*). \quad (4)$$

3. (x^*, y^*) is a stationary point, if :

$$\nabla_x f(x^*, y^*) = \nabla_y f(x^*, y^*) = 0. \quad (5)$$

For general nonconvex-nonconcave minimax problems, these three notions of optimality are not necessarily equivalent. A stationary point may not be a saddle point or a global minimax point; a global minimax point may not be a saddle point or a stationary point. Note that generally speaking, for minimax problems, a saddle point or a global minimax point may not always exist. However, since our goal in this paper is to find global optima, in the remainder of the paper, we assume that a saddle point always exists.

Assumption 2 (Existence of saddle point). *The objective function f has at least one saddle point. We also assume that for any fixed y , $\min_{x \in \mathbb{R}^{d_1}} f(x, y)$ has a nonempty solution set and a optimal value, and for any fixed x , $\max_{y \in \mathbb{R}^{d_2}} f(x, y)$ has a nonempty solution set and a finite optimal value.*

For unconstrained minimization problems: $\min_{x \in \mathbb{R}^n} f(x)$, Polyak [1963] proposed Polyak-Łojasiewicz (PL) condition, which is sufficient to show global linear convergence for gradient descent without assuming convexity. Specifically, a function $f(\cdot)$ satisfies PL condition if it has a nonempty solution set and a finite optimal value f^* , and there exists some $\mu > 0$ such that $\frac{1}{2} \|\nabla f(x)\|^2 \geq \mu(f(x) - f^*)$, $\forall x$. As discussed in Karimi et al. [2016], PL condition is weaker, or not stronger, than other well-known conditions that guarantee linear convergence for gradient descent, such as error bounds (EB) [Luo and Tseng, 1993], weak strong convexity (WSC) [Necoara et al., 2018] and restricted secant inequality (RSI) [Zhang and Yin, 2013].

We introduce a straightforward generalization of the PL condition to the minimax problem: function $f(x, y)$ satisfies the PL condition with constant μ_1 with respect to x , and $-f$ satisfies PL condition with constant μ_2 with respect to y . We formally state this in the following definition.

Definition 2 (Two-sided PL condition). *A continuously differentiable function $f(x, y)$ satisfies the two-sided PL condition if there exist constants $\mu_1, \mu_2 > 0$ such that:*

$$\begin{aligned} \|\nabla_x f(x, y)\|^2 &\geq 2\mu_1[f(x, y) - \min_x f(x, y)], \forall x, y, \\ \|\nabla_y f(x, y)\|^2 &\geq 2\mu_2[\max_y f(x, y) - f(x, y)], \forall x, y. \end{aligned}$$

The two-sided PL condition does not imply convexity-concavity, and it is a much weaker condition than strong-convexity-strong-concavity. In Lemma 2.1, we show that three notions of optimality are equivalent under the two-sided PL condition. Note that they may not be unique.

Lemma 2.1. *If the objective function $f(x, y)$ satisfies the two-sided PL condition, then the following holds true:*

$$(\text{saddle point}) \Leftrightarrow (\text{global minimax}) \Leftrightarrow (\text{stationary point}).$$

Below we give some examples that satisfy this condition.

Example 1. *The nonconvex-nonconcave function in the introduction, $f(x, y) = x^2 + 3 \sin^2 x \sin^2 y - 4y^2 - 10 \sin^2 y$ satisfies the two-sided PL condition with $\mu_1 = 1/16, \mu_2 = 1/11$ (see Appendix A).*

Example 2. *$f(x, y) = F(Ax, By)$, where $F(\cdot, \cdot)$ is strongly-convex-strongly-concave and A and B are arbitrary matrices, satisfies the two-sided PL condition.*

Example 3. *The generative adversarial imitation learning for LQR can be formulated as $\min_K \min_\theta m(K, \theta)$, where m is strongly-concave in terms of θ and satisfies PL condition in terms of K (see [Cai et al., 2019] for more details), thus satisfying the two-sided PL condition.*

Under the two-sided PL condition, the function $g(x) := \max_y f(x, y)$ can be shown to satisfy PL condition with μ_1 (see Appendix A). Moreover, it holds that g is also L -smooth with $L := l + l^2/\mu_2$ [Nouiehed et al., 2019]. Finally, we denote $\mu = \min(\mu_1, \mu_2)$ and $\kappa = \frac{L}{\mu}$, which represents the condition number of the problem.

3 Global convergence of AGDA and Stoc-AGDA

In this section, we establish the convergence rate of the stochastic alternating gradient descent ascent (Stoc-AGDA) algorithm, which we present in Algorithm 1, under the two-sided PL condition. Stoc-AGDA updates variables x and y sequentially using stochastic gradient descent/ascent steps. Here we make standard assumptions about stochastic gradients $G_x(x, y, \xi)$ and $G_y(x, y, \xi)$.

Assumption 3 (Bounded variance). *$G_x(x, y, \xi)$ and $G_y(x, y, \xi)$ are unbiased stochastic estimators of $\nabla_x f(x, y)$ and $\nabla_y f(x, y)$ and have variances bounded by $\sigma^2 > 0$.*

Algorithm 1 Stoc-AGDA

```

1: Input:  $(x_0, y_0)$ , step sizes  $\{\tau_1^t\}_t > 0, \{\tau_2^t\}_t > 0$ 
2: for all  $t = 0, 1, 2, \dots$  do
3:   Draw two i.i.d. samples  $\xi_{t1}, \xi_{t2} \sim P(\xi)$ 
4:    $x_{t+1} \leftarrow x_t - \tau_1^t G_x(x_t, y_t, \xi_{t1})$ 
5:    $y_{t+1} \leftarrow y_t + \tau_2^t G_y(x_{t+1}, y_t, \xi_{t2})$ 
6: end for
```

Note that Stoc-AGDA with constant stepsizes (i.e., $\tau_1^t = \tau_1$ and $\tau_2^t = \tau_2$) and noiseless stochastic gradient (i.e., $\sigma^2 = 0$) reduces to AGDA:

$$x_{t+1} = x_t - \tau_1 \nabla_x f(x_t, y_t), \quad (6)$$

$$y_{t+1} = y_t - \tau_2 \nabla_y f(x_{t+1}, y_t). \quad (7)$$

We will measure the inaccuracy of (x_t, y_t) through the potential function

$$P_t := a_t + \lambda \cdot b_t, \quad (8)$$

where $a_t = \mathbb{E}[g(x_t) - g^*]$, $b_t = \mathbb{E}[g(x_t) - f(x_t, y_t)]$ and $\lambda > 0$ to be specified later in the theorems. Recall that $g(x) := \max_y f(x, y)$ and $g^* = \min_x g(x)$. This metric is driven by the definition of minimax point, because $g(x) - g^*$ and $g(x) - f(x, y)$ are non-negative for any (x, y) , and both equal to 0 if and only if (x, y) is a minimax point.

Stoc-AGDA with constant stepsizes We first consider Stoc-AGDA with constant stepsizes. We show that $\{(x_t, y_t)\}_t$ will converge linearly to a neighbourhood of the optimal set.

Theorem 3.1. *Suppose Assumptions 1, 2, 3 hold and $f(x, y)$ satisfies the two-sided PL condition with μ_1 and μ_2 . Define $P_t := a_t + \frac{1}{10}b_t$. If we run Algorithm 1 with $\tau_2^t = \tau_2 \leq \frac{1}{l}$ and $\tau_1^t = \tau_1 \leq \frac{\mu_2^2 \tau_2}{18l^2}$, then*

$$P_t \leq (1 - \frac{1}{2}\mu_1\tau_1)^t P_0 + \delta, \quad (9)$$

where $\delta = \frac{(1-\mu_2\tau_2)(L+l)\tau_1^2 + l\tau_2^2 + 10L\tau_1^2}{10\mu_1\tau_1}\sigma^2$.

Remark 1. In the theorem above, we choose τ_1 smaller than τ_2 , $\tau_1/\tau_2 \leq \mu_2^2/(18l^2)$, because our potential function is not symmetric about x and y . Another reason is because we want y_t to approach $y^*(x_t) \in \arg \max_y f(x_t, y)$ faster so that $\nabla_x f(x_t, y_t)$ is a better approximation for $\nabla g(x_t)$ ($\nabla g(x) = \nabla_x f(x, y^*(x))$, see Nowieh et al. [2019]). Indeed, it is common to use different learning rates for x and y in GDA algorithms for nonconvex minimax problems; see e.g., Jin et al. [2019] and Lin et al. [2019]. Note that the ratio between these two learning rates is quite crucial here. We also observe empirically when the same learning rate is used, even if small, the algorithm may not converge to saddle points.

Remark 2. When $t \rightarrow \infty$, $P_t \rightarrow \delta$. If $\tau_1 \rightarrow 0$ and $\tau_2^2/\tau_1 \rightarrow 0$, the error term δ will go to 0. When using smaller stepsizes, the algorithm reaches a smaller neighbour of the saddle point yet at the cost of a slower rate, as the contraction factor also deteriorates.

Linear convergence of AGDA Setting $\sigma^2 = 0$, it follows immediately from the previous theorem that AGDA converges linearly under the two-sided PL condition. Moreover, we have

Theorem 3.2. *Suppose Assumptions 1, 2 hold and $f(x, y)$ satisfies the two-sided PL condition with μ_1 and μ_2 . Define $P_t := a_t + \frac{1}{10}b_t$. If we run AGDA with $\tau_1 = \frac{\mu_2^2}{18l^3}$ and $\tau_2 = \frac{1}{l}$, then*

$$P_t \leq \left(1 - \frac{\mu_1\mu_2^2}{36l^3}\right)^t P_0. \quad (10)$$

Furthermore, $\{(x_t, y_t)\}_t$ converges to some saddle point (x^*, y^*) , and

$$\|x_t - x^*\|^2 + \|y_t - y^*\|^2 \leq \alpha \left(1 - \frac{\mu_1\mu_2^2}{36l^3}\right)^t P_0, \quad (11)$$

where α is a constant depending on μ_1, μ_2 and l .

The above theorem implies that the limit point of $\{(x_t, y_t)\}_t$ is a saddle point and the distance to the saddle point decreases in the order of $\mathcal{O}((1 - \kappa^{-3})^t)$. Note that in the special case when the objective is strongly-convex-strongly-concave, it is known that SGDA (GDA with simultaneous updates) achieves an $\mathcal{O}(\kappa^2 \log(1/\epsilon))$ iteration complexity (see, e.g., Facchinei and Pang [2007]) and this can be further improved to $\mathcal{O}(\kappa \log(1/\epsilon))$ by extragradient methods [Korpelevich, 1976], Nesterov's dual extrapolation [Nesterov and Scramali, 2006] or accelerated proximal point algorithm [Lin et al., 2020]. However, these result relies heavily on the strong monotonicity of the corresponding variational inequality. For the general two-sided PL condition, we may not achieve the same dependency on κ .

Stoc-AGDA with diminishing stepsizes While Stoc-AGDA with constant stepsizes only converges linearly to a neighbourhood of the saddle point, Stoc-AGDA with diminishing stepsizes converges to the saddle point but at a sublinear rate $\mathcal{O}(1/t)$.

Theorem 3.3. Suppose Assumptions 1, 2, 3 hold and $f(x, y)$ satisfies the two-sided PL condition with μ_1 and μ_2 . Define $P_t = a_t + \frac{1}{10}b_t$. If we run algorithm 1 with stepsizes $\tau_1^t = \frac{\beta}{\gamma+t}$ and $\tau_2^t = \frac{18l^2\beta}{\mu_2^2(\gamma+t)}$ for some $\beta > 2/\mu_1$ and $\gamma > 0$ such that $\tau_1^1 \leq \min\{1/L, \mu_2^2/18l^2\}$, then we have

$$P_t \leq \frac{\nu}{\gamma+t}, \quad (12)$$

where $\nu :=$

$$\max\left\{\gamma P_0, \frac{[(L+l)\beta^2 + 18^2 l^5 \beta^2 / \mu_2^4 + 10L\beta^2]\sigma^2}{10\mu_1\beta - 20}\right\}.$$

Remark 3. Note the rate is affected by ν , and the first term in the definition of ν is controlled by the initial point. In practice, we can find a good initial point by running Stoc-AGDA with constant stepsizes so that only the second term in the definition of ν matters. Then by choosing $\beta = 3/\mu_1$, we have $\nu = \mathcal{O}\left(\frac{l^5\sigma^2}{\mu_1^2\mu_2^4}\right)$. Thus, the convergence rate of Stoc-AGDA is $\mathcal{O}\left(\frac{\kappa^5\sigma^2}{\mu t}\right)$.

4 Stochastic variance reduced algorithm

In this section, we study the minimax problem in (2) with the finite-sum structure:

$$\min_x \max_y f(x, y) = \frac{1}{n} \sum_{i=1}^n f_i(x, y),$$

which arises ubiquitously in machine learning. We are especially interested in the case when n is large. We assume the overall objective function $f(x, y)$ still satisfies the two-sided PL condition with μ_1 and μ_2 , but we do not assume each f_i to satisfy the two-sided PL condition. Instead of Assumption 1, we now assume each component f_i has Lipschitz gradients.

Assumption 4. Each f_i has l -Lipschitz gradients.

If we run AGDA with full gradients to solve the finite-sum minimax problem, the total complexity for finding an ϵ -optimal solution is $\mathcal{O}(n\kappa^3 \log(1/\epsilon))$ by Theorem 3.2. Despite the linear convergence, the per-iteration cost is high and the complexity can be huge when the number of components n and condition number κ are large. Instead, if we run Stoc-AGDA, this leads to the total complexity $\mathcal{O}\left(\frac{\kappa^5\sigma^2}{\mu_2\epsilon}\right)$ by Remark 3, which has worse dependence on ϵ .

Motivated by the recent success of stochastic variance reduced gradient (SVRG) technique [Johnson and Zhang, 2013, Reddi et al., 2016a, Palaniappan and Bach, 2016], we introduce the VR-AGDA algorithm (presented in Algorithm 2), that combines AGDA with SVRG so that the linear convergence is preserved while improving the dependency on n and κ . VR-AGDA can be viewed as the applying SVRG to AGDA with restarting: at every epoch k , we restart the SVRG subroutine (with T outer iterations, N inner steps) by initializing it with (x^k, y^k) , which is randomly selected from previous SVRG subroutine. This is partly inspired by the GD-SVRG algorithm for minimizing PL functions [Reddi et al., 2016a]. Notice when $T = 1$, VR-AGDA reduces to a double-loop algorithm which is similar to the SVRG for saddle point problems proposed by Palaniappan and Bach [2016], except for several notable differences: (i) we are using the alternating updates rather than simultaneous updates, (ii) as a result, we require to sample two independent indices rather than one at each iteration, and (iii) most importantly, we are dealing with possibly nonconvex-nonconcave objectives that satisfy the two-sided PL condition.

The following two theorems capture the convergence of VR-AGDA.

Algorithm 2 VR-AGDA

```
1: input:  $(\tilde{x}_0, \tilde{y}_0)$ , stepsizes  $\tau_1, \tau_2$ , iteration numbers  $N, T$ 
2: for all  $k = 0, 1, 2, \dots$  do
3:   for all  $t = 0, 1, 2, \dots, T - 1$  do
4:      $x_{t,0} = \tilde{x}_t, \quad y_{t,0} = \tilde{y}_t,$ 
5:     compute  $\nabla_x f(\tilde{x}_t, \tilde{y}_t) = \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(\tilde{x}_t, \tilde{y}_t)$ 
6:     compute  $\nabla_y f(\tilde{x}_t, \tilde{y}_t) = \frac{1}{n} \sum_{i=1}^n \nabla_y f_i(\tilde{x}_t, \tilde{y}_t)$ 
7:     for all  $j = 0$  to  $N - 1$  do
8:       sample i.i.d. indices  $i_j^1, i_j^2$  uniformly from  $[n]$ 
9:        $x_{t,j+1} = x_{t,j} - \tau_1 [\nabla_x f_{i_j^1}(x_{t,j}, y_{t,j}) - \nabla_x f_{i_j^1}(\tilde{x}_t, \tilde{y}_t) + \nabla_x f(\tilde{x}_t, \tilde{y}_t)]$ 
10:       $y_{t,j+1} = y_{t,j} + \tau_2 [\nabla_y f_{i_j^2}(x_{t,j+1}, y_{t,j}) - \nabla_y f_{i_j^2}(\tilde{x}_t, \tilde{y}_t) + \nabla_y f(\tilde{x}_t, \tilde{y}_t)]$ 
11:    end for
12:     $\tilde{x}_{t+1} = x_{t,N}, \quad \tilde{y}_{t+1} = y_{t,N}$ 
13:  end for
14:  choose  $(x^k, y^k)$  from  $\{(x_{t,j}, y_{t,j})\}_{j=0}^{N-1}\}_{t=0}^{T-1}$  uniformly at random
15:   $\tilde{x}_0 = x^k, \quad \tilde{y}_0 = y^k$ 
16: end for
```

Theorem 4.1. Suppose Assumptions 2 and 4 hold and $f(x, y)$ satisfies the two-sided PL condition with μ_1 and μ_2 . Define $P_k = a^k + \frac{1}{20}b^k$, where $a^k = \mathbb{E}[g(x^k) - g^*]$ and $b^k = \mathbb{E}[g(x^k) - f(x^k, y^k)]$. If we run VR-AGDA with $\tau_1 = \beta/(28\kappa^8 l)$, $\tau_2 = \beta/(l\kappa^6)$, $N = \lfloor \alpha\beta^{-2/3}\kappa^9(2 + 4\beta^{1/2}\kappa^{-3})^{-1} \rfloor$ and $T = 1$, where α, β are constants irrelevant to l, n, μ_1, μ_2 , then $P_{k+1} \leq \frac{1}{2}P_k$. This further implies a total complexity of

$$\mathcal{O}((n + \kappa^9) \log(1/\epsilon))$$

for VR-AGDA to achieve an ϵ -optimal solution.

Theorem 4.2. Under the same assumptions in Theorem 4.1 and further assuming $n \leq \kappa^9$, if we run VR-AGDA with $\tau_1 = \beta/(28\kappa^2 l n^{2/3})$, $\tau_2 = \beta/(l n^{2/3})$, $N = \lfloor \alpha\beta^{-2/3}n(2 + 4\beta^{1/2}n^{-1/3})^{-1} \rfloor$, and $T = \lceil \kappa^3 n^{-1/3} \rceil$, where α, β are constants irrelevant to l, n, μ_1, μ_2 , then $P_{k+1} \leq \frac{1}{2}P_k$. This further implies a total complexity of

$$\mathcal{O}(n^{2/3}\kappa^3 \log(1/\epsilon))$$

for VR-AGDA to achieve an ϵ -optimal solution.

Remark 4. Theorems 4.1 and 4.2 are different in their choices of stepsizes and iteration numbers, which gives rise to different complexities. Another difference is that Theorem 4.2 only works in the regime where the number of components n is not “too large” compared to the condition number, i.e., $n \leq \kappa^9$, which naturally guarantees $T = \lceil \kappa^3 n^{-1/3} \rceil \geq 1$.

Remark 5. Since AGDA has complexity $\mathcal{O}(n\kappa^3 \log(1/\epsilon))$, VR-AGDA with the setting in Theorem 4.1 is better than AGDA when $n \geq \kappa^6$. With the setting in Theorem 4.2, VR-AGDA outperforms AGDA as long as the assumption $n \leq \kappa^9$ holds. As a result of these two theorems, VR-AGDA always improves over AGDA. Furthermore, VR-AGDA with the second setting has a lower complexity than the first setting in the regime $n \leq \kappa^9$, although the first setting allows a simpler double-loop algorithm. Figure 3 summarizes the performance of VR-AGDA compared to AGDA in different regimes of n and κ .




Figure 3: Comparison of complexities of AGDA and VR-AGDA, where VR-AGDA1, VR-AGDA2 correspond to the two settings in Theorems 4.1 and 4.2. In the regime $n \leq \kappa^9$, VR-AGDA2 performs best; in the regime $n \geq \kappa^9$, VR-AGDA1 performs best.

5 Experiments

In the introduction, we already presented the convergence results of AGDA on a two-dimensional nonconvex-nonconcave function that satisfies the two-sided PL condition. In this section, we will present numerical experiments on machine learning applications: robust least square and imitation learning for linear quadratic regulators (LQR). Particularly, we focus on the comparison between AGDA, Stoc-AGDA, and VR-AGDA.

5.1 Robust least square

We consider the least square problems with coefficient matrix $A \in \mathbb{R}^{n \times m}$ and noisy vector $y_0 \in \mathbb{R}^n$. We assume that y_0 is subject to bounded deterministic perturbation δ . Robust least square (RLS) minimizes the worst case residual, and can be formulated as [El Ghaoui and Lebret, 1997]:

$$\min_x \max_{\delta: \|\delta\| \leq \rho} \|Ax - y\|^2, \text{ where } \delta = y_0 - y.$$

We consider RLS with soft constraint:

$$\min_x \max_y F(x, y) := \|Ax - y\|_M^2 - \lambda \|y - y_0\|_M^2, \quad (13)$$

where we also adopt the general M-(semi-)norm in (13): $\|x\|_M^2 = x^T M x$ and M is positive semi-definite. $F(x, y)$ satisfies the two-sided PL condition when $\lambda > 1$, because it can be written as the composition of a strongly-convex-strongly-concave function and an affine function (Example 2). However, $F(x, y)$ is not strongly convex about x , and when M is not full-rank, it is not strongly concave about y .

Datasets. We use three datasets in the experiments, and two of them are generated in the same way as in Du and Hu [2019]. We generate the first dataset with $n = 1000$ and $m = 500$ by sampling rows of A from a Gaussian $\mathcal{N}(0, I_n)$ distribution and setting $y_0 = Ax^* + \epsilon$ with x^* from Gaussian $\mathcal{N}(0, 1)$ and ϵ from Gaussian $\mathcal{N}(0, 0.01)$. We set $M = I_n$ and $\lambda = 3$. The second dataset is the rescaled aquatic toxicity dataset by Cassotti et al. [2014], which uses 8 molecular descriptors of 546 chemicals to predict quantitative acute aquatic toxicity towards *Daphnia Magna*. We use $M = I$ and $\lambda = 2$ for this dataset. The third dataset is generated with $A \in \mathbb{R}^{1000 \times 500}$ from Gaussian $\mathcal{N}(0, \Sigma)$ where $\Sigma_{i,j} = 2^{-|i-j|/10}$, M being rank-deficit with positive eigenvalues sampled from $[0.2, 1.8]$ and $\lambda = 1.5$. These three datasets represent cases with low, median, and high condition numbers, respectively.




Figure 4: Comparison of the convergences of AGDA, Stoc-AGDA and SVRG-AGDA on three datasets based on two inaccuracy measures: (i) $\|x_t - x^*\|^2 + \|y_t - y^*\|^2$ (as shown in the first row), and (ii) $P_t = (g(x_t) - g^*) + (g(x_t) - f(x_t, y_t))$ (as shown in the second row).

Evaluation. For each dataset, we compare three algorithms: AGDA, Stoc-AGDA, and VR-AGDA. We tune the stepsizes of all algorithms to achieve the best convergence. For Stoc-AGDA, we choose constant stepsizes to form a fair comparison with the other two. We report the potential function value, i.e., P_t described in our theorems, and distance to the limit point $\|(x_t, y_t) - (x^*, y^*)\|^2$. These errors are plotted against the number of gradient evaluations normalized by n (i.e., number of full gradients). Results are reported in Figure 4. We observe that VR-AGDA and AGDA both exhibit linear convergence, and the speedup of VR-AGDA is fairly significant when the condition number is large, whereas Stoc-AGDA progresses fast at the beginning and stagnates later on. These numerical results clearly validate our theoretical findings.

5.2 Generative adversarial imitation learning for LQR

The optimal control problem for LQR can be formulated as:

$$\begin{aligned} \underset{\pi_t}{\text{minimize}} \quad & \mathbb{E}_{x_0 \sim \mathcal{D}} \sum_{t=0}^{\infty} x_t^\top Q x_t + u_t^\top R u_t \\ \text{such that} \quad & x_{t+1} = A x_t + B u_t, u_t = \pi_t(x_t) \end{aligned}$$

where $x_t \in \mathbb{R}^d$ is a state, $u_t \in \mathbb{R}^k$ is a control, $\mathcal{D}$ is the distribution of initial state x_0 , and π_t is a policy. It is known that the optimal policy is linear: $u_t = -K^* x_t$, where $K^* \in \mathbb{R}^{k \times d}$. If we parametrize the policy in the linear form, $u_t = -K x_t$, the problem can be written as:

$$\min_K C(K; Q, R) := \mathbb{E}_{x_0 \sim \mathcal{D}} \left[\sum_{t=0}^{\infty} (x_t^\top Q x_t + u_t^\top R u_t) \right]$$




Figure 5: AGDA and VR-AGDA on generative adversarial learning for LQR

where the trajectory is induced by LQR dynamics and policy K . In generative adversarial imitation learning for LQR, the trajectories induced by an expert policy K_E are observed and part of the goal is to learn the cost function parameters Q and R from the expert. This can be formulated as a minimax problem [Cai et al., 2019]:

$$\min_K \max_{(Q,R) \in \Theta} m(K, Q, R)$$

where $m(K, Q, R) := C(K; Q, R) - C(K_E; Q, R) - \Phi(Q, R)$, $\Theta = \{(Q, R) : \alpha_Q I \preceq Q \preceq \beta_Q I, \alpha_R I \preceq R \preceq \beta_R I\}$ and Φ is a strongly-convex regularizer. We sample n initial points $x_0^{(1)}, x_0^{(2)}, \dots, x_0^{(n)}$ from $\mathcal{D}$ and approximate $C(K; Q, R)$ by sample average

$$C_n(K; Q, R) := \frac{1}{n} \sum_{i=1}^n \left[\sum_{t=0}^{\infty} (x_t^\top Q x_t + u_t^\top R u_t) \right]_{x_0=x_0^{(i)}}.$$

We then consider

$$m_n(K, Q, R) = C_n(K; Q, R) - C_n(K_E; Q, R) - \Phi(Q, R).$$

Note that m_n satisfies the PL condition in terms of K [Fazel et al., 2018], and m_n is strongly-concave in terms of (Q, R) , so the function satisfies the two-sided PL condition.

In our experiment, we use $\Phi(Q, R) = \lambda(\|Q - \bar{Q}\|^2 + \|R - \bar{R}\|^2)$ for some $\bar{Q}, \bar{R}$ and $\lambda = 1$. We generate three datasets with different dimensions: (1) $d = 3, k = 2$; (2) $d = 20, k = 10$; (3) $d = 30, k = 20$. The initial distribution $\mathcal{D}$ is $\mathcal{N}(0, I_d)$ and we sample $n = 100$ initial points. The exact gradients can be computed based on the compact forms established in Fazel et al. [2018], Cai et al. [2019]. We compare AGDA and VR-AGDA under fine-tuned stepsizes, and track their errors in terms of $\|K_t - K^*\|^2 + \|Q_t - Q^*\|_F^2 + \|R_t - R^*\|_F^2$. The result is reported in Figure 5, which again indicates that VR-AGDA significantly outperforms AGDA.

6 Conclusion

In this paper, we identify a subclass of nonconvex-nonconcave minimax problems, represented by the so-called two-side PL condition, for which AGDA and Stoc-AGDA can converge to *global* saddle points. We also propose the first linearly-convergent variance-reduced AGDA algorithm that is provably always faster than AGDA, for this subclass of minimax problems. We hope this work can shed some light on the understanding of nonconvex-nonconcave minimax optimization: (1) different learning rates for two players are essential in GDA algorithms with alternating updates; (2) convexity-concavity is not a watershed to

guarantee global convergence of GDA algorithms; (3) the complexity of solving minimax problems under PL conditions may have high-order dependence on the condition number in contrast to problems with strong convex-concavity conditions. It remains interesting to explore whether similar results apply to GDA algorithms with simultaneous updates and whether these algorithms can be further accelerated with momentum or catalyst schemes.

References

- Jacob Abernethy, Kevin A Lai, and Andre Wibisono. Last-iterate convergence rates for min-max optimization. *arXiv preprint arXiv:1906.02027*, 2019.
- James P Bailey, Gauthier Gidel, and Georgios Piliouras. Finite regret and cycles with fixed step-size via alternating gradient descent-ascent. *arXiv preprint arXiv:1907.04392*, 2019.
- Qi Cai, Mingyi Hong, Yongxin Chen, and Zhaoran Wang. On the global convergence of imitation learning: A case for linear quadratic regulator. *arXiv preprint arXiv:1901.03674*, 2019.
- Matteo Cassotti, Davide Ballabio, Viviana Consonni, Andrea Mauri, Igor V Tetko, and Roberto Todeschini. Prediction of acute aquatic toxicity toward daphnia magna by using the ga-k nn method. *Alternatives to Laboratory Animals*, 42(1):31–41, 2014.
- Robert S Chen, Brendan Lucier, Yaron Singer, and Vasilis Syrgkanis. Robust optimization for non-convex objectives. In *Advances in Neural Information Processing Systems*, pages 4705–4714, 2017.
- Yichen Chen and Mengdi Wang. Stochastic primal-dual methods and sample complexity of reinforcement learning. *arXiv preprint arXiv:1612.02516*, 2016.
- Bo Dai, Niao He, Yunpeng Pan, Byron Boots, and Le Song. Learning from conditional distributions via dual embeddings. In *Artificial Intelligence and Statistics*, pages 1458–1467, 2017.
- Bo Dai, Albert Shaw, Lihong Li, Lin Xiao, Niao He, Zhen Liu, Jianshu Chen, and Le Song. SBEED: Convergent reinforcement learning with nonlinear function approximation. 80:1125–1134, 10–15 Jul 2018.
- Constantinos Daskalakis, Andrew Ilyas, Vasilis Syrgkanis, and Haoyang Zeng. Training gans with optimism. In *International Conference on Learning Representations (ICLR 2018)*, 2018.
- Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. In *International Conference on Machine Learning*, pages 1675–1685, 2019.
- Simon S Du and Wei Hu. Linear convergence of the primal-dual gradient method for convex-concave saddle point problems without strong convexity. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 196–205, 2019.
- Laurent El Ghaoui and Hervé Lebret. Robust solutions to least-squares problems with uncertain data. *SIAM Journal on matrix analysis and applications*, 18(4):1035–1064, 1997.
- Francisco Facchinei and Jong-Shi Pang. *Finite-dimensional variational inequalities and complementarity problems*. Springer Science & Business Media, 2007.
- Maryam Fazel, Rong Ge, Sham Kakade, and Mehran Mesbahi. Global convergence of policy gradient methods for the linear quadratic regulator. In *International Conference on Machine Learning*, pages 1467–1476, 2018.

- Gauthier Gidel, Reyhane Askari Hemmat, Mohammad Pezeshki, Rémi Le Priol, Gabriel Huang, Simon Lacoste-Julien, and Ioannis Mitliagkas. Negative momentum for improved game dynamics. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1802–1811, 2019.
- Ian Goodfellow, Yoshua Bengio, and Aaron Courville. *Deep learning*. MIT press, 2016.
- Chi Jin, Praneeth Netrapalli, and Michael I Jordan. What is local optimality in nonconvex-nonconcave minimax optimization? *arXiv preprint arXiv:1902.00618*, 2019.
- Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in neural information processing systems*, pages 315–323, 2013.
- Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-lojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.
- GM Korpelevich. The extragradient method for finding saddle points and other problems. *Matecon*, 12: 747–756, 1976.
- Qihang Lin, Mingrui Liu, Hassan Rafique, and Tianbao Yang. Solving weakly-convex-weakly-concave saddle-point problems as successive strongly monotone variational inequalities. *arXiv preprint arXiv:1810.10207*, 2018.
- Tianyi Lin, Chi Jin, and Michael I Jordan. On gradient descent ascent for nonconvex-concave minimax problems. *arXiv preprint arXiv:1906.00331*, 2019.
- Tianyi Lin, Chi Jin, Michael Jordan, et al. Near-optimal algorithms for minimax optimization. *arXiv preprint arXiv:2002.02417*, 2020.
- Luo Luo, Cheng Chen, Yujun Li, Guangzeng Xie, and Zhihua Zhang. A stochastic proximal point algorithm for saddle-point problems. *arXiv preprint arXiv:1909.06946*, 2019.
- Luo Luo, Haishan Ye, and Tong Zhang. Stochastic recursive gradient descent ascent for stochastic nonconvex-strongly-concave minimax problems. *arXiv preprint arXiv:2001.03724*, 2020.
- Zhi-Quan Luo and Paul Tseng. Error bounds and convergence analysis of feasible descent methods: a general approach. *Annals of Operations Research*, 46(1):157–178, 1993.
- Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. *arXiv preprint arXiv:1706.06083*, 2017.
- Eric Mazumdar and Lillian J Ratliff. On the convergence of gradient-based learning in continuous games. *ArXiv e-prints*, 2018.
- Lars Mescheder, Andreas Geiger, and Sebastian Nowozin. Which training methods for gans do actually converge? In *International Conference on Machine Learning*, pages 3481–3490, 2018.
- Hongseok Namkoong and John C Duchi. Stochastic gradient methods for distributionally robust optimization with f-divergences. In *Advances in Neural Information Processing Systems*, pages 2208–2216, 2016.
- Hongseok Namkoong and John C Duchi. Variance-based regularization with convex objectives. In *Advances in Neural Information Processing Systems*, pages 2971–2980, 2017.

- John Nash. Two-person cooperative games. *Econometrica: Journal of the Econometric Society*, pages 128–140, 1953.
- Ion Necoara, Yu Nesterov, and Francois Glineur. Linear convergence of first order methods for non-strongly convex optimization. *Mathematical Programming*, pages 1–39, 2018.
- Yurii Nesterov and Laura Scrimali. Solving strongly monotone variational and quasi-variational inequalities. *Available at SSRN 970903*, 2006.
- Maher Nouiehed, Maziar Sanjabi, Tianjian Huang, Jason D Lee, and Meisam Razaviyayn. Solving a class of non-convex min-max games using iterative first order methods. In *Advances in Neural Information Processing Systems*, pages 14905–14916, 2019.
- Balamurugan Palaniappan and Francis Bach. Stochastic variance reduction methods for saddle-point problems. In *Advances in Neural Information Processing Systems*, pages 1416–1424, 2016.
- Boris Teodorovich Polyak. Gradient methods for minimizing functionals. *Zhurnal Vychislitel’noi Matematiki i Matematicheskoi Fiziki*, 3(4):643–653, 1963.
- Qi Qian, Shenghuo Zhu, Jiasheng Tang, Rong Jin, Baigui Sun, and Hao Li. Robust optimization over multiple domains. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 33, pages 4739–4746, 2019.
- Sashank J Reddi, Ahmed Hefny, Suvrit Sra, Barnabas Poczos, and Alex Smola. Stochastic variance reduction for nonconvex optimization. In *International conference on machine learning*, pages 314–323, 2016a.
- Sashank J Reddi, Suvrit Sra, Barnabás Póczos, and Alex Smola. Fast incremental method for smooth nonconvex optimization. In *2016 IEEE 55th Conference on Decision and Control (CDC)*, pages 1971–1977. IEEE, 2016b.
- Aman Sinha, Hongseok Namkoong, and John Duchi. Certifiable distributional robustness with principled adversarial training. *stat*, 1050:29, 2017.
- Ju Sun, Qing Qu, and John Wright. A geometric analysis of phase retrieval. *Foundations of Computational Mathematics*, 18(5):1131–1198, 2018.
- Kiran K Thekumparampil, Prateek Jain, Praneeth Netrapalli, and Sewoong Oh. Efficient algorithms for smooth minimax optimization. In *Advances in Neural Information Processing Systems*, pages 12659–12670, 2019.
- John Von Neumann, Oskar Morgenstern, and Harold William Kuhn. *Theory of games and economic behavior (commemorative edition)*. Princeton university press, 2007.
- Lin Xiao and Tong Zhang. A proximal stochastic gradient method with progressive variance reduction. *SIAM Journal on Optimization*, 24(4):2057–2075, 2014.
- Linli Xu, James Neufeld, Bryce Larson, and Dale Schuurmans. Maximum margin clustering. In *Advances in neural information processing systems*, pages 1537–1544, 2005.
- Hui Zhang and Wotao Yin. Gradient methods for convex minimization: better rates under weaker conditions. *arXiv preprint arXiv:1303.4645*, 2013.
- Yi Zhou, Huishuai Zhang, and Yingbin Liang. Geometrical properties and accelerated gradient solvers of non-convex phase retrieval. In *2016 54th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 331–335. IEEE, 2016.

Appendix

A Proofs for Section 2

We first present several key lemmas.

Lemma A.1 (Karimi et al. [2016]). *If $f(\cdot)$ is l -smooth and it satisfies PL with constant μ , then it also satisfies error bound (EB) condition with μ , i.e.*

$$\|\nabla f(x)\| \geq \mu \|x_p - x\|, \forall x,$$

where x_p is the projection of x onto the optimal set, also it satisfies quadratic growth (QG) condition with μ , i.e.

$$f(x) - f^* \geq \frac{\mu}{2} \|x_p - x\|^2, \forall x.$$

Conversely, if $f(\cdot)$ is l -smooth and it satisfies EB with constant μ , then it satisfies PL with constant μ/l .

From the above lemma, we easily derive that $l \geq \mu$.

Lemma A.2 (Nouiehed et al. [2019]). *In the minimax problem, when $-f(x, \cdot)$ satisfies PL condition with constant μ_2 for any x and f satisfies Assumption 1, then the function $g(x) := \max_y f(x, y)$ is L -smooth with $L := l + l^2/\mu_2$ and $\nabla g(x) = \nabla_x f(x, y^*(x))$ for any $y^*(x) \in \arg \max_y f(x, y)$.*

Lemma A.3. *In the minimax problem 1, when the objective function f satisfies Assumption 1 (Lipschitz gradient) and the two-sided PL condition with constant μ_1 and μ_2 , then function $g(x) := \max_y f(x, y)$ satisfies the PL condition with μ_1 .*

Proof. From Lemma A.2,

$$\|\nabla g(x)\|^2 = \|\nabla_x f(x, y^*(x))\|^2.$$

Since $f(\cdot, y)$ satisfies PL condition with constant μ_1 , we get

$$\|\nabla g(x)\|^2 \geq 2\mu_1 [f(x, y^*(x)) - \min_{x'} f(x', y^*(x))]. \quad (14)$$

Also,

$$f(x', y^*(x)) \leq \max_y f(x', y) \implies \min_{x'} f(x', y^*(x)) \leq \min_{x'} \max_y f(x', y) = g^*. \quad (15)$$

Combining equation (14) and (15), we obtain,

$$\|\nabla g(x)\|^2 \geq 2\mu_1 (g(x) - g^*).$$

□

The following lemma states that stochastic gradient descent converges linearly to the neighbourhood of the optimal set under PL condition. The proof is based on [Karimi et al., 2016].

Lemma A.4. *Consider the optimization problem $\min_x f(x) = \mathbb{E}[F(x; \xi)]$, where f is l -smooth and satisfies PL condition with constant μ . Using the stochastic gradient descent with step size $\tau \leq 1/l$,*

$$x_{t+1} = x_t - \tau G(x_t, \xi_t),$$

where

$$\mathbb{E}[G(x, \xi) - \nabla f(x)] = 0, \quad \mathbb{E}[\|G(x, \xi) - \nabla f(x)\|^2] \leq \sigma^2,$$

then we have

$$\mathbb{E}[f(x_{t+1}) - f^*] \leq (1 - \mu\tau)\mathbb{E}[f(x_t) - f^*] + \frac{l\tau^2}{2}\sigma^2.$$

Proof. By smoothness of f we have

$$\begin{aligned} f(x_{t+1}) - f^* &\leq f(x_t) + \langle \nabla f(x_t), x_{t+1} - x_t \rangle + \frac{l}{2}\|x_{t+1} - x_t\|^2 - f^* \\ &= f(x_t) - \tau \langle \nabla f(x_t), G(x_t, \xi_t) \rangle + \frac{l\tau^2}{2}\|G(x_t, \xi_t)\|^2 - f^*. \end{aligned}$$

Taking expectation of both sides, we get

$$\begin{aligned} \mathbb{E}[f(x_{t+1}) - f^*] &\leq \mathbb{E}[f(x_t) - f^*] - \tau \mathbb{E}[\|\nabla f(x_t)\|^2] + \frac{l\tau^2}{2}\mathbb{E}[\|G(x_t, \xi_t)\|^2] \\ &= \mathbb{E}[f(x_t) - f^*] - \tau \mathbb{E}[\|\nabla f(x_t)\|^2] + \frac{l\tau^2}{2}\mathbb{E}[\|\nabla f(x_t)\|^2] \\ &\quad + \frac{l\tau^2}{2}\mathbb{E}[\|\nabla f(x_t) - G(x_t, \xi_t)\|^2] \\ &\leq \mathbb{E}[f(x_t) - f^*] - \frac{\tau}{2}\mathbb{E}[\|\nabla f(x_t)\|^2] + \frac{l\tau^2}{2}\sigma^2 \\ &\leq (1 - \mu\tau)\mathbb{E}[f(x_t) - f^*] + \frac{l\tau^2}{2}\sigma^2, \end{aligned}$$

where in the equality we use $\mathbb{E}[G(x_t, \xi_t)] = \nabla f(x_t)$, in the second inequality we use $\tau \leq 1/l$, and we use PL condition in the last inequality. $\square$

Proof for Lemma 2.1.

Proof. • (stationary point) $\implies$ (saddle point): From the definition of PL condition, if (x^*, y^*) is a stationary point,

$$\begin{aligned} \max_y f(x^*, y) - f(x^*, y^*) &\leq \frac{1}{2\mu_2} \|\nabla_y f(x^*, y^*)\|^2 = 0, \\ f(x^*, y^*) - \min_x f(x, y^*) &\leq \frac{1}{2\mu_1} \|\nabla_x f(x^*, y^*)\|^2 = 0, \end{aligned}$$

so $\max_y f(x^*, y) = f(x^*, y^*) = \min_x f(x, y^*)$, and therefore $f(x^*, y^*)$ is a saddle point.

- (saddle point) $\implies$ (global minimax point): Follow from definitions.
- (global minimax point) $\implies$ (stationary point): If (x^*, y^*) is a global minimax point, then by definition,

$$y^* \in \arg \max_y f(x^*, y), x^* \in \arg \min_x g(x),$$

Then by first order necessary condition, we have,

$$\nabla_y f(x^*, y^*) = 0, \nabla_x g(x^*) = 0,$$

Further with Lemma A.2,

$$\nabla g(x^*) = \nabla_x f(x^*, y^*) = 0$$

Thus, (x^*, y^*) is a stationary point. □

Proposition 1. *The function*

$$f(x, y) = x^2 + 3 \sin^2 x \sin^2 y - 4y^2 - 10 \sin^2 y,$$

satisfies the two-sided PL condition with $\mu_1 = 1/16, \mu_2 = 1/14$.

Proof. It is not hard to derive that $\arg \min_x f(x, y) = 0, \forall y$, and $\arg \max_y f(x, y) = 0, \forall x$, i.e. $x^*(y) = y^*(x) = 0, \forall x, y$. Therefore, $(0, 0)$ is the only saddle point. Then compute the gradients:

$$\begin{aligned} \nabla_x f(x, y) &= 2x + 3 \sin^2(y) \sin(2x), \\ \nabla_y f(x, y) &= -8y + 3 \sin^2(x) \sin(2y) - 10 \sin(2y). \end{aligned}$$

and

$$\begin{aligned} |\nabla_x^2 f(x, y)| &= |2 + 6 \sin^2(y) \cos(2x)| \leq 8, \\ |\nabla_y^2 f(x, y)| &= |-8 + 6 \sin^2(x) \cos(2y) - 20 \cos(2y)| \leq 28. \end{aligned}$$

so $f(\cdot, y)$ is L_1 -smooth with $L_1 = 8$ for any x and $f(x, \cdot)$ is L_2 -smooth with $L_2 = 28$ for any y . Then note that:

$$\begin{aligned} \frac{|\nabla_x f(x, y)|}{|x - x^*(y)|} &= \frac{|\nabla_x f(x, y)|}{|x|} = \frac{|2x + 3 \sin^2(y) \sin(2x)|}{|x|} \geq \frac{1}{2}, \\ \frac{|\nabla_y f(x, y)|}{|y - y^*(x)|} &= \frac{|\nabla_y f(x, y)|}{|y|} = \frac{|-8y + 3 \sin^2(x) \sin(2y) - 10 \sin(2y)|}{|y|} \geq 2 \end{aligned}$$

So $f(\cdot, y)$ satisfies EB with $\mu_{EB1} = 1/2$, and $-f(x, \cdot)$ satisfies EB with $\mu_{EB2} = 2$. By Lemma A.1, we have $f(\cdot, y)$ satisfies PL with constant $\mu_1 = 1/16$ and $-f(x, \cdot)$ satisfies PL with constant $\mu_1 = 1/14$. □

B Proofs for Section 3

Before we step into proofs for Theorem 3.1, 3.2 and 3.3, we first present a contraction theorem for each iteration.

Theorem B.1. *Assume Assumption 1, 2, 3 hold and $f(x, y)$ satisfies the two-sided PL condition with μ_1 and μ_2 . Define $a_t = \mathbb{E}[g(x_t) - g^*]$ and $b_t = \mathbb{E}[g(x_t) - f(x_t, y_t)]$. If we run one iteration of Algorithm 1 with $\tau_1^t = \tau_1 \leq 1/L$ (L is specified in Lemma A.2) and $\tau_2^t = \tau_2 \leq 1/l$, then*

$$a_{t+1} + \lambda b_{t+1} \leq \max\{k_1, k_2\}(a_t + \lambda b_t) + \lambda(1 - \mu_2 \tau_2) \frac{L+l}{2} \tau_1^2 \sigma^2 + \frac{l}{2} \lambda \tau_2^2 \sigma^2 + \frac{L}{2} \tau_1^2 \sigma^2,$$

where

$$k_1 := 1 - \mu_1 [\tau_1 + \lambda(1 - \mu_2 \tau_2) \tau_1 - \lambda(1 + \beta)(1 - \mu_2 \tau_2)(2\tau_1 + l\tau_1^2)], \quad (16)$$

$$k_2 := 1 - \mu_2 \tau_2 + \frac{l^2 \tau_1}{\mu_2 \lambda} + (1 - \mu_2 \tau_2) \frac{l^2}{\mu_2} \tau_1 + (1 + \frac{1}{\beta})(1 - \mu_2 \tau_2) \frac{l^2}{\mu_2} (2\tau_1 + l\tau_1^2), \quad (17)$$

and $\lambda, \beta > 0$ such that $k_1 \leq 1$.

Proof. Because g is L -smooth by Lemma A.2, we have

$$\begin{aligned} g(x_{t+1}) - g^* &\leq g(x_t) - g^* + \langle \nabla g(x_t), x_{t+1} - x_t \rangle + \frac{L}{2} \|x_{t+1} - x_t\|^2 \\ &= g(x_t) - g^* - \tau_1 \langle \nabla g(x_t), G_x(x_t, y_t, \xi_{t1}) \rangle + \frac{L}{2} \tau_1^2 \|G_x(x_t, y_t, \xi_{t1})\|^2. \end{aligned}$$

Taking expectation of both side and use Assumption 3, we get

$$\begin{aligned} \mathbb{E}[g(x_{t+1}) - g^*] &\leq \mathbb{E}[g(x_t) - g^*] - \tau_1 \mathbb{E}[\langle \nabla g(x_t), \nabla_x f(x_t, y_t) \rangle] + \frac{L}{2} \tau_1^2 \mathbb{E}[\|G_x(x_t, y_t, \xi_{t1})\|^2] \\ &\leq \mathbb{E}[g(x_t) - g^*] - \tau_1 \mathbb{E}[\langle \nabla g(x_t), \nabla_x f(x_t, y_t) \rangle] + \frac{L}{2} \tau_1^2 \mathbb{E}[\|\nabla_x f(x_t, y_t)\|^2] + \frac{L}{2} \tau_1^2 \sigma^2 \\ &\leq \mathbb{E}[g(x_t) - g^*] - \tau_1 \mathbb{E}[\langle \nabla g(x_t), \nabla_x f(x_t, y_t) \rangle] + \frac{\tau_1}{2} \mathbb{E}[\|\nabla_x f(x_t, y_t)\|^2] + \frac{L}{2} \tau_1^2 \sigma^2 \\ &\leq \mathbb{E}[g(x_t) - g^*] - \frac{\tau_1}{2} \mathbb{E}[\|\nabla g(x_t)\|^2] + \frac{\tau_1}{2} \mathbb{E}[\|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2] + \frac{L}{2} \tau_1^2 \sigma^2, \end{aligned} \quad (18)$$

where in the second inequality we use Assumption 3, and in the third inequality we use $\tau_1 \leq 1/L$. Because $-f(x_{t+1}, y)$ is l -smooth and μ_1 -PL, by Lemma A.4, when $\tau_1 \leq 1/l$ we have

$$\begin{aligned} \mathbb{E}[g(x_{t+1}) - f(x_{t+1}, y_{t+1})] &\leq (1 - \mu_2 \tau_2) \mathbb{E}[g(x_{t+1}) - f(x_{t+1}, y_t)] + \frac{l}{2} \tau_2^2 \sigma^2 \\ &\leq (1 - \mu_2 \tau_2) \mathbb{E}[g(x_t) - f(x_t, y_t) + f(x_t, y_t) - f(x_{t+1}, y_t) + g(x_{t+1}) - g(x_t)] + \frac{l}{2} \tau_2^2 \sigma^2 \end{aligned} \quad (19)$$

Because of lipschitz continuity of the gradient, we can bound $f(x_t, y_t) - f(x_{t+1}, y_t)$ as

$$\begin{aligned} f(x_t, y_t) - f(x_{t+1}, y_t) &\leq -\langle \nabla_x f(x_t, y_t), x_{t+1} - x_t \rangle + \frac{l}{2} \|x_{t+1} - x_t\|^2 \\ &\leq \tau_1 \langle \nabla_x f(x_t, y_t), G_x(x_t, y_t, \xi_{t1}) \rangle + \frac{l}{2} \tau_1^2 \|G_x(x_t, y_t, \xi_{t1})\|^2. \end{aligned}$$

Taking expectation of both side and use Assumption 3,

$$\mathbb{E}[f(x_t, y_t) - f(x_{t+1}, y_t)] \leq (\tau_1 + \frac{l}{2} \tau_1^2) \mathbb{E}[\|\nabla_x f(x_t, y_t)\|^2] + \frac{l}{2} \tau_1^2 \sigma^2. \quad (20)$$

Also from (18) ,

$$\mathbb{E}[g(x_{t+1}) - g(x_t)] \leq -\frac{\tau_1}{2} \mathbb{E}[\|\nabla g(x_t)\|^2] + \frac{\tau_1}{2} \mathbb{E}[\|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2] + \frac{L}{2} \tau_1^2 \sigma^2. \quad (21)$$

Combining (19), (20) and (21),

$$\begin{aligned} \mathbb{E}[g(x_{t+1}) - f(x_{t+1}, y_{t+1})] &\leq (1 - \mu_2 \tau_2) \mathbb{E}[g(x_t) - f(x_t, y_t)] + (1 - \mu_2 \tau_2) (\tau_1 + \frac{l}{2} \tau_1^2) \mathbb{E}[\|\nabla_x f(x_t, y_t)\|^2] - \\ &\quad (1 - \mu_2 \tau_2) \frac{\tau_1}{2} \mathbb{E}[\|\nabla g(x_t)\|^2] + (1 - \mu_2 \tau_2) \frac{\tau_1}{2} \mathbb{E}[\|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2] + \\ &\quad (1 - \mu_2 \tau_2) \frac{L + l}{2} \tau_1^2 \sigma^2 + \frac{l}{2} \tau_2^2 \sigma^2. \end{aligned} \quad (22)$$

Combining (18) and (22), we have for $\forall \lambda > 0$

$$\begin{aligned}
a_{t+1} + \lambda b_{t+1} &\leq a_t - \left[\frac{\tau_1}{2} + \lambda(1 - \mu_2 \tau_1) \frac{\tau_1}{2} \right] \mathbb{E} \|\nabla g(x_t)\|^2 + \lambda(1 - \mu_2 \tau_2) b_t + \\
&\quad \left[\frac{\tau_1}{2} + \lambda(1 - \mu_2 \tau_2) \frac{\tau_1}{2} \right] \mathbb{E} \|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 + \lambda(1 - \mu_2 \tau_2) \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \mathbb{E} \|\nabla_x f(x_t, y_t)\|^2 + \\
&\quad \lambda(1 - \mu_2 \tau_2) \frac{L+l}{2} \tau_1^2 \sigma^2 + \frac{l}{2} \lambda \tau_2^2 \sigma^2 + \frac{L}{2} \tau_1^2 \sigma^2 \\
&\leq a_t - \left[\frac{\tau_1}{2} + \lambda(1 - \mu_2 \tau_1) \frac{\tau_1}{2} - \lambda(1 + \beta)(1 - \mu_2 \tau_2) \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right] \mathbb{E} \|\nabla g(x_t)\|^2 + \\
&\quad \lambda(1 - \mu_2 \tau_2) b_t + \left[\frac{\tau_1}{2} + \lambda(1 - \mu_2 \tau_2) \frac{\tau_1}{2} + \lambda \left(1 + \frac{1}{\beta} \right) (1 - \mu_2 \tau_2) \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right] \mathbb{E} \|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 + \\
&\quad \lambda(1 - \mu_2 \tau_2) \frac{L+l}{2} \tau_1^2 \sigma^2 + \frac{l}{2} \lambda \tau_2^2 \sigma^2 + \frac{L}{2} \tau_1^2 \sigma^2, \tag{23}
\end{aligned}$$

where in the second inequality we use Young's Inequality and $\beta > 0$. Now it suffices to bound $\|g(x_t)\|^2$ and $\|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2$ by a_t and b_t . With Lemma A.2, we have:

$$\|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 = \|\nabla_x f(x_t, y_t) - \nabla_x f(x_t, y^*(x_t))\|^2 \leq l^2 \|y^*(x_t) - y_t\|^2, \tag{24}$$

for any $y^*(x_t) \in \arg \max_y f(x_t, y)$. Now we fix $y^*(x_t)$ to be the projection of y_t on the the set $\arg \max_y f(x_t, y)$. Because $-f(\mathbf{x}_t, \cdot)$ satisfies PL condition with μ_2 , and Lemma A.1 therefore indicates it also satisfies quadratic growth condition with μ_2 , i.e.

$$\|y^*(x_t) - y_t\|^2 \leq \frac{2}{\mu_2} [g(x_t) - f(x_t, y_t)], \tag{25}$$

along with (24), we get

$$\|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 \leq \frac{2l^2}{\mu_2} [g(x_t) - f(x_t, y_t)]. \tag{26}$$

Because g satisfies PL condition with μ_1 by Lemma A.3,

$$\|\nabla g(x_t)\|^2 \geq 2\mu_1 [g(x_t) - g^*]. \tag{27}$$

Plug (26) and (27) into (23), we can get

$$\begin{aligned}
a_{t+1} + \lambda b_{t+1} &\leq \left\{ 1 - \mu_1 [\tau_1 + \lambda(1 - \mu_2 \tau_2) \tau_1 - \lambda(1 + \beta)(1 - \mu_2 \tau_2)(2\tau_1 + l\tau_1^2)] \right\} a_t + \\
&\quad \lambda \left\{ 1 - \mu_2 \tau_2 + \frac{l^2 \tau_1}{\mu_2 \lambda} + (1 - \mu_2 \tau_2) \frac{l^2}{\mu_2} \tau_1 + (1 + \frac{1}{\beta})(1 - \mu_2 \tau_2) \frac{l^2}{\mu_2} (2\tau_1 + l\tau_1^2) \right\} b_t + \\
&\quad \lambda(1 - \mu_2 \tau_2) \frac{L+l}{2} \tau_1^2 \sigma^2 + \frac{l}{2} \lambda \tau_2^2 \sigma^2 + \frac{L}{2} \tau_1^2 \sigma^2. \tag{28}
\end{aligned}$$

□

Proof of Theorem 3.1

Proof. In the setting of Theorem 1, $\tau_1^t = \tau_1$ and $\tau_2^t = \tau_2, \forall t$. By Thoerem B.1, We only need to choose τ_1, τ_2, λ and β to let $k_1, k_2 < 1$. Here we first choose $\beta = 1$ and $\lambda = 1/10$. Then

$$\begin{aligned}
k_1 &= 1 - \mu_1 [\tau_1 + \lambda(1 - \mu_2 \tau_2) \tau_1 - \lambda(1 + \beta)(1 - \mu_2 \tau_2)(2\tau_1 + l\tau_1^2)] \\
&\leq 1 - \mu_1 \{ \tau_1 - \lambda(1 - \mu_2 \tau_2) \tau_1 [(1 + \beta)(2 + l\tau_1) - 1] \} \leq 1 - \frac{1}{2} \tau_1 \mu_1, \tag{29}
\end{aligned}$$

where in the last inequality we just plug in β and λ and use $l\tau_1 \leq 1$. Also,

$$\begin{aligned}
k_2 &= 1 - \mu_2\tau_2 + \frac{l^2\tau_1}{\mu_2\lambda} + (1 - \mu_2\tau_2)\frac{l^2}{\mu_2}\tau_1 + (1 + \frac{1}{\beta})(1 - \mu_2\tau_2)\frac{l^2}{\mu_2}(2\tau_1 + l\tau_1^2) \\
&\leq 1 - \frac{l^2\tau_1}{\mu_2} \left\{ \frac{\mu_2^2\tau_2}{\tau_1 l^2} - \frac{1}{\lambda} - (1 - \mu_2\tau_2) \left[1 + \left(1 + \frac{1}{\beta} \right) (2 + l\tau_1) \right] \right\} \\
&\leq 1 - \frac{l^2\tau_1}{\mu_2},
\end{aligned} \tag{30}$$

where in the last inequality we plug in β and λ and we use $\frac{\mu_2^2\tau_2}{\tau_1 l^2} \leq 18$ by our choice of τ_1 . Note that $\frac{1}{2}\tau_1\mu_1 < \frac{l^2\tau_1}{\mu_2}$, because $(\frac{1}{2}\tau_1\mu_1) / (\frac{l^2\tau_1}{\mu_2}) = \frac{\mu_1\mu_2}{2l^2} < 1$. Define $P_t := a_t + \frac{1}{10}b_t$, and by Theorem B.1,

$$P_{t+1} \leq \left(1 - \frac{1}{2}\tau_1\mu_1\right) P_t + \frac{(1 - \mu_2\tau_2)(L + l)\tau_1^2}{20}\sigma^2 + \frac{l\tau_2^2}{20}\sigma^2 + \frac{L\tau_1^2}{2}\sigma^2.$$

With some simple computation,

$$P_t \leq (1 - \frac{1}{2}\mu_1\tau_1)^t P_0 + \frac{(1 - \mu_2\tau_2)(L + l)\tau_1^2 + l\tau_2^2 + 10L\tau_1^2}{10\mu_1\tau_1}\sigma^2.$$

We verify that $\tau_1 \leq 1/L$ by noting: $\tau_1 \leq \frac{\mu_2^2\tau_2}{18l^2} \leq \frac{\mu_2^2}{18l^3} \leq \frac{\mu_2}{2l^2}$ and $L = l + \frac{l^2}{\mu_2} \leq \frac{2l^2}{\mu_2}$. $\square$

Proof of Theorem 3.2

Proof. The first part of Theorem 3.2 is a direct corollary of Theorem 3.1 by setting $\sigma = 0$. We show the second part by noting that

$$\|x_{t+1} - x_t\|^2 = \tau_1^2 \|\nabla_x f(x_t, y_t)\|^2, \text{ and } \|y_{t+1} - y_t\|^2 = \tau_2^2 \|\nabla_y f(x_{t+1}, y_t)\|^2. \tag{31}$$

Also,

$$\begin{aligned}
\|\nabla_y f(x_{t+1}, y_t)\|^2 &\leq \|\nabla_y f(x_t, y_t)\|^2 + \|\nabla_y f(x_{t+1}, y_t) - \nabla_y f(x_t, y_t)\|^2 \\
&\leq \|\nabla_y f(x_t, y_t) - \nabla_y f(x_t, y^*(x_t))\|^2 + l^2 \|x_{t+1} - x_t\|^2 \\
&\leq l^2 \|y_t - y^*(x_t)\|^2 + l^2 \|x_{t+1} - x_t\|^2 \\
&\leq \frac{2l^2}{\mu_2} b_t + l^2 \|x_{t+1} - x_t\|^2 = \frac{2l^2}{\mu_2} b_t + l^2 \tau_1^2 \|\nabla_x f(x_t, y_t)\|^2,
\end{aligned} \tag{32}$$

where in the second inequality $y^*(x_t)$ is the projection of y_t on the the set $\arg\max_y f(x_t, y)$ and $\nabla_y f(x_t, y^*(x_t)) = 0$, in the third inequality we use lipschitz continuity of gradient, and in the last inequality we use quadratic growth condition. Also,

$$\begin{aligned}
\|\nabla_x f(x_t, y_t)\|^2 &\leq \|\nabla g(x_t)\|^2 + \|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 \\
&= \|\nabla g(x_t) - \nabla g(x^*)\|^2 + \|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 \\
&\leq L^2 \|x_t - x^*\|^2 + l^2 \|y^*(x_t) - y_t\|^2 \\
&\leq \frac{2L^2}{\mu_1} a_t + \frac{2l^2}{\mu_2} b_t,
\end{aligned} \tag{33}$$

where in the first equality x^* is the projection of x_t on the set $\arg \min_x g(x)$ and $\nabla g(x^*) = 0$, in the second inequality $y^*(x_t)$ is the projection of y_t on the the set $\arg \max_y f(x_t, y)$ and $\nabla g(x_t) = \nabla_x f(x_t, y_t)$, and in the last inequality we use quadratic growth condition. Therefore with (32) and (33),

$$\begin{aligned} \|x_t - x^*\|^2 + \|y_t - y^*\|^2 &\leq \tau_1^2 \|\nabla_x f(x_t, y_t)\|^2 + \tau_2^2 \|\nabla_y f(x_{t+1}, y_t)\|^2 \\ &\leq (1 + \tau_2^2 l^2) \tau_1^2 \|\nabla_x f(x_t, y_t)\|^2 + \frac{2l^2}{\mu_2} \tau_2^2 b_t \\ &\leq \frac{2(1 + \tau_2^2 l^2) \tau_1^2 L^2}{\mu_1} a_t + \frac{2(1 + \tau_2^2 l^2) \tau_1^2 l^2 + 2l^2 \tau_2^2}{\mu_2} b_t \\ &\leq \left[\frac{2(1 + \tau_2^2 l^2) \tau_1^2 L^2}{\mu_1} + \frac{20(1 + \tau_2^2 l^2) \tau_1^2 l^2 + 20l^2 \tau_2^2}{\mu_2} \right] P_0 c^t, \end{aligned}$$

where $c = 1 - \frac{\mu_1 \mu_2^2}{36l^3}$. Letting $\alpha_1 = \left[\frac{2(1 + \tau_2^2 l^2) \tau_1^2 L^2}{\mu_1} + \frac{20(1 + \tau_2^2 l^2) \tau_1^2 l^2 + 20l^2 \tau_2^2}{\mu_2} \right] P_0$, we have

$$\|x_{t+1} - x_t\| + \|y_{t+1} - y_t\| \leq \sqrt{2\alpha_1} c^{t/2}.$$

For $n \geq t$,

$$\|x_n - x_t\| + \|y_n - y_t\| \leq \sum_{i=t}^{n-1} \|x_{i+1} - x_i\| + \|y_{i+1} - y_i\| \leq \sqrt{2\alpha_1} \sum_{i=t}^{\infty} c^{i/2} \leq \frac{\sqrt{2\alpha_1} c^{t/2}}{1 - \sqrt{c}},$$

so $\{(x_t, y_t)\}_t$ converges and by first part of this theorem the limit (x^*, y^*) must be a saddle point. Thus we have

$$\|x_t - x^*\|^2 + \|y_t - y^*\|^2 \leq \frac{2\alpha_1}{(1 - \sqrt{c})^2} c^t = \alpha c^t P_0,$$

with $\alpha = 2 \left[\frac{2(1 + \tau_2^2 l^2) \tau_1^2 L^2}{\mu_1} + \frac{20(1 + \tau_2^2 l^2) \tau_1^2 l^2 + 20l^2 \tau_2^2}{\mu_2} \right] / (1 - \sqrt{c})^2$. $\square$

Proof of Theorem 3.3

Proof. First note that since $\tau_1^t \leq \mu_2^2 / 18l^2$, $\tau_2^t = \frac{18l^2 \beta}{\mu_2^2 (\gamma + t)} = \frac{18l^2 \tau_1^t}{\mu_2^2} \leq \frac{1}{l}$. Similar to the proof of Theorem 3.1, by choosing $\beta = 1$ and $\lambda = 1/10$ in the Theorem B.1, we have $\min\{k_1, k_2\} = \frac{1}{2} \mu_1 \tau_1^t$. We prove the theorem by induction. When $t = 1$, it is naturally satisfied by definition of ν . We assume that $P_t \leq \frac{\nu}{\gamma + t}$. Then by Theorem B.1,

$$\begin{aligned} P_{t+1} &\leq \left(1 - \frac{1}{2} \mu_1 \tau_1\right) P_t + \lambda(1 - \mu_2 \tau_2^t) \frac{L + l}{2} (\tau_1^t)^2 \sigma^2 + \frac{l}{2} \lambda (\tau_2^t)^2 \sigma^2 + \frac{L}{2} (\tau_1^t)^2 \sigma^2 \\ &\leq \frac{\gamma + t - \frac{1}{2} \mu_1 \beta}{\gamma + t} \frac{\nu}{\gamma + t} + \left[\frac{(L + l) \beta^2}{20(\gamma + t)^2} + \frac{18^2 l^5 \beta^2}{20 \mu_2^4 (\gamma + t)^2} + \frac{L \beta^2}{2(\gamma + t)^2} \right] \sigma^2 \\ &\leq \frac{\gamma + t - 1}{(\gamma + t)^2} \nu - \frac{\frac{1}{2} \mu_1 \beta - 1}{(\gamma + t)^2} \nu + \left[\frac{(L + l) \beta^2}{20(\gamma + t)^2} + \frac{18^2 l^5 \beta^2}{20 \mu_2^4 (\gamma + t)^2} + \frac{L \beta^2}{2(\gamma + t)^2} \right] \sigma^2 \\ &\leq \frac{\nu}{\gamma + t + 1}, \end{aligned} \tag{34}$$

where in the second inequality we plug in τ_1^t and τ_2^t , in the last inequality we use $(\gamma + t + 1)(\gamma + t - 1) \leq (\gamma + t)^2$ and the fact that sum of last two terms in (34) is no greater than 0 by our choice of ν . $\square$

C Proofs for Section 4

Proof of Theorem 4.1

Proof. Because the proof is long, we break the proof into three parts for the convenience of understanding the intuition behind it.

Part 1.

Consider in one outer loop k . Define $a_{t,j} = \mathbb{E}[g(x_{t,j}) - g^*]$, $b_{t,j} = \mathbb{E}[g(x_{t,j}) - f(x_{t,j}, y_{t,j})]$, $\tilde{a}_t = \mathbb{E}[g(\tilde{x}_t) - g^*]$ and $\tilde{b}_t = \mathbb{E}[g(\tilde{x}_t) - f(\tilde{x}_t, \tilde{y}_t)]$. We omit the subscript t for now. We denote the stochastic gradients as

$$\begin{aligned} G_x(x_j, y_j) &= \nabla_x f_{i_j}(x_j, y_j) - \nabla_x f_{i_j}(\tilde{x}, \tilde{y}) + \nabla_x f(\tilde{x}, \tilde{y}), \\ G_y(x_j, y_{j+1}) &= \nabla_y f_{i_j}(x_{j+1}, y_j) - \nabla_y f_{i_j}(\tilde{x}, \tilde{y}) + \nabla_y f(\tilde{x}, \tilde{y}). \end{aligned}$$

Note that these are unbiased stochastic gradients. Similar to the proof of Theorem B.1 (replace σ^2 in (18)), with $\tau_1 \leq 1/L$, we have

$$a_{j+1} \leq a_j - \frac{\tau_1}{2} \mathbb{E} \|\nabla g(x_j)\|^2 + \frac{\tau_1}{2} \mathbb{E} \|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \frac{L}{2} \tau_1^2 \mathbb{E} \|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 \quad (35)$$

By Lemma A.4, with $\tau_2 \leq 1/l$,

$$b_{j+1} \leq \mathbb{E}[g(x_{j+1}) - f(x_{j+1}, y_j)] - \frac{\tau_2}{2} \mathbb{E} \|\nabla_y f(x_{j+1}, y_j)\|^2 + \frac{l}{2} \tau_2^2 \mathbb{E} \|G_y(x_{j+1}, y_j) - \nabla_y f(x_{j+1}, y_j)\|^2 \quad (36)$$

Furthermore, we bound the distance to the $\tilde{x} = x_0$ as

$$\begin{aligned} \mathbb{E} \|x_{j+1} - \tilde{x}\|^2 &= \mathbb{E} \|x_j - \tau_1 G_x(x_j, y_j) - \tilde{x}\|^2 \\ &= \mathbb{E} \|x_j - \tilde{x}\|^2 + 2\mathbb{E} \langle x_j - \tilde{x}, \tau_1 \nabla_x f(x_j, y_j) \rangle + \tau_1^2 \mathbb{E} \|\nabla_x f(x_j, y_j)\|^2 + \tau_1^2 \mathbb{E} \|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 \\ &\leq (1 + \tau_1 \beta_1) \mathbb{E} \|x_j - \tilde{x}\|^2 + \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) \mathbb{E} \|\nabla_x f(x_j, y_j)\|^2 + \tau_1^2 \mathbb{E} \|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2, \end{aligned} \quad (37)$$

where in the last inequality we use Young's inequality to the inner product and $\beta_1 > 0$ is a constant which we will determine later. Similarly,

$$\mathbb{E} \|y_{j+1} - \tilde{y}\|^2 \leq (1 + \tau_2 \beta_2) \mathbb{E} \|y_j - \tilde{y}\|^2 + \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mathbb{E} \|\nabla_y f(x_{j+1}, y_j)\|^2 + \tau_2^2 \mathbb{E} \|G_y(x_{j+1}, y_j) - \nabla_y f(x_{j+1}, y_j)\|^2 \quad (38)$$

where in the last inequality we use Young's inequality to the inner product and $\beta_2 > 0$ is a constant. We are going to construct a potential function

$$R_j = a_j + \lambda b_j + c_j \|x_j - \tilde{x}\|^2 + d_j \|y_j - \tilde{y}\|^2, \quad (39)$$

and we will determine λ, c_j and d_j later. Combine (35), (36) and (38),

$$\begin{aligned} R_{j+1} &\leq a_j - \frac{\tau_1}{2} \mathbb{E} \|\nabla g(x_j)\|^2 + \frac{\tau_1}{2} \mathbb{E} \|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \frac{L}{2} \tau_1^2 \mathbb{E} \|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 + \\ &\quad \lambda \mathbb{E}[g(x_{j+1}) - f(x_{j+1}, y_j)] - \frac{\lambda \tau_2}{2} \mathbb{E} \|\nabla_y f(x_{j+1}, y_j)\|^2 + \\ &\quad c_{j+1} \mathbb{E} \|x_{j+1} - \tilde{x}\|^2 + \left(d_{j+1} + \frac{\lambda l}{2} \right) \tau_2^2 \mathbb{E} \|G_y(x_{j+1}, y_j) - \nabla_y f(x_{j+1}, y_j)\|^2 + \\ &\quad d_{j+1} (1 + \tau_2 \beta_2) \mathbb{E} \|y_j - \tilde{y}\|^2 + d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mathbb{E} \|\nabla_y f(x_{j+1}, y_j)\|^2 \end{aligned} \quad (40)$$

Then we bound the variance of the stochastic gradients,

$$\begin{aligned}\mathbb{E}\|G_y(x_{j+1}, y_j) - \nabla_y f(x_{j+1}, y_j)\|^2 &= \mathbb{E}\|\nabla_y f_{i_j}(x_{j+1}, y_j) - \nabla_y f_{i_j}(\tilde{x}, \tilde{y}) + \nabla_y f(\tilde{x}, \tilde{y}) - \nabla_y f(x_{j+1}, y_j)\|^2 \\ &\leq \mathbb{E}\|\nabla_y f_{i_j}(x_{j+1}, y_j) - \nabla_y f_{i_j}(\tilde{x}, \tilde{y})\|^2 \leq l^2 \mathbb{E}\|x_{j+1} - \tilde{x}\|^2 + l^2 \mathbb{E}\|y_j - \tilde{y}\|^2\end{aligned}\quad (41)$$

where in the first inequality we use $\mathbb{E}[\nabla_y f_{i_j}(x_{j+1}, y_j) - \nabla_y f_{i_j}(\tilde{x}, \tilde{y})] = \nabla_y f(x_{j+1}, y_j) - \nabla_y f(\tilde{x}, \tilde{y})$. Similarly,

$$\mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 \leq l^2 \mathbb{E}\|x_j - \tilde{x}\|^2 + l^2 \mathbb{E}\|y_j - \tilde{y}\|^2. \quad (42)$$

Plugging (41) into (40),

$$\begin{aligned}R_{j+1} &\leq a_j - \frac{\tau_1}{2} \mathbb{E}\|\nabla g(x_j)\|^2 + \frac{\tau_1}{2} \mathbb{E}\|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \frac{L}{2} \tau_1^2 \mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 + \\ &\quad \lambda \mathbb{E}[g(x_{j+1}) - f(x_{j+1}, y_j)] - \frac{\lambda \tau_2}{2} \mathbb{E}\|\nabla_y f(x_{j+1}, y_j)\|^2 + \\ &\quad \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \mathbb{E}\|x_{j+1} - \tilde{x}\|^2 + \\ &\quad \left[d_{j+1}(1 + \tau_2 \beta_2) + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \mathbb{E}\|y_j - \tilde{y}\|^2 + d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mathbb{E}\|\nabla_y f(x_{j+1}, y_j)\|^2.\end{aligned}\quad (43)$$

Then we plug in (37) and rearrange,

$$\begin{aligned}R_{j+1} &\leq a_j - \frac{\tau_1}{2} \mathbb{E}\|\nabla g(x_j)\|^2 + \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) \mathbb{E}\|\nabla_x f(x_j, y_j)\|^2 + \frac{\tau_1}{2} \mathbb{E}\|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \\ &\quad \lambda \mathbb{E}[g(x_{j+1}) - f(x_{j+1}, y_j)] - \left[\frac{\lambda \tau_2}{2} - d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \right] \mathbb{E}\|\nabla_y f(x_{j+1}, y_j)\|^2 + \\ &\quad \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] (1 + \tau_1 \beta_1) \mathbb{E}\|x_j - \tilde{x}\|^2 + \left[d_{j+1}(1 + \tau_2 \beta_2) + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \mathbb{E}\|y_j - \tilde{y}\|^2 + \\ &\quad \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \tau_1^2 \mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2\end{aligned}\quad (44)$$

Consider the second line. Using PL condition $\|\nabla_y f(x_{j+1}, y_j)\|^2 \geq 2\mu_2[g(x_{j+1}) - f(x_{j+1}, y_j)]$ and assuming $\lambda \geq d_{j+1}(\tau_2 + 1/\beta_2)$, which we will justify later by our choices of d_{j+1} and β_2 , we have

$$\begin{aligned}\text{the second line} &\leq \lambda \left[1 - \tau_2 \mu_2 + \frac{\lambda}{2} d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 \right] \mathbb{E}[g(x_{j+1}) - f(x_{j+1}, y_j)] \\ &\leq \lambda \left[1 - \tau_2 \mu_2 + \frac{\lambda}{2} d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 \right] \left\{ b_j + \mathbb{E}(f(x_j, y_j) - f(x_{j+1}, y_j)) + (a_{j+1} - a_j) \right\} \\ &\leq \lambda \left[1 - \tau_2 \mu_2 + \frac{\lambda}{2} d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 \right] \left\{ b_j + \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \mathbb{E}\|\nabla_x f(x_j, y_j)\|^2 + \right. \\ &\quad \left. \frac{l}{2} \tau_1^2 \mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 - \frac{\tau_1}{2} \mathbb{E}\|\nabla g(x_j)\|^2 + \right. \\ &\quad \left. \frac{\tau_1}{2} \mathbb{E}\|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \frac{L}{2} \tau_1^2 \mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2 \right\}\end{aligned}$$

where in the last inequality we use (35) and (20). Now we plug this into R_{j+1} ,

$$\begin{aligned}
R_{j+1} \leq & a_j - \frac{\tau_1}{2}(1 + \lambda\zeta)\mathbb{E}\|\nabla g(x_j)\|^2 + \left\{ \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) + \lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right\} \mathbb{E}\|\nabla_x f(x_j, y_j)\|^2 + \\
& \frac{\tau_1}{2}(1 + \lambda\zeta)\mathbb{E}\|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \lambda\zeta b_j + \\
& \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] (1 + \tau_1 \beta_1) \mathbb{E}\|x_j - \tilde{x}\|^2 + \left[d_{j+1}(1 + \tau_2 \beta_2) + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \mathbb{E}\|y_j - \tilde{y}\|^2 + \\
& \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 + \lambda\zeta \frac{L+l}{2} \right] \tau_1^2 \mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2, \tag{45}
\end{aligned}$$

where we define $\zeta = 1 - \tau_2 \mu_2 + \frac{\lambda}{2} d_{j+1} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2$ and $\psi = 1 - \zeta$. With $\|\nabla_x f(x_j, y_j)\|^2 \leq 2\|\nabla g(x_j)\|^2 + 2\|\nabla g(x_j) - \nabla_x f(x_j, y_j)\|^2$,

$$\begin{aligned}
R_{j+1} \leq & a_j - \left\{ \frac{\tau_1}{2}(1 + \lambda\zeta) - 2 \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - 2\lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right\} \mathbb{E}\|\nabla g(x_j)\|^2 + \\
& \lambda\zeta b_j + \left\{ \frac{\tau_1}{2}(1 + \lambda\zeta) + 2 \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - 2\lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right\} \mathbb{E}\|\nabla_x f(x_j, y_j) - \nabla g(x_j)\|^2 + \\
& \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] (1 + \tau_1 \beta_1) \mathbb{E}\|x_j - \tilde{x}\|^2 + \left[d_{j+1}(1 + \tau_2 \beta_2) + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \mathbb{E}\|y_j - \tilde{y}\|^2 + \\
& \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 + \lambda\zeta \frac{L+l}{2} \right] \tau_1^2 \mathbb{E}\|G_x(x_j, y_j) - \nabla_x f(x_j, y_j)\|^2. \tag{46}
\end{aligned}$$

Then plugging in (26), (27) and (42), we get

$$\begin{aligned}
R_{j+1} \leq & a_j - \left\{ \tau_1(1 + \lambda\zeta) - 4 \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - 4\lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right\} \mu_1 a_j + \\
& \lambda b_j - \lambda \frac{1}{\lambda} \left\{ \lambda\psi - \frac{l^2 \tau_1}{\mu_2}(1 + \lambda\zeta) - \frac{4l^2}{\mu_2} \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - \frac{4l^2}{\mu_2} \lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right\} b_j + \\
& \left\{ \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] (1 + \tau_1 \beta_1) + \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 + \lambda\zeta \frac{L+l}{2} \right] \tau_1^2 l^2 \right\} \mathbb{E}\|x_j - \tilde{x}\|^2 + \\
& \left\{ \left[d_{j+1}(1 + \tau_2 \beta_2) + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] + \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 + \lambda\zeta \frac{L+l}{2} \right] \tau_1^2 l^2 \right\} \mathbb{E}\|y_j - \tilde{y}\|^2. \tag{47}
\end{aligned}$$

Now we are ready to define sequences $\{c_j\}_j$ and $\{d_j\}_j$. Let $c_N = d_N = 0$, and

$$\begin{aligned}
c_j &= \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] (1 + \tau_1 \beta_1) + \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 + \lambda\zeta \frac{L+l}{2} \right] \tau_1^2 l^2, \\
d_j &= \left[d_{j+1}(1 + \tau_2 \beta_2) + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] + \left[\frac{L}{2} + c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 + \lambda\zeta \frac{L+l}{2} \right] \tau_1^2 l^2.
\end{aligned}$$

We further define

$$m_j^1 := \tau_1(1 + \lambda\zeta) - 4 \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - 4\lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right), \tag{48}$$

$$m_j^2 := \frac{1}{\lambda} \left\{ \lambda\psi - \frac{l^2 \tau_1}{\mu_2}(1 + \lambda\zeta) - \frac{4l^2}{\mu_2} \left[c_{j+1} + \left(d_{j+1} + \frac{\lambda l}{2} \right) l^2 \tau_2^2 \right] \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - \frac{4l^2}{\mu_2} \lambda\zeta \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \right\}. \tag{49}$$

Then we can write (47) as

$$R_{j+1} \leq R_j - m_j^1 a_j - \lambda m_j^2 b_j \quad (50)$$

Now we bring back the subscript t . Summing the equation from 0 to $N-1$,

$$\sum_{j=0}^{N-1} a_{t,j} + \lambda b_{t,j} \leq \frac{R_0 - R_N}{N\gamma} = \frac{a_{t,0} + \lambda b_{t,0} - a_{t,N} - \lambda b_{t,N}}{N\gamma} = \frac{\tilde{a}_t + \lambda \tilde{b}_t - \tilde{a}_{t+1} - \lambda \tilde{b}_{t+1}}{N\gamma}, \quad (51)$$

where $\gamma := \min_j \{m_j^1, m_j^2\}$, and the first equality is due to $c_N = d_N = 0$ and $(x_{t,0}, y_{t,0}) = (\tilde{x}_t, \tilde{y}_t)$. Summing t from 0 to $T-1$, we get

$$\frac{1}{NT} \sum_{t=0}^{T-1} \sum_{j=0}^{N-1} a_{t,j} + \lambda b_{t,j} \leq \frac{\tilde{a}_0 + \lambda \tilde{b}_0}{NT\gamma} = \frac{a^k + \lambda b^k}{NT\gamma}. \quad (52)$$

The left hand side is exactly $a^{k+1} + \lambda b^{k+1}$, because (x_k, y_k) is sampled uniformly from $\{(x_{t,j}, y_{t,j})\}_{j=0}^{N-1}\}_{t=0}^{T-1}$.

Part 2.

It suffices to choose proper τ_1, τ_2, N and T such that $NT\gamma > 1$. Driven by the proof, we choose

$$\tau_1 = \frac{k_1}{\kappa^2 l}, \quad \beta_1 = k_2 \kappa^2 l, \quad \tau_2 = \frac{k_3}{l}, \quad \beta_2 = l k_4.$$

We will choose k_1, k_2, k_3 and k_4 later and we let $k_1, k_2, k_3, k_4 \leq 1$. Plug back to c_j and d_j , we have

$$\begin{aligned} c_j &= \left(1 + k_1 k_2 + \frac{k_1^2}{\kappa^4}\right) c_{j+1} + \left[k_3^2(1 + k_1 k_2) + \frac{k_1^2 k_3^2}{\kappa^4} + (L + l) \frac{k_1^2}{\kappa^4} \left(\frac{k_3^2}{l^2} + \frac{k_3}{l^2 k_4}\right) \mu_2\right] d_{j+1} + \\ &\quad \frac{\lambda}{2} l k_3^2 (1 + k_1 k_2) + \frac{L}{2\kappa^4} k_1^2 + \frac{\lambda}{2\kappa^4} l k_1^2 k_3^2 + \frac{\lambda}{2\kappa^4} (L + l) k_1^2 (1 - k_3 k_4) \\ &\leq \left(1 + k_1 k_2 + \frac{k_1^2}{\kappa^4}\right) c_{j+1} + \left(3k_3^2 + 3\frac{1}{\kappa^3} k_1^2\right) d_{j+1} + 2\lambda l k_3^2 + (1 + 2\lambda) \frac{l}{\kappa^3} k_1^2, \end{aligned} \quad (53)$$

where in the last inequality we assume $k_3^2 + \frac{k_3}{k_4} \leq 1$.

$$\begin{aligned} d_j &= \frac{k_1^2}{\kappa^4} c_{j+1} + \left[1 + k_3 k_4 + k_3^2 + (L + l) \frac{k_1^2}{\kappa^4} \left(\frac{k_3^2}{l^2} + \frac{k_3}{l^2 k_4}\right) \mu_2 + \frac{1}{\kappa^4} k_1^2 k_3^2\right] d_{j+1} + \\ &\quad \frac{\lambda}{2} l k_3^2 + \frac{L}{2\kappa^4} k_1^2 + \frac{\lambda}{2\kappa^4} l k_1^2 k_3^2 + \frac{\lambda}{2\kappa^4} (L + l) k_1^2 (1 - k_3 k_4) \\ &\leq \frac{k_1^2}{\kappa^4} c_{j+1} + \left(1 + k_3 k_4 + 2k_3^2 + \frac{3}{\kappa^3} k_1^2\right) d_{j+1} + \lambda l k_3^2 + (1 + 2\lambda) \frac{l}{\kappa^3} k_1^2. \end{aligned} \quad (54)$$

We define $e_j = \max\{c_j, d_j\}$. Then combining (53) and (54), we easily get

$$e_j \leq \left(1 + k_1 k_2 + k_3 k_4 + 3k_3^2 + \frac{4}{\kappa^3} k_1^2\right) e_{j+1} + 2\lambda l k_3^2 + (1 + 2\lambda) \frac{l}{\kappa^3} k_1^2.$$

As $e_N = 0$, we have

$$e_0 \leq \left[2\lambda l k_3^2 + (1 + 2\lambda) \frac{l}{\kappa^3} k_1^2\right] \frac{\left(1 + k_1 k_2 + k_3 k_4 + 3k_3^2 + \frac{4}{\kappa^3} k_1^2\right)^N - 1}{k_1 k_2 + k_3 k_4 + 3k_3^2 + \frac{4}{\kappa^3} k_1^2}, \quad (55)$$

and note that $e_j > e_{j+1}$ so $e_j \leq e_0, \forall j$. Then we want to lower bound γ . Rearrange (48),

$$\begin{aligned} m_j^1 = & \mu_1 \left\{ \tau_1 (1 + \lambda - \lambda \tau_2 \mu_2) - 2\lambda l^3 \tau_2^2 \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) - 4\lambda \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) (1 - \tau_2 \mu_2) - \right. \\ & \left[-2\tau_1 \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 + 4 \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) l^2 \tau_2^2 + 8 \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 \right] d_{j+1} - \\ & \left. 4 \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) c_{j+1} \right\} \\ \geq & \frac{1}{2} \tau_1 \mu_1 - \left[\frac{4}{\kappa^4} k_3^2 \left(k_1^2 + \frac{k_1}{k_2} \right) + \frac{10\mu_2}{\kappa^2 l} k_1 \left(k_3^2 + \frac{k_3}{k_4} \right) \right] \frac{\mu_1}{l^2} d_{j+1} - \frac{4}{\kappa^4} \left(k_1^2 + \frac{k_1}{k_2} \right) \frac{\mu_1}{l^2} c_{j+1}, \end{aligned} \quad (56)$$

where in the inequality, we use $\lambda = 1/20$ and assume that $\frac{1}{\kappa^2} k_3^2 (k_1 + \frac{1}{k_2}) \leq 10$. Rearranging (49),

$$\begin{aligned} m_j^2 = & \tau_2 \mu_2 - \frac{l^2 \tau_1}{\mu_2} \left(\frac{1}{\lambda} + 1 - \tau_2 \mu_2 \right) - \frac{2l^5}{\mu_2} \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) \tau_2^2 - \frac{4l^2}{\mu_2} \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) (1 - \tau_2 \mu_2) - \\ & \left[\frac{2}{\lambda} \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 + \frac{2}{\lambda} l^2 \tau_1 \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) + \frac{4}{\lambda} \frac{l^4}{\mu_2} \tau_2^2 \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) + \frac{8l^2}{\lambda \mu_2} \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \left(\tau_2^2 + \frac{\tau_2}{\beta_2} \right) \mu_2 \right] d_{j+1} - \\ & \frac{4}{\lambda} \frac{l^2}{\mu_2} \left(\tau_1^2 + \frac{\tau_1}{\beta_1} \right) c_{j+1} \\ \geq & \frac{l^2 \tau_1}{2 \min\{\mu_1, \mu_2\}} - \left[200 \left(k_3^2 + \frac{k_3}{k_4} \right) + \frac{80}{\kappa^2} \left(k_1^2 + \frac{k_1}{k_2} \right) \right] \frac{\mu_2}{l^2} d_{j+1} - \frac{80}{\kappa^2} \left(k_1^2 + \frac{k_1}{k_2} \right) \frac{\mu_2}{l^2} c_{j+1}, \end{aligned} \quad (57)$$

where in the inequality we use $\lambda = 1/20$ and assume $k_1 \leq k_3/28$ and $\frac{1}{\kappa^2} k_3^2 (k_1 + \frac{1}{k_2}) \leq 1/4$. Note that $\frac{1}{2} \tau_1 \mu_1 = \frac{\mu_1}{2\kappa^2 l} k_1$ and $\frac{l^2 \tau_1}{2 \min\{\mu_1, \mu_2\}} = \frac{l}{2\kappa^2 \min\{\mu_1, \mu_2\}} k_1$. Then we have

$$m_j^1 \geq \frac{1}{\kappa^3} \left\{ \frac{1}{2} k_1 - \left[\frac{4}{\kappa^2} k_3^2 \left(k_1^2 + \frac{k_1}{k_2} \right) + \frac{10\mu_2}{l} k_1 \left(k_3^2 + \frac{k_3}{k_4} \right) \right] \frac{d_{j+1}}{l} - \frac{4}{\kappa^2} \left(k_1^2 + \frac{k_1}{k_2} \right) \frac{c_{j+1}}{l} \right\}, \quad (58)$$

$$m_j^2 \geq \frac{1}{\kappa} \left\{ \frac{1}{2} k_1 - \left[\frac{80}{\kappa^2} \left(k_1^2 + \frac{k_1}{k_2} \right) + 200 \left(k_3^2 + \frac{k_3}{k_4} \right) \right] \frac{d_{j+1}}{l} - \frac{80}{\kappa^2} \left(k_1^2 + \frac{k_1}{k_2} \right) \frac{c_{j+1}}{l} \right\}. \quad (59)$$

Letting $k_1/k_2 = k_3/k_4$ and $k_1 = \frac{1}{28} k_3$, we have

$$\gamma \geq \frac{1}{\kappa^3} \left\{ \frac{1}{56} k_3 - 360 \left(k_3^2 + \frac{k_3}{k_4} \right) \frac{e_0}{l} \right\}, \quad (60)$$

where we use $c_j, d_j \leq e_0, \forall j$. By plugging in $k_1 = k_3/28$ and $\lambda = 1/20$ into (55), we have

$$e_0 \leq l \frac{(1 + 2k_3 k_4 + 4k_3^2)^N - 1}{k_4/k_3 + 3}. \quad (61)$$

Plugging this into (60), we have

$$\gamma \geq \frac{1}{\kappa^3} \left[\frac{k_3}{56} - 360 \frac{(1 + 2k_3 k_4 + 4k_3^2)^N - 1}{k_4/k_3 + 3} (k_3^2 + k_3/k_4) \right]. \quad (62)$$

We choose $k_4 = k_3^{1/2}$, then

$$NT\gamma \geq \frac{1}{\kappa^3} \left[\frac{k_3}{56} - 360 \left((1 + 2k_3^{3/2} + 4k_3^2)^N - 1 \right) \left(\frac{k_3^2 + k_3^{1/2}}{k_3^{-1/2} + 3} \right) \right] NT. \quad (63)$$

Part 3.

We choose $T = 1$, $k_3 = \beta\kappa^{-6}$ and $N = \alpha(2k_3^{3/2} + 4k_3^2)^{-1} \geq \frac{\alpha}{2}k_2^{-3/2}$, where α, β is irrelevant to n, l, μ_1, μ_2 . Then since $(1 + 2k_3^{3/2} + 4k_3^2)^N \leq e^\alpha$, after plugging in N and k_3 , we have

$$NT\gamma \geq \frac{1}{\kappa^3} \left[\frac{k_3}{56} - 360(e^\alpha - 1)(2k_3) \right] \frac{\alpha}{2} k_2^{-3/2} \geq \frac{1}{2} \left[\frac{1}{56} - 2 \times 360(e^\alpha - 1) \right] \alpha \beta^{-1/2}. \quad (64)$$

Therefore, for choosing α small enough and β small enough, we have $NT\gamma \geq 2$. Now it remains to verify several assumptions we made in the proof. The first is $\frac{k_3}{k_4} + k_3^2 \leq 1$. Since $\frac{k_3}{k_4} + k_3^2 = k_3^{1/2} + k_3^2$, this assumption easily holds when $\beta \leq 1/4$. The second assumption we want to verify is $\frac{1}{\kappa^2} k_3^2 \left(k_1 + \frac{1}{k_2} \right) \leq 1/4$. Note that

$$\frac{1}{\kappa^2} k_3^2 \left(k_1 + \frac{1}{k_2} \right) = \frac{1}{\kappa^2} k_3^2 \left(k_1 + \frac{k_3}{k_4 k_1} \right) = \frac{1}{\kappa^2} k_3^2 \left(\frac{1}{28} k_3 + 28k_3^{-1/2} \right).$$

So this assumption can also be easily satisfied when β is small. The last assumption we need to verify is $\lambda \geq d_{j+1} \left(\tau_2 + \frac{1}{\beta_2} \right)$. Because $d_{j+1} \leq e_0$ and (61),

$$\begin{aligned} d_{j+1} \left(\tau_2 + \frac{1}{\beta_2} \right) &\leq l \frac{(1 + 2k_3 k_4 + 4k_3^2)^N - 1}{k_4/k_3 + 3} \left(\frac{k_3}{l} + \frac{1}{k_4 l} \right) \\ &\leq ((1 + 2k_3 k_4 + 4k_3^2)^N - 1) \left(\frac{k_3^2 + k_3^{1/2}}{k_3^{-1/2} + 3} \right) \\ &\leq 2(e^\alpha - 1)k_3. \end{aligned}$$

So this assumption holds when α and β are small. □

Proof of Theorem 4.2

Proof. We start from Part 3 of the proof of Theorem 4.1. We now choose $k_3 = \beta n^{-2/3}$, $N = \alpha(2k_3^{3/2} + 4k_3^2)^{-1}$, and $T = \kappa^3 n^{-1/3}$ then

$$NT\gamma \geq \frac{1}{2} \left[\frac{1}{56} - 2 \times 360(e^\alpha - 1) \right] \alpha \beta^{-1/2} \quad (65)$$

Therefore, for choosing α small enough and β small enough, we have $NT\gamma \geq 2$. Other assumptions can be easily verified by the same way as in the proof of Theorem 4.1. □

D AGDA for minimax problems under one-sided PL condition

We are here to show that if $-f(x, \cdot)$ satisfies PL condition with constant μ and $f(\cdot, y)$ may be nonconvex (referred to as PL game by Nouiehed et al. [2019]), AGDA as presented in Algorithm 3 can find ϵ -stationary point of $g(x) := \max_y f(x, y)$ within $\mathcal{O}(\epsilon^{-2})$ iterations. Note that GDmax has complexity $\mathcal{O}(\epsilon^{-2} \log(1/\epsilon))$ on minimax problems under the one-sided PL condition [Nouiehed et al., 2019]; SGDA has complexity $\mathcal{O}(\epsilon^{-2})$ on nonconvex-strongly-concave minimax problems [Lin et al., 2019]. Here we define condition number $\kappa = \frac{\mu}{l}$ and L is still defined the same as before. The proof is based on our previous analysis and Lin et al. [2019].

Definition 3. x is ϵ -stationary point of a differential function f if $\mathbb{E} \|\nabla f(x)\| \leq \epsilon$.

Algorithm 3 AGDA

1: Input: (x_0, y_0) , step sizes $\tau_1 > 0, \tau_2^t > 0$
2: **for all** $t = 0, 1, 2, \dots, T-1$ **do**
3: $x_{t+1} \leftarrow x_t - \tau_1 \nabla f_x(x_t, y_t)$
4: $y_{t+1} \leftarrow y_t + \tau_2 \nabla f_y(x_{t+1}, y_t)$
5: **end for**
6: choose (x^T, y^T) uniformly from $\{(x_t, y_t)\}_{t=0}^T$

Theorem D.1. Suppose Assumption 1 holds and $-f(x, \cdot)$ satisfies PL condition with constant μ for any x . If we run Algorithm 3 with $\tau_1 = \frac{1}{20\kappa^2 l}$ and $\tau_2 = \frac{1}{l}$, then

$$\mathbb{E} \|\nabla g(x^T)\|^2 \leq \frac{8}{T+1} [10\kappa^2 l a_0 + \kappa^2 l b_0], \quad (66)$$

where $a_0 = g(x_0) - g^*$ and $b_0 = g(x_0) - f(x_0, y_0)$.

Proof. For convenience, we still define $b_t = g(x_t) - f(x_t, y_t)$. Since it can be easily verified that $\tau_1 \leq 1/L$, by (18) and (26), we have

$$g(x_{t+1}) \leq g(x_t) - \frac{\tau_1}{2} \|\nabla g(x_t)\|^2 + \frac{\tau_1 l^2}{\mu_2} b_t. \quad (67)$$

By (22), we have

$$\begin{aligned} b_{t+1} &\leq (1 - \mu_2 \tau_2) b_t + (1 - \mu_2 \tau_2) \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) \|\nabla_x f(x_t, y_t)\|^2 - \\ &\quad (1 - \mu_2 \tau_2) \frac{\tau_1}{2} \|\nabla g(x_t)\|^2 + (1 - \mu_2 \tau_2) \frac{\tau_2}{2} \|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 \\ &\leq (1 - \mu_2 \tau_2) b_t + \left[2(1 - \mu_2 \tau_2) \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) - (1 - \mu_2 \tau_2) \frac{\tau_2}{2} \right] \|\nabla g(x_t)\|^2 + \\ &\quad \left[2(1 - \mu_2 \tau_2) \left(\tau_1 + \frac{l}{2} \tau_1^2 \right) + (1 - \mu_2 \tau_2) \frac{\tau_2}{2} \right] \|\nabla_x f(x_t, y_t) - \nabla g(x_t)\|^2 \\ &\leq (1 - \mu_2 \tau_2) \left[1 + (5\tau_1 + 2l\tau_1^2) \frac{l^2}{\mu_2} \right] b_t + (1 - \mu_2 \tau_2) \left[\frac{3}{2} \tau_1 + l\tau_1^2 \right] \|\nabla g(x_t)\|^2, \end{aligned} \quad (68)$$

where in the second inequality we use Young's inequality, and in third inequality we use (26). We write

$$b_{t+1} = \alpha b_t + \beta \|\nabla g(x_k)\|^2 \quad (69)$$

with

$$\alpha = (1 - \mu_2 \tau_2) \left[1 + (5\tau_1 + 2l\tau_1^2) \frac{l^2}{\mu_2} \right], \quad \beta = (1 - \mu_2 \tau_2) \left[\frac{3}{2} \tau_1 + l\tau_1^2 \right].$$

Then

$$b_t \leq \alpha^t b_0 + \beta \sum_{k=0}^{t-1} \alpha^{t-1-k} \|\nabla g(x_k)\|^2, \quad t \geq 1.$$

Plugging into (67), we have

$$g(x_{t+1}) \leq g(x_t) - \frac{\tau_1}{2} \|\nabla g(x_t)\|^2 + \frac{\tau_1 l^2}{\mu_2} \alpha^t b_0 + \frac{\tau_1 l^2 \beta}{\mu_2} \sum_{k=0}^{t-1} \alpha^{t-1-k} \|\nabla g(x_k)\|^2, \quad t \geq 1. \quad (70)$$

Telescoping and rearranging,

$$\frac{\tau_1}{2} \sum_{t=0}^T \|\nabla g(x_t)\|^2 - \frac{\tau_1 l^2 \beta}{\mu_2} \sum_{t=1}^T \sum_{k=0}^{t-1} \alpha^{t-1-k} \|\nabla g(x_k)\|^2 \leq g(x_0) - g(x_{T+1}) + \frac{\tau_1 l^2}{\mu_2} b_0 \sum_{t=0}^T \alpha^t \leq a_0 + \frac{\tau_1 l^2}{\mu_2(1-\alpha)} b_0 \quad (71)$$

Considering the left hand side of (71),

$$\sum_{t=1}^T \sum_{k=0}^{t-1} \alpha^{t-1-k} \|\nabla g(x_k)\|^2 = \sum_{k=0}^{T-1} \sum_{t=k+1}^T \alpha^{t-1-k} \|\nabla g(x_k)\|^2 \leq \sum_{k=0}^{T-1} \frac{1}{1-\alpha} \|\nabla g(x_k)\|^2, \quad (72)$$

and therefore,

$$\frac{\tau_1}{2} \sum_{t=0}^T \|\nabla g(x_t)\|^2 - \frac{\tau_1 l^2 \beta}{\mu_2} \sum_{t=0}^T \sum_{k=0}^{t-1} \alpha^{t-1-k} \|\nabla g(x_k)\|^2 \geq \sum_{t=0}^T \left\{ \frac{1}{2} - \frac{l^2 \beta}{\mu_2(1-\alpha)} \right\} \tau_1 \|\nabla g(x_t)\|^2. \quad (73)$$

We note that $\beta = (1 - \mu_2 \tau_2) \left[\frac{3}{2} \tau_1 + l \tau_1^2 \right] \leq \frac{5}{2} \tau_1$ because $l/\tau_1 \leq 1$ by our choice of τ_1 . Also,

$$1 - \alpha = \mu_2 \tau_2 - (1 - \mu_2 \tau_2) (5\tau_1 + 2l\tau_1^2) \frac{l^2}{\mu_2} \geq \mu_2 \tau_2 - 7(1 - \mu_2 \tau_2) \frac{\tau_1 l^2}{\mu_2} \geq \frac{1}{2\kappa}, \quad (74)$$

where in the last inequality we use $\mu_2 \tau_2 = 1/\kappa$ and $(1 - \mu_2 \tau_2) \frac{\tau_1 l^2}{\mu_2} = (1 - 1/\kappa)/(20\kappa) \leq 1/(20\kappa)$. Plugging into (73),

$$\frac{\tau_1}{2} \sum_{t=0}^T \|\nabla g(x_t)\|^2 - \frac{\tau_1 l^2 \beta}{\mu_2} \sum_{t=1}^T \sum_{k=0}^{t-1} \alpha^{t-1-k} \|\nabla g(x_k)\|^2 \geq \frac{\tau_1}{4} \sum_{t=0}^T \|\nabla g(x_t)\|^2. \quad (75)$$

Combining with (71), we have

$$\frac{1}{T+1} \sum_{t=0}^T \|\nabla g(x_t)\|^2 \leq \frac{4}{(T+1)\tau_1} \left[a_0 + \frac{\tau_1 l^2}{\mu_2(1-\alpha)} b_0 \right] \leq \frac{8}{T+1} [10\kappa^2 l a_0 + \kappa^2 l b_0], \quad (76)$$

where in the inequality we use $1 - \alpha \geq 1/(2\kappa)$ again.

□