

Handling the Positive-Definite Constraint in the Bayesian Learning Rule

Wu Lin¹ Mark Schmidt^{1,2} Mohammad Emtiyaz Khan³

Abstract

The Bayesian learning rule is a natural-gradient variational inference method, which not only contains many existing learning algorithms as special cases but also enables the design of new algorithms. Unfortunately, when variational parameters lie in an open constraint set, the rule may not satisfy the constraint and requires line-searches which could slow down the algorithm. In this work, we address this issue for positive-definite constraints by proposing an improved rule that naturally handles the constraints. Our modification is obtained by using Riemannian gradient methods, and is valid when the approximation attains a *block-coordinate natural parameterization* (e.g., Gaussian distributions and their mixtures). We propose a principled way to derive Riemannian gradients and retractions from scratch. Our method outperforms existing methods without any significant increase in computation. Our work makes it easier to apply the rule in the presence of positive-definite constraints in parameter spaces.

1. Introduction

The Bayesian learning rule, a recently proposed method, enables derivation of learning algorithms from Bayesian principles (Khan & Rue, 2020). It is a natural-gradient variational inference method (Khan & Lin, 2017) where, by carefully choosing a posterior approximation, we can derive many algorithms in fields such as probabilistic graphical models, continuous optimization, and deep learning. Khan & Lin (2017) derive approximate inference methods, such as stochastic variational inference and variational message passing; Khan et al. (2018) derive connections to deep-learning algorithms; and Khan & Rue (2020) derive many classical algorithms such as least-squares, gradient descent,

Newton’s method, and the forward-backward algorithm. We can also design new algorithms using this rule such as uncertainty estimation in deep learning (Osawa et al., 2019) and the ensemble of Newton methods (Lin et al., 2019a).

An issue with the rule is that when parameters of a posterior approximation lie in an open constraint set, the update may not always satisfy the constraints. For Gaussian approximations, the posterior covariance needs to be positive definite but the rule may violate this; see Appendix D.1 in Khan et al. (2018) for detail. A straightforward solution is to use a backtracking line-search to keep the updates within the constraint set (Khan & Lin, 2017), but this can lead to slow convergence. In some cases, we can find an approximate update which always satisfies the constraints, e.g., for Gaussian approximations (Khan et al., 2018). However, in general, it is difficult to come up with such approximations that are both fast and reasonably accurate. Our goal in this paper is to modify the Bayesian learning rule so that it can naturally handle such constraints.

We propose an improved Bayesian learning rule to handle the positive-definite constraints. This is obtained by using a generalization of natural-gradient methods called Riemannian-gradient methods. We show that, for many useful approximations with a specific block-diagonal structure on the Fisher information matrix, the constraints are satisfied after an additional term is added to the rule. Such a structure is possible when the parameters of the approximation are partitioned in what we call the *block-coordinate natural (BCN) parameterizations*. Fortunately, for many approximations with such parameterizations, the improved rule requires almost the same computation as the original rule. An example is shown in Figure 1 where our improved rule fixes an implementation issue with an algorithm proposed by Osawa et al. (2019) for deep learning. We present examples where the improved rule converges faster than the original rule and many existing baseline methods.

2. Bayesian Learning Rule

Given a dataset \mathcal{D} , it is common to estimate unknown variables \mathbf{z} of a statistical model by minimizing¹ $\bar{\ell}(\mathbf{z}) \equiv \ell(\mathcal{D}, \mathbf{z}) + R(\mathbf{z})$ where $\ell(\mathcal{D}, \mathbf{z})$ is a loss function and $R(\mathbf{z})$

¹University of British Columbia, Vancouver, Canada. ²CIFAR AI Chair, Alberta Machine Intelligence Institute, Canada. ³RIKEN Center for Advanced Intelligence Project, Tokyo, Japan. Correspondence to: Wu Lin <wlin2018@cs.ubc.ca>.

¹We assume $\nabla_{\mathbf{z}} \bar{\ell}(\mathbf{z})$ and $\nabla_{\mathbf{z}}^2 \bar{\ell}(\mathbf{z})$ exist almost surely whenever they are needed. ∇ denotes the standard derivative in this paper.

Variational Online Gauss-Newton (VOGN) Algorithm

- 1: $\mathbf{z} \leftarrow \boldsymbol{\mu} + (N\hat{\mathbf{s}})^{-1/2} \odot \boldsymbol{\epsilon}$, where $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$
- 2: Randomly sample a minibatch \mathcal{M} of size M
- 3: **Compute and store individual gradients $\mathbf{g}_i, \forall i \in \mathcal{M}$**
- 4: $\mathbf{g}_\mu \leftarrow \frac{\lambda}{N} \boldsymbol{\mu} + \frac{1}{M} \sum_{i=1}^M \mathbf{g}_i$
- 5: $\mathbf{m} \leftarrow r_1 \mathbf{m} + (1 - r_1) \mathbf{g}_\mu$, $\bar{\mathbf{m}} \leftarrow \mathbf{m} / (1 - r_1^k)$
- 6: $\mathbf{g}_s \leftarrow \frac{\lambda}{N} - \hat{\mathbf{s}} + \frac{1}{M} \sum_{i=1}^M (\mathbf{g}_i \odot \mathbf{g}_i)$
- 7: $\hat{\mathbf{s}} \leftarrow \hat{\mathbf{s}} + (1 - r_2) \mathbf{g}_s$
- 8: $\boldsymbol{\mu} \leftarrow \boldsymbol{\mu} - t \bar{\mathbf{m}} / \bar{\mathbf{s}}$, where $\bar{\mathbf{s}} \leftarrow \hat{\mathbf{s}} / (1 - r_2^k)$

Our Adam-like Optimizer

- 1: $\mathbf{z} \leftarrow \boldsymbol{\mu} + (N\hat{\mathbf{s}})^{-1/2} \odot \boldsymbol{\epsilon}$, where $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$
- 2: Randomly sample a minibatch \mathcal{M} of size M
- 3: **Compute a mini-batch gradient $\bar{\mathbf{g}} \leftarrow \frac{1}{M} \sum_{i=1}^M \mathbf{g}_i$**
- 4: $\mathbf{g}_\mu \leftarrow \frac{\lambda}{N} \boldsymbol{\mu} + \bar{\mathbf{g}}$
- 5: $\mathbf{m} \leftarrow r_1 \mathbf{m} + (1 - r_1) \mathbf{g}_\mu$, $\bar{\mathbf{m}} \leftarrow \mathbf{m} / (1 - r_1^k)$
- 6: $\mathbf{g}_s \leftarrow \frac{\lambda}{N} - \hat{\mathbf{s}} + [(N\hat{\mathbf{s}}) \odot (\mathbf{z} - \boldsymbol{\mu})] \odot \bar{\mathbf{g}}$
- 7: $\boldsymbol{\mu} \leftarrow \boldsymbol{\mu} - t \bar{\mathbf{m}} / \bar{\mathbf{s}}$, where $\bar{\mathbf{s}} \leftarrow \hat{\mathbf{s}} / (1 - r_2^k)$
- 8: $\hat{\mathbf{s}} \leftarrow \hat{\mathbf{s}} + (1 - r_2) \mathbf{g}_s + \frac{1}{2} (1 - r_2)^2 \mathbf{g}_s \odot \hat{\mathbf{s}}^{-1} \odot \mathbf{g}_s$

Figure 1. Our improved Bayesian learning rule solves an implementation issue with an existing algorithm known as VOGN (Khan et al., 2018) (shown in the left). VOGN is an Adam-like optimizer which gives state-of-the-art results on large deep learning problems (Osawa et al., 2019). However, it requires us to store individual gradients in a minibatch which makes the algorithm slow (shown with blue in line 3 and 6). This is necessary for the scaling vector $\hat{\mathbf{s}}$ to obtain a good estimate of uncertainty. Our work in this paper fixes this issue using the improved Bayesian learning rule. Our Adam-like optimizer (shown in the right) only requires average over the minibatch (see line 3). Line 6 is simply changed to use the re-parametrization trick with the averaged gradient. The additional terms added to the Bayesian learning rule is shown in red in line 8. These changes do not increase the computation cost significantly while fixing the implementation issue of VOGN. Due to our modification, the scaling vector $\hat{\mathbf{s}}$ always remains positive. A small difference is that the mean $\boldsymbol{\mu}$ is updated before in our optimizer (see line 7 and 8), while in VOGN it is the opposite. The difference shows that NGD depends on parameterization.

is a regularizer. Many estimation strategies can be used, giving rise to various learning algorithms. E.g., maximum-likelihood approaches use gradient-based methods such as gradient descent and Newton’s method, while Bayesian approaches use inference algorithms such as message passing.

Khan & Rue (2020) showed that many learning algorithms can be obtained from Bayesian principles. The key idea is to use the following Bayesian formulation where, instead of minimizing over \mathbf{z} , we minimize over a distribution $q(\mathbf{z})$:

$$\min_{q(\mathbf{z}) \in \mathcal{Q}} \mathbb{E}_{q(\mathbf{z})}[\ell(\mathcal{D}, \mathbf{z})] + \mathbb{D}_{KL}[q(\mathbf{z}) \parallel p(\mathbf{z})] \equiv \mathcal{L}(q). \quad (1)$$

Here, $q(\mathbf{z})$ is an approximation of the posterior of \mathbf{z} given \mathcal{D} , \mathcal{Q} is the set of approximation distributions, $p(\mathbf{z}) \propto \exp(-R(\mathbf{z}))$ is the prior, and \mathbb{D}_{KL} denotes the Kullback-Leibler divergence. To obtain existing learning algorithms from the above formulation, we need to carefully choose the approximation family \mathcal{Q} . Khan & Rue (2020) consider the following *minimal* exponential family (EF) distribution:

$$q(\mathbf{z}|\boldsymbol{\lambda}) := h(\mathbf{z}) \exp[\langle \phi(\mathbf{z}), \boldsymbol{\lambda} \rangle - A(\boldsymbol{\lambda})]$$

where $\phi(\mathbf{z})$ is a vector containing sufficient statistics, $h(\mathbf{z})$ is the base measure, $\boldsymbol{\lambda} \in \Omega$ is the natural parameter, Ω is the set of valid natural-parameters so that the log-partition function $A(\boldsymbol{\lambda})$ is finite, and $\langle \cdot, \cdot \rangle$ denotes an inner product.

Khan & Rue (2020) present the Bayesian learning rule to optimize (1), which is a natural-gradient descent (NGD) update originally proposed by Khan & Lin (2017) for variational inference. The update takes the following form:

$$\text{NGD: } \boldsymbol{\lambda} \leftarrow \boldsymbol{\lambda} - t \hat{\mathbf{g}}, \text{ with } \hat{\mathbf{g}} := \mathbf{F}^{-1} \partial_{\boldsymbol{\lambda}} \mathcal{L}(\boldsymbol{\lambda}) \quad (2)$$

where $t > 0$ is a scalar step-size and $\hat{\mathbf{g}}$ is the natural gradient defined using the Fisher information matrix (FIM) $\mathbf{F} :=$

$-\mathbb{E}_q[\partial_{\boldsymbol{\lambda}}^2 \log q(\mathbf{z}|\boldsymbol{\lambda})]$ of q and $\mathcal{L}(\boldsymbol{\lambda})$ which is equal to $\mathcal{L}(q)$ but defined in terms of $\boldsymbol{\lambda}$. Khan & Rue (2020) proposed further simplifications, e.g., for approximations with base measure $h(\mathbf{z}) \equiv 1$, we can write (2) as

$$\boldsymbol{\lambda} \leftarrow (1 - t) \boldsymbol{\lambda} - t \partial_m \mathbb{E}_q[\bar{\ell}(\mathbf{z})] \quad (3)$$

where $\mathbf{m} := \mathbb{E}_{q(\mathbf{z})}[\phi(\mathbf{z})]$ denotes the expectation parameter.

Existing learning algorithms can be derived as special cases by choosing an approximate form for $q(\mathbf{z})$. For example, when $q(\mathbf{z}) := \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}, \mathbf{S}^{-1})$ is a multivariate Gaussian approximation with the mean $\boldsymbol{\mu}$ and the precision matrix \mathbf{S} , the learning rule (3) can be expressed as follows:

$$\mathbf{S} \leftarrow (1 - t) \mathbf{S} + t \mathbb{E}_q[\nabla_{\mathbf{z}}^2 \bar{\ell}(\mathbf{z})] \quad (4)$$

$$\boldsymbol{\mu} \leftarrow \boldsymbol{\mu} - t \mathbf{S}^{-1} \mathbb{E}_q[\nabla_{\mathbf{z}} \bar{\ell}(\mathbf{z})] \quad (5)$$

This algorithm uses the Hessian to update \mathbf{S} which is then used to scale the update for $\boldsymbol{\mu}$, in a similar fashion as Newton’s method. The main difference here is that the gradient and Hessian are obtained at samples from $q(\mathbf{z})$ instead of the current iterate $\boldsymbol{\mu}$. Khan & Rue (2020) approximate the expectation at $\boldsymbol{\mu}$ to obtain an online Newton method. This algorithm is closely related to deep-learning optimizers, such as, RMSprop and Adam (Khan et al., 2018; Zhang et al., 2018). A simplified version of this algorithm obtains state-of-the-art results on large deep-learning problems for uncertainty estimation as shown by Osawa et al. (2019).

Many other examples are discussed in Khan & Rue (2020), including algorithms such as stochastic gradient descent. The relationship to message passing algorithms and stochastic variational inference is shown in Khan & Lin (2017). In summary, the Bayesian learning rule is a generic learning rule that can be used not only to derive existing algorithms, but also to improve them and design new ones.

2.1. Positive-Definite Constraints

An issue with updates (2) and (3) is that the constraint $\lambda \in \Omega$ is not taken into account, where Ω is the set of valid parameters. The update is valid when Ω is unconstrained (e.g., a Euclidean space), but otherwise it may violate the constraint. An example is the multivariate Gaussian of dimension d where the precision matrix $\mathbf{S} \in \mathbb{S}_{++}^{d \times d}$ is required to be real and positive-definite, while the mean $\mu \in \mathbb{R}^d$ is unconstrained. In such cases, the update may violate the constraint. E.g., in the update (4), \mathbf{S} can be indefinite, when the loss $\bar{\ell}(\mathbf{z})$ is nonconvex. A similar issue appears when flexible approximations are used such as Gaussian mixtures.

Another example is a gamma distribution: $q(z|\alpha, \beta) \propto z^{\alpha-1}e^{-z\beta}$ where both $\alpha, \beta > 0$. We denote the positivity constraint using \mathbb{S}_{++}^1 . The rule takes the following form:

$$\alpha \leftarrow (1-t)\alpha - t\hat{g}_\alpha, \quad \beta \leftarrow (1-t)\beta - t\hat{g}_\beta \quad (6)$$

where \hat{g}_α and \hat{g}_β are gradient of $\mathbb{E}_{q(z)}[\bar{\ell}(z) - \log z]$ with respect to the expectation parameters $m_\alpha = \mathbb{E}_{q(z)}[\log z]$ and $m_\beta = \mathbb{E}_{q(z)}[-z]$ respectively; see a detailed derivation in Appendix E.3 in Khan & Lin (2017). Here again the learning rule does not ensure that α and β are always positive.

In general, a backtracking line search proposed in Khan & Lin (2017) can be used so that the iterates stay within the constraint set. However, this could be slow in practice. Khan et al. (2018) discuss this issue for the Gaussian case; see Appendix D.1 in their paper. They found that using line-search is computationally expensive and non-trivial to implement for deep-learning problems. They address this issue by approximating the Hessian in (4) with a positive-definite matrix. This ensures that \mathbf{S} is always positive-definite. However, such approximations are difficult to come up with for general cases. E.g., for the gamma case, there is no such straight-forward approximation in update (6) to ensure positivity of α and β . It is also possible to use an unconstrained transformation (e.g., a Cholesky factor). This approach uses automatic-differentiation (Auto-Diff), which can be much slower than explicit gradient forms (see the discussion in Section 4). Handling constraints within the Bayesian learning rule is an open issue which limits its applications.

In this paper, we focus on positive-definite constraints and show that, in many cases, such constraints can be naturally handled by adding an additional term to the Bayesian learning rule. We show that, for this to happen, the approximation needs to follow a specific parameterization. We will now describe the modification in the next section, and later give its derivation using Riemannian gradient methods.

3. Improved Bayesian Learning Rule

We will give a new rule to handle the positive-definite constraints. Our idea is to partition the parameter into blocks

so that each constraint is isolated in an individual block.

Assumption 1 [Mutually-Exclusive Constraints] : We assume parameter $\lambda = \{\lambda^{[1]}, \dots, \lambda^{[m]}\}$ can be partitioned into m blocks with mutually-exclusive constraints $\Omega = \Omega_1 \times \dots \times \Omega_m$, where square bracket $[i]$ denotes the i -th block and each block $\lambda^{[i]}$ is either unconstrained or positive-definite.

For example, consider multivariate Gaussian approximations with the two blocks: one block containing the mean μ and another containing the full precision \mathbf{S} . This satisfies the above assumption because the first block is unconstrained and the second block is positive definite. In d -dimensional diagonal Gaussian cases, we consider $2d$ blocks: one block containing the mean μ_i and one block containing the precision s_i for each dimension i , where each s_i is positive. Other examples such as gammas and inverse Gaussians can be partitioned to two blocks, where each block is positive.

Assumption 2 [Block Coordinate Parameterization] : A parameterization satisfied Assumption 1 is block coordinate (BC) if the FIM is block-diagonal according to the block structure of the parameterization.

For Gaussians, using the mean and the covariance/precision as two blocks is a BC parameterization (see Appendix E), while the natural parameterization is not (Malagò & Pistone, 2015). For EFs, we could use the Crouzeix identity (Nielsen, 2019) to identify a BC parameterization.

Assumption 3 [Block Natural Parameterization for EF] : For $q(\mathbf{z}|\lambda)$ and each block $\lambda^{[i]}$, there exist function ϕ_i and h_i such that $q(\mathbf{z}|\lambda)$ can be re-expressed as a minimal EF distribution given that the rest of blocks $\lambda^{[-i]}$ are known.

$$q(\mathbf{z}|\lambda) \equiv h_i(\mathbf{z}, \lambda^{[-i]}) \exp \left[\left\langle \phi_i(\mathbf{z}, \lambda^{[-i]}), \lambda^{[i]} \right\rangle - A(\lambda) \right]$$

Lin et al. (2019a) originally use Assumption 3 to define a multilinear EF. We illustrate this assumption on the Gaussian distribution which can be written as the following exponential form, where $A(\mu, \mathbf{S}) = \frac{1}{2} [\mu^T \mathbf{S} \mu - \log |\mathbf{S}| / (2\pi)]$ is the log-partition function.

$$q(\mathbf{z}|\mu, \mathbf{S}) = \exp \left(-\frac{1}{2} \mathbf{z}^T \mathbf{S} \mathbf{z} + \mathbf{z}^T \mathbf{S} \mu - A(\mu, \mathbf{S}) \right)$$

Considering two blocks with μ and \mathbf{S} respectively, we can express this distribution as follows, where the first equation is for μ while the second equation is for \mathbf{S} :

$$\begin{aligned} q(\mathbf{z}|\mu, \mathbf{S}) &= \underbrace{\exp(-\frac{1}{2} \mathbf{z}^T \mathbf{S} \mathbf{z})}_{h_1(\mathbf{z}, \mathbf{S})} \exp \left(\underbrace{\langle \mathbf{S} \mathbf{z}, \mu \rangle}_{\phi_1(\mathbf{z}, \mathbf{S})} - A(\mu, \mathbf{S}) \right) \\ &= \underbrace{1}_{h_2(\mathbf{z}, \mu)} \exp \left(\underbrace{\langle -\frac{1}{2} \mathbf{z} \mathbf{z}^T + \mu \mathbf{z}^T, \mathbf{S} \rangle}_{\phi_2(\mathbf{z}, \mu)} - A(\mu, \mathbf{S}) \right) \end{aligned}$$

We define the *block-coordinate natural (BCN) parameterization* for an EF distribution as the parameterization which

satisfies Assumptions from 1 to 3. Therefore, Gaussian distribution with μ and S can be expressed in a BCN parameterization $\lambda = \{\lambda^{[1]}, \lambda^{[2]}\}$, where $\lambda^{[1]} = \mu$ and $\lambda^{[2]} = S$. Let λ^{a_i} denote the a_i -th entry of the i -th block parameter $\lambda^{[i]}$, where a_i is a local index for block i . \hat{g}^{c_i} is the c -th entry of natural gradient $\hat{g}^{[i]}$ with respect to $\lambda^{[i]}$.

We now present the rule (see Section 5 for a derivation). Under a BCN parameterization λ , our rule for block i takes the following form with an extra term shown in red:

$$\lambda^{c_i} \leftarrow \lambda^{c_i} - t\hat{g}^{c_i} - \frac{t^2}{2} \sum_{a_i} \sum_{b_i} \Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i}, \quad (7)$$

where each summation is to sum over all entries of the i -th block, $\Gamma_{a_i b_i}^{c_i} := \frac{1}{2} \partial_{m_{c_i}} \partial_{\lambda^{a_i}} \partial_{\lambda^{b_i}} A(\lambda)$, and m_{c_i} is the c -th entry of the BC expectation parameter $m_{[i]} := \mathbb{E}_q[\phi_i(\mathbf{z}, \lambda^{[-i]})] = \partial_{\lambda^{[i]}} A(\lambda)$.

The modification involves computation of the third-order term of the log-partition function² $A(\lambda)$. In the following Section 3.1 and 3.2, we discuss two examples where this computation is simplified and can be carried out like the original rule with minimal computational increase.

Table 2 in Appendix C lists more examples satisfying Assumption 1-3, where our rule can be applied and simplified.

3.1. Example: Online Newton using Gaussian Approximation

The original rule for Gaussian approximations gives the update (4)-(5), where the natural parameterization of Gaussian is used. We consider the parameterization μ and S , in which the improved rule takes the form (a detailed simplification is in Appendix E) with an extra non-zero term shown in red:

$$\mu \leftarrow \mu - tS^{-1} \mathbb{E}_q[\nabla_z \bar{\ell}(\mathbf{z})] + \mathbf{0} \quad (8)$$

$$S \leftarrow (1-t)S + t\mathbb{E}_q[\nabla_z^2 \bar{\ell}(\mathbf{z})] + \frac{t^2}{2} \hat{G} S^{-1} \hat{G}, \quad (9)$$

where $\hat{G} := S - \mathbb{E}_q[\nabla_z^2 \bar{\ell}(\mathbf{z})]$. The extra term ensures that the positive definite constraint is satisfied due to Theorem 1.

Theorem 1 *The updated S in (9) is positive definite if the initial S is positive-definite.*

The proof of Theorem 1 can be found in Appendix E.1.

Although (8)-(9) appear similar to (4)-(5), there is one difference – the old S is used as a preconditioner to update μ . Note that (8)-(9) becomes a natural-gradient descent (NGD) update if we ignore the additional term. Though natural gra-

²We assume $A(\lambda)$ is (jointly) C^3 -smooth. Note that $A(\lambda)$ is block-wisely C^3 -smooth as shown in Johansen (1979). Approximations considered in this paper satisfy this assumption.

dient³ is invariant to parameterization, NGD update depends on parameterization as shown by the difference. However, we expect the difference to make a small change in practice.

Like Khan & Rue (2020), an online Newton method can be obtained by approximating the expectations at μ , e.g., $\mathbb{E}_q[\nabla_z \bar{\ell}(\mathbf{z})] \approx \nabla_\mu \bar{\ell}(\mu)$ and $\mathbb{E}_q[\nabla_z^2 \bar{\ell}(\mathbf{z})] \approx \nabla_\mu^2 \bar{\ell}(\mu)$. In this case, the algorithm converges to a local minimal of the loss $\bar{\ell}(\mathbf{z})$. A key point is that, unlike Newton’s method where the preconditioner may not be positive-definite for nonconvex functions, S is guaranteed to be positive definite.

When applied to factorized Gaussians, these updates give an improved version of the Variational Online Gauss-Newton (VOGN) algorithm in Osawa et al. (2019). It is shown in Figure 1 where the differences in our algorithm are shown in red. Our algorithm uses the reparameterization trick to avoid computing $\nabla_z^2 \bar{\ell}(\mathbf{z})$ in (9). A derivation is given in Appendix E.3. Our algorithm fixes an implementation issue in VOGN without comprising its performance and speed, where our update only stores a mini-batch gradient while VOGN has to store all individual gradients in a mini-batch.

3.2. Example: Gamma Approximation

Let’s consider gamma cases. We use a BCN parameterization $\lambda = \{\lambda^{[1]}, \lambda^{[2]}\}$ (see Appendix F for detail), where $\lambda^{[1]} = \alpha$ and $\lambda^{[2]} = \frac{\beta}{\alpha}$. The constraint is $\Omega = \mathbb{S}_{++}^1 \times \mathbb{S}_{++}^1$. Since each block contains a scalar, we use global indexes as $\lambda^{(i)} = \lambda^{[i]}$ and $\hat{g}^{(i)} = \hat{g}^{[i]}$. Moreover, we use Γ_{ii}^i to denote $\Gamma_{a_i b_i}^{c_i}$. Let $\text{Ga}(\cdot)$ be the gamma function. Under this parameterization, a gamma distribution is expressed as:

$$q(z|\lambda) = \frac{1}{z} \exp\left(\lambda^{(1)} \log z - z\lambda^{(1)}\lambda^{(2)} - A(\lambda)\right),$$

where $A(\lambda) = \log \text{Ga}(\lambda^{(1)}) - \lambda^{(1)} (\log \lambda^{(1)} + \log \lambda^{(2)})$.

Let $\psi(\cdot)$ be the digamma function. We can compute the third derivatives (see Appendix F for a derivation) as:

$$\Gamma_{11}^1 = \frac{\frac{1}{\lambda^{(1)} \times \lambda^{(1)}} + \partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)})}{2 \left(-\frac{1}{\lambda^{(1)}} + \partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) \right)}, \quad \Gamma_{22}^2 = -\frac{1}{\lambda^{(2)}}$$

The proposed rule in this case is

$$\lambda^{(i)} \leftarrow \lambda^{(i)} - t\hat{g}^{(i)} - \frac{t^2}{2} (\Gamma_{ii}^i) \hat{g}^{(i)} \times \hat{g}^{(i)}, \quad i = 1, 2 \quad (10)$$

where each $\hat{g}^{(i)}$ is a natural gradient computed via the implicit re-parameterization trick as shown in Appendix F.2.

Theorem 2 *The updated $\lambda^{(i)}$ in (10) is positive if the initial $\lambda^{(i)}$ is positive.*

The proof of Theorem 2 can be found in Appendix F.1.

³It is a representation of an abstract (parameterization-free) tangent vector in a Riemannian manifold under a parameterization.

3.3. Extension to EF Mixtures

Our learning rule can be extended to mixture approximations, such as finite mixture of Gaussians (MOG) (shown in Appendix J) and skew Gaussian approximations (given in Appendix K) using the joint FIM – the FIM of the joint distribution of a mixture – as suggested by Lin et al. (2019a). By extending the definition of the BCN parameterization to the joint distribution, our rule can be easily applied to mixture cases (see Appendix I for detail). For example, our update for MOG approximation can be found at (27) in Appendix J, where our rule handles the positive-definite constraints in MOG. Our update can be viewed as an improved version of the ensemble of Newton methods proposed by Lin et al. (2019a). We also discuss why it is non-trivial to extend VOGN to MOG cases in Appendix J.1.

4. Related Works

In d -dim Gaussian cases, we can use unconstrained transformations (e.g., a Cholesky factor). However, the natural-gradient computation becomes complicated. Eq (2) gives $O(d^6)$ for direct computation. Salimbeni et al. (2018) propose an indirect approach via additional vector-Jacobian products (VJPs), which could give $O(d^4)$. For some parameterizations, their method gives an implicit $O(d^3)$ update, where Auto-Diff is needed to track non-zero terms in the additional Jacobians and to simplify the VJPs. Contrarily, our method gives a simple and explicit $O(d^3)$ update and builds a direct connection to Newton’s method. In practice, our update is more numerically stable and much faster (see Figure 7 in Appendix L) than theirs if both use Auto-Diff. Our approach is also easily extended to EFs and mixtures.

Our work is closely related to the method of Tran et al. (2019). They propose a method based on a retraction map in Gaussian cases, which is a special case of ours (see Appendix E.4). However, their retraction map does not directly generalize to other distributions, while ours does. They do not provide a justification or derivation of the map. We fix this gap by deriving the map from first principles, justifying its use, and obtaining an Adam-like update by choosing a proper parametrization for Gaussian cases (see Appendix E). They also do not distinguish the difference between the Riemannian gradient for a positive-definite matrix and the natural gradient for a covariance matrix (see Footnote 20 in Appendix E.4). Moreover, the retraction and Riemannian gradients used in neural network cases are not derived from the same Riemannian metric. In our work, retractions are implicitly induced by our rule. Our retractions and Riemannian gradients are naturally derived from the same metric.

Song et al. (2018) give a similar update in non-Bayesian contexts, but the update does not always satisfy the constraints for univariate Gaussians (see Appendix A). Their

update is neither simple nor efficient for multivariate cases such as multivariate Gaussians and MOGs (see Section 5.3).

Hosseini & Sra (2015) use a similar approach to ours but for parameter estimation of Gaussian mixtures. They propose a transformation for each Gaussian component so that the mean and the covariance together can be re-parameterized as an augmented positive-definite matrix with an extra constraint.⁴ They show that a local minimum of the negative log-likelihood of a mixture automatically satisfies the extra constraint. Thus, they can employ Riemannian-gradient descent with a retraction map to update the augmented positive-definite matrix, where the extra constraint can be safely ignored. It is unclear if this approach generalizes to variational inference (VI) since generating samples from MOG requires the extra constraint to be satisfied. Thus, the constraint cannot be omitted at each iteration in VI. Moreover, it is unclear whether Riemannian gradients and the retraction are derived from the same metric for the mixture. Hosseini & Sra (2015) use the gradients and the retraction designed for positive-definite matrices while we derive them from the joint Fisher metric for MOGs in a principled way.

5. Derivation of the Improved Rule

5.1. Gradient Descent

We first review gradient descent in Euclidean spaces and generalize it to Riemannian manifolds, where we derive our rule. Recall that we want to minimize (1) in terms of λ as:

$$\min_{\lambda \in \Omega} \mathbb{E}_{q(z|\lambda)}[\ell(\mathcal{D}, \mathbf{z})] + \mathbb{D}_{KL}[q(\mathbf{z}|\lambda) \| p(\mathbf{z})] \equiv \mathcal{L}(\lambda).$$

If $\Omega = \mathbb{R}^d$ is a Euclidean space,⁵ we can solve the minimization problem using gradient descent (GD) as:

$$\text{GD} : \lambda \leftarrow \lambda - t\mathbf{g}$$

where $\mathbf{g} = \partial_{\lambda} \mathcal{L}(\lambda)$ denotes a Euclidean gradient and $t > 0$ is a scalar step-size. We can view the update as a line (the shortest curve) $\mathbf{L}(t)$ in the Euclidean space \mathbb{R}^d as t varies. Given a starting point λ and a Euclidean direction $-\mathbf{g}$, the line is a differentiable map $\mathbf{L}(t)$ so that the following ordinary differential equation⁶ (ODE) is satisfied.

$$\dot{\mathbf{L}}(0) = -\mathbf{g} ; \quad \mathbf{L}(0) = \lambda ; \quad \ddot{\mathbf{L}}(t) = \mathbf{0} \quad (11)$$

where $\dot{\mathbf{L}}(x) := \frac{d\mathbf{L}(t)}{dt} \Big|_{t=x}$, $\ddot{\mathbf{L}}(x) := \frac{d^2\mathbf{L}(t)}{dt^2} \Big|_{t=x}$. The solution of the ODE is $\mathbf{L}(t) = \lambda - t\mathbf{g}$, which is the GD update.

5.2. Exact Riemannian Gradient Descent (RGD)

Unfortunately, Ω usually is not a Euclidean space but a Riemannian manifold with a metric. We use a metric to characterize distances in the manifold. A useful metric for

⁴The parameterization violates Assumption 1 since the positive-definite constraint of the augmented matrix and the extra constraint can not be partitioned into two blocks.

⁵In this paper, it always uses a Cartesian coordinate system.

⁶It is also known as an initial value problem.

Table 1. Table of Index Notation

$\lambda^{[i]}$	i -th block parameter of parameterization λ .
λ^{a_i}	a -th entry of block parameter $\lambda^{[i]}$.
$\lambda^a, \lambda^{(a)}$	a -th entry of parameterization λ .
g_a	a -th entry of Euclidean gradient \mathbf{g} .
$\hat{g}^a, \hat{g}^{(a)}$	a -th entry of Riemannian/natural gradient $\hat{\mathbf{g}}$.
F_{ab}	entry of \mathbf{F} with global index (a, b) .
F^{ab}	entry of \mathbf{F}^{-1} with global index (a, b) .
Γ_{ab}^c	entry with global index (c, a, b) .
$F^{a_i b_i}$	entry with local index (a, b) in block i .
$\Gamma_{a_i b_i}^{c_i}$	entry with local index (c, a, b) in block i .

statistical manifolds is the FIM (Fisher, 1922; Rao, 1945).

Now, we generalize gradient descent in a manifold. First, we introduce the index convention and the Einstein summation notation used in Riemannian geometry. The notation is summarized in Table 1. We denote a Euclidean gradient \mathbf{g} using a subscript. A Riemannian gradient $\hat{\mathbf{g}}$ is denoted by a superscript. A metric⁷ is used to characterize inner products and arc length in a manifold. Given a metric \mathbf{F} , let F_{ab} denote the element of \mathbf{F} at position (a, b) and F^{ca} denote the element of \mathbf{F}^{-1} at position (c, a) . We use the Einstein notation to omit summation symbols such as $F^{ca}F_{ab} := \sum_a F^{ca}F_{ab}$. Therefore, we have $F^{ca}F_{ab} = I_b^c$, where I_b^c is the element of an identity matrix at position (c, b) . A Riemannian gradient is defined as $\hat{g}^c := F^{ca}g_a$, where g_a is the a -th element of a Euclidean gradient \mathbf{g} . When \mathbf{F} is the FIM, a Riemannian gradient becomes a natural gradient. If the metric \mathbf{F} is positive-definite for all⁸ $\lambda \in \Omega$, an approximation family $q(\mathbf{z}|\lambda)$ induces a Riemannian manifold denoted by (Ω, \mathbf{F}) where λ is a coordinate system.

Like GD, RGD can be derived from a geodesic,⁹ which is a generalization of the “shortest” curve¹⁰ to a manifold. Given a starting point $\lambda \in \Omega$ and a Riemannian direction $-\hat{\mathbf{g}} = -\mathbf{F}^{-1}\mathbf{g}$, a geodesic is a differentiable map $\mathbf{L}(t)$ so that the following geodesic ODE¹¹ is satisfied.

$$\dot{\mathbf{L}}^c(0) = -F^{ca}g_a; \quad \mathbf{L}^c(0) = \lambda^c \quad (12)$$

$$\ddot{\mathbf{L}}^c(t) = -\Gamma_{ab}^c(t)\dot{\mathbf{L}}^a(t)\dot{\mathbf{L}}^b(t) \quad (13)$$

where $\mathbf{L}^c(t)$ is the c -th element of $\mathbf{L}(t)$, $\dot{\mathbf{L}}^c(x) := \frac{d\mathbf{L}^c(t)}{dt}\big|_{t=x}$, $\ddot{\mathbf{L}}^c(x) := \frac{d^2\mathbf{L}^c(t)}{dt^2}\big|_{t=x}$, $\Gamma_{ab}^c(t) := \Gamma_{ab}^c|_{\lambda=\mathbf{L}(t)}$. Γ_{ab}^c is the Christoffel symbol of the 2nd kind defined by

$$\Gamma_{ab}^c := F^{cd}\Gamma_{d,ab}; \quad \Gamma_{d,ab} := \frac{1}{2}[\partial_a F_{bd} + \partial_b F_{ad} - \partial_d F_{ab}]$$

where $\partial_a := \partial_{\lambda^a}$ is for notation simplicity and $\Gamma_{d,ab}$ is the Christoffel symbol of the 1st kind. $\ddot{\mathbf{L}}$ characterizes the

⁷A metric is well-defined if it is positive definite everywhere.

⁸Such assumption is valid for minimal EF.

⁹The geodesic induces an exponential map used in exact RGD.

¹⁰Due to the Euler-Lagrange equation, a geodesic is a stationary curve. However, a geodesic may not be the shortest curve.

¹¹The domain of $\mathbf{L}(t)$ is \mathbb{R} for a complete manifold.

curvature of a geodesic since a manifold is not flat in general. In Euclidean cases, the metric $\mathbf{F} = \mathbf{I}$ is a constant identity matrix and (13) vanishes since $\Gamma_{d,ab}$ and Γ_{ab}^c are zeros, which implies Euclidean spaces are flat. Therefore, we recover the GD update in (11) since $\hat{\mathbf{g}} = \mathbf{I}^{-1}\mathbf{g} = \mathbf{g}$.

Given any parameterization with the FIM, we can compute $\Gamma_{d,ab}$ by using Eq (17) in Appendix D.1, which involves extra integrations. We will show that a BCN parameterization can get rid of the extra integrations (see Theorem 3).

5.3. Our Rule as an Inexact RGD Update

However, it is hard to exactly solve the geodesic ODE. Inexact RGD is derived by approximating the geodesic.¹² Recall that the original rule is a natural gradient descent (NGD) update. NGD can be derived by the first-order approximation of the geodesic $\mathbf{L}(t)$ at $t_0 = 0$ with the FIM.

$$\text{NGD} : \quad \lambda \leftarrow \mathbf{L}(t_0) + \dot{\mathbf{L}}(t_0)(t - t_0) = \lambda - t\hat{\mathbf{g}}$$

Unfortunately, this approximation is only well-defined in a small neighborhood at t_0 with radius t . For a stochastic NGD update, the step-size t is very small, which often result in slow convergence. Our learning rule addresses this issue, which is indeed a new inexact RGD update. Moreover, our update can use a bigger step-size and often converges faster than NGD without introducing significant computational overhead in useful cases such as gamma, Gaussian, MOG.

Consider cases when $\lambda = \{\lambda^{[1]}, \dots, \lambda^{[m]}\}$ has m blocks. We can express a Riemannian gradient as $\hat{\mathbf{g}} = \{\hat{g}^{[1]}, \dots, \hat{g}^{[m]}\}$. We use the block summation notation to omit the summation signs in (7) as $\Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i} := \sum_{a_i} \sum_{b_i} \Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i}$. By the global index notation, we have $\Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i} = \sum_{a \in [i]} \sum_{b \in [i]} \Gamma_{ab}^{(c_i)} \hat{g}^a \hat{g}^b$, where $[i]$ is the index set for block i , (c_i) is the corresponding global index of local index c_i , and a and b are global indexes.

We can extend the definition of a BC parameterization to any Riemannian metric \mathbf{F} . Given a metric \mathbf{F} , we have Lemma 1 for any block i (see Appendix B.1 for a proof) :

Lemma 1 When λ is a BC parameterization of metric \mathbf{F} , we have $\hat{g}^{a_i} = F^{a_i b_i} g_{b_i}$ and $\Gamma_{a_i b_i}^{c_i} = F^{c_i d_i} \Gamma_{d_i, a_i b_i}$.

Given a manifold equipped with metric \mathbf{F} and a BC parameterization λ , consider the solution of the block-wise (geodesic) ODE¹³ denoted by $\mathbf{R}^{[i]}(t)$ for block i :

$$\dot{\mathbf{R}}^{c_i}(0) = -F^{c_i a_i} g_{a_i}; \quad \mathbf{R}^{c_i}(0) = \lambda^{c_i} \quad (14)$$

$$\ddot{\mathbf{R}}^{c_i}(t) = -\Gamma_{a_i b_i}^{c_i}(t) \dot{\mathbf{R}}^{a_i}(t) \dot{\mathbf{R}}^{b_i}(t) \quad (15)$$

¹²A retraction map can be derived by approximating the geodesic. An exact RGD update is invariant under parameterization while inexact RGD updates including NGD often are not.

¹³ $\mathbf{R}^{[i]}(t)$ is easier to solve compared to $\mathbf{L}(t)$. Note that (14) is the minimum requirement of a retraction map (Absil et al., 2009).

where $R^{c_i}(0)$, $\dot{R}^{c_i}(0)$, $\ddot{R}^{c_i}(t)$ respectively denote the c -th entry of $\mathbf{R}^{[i]}(0)$, $\dot{\mathbf{R}}^{[i]}(0)$, and $\ddot{\mathbf{R}}^{[i]}(t)$; $\Gamma_{a_i b_i}^{c_i}(t) := \Gamma_{a_i b_i}^{c_i} \big|_{\lambda^{[i]} = R^{[i]}(t)}$.

We use $\hat{q}(\mathbf{z}|\lambda^{[i]})$ to denote $q(\mathbf{z}|\lambda)$ when $\lambda^{[j]} = \mathbf{R}^{[j]}(0)$ is known for each block j except the i -th block. $F^{c_i a_i}$ is the entry of $(\mathbf{F}^{[i]})^{-1}$ at position (c, a) , where $\mathbf{F}^{[i]}$ is the sub-block matrix of \mathbf{F} for $\lambda^{[i]}$. In fact, $\mathbf{F}^{[i]}$ is the induced metric for $\hat{q}(\mathbf{z}|\lambda^{[i]})$, and $\mathbf{R}^{[i]}(t)$ is a geodesic for $\hat{q}(\mathbf{z}|\lambda^{[i]})$ under BC parameterization λ . Moreover, if λ is a BCN parameterization and \mathbf{F} is the FIM, we have $F_{a_i b_i} = \partial_{\lambda^{a_i}} \partial_{\lambda^{b_i}} A(\lambda)$.

We define a curve $\mathbf{R}(t) := \{\mathbf{R}^{[1]}(t), \dots, \mathbf{R}^{[m]}(t)\}$. By Lemma 1, we can show that the first-order approximation of $\mathbf{R}(t)$ at $t_0 = 0$ induces NGD if \mathbf{F} is the FIM and λ is a BC parameterization. Appendix B.2 shows this in detail.

We propose to use the second-order approximation¹⁴ of $\mathbf{R}(t)$ at $t_0 = 0$ for block i , where λ is a BC parameterization.

$$\begin{aligned} \text{Our : } \lambda^{c_i} &\leftarrow R^{c_i}(t_0) + \dot{R}^{c_i}(t_0)(t - t_0) + \frac{1}{2} \ddot{R}^{c_i}(t_0)(t - t_0)^2 \\ &= \lambda^{c_i} - t \hat{g}^{c_i} - \frac{t^2}{2} \Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i} \end{aligned} \quad (16)$$

where $\Gamma_{a_i b_i}^{c_i}$ is computed at $t_0 = 0$, $\hat{g}^{c_i} = F^{c_i a_i} g_{a_i}$, and c_i denotes the c -th element of the i -th block. Our rule works for both a BCN parameterization and a BC parameterization.

Song et al. (2018) suggest using the second-order approximation of $\mathbf{L}(t)$ at $t_0 = 0$, which has to compute the whole Christoffel symbol $\Gamma_{ab}^{(c_i)}$. However, their proposal does not guarantee the update stays in the constraint set even in univariate Gaussian cases (see Appendix A). Moreover, it is inefficient to compute the whole Christoffel symbol since all cross terms between any two blocks are needed in $\Gamma_{ab}^{(c_i)} \hat{g}^a \hat{g}^b$. When a parameterization has m blocks, $\Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i} \neq \Gamma_{ab}^{(c_i)} \hat{g}^a \hat{g}^b$ since the hidden summations are taken over entries only in block i on the left while the summations are taken over entries in all m blocks on the right. This is the key difference between our method and their method. In our method, only the block-wise $\Gamma_{a_i b_i}^{c_i}$ is computed, which makes our method efficient in many cases such as multivariate Gaussians and MOGs. Moreover, a BCN parameterization can further simplify the computation of our rule due to Theorem 3 (see Appendix D for a proof).

Theorem 3 *Under a BCN parameterization of EF with the FIM, natural gradients and the Christoffel symbol for each block i can be simplified as*

$$\hat{g}^{a_i} = \partial_{m_{a_i}} \mathcal{L}(\lambda) ; \Gamma_{d_i, a_i b_i} = \frac{1}{2} \partial_{\lambda^{a_i}} \partial_{\lambda^{b_i}} \partial_{\lambda^{d_i}} A(\lambda)$$

¹⁴Our approximation allows us to use a bigger step-size than NGD. In many cases, the underlying parameterization constraints are satisfied regardless of the choice of the step-size and therefore, a line search for the constraint satisfaction is no longer required.

where λ^{a_i} is the a -th entry of $\lambda^{[i]}$; m_{a_i} is the a -th entry of the BC expectation parameter¹⁵ $\mathbf{m}^{[i]} := \mathbb{E}_q[\phi_i(\mathbf{z}, \lambda^{[i]})] = \partial_{\lambda^{[i]}} A(\lambda)$.

Since $A(\lambda)$ is C^3 -smooth for block $\lambda^{[i]}$ (Johansen, 1979), we have $\partial_{\lambda^{a_i}} \partial_{\lambda^{b_i}} \partial_{\lambda^{d_i}} A(\lambda) = \partial_{\lambda^{d_i}} \partial_{\lambda^{a_i}} \partial_{\lambda^{b_i}} A(\lambda)$. Thus, by Theorem 3, we have $\Gamma_{a_i b_i}^{c_i} = \frac{1}{2} \partial_{m_{c_i}} \partial_{\lambda^{a_i}} \partial_{\lambda^{b_i}} A(\lambda)$.

A similar theorem for EF mixtures is in Appendix I.

To sum up, our rule is an instance of RGD with a retraction map. We give a principled way to derive Riemannian gradients and retractions from scratch (see Footnote 12,13). The convergence analysis could be obtained by existing works (Bonnabel, 2013) if the retraction satisfies some properties.

6. Numerical Results¹⁶

6.1. Results on Synthetic Examples

To validate our rule, we visualize our approximations in 2-dimensional toy examples, where we use the re-parametrization trick suggested by Lin et al. (2019a;b) (see (20) in Appendix E for full Gaussian and (28) in Appendix J for mixture of Gaussians (MOG)) to compute gradients. Due to the trick, $\nabla_z^2 \bar{\ell}(\mathbf{z})$ is not needed. See Figure 5-6 in Appendix L for more visualization examples such as the banana distribution (Haario et al., 2001) and a BNN example taken from Au et al. (2020). We then compare our method to baseline methods in a higher dimensional example.

We first visualize Gaussian approximations with full covariance structures for the Bayesian Logistic regression example taken from Murphy (2013) ($N = 60, d = 2$). Figure 2(a) shows posterior approximations obtained from various methods. From the figure, our approximation matches the exact variational Gaussian approximation. For skew-Gaussian approximations (Lin et al., 2019a) and mean-field Gaussian approximations, see Figure 6 in Appendix L.

In the second example, we approximate the beta-binomial model (Salimans & Knowles, 2013) ($N = 20, d = 2$) by MOG. The exact posterior is skewed. From Figure 2(b), we see that the approximation matches the exact posterior better and better as we increase the number of mixtures.

In the third example, we approximate a correlated Laplace distribution $\exp(-\bar{\ell}(\mathbf{z})) = \text{Lap}(z_1|0, 1)\text{Lap}(z_2|z_1, 1)$ by using MOG, where $\text{Lap}(z_2|z_1, 1) = \frac{1}{2} \exp(-|z_2 - z_1|)$. The distribution is non-smooth and thus $\nabla_z^2 \bar{\ell}(\mathbf{z})$ does not exist. However, we can use the re-parametrization trick since $\nabla_z \bar{\ell}(\mathbf{z})$ exists almost surely. From Figure 2(c), we see that our method gives smooth approximations of the target.

¹⁵Instead of using $\mathbf{m}^{[i]}$, we use $\mathbf{m}_{[i]}$ to emphasize that Euclidean gradient for $\mathbf{m}_{[i]}$ is equivalent to natural gradient for $\lambda^{[i]}$.

¹⁶Our implementation: github.com/yorkerlin/iBayesLRrule




Figure 2. Visualization of posterior approximations on 2-D toy examples. Figure 2(a) shows the Gaussian approximation to fit a Bayesian logistic model, where our approximation matches the exact variational Gaussian approximation. Figure 2(b) shows MOG approximation fit to a beta-binomial model in a 2-D problem. The number indicates the number of mixture components. By increasing the number of components, we get better results. Figure 2(c) shows MOG approximation fit to a correlated 2-D Laplace distribution. The number indicates the number of mixtures. We get smooth approximations of the non-smooth distribution. Figure 2(d) shows MOG approximation fit to a double banana distribution. The number indicates the number of mixtures, where we only show the last 8 MOG approximations. The complete MOG approximations can be found in Appendix L. As the number of components increases, we get better results.




Figure 3. Comparison results on a 20-D mixture of Student’s t distributions with 10 components by MOG approximations. The leftmost figure shows the performance of each method, where our method outperforms existing methods. The first 9 dimensions obtained by our method are shown in the figure where MOG approximation fits the marginals well. We also test a 300-D mixture problem in Appendix L.

In the fourth example, we approximate the double banana distribution constructed by [Detommaso et al. \(2018\)](#). The true distribution has two modes and is skewed. As shown in Figure 2(d), our MOG approximation approximates the target better and better when we increase the number of mixtures. For a complete plot using the approximation with various components, see Figure 5(b) in Appendix L.

Finally, we conduct a comparison study on approximations for a mixture of Student’s t distributions $\exp(-\ell(\mathbf{z})) = \frac{1}{C} \sum_{k=1}^C \mathcal{T}(\mathbf{z}|\mathbf{u}_k, \mathbf{V}_k, \alpha)$ with degrees of freedom $\alpha = 2$, where $\mathbf{z} \in \mathbb{R}^d$. We generate each entry of location vector \mathbf{u}_k uniformly in an interval $(-s, s)$. Each shape matrix \mathbf{V}_k is taken a form of $\mathbf{V}_k = \mathbf{A}_k^T \mathbf{A}_k + \mathbf{I}_d$, where each entry of the $d \times d$ matrix \mathbf{A}_k is independently drawn from a Gaussian distribution with mean 0 and standard deviation $0.1d$. We approximate the posterior distribution by MOG with K components and use the importance sampling technique to compute gradients as suggested by [Lin et al. \(2019a\)](#) so that the number of Monte Carlo (MC) samples does not depend the number of Gaussian components K . We compare our

method to existing methods, where the BayesLRule for MOG is proposed by [Lin et al. \(2019a\)](#).

We consider a case with $K = 25, C = 10, d = 20, s = 20$. For simplicity, we fix the mixing weight to be $\frac{1}{K}$ and only update each Gaussian component with the precision \mathbf{S}_c and the mean μ_c during training. We use 10 MC samples to compute gradients, where gradients are computed using either the re-parametrization trick (referred to as “-rep”) as shown in (28) in Appendix J or the Hessian trick (referred to as “-hess”) as shown in (29) in Appendix J. Note that BayesLRule with either the re-parametrization trick or the Hessian trick does not stay in the constraint set. We use the same initialization and tune the step size by grid search for each method. The leftmost plot of Figure 3 shows the performance. We clearly see that our methods converge fastest, when we use the maximum mean discrepancy (MMD) to measure the difference between an approximation and the ground-truth. The remaining plots of Figure 3 show the first 9 marginal distributions of the true distribution and our approximations with two kinds of gradient estimation, where




Figure 4. Results on real-world datasets showing the performances of our method (iBayesLRule) highlighted in red compared to BBVI, BayesLRule with a line search, and VOGN. Figure 4(a) and 4(b) show the performances using Gaussian approximations with full covariance structure to fit a Bayesian linear regression and a Bayesian logistic regression, respectively, where our method converges faster than BayesLRule and BBVI and gives a more accurate approximation than VOGN. Figure 4(c) shows the performances using Gamma approximations to fit a Gamma factor model, where our method converges faster. Figure 4(d) shows the performances of methods in a Bayesian MLP network with diagonal Gaussian approximations, where our method performs comparably to VOGN.

MOG closely matches the marginals. All 20 marginal distributions are in Figure 8 in Appendix L. We also consider a more difficult case with $K = 60$, $C = 20$, $d = 300$, $s = 25$ using 10 MC samples. Figure 9-13 in Appendix L show all 300 marginal distributions obtained by our method.

6.2. Results on Real Data

Now, we show results on real-world datasets. We consider four models in our experiments. The first model is the Bayesian linear regression, where we can obtain the exact solution and the optimal negative ELBO denoted by \mathcal{L}^* . We present results for full Gaussian approximations on the “Abalone” dataset ($N = 4,177$, $d = 8$) with 3341 chosen for training. We train the model with mini-batch size 168. In Figure 4(a), we plot the difference of ELBO between the exact and an approximation. We compare our method (referred to as “iBayesLRule”) to the black-box gradient method (referred to as “BBVI”) using the Adam optimizer (Kingma & Ba, 2015) and the original Bayesian learning rule (referred to as “BayesLRule”) with the re-parametrization trick (referred to as “-rep”) and the VOGN method. BBVI requires us to use an unconstrained parametrization. BayesLRule with the re-parametrization trick does not stay in the constraint set so a line search has to be used in BayesLRule. We can see that our method converges faster than BayesLRule and BBVI and is more accurate than VOGN.

Next, we consider the Bayesian logistic regression and present results for full Gaussian approximations on the “Ionosphere” dataset ($N = 351$, $d = 34$) with 175 chosen for training. We train the model with mini-batch size 17. In Figure 4(b), we plot the test log-loss and compare our method to BBVI and BayesLRule with the re-parametrization trick (referred to as “-rep”). We also consider the VOGN method proposed for Gaussian approximations. Note that BayesLRule using the re-parametrization trick does not stay in the constraint set and a line search is

used. From the plot, we can see our method outperforms BayesLRule and performs comparably to VOGN.

Then, we consider the Gamma factor model (Knowles, 2015; Khan & Lin, 2017) using Gamma approximations on the “CyTOF” dataset ($N = 522,656$, $d = 40$) with 300,000 chosen for training, where gradients are computed using the implicit re-parametrization trick (Furnov et al., 2018) (referred to as “-rep”). We train the model with mini-batch size 39 and tune the step size for all methods. In Figure 4(c), we plot the test log-loss and compare our to BayesLRule and BBVI. BayesLRule uses a line search since the updates using the re-parametrization trick do not satisfy the constraint. Our method outperforms BayesLRule and BBVI.

Finally, we consider a Bayesian MLP network with 2 hidden layers, where we use 1000 units for each layer. We train the network with diagonal Gaussian approximations on the “CIFAR-10” dataset ($N = 60,000$, $d = 3 \times 32 \times 32$) with 50,000 images for training and 10,000 images for validation. We train the model with mini-batch size 128 and compare our Adam-like update (referred to as “iBayesLRule-adam”) to VOGN. We use the same initialization and hyperparameters in both methods. In Figure 4(d), we plot the validation accuracy. Our method performs similarly to VOGN.

7. Discussion

We present an improved learning rule to handle positive-definite parameterization constraints. We propose a BCN parameterization so that natural gradients and the extra terms are easy to compute. Under this parameterization, the Fisher matrix and the Christoffel symbols admit a closed-form via differentiation without introducing extra integrations.

Our main focus is on the derivation of simple and efficient updates that naturally handle positive-definite constraints. We give examples where our updates have low iteration cost. We hope to perform large-scale experiments in the future.

Acknowledgements

We would like to thank Hiroyuki Kasai (Waseda University) for useful discussions at ICML 2019 at the early stage of this project. WL is supported by a UBC International Doctoral Fellowship.

References

- Absil, P.-A., Mahony, R., and Sepulchre, R. *Optimization algorithms on matrix manifolds*. Princeton University Press, 2009.
- Au, K. X., Graham, M. M., and Thiery, A. H. Manifold lifting: scaling MCMC to the vanishing noise regime. *arXiv preprint arXiv:2003.03950*, 2020.
- Batir, N. Some new inequalities for gamma and polygamma functions. *J. Inequal. Pure Appl. Math*, 6(4):1–9, 2005.
- Bonnabel, S. Stochastic gradient descent on Riemannian manifolds. *IEEE Transactions on Automatic Control*, 58(9):2217–2229, 2013.
- Detommaso, G., Cui, T., Marzouk, Y., Spantini, A., and Scheichl, R. A Stein variational Newton method. In *Advances in Neural Information Processing Systems*, pp. 9169–9179, 2018.
- Figurnov, M., Mohamed, S., and Mnih, A. Implicit reparameterization gradients. In *Advances in Neural Information Processing Systems*, pp. 441–452, 2018.
- Fisher, R. A. On the mathematical foundations of theoretical statistics. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, 222(594-604):309–368, 1922.
- Fletcher, P. T. and Joshi, S. Principal geodesic analysis on symmetric spaces: Statistics of diffusion tensors. In *Computer Vision and Mathematical Methods in Medical and Biomedical Image Analysis*, pp. 87–98. Springer, 2004.
- Haario, H., Saksman, E., Tamminen, J., et al. An adaptive Metropolis algorithm. *Bernoulli*, 7(2):223–242, 2001.
- Hosseini, R. and Sra, S. Matrix manifold optimization for Gaussian mixtures. In *Advances in Neural Information Processing Systems*, pp. 910–918, 2015.
- Johansen, S. Introduction to the theory of regular exponential families. 1979.
- Khan, M. and Lin, W. Conjugate-computation variational inference: Converting variational inference in non-conjugate models to inferences in conjugate models. In *Artificial Intelligence and Statistics*, pp. 878–887, 2017.
- Khan, M. E. and Rue, H. Learning-algorithms from Bayesian principles. 2020. https://emtiyaz.github.io/papers/learning_from_bayes.pdf.
- Khan, M. E., Nielsen, D., Tangkaratt, V., Lin, W., Gal, Y., and Srivastava, A. Fast and scalable Bayesian deep learning by weight-perturbation in Adam. In *Proceedings of the 35th International Conference on Machine Learning*, pp. 2611–2620, 2018.
- Kingma, D. P. and Ba, J. Adam: A method for stochastic optimization. In *International Conference on Learning Representations*, 2015.
- Knowles, D. A. Stochastic gradient variational Bayes for gamma approximating distributions. *arXiv preprint arXiv:1509.01631*, 2015.
- Koumandos, S. Monotonicity of some functions involving the gamma and psi functions. *Mathematics of Computation*, 77(264):2261–2275, 2008.
- Lin, W., Khan, M. E., and Schmidt, M. Fast and simple natural-gradient variational inference with mixture of exponential-family approximations. In *International Conference on Machine Learning*, pp. 3992–4002, 2019a.
- Lin, W., Khan, M. E., and Schmidt, M. Stein’s Lemma for the Reparameterization Trick with Exponential-family Mixtures. *arXiv preprint arXiv:1910.13398*, 2019b.
- Malagò, L. and Pistone, G. Information geometry of the Gaussian distribution in view of stochastic optimization. In *Proceedings of the 2015 ACM Conference on Foundations of Genetic Algorithms XIII*, pp. 150–162, 2015.
- Minh, H. Q. and Murino, V. Covariances in computer vision and machine learning. *Synthesis Lectures on Computer Vision*, 7(4):1–170, 2017.
- Murphy, K. P. *Machine learning : a probabilistic perspective*. MIT Press, Cambridge, Mass., 2013. ISBN 9780262018029 0262018020.
- Nielsen, F. On geodesic triangles with right angles in a dually flat space. *arXiv preprint arXiv:1910.03935*, 2019.
- Opper, M. and Archambeau, C. The variational Gaussian approximation revisited. *Neural computation*, 21(3):786–792, 2009.
- Osawa, K., Swaroop, S., Khan, M. E. E., Jain, A., Eschenhagen, R., Turner, R. E., and Yokota, R. Practical deep learning with Bayesian principles. In *Advances in neural information processing systems*, pp. 4287–4299, 2019.

- Pennec, X., Fillard, P., and Ayache, N. A Riemannian framework for tensor computing. *International Journal of computer vision*, 66(1):41–66, 2006.
- Rao, C. R. Information and accuracy attainable in the estimation of statistical parameters. *Bulletin of the Calcutta Mathematical Society*, 37(3):81–91, 1945.
- Salimans, T. and Knowles, D. Fixed-form variational posterior approximation through stochastic linear regression. *Bayesian Analysis*, 8(4):837–882, 2013.
- Salimbeni, H., Eleftheriadis, S., and Hensman, J. Natural gradients in practice: Non-conjugate variational inference in gaussian process models. *arXiv preprint arXiv:1803.09151*, 2018.
- Särkkä, S. *Bayesian filtering and smoothing*, volume 3. Cambridge University Press, 2013.
- Song, Y., Song, J., and Ermon, S. Accelerating natural gradient with higher-order invariance. In *International Conference on Machine Learning*, pp. 4720–4729, 2018.
- Tran, M.-N., Nguyen, D. H., and Nguyen, D. Variational Bayes on manifolds. *arXiv preprint arXiv:1908.03097v2*, 2019.
- Tseng, P.-H. and Lee, T.-C. Numerical evaluation of exponential integral: Theis well function approximation. *Journal of Hydrology*, 205(1-2):38–51, 1998.
- Zhang, G., Sun, S., Duvenaud, D., and Grosse, R. Noisy natural gradient as variational inference. In *International Conference on Machine Learning*, pp. 5847–5856, 2018.

Appendices

In the appendices, we will use the index notation and the Einstein summation notation introduced in Section 5.2.

A. A counter-example for Song et al. (2018)

We show that the update suggested by Song et al. (2018) does not stay in the constraint set while ours does.

Let's consider the following univariate Gaussian distribution under a BC parameterization $\lambda = \{\mu, \sigma\}$, where σ denotes the standard deviation¹⁷. The constraint is $\Omega_1 = \mathbb{R}$ and $\Omega_2 = \mathbb{S}_{++}^1$. $\hat{g}^{(1)}$ and $\hat{g}^{(2)}$ are natural gradients for μ and σ , respectively.

$$q(z|\lambda) = \exp \left\{ -\frac{1}{2} \left(\frac{z - \mu}{\sigma} \right)^2 - \frac{1}{2} \log(2\pi) - \log(\sigma) \right\}$$

Recall that the Christoffel symbols of the second kind can be computed as $\Gamma_{ab}^c = F^{cd} \Gamma_{d,ab}$ where $\Gamma_{d,ab}$ is the Christoffel symbols of the first kind and F^{cd} is the entry of the inverse the FIM, \mathbf{F}^{-1} , at position (c, d) .

Under this parameterization, the FIM and the Christoffel symbols of the second kind are given below, where the Christoffel symbols of the first kind are computed by using Eq. (17). The computation of the Christoffel symbols can be difficult since the parameterization is not a BCN parameterization.

$$F_{ab} = \begin{bmatrix} \frac{1}{\sigma^2} & 0 \\ 0 & \frac{2}{\sigma^2} \end{bmatrix}, \quad \Gamma^1_{ab} = \begin{bmatrix} 0 & -\frac{1}{\sigma} \\ -\frac{1}{\sigma} & 0 \end{bmatrix}, \quad \Gamma^2_{ab} = \begin{bmatrix} \frac{1}{2\sigma} & 0 \\ 0 & -\frac{1}{\sigma} \end{bmatrix}$$

The update suggested by Song et al. (2018) is

$$\begin{aligned} \mu &\leftarrow \mu - t\hat{g}^{(1)} - t\hat{g}^{(1)} - \frac{t \times t}{2} \Gamma^1_{ab} \hat{g}^{(a)} \hat{g}^{(b)} = \mu - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{2\hat{g}^{(1)}\hat{g}^{(2)}}{\sigma} \right) \\ \sigma &\leftarrow \sigma - t\hat{g}^{(2)} - t\hat{g}^{(2)} - \frac{t \times t}{2} \Gamma^2_{ab} \hat{g}^{(a)} \hat{g}^{(b)} = \sigma - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{2(\hat{g}^{(2)})^2 - (\hat{g}^{(1)})^2}{2\sigma} \right) \end{aligned}$$

Clearly, the updated σ does not always satisfy the positivity constraint \mathbb{S}_{++}^1 .

As shown in Eq. (16), our rule can be used in not only a BCN parameterization but also a BC parameterization. Since every block contains only a scalar, we use global indexes such as $\lambda^{(i)} = \lambda^{a_i}$, $\hat{g}^{(i)} = \hat{g}^{[i]}$ and $\Gamma_{i,ii} = \Gamma_{a_i,b_i c_i}$ for notation simplicity. Note that $\Gamma^1_{11} = 0$ is the entry at the upper-left corner of Γ^1_{ab} and $\Gamma^2_{22} = -\frac{1}{\sigma}$ is the entry at the lower-right corner of Γ^2_{ab} . In our update (see Eq. (16)), we can see the update automatically satisfies the constraint as shown below.

$$\begin{aligned} \underbrace{\lambda^{(1)}}_{\mu} &\leftarrow \underbrace{\lambda^{(1)}}_{\mu} - \frac{t^2}{2} \Gamma^1_{11} \hat{g}^{(1)} \hat{g}^{(1)} = \mu - t\hat{g}^{(1)} \\ \underbrace{\sigma}_{\lambda^{(2)}} &\leftarrow \underbrace{\sigma}_{\lambda^{(2)}} - \frac{t^2}{2} \Gamma^2_{22} \hat{g}^{(2)} \hat{g}^{(2)} = \sigma - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{(\hat{g}^{(2)})^2}{\sigma} \right) = \underbrace{\frac{1}{2\sigma}}_{>0} \left[\underbrace{\sigma^2}_{>0} + \underbrace{(\sigma - t\hat{g}^{(2)})^2}_{\geq 0} \right] \end{aligned}$$

As we discuss at Section 5.3 of the main text, only the block-wise Christoffel symbol $\Gamma_{i,ii}$ for each block i is required, which becomes essential for multivariate Gaussians and mixture of Gaussians.

Let's consider another BC parameterization $\lambda = \{\mu, v\}$ for the Gaussian distribution, where $v = \sigma^2$ denotes the variance. Note that we consider the parameterization for univariate Gaussian. For multivariate Gaussian, see Appendix E.4. The

¹⁷It is also used as an unconstrained parameterization of Gaussian distributions for BBVI. Technically, this parameterization has a positivity constraint, which is often ignored in practice. In multivariate cases, the Cholesky factor is used as an unconstrained parameterization, where the positivity constraint in the diagonal elements is often ignored.

underlying constraint is $\Omega = \mathbb{R} \times \mathbb{S}_{++}^1$. $\hat{g}^{(1)}$ and $\hat{g}^{(2)}$ are natural gradients for μ and v , respectively.

$$q(z|\boldsymbol{\lambda}) = \exp \left\{ -\frac{1}{2} \frac{(z - \mu)^2}{v} - \frac{1}{2} \log(2\pi) - \frac{1}{2} \log(v) \right\}$$

Under this parameterization, the FIM and the Christoffel symbols of the second kind are given below, where the Christoffel symbols of the first kind are computed by using Eq. (17). The computation of the Christoffel symbols can be difficult since the parameterization is not a BCN parameterization.

$$F_{ab} = \begin{bmatrix} \frac{1}{v} & 0 \\ 0 & \frac{1}{2v^2} \end{bmatrix}, \quad \Gamma^1_{ab} = \begin{bmatrix} 0 & -\frac{1}{2v} \\ -\frac{1}{2v} & 0 \end{bmatrix}, \quad \Gamma^2_{ab} = \begin{bmatrix} 1 & 0 \\ 0 & -\frac{1}{v} \end{bmatrix}$$

The update suggested by Song et al. (2018) is

$$\begin{aligned} \mu &\leftarrow \mu - t\hat{g}^{(1)} - \frac{t^2}{2} \Gamma^1_{ab} \hat{g}^{(a)} \hat{g}^{(b)} = \mu - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{\hat{g}^{(1)} \hat{g}^{(2)}}{v} \right) \\ v &\leftarrow v - t\hat{g}^{(2)} - \frac{t^2}{2} \Gamma^2_{ab} \hat{g}^{(a)} \hat{g}^{(b)} = v - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{(\hat{g}^{(2)})^2}{v} - (\hat{g}^{(1)})^2 \right) \end{aligned}$$

Obviously, the above updated v does not always satisfy the positivity constraint.

Similarly, we use global indexes such as $\lambda^{(i)} = \lambda^{a_i}$, $\hat{g}^{(i)} = \hat{g}^{[i]}$ and $\Gamma_{i,ii} = \Gamma_{a_i,b_i c_i}$ for notation simplicity since every block contains only a scalar. Note that $\Gamma^1_{11} = 0$ is the entry at the upper-left corner of Γ^1_{ab} and $\Gamma^2_{22} = -\frac{1}{v}$ is the entry at the lower-right corner of Γ^2_{ab} . In our update (see Eq. (16)), we can see the update automatically satisfies the constraint as shown below.

$$\begin{aligned} \mu &\leftarrow \mu - t\hat{g}^{(1)} - \frac{t^2}{2} \Gamma^1_{11} \hat{g}^{(1)} \hat{g}^{(1)} = \mu - t\hat{g}^{(1)} \\ v &\leftarrow v - t\hat{g}^{(2)} - \frac{t^2}{2} \Gamma^2_{22} \hat{g}^{(2)} \hat{g}^{(2)} = v - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{(\hat{g}^{(2)})^2}{v} \right) = \underbrace{\frac{1}{2v}}_{>0} \left[\underbrace{v^2}_{>0} + \underbrace{(v - t\hat{g}^{(2)})^2}_{\geq 0} \right] \end{aligned}$$

B. Riemannian Optimization

B.1. Proof of Lemma 1

Let's consider a parameterization $\boldsymbol{\lambda} := \{\boldsymbol{\lambda}^{[1]}, \dots, \boldsymbol{\lambda}^{[m]}\}$ with m blocks for a statistical manifold with metric \mathbf{F} . We first define a BC parameterization $\boldsymbol{\lambda}$ for a general metric \mathbf{F} .

Definition 1 Block Coordinate Parameterization: A parameterization is block coordinate (BC) if the metric \mathbf{F} under this parameterization is block-diagonal according to the block structure of the parameterization.

Recall that we use the following block notation: $\Gamma^{c_i}_{a_i b_i} \hat{g}^{a_i} \hat{g}^{b_i} := \sum_{a \in [i]} \sum_{b \in [i]} \Gamma^{(c_i)}_{ab} \hat{g}^a \hat{g}^b$ where $[i]$ denotes the index set of block i , (c_i) is the corresponding global index of c_i , and a and b are global indexes.

Now, we prove Lemma 1.

Proof: By the definition of a Riemannian gradient \hat{g} , we have

$$\hat{g}^{a_i} = \sum_b F^{(a_i)b} g_b = \sum_{b \in [i]} F^{(a_i)b} g_b + \sum_{b \notin [i]} \underbrace{F^{(a_i)b}}_0 g_b = \sum_{b \in [i]} F^{(a_i)b} g_b = F^{a_i b_i} g_{b_i},$$

where in the second step, $F^{(a_i)b} = 0$ for any $b \notin [i]$ (see (18) for visualization) since the parameterization is BC, and we use the definition of the block summation notation in the last step.

Similarly, we have

$$\Gamma^{c_i}_{a_i b_i} = \sum_d F^{(c_i)d} \Gamma_{d, (a_i)(b_i)} = \sum_{d \in [i]} F^{(c_i)d} \Gamma_{d, (a_i)(b_i)} + \sum_{d \notin [i]} \underbrace{F^{(c_i)d}}_0 \Gamma_{d, (a_i)(b_i)} = \sum_{d \in [i]} F^{(c_i)d} \Gamma_{d, (a_i)(b_i)} = F^{c_i d_i} \Gamma_{d_i, a_i b_i}$$

□

B.2. NGD is a First-order Approximation of $\mathbf{R}(t)$

Now, we assume parameterization $\lambda = \{\lambda^{[1]}, \dots, \lambda^{[m]}\}$ is a BC parameterization with m blocks. Recall that we define the curve $\mathbf{R}(t)$ as $\mathbf{R}(t) := \{\mathbf{R}^{[1]}(t), \dots, \mathbf{R}^{[m]}(t)\}$, where $\mathbf{R}^{[i]}(t)$ is the solution of following ODE for block i .

$$\begin{aligned}\dot{R}^{c_i}(0) &= -F^{c_i a_i} g_{a_i} ; \quad R^{c_i}(0) = \lambda^{c_i} \\ \ddot{R}^{c_i}(t) &= -\Gamma_{a_i b_i}^{c_i}(t) \dot{R}^{a_i}(t) \dot{R}^{b_i}(t)\end{aligned}$$

where $R^{c_i}(0)$, $\dot{R}^{c_i}(0)$, $\ddot{R}^{c_i}(t)$ respectively denote the c -th entry of $\mathbf{R}^{[i]}(0)$, $\dot{\mathbf{R}}^{[i]}(0)$, and $\ddot{\mathbf{R}}^{[i]}(t)$; $\Gamma_{a_i b_i}^{c_i}(t) := \Gamma_{a_i b_i}^{c_i} \big|_{\lambda^{[-i]} = \mathbf{R}^{[-i]}(0)}^{\lambda^{[-i]} = \mathbf{R}^{[-i]}(t)}$.

Recall that $F^{c_i a_i}$ is the entry of $(\mathbf{F}^{[i]})^{-1}$ at position (c, a) , where $\mathbf{F}^{[i]}$ is the i -th block of \mathbf{F} . Note that \mathbf{F} and $\hat{\mathbf{g}}$ are computed at $\lambda = \mathbf{R}(0)$. Since λ is a BC parameterization, by Lemma 1, we have $F^{c_i a_i} g_{a_i} = \hat{g}^{c_i}$.

Therefore, when \mathbf{F} is the FIM, the first-order approximation of $\mathbf{R}(t)$ at $t_0 = 0$ is also a NGD update as shown below.

$$\begin{aligned}\lambda^{c_i} &\leftarrow R^{c_i}(t_0) + \dot{R}^{c_i}(t_0)(t - t_0) \\ &= \lambda^{c_i} - t \hat{g}^{c_i}\end{aligned}$$

C. Summary of Approximations Considered in This Work

Table 2. Summary of the Proposed Updates Induced by Our Rule in Various Approximations

Approximation	Parameterization (λ)	Constraints	Additional Term
Inverse Gaussian (Appendix H)	$\lambda^{(1)} = \beta^2$ $\lambda^{(2)} = \alpha$	$\lambda^{(1)} \in \mathbb{S}_{++}^1$ $\lambda^{(2)} \in \mathbb{S}_{++}^1$	$\frac{t^2}{2} \left(\frac{3}{4\lambda^{(1)}} \right) \left(\hat{g}^{(1)} \right)^2$ $\frac{t^2}{2} \left(\frac{1}{\lambda^{(2)}} \right) \left(\hat{g}^{(2)} \right)^2$
Gamma (Appendix F)	$\lambda^{(1)} = \alpha$ $\lambda^{(2)} = \frac{\beta}{\alpha}$	$\lambda^{(1)} \in \mathbb{S}_{++}^1$ $\lambda^{(2)} \in \mathbb{S}_{++}^1$	$-\frac{t^2}{2} \frac{\partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)}) + \frac{1}{(\lambda^{(1)})^2}}{\partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}}} \left(\hat{g}^{(1)} \right)^2$ $\frac{t^2}{2} \left(\frac{1}{\lambda^{(2)}} \right) \left(\hat{g}^{(2)} \right)^2$
Exponential (Appendix G)	$\lambda^{(1)} = \lambda$	$\lambda^{(1)} \in \mathbb{S}_{++}^1$	$\frac{t^2}{2} \left(\frac{1}{\lambda^{(1)}} \right) \left(\hat{g}^{(1)} \right)^2$
Multivariate Gaussian (Appendix E)	$\lambda^{[1]} = \mu$ $\lambda^{[2]} = \Sigma^{-1}$	$\lambda^{[1]} \in \mathbb{R}^d$ $\lambda^{[2]} \in \mathbb{S}_{++}^{d \times d}$	$\mathbf{0}$ $\frac{t^2}{2} \hat{\mathbf{g}}^{[2]} \left(\lambda^{[2]} \right)^{-1} \hat{\mathbf{g}}^{[2]}$
Mixture of Gaussians (Appendix J)	$\{\lambda_c^{[1]}\}_{c=1}^K = \{\mu_c\}_{c=1}^K$ $\{\lambda_c^{[2]}\}_{c=1}^K = \{\Sigma_c^{-1}\}_{c=1}^K$ $\lambda_w = \{\log(\pi_c / (1 - \sum_{k=1}^{K-1} \pi_k))\}_{c=1}^{K-1}$	$\lambda_c^{[1]} \in \mathbb{R}^d$ $\lambda_c^{[2]} \in \mathbb{S}_{++}^{d \times d}$ $\lambda_w \in \mathbb{R}^{K-1}$	$\mathbf{0}$ $\frac{t^2}{2} \hat{\mathbf{g}}_c^{[2]} \left(\lambda_c^{[2]} \right)^{-1} \hat{\mathbf{g}}_c^{[2]}$ $\mathbf{0}^{18}$
Skew Gaussian (Appendix K)	$\lambda^{[1]} = \begin{bmatrix} \mu \\ \alpha \end{bmatrix}$ $\lambda^{[2]} = \Sigma^{-1}$	$\lambda^{[1]} \in \mathbb{R}^{2d}$ $\lambda^{[2]} \in \mathbb{S}_{++}^{d \times d}$	$\mathbf{0}$ $\frac{t^2}{2} \hat{\mathbf{g}}^{[2]} \left(\lambda^{[2]} \right)^{-1} \hat{\mathbf{g}}^{[2]}$

Recall that we give Assumption 1-3 for exponential family distributions in Section 3. We also extend Assumption 1-3 to exponential family mixtures as shown in Appendix I.

In Appendix H, F, G, E, J, K, we show that Assumption 1-3 are satisfied and the additional term for each approximation is simplified. In the corresponding appendix, we also show how to compute natural gradients with the (implicit) reparameterization trick for each approximation listed in Table 2.

¹⁸We do not compute the additional term in MOG since $\lambda_w \in \mathbb{R}^{K-1}$ is unconstrained.

D. Exponential Family (EF) Approximation

D.1. Christoffel Symbols

We first show how to simplify the Christoffel symbols of the first kind. The FIM and the corresponding Christoffel symbols of the first kind are defined as follows.

$$F_{ab} := -\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_a \partial_b \log q(\mathbf{z}|\boldsymbol{\lambda})]; \quad \Gamma_{d,ab} := \frac{1}{2} [\partial_a F_{bd} + \partial_b F_{ad} - \partial_d F_{ab}]$$

where we denote $\partial_a = \partial_{\lambda^a}$ for notation simplicity.

Since $\partial_a F_{bd} = -\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_b \partial_d \log q(\mathbf{z}|\boldsymbol{\lambda}) \partial_a \log q(\mathbf{z}|\boldsymbol{\lambda})] - \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_a \partial_b \partial_d \log q(\mathbf{z}|\boldsymbol{\lambda})]$, the Christoffel symbols of the first kind induced by the FIM can be computed as follows, where $\boldsymbol{\lambda}$ can be any parameterization.

$$\begin{aligned} \Gamma_{d,ab} = \frac{1}{2} & \left[\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_a \partial_b \log q(\mathbf{z}|\boldsymbol{\lambda}) \partial_d \log q(\mathbf{z}|\boldsymbol{\lambda})] - \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_b \partial_d \log q(\mathbf{z}|\boldsymbol{\lambda}) \partial_a \log q(\mathbf{z}|\boldsymbol{\lambda})] \right. \\ & \left. - \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_a \partial_d \log q(\mathbf{z}|\boldsymbol{\lambda}) \partial_b \log q(\mathbf{z}|\boldsymbol{\lambda})] - \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_a \partial_b \partial_d \log q(\mathbf{z}|\boldsymbol{\lambda})] \right] \end{aligned} \quad (17)$$

Note that Eq 17 is also applied to a general distribution beyond exponential family. However, the Christoffel symbol is not easy to compute due to extra integrations in Eq 17 and the FIM can be singular in general. The Christoffel symbol could be easy to compute for an exponential family distribution under a BCN parameterization since we compute the symbol via differentiation without the extra integrations. Moreover, the FIM is always positive-definite under a BCN parameterization. Theorem 3 show this.

D.2. Proof of Theorem 3

In this case, $q(\mathbf{z}|\boldsymbol{\lambda})$ is an EF distribution. Since $\boldsymbol{\lambda}$ is a BCN parameterization, given that $\boldsymbol{\lambda}^{[-i]}$ is known, $q(\mathbf{z}|\boldsymbol{\lambda})$ is a one-parameter EF distribution as

$$q(\mathbf{z}|\boldsymbol{\lambda}) = h_i(\mathbf{z}, \boldsymbol{\lambda}^{[-i]}) \exp \left[\langle \phi_i(\mathbf{z}, \boldsymbol{\lambda}^{[-i]}), \boldsymbol{\lambda}^{[i]} \rangle - A(\boldsymbol{\lambda}) \right]$$

Therefore, we have the following identities given $\boldsymbol{\lambda}^{[-i]}$ is known.

$$\partial_{a_i} \partial_{b_i} \log q(\mathbf{z}|\boldsymbol{\lambda}) = -\partial_{a_i} \partial_{b_i} A(\boldsymbol{\lambda}); \quad \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{a_i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = 0$$

where $\partial_{a_i} = \partial_{\lambda^{a_i}}$ for notation simplicity.

Using the above identities, we have

$$\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{a_i} \partial_{b_i} \log q(\mathbf{z}|\boldsymbol{\lambda}) \partial_{d_i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = -\partial_{a_i} \partial_{b_i} A(\boldsymbol{\lambda}) \underbrace{\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{d_i} \log q(\mathbf{z}|\boldsymbol{\lambda})]}_0 = 0$$

Therefore, by Eq. (17), $\Gamma_{d_i, a_i b_i}$ can be computed as follows

$$\Gamma_{d_i, a_i b_i} = -\frac{1}{2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{a_i} \partial_{b_i} \partial_{d_i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = \frac{1}{2} \partial_{a_i} \partial_{b_i} \partial_{d_i} A(\boldsymbol{\lambda})$$

Let $\mathbf{m}_{[i]} = \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\phi_i(\mathbf{z})]$ denote the block coordinate expectation (BCE) parameter. We have

$$0 = \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{a_i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = m_{a_i} - \partial_{a_i} A(\boldsymbol{\lambda})$$

where m_{a_i} denotes the a -th element of $\mathbf{m}_{[i]}$.

Therefore, we know that $m_{a_i} = \partial_{a_i} A(\boldsymbol{\lambda})$

Recall that the i -th block of \mathbf{F} denoted by $\mathbf{F}^{[i]}$, can be computed as

$$F_{a_i b_i} = -\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{b_i} \partial_{a_i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = \partial_{b_i} \partial_{a_i} A(\boldsymbol{\lambda}) = \partial_{b_i} [\partial_{a_i} A(\boldsymbol{\lambda})] = \partial_{\lambda^{b_i}} m_{a_i}$$

where $\partial_{b_i} = \partial_{\lambda^{b_i}}$ is for notation simplicity.

Recall that λ is a BC parameterization with n blocks and \mathbf{F} is block diagonal as shown below.

$$\mathbf{F} = \begin{bmatrix} \mathbf{F}^{[1]} & \dots & \mathbf{0} \\ \vdots & \ddots & \vdots \\ \mathbf{0} & \dots & \mathbf{F}^{[n]} \end{bmatrix} \quad (18)$$

Recall that F^{ab} denotes the element of \mathbf{F}^{-1} with global index (a, b) and $F^{a_i b_i}$ denotes the element of $(\mathbf{F}^{[i]})^{-1}$ with local index (a, b) in block i .

If $\mathbf{F}^{[i]}$ is positive definite everywhere, we have

$$F^{a_i b_i} = \partial_{m_{a_i}} \lambda^{b_i}$$

Note that $\mathbf{F}^{[i]}$ is positive definite everywhere when $q(\mathbf{z}|\lambda^{[i]}, \lambda^{[-i]})$ is a one-parameter minimal EF distribution given $\lambda^{[-i]}$ is known (See Theorem 1 of Lin et al. (2019a)).

By Lemma 1, Riemannian gradient \hat{g}^{a_i} can be computed as

$$\hat{g}^{a_i} = F^{a_i b_i} g_{b_i} = [\partial_{m_{a_i}} \lambda^{b_i}] [\partial_{\lambda^{b_i}} \mathcal{L}] = \partial_{m_{a_i}} \mathcal{L}$$

where $g_{b_i} = \partial_{\lambda^{b_i}} \mathcal{L}$ is a Euclidean gradient.

E. Example: Gaussian Approximation

We consider the following parameterization $\lambda = \{\mu, \mathbf{S}\}$, where μ is the mean and \mathbf{S} is the precision. The open-set constraint is $\Omega_1 = \mathbb{R}^d$ and $\Omega_2 = \mathbb{S}_{++}^{d \times d}$. Under this parameterization, the distribution can be expressed as below.

$$q(\mathbf{z}|\lambda) = \exp \left(-\frac{1}{2} \mathbf{z}^T \mathbf{S} \mathbf{z} + \mathbf{z}^T \mathbf{S} \mu - A(\lambda) \right)$$

where $A(\lambda) = \frac{1}{2} [\mu^T \mathbf{S} \mu - \log |\mathbf{S}|/(2\pi)]$

Lemma 2 *The Fisher information matrix under this parameterization is block diagonal with two blocks*

$$\mathbf{F} = \begin{bmatrix} \mathbf{F}_\mu & \mathbf{0} \\ \mathbf{0} & \mathbf{F}_S \end{bmatrix},$$

$\mathbf{F}_{\mu S}$

where $\mathbf{F}_{\mu S} = -\mathbb{E}_{q(\mathbf{z})} [\partial_{\text{vec}(\mathbf{S})} \partial_\mu \log q(\mathbf{z}|\mu, \mathbf{S})]$ and $\mathbf{F}_S = -\mathbb{E}_{q(\mathbf{z})} [\partial_{\text{vec}(\mathbf{S})}^2 \log q(\mathbf{z}|\mu, \mathbf{S})]$.

Therefore, $\lambda = \{\mu, \mathbf{S}\}$ is a BC parameterization.

Proof: We denote the i -th element of μ using μ^i . Similarly, we denote the element of \mathbf{S} at position (j, k) using S^{jk} . We prove this statement by showing cross terms in the Fisher information matrix denoted by $\mathbf{F}_{\mu S}$ are all zeros. To show $\mathbf{F}_{\mu S} = \mathbf{0}$, it is equivalent to show $-\mathbb{E}_{q(\mathbf{z}|\lambda)} [\partial_{S^{jk}} \partial_{\mu^i} \log q(\mathbf{z}|\lambda)] = 0$ each μ^i and S^{jk} .

Notice that $\mathbb{E}_{q(\mathbf{z}|\lambda)} [\mathbf{z}] = \mu$. We can obtain the above expression since

$$\begin{aligned} \mathbb{E}_{q(\mathbf{z}|\lambda)} [\partial_{S^{jk}} \partial_{\mu^i} \log q(\mathbf{z}|\lambda)] &= \mathbb{E}_{q(\mathbf{z}|\lambda)} [\partial_{S^{jk}} (\mathbf{z}^T \mathbf{S} \mathbf{e}_i - \mathbf{e}_i^T \mathbf{S} \mu)] \\ &= \mathbb{E}_{q(\mathbf{z}|\lambda)} [(\mathbf{z}^T \mathbf{I}_{jk} \mathbf{e}_i - \mathbf{e}_i^T \mathbf{I}_{jk} \mu)] \\ &= \mathbb{E}_{q(\mathbf{z}|\lambda)} [(\mathbf{e}_i^T \mathbf{I}_{jk} (\mathbf{z} - \mu))] \\ &= \mathbf{e}_i^T \mathbf{I}_{jk} \underbrace{\mathbb{E}_{q(\mathbf{z}|\lambda)} [\mathbf{z} - \mu]}_{\mathbf{0}} = 0 \end{aligned}$$

where \mathbf{e}_i denotes an one-hot vector where all entries are zeros except the i -th entry with value 1, and \mathbf{I}_{jk} denotes an one-hot matrix where all entries are zeros except the entry at position (j, k) with value 1.

The above expression also implies that $\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_S \partial_{\mu^i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = \mathbf{0}$. \square

Now, we show that $\boldsymbol{\lambda} = \{\boldsymbol{\mu}, \boldsymbol{\Sigma}\}$ is also a BC parameterization. Note that

$$-\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\Sigma^{jk}} \partial_{\mu^i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = -\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\text{Tr}\{(\partial_{\Sigma^{jk}} \mathbf{S}) \partial_S \partial_{\mu^i} \log q(\mathbf{z}|\boldsymbol{\lambda})\}] = -\text{Tr}\{(\partial_{\Sigma^{jk}} \mathbf{S}) \underbrace{\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_S \partial_{\mu^i} \log q(\mathbf{z}|\boldsymbol{\lambda})]}_{\mathbf{0}}\} = 0.$$

Since $\mathbf{F}_{\mu\Sigma} = -\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\text{vec}(\Sigma)} \partial_{\mu} \log q(\mathbf{z}|\boldsymbol{\lambda})]$ and $-\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\Sigma^{jk}} \partial_{\mu^i} \log q(\mathbf{z}|\boldsymbol{\lambda})] = 0$ from above expression for any i, j , and k , we have $\mathbf{F}_{\mu\Sigma} = \mathbf{0}$. Therefore, $\boldsymbol{\lambda} = \{\boldsymbol{\mu}, \boldsymbol{\Sigma}\}$ is also a BC parameterization since the cross terms of FIM under this new parameterization denoted by $\mathbf{F}_{\mu\Sigma}$ are zeros.

We denote the Christoffel symbols of the first kind and the second kind for $\boldsymbol{\mu}$ as $\Gamma_{a_1, b_1 c_1}$ and $\Gamma^{a_1}_{b_1 c_1}$, respectively.

Lemma 3 *All entries of $\Gamma^{a_1}_{b_1 c_1}$ are zeros.*

Proof: We will prove this by showing that all entries of $\Gamma_{a_1, b_1 c_1}$ are zeros. For notation simplicity, we use $\Gamma_{a, bc}$ to denote $\Gamma_{a_1, b_1 c_1}$ in the proof. Let μ^a denote the a -th element of $\boldsymbol{\mu}$. The following expression holds for any valid a, b , and c .

$$\Gamma_{a, bc} = \frac{1}{2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\mu^b} \partial_{\mu^c} \partial_{\mu^a} A(\boldsymbol{\lambda})] = 0$$

We can obtain the above expression since

$$\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\mu^b} \partial_{\mu^c} \partial_{\mu^a} A(\boldsymbol{\lambda})] = \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\mu^b} \partial_{\mu^c} (\mathbf{e}_a^T \mathbf{S} \boldsymbol{\mu})] = \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\partial_{\mu^b} (\mathbf{e}_a^T \mathbf{S} \mathbf{e}_c)] = 0$$

where in the last step we use the fact that \mathbf{S} , \mathbf{e}_a , and \mathbf{e}_c do not depend on $\boldsymbol{\mu}$. \square

Similarly, we denote the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S})$ as $\Gamma^{a_2}_{b_2 c_2}$. Note that \mathbf{S} is now a matrix. It is possible but tedious to directly compute the Christoffel symbol and element-wisely validate the expression of the additional term for \mathbf{S} . Below, we give an alternative approach to identify the additional term for \mathbf{S} as shown in the proof of Lemma 4.

Recall that $\mathbf{R}^{[2]}(t)$ is the solution of the following ODE for block $\text{vec}(\mathbf{S})$:

$$\begin{aligned} \dot{R}^{a_2}(0) &= -\hat{g}^{a_2} ; \quad R^{a_2}(0) = S^{a_2} \\ \ddot{R}^{a_2}(t) &= -\Gamma^{a_2}_{b_2 c_2}(t) \dot{R}^{b_2}(t) \dot{R}^{c_2}(t), \end{aligned}$$

where $R^{a_2}(t)$ denotes the a -th element of $\mathbf{R}^{[2]}(t)$ and S^{a_2} denotes the a -th entry of $\text{vec}(\mathbf{S})$.

Lemma 4 *The additional term for \mathbf{S} is $\text{Mat}(\Gamma^{a_2}_{b_2 c_2} \hat{g}^{b_2} \hat{g}^{c_2}) = -\hat{\mathbf{g}}^{[2]} \mathbf{S}^{-1} \hat{\mathbf{g}}^{[2]}$ where \hat{g}^{a_2} denotes the a -th element of $\text{vec}(\hat{\mathbf{g}}^{[2]})$.*

Proof: As discussed in Sec 5, $\mathbf{R}^{[i]}(t)$ is a (block coordinate) geodesic given $\boldsymbol{\lambda}^{[-i]}$ is known. In this case, given that $\boldsymbol{\mu}$ is known, $\mathbf{R}^{[2]}(t)$ has the following closed-form expression (Pennec et al., 2006; Fletcher & Joshi, 2004; Minh & Murino, 2017).

$$\text{Mat}(\mathbf{R}^{[2]}(t)) = \mathbf{U} \text{Exp}(t \mathbf{U}^{-1} \hat{\mathbf{g}}^{[2]} \mathbf{U}^{-1}) \mathbf{U}$$

where $\mathbf{U} = \mathbf{S}^{\frac{1}{2}}$ denotes the matrix square root and $\text{Exp}(\mathbf{X}) := \mathbf{I} + \sum_{n=1}^{\infty} \frac{\mathbf{X}^n}{n!}$ denotes the matrix exponential function.¹⁹

¹⁹The function is well-defined since the matrix series is absolutely convergent element-wisely.

The additional term for \mathbf{S} can be obtained as follows.

$$\begin{aligned}
 -\text{Mat}(\Gamma^{a_2}_{b_2 c_2} \hat{g}^{b_2} \hat{g}^{c_2}) &= \text{Mat}(\ddot{\mathbf{R}}^{[2]}(0)) \\
 &= \text{Mat}(\nabla_t^2 \mathbf{R}^{[2]}(t)|_{t=0}) \\
 &= \nabla_t^2 \text{Mat}(\mathbf{R}^{[2]}(t))|_{t=0} \\
 &= \nabla_t^2 \left(\mathbf{U} \text{Exp}(\mathbf{U}^{-1} t \hat{\mathbf{g}}^{[2]} \mathbf{U}^{-1}) \mathbf{U} \right)|_{t=0} \\
 &= \mathbf{U} \nabla_t^2 \left(\text{Exp}(\mathbf{U}^{-1} t \hat{\mathbf{g}}^{[2]} \mathbf{U}^{-1}) \right)|_{t=0} \mathbf{U} \\
 &= \mathbf{U} (\mathbf{U}^{-1} \hat{\mathbf{g}}^{[2]} \mathbf{U}^{-1}) (\mathbf{U}^{-1} \hat{\mathbf{g}}^{[2]} \mathbf{U}^{-1}) \mathbf{U} \\
 &= \mathbf{U} (\mathbf{U}^{-1} \hat{\mathbf{g}}^{[2]} \mathbf{S}^{-1} \hat{\mathbf{g}}^{[2]} \mathbf{U}^{-1}) \mathbf{U} \\
 &= \hat{\mathbf{g}}^{[2]} \mathbf{S}^{-1} \hat{\mathbf{g}}^{[2]}
 \end{aligned}$$

where we use the following expression to move from step 5 to step 6.

$$\nabla_t^2 \text{Exp}(t\mathbf{X})|_{t=0} = \nabla_t^2 \left(\mathbf{I} + \sum_{n=1}^{\infty} \frac{(t\mathbf{X})^n}{n!} \right)|_{t=0} = \mathbf{X}^2$$

□

Finally, by Lemma 3 and 4, the update induced by the proposed rule is

$$\begin{aligned}
 \mu^c &\leftarrow \mu^c - t \hat{g}^{c_1} - \frac{t \times t}{2} \overbrace{\Gamma^{c_1}_{a_1 b_1}}^0 \hat{g}^{a_1} \hat{g}^{b_1} \\
 s^c &\leftarrow s^c - t \hat{g}^{c_2} - \frac{t \times t}{2} \Gamma^{c_2}_{a_2 b_2} \hat{g}^{a_2} \hat{g}^{b_2}
 \end{aligned}$$

where s^c is the c -th element of $\text{vec}(\mathbf{S})$.

Therefore, we have

$$\begin{aligned}
 \boldsymbol{\mu} &\leftarrow \underbrace{\boldsymbol{\mu}}_{\text{vec}(\boldsymbol{\mu}^c)} - t \underbrace{\hat{\mathbf{g}}^{[1]}}_{\text{vec}(\hat{\mathbf{g}}^{c_1})} \\
 \mathbf{S} &\leftarrow \underbrace{\mathbf{S}}_{\text{Mat}(s^c)} - t \underbrace{\hat{\mathbf{g}}^{[2]}}_{\text{Mat}(\hat{\mathbf{g}}^{c_2})} + \frac{t \times t}{2} \underbrace{\hat{\mathbf{g}}^{[2]} \mathbf{S}^{-1} \hat{\mathbf{g}}^{[2]}}_{-\text{Mat}(\Gamma^{c_2}_{a_2 b_2} \hat{g}^{a_2} \hat{g}^{b_2})}
 \end{aligned}$$

E.1. Proof of Theorem 1

Now, we give a proof of Theorem 1.

Proof: First note that $\hat{\mathbf{G}} = \mathbf{S} - \mathbb{E}_q[\nabla_z^2 \bar{\ell}(\mathbf{z})]$ is a symmetric matrix. Let \mathbf{L} be the Cholesky of the current $\mathbf{S} = \mathbf{L}\mathbf{L}^T$. We can simplify the right hand side of (9) as follows:

$$(1-t)\mathbf{S} + t\mathbb{E}_q[\nabla_z^2 \bar{\ell}(\mathbf{z})] + \frac{t^2}{2} \hat{\mathbf{G}} \mathbf{S}^{-1} \hat{\mathbf{G}} = \mathbf{S} - t\hat{\mathbf{G}} + \frac{t^2}{2} \hat{\mathbf{G}} \mathbf{S}^{-1} \hat{\mathbf{G}} = \frac{1}{2} \left(\mathbf{S} + \left(\mathbf{L} - t\hat{\mathbf{G}}\mathbf{L}^{-T} \right) \left(\mathbf{L}^T - t\mathbf{L}^{-1}\hat{\mathbf{G}} \right) \right) = \frac{1}{2} \left(\mathbf{S} + \mathbf{U}^T \mathbf{U} \right),$$

where $\mathbf{U} := \mathbf{L}^T - t\mathbf{L}^{-1}\hat{\mathbf{G}}$. Since the current \mathbf{S} is positive-definite, and $\mathbf{U}^T \mathbf{U}$ is positive semi-definite, we know that the update for \mathbf{S} is positive-definite. □

E.2. Natural Gradients and the Reparameterization Trick

Since $\boldsymbol{\lambda} = \{\boldsymbol{\mu}, \mathbf{S}\}$ is a BCN parameterization of an exponential family distribution, gradients w.r.t. BC expectation parameters are natural gradients for BC natural parameters as shown in Theorem 3.

Given that \mathbf{S} is known, the BC expectation parameter is $\mathbf{m}_{[1]} = \mathbb{E}_{q(\mathbf{z})} [\mathbf{S}\mathbf{z}] = \mathbf{S}\boldsymbol{\mu}$. In this case, we know that $\partial_{\boldsymbol{\mu}}\mathcal{L} = \mathbf{S}\partial_{\mathbf{m}_{[1]}}\mathcal{L}$. Therefore, the natural gradient w.r.t. $\boldsymbol{\mu}$ is $\hat{\mathbf{g}}^{[1]} = \partial_{\mathbf{m}_{[1]}}\mathcal{L} = \mathbf{S}^{-1}\partial_{\boldsymbol{\mu}}\mathcal{L} = \boldsymbol{\Sigma}\partial_{\boldsymbol{\mu}}\mathcal{L}$.

Likewise, given that $\boldsymbol{\mu}$ is known, the BC expectation parameter is $\mathbf{m}_{[2]} = \mathbb{E}_{q(\mathbf{z})} [-\frac{1}{2}\mathbf{z}\mathbf{z}^T + \boldsymbol{\mu}\mathbf{z}^T] = \frac{1}{2}(\boldsymbol{\mu}\boldsymbol{\mu}^T - \mathbf{S}^{-1})$. Therefore, the natural gradient w.r.t. \mathbf{S} is $\hat{\mathbf{g}}^{[2]} = \partial_{\mathbf{m}_{[2]}}\mathcal{L} = -2\partial_{\mathbf{S}^{-1}}\mathcal{L} = -2\partial_{\boldsymbol{\Sigma}}\mathcal{L}$.

Recall that $\mathcal{L}(\boldsymbol{\lambda}) = \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z}) + \log q(\mathbf{z}|\boldsymbol{\lambda})]$, by the Gaussian identities (Opper & Archambeau, 2009; Särkkä, 2013) (see Lin et al. (2019b) for a derivation of these identities), we have

$$\begin{aligned}\partial_{\boldsymbol{\mu}}\mathcal{L}(\boldsymbol{\lambda}) &= \partial_{\boldsymbol{\mu}} \left[\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})] - \frac{1}{2} \log |2\pi e \boldsymbol{\Sigma}| \right] \\ &= \partial_{\boldsymbol{\mu}} [\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]] \\ &= \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\nabla_{\mathbf{z}} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]]\end{aligned}\tag{19}$$

$$\begin{aligned}\partial_{\boldsymbol{\Sigma}}\mathcal{L}(\boldsymbol{\lambda}) &= \partial_{\boldsymbol{\Sigma}} \left[\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})] - \frac{1}{2} \log |2\pi e \boldsymbol{\Sigma}| \right] \\ &= \partial_{\boldsymbol{\Sigma}} [\mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]] - \frac{1}{2} \boldsymbol{\Sigma}^{-1} \\ &= \frac{1}{2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\boldsymbol{\Sigma}^{-1} (\mathbf{z} - \boldsymbol{\mu}) \nabla_{\mathbf{z}}^T [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]] - \frac{1}{2} \boldsymbol{\Sigma}^{-1}\end{aligned}\tag{20}$$

$$= \frac{1}{2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\lambda})} [\nabla_{\mathbf{z}}^2 [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]] - \frac{1}{2} \boldsymbol{\Sigma}^{-1}\tag{21}$$

where (19) is also known as the reparameterization trick for the mean, (20) is also known as the reparameterization trick for the covariance, and we call (21) the Hessian trick.

Using Monte Carlo approximation, we have

$$\begin{aligned}\partial_{\boldsymbol{\mu}}\mathcal{L} &\approx \nabla_{\mathbf{z}} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})] \\ \partial_{\boldsymbol{\Sigma}}\mathcal{L} &\approx \frac{1}{4} [\bar{\mathbf{S}} + \bar{\mathbf{S}}^T] - \frac{1}{2} \boldsymbol{\Sigma}^{-1} \quad \text{referred to as “-rep”} \\ \partial_{\boldsymbol{\Sigma}}\mathcal{L} &\approx \frac{1}{2} [\nabla_{\mathbf{z}}^2 [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]] - \frac{1}{2} \boldsymbol{\Sigma}^{-1} \quad \text{referred to as “-hess”}\end{aligned}$$

where $\bar{\mathbf{S}} := \boldsymbol{\Sigma}^{-1}(\mathbf{z} - \boldsymbol{\mu}) \nabla_{\mathbf{z}}^T [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z})]$ and $\mathbf{z} \sim q(\mathbf{z}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}, \boldsymbol{\Sigma})$.

E.3. Adam-like Update

We consider to solve the following problem, where we use a diagonal Gaussian approximation $q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})$ and $\mathbf{s} = \boldsymbol{\sigma}^{-2}$.

$$\min_{\boldsymbol{\mu}, \mathbf{s}} \mathcal{L}(\boldsymbol{\mu}, \mathbf{s}) = \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} \left[\left(\sum_{i=1}^N \ell_i(\mathbf{z}) \right) - \log \mathcal{N}(\mathbf{z}|\mathbf{0}, \lambda^{-1} \mathbf{I}) + \log q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s}) \right]$$

Note that

$$\begin{aligned}\partial_{\boldsymbol{\mu}}\mathcal{L}(\boldsymbol{\mu}, \mathbf{s}) &:= \sum_{i=1}^N \partial_{\boldsymbol{\mu}} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})] + \lambda \boldsymbol{\mu} \\ \partial_{\sigma^2}\mathcal{L}(\boldsymbol{\mu}, \mathbf{s}) &:= \sum_{i=1}^N \partial_{\sigma^2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})] + \frac{1}{2} \lambda - \frac{1}{2} \mathbf{s}\end{aligned}$$

where $\partial_{\boldsymbol{\mu}} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})]$ and $\partial_{\sigma^2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})]$ can be computed by the reparameterization trick with MC approximations where $\mathbf{z} \sim \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})$.

$$\begin{aligned}\partial_{\boldsymbol{\mu}} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})] &= \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\nabla_{\mathbf{z}} \ell_i(\mathbf{z})] \approx \nabla_{\mathbf{z}} \ell_i(\mathbf{z}) \\ \partial_{\sigma^2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})] &= \frac{1}{2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\mathbf{s} \odot (\mathbf{z} - \boldsymbol{\mu}) \odot \nabla_{\mathbf{z}} \ell_i(\mathbf{z})] \approx \frac{1}{2} [\mathbf{s} \odot (\mathbf{z} - \boldsymbol{\mu})] \odot \nabla_{\mathbf{z}} \ell_i(\mathbf{z})\end{aligned}$$

The natural gradients can be computed as follows.

$$\begin{aligned}\hat{\mathbf{g}}_k^{[1]} &= \boldsymbol{\sigma}_k^2 \left(\partial_{\boldsymbol{\mu}} \mathcal{L}(\boldsymbol{\mu}, \mathbf{s}) \Big|_{\boldsymbol{\mu}=\boldsymbol{\mu}_k, \mathbf{s}=\mathbf{s}_k} \right) \\ \hat{\mathbf{g}}_k^{[2]} &= -2 \partial_{\sigma^2} \mathcal{L}(\boldsymbol{\mu}, \mathbf{s}) \Big|_{\boldsymbol{\mu}=\boldsymbol{\mu}_k, \mathbf{s}=\mathbf{s}_k}\end{aligned}$$

The update induced by our rule with exponential decaying step-sizes and the natural momentum (Khan et al., 2018) shown in blue is given as follows.

$$\begin{aligned}\boldsymbol{\mu}_{k+1} &= \boldsymbol{\mu}_k - t_1 \hat{\mathbf{g}}_k^{[1]} + t_2 \boldsymbol{\sigma}_k^2 \odot \boldsymbol{\sigma}_{k-1}^{-2} \odot (\boldsymbol{\mu}_k - \boldsymbol{\mu}_{k-1}) \\ \boldsymbol{\sigma}_{k+1}^{-2} &= \boldsymbol{\sigma}_k^{-2} - t_3 \hat{\mathbf{g}}_k^{[2]} + \frac{t_3^2}{2} \hat{\mathbf{g}}_k^{[2]} \odot \boldsymbol{\sigma}_k^2 \odot \hat{\mathbf{g}}_k^{[2]}\end{aligned}$$

where $t_1 = t(1 - r_1) \frac{1-r_2^k}{1-r_1^k}$, $t_2 = r_1 \frac{1-r_2^k}{1-r_1^k} \frac{1-r_1^{k-1}}{1-r_2^{k-1}}$, and $t_3 = (1 - r_2)$.

Recall that $\mathbf{s} = \boldsymbol{\sigma}^{-2}$. The proposed update can be expressed as

$$\begin{aligned}\boldsymbol{\mu}_{k+1} &= \boldsymbol{\mu}_k - t(1 - r_1) \frac{1 - r_2^k}{1 - r_1^k} \hat{\mathbf{s}}_k^{-1} \odot \mathbf{g}_k + r_1 \frac{1 - r_2^k}{1 - r_1^k} \frac{1 - r_1^{k-1}}{1 - r_2^{k-1}} \hat{\mathbf{s}}_k^{-1} \odot \hat{\mathbf{s}}_{k-1} \odot (\boldsymbol{\mu}_k - \boldsymbol{\mu}_{k-1}) \\ \hat{\mathbf{s}}_{k+1} &= \hat{\mathbf{s}}_k + (1 - r_2) \mathbf{h}_k + \frac{(1 - r_2)^2}{2} \mathbf{h}_k \odot \hat{\mathbf{s}}_k^{-1} \odot \mathbf{h}_k \\ \mathbf{s}_{k+1} &= N \hat{\mathbf{s}}_{k+1}\end{aligned}$$

where $\mathbf{g}_k := \frac{1}{N} \sum_{i=1}^N \partial_{\boldsymbol{\mu}} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})] \big|_{\boldsymbol{\mu}=\boldsymbol{\mu}_k, \mathbf{s}=\mathbf{s}_k} + \frac{\lambda}{N} \boldsymbol{\mu}_k$ and $\mathbf{h}_k := \frac{2}{N} \sum_{i=1}^N \partial_{\boldsymbol{\sigma}^2} \mathbb{E}_{q(\mathbf{z}|\boldsymbol{\mu}, \mathbf{s})} [\ell_i(\mathbf{z})] \big|_{\boldsymbol{\mu}=\boldsymbol{\mu}_k, \mathbf{s}=\mathbf{s}_k} + \frac{\lambda}{N} - \hat{\mathbf{s}}_k$.

Let's define $\mathbf{m}_k := \frac{1-r_1^{k-1}}{t(1-r_2^{k-1})} \hat{\mathbf{s}}_{k-1} \odot (\boldsymbol{\mu}_{k-1} - \boldsymbol{\mu}_k)$. We can further simplify the above update as shown below.

$$\begin{aligned}\boldsymbol{\mu}_{k+1} &= \boldsymbol{\mu}_k - t(1 - r_1) \frac{1 - r_2^k}{1 - r_1^k} \hat{\mathbf{s}}_k^{-1} \odot \mathbf{g}_k + tr_1 \frac{1 - r_2^k}{1 - r_1^k} \hat{\mathbf{s}}_k^{-1} \odot \left(\frac{1 - r_1^{k-1}}{t(1 - r_2^{k-1})} \hat{\mathbf{s}}_{k-1} \odot (\boldsymbol{\mu}_k - \boldsymbol{\mu}_{k-1}) \right) \\ &= \boldsymbol{\mu}_k - t \frac{1 - r_2^k}{1 - r_1^k} \hat{\mathbf{s}}_k^{-1} \odot [(1 - r_1) \mathbf{g}_k + r_1 \mathbf{m}_k] \\ \mathbf{m}_{k+1} &= \frac{1 - r_1^k}{t(1 - r_2^k)} \hat{\mathbf{s}}_k \odot (\boldsymbol{\mu}_k - \boldsymbol{\mu}_{k+1}) \\ &= \frac{1 - r_1^k}{t(1 - r_2^k)} t \frac{1 - r_2^k}{1 - r_1^k} [(1 - r_1) \mathbf{g}_k + r_1 \mathbf{m}_k] \\ &= (1 - r_1) \mathbf{g}_k + r_1 \mathbf{m}_k \\ \hat{\mathbf{s}}_{k+1} &= \hat{\mathbf{s}}_k + (1 - r_2) \mathbf{h}_k + \frac{(1 - r_2)^2}{2} \mathbf{h}_k \odot \hat{\mathbf{s}}_k^{-1} \odot \mathbf{h}_k \\ &= \frac{1}{2} [\hat{\mathbf{s}}_k + (\hat{\mathbf{s}}_k + (1 - r_2) \mathbf{h}_k) \odot \hat{\mathbf{s}}_k^{-1} \odot (\hat{\mathbf{s}}_k + (1 - r_2) \mathbf{h}_k)] \\ \mathbf{s}_{k+1} &= N \hat{\mathbf{s}}_{k+1}\end{aligned}$$

where $\mathbf{z} \sim q(\mathbf{z}|\boldsymbol{\mu}_k, \mathbf{s}_k)$, $\mathbf{g}_k \approx \nabla_{\boldsymbol{\mu}} \ell_i(\mathbf{z}) + \frac{\lambda}{N} \boldsymbol{\mu}_k$, and $\mathbf{h}_k \approx [(N \hat{\mathbf{s}}_k) \odot (\mathbf{z} - \boldsymbol{\mu})] \odot \nabla_{\mathbf{z}} \ell_i(\mathbf{z}) + \frac{\lambda}{N} - \hat{\mathbf{s}}_k$.

E.4. Tran et al. (2019) is a special case of our update

In the Gaussian case, Tran et al. (2019) consider the following update by using parameterization $\boldsymbol{\lambda} = \{\boldsymbol{\mu}, \boldsymbol{\Sigma}\}$, where $\boldsymbol{\Sigma}$ is the covariance matrix.

$$\boldsymbol{\mu} \leftarrow \boldsymbol{\mu} - t \boldsymbol{\Sigma} (\partial_{\boldsymbol{\mu}} \mathcal{L}) \quad (22)$$

$$\boldsymbol{\Sigma} \leftarrow \boldsymbol{\Sigma} - t \hat{\mathbf{g}}^{[2]} + \frac{t \times t}{2} \hat{\mathbf{g}}^{[2]} \boldsymbol{\Sigma}^{-1} \hat{\mathbf{g}}^{[2]} = \text{Ret}(\boldsymbol{\Sigma}, -t \hat{\mathbf{g}}^{[2]}). \quad (23)$$

where the natural gradient²⁰ for $\boldsymbol{\Sigma}$ is $\hat{\mathbf{g}}^{[2]} := 2\boldsymbol{\Sigma}(\partial_{\boldsymbol{\Sigma}} \mathcal{L})\boldsymbol{\Sigma}$ and the retraction map is $\text{Ret}(\boldsymbol{\Sigma}, \mathbf{b}) := \boldsymbol{\Sigma} + \mathbf{b} + \frac{1}{2} \mathbf{b} \boldsymbol{\Sigma}^{-1} \mathbf{b}$.

However, Tran et al. (2019) do not justify the use of the retraction map, which is just one of retraction maps developed for positive definite matrices. In this section, we show that how to derive this update from our rule.

²⁰There is a typo in Algorithm 2 of Tran et al. (2019). The natural gradient for $\boldsymbol{\Sigma}$ should be $2\boldsymbol{\Sigma}(\partial_{\boldsymbol{\Sigma}} \mathcal{L})\boldsymbol{\Sigma}$ instead of $\boldsymbol{\Sigma}(\partial_{\boldsymbol{\Sigma}} \mathcal{L})\boldsymbol{\Sigma}$. Note that the Riemannian gradient for a positive-definite matrix is $\boldsymbol{\Sigma}(\partial_{\boldsymbol{\Sigma}} \mathcal{L})\boldsymbol{\Sigma}$ (see Table 1 of Hosseini & Sra (2015)) while the natural/Riemannian gradient for Gaussian distribution w.r.t. $\boldsymbol{\Sigma}$ is $2\boldsymbol{\Sigma}(\partial_{\boldsymbol{\Sigma}} \mathcal{L})\boldsymbol{\Sigma}$.

As shown in Eq. (16), our rule can be used under not only a BCN parameterization but also a BC parameterization. Now, we show that our rule can recover the above update using the parameterization $\lambda = \{\mu, \Sigma\}$. Recall that this parameterization is a BC parameterization. It only requires us to show that natural gradients and the additional terms are described in Eq. (23).

Given that Σ is known, μ is the natural parameter and the expectation parameter is $\mathbf{m}_{[1]} = \mathbb{E}_{q(z)} [\Sigma^{-1} \mathbf{z}] = \Sigma^{-1} \mu$ as shown in Appendix E.2. Therefore, the natural gradient w.r.t. μ is $\hat{\mathbf{g}}^{[1]} = \partial_{\mathbf{m}_{[1]}} \mathcal{L} = \Sigma \partial_{\mu} \mathcal{L}$.

Now, we show that the natural gradients w.r.t. Σ is

$$\hat{\mathbf{g}}^{[2]} = 2\Sigma(\partial_{\Sigma} \mathcal{L})\Sigma$$

A proof using matrix calculus is provided below. See Malagò & Pistone (2015) for alternative proofs. By matrix calculus, we have

$$\begin{aligned} & -\mathbb{E}_{q(z)} [\partial_{\Sigma^{ij}} \partial_{\Sigma} [\log q(\mathbf{z}|\mu, \Sigma)]] \\ &= \mathbb{E}_{q(z)} [\partial_{\Sigma^{ij}} \partial_{\Sigma} [\frac{1}{2}(\mathbf{z} - \mu)^T \Sigma^{-1} (\mathbf{z} - \mu) + \frac{1}{2} \log |\Sigma|/(2\pi)]] \\ &= \frac{1}{2} \mathbb{E}_{q(z)} [\partial_{\Sigma^{ij}} [-\Sigma^{-1} (\mathbf{z} - \mu)(\mathbf{z} - \mu)^T \Sigma^{-1} + \Sigma^{-1}]] \\ &= \frac{1}{2} \mathbb{E}_{q(z)} [-\partial_{\Sigma^{ij}} [\Sigma^{-1}] (\mathbf{z} - \mu)(\mathbf{z} - \mu)^T \Sigma^{-1} - \Sigma^{-1} (\mathbf{z} - \mu)(\mathbf{z} - \mu)^T \partial_{\Sigma^{ij}} [\Sigma^{-1}] + \partial_{\Sigma^{ij}} [\Sigma^{-1}]] \\ &= \frac{1}{2} \mathbb{E}_{q(z)} [-\partial_{\Sigma^{ij}} [\Sigma^{-1}] (\mathbf{z} - \mu)(\mathbf{z} - \mu)^T \Sigma^{-1} - \Sigma^{-1} (\mathbf{z} - \mu)(\mathbf{z} - \mu)^T \partial_{\Sigma^{ij}} [\Sigma^{-1}] + \partial_{\Sigma^{ij}} [\Sigma^{-1}]] \\ &= -\frac{1}{2} \partial_{\Sigma^{ij}} [\Sigma^{-1}] \underbrace{\mathbb{E}_{q(z)} [(\mathbf{z} - \mu)(\mathbf{z} - \mu)^T]}_{\Sigma} \Sigma^{-1} - \frac{1}{2} \Sigma^{-1} \underbrace{\mathbb{E}_{q(z)} [(\mathbf{z} - \mu)(\mathbf{z} - \mu)^T]}_{\Sigma} \partial_{\Sigma^{ij}} [\Sigma^{-1}] + \frac{1}{2} \partial_{\Sigma^{ij}} [\Sigma^{-1}] \\ &= \frac{1}{2} [-\partial_{\Sigma^{ij}} [\Sigma^{-1}] \mathbf{I} - \mathbf{I} \partial_{\Sigma^{ij}} [\Sigma^{-1}] + \partial_{\Sigma^{ij}} [\Sigma^{-1}]] \\ &= -\frac{1}{2} \partial_{\Sigma^{ij}} [\Sigma^{-1}] \end{aligned}$$

Therefore, the block matrix of the FIM related to Σ is $\mathbf{F}_{\Sigma} := -\mathbb{E}_{q(z)} [\partial_{\text{vec}(\Sigma)}^2 [\log q(\mathbf{z}|\mu, \Sigma)]] = -\frac{1}{2} \partial_{\text{vec}(\Sigma)} [\text{vec}(\Sigma^{-1})]$ due to the above expression. Note that $\mathbf{F}_{\Sigma}^{-1} = -2\partial_{\text{vec}(\Sigma^{-1})} [\text{vec}(\Sigma)]$.

Note that $\hat{\mathbf{g}}^{[2]}$ is the natural gradient for Σ . Since $\lambda = \{\mu, \Sigma\}$ is a BC parameterization, by Lemma 1, the natural gradient w.r.t. $\text{vec}(\Sigma)$ is

$$\begin{aligned} \text{vec}(\hat{\mathbf{g}}^{[2]}) &:= \mathbf{F}_{\Sigma}^{-1} \text{vec}(\partial_{\Sigma} \mathcal{L}) \\ &= -2\partial_{\text{vec}(\Sigma^{-1})} [\text{vec}(\Sigma)] \text{vec}(\partial_{\Sigma} \mathcal{L}) \\ &= -2\partial_{\text{vec}(\Sigma^{-1})} [\text{vec}(\Sigma)] \partial_{\text{vec}(\Sigma)} \mathcal{L} \\ &= -2\partial_{\text{vec}(\Sigma^{-1})} \mathcal{L} \\ &= -2\text{vec}(\partial_{\Sigma^{-1}} \mathcal{L}) \end{aligned}$$

where we obtain the fourth step using the chain rule.

Therefore, we have $\hat{\mathbf{g}}^{[2]} = -2\partial_{\Sigma^{-1}} \mathcal{L}$. By matrix calculus, we have

$$\partial_{\Sigma^{-1}} \mathcal{L} = -\Sigma(\partial_{\Sigma} \mathcal{L})\Sigma$$

Finally, we have

$$\hat{\mathbf{g}}^{[2]} = 2\Sigma(\partial_{\Sigma} \mathcal{L})\Sigma$$

Now, we show that the additional term for μ is $\mathbf{0}$ under parameterization $\lambda = \{\mu, \Sigma\}$. Since λ is a BC parameterization, by Lemma 3, all entries of $\Gamma_{b_1 c_1}^{a_1}$ for μ are zeros. Therefore, the additional term for μ is $\mathbf{0}$.

We denote the Christoffel symbol of the second kind for $\text{vec}(\Sigma)$ as $\Gamma_{b_2 c_2}^{a_2}$. Now, we show that the additional term for Σ is $\frac{t \times t}{2} \hat{\mathbf{g}}^{[2]} \Sigma^{-1} \hat{\mathbf{g}}^{[2]}$. It is equivalent to show $\text{Mat}(\Gamma_{b_2 c_2}^{a_2} \hat{g}^{b_2} \hat{g}^{c_2}) = -\hat{\mathbf{g}}^{[2]} \Sigma^{-1} \hat{\mathbf{g}}^{[2]}$.

Recall that the natural gradient for $\mathbf{S} = \Sigma^{-1}$ is $\mathbf{G} = -2\partial_{\Sigma}\mathcal{L}$. Under parameterization $\bar{\lambda} = \{\mu, \mathbf{S}\}$, $\bar{\mathbf{R}}^{[2]}(t)$ has the following closed-form expression, which is used in the proof of Lemma 4.

$$\text{Mat}(\bar{\mathbf{R}}^{[2]}(t)) = \mathbf{U}\text{Exp}(t\mathbf{U}^{-1}\mathbf{G}\mathbf{U}^{-1})\mathbf{U}$$

where $\mathbf{U} = \mathbf{S}^{\frac{1}{2}}$ and $\text{Exp}(\mathbf{X}) := \mathbf{I} + \sum_{n=1}^{\infty} \frac{\mathbf{X}^n}{n!}$.

Note that $\Sigma = \mathbf{S}^{-1}$. Therefore, under parameterization $\lambda = \{\mu, \Sigma\}$, we have

$$\begin{aligned} \overbrace{\text{Mat}(\mathbf{R}^{[2]}(t))}^{\Sigma_{\text{new}}} &= \left[\overbrace{\text{Mat}(\bar{\mathbf{R}}^{[2]}(t))}^{\mathbf{S}_{\text{new}}} \right]^{-1} \\ &= (\mathbf{U}\text{Exp}(t\mathbf{U}^{-1}\mathbf{G}\mathbf{U}^{-1})\mathbf{U})^{-1} \\ &= \mathbf{U}^{-1}\text{Exp}(-t\mathbf{U}^{-1}\mathbf{G}\mathbf{U}^{-1})\mathbf{U}^{-1} \\ &= \Sigma^{1/2}\text{Exp}(-t\Sigma^{1/2}\mathbf{G}\Sigma^{1/2})\Sigma^{1/2} \\ &= \Sigma^{1/2}\text{Exp}(t\Sigma^{1/2}(2\partial_{\Sigma}\mathcal{L})\Sigma^{1/2})\Sigma^{1/2} \\ &= \Sigma^{1/2}\text{Exp}(t\Sigma^{-1/2}\underbrace{[2\Sigma(\partial_{\Sigma}\mathcal{L})\Sigma]}_{\hat{\mathbf{g}}^{[2]}}\Sigma^{-1/2})\Sigma^{1/2} \\ &= \Sigma^{1/2}\text{Exp}(t\Sigma^{-1/2}\hat{\mathbf{g}}^{[2]}\Sigma^{-1/2})\Sigma^{1/2}, \end{aligned}$$

where we use the identity $(\text{Exp}(t\mathbf{U}^{-1}\mathbf{G}\mathbf{U}^{-1}))^{-1} = \text{Exp}(-t\mathbf{U}^{-1}\mathbf{G}\mathbf{U}^{-1})$.

Note that a geodesic is invariant under parameterization. Alternatively, we can obtain the above equation by using the fact that $\mathbf{R}^{[2]}(t)$ is a geodesic of Gaussian distribution with a constant mean.

Using a similar proof as shown in Lemma 4, the additional term for Σ is

$$\text{Mat}(\Gamma_{b_2c_2}^{a_2}\hat{g}^{b_2}\hat{g}^{c_2}) = -\hat{\mathbf{g}}^{[2]}\Sigma^{-1}\hat{\mathbf{g}}^{[2]}$$

where $\Gamma_{b_2c_2}^{a_2}$ is the Christoffel symbol of the second kind for $\text{vec}(\Sigma)$ and \hat{g}^{a_2} denotes the a -th element of $\text{vec}(\hat{\mathbf{g}}^{[2]})$.

F. Example: Gamma Approximation

We consider the gamma distribution under the parameterization $\lambda = \{\lambda^{[1]}, \lambda^{[2]}\}$, where $\lambda^{[1]} = \alpha$ and $\lambda^{[2]} = \frac{\beta}{\alpha}$.

Since every block contains only a scalar, we use global indexes such as $\lambda^{(i)} = \lambda^{[i]}$, $\lambda^{(i)} = \lambda^{a_i}$ and $\Gamma_{i,ii} = \Gamma_{a_i,b_i c_i}$ for notation simplicity. The open-set constraint is $\Omega_1 = \mathbb{S}_{++}^1$ and $\Omega_2 = \mathbb{S}_{++}^1$. Under this parameterization, we can express the distribution as below.

$$q(z|\lambda) = z^{-1} \exp\left(\lambda^{(1)} \log z - z\lambda^{(1)}\lambda^{(2)} - A(\lambda)\right)$$

where $A(\lambda) = \log \text{Ga}(\lambda^{(1)}) - \lambda^{(1)}(\log \lambda^{(1)} + \log \lambda^{(2)})$ and $\text{Ga}(\cdot)$ is the gamma function.

Lemma 5 *The Fisher information matrix is diagonal under this parameterization. It implies that this parameterization is a BC parameterization.*

Proof: Notice that $\mathbb{E}_{q(z|\lambda)}[z] = \frac{1}{\lambda^{(2)}}$. The Fisher information matrix is diagonal as shown below.

$$\begin{aligned} \mathbf{F} &= -\mathbb{E}_{q(z|\lambda)}\left[\partial_{\lambda}^2 \log q(z|\lambda)\right] \\ &= -\mathbb{E}_{q(z|\lambda)}\left[\begin{array}{cc} -\partial_{\lambda^{(1)}}^2 A(\lambda) & (-z + \frac{1}{\lambda^{(2)}}) \\ (-z + \frac{1}{\lambda^{(2)}}) & -\partial_{\lambda^{(2)}}^2 A(\lambda) \end{array}\right] \\ &= \mathbb{E}_{q(z|\lambda)}\left[\begin{array}{cc} \partial_{\lambda^{(1)}}^2 A(\lambda) & 0 \\ 0 & \partial_{\lambda^{(2)}}^2 A(\lambda) \end{array}\right] \\ &= \begin{bmatrix} \partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}} & 0 \\ 0 & \frac{\lambda^{(1)}}{(\lambda^{(2)})^2} \end{bmatrix} \end{aligned}$$

where $\psi(\cdot)$ denotes the digamma function. □

Lemma 6 λ is a BCN parameterization.

Proof: By Lemma 5, we know that λ is a BC parameterization. Now, we show that $\lambda = \{\lambda^{(1)}, \lambda^{(2)}\}$ is a BCN parameterization. Clearly, each $\lambda^{(i)} \in \mathbb{S}_{++}^1$ has all degrees of freedom.

The gamma distribution which can be written as following exponential form:

$$q(z|\lambda^{(1)}, \lambda^{(2)}) = z^{-1} \exp \left(\lambda^{(1)} \log z - z\lambda^{(1)}\lambda^{(2)} - A(\lambda) \right)$$

Considering two blocks with $\lambda^{(1)}$ and $\lambda^{(2)}$ respectively, we can express this distribution in the following two ways where the first equation is for the $\lambda^{(1)}$ block while the second equation is for the $\lambda^{(2)}$ block:

$$\begin{aligned} q(z|\lambda^{(1)}, \lambda^{(2)}) &= \underbrace{z^{-1}}_{h_1(z, \lambda^{(2)})} \exp \left(\underbrace{\langle \log z - z\lambda^{(2)}, \lambda^{(1)} \rangle}_{\phi_1(z, \lambda^{(2)})} - A(\lambda) \right) \\ &= \underbrace{z^{-1} \exp(\lambda^{(1)} \log z)}_{h_2(z, \lambda^{(1)})} \exp \left(\underbrace{\langle -z\lambda^{(1)}, \lambda^{(2)} \rangle}_{\phi_2(z, \lambda^{(1)})} - A(\lambda) \right) \end{aligned}$$

Therefore, by the definition of BCN, we know that λ is a BCN parameterization. □

Using this BCN parameterization, the Christoffel symbols can be readily computed as below.

$$\begin{aligned} \Gamma_{1,11} &= \frac{1}{2} \partial_{\lambda^{(1)}}^3 A(\lambda) = \frac{1}{2} \left(\partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)}) + \frac{1}{(\lambda^{(1)})^2} \right), \quad \Gamma_{2,22} = \frac{1}{2} \partial_{\lambda^{(2)}}^3 A(\lambda) = -\frac{\lambda^{(1)}}{(\lambda^{(2)})^3} \\ \Gamma^1_{11} &= \frac{\Gamma_{1,11}}{F_{11}} = \frac{\partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)}) + \frac{1}{(\lambda^{(1)})^2}}{2 \left(\partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}} \right)}, \quad \Gamma^2_{22} = \frac{\Gamma_{2,22}}{F_{22}} = -\frac{1}{\lambda^{(2)}} \end{aligned}$$

F.1. Proof of Theorem 2

We first prove the following lemma.

Lemma 7 $\Gamma^1_{11} < -\frac{1}{\lambda^{(1)}}$ when $\lambda^{(1)} > 0$.

Proof: By Eq 1.4 at Batir (2005) and the last inequality at page 13 of Koumandos (2008), we have the following inequalities when $\lambda^{(1)} > 0$.

$$\partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}} > \frac{1}{2(\lambda^{(1)})^2} > 0 \quad \text{Batir (2005)} \quad (24)$$

$$\partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)}) < \frac{1}{(\lambda^{(1)})^2} - \frac{2\partial_{\lambda^{(1)}} \psi(\lambda^{(1)})}{\lambda^{(1)}} \quad \text{Koumandos (2008)} \quad (25)$$

By (25), we have

$$\partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)}) + \frac{1}{(\lambda^{(1)})^2} < \frac{2}{(\lambda^{(1)})^2} - \frac{2\partial_{\lambda^{(1)}} \psi(\lambda^{(1)})}{\lambda^{(1)}} = \frac{2}{\lambda^{(1)}} \left(\frac{1}{\lambda^{(1)}} - \partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) \right)$$

Since $\partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}} > 0$, we have

$$2\Gamma^1_{11} = \frac{\partial_{\lambda^{(1)}}^2 \psi(\lambda^{(1)}) + \frac{1}{(\lambda^{(1)})^2}}{\partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}}} < -\frac{2}{\lambda^{(1)}}$$

which shows $\Gamma^1_{11} < -\frac{1}{\lambda^{(1)}}$. □

Now, We give a proof for Theorem 2.

Proof: The proposed update for $\lambda^{(1)}$ with step-size t is given below.

$$\begin{aligned}
 \lambda^{(1)} &\leftarrow \lambda^{(1)} - t\hat{g}^{(1)} - \frac{t^2}{2} (\Gamma_{11}^1) (\hat{g}^{(1)})^2 \\
 &> \lambda^{(1)} - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{1}{\lambda^{(1)}} \right) (\hat{g}^{(1)})^2 \\
 &= \frac{1}{2\lambda^{(1)}} \left[2(\lambda^{(1)})^2 - 2t\hat{g}^{(1)}\lambda^{(1)} + (t\hat{g}^{(1)})^2 \right] \\
 &= \underbrace{\frac{1}{2\lambda^{(1)}}}_{>0} \left[\underbrace{(\lambda^{(1)})^2}_{>0} + \underbrace{(\lambda^{(1)} - t\hat{g}^{(1)})^2}_{\geq 0} \right]
 \end{aligned}$$

where in the second step we use the inequality $\Gamma_{11}^1 < -\frac{1}{\lambda^{(1)}}$ shown in Lemma 7 since the current/old $\lambda^{(1)} > 0$.

Similarly, we can show the update for $\lambda^{(2)}$ also satisfies the constraint.

$$\begin{aligned}
 \lambda^{(2)} &\leftarrow \lambda^{(2)} - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{1}{\lambda^{(2)}} \right) (\hat{g}^{(2)})^2 \\
 &= \frac{1}{2\lambda^{(2)}} \left[\underbrace{(\lambda^{(2)})^2}_{>0} + \underbrace{(\lambda^{(2)} - t\hat{g}^{(2)})^2}_{\geq 0} \right]
 \end{aligned}$$

It is obvious to see that the proposed update satisfies the underlying constraint. □

F.2. Natural Gradients

Recall that \hat{g} are the natural-gradients, which can be computed as shown below.

$$\hat{g}^{(1)} = \frac{\partial_{\lambda^{(1)}} \mathcal{L}}{\partial_{\lambda^{(1)}} \psi(\lambda^{(1)}) - \frac{1}{\lambda^{(1)}}}, \quad \hat{g}^{(2)} = \frac{(\lambda^{(2)})^2}{\lambda^{(1)}} \partial_{\lambda^{(2)}} \mathcal{L}$$

Recall that $\lambda^{(1)} = \alpha$ and $\lambda^{(2)} = \frac{\beta}{\alpha}$. Using the chain rule, we know that

$$\partial_{\lambda^{(1)}} \mathcal{L} = \partial_{\alpha} \mathcal{L} + \frac{\beta}{\alpha} \partial_{\beta} \mathcal{L}, \quad \partial_{\lambda^{(2)}} \mathcal{L} = \alpha \partial_{\beta} \mathcal{L}$$

$\partial_{\alpha} \mathcal{L}$ and $\partial_{\beta} \mathcal{L}$ can be computed by the implicit reparameterization trick (Salimans & Knowles, 2013; Figurnov et al., 2018).

G. Example: Exponential Approximation

In this case, there is only one block with a scalar. We use global indexes such as $\lambda^{(1)} = \lambda^{[1]}$ and $\Gamma_{1,11} = \Gamma_{a_1, b_1 c_1}$ for notation simplicity. We consider an exponential distribution under the natural parameterization $\lambda = \lambda^{(1)}$ with the open-set constraint $\Omega = \mathbb{S}_{++}^1$:

$$q(z|\lambda) = \exp \left(-\lambda^{(1)} z - A(\lambda) \right)$$

where $A(\lambda) = -\log \lambda^{(1)}$. The FIM is a scalar $F_{11} = \frac{1}{(\lambda^{(1)})^2}$. It is obvious that λ is a BCN parameterization. the Christoffel symbols can be readily computed as below.

$$\Gamma_{1,11} = \frac{1}{2} \partial_{\lambda^{(1)}}^3 A(\lambda) = -\frac{1}{(\lambda^{(1)})^3}, \quad \Gamma_{11}^1 = \frac{\Gamma_{1,11}}{F_{11}} = -\frac{1}{\lambda^{(1)}}$$

The proposed natural-gradient update with step-size t is

$$\lambda^{(1)} = \lambda^{(1)} - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{1}{\lambda^{(1)}} \right) \left(\hat{g}^{(1)} \right)^2$$

where $\hat{g}^{(1)}$ is the natural-gradient. Note that $\hat{g}^{(1)}$ is the natural-gradient, which can be computed as shown below.

$$\hat{g}^{(1)} = \left(\lambda^{(1)} \right)^2 \partial_{\lambda^{(1)}} \mathcal{L}.$$

where $\partial_{\lambda^{(1)}} \mathcal{L}$ can be computed by the implicit reparameterization trick as $\partial_{\lambda^{(1)}} \mathcal{L} \approx [\partial_{\lambda} z] [\partial_z b(z)]$, where $z \sim q(z|\lambda^{(1)})$ and $b(z) := \bar{\ell}(z) + \log q(z|\lambda^{(1)})$

Lemma 8 *The proposed update satisfies the underlying constraint.*

Proof: The proposed natural-gradient update with step-size t is given below.

$$\begin{aligned} \lambda^{(1)} &\leftarrow \lambda^{(1)} - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{1}{\lambda^{(1)}} \right) \left(\hat{g}^{(1)} \right)^2 \\ &= \frac{1}{2\lambda^{(1)}} \left[2 \left(\lambda^{(1)} \right)^2 - 2t\hat{g}^{(1)}\lambda^{(1)} + \left(t\hat{g}^{(1)} \right)^2 \right] \\ &= \frac{1}{2\lambda^{(1)}} \left[\left(\lambda^{(1)} \right)^2 + \left(\lambda^{(1)} - t\hat{g}^{(1)} \right)^2 \right] \end{aligned}$$

It is obvious to see that the proposed update satisfies the underlying constraint. □

G.1. Implicit reparameterization gradient

Now, we discuss how to compute the gradients w.r.t. λ using the implicit reparameterization trick. To use the implicit reparameterization trick, we have to compute the following term.

$$\partial_{\lambda} z = -\frac{\partial_{\lambda} Q(z|\lambda)}{q(z|\lambda)} = -\frac{\partial_{\lambda} (1 - \exp(-\lambda z))}{\lambda \exp(-\lambda z)} = -\frac{z \exp(-\lambda z)}{\lambda \exp(-\lambda z)} = -\frac{z}{\lambda}$$

where $Q(z|\lambda)$ is the C.D.F. of $q(z|\lambda)$.

H. Example: Inverse Gaussian Approximation

We consider the following distribution.

$$q(z|\alpha, \beta) = \sqrt{\frac{1}{2\pi z^3}} \exp \left(-\frac{z\alpha\beta^2}{2} - \frac{\alpha}{2z} + \frac{\log \alpha}{2} + \alpha\beta \right)$$

where $\{\frac{1}{\beta}, \alpha\}$ is a BC parameterization.

We consider a BCN parameterization $\boldsymbol{\lambda} = \{\lambda^{[1]}, \lambda^{[2]}\}$, where $\lambda^{[1]} = \beta^2$ and $\lambda^{[2]} = \alpha$ and the open-set constraint is $\Omega_1 = \mathbb{S}_{++}^1$ and $\Omega_2 = \mathbb{S}_{++}^1$. Since every block contains only a scalar, we use global indexes such as $\lambda^{(i)} = \lambda^{[i]}$ and $\Gamma_{i,ii} = \Gamma_{a_i, b_i c_i}$ for notation simplicity. Under this parameterization, we can re-express the distribution as

$$q(z|\boldsymbol{\lambda}) = \sqrt{\frac{1}{2\pi z^3}} \exp \left(-\frac{z}{2} \lambda^{(1)} \lambda^{(2)} - \frac{\lambda^{(2)}}{2z} - A(\boldsymbol{\lambda}) \right)$$

where $A(\boldsymbol{\lambda}) = -\frac{\log \lambda^{(2)}}{2} - \lambda^{(2)} \sqrt{\lambda^{(1)}}$.

Lemma 9 *The FIM is (block) diagonal under this parameterization.*

Proof: Notice that $\mathbb{E}_{q(z|\lambda)}[z] = \frac{1}{\sqrt{\lambda^{(1)}}}$. The FIM is (block) diagonal as shown below.

$$\begin{aligned} \mathbf{F} &= -\mathbb{E}_{q(z|\lambda)} [\partial_{\lambda}^2 \log q(z|\lambda)] \\ &= -\mathbb{E}_{q(z|\lambda)} \begin{bmatrix} -\partial_{\lambda^{(1)}}^2 A(\lambda) & \frac{1}{2} \left(-z + \frac{1}{\sqrt{\lambda^{(1)}}} \right) \\ \frac{1}{2} \left(-z + \frac{1}{\sqrt{\lambda^{(1)}}} \right) & -\partial_{\lambda^{(2)}}^2 A(\lambda) \end{bmatrix} \\ &= \mathbb{E}_{q(z|\lambda)} \begin{bmatrix} \partial_{\lambda^{(1)}}^2 A(\lambda) & 0 \\ 0 & \partial_{\lambda^{(2)}}^2 A(\lambda) \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{4} (\lambda^{(1)})^{-3/2} \lambda^{(2)} & 0 \\ 0 & \frac{1}{2} (\lambda^{(2)})^{-2} \end{bmatrix} \end{aligned}$$

□

It is easy to show that λ is a BCN parameterization since λ satisfies Assumption 1 to 3.

Due to the BCN parameterization, the Christoffel symbols can be readily computed as below.

$$\begin{aligned} \Gamma_{1,11} &= \frac{1}{2} \partial_{\lambda^{(1)}}^3 A(\lambda) = -\frac{3}{16} (\lambda^{(1)})^{-5/2} \lambda^{(2)}, \quad \Gamma_{2,22} = \frac{1}{2} \partial_{\lambda^{(2)}}^3 A(\lambda) = -\frac{1}{2} (\lambda^{(2)})^{-3} \\ \Gamma^1_{11} &= \frac{\Gamma_{1,11}}{F_{11}} = -\frac{3}{4\lambda^{(1)}}, \quad \Gamma^2_{22} = \frac{\Gamma_{2,22}}{F_{22}} = -\frac{1}{\lambda^{(2)}} \end{aligned}$$

The proposed natural-gradient update with step-size t is

$$\begin{aligned} \lambda^{(1)} &\leftarrow \lambda^{(1)} - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{3}{4\lambda^{(1)}} \right) (\hat{g}^{(1)})^2 \\ \lambda^{(2)} &\leftarrow \lambda^{(2)} - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{1}{\lambda^{(2)}} \right) (\hat{g}^{(2)})^2 \end{aligned}$$

Lemma 10 *The update above satisfies the underlying constraint.*

Proof: The proposed natural-gradient update with step-size t is given below.

$$\begin{aligned} \lambda^{(1)} &\leftarrow \lambda^{(1)} - t\hat{g}^{(1)} + \frac{t^2}{2} \left(\frac{3}{4\lambda^{(1)}} \right) (\hat{g}^{(1)})^2 \\ &= \frac{1}{4\lambda^{(1)}} \left[4(\lambda^{(1)})^2 - 4t\hat{g}^{(1)}\lambda^{(1)} + \frac{3}{2} (t\hat{g}^{(1)})^2 \right] \\ &= \frac{1}{4\lambda^{(1)}} \left[\underbrace{(2\lambda^{(1)} - t\hat{g}^{(1)})^2}_{\text{Term I}} + \underbrace{\frac{1}{2} (t\hat{g}^{(1)})^2}_{\text{Term II}} \right] \\ \lambda^{(2)} &\leftarrow \lambda^{(2)} - t\hat{g}^{(2)} + \frac{t^2}{2} \left(\frac{1}{\lambda^{(2)}} \right) (\hat{g}^{(2)})^2 \\ &= \frac{1}{2\lambda^{(2)}} \left[2(\lambda^{(2)})^2 - 2t\hat{g}^{(2)}\lambda^{(2)} + (t\hat{g}^{(2)})^2 \right] \\ &= \frac{1}{2\lambda^{(2)}} \left[(\lambda^{(2)})^2 + (\lambda^{(2)} - t\hat{g}^{(2)})^2 \right] \end{aligned}$$

Note that Term I and Term II cannot be both zero at the same time when $\lambda^{(1)} > 0$. A similar argument can be made for the update about $\lambda^{(2)}$. Therefore, the proposed update satisfies the underlying constraint. □

Recall that \hat{g} are the natural-gradients, which can be computed as shown below.

$$\hat{g}^{(1)} = \frac{4}{\lambda^{(2)}} (\lambda^{(1)})^{3/2} \partial_{\lambda^{(1)}} \mathcal{L}, \quad \hat{g}^{(2)} = 2 (\lambda^{(2)})^2 \partial_{\lambda^{(2)}} \mathcal{L}$$

Using the chain rule, we know that

$$\partial_{\lambda^{(1)}} \mathcal{L} = \frac{1}{2\beta} \partial_{\beta} \mathcal{L}, \quad \partial_{\lambda^{(2)}} \mathcal{L} = \partial_{\alpha} \mathcal{L}$$

$\partial_{\alpha} \mathcal{L}$ and $\partial_{\beta} \mathcal{L}$ can be computed by the implicit reparameterization trick (Salimans & Knowles, 2013; Figurnov et al., 2018) as $\partial_{\eta} \mathcal{L} \approx [\partial_{\eta} z] [\nabla_z b(z)]$, where $\eta = \{\alpha, \beta\}$, $z \sim q(z|\alpha, \beta)$ and $b(z) := \bar{\ell}(z) + \log q(z|\alpha, \beta)$

H.1. Implicit reparameterization gradient

Now, we discuss how to compute the gradients w.r.t. α and β using the implicit reparameterization trick. To use the implicit reparameterization trick, we have to compute the following term.

$$\begin{aligned} \partial_{\eta} z &= -\frac{\partial_{\eta} Q(z|\eta)}{q(z|\eta)} \\ &= -\frac{\partial_{\eta} [\Phi(\sqrt{\frac{\alpha}{z}}(z\beta - 1)) + \exp(2\alpha\beta)\Phi(-\sqrt{\frac{\alpha}{z}}(z\beta + 1))]}{\sqrt{\frac{1}{2\pi z^3}} \exp\left(-\frac{z\alpha\beta^2}{2} - \frac{\alpha}{2z} + \frac{\log \alpha}{2} + \alpha\beta\right)} \end{aligned}$$

where $\eta = \{\alpha, \beta\}$, $Q(z|\eta)$ is the C.D.F. of the inverse Gaussian distribution, and $\Phi(x) = \int_{-\infty}^x \mathcal{N}(t|0, 1)dt$ is the C.D.F. of the standard Gaussian distribution. We use the following fact to simplify the above expression.

$$\delta(z, \alpha, \beta) := \frac{\exp(2\alpha\beta)\Phi(-\sqrt{\frac{\alpha}{z}}(z\beta + 1))}{\mathcal{N}(\sqrt{\frac{\alpha}{z}}(z\beta - 1)|0, 1)} = \frac{\Phi(-\sqrt{\frac{\alpha}{z}}(z\beta + 1))}{\mathcal{N}(-\sqrt{\frac{\alpha}{z}}(z\beta + 1)|0, 1)}$$

where $\delta(z, \alpha, \beta)$ is known as the Mills ratio of Gaussian distribution. Using this fact, we can get the simplified expressions as follows.

$$\begin{aligned} \partial_{\alpha} z &= \frac{z}{\alpha} - 2\beta z^{3/2} \alpha^{-1/2} \delta(z, \alpha, \beta) \\ \partial_{\beta} z &= -2z^{3/2} \alpha^{1/2} \delta(z, \alpha, \beta) \end{aligned}$$

where we compute $\log(\delta(z, \alpha, \beta))$ for numerical stability since the logarithm of Gaussian cumulative distribution function can be computed by using existing libraries, such as the `scipy.special.log_ndtr()` function.

In fact, we have closed-form expressions of gradients of the entropy term as shown below.

$$\begin{aligned} \mathbb{E}_{q(z|\eta)} [-\log q(z|\eta)] &= \frac{1}{2} [-\log \alpha - 3(\log \beta + \exp(2\alpha\beta)E_1(2\alpha\beta)) + 1 + \log(2\pi)] \\ \partial_{\alpha} \mathbb{E}_{q(z|\eta)} [-\log q(z|\eta)] &= \frac{1}{\alpha} - 3\beta \exp(2\alpha\beta)E_1(2\alpha\beta) \\ \partial_{\beta} \mathbb{E}_{q(z|\eta)} [-\log q(z|\eta)] &= -3\alpha \exp(2\alpha\beta)E_1(2\alpha\beta) \end{aligned}$$

where $E_1(x) := \int_x^{\infty} \frac{e^{-t}}{t} dt$ is the exponential integral. It is not numerical stable to compute the product $\exp(x)E_1(x)$ when $x > 100$. In this case, we can use the asymptotic expansion (see Eq 3 at Tseng & Lee (1998)) for the exponential integral to approximate the product as shown below.

$$\exp(x)E_1(x) \approx \frac{1}{x} \left[1 + \sum_{n=1}^N \frac{(-1)^n n!}{x^n} \right] \quad \text{when } x > 100,$$

where N is an integer such as $N \leq x < N + 1$.

I. Mixture of Exponential Family Distributions

Let's consider the following mixture of exponential family distributions $q(\mathbf{z}) = \int q(\mathbf{z}, \mathbf{w})d\mathbf{w}$. The joint distribution $q(\mathbf{z}, \mathbf{w}|\boldsymbol{\lambda}) = q(\mathbf{w}|\boldsymbol{\lambda}_w)q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$ is known as the conditional exponential family (CEF) defined by Lin et al. (2019a).

$$\begin{aligned} q(\mathbf{w}|\boldsymbol{\lambda}_w) &:= h_w(\mathbf{w}) \exp[\langle \boldsymbol{\phi}_w(\mathbf{w}), \boldsymbol{\lambda}_w \rangle - A_w(\boldsymbol{\lambda}_w)] \\ q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z) &:= h_z(\mathbf{w}, \mathbf{z}) \exp[\langle \boldsymbol{\phi}_z(\mathbf{w}, \mathbf{z}), \boldsymbol{\lambda}_z \rangle - A_z(\boldsymbol{\lambda}_z, \mathbf{w})] \end{aligned}$$

where $\lambda = \{\lambda_z, \lambda_w\}$.

We will use the joint Fisher information matrix(FIM) suggested by Lin et al. (2019a) as the metric \mathbf{F} to derive our improved learning rule for mixture approximations.

I.1. The Joint Fisher Information Matrix and the Christoffel Symbol

Lin et al. (2019a) propose to use the FIM of the joint distribution $q(\mathbf{w}, \mathbf{z}|\lambda)$, where they refer this FIM as the joint FIM. The joint FIM and the corresponding Christoffel symbol of the first kind are defined as follows.

$$F_{ab} := -\mathbb{E}_{q(\mathbf{w}, \mathbf{z}|\lambda)} [\partial_a \partial_b \log q(\mathbf{w}, \mathbf{z}|\lambda)]$$

$$\Gamma_{d,ab} := \frac{1}{2} [\partial_a F_{bd} + \partial_b F_{ad} - \partial_d F_{ab}]$$

where we denote $\partial_a = \partial_{\lambda^a}$ for notation simplicity.

Like the exponential family cases as shown in Eq. (17), the Christoffel symbol of the first kind can be computed as

$$\begin{aligned} \Gamma_{d,ab} = \frac{1}{2} & \left[\mathbb{E}_{q(\mathbf{w}, \mathbf{z}|\lambda)} [\partial_a \partial_b \log q(\mathbf{w}, \mathbf{z}|\lambda) \partial_d \log q(\mathbf{w}, \mathbf{z}|\lambda)] - \mathbb{E}_{q(\mathbf{w}, \mathbf{z}|\lambda)} [\partial_b \partial_d \log q(\mathbf{w}, \mathbf{z}|\lambda) \partial_a \log q(\mathbf{w}, \mathbf{z}|\lambda)] \right. \\ & \left. - \mathbb{E}_{q(\mathbf{w}, \mathbf{z}|\lambda)} [\partial_a \partial_d \log q(\mathbf{w}, \mathbf{z}|\lambda) \partial_b \log q(\mathbf{w}, \mathbf{z}|\lambda)] - \mathbb{E}_{q(\mathbf{w}, \mathbf{z}|\lambda)} [\partial_a \partial_b \partial_d \log q(\mathbf{w}, \mathbf{z}|\lambda)] \right] \end{aligned} \quad (26)$$

I.2. The BCN Parameterization

Now, we show that how to simplify the computation of the Christoffel symbol by extending the BCN parameterization for this kind of mixtures.

To this end, we first assume that λ can be partitioned with $(m+n)$ blocks to satisfy Assumption 1 in the main text.

$$\lambda = \underbrace{\{\lambda_z^{[1]}, \dots, \lambda_z^{[m]}\}}_{\lambda_z}, \underbrace{\{\lambda_w^{[m+1]}, \dots, \lambda_w^{[m+n]}\}}_{\lambda_w}$$

Then, we extend the definition of BC parameterization to the conditional exponential family, which is similar to Assumption 2 in the main text and a concrete example of Definition 1 in Appendix B.1.

Assumption 2 [Block Coordinate Parameterization] : A parameterization satisfied Assumption 1 is block coordinate (BC) if the *joint FIM* under this parameterization is block-diagonal according to the block structure of the parameterization.

Lin et al. (2019a) show that, for any parameterization $\lambda = \{\lambda_z, \lambda_w\}$, the joint FIM has the following two blocks: \mathbf{F}_z for block λ_z and \mathbf{F}_w for block λ_w as given below.

$$\mathbf{F} = \begin{bmatrix} \mathbf{F}_z & \mathbf{0} \\ \mathbf{0} & \mathbf{F}_w \end{bmatrix}$$

Assumption 2 implies that \mathbf{F}_w and \mathbf{F}_z are both block-diagonal according to the block structure of λ_w and λ_z , respectively. The block diagonal structure is given below if $\lambda = \{\lambda_z^{[1]}, \dots, \lambda_z^{[m]}, \lambda_w^{[m+1]}, \dots, \lambda_w^{[m+n]}\}$ is a BC parameterization.

$$\mathbf{F} = \begin{bmatrix} \overbrace{\begin{bmatrix} \mathbf{F}_z^{[1]} & \dots & \mathbf{0} \\ \vdots & \ddots & \vdots \\ \mathbf{0} & \dots & \mathbf{F}_z^{[m]} \end{bmatrix}}^{\mathbf{F}_z} & \mathbf{0} \\ \mathbf{0} & \underbrace{\begin{bmatrix} \mathbf{F}_w^{[m+1]} & \dots & \mathbf{0} \\ \vdots & \ddots & \vdots \\ \mathbf{0} & \dots & \mathbf{F}_w^{[m+n]} \end{bmatrix}}_{\mathbf{F}_w} \end{bmatrix}$$

Assumption 3 [Block Natural Parameterization for the Conditional Exponential-Family] : For a conditional exponential-family distribution $q(\mathbf{w}, \mathbf{z}|\boldsymbol{\lambda}) = q(\mathbf{w}|\boldsymbol{\lambda}_w)q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$, a parameterization $\boldsymbol{\lambda} = \{\boldsymbol{\lambda}_z, \boldsymbol{\lambda}_w\}$ has the following properties.

- $\boldsymbol{\lambda}_w$ is a BCN parameterization of the exponential family distribution $q(\mathbf{w}|\boldsymbol{\lambda}_w)$ as defined in the main text.
- $\boldsymbol{\lambda}_z$ is a parameterization of $q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$, where there exist function ϕ_{z_i} and h_{z_i} for each block $\boldsymbol{\lambda}_z^{[i]}$ such that conditioning on \mathbf{w} , $q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$ can be re-expressed as a minimal conditional exponential family distribution (see Lin et al. (2019a) for the definition of the minimality) given that the rest of blocks $\boldsymbol{\lambda}_z^{[-i]}$ are known.

$$q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z) \equiv h_{z_i}(\mathbf{w}, \mathbf{z}, \boldsymbol{\lambda}_z^{[-i]}) \exp [\langle \phi_{z_i}(\mathbf{w}, \mathbf{z}, \boldsymbol{\lambda}_z^{[-i]}), \boldsymbol{\lambda}_z^{[i]} \rangle - A_z(\boldsymbol{\lambda}_z, \mathbf{w})]$$

We say $\boldsymbol{\lambda} = \{\boldsymbol{\lambda}_z, \boldsymbol{\lambda}_w\}$ is a BCN parameterization for the mixture if it satisfies Assumption 1 to 3.

Many mixture approximations studied in Lin et al. (2019a) have a BCN parameterization. For concrete examples, see Appendix J and K.

I.3. Our Learning Rule for Mixture Approximations

Now, we are ready to discuss the learning rule for mixture approximations. Under a BC parameterization $\boldsymbol{\lambda} = \{\boldsymbol{\lambda}_z, \boldsymbol{\lambda}_w\}$, our learning rule remains the same as shown below.

$$\lambda^{c_i} \leftarrow \lambda^{c_i} - t \hat{g}^{c_i} - \frac{t^2}{2} \Gamma_{a_i b_i}^{c_i} \hat{g}^{a_i} \hat{g}^{b_i}$$

where block i can be either a block of $\boldsymbol{\lambda}_w$ or $\boldsymbol{\lambda}_z$.

First, note that the sub-block matrix \mathbf{F}_w of the joint FIM is indeed the FIM of $q(\mathbf{w}|\boldsymbol{\lambda}_w)$. Furthermore, $q(\mathbf{w}|\boldsymbol{\lambda}_w)$ is an exponential family distribution. If $\boldsymbol{\lambda} = \{\boldsymbol{\lambda}_z, \boldsymbol{\lambda}_w\}$ is a BCN parameterization, it is easy to see that the computation of the Christoffel symbol for $\boldsymbol{\lambda}_w$ is exactly the same as the exponential family cases as discussed in Appendix D.

Furthermore, we can simplify the Christoffel symbol for $\boldsymbol{\lambda}_z$ due to the following Theorem.

Theorem 4 If $\boldsymbol{\lambda}$ is a BCN parameterization of a conditional exponential family (CEF) with the joint FIM, natural gradient and the Christoffel symbol of the first kind for block $\boldsymbol{\lambda}_z^{[i]}$ can be simplified as

$$\hat{g}^{a_i} = \partial_{m_{z_{a_i}}} \mathcal{L} ; \Gamma_{d_i, a_i b_i} = \frac{1}{2} \mathbb{E}_{q(\mathbf{w}|\boldsymbol{\lambda}_w)} \left[\partial_{\lambda_z^{a_i}} \partial_{\lambda_z^{b_i}} \partial_{\lambda_z^{d_i}} A_z(\boldsymbol{\lambda}_z, \mathbf{w}) \right]$$

where $\lambda_z^{a_i}$ is the a -th element of $\boldsymbol{\lambda}_z^{[i]}$; $m_{z_{a_i}}$ denotes the a -th element of the block coordinate expectation parameter $\mathbf{m}_{z_{[i]}} = \mathbb{E}_{q(\mathbf{w}, \mathbf{z}|\boldsymbol{\lambda})} [\phi_{z_i}(\mathbf{w}, \mathbf{z}, \boldsymbol{\lambda}_z^{[-i]})] = \mathbb{E}_{q(\mathbf{w}|\boldsymbol{\lambda}_w)} [\partial_{\lambda_z^{[i]}} A_z(\boldsymbol{\lambda}_z, \mathbf{w})]$.

I.4. Proof of Theorem 4

Proof: We assume $\boldsymbol{\lambda}_z = \{\boldsymbol{\lambda}_z^{[1]}, \dots, \boldsymbol{\lambda}_z^{[m]}\}$ is partitioned with m blocks.

Since $\boldsymbol{\lambda}$ is a BCN parameterization, conditioning on \mathbf{w} and given $\boldsymbol{\lambda}_z^{[-i]}$ and $\boldsymbol{\lambda}_w$ are known, we can re-express $q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$ as

$$q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z) = h_{z_i}(\mathbf{z}, \mathbf{w}, \boldsymbol{\lambda}_z^{[-i]}) \exp [\langle \phi_{z_i}(\mathbf{z}, \mathbf{w}, \boldsymbol{\lambda}_z^{[-i]}), \boldsymbol{\lambda}_z^{[i]} \rangle - A_z(\boldsymbol{\lambda}_z, \mathbf{w})]$$

where $q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$ is also a one-parameter EF distribution conditioning on $\boldsymbol{\lambda}_z^{[-i]}$ and \mathbf{w} . Similarly, we have the following results.

$$\begin{aligned} \partial_{a_i} \partial_{b_i} \log q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z) &= -\partial_{a_i} \partial_{b_i} A_z(\boldsymbol{\lambda}_z, \mathbf{w}) \\ \mathbb{E}_{q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)} [\partial_{a_i} \log q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)] &= 0 \end{aligned}$$

where $\partial_{a_i} = \partial_{\lambda_z^{a_i}}$ is for notation simplicity. Using the above identities, we have

$$\begin{aligned} \mathbb{E}_{q(z,w|\lambda)} [\partial_{a_i} \partial_{b_i} \log q(\mathbf{z}, \mathbf{w}|\lambda) \partial_{d_i} \log q(\mathbf{z}, \mathbf{w}|\lambda)] &= \mathbb{E}_{q(z,w|\lambda)} [\partial_{a_i} \partial_{b_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z) \partial_{d_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z)] \\ &= \mathbb{E}_{q(w|\lambda_w)} [\mathbb{E}_{q(z|w, \lambda_z)} [\partial_{a_i} \partial_{b_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z) \partial_{d_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z)]] \\ &= -\mathbb{E}_{q(w|\lambda_w)} \left[\partial_{a_i} \partial_{b_i} A_z(\lambda_z, \mathbf{w}) \underbrace{\mathbb{E}_{q(z|w, \lambda_z)} [\partial_{d_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z)]}_0 \right] \\ &= 0 \end{aligned}$$

Therefore, by Eq. (26), we can simplify the Christoffel symbol for $\lambda_z^{[i]}$ as follows.

$$\begin{aligned} \Gamma_{d_i, a_i b_i} &= -\frac{1}{2} \mathbb{E}_{q(z,w|\lambda)} [\partial_{a_i} \partial_{b_i} \partial_{d_i} \log q(\mathbf{z}, \mathbf{w}|\lambda)] \\ &= -\frac{1}{2} \mathbb{E}_{q(w|\lambda_w)} [\partial_{a_i} \partial_{b_i} \partial_{d_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z)] \\ &= \frac{1}{2} \mathbb{E}_{q(w|\lambda_w)} [\partial_{a_i} \partial_{b_i} \partial_{d_i} A_z(\lambda_z, \mathbf{w})] \end{aligned}$$

where we use d_i to denote the d -th entry of block $\lambda_z^{[i]}$.

Likewise, let $\mathbf{m}_{z[i]} = \mathbb{E}_{q(z,w|\lambda)} [\phi_{z[i]}(\mathbf{z}, \mathbf{w}, \lambda_z^{[-i]})]$ denote the block coordinate expectation parameter. We have

$$0 = \mathbb{E}_{q(w|\lambda_w)} \left[\underbrace{\mathbb{E}_{q(z|w, \lambda_z)} [\partial_{a_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z)]}_0 \right] = m_{z_{a_i}} - \mathbb{E}_{q(w|\lambda_w)} [\partial_{a_i} A_z(\lambda_z, \mathbf{w})]$$

where $m_{z_{a_i}}$ denotes the a -th element of $\mathbf{m}_{z[i]}$.

Therefore, we know that $m_{z_{a_i}} = \mathbb{E}_{q(w|\lambda_w)} [\partial_{a_i} A_z(\lambda_z, \mathbf{w})]$.

Recall that the sub-block of the joint FIM for $\lambda_z^{[i]}$ denoted by $\mathbf{F}_z^{[i]}$ can be computed as

$$\begin{aligned} F_{a_i b_i} &= -\mathbb{E}_{q(z,w|\lambda)} [\partial_{b_i} \partial_{a_i} \log q(\mathbf{z}, \mathbf{w}|\lambda)] \\ &= -\mathbb{E}_{q(z,w|\lambda)} [\partial_{b_i} \partial_{a_i} \log q(\mathbf{z}|\mathbf{w}, \lambda_z)] \\ &= -\mathbb{E}_{q(z,w|\lambda)} [-\partial_{b_i} \partial_{a_i} A_z(\lambda_z, \mathbf{w})] \\ &= \mathbb{E}_{q(w|\lambda_w)} [\partial_{b_i} \partial_{a_i} A_z(\lambda_z, \mathbf{w})] \\ &= \partial_{b_i} \mathbb{E}_{q(w|\lambda_w)} [\partial_{a_i} A_z(\lambda_z, \mathbf{w})] \\ &= \partial_{b_i} m_{z_{a_i}} \end{aligned}$$

where we use the fact that λ_w does not depend on $\lambda_z^{b_i} \in \lambda_z$ and $\partial_{b_i} = \partial_{\lambda_z^{b_i}}$ to move from the fourth step to the fifth step.

Recall that when λ is a BC parameterization, the joint FIM \mathbf{F} is block-diagonal as shown below.

$$\mathbf{F} = \begin{bmatrix} \overbrace{\begin{bmatrix} \mathbf{F}_z^{[1]} & \cdots & \mathbf{0} \\ \vdots & \ddots & \vdots \\ \mathbf{0} & \cdots & \mathbf{F}_z^{[m]} \end{bmatrix}}^{\mathbf{F}_z} & & \\ & \mathbf{0} & \\ & & \underbrace{\begin{bmatrix} \mathbf{F}_w^{[m+1]} & \cdots & \mathbf{0} \\ \vdots & \ddots & \vdots \\ \mathbf{0} & \cdots & \mathbf{F}_w^{[m+n]} \end{bmatrix}}_{\mathbf{F}_w} \end{bmatrix}$$

If $\mathbf{F}_z^{[i]}$ is positive definite everywhere, we have

$$F_{a_i b_i} = \partial_{m_{z_{a_i}}} \lambda_z^{b_i}$$

The above assumption is true if given that $\lambda_z^{[-i]}$ and λ_w are known, $q(\mathbf{w}, \mathbf{z}|\lambda)$ is a one-parameter minimal CEF distribution (See Theorem 2 of Lin et al. (2019a)).

The above result implies that we can compute natural gradients as follows.

$$\hat{g}^{a_i} = F^{a_i b_i} g_{b_i} = \left[\partial_{m_{z_{a_i}}} \lambda_z^{b_i} \right] \left[\partial_{\lambda_z^{b_i}} \mathcal{L} \right] = \partial_{m_{z_{a_i}}} \mathcal{L}$$

where $g_{b_i} = \partial_{\lambda_z^{b_i}} \mathcal{L}$.

□

If we can interchange the differentiations and the integration, we can show, by Theorem 4, we have $\Gamma_{a_i b_i}^{c_i} = \frac{1}{2} \partial_{m_{z_{c_i}}} \partial_{\lambda_z^{a_i}} \partial_{\lambda_z^{b_i}} \mathbb{E}_{q(w|\lambda_w)} [A_z(\lambda_z, \mathbf{w})]$ since $A_z(\lambda_z, \mathbf{w})$ is C^3 -smooth w.r.t. $\lambda_z^{[i]}$.

J. Example: Finite Mixture of Gaussians Approximation

We consider a K-mixture of Gaussians under this parameterization $\lambda = \{\{\mu_c, \mathbf{S}_c\}_{c=1}^K, \lambda_w\}$

$$q(\mathbf{z}|\pi, \{\mu_c, \mathbf{S}_c\}_{c=1}^K) = \sum_{c=1}^K \pi_c \mathcal{N}(\mathbf{z}|\mu_c, \mathbf{S}_c)$$

where π_c is the mixing weight so that $\sum_{c=1}^K \pi_c = 1$, $\mathbf{S}_c = \Sigma_c^{-1}$, $\lambda_w = \{\log(\pi_c/\pi_K)\}_{c=1}^{K-1}$ and $\pi_K = 1 - \sum_{c=1}^{K-1} \pi_c$. The constraints are $\lambda_w \in \mathbb{R}^{K-1}$, $\mu_c \in \mathbb{R}^d$, and $\mathbf{S}_c \in \mathbb{S}_{++}^{d \times d}$.

Under this parameterization, the joint distribution can be expressed as below.

$$\begin{aligned} q(\mathbf{z}, w|\lambda) &= q(w|\lambda_w) q(\mathbf{z}|w, \{\mu_c, \mathbf{S}_c\}_{c=1}^K) \\ q(w|\lambda_w) &= \exp\left(\sum_{c=1}^{K-1} \mathbb{I}(w=c) \lambda_{w_c} - A_w(\lambda_w)\right) \\ q(\mathbf{z}|w, \{\mu_c, \mathbf{S}_c\}_{c=1}^K) &= \exp\left(\sum_{c=1}^K \mathbb{I}(w=c) \left[-\frac{1}{2} \mathbf{z}^T \mathbf{S}_c \mathbf{z} + \mathbf{z}^T \mathbf{S}_c \mu_c\right] - A_z(\{\mu_c, \mathbf{S}_c\}_{c=1}^K, w)\right) \end{aligned}$$

where $B(\mu_c, \mathbf{S}_c) = \frac{1}{2} [\mu_c^T \mathbf{S}_c \mu_c - \log |\mathbf{S}_c|/(2\pi)]$, $A_z(\{\mu_c, \mathbf{S}_c\}_{c=1}^K, w) = \sum_{c=1}^K \mathbb{I}(w=c) B(\mu_c, \mathbf{S}_c)$, $\lambda_{w_c} = \log(\frac{\pi_c}{\pi_K})$, $A_w(\lambda_w) = \log(1 + \sum_{c=1}^{K-1} \exp(\lambda_{w_c}))$.

Lemma 11 *The joint FIM is block diagonal under this parameterization.*

$$\mathbf{F} = \begin{bmatrix} \begin{bmatrix} \mathbf{F}_{\mu_1} & \mathbf{0} \\ \mathbf{0} & \mathbf{F}_{S_1} \end{bmatrix} & \cdots & \mathbf{0} & \mathbf{0} \\ \vdots & \ddots & \vdots & \vdots \\ \mathbf{0} & \cdots & \begin{bmatrix} \mathbf{F}_{\mu_K} & \mathbf{0} \\ \mathbf{0} & \mathbf{F}_{S_K} \end{bmatrix} & \mathbf{0} \\ \mathbf{0} & \cdots & \mathbf{0} & \mathbf{F}_w \end{bmatrix}$$

Therefore, this parameterization is a BC parameterization.

Proof: We will prove this lemma by showing that all cross terms are zeros.

Case 1: First, we will show that cross terms (shown in red) between λ_w and $\lambda_z := \{\mu_c, \mathbf{S}_c\}_{c=1}^K$ are zeros.

Let's denote λ_w^i be an element of λ_w and λ_z^j be an element of λ_z . By the definition, each cross term in this case is defined as belows.

$$-\mathbb{E}_{q(\mathbf{z}, w|\lambda)} \left[\partial_{\lambda_w^i} \partial_{\lambda_z^j} \log q(\mathbf{z}, w|\lambda) \right] = -\mathbb{E}_{q(\mathbf{z}, w|\lambda)} \left[\partial_{\lambda_w^i} \partial_{\lambda_z^j} (\log q(w|\lambda_w) + \log q(\mathbf{z}|w, \lambda_z)) \right] = 0$$

Case 2: Next, we will show that cross terms between (shown in blue) any two Gaussian components are zeros.

Let's denote λ_a^i be an element of $\{\boldsymbol{\mu}_a, \mathbf{S}_a\}$ and λ_b^j be an element of $\{\boldsymbol{\mu}_b, \mathbf{S}_b\}$, where $a \neq b$.

By the definition, each cross term in this case is defined as belows.

$$\begin{aligned}
 & -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\partial_{\lambda_a^i} \partial_{\lambda_b^j} \log q(\mathbf{z}, w|\boldsymbol{\lambda}) \right] \\
 &= -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\partial_{\lambda_a^i} \partial_{\lambda_b^j} \left(\log q(\mathbf{z}|w, \{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K) \right) \right] \\
 &= -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\mathbb{I}(w=b) \partial_{\lambda_a^i} \underbrace{\left(\partial_{\lambda_b^j} \left[-\frac{1}{2} \mathbf{z}^T \mathbf{S}_b \mathbf{z} + \mathbf{z}^T \mathbf{S}_b \boldsymbol{\mu}_b - B(\boldsymbol{\mu}_b, \mathbf{S}_b) \right] \right)}_{u(\mathbf{z}, \boldsymbol{\mu}_b, \mathbf{S}_b)} \right] = 0
 \end{aligned}$$

It is obvious that the above expression is 0 since $\partial_{\lambda_a^i} u(\mathbf{z}, \boldsymbol{\mu}_b, \mathbf{S}_b) = 0$ when $a \neq b$.

Case 3: Finally, we will show that for each component a , cross terms (shown in green) between $\boldsymbol{\mu}_a$ and \mathbf{S}_a are zeros.

Let's denote μ_a^i be the i -th element of $\boldsymbol{\mu}_a$ and S_a^{jk} be the element of \mathbf{S}_a at position (j, k) . Furthermore, \mathbf{e}_i denotes an one-hot vector where all entries are zeros except the i -th entry with value 1, and \mathbf{I}_{jk} denotes an one-hot matrix where all entries are zeros except the entry at position (j, k) with value 1. By the definition, the cross term is defined as belows.

$$\begin{aligned}
 & -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\partial_{\mu_a^i} \partial_{S_a^{jk}} \log q(\mathbf{z}, w|\boldsymbol{\lambda}) \right] \\
 &= -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\partial_{\mu_a^i} \partial_{S_a^{jk}} \left(\log q(\mathbf{z}|w, \{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K) \right) \right] \\
 &= -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\partial_{\mu_a^i} \partial_{S_a^{jk}} \left(\mathbb{I}(w=a) \left[-\frac{1}{2} \mathbf{z}^T \mathbf{S}_a \mathbf{z} + \mathbf{z}^T \mathbf{S}_a \boldsymbol{\mu}_a - B(\boldsymbol{\mu}_a, \mathbf{S}_a) \right] \right) \right] \\
 &= -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\mathbb{I}(w=a) \left[\mathbf{e}_i^T \mathbf{I}_{jk} \mathbf{z} - \mathbf{e}_i^T \mathbf{I}_{jk} \boldsymbol{\mu}_a \right] \right] \\
 &= -\mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\mathbb{I}(w=a) \mathbf{e}_i^T \mathbf{I}_{jk} \mathbf{z} \right] + \mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} \left[\mathbb{I}(w=a) \mathbf{e}_i^T \mathbf{I}_{jk} \boldsymbol{\mu}_a \right] \\
 &= -\pi_a \mathbf{e}_i^T \mathbf{I}_{jk} \boldsymbol{\mu}_a + \pi_a \mathbf{e}_i^T \mathbf{I}_{jk} \boldsymbol{\mu}_a = 0
 \end{aligned}$$

where we use the following fact in the last step.

$$\begin{aligned}
 \mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} [\mathbb{I}(w=a) \mathbf{z}] &= \pi_a \boldsymbol{\mu}_a \\
 \mathbb{E}_{q(\mathbf{z}, w|\boldsymbol{\lambda})} [\mathbb{I}(w=a)] &= \pi_a
 \end{aligned}$$

□

Lemma 12 The parameterization $\boldsymbol{\lambda} = \{\{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K, \boldsymbol{\lambda}_w\}$ is a BCN parameterization.

Proof: Clearly, this parameterization satisfies Assumption 1 described in the main text. By Lemma 11, we know that this parameterization is a BC parameterization. Now, we will show that this parameterization also satisfies Assumption 3 in Appendix I.2.

First note that $\boldsymbol{\lambda}_w$ has only one block and it is the natural parameterization of exponential family distribution $q(w|\boldsymbol{\lambda}_w)$, which implies that $\boldsymbol{\lambda}_w$ is a BCN parameterization for $q(w|\boldsymbol{\lambda}_w)$.

Note that given the rest blocks are known and conditioning on w , $q(\mathbf{z}|w, \boldsymbol{\lambda}_z)$ can be re-expressed as follows in terms of block $\boldsymbol{\mu}_k$.

$$\begin{aligned}
 q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z) &= \exp \left(\sum_{c=1}^K \mathbb{I}(w=c) \left[-\frac{1}{2} \mathbf{z}^T \mathbf{S}_c \mathbf{z} + \mathbf{z}^T \mathbf{S}_c \boldsymbol{\mu}_c \right] - A_z(\{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K, w) \right) \\
 &= \exp \left(\underbrace{\sum_{c \neq k} \left[\mathbb{I}(w=c) \left[-\frac{1}{2} \mathbf{z}^T \mathbf{S}_c \mathbf{z} + \mathbf{z}^T \mathbf{S}_c \boldsymbol{\mu}_c \right] \right]}_{h_{z_{k1}}(w, \mathbf{z}, \boldsymbol{\lambda}_z^{[-k1]})} + \mathbb{I}(w=k) \left[-\frac{1}{2} \mathbf{z}^T \mathbf{S}_k \mathbf{z} \right] \right) \exp \left(\underbrace{\left\langle \mathbb{I}(w=k) \mathbf{S}_k \mathbf{z}, \boldsymbol{\mu}_k \right\rangle}_{\phi_{z_{k1}}(w, \mathbf{z}, \boldsymbol{\lambda}_z^{[-k1]})} - A_z(\{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K, w) \right)
 \end{aligned}$$

Similarly, for block \mathbf{S}_k , $q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z)$ can be re-expressed as follows

$$q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda}_z) = \exp \left(\underbrace{\sum_{c \neq k} [\mathbb{I}(w = c) [-\frac{1}{2} \mathbf{z}^T \mathbf{S}_c \mathbf{z} + \mathbf{z}^T \mathbf{S}_c \boldsymbol{\mu}_c]]}_{h_{z_{k_2}}(w, \mathbf{z}, \boldsymbol{\lambda}_z^{[-k_2]})} \right) \exp \left(\underbrace{\langle \mathbb{I}(w = k) [-\frac{1}{2} \mathbf{z} \mathbf{z}^T + \boldsymbol{\mu}_k \mathbf{z}^T], \mathbf{S}_k \rangle}_{\phi_{z_{k_2}}(w, \mathbf{z}, \boldsymbol{\lambda}_z^{[-k_2]})} - \underbrace{A_z(\{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K, w)}_{\lambda_z^{k_2}} \right)$$

Since this parameterization satisfies Assumption 1 to 3, this parameterization is a BCN parameterization. \square

We denote the Christoffel symbols of the first kind and the second kind for $\boldsymbol{\mu}_k$ as $\Gamma_{a_{k_1}, b_{k_1} c_{k_1}}$ and $\Gamma^{a_{k_1}}_{b_{k_1} c_{k_1}}$ respectively.

Lemma 13 For each component k , all entries of $\Gamma^{a_{k_1}}_{b_{k_1} c_{k_1}}$ for $\boldsymbol{\mu}_k$ are zeros.

Proof: The proof is very similar to the proof of Lemma 3. We will prove this by showing that all entries of $\Gamma_{a_{k_1}, b_{k_1} c_{k_1}}$ are zeros. For notation simplicity, we use $\Gamma_{a, bc}$ to denote $\Gamma_{a_{k_1}, b_{k_1} c_{k_1}}$. Let μ_k^a denote the a -th element of $\boldsymbol{\mu}_k$.

The following expression holds for any valid a, b , and c .

$$\begin{aligned} \Gamma_{a, bc} &= \frac{1}{2} \mathbb{E}_{q(z, w | \boldsymbol{\lambda})} \left[\partial_{\mu_k^b} \partial_{\mu_k^c} \partial_{\mu_k^a} A_z(\{\boldsymbol{\mu}_j, \mathbf{S}_j\}_{j=1}^K, w) \right] \\ &= \frac{1}{2} \mathbb{E}_{q(z, w | \boldsymbol{\lambda})} \left[\mathbb{I}(w = k) \partial_{\mu_k^b} \partial_{\mu_k^c} \partial_{\mu_k^a} B(\boldsymbol{\mu}_k, \mathbf{S}_k) \right] \\ &= \frac{1}{2} \mathbb{E}_{q(z, w | \boldsymbol{\lambda})} \left[\mathbb{I}(w = k) \partial_{\mu_k^b} \partial_{\mu_k^c} (\mathbf{e}_a^T \mathbf{S}_k \boldsymbol{\mu}_k) \right] \\ &= \frac{1}{2} \mathbb{E}_{q(z, w | \boldsymbol{\lambda})} \left[\mathbb{I}(w = k) \underbrace{\partial_{\mu_k^b} (\mathbf{e}_a^T \mathbf{S}_k \mathbf{e}_c)}_0 \right] = 0 \end{aligned}$$

where in the last step we use the fact that \mathbf{S}_k , \mathbf{e}_a , and \mathbf{e}_c do not depend on $\boldsymbol{\mu}_k$. \square

Similarly, we denote the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S}_k)$ as $\Gamma^{a_{k_2}}_{b_{k_2} c_{k_2}}$.

Lemma 14 For each component k , the additional term for \mathbf{S}_k is $-\hat{\mathbf{g}}_k^{[2]} \mathbf{S}_k^{-1} \hat{\mathbf{g}}_k^{[2]}$

Proof: Recall that, in the Gaussian case $\mathcal{N}(\bar{\boldsymbol{\mu}}, \bar{\mathbf{S}})$, the additional term for $\bar{\mathbf{S}}$ is $\text{Mat}(\bar{\Gamma}^{a_2}_{b_2 c_2} \hat{g}^{b_2} \hat{g}^{c_2}) = \hat{\mathbf{g}}^{[2]} \bar{\mathbf{S}}^{-1} \hat{\mathbf{g}}^{[2]}$, where $\bar{\Gamma}^{a_2}_{b_2 c_2}$ denotes the Christoffel symbols of the second kind for $\text{vec}(\bar{\mathbf{S}})$.

To prove the statement, we will show that the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S}_k)$ is exactly the same as the Gaussian case, when $\bar{\mathbf{S}} = \mathbf{S}_k$. In other words, when $\bar{\mathbf{S}} = \mathbf{S}_k$, we will show $\Gamma^{a_{k_2}}_{b_{k_2} c_{k_2}} = \bar{\Gamma}^{a_2}_{b_2 c_2}$.

We denote the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S}_k)$ using $\Gamma^{a_{k_2}}_{b_{k_2} c_{k_2}}$. By definition, the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S}_k)$ is defined as follows since $\boldsymbol{\lambda}$ is a BC parameterization.

$$\Gamma^{a_{k_2}}_{b_{k_2} c_{k_2}} = F^{a_{k_2} d_{k_2}} \Gamma_{d_{k_2}, b_{k_2} c_{k_2}}$$

We will first show that $\Gamma_{d_{k_2}, b_{k_2} c_{k_2}} = \pi_k \bar{\Gamma}_{d_2, b_2 c_2}$.

In the Gaussian case, by definition, we have

$$\bar{\Gamma}_{d_2, b_2 c_2} = \frac{1}{2} \mathbb{E}_{q(z | \bar{\boldsymbol{\lambda}})} \left[\partial_{\bar{\mathbf{S}}^b} \partial_{\bar{\mathbf{S}}^c} \partial_{\bar{\mathbf{S}}^d} A(\bar{\boldsymbol{\mu}}, \bar{\mathbf{S}}) \right] = -\frac{1}{4} \partial_{\bar{\mathbf{S}}^b} \partial_{\bar{\mathbf{S}}^c} \partial_{\bar{\mathbf{S}}^d} (\log |\bar{\mathbf{S}}|)$$

where $A(\bar{\boldsymbol{\mu}}, \bar{\mathbf{S}}) = \frac{1}{2} [\bar{\boldsymbol{\mu}}^T \bar{\mathbf{S}} \bar{\boldsymbol{\mu}} - \log |\bar{\mathbf{S}}/(2\pi)|]$ is the log partition function of the Gaussian distribution and $\bar{\mathbf{S}}^d$ denotes the d -th element of $\text{vec}(\bar{\mathbf{S}})$ in the Gaussian case.

Therefore, we have the following result in the MOG case when $\mathbf{S}_k = \bar{\mathbf{S}}$.

$$\begin{aligned}
 \Gamma_{d_{k_2}, b_{k_2} c_{k_2}} &= \frac{1}{2} \mathbb{E}_{q(z, w | \lambda)} \left[\partial_{S_k^b} \partial_{S_k^c} \partial_{S_k^d} A_z(\{\boldsymbol{\mu}_j, \mathbf{S}_j\}_{j=1}^K, w) \right] \\
 &= \frac{1}{2} \mathbb{E}_{q(z, w | \lambda)} \left[\mathbb{I}(w = k) \partial_{S_k^b} \partial_{S_k^c} \partial_{S_k^d} B(\boldsymbol{\mu}_k, \mathbf{S}_k) \right] \\
 &= \frac{1}{2} \mathbb{E}_{q(z, w | \lambda)} \left[\mathbb{I}(w = k) \partial_{S_k^b} \partial_{S_k^c} \partial_{S_k^d} \left(-\frac{1}{2} \log |\mathbf{S}_k| / (2\pi) \right) \right] \\
 &= -\frac{\pi_k}{4} \partial_{S_k^b} \partial_{S_k^c} \partial_{S_k^d} (\log |\mathbf{S}_k|) \\
 &= \pi_k \bar{\Gamma}_{d_2, b_2 c_2}
 \end{aligned}$$

where S_k^a denotes the a -th element of $\text{vec}(\mathbf{S}_k)$ and $\mathbb{E}_{q(z, w | \lambda)} [\mathbb{I}(w = k)] = \pi_k$.

Let $F_{a_{k_2} d_{k_2}}$ denote the element at position (a, d) of the sub-block matrix of the joint FIM for block $\text{vec}(\mathbf{S}_k)$ in the MOG case. Similarly, when $\mathbf{S}_k = \bar{\mathbf{S}}$, we can show that $F_{a_{k_2} d_{k_2}} = \pi_k \bar{F}_{a_2 d_2}$, where $\bar{F}_{a_2 d_2}$ denotes the element at position (a, d) of the sub-block matrix of the FIM for block $\text{vec}(\bar{\mathbf{S}})$ in the Gaussian case.

Therefore, $F^{a_{k_2} d_{k_2}} = \pi_k^{-1} \bar{F}^{a_2 d_2}$ when $\bar{\mathbf{S}} = \mathbf{S}_k$.

Finally, when $\bar{\mathbf{S}} = \mathbf{S}_k$, we obtain the desired result since

$$\Gamma_{b_{k_2} c_{k_2}}^{a_{k_2}} = F^{a_{k_2} d_{k_2}} \Gamma_{d_{k_2}, b_{k_2} c_{k_2}} = (\pi_k^{-1} \bar{F}^{a_2 d_2}) (\pi_k \bar{\Gamma}_{d_2, b_2 c_2}) = \bar{F}^{a_2 d_2} \bar{\Gamma}_{d_2, b_2 c_2} = \bar{\Gamma}_{b_2 c_2}^{a_2}$$

where $\bar{\Gamma}_{b_2 c_2}^{a_2}$ denotes the Christoffel symbols of the second kind for $\text{vec}(\bar{\mathbf{S}})$ in the Gaussian case. □

J.1. Natural Gradients

Recall that $\mathcal{L}(\boldsymbol{\lambda}) = \mathbb{E}_{q(\mathbf{z} | \boldsymbol{\lambda})} [\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z}) + \log q(\mathbf{z} | \boldsymbol{\lambda})]$, where $q(\mathbf{z} | \boldsymbol{\lambda}) = \int q(\mathbf{z}, w | \boldsymbol{\lambda}) dw$.

Lin et al. (2019a) propose to use the importance sampling technique so that the number of Monte Carlo gradient evaluations is independent of the number of mixing components K .

Note that $\boldsymbol{\lambda}_w$ is the natural parameter of exponential family distribution $q(w | \boldsymbol{\lambda}_w)$, we can obtain the natural gradient by computing the gradient w.r.t. the mean parameter as shown by Lin et al. (2019a).

$$\hat{g}_w = \partial_{\pi} \mathcal{L}.$$

where $\pi_c := \mathbb{E}_{q(w)} [\mathbb{I}(w = c)]$, $\partial_{\pi_c} \mathcal{L}$ denotes the c -th element of $\partial_{\pi} \mathcal{L}$, and the gradient $\partial_{\pi_c} \mathcal{L}$ can be computed as below as suggested by Lin et al. (2019a).

$$\partial_{\pi_c} \mathcal{L} = \mathbb{E}_{q(z)} [(\delta_c - \delta_K) b(\mathbf{z})]$$

where $b(\mathbf{z}) := \ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z}) + \log q(\mathbf{z} | \boldsymbol{\lambda})$, and $\delta_c := \mathcal{N}(\mathbf{z} | \boldsymbol{\mu}_c, \mathbf{S}_c) / \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{z} | \boldsymbol{\mu}_k, \mathbf{S}_k)$.

Recall that $\boldsymbol{\lambda}_w$ is unconstrained in this case, there is no need to compute the addition term for $\boldsymbol{\lambda}_w$.

Now, we discuss how to compute the natural gradients $\{\hat{g}_c^{[1]}, \hat{g}_c^{[2]}\}_{c=1}^K$. Since $\{\boldsymbol{\mu}_c, \mathbf{S}_c\}_{c=1}^K$ are BCN parameters, we can obtain the natural gradients by computing gradients w.r.t. its BC expectation parameter due to Theorem 4.

Given the rest of blocks are known, the BC expectation parameter for block $\boldsymbol{\mu}_k$ is

$$\mathbf{m}_{k_1} = \mathbb{E}_{q(w, z)} [\mathbb{I}(w = k) (\mathbf{S}_k \mathbf{z})] = \pi_k \mathbf{S}_k \boldsymbol{\mu}_k$$

In this case, we know that $\partial_{\mu_k} \mathcal{L} = \pi_k \mathbf{S}_k \partial_{m_{k_1}} \mathcal{L}$. Therefore, the natural gradient w.r.t. $\boldsymbol{\mu}_k$ is $\hat{g}_k^{[1]} = \partial_{m_{k_1}} \mathcal{L} = \pi_k^{-1} \mathbf{S}_k^{-1} \partial_{\mu_k} \mathcal{L} = \pi_k^{-1} \boldsymbol{\Sigma}_k \partial_{\mu_k} \mathcal{L}$, where the gradient $\partial_{\mu_k} \mathcal{L}$ can be computed as belows as suggested by Lin et al. (2019a).

$$\partial_{\mu_k} \mathcal{L} = \mathbb{E}_{q(z)} [\pi_k \delta_k \nabla_z b(\mathbf{z})]$$

Likewise, given the rest of blocks are known, the BC expectation parameter for block \mathbf{S}_k is

$$\mathbf{m}_{k_2} = \mathbb{E}_{q(w,z)} [\mathbb{I}(w = k) (-\frac{1}{2} \mathbf{z} \mathbf{z}^T + \boldsymbol{\mu}_k \mathbf{z}^T)] = \frac{\pi_k}{2} (\boldsymbol{\mu}_k \boldsymbol{\mu}_k^T - \mathbf{S}_k^{-1})$$

Therefore, the natural gradient w.r.t. \mathbf{S}_k is $\hat{\mathbf{g}}_k^{[2]} = \partial_{m_{k_2}} \mathcal{L} = -\frac{2}{\pi_k} \partial_{\mathbf{S}_k^{-1}} f = -\frac{2}{\pi_k} \partial_{\Sigma_k} f$, where the gradient $\partial_{\Sigma_k} f$ can be computed as belows as suggested by Lin et al. (2019a).

$$\partial_{\Sigma_k} \mathcal{L} = \frac{1}{2} \mathbb{E}_{q(z)} [\pi_k \delta_k \nabla_z^2 b(\mathbf{z})]$$

Alternatively, we can use the re-parametrization trick to compute the gradient as below.

$$\partial_{\Sigma_k} \mathcal{L} = \frac{1}{2} \mathbb{E}_{q(z)} [\pi_k \delta_k \mathbf{S}_k (\mathbf{z} - \boldsymbol{\mu}_k) \nabla_z^T b(\mathbf{z})]$$

By Lemma 13 and 14, the proposed update induced by our rule is

$$\begin{aligned} \log(\pi_c/\pi_K) &\leftarrow \log(\pi_c/\pi_K) - t \mathbb{E}_{q(z)} [(\delta_c - \delta_K) b(\mathbf{z})] \\ \boldsymbol{\mu}_c &\leftarrow \boldsymbol{\mu}_c - t \mathbf{S}_c^{-1} \mathbb{E}_{q(z)} [\delta_c \nabla_z b(\mathbf{z})] + \mathbf{0} \\ \mathbf{S}_c &\leftarrow \mathbf{S}_c - t \hat{\mathbf{G}}_c + \frac{t^2}{2} \hat{\mathbf{G}}_c (\mathbf{S}_c)^{-1} \hat{\mathbf{G}}_c \end{aligned} \quad (27)$$

where we do not compute the additional term for $\boldsymbol{\lambda}_w$ since $\boldsymbol{\lambda}_w$ is unconstrained, $\delta_c := \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_c, \mathbf{S}_c) / \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_k, \mathbf{S}_k)$, $b(\mathbf{z}) := \ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z}) + \log q(\mathbf{z}|\boldsymbol{\lambda})$ and $\hat{\mathbf{G}}_c$ can be computed as below.

Note that $b(\mathbf{z})$ can be the logarithm of an unnormalized target function as such $b(\mathbf{z}) = \bar{\ell}(\mathbf{z}) + \text{Constant} + \log q(\mathbf{z}|\boldsymbol{\lambda})$. Recall that $\ell(\mathcal{D}, \mathbf{z}) - \log p(\mathbf{z}) = \bar{\ell}(\mathbf{z}) + \text{Constant}$. Lin et al. (2019a) suggest using the Hessian trick to compute $\hat{\mathbf{G}}_c$ as shown in (29). We can also use the re-parameterization trick to compute $\hat{\mathbf{G}}_c$ as shown in (28).

$$\hat{\mathbf{G}}_c = -\mathbb{E}_{q(z)} [\delta_c \mathbf{S}_c (\mathbf{z} - \boldsymbol{\mu}_c) \nabla_z^T b(\mathbf{z})] = -\mathbb{E}_{q(z)} [\delta_c \mathbf{S}_c (\mathbf{z} - \boldsymbol{\mu}_c) \nabla_z^T \bar{\ell}(\mathbf{z})] - \mathbb{E}_{q(z)} [\delta_c \nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda})] \quad (28)$$

$$= -\mathbb{E}_{q(z)} [\delta_c \nabla_z^2 b(\mathbf{z})] = -\mathbb{E}_{q(z)} [\delta_c \nabla_z^2 \bar{\ell}(\mathbf{z})] - \mathbb{E}_{q(z)} [\delta_c \nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda})]. \quad (29)$$

We use the MC approximation to compute $\hat{\mathbf{G}}_c$ as below.

$$\begin{aligned} \hat{\mathbf{G}}_c &\approx -\delta_c \left(\frac{\bar{\mathbf{S}}_c + \bar{\mathbf{S}}_c^T}{2} + \nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda}) \right) && \text{referred to as “-rep”} \\ \hat{\mathbf{G}}_c &\approx -\delta_c \left(\nabla_z^2 \bar{\ell}(\mathbf{z}) + \nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda}) \right) && \text{referred to as “-hess”} \end{aligned}$$

where $\mathbf{z} \sim q(\mathbf{z}|\boldsymbol{\lambda})$, $\bar{\mathbf{S}}_c := \mathbf{S}_c (\mathbf{z} - \boldsymbol{\mu}_c) \nabla_z^T \bar{\ell}(\mathbf{z})$ and $\nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda})$ can be manually coded or computed by Auto-Diff.

Recall that when $q(\mathbf{z}|\boldsymbol{\lambda})$ is Gaussian, $-\mathbb{E}_{q(z)} [\nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda})] = \boldsymbol{\Sigma}^{-1}$, which is positive definite. VOGN is proposed to approximate $\mathbb{E}_{q(z)} [\nabla_z^2 \bar{\ell}(\mathbf{z})]$ by a positive definite matrix when $q(\mathbf{z}|\boldsymbol{\lambda})$ is Gaussian. In MOG cases, $-\mathbb{E}_{q(z)} [\nabla_z^2 \log q(\mathbf{z}|\boldsymbol{\lambda})]$ is no longer a positive definite matrix. VOGN does not guarantee that the update for \mathbf{S}_c stays in the constraint set. Furthermore, directly approximating $-\hat{\mathbf{G}}_c$ by naively extending the idea of VOGN does not give a good posterior approximation. Unlike VOGN, our update satisfies the constraint without the loss of the approximation accuracy for both Gaussian and MOG cases.

K. Example: Skew Gaussian Approximation

We consider the skew Gaussian approximation proposed by Lin et al. (2019a). The joint distribution is given below.

$$q(\mathbf{z}, w|\boldsymbol{\alpha}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = q(\mathbf{z}|w, \boldsymbol{\alpha}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) \mathcal{N}(w|0, 1)$$

$$q(\mathbf{z}|w, \boldsymbol{\alpha}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu} + |w|\boldsymbol{\alpha}, \boldsymbol{\Sigma})$$

$$= \exp\left(\left\{ \text{Tr}\left(-\frac{1}{2} \boldsymbol{\Sigma}^{-1} \mathbf{z} \mathbf{z}^T\right) + |w| \boldsymbol{\alpha}^T \boldsymbol{\Sigma}^{-1} \mathbf{z} + \boldsymbol{\mu}^T \boldsymbol{\Sigma}^{-1} \mathbf{z} - \frac{1}{2} ((\boldsymbol{\mu} + |w|\boldsymbol{\alpha})^T \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu} + |w|\boldsymbol{\alpha}) + \log |2\pi \boldsymbol{\Sigma}|) \right\}\right)$$

We consider the parameterization $\lambda = \left\{ \begin{bmatrix} \mu \\ \alpha \end{bmatrix}, \mathbf{S} \right\}$, where $\mathbf{S} = \Sigma^{-1}$, $\lambda^{[1]} = \begin{bmatrix} \mu \\ \alpha \end{bmatrix}$, and $\lambda^{[2]} = \mathbf{S}$. The open-set constraint is $\lambda \in \mathbb{R}^{2d} \times \mathbb{S}_{++}^{d \times d}$. Under this parameterization, the distribution $q(\mathbf{z}|w)$ can be re-expressed as below.

$$q(\mathbf{z}|w, \lambda) = \exp \left\{ \text{Tr} \left(-\frac{1}{2} \mathbf{S} \mathbf{z} \mathbf{z}^T \right) + \mathbf{z}^T \mathbf{S} (\mathbf{Q}(w))^T \lambda^{[1]} - A_z(\lambda, w) \right\}$$

where $\mathbf{Q}(w) := \begin{bmatrix} \mathbf{I}_d \\ |w| \mathbf{I}_d \end{bmatrix}$ is a $2d$ -by- d matrix and $A_z(\lambda, w) = \frac{1}{2} \left[\begin{bmatrix} \mu^T & \alpha^T \end{bmatrix} \mathbf{Q}(w) \mathbf{S} (\mathbf{Q}(w))^T \begin{bmatrix} \mu \\ \alpha \end{bmatrix} - \log |\mathbf{S}|/(2\pi) \right]$.

Lemma 15 *The joint FIM is block diagonal with two blocks under this parameterization.*

$$\mathbf{F} = \begin{bmatrix} \mathbf{F}^{[1]} & \mathbf{0} \\ \mathbf{0} & \mathbf{F}^{[2]} \end{bmatrix}$$

Therefore, this parameterization is a BC parameterization.

Proof: We will prove this lemma by showing that all cross terms shown in red are zeros.

Let's denote λ^{a1} be the a -th element of $\lambda^{[1]}$ and S^{bc} be the element of \mathbf{S} at position (b, c) . Furthermore, \mathbf{e}_a denotes an one-hot vector where all entries are zeros except the a -th entry with value 1, and \mathbf{I}_{bc} denotes an one-hot matrix where all entries are zeros except the entry at position (b, c) with value 1.

By definition, the cross term is defined as belows.

$$\begin{aligned} & -\mathbb{E}_{q(\mathbf{z}, w|\lambda)} [\partial \lambda^{a1} \partial S^{bc} \log q(\mathbf{z}, w|\lambda)] \\ &= -\mathbb{E}_{q(\mathbf{z}, w|\lambda)} \left[\mathbf{z}^T \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a - \left(\lambda^{[1]} \right)^T \mathbf{Q}(w) \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a \right] \\ &= -\mathbb{E}_{q(w)} \left[\mathbb{E}_{q(\mathbf{z}|w, \lambda)} \left[\mathbf{z}^T \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a - \left(\lambda^{[1]} \right)^T \mathbf{Q}(w) \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a \right] \right] \\ &= -\mathbb{E}_{q(w)} \left[\mathbb{E}_{q(\mathbf{z}|w, \lambda)} \left[\mathbf{z}^T \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a \right] - \left(\lambda^{[1]} \right)^T \mathbf{Q}(w) \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a \right] \\ &= -\mathbb{E}_{q(w)} \left[\left(\lambda^{[1]} \right)^T \mathbf{Q}(w) \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a - \left(\lambda^{[1]} \right)^T \mathbf{Q}(w) \mathbf{I}_{bc} (\mathbf{Q}(w))^T \mathbf{e}_a \right] = 0 \end{aligned}$$

where we use the following expression in the last step.

$$\mathbb{E}_{q(\mathbf{z}|w, \lambda)} [\mathbf{z}] = |w| \alpha + \mu = (\mathbf{Q}(w))^T \lambda^{[1]}$$

□

Note that another parameterization $\{\mu, \alpha, \mathbf{S}\}$ is **not** a BC parameterization since the joint FIM is not block-diagonal under this parameterization.

Lemma 16 *Parameterization λ is a BCN parameterization.*

Proof: Clearly, this parameterization satisfies Assumption 1 described in the main text. By Lemma 15, we know that this parameterization is a BC parameterization. Now, we will show that this parameterization also satisfies Assumption 3 in Appendix I.2.

Note that given the rest blocks are known and conditioning on w , $q(\mathbf{z}|w, \lambda)$ can be re-expressed as follows in terms of block $\lambda^{[1]}$.

$$\begin{aligned} q(\mathbf{z}|w, \lambda) &= \exp \left\{ \text{Tr} \left(-\frac{1}{2} \mathbf{S} \mathbf{z} \mathbf{z}^T \right) + \mathbf{z}^T \mathbf{S} (\mathbf{Q}(w))^T \lambda^{[1]} - A_z(\lambda, w) \right\} \\ &= \underbrace{\exp \left\{ \text{Tr} \left(-\frac{1}{2} \mathbf{S} \mathbf{z} \mathbf{z}^T \right) \right\}}_{h_1(w, \mathbf{z}, \lambda^{[-1]})} \exp \left[\underbrace{\langle \mathbf{Q}(w) \mathbf{S} \mathbf{z}, \lambda^{[1]} \rangle}_{\phi_1(w, \mathbf{z}, \lambda^{[-1]})} - A_z(\lambda, w) \right] \end{aligned}$$

Similarly, for block \mathbf{S} , $q(\mathbf{z}|\mathbf{w}, \boldsymbol{\lambda})$ can be re-expressed as follows

$$q(\mathbf{z}|w, \boldsymbol{\lambda}) = \underbrace{1}_{h_2(w, \mathbf{z}, \boldsymbol{\lambda}^{[-2]})} \exp \left[\underbrace{\left\langle -\frac{1}{2} \mathbf{z} \mathbf{z}^T + \mathbf{z} \left(\boldsymbol{\lambda}^{[1]} \right)^T \mathbf{Q}(w), \mathbf{S} \right\rangle}_{\phi_2(w, \mathbf{z}, \boldsymbol{\lambda}^{[-2]})} - A_z(\boldsymbol{\lambda}, w) \right]$$

Since this parameterization satisfies Assumption 1 to 3, this parameterization is a BCN parameterization. \square

We denote the Christoffel symbols of the first kind and the second kind for $\boldsymbol{\lambda}^{[1]}$ as $\Gamma_{a_1, b_1 c_1}$ and $\Gamma^{a_1}_{b_1 c_1}$ respectively.

Lemma 17 All entries of $\Gamma^{a_1}_{b_1 c_1}$ for $\boldsymbol{\lambda}^{[1]}$ are zeros.

Proof: We will prove this by showing that all entries of $\Gamma^{a_1}_{b_1 c_1}$ are zeros. Let λ^{a_1} denote the a -th element of $\boldsymbol{\lambda}^{[1]}$.

The following expression holds for any valid a , b , and c .

$$\begin{aligned} \Gamma_{a_1, b_1 c_1} &= \frac{1}{2} \mathbb{E}_{q(z, w|\boldsymbol{\lambda})} [\partial_{\lambda^{b_1}} \partial_{\lambda^{c_1}} \partial_{\lambda^{a_1}} A_z(\boldsymbol{\lambda}, w)] \\ &= \frac{1}{2} \mathbb{E}_{q(z, w|\boldsymbol{\lambda})} \left[\partial_{\lambda^{b_1}} \partial_{\lambda^{c_1}} \left((\mathbf{e}_a)^T \mathbf{Q}(w) \mathbf{S} (\mathbf{Q}(w))^T \boldsymbol{\lambda}^{[1]} \right) \right] \\ &= \frac{1}{2} \mathbb{E}_{q(z, w|\boldsymbol{\lambda})} \left[\partial_{\lambda^{b_1}} \left(\mathbf{e}_a^T \mathbf{Q}(w) \mathbf{S} (\mathbf{Q}(w))^T \mathbf{e}_c \right) \right] = 0 \end{aligned}$$

where in the last step we use the fact that \mathbf{S} , $\mathbf{Q}(w)$, \mathbf{e}_a , and \mathbf{e}_c do not depend on $\boldsymbol{\lambda}^{[1]}$. \square

We denote the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S})$ as $\Gamma^{a_2}_{b_2 c_2}$.

Lemma 18 The additional term for \mathbf{S} is $-\hat{\mathbf{g}}^{[2]} \mathbf{S}^{-1} \hat{\mathbf{g}}^{[2]}$

Proof: Recall that, in the Gaussian case $\mathcal{N}(\bar{\boldsymbol{\mu}}, \bar{\mathbf{S}})$, the additional term for $\bar{\mathbf{S}}$ is $\text{Mat}(\bar{\Gamma}^{a_2}_{b_2 c_2} \hat{g}^{b_2} \hat{g}^{c_2}) = \hat{\mathbf{g}}^{[2]} \bar{\mathbf{S}}^{-1} \hat{\mathbf{g}}^{[2]}$, where $\bar{\Gamma}^{a_2}_{b_2 c_2}$ denotes the Christoffel symbols of the second kind for $\text{vec}(\bar{\mathbf{S}})$.

To prove the statement, we will show that the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S})$ is exactly the same as the Gaussian case, when $\bar{\mathbf{S}} = \mathbf{S}$.

We denote the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S})$ as $\Gamma^{a_2}_{b_2 c_2}$. By definition, the Christoffel symbols of the second kind for $\text{vec}(\mathbf{S})$ is defined as follows.

$$\Gamma^{a_2}_{b_2 c_2} = F^{a_2 d_2} \Gamma_{d_2, b_2 c_2}$$

We will show that $\Gamma_{a_2, b_2 c_2} = \bar{\Gamma}_{a_2, b_2 c_2}$.

In the Gaussian case, we have

$$\bar{\Gamma}_{d_2, b_2 c_2} = -\frac{1}{4} \partial_{\bar{S}^b} \partial_{\bar{S}^c} \partial_{\bar{S}^d} (\log |\bar{\mathbf{S}}|)$$

where $A(\bar{\boldsymbol{\mu}}, \bar{\mathbf{S}}) = \frac{1}{2} [\bar{\boldsymbol{\mu}}^T \bar{\mathbf{S}} \bar{\boldsymbol{\mu}} - \log |\bar{\mathbf{S}}|/(2\pi)]$ is the log partition function of the Gaussian distribution and \bar{S}^a is the a -th element of $\text{vec}(\bar{\mathbf{S}})$ in the Gaussian case.

Therefore, we have the following result when $\bar{\mathbf{S}} = \mathbf{S}$.

$$\Gamma_{d_2, b_2 c_2} = \frac{1}{2} \mathbb{E}_{q(z, w|\boldsymbol{\lambda})} [\partial_{S^b} \partial_{S^c} \partial_{S^d} A_z(\boldsymbol{\lambda}, w)] = -\frac{1}{4} \partial_{S^b} \partial_{S^c} \partial_{S^d} \log |\mathbf{S}| = \bar{\Gamma}_{d_2, b_2 c_2}$$

where S^a denotes the a -th element of $\text{vec}(\mathbf{S})$.

Let $F_{a_2 d_2}$ denote the element at position (a, d) of the sub-block matrix of the joint FIM for $\text{vec}(\mathbf{S})$. Similarly, we can show that $F_{a_2 d_2} = \bar{F}_{a_2 d_2}$, where $\bar{F}_{a_2 d_2}$ denotes the element at position (a, d) of the FIM for $\text{vec}(\bar{\mathbf{S}})$ in the Gaussian case. Therefore, $F^{a_2 d_2} = \bar{F}^{a_2 d_2}$.

Finally, when $\bar{\mathbf{S}} = \mathbf{S}$, we obtain the desired result since

$$\Gamma_{b_2 c_2}^{a_2} = F^{a_2 d_2} \Gamma_{d_2, b_2 c_2} = \bar{F}^{a_2 d_2} \bar{\Gamma}_{d_2, b_2 c_2} = \bar{\Gamma}_{b_2 c_2}^{a_2}$$

where $\bar{\Gamma}_{b_2 c_2}^{a_2}$ denotes the Christoffel symbols of the second kind for $\text{vec}(\bar{\mathbf{S}})$ in the Gaussian case. \square

Using these lemmas, the proposed update induced by our rule is

$$\begin{aligned} \begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\alpha} \end{bmatrix} &\leftarrow \begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\alpha} \end{bmatrix} - t \hat{\mathbf{g}}^{[1]} + \mathbf{0} \\ \mathbf{S} &\leftarrow \mathbf{S} - t \hat{\mathbf{g}}^{[2]} + \frac{t^2}{2} \hat{\mathbf{g}}^{[2]} \mathbf{S}^{-1} \hat{\mathbf{g}}^{[2]} \end{aligned}$$

where $\hat{\mathbf{g}}^{[1]}$ and $\hat{\mathbf{g}}^{[2]}$ are natural gradients.

Similarly, it can be shown that the above update satisfies the underlying constraints.

K.1. Natural Gradients

Now, we discuss how to compute the natural gradients. Since the parameterization is a BCN parameterization, gradients w.r.t. BC expectation parameters are natural gradients for BCN parameters due to Theorem 4.

Recall that $\boldsymbol{\lambda}^{[1]} = \begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\alpha} \end{bmatrix}$. Let $\mathbf{m}_{[1]} = \begin{bmatrix} \mathbf{m}_\mu \\ \mathbf{m}_\alpha \end{bmatrix}$ denote the BC expectation parameter for $\boldsymbol{\lambda}^{[1]}$. Given \mathbf{S} is known, the BC expectation parameter is

$$\begin{aligned} \begin{bmatrix} \mathbf{m}_\mu \\ \mathbf{m}_\alpha \end{bmatrix} &= \mathbb{E}_{q(w, z)} [\mathbf{Q}(w) \mathbf{S} \mathbf{z}] \\ &= \mathbb{E}_{q(w)} [\mathbf{Q}(w) \mathbf{S} (\mathbf{Q}(w))^T \boldsymbol{\lambda}^{[1]}] \\ &= \mathbb{E}_{q(w)} \left[\begin{bmatrix} \mathbf{S} & |w| \mathbf{S} \\ |w| \mathbf{S} & w^2 \mathbf{S} \end{bmatrix} \begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\alpha} \end{bmatrix} \right] \\ &= \begin{bmatrix} \mathbf{S} & c\mathbf{S} \\ c\mathbf{S} & \mathbf{S} \end{bmatrix} \begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\alpha} \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{S}\boldsymbol{\mu} + c\mathbf{S}\boldsymbol{\alpha} \\ c\mathbf{S}\boldsymbol{\mu} + \mathbf{S}\boldsymbol{\alpha} \end{bmatrix} \end{aligned}$$

where $c = \mathbb{E}_{q(w)} [|w|] = \sqrt{\frac{2}{\pi}}$.

Since $\mathbf{S} = \boldsymbol{\Sigma}^{-1}$, we have the following expressions.

$$\boldsymbol{\mu} = \frac{1}{1 - c^2} \boldsymbol{\Sigma} (\mathbf{m}_\mu - c\mathbf{m}_\alpha), \quad \boldsymbol{\alpha} = \frac{1}{1 - c^2} \boldsymbol{\Sigma} (\mathbf{m}_\alpha - c\mathbf{m}_\mu)$$

By the chain rule, we have

$$\partial_{m_\mu} \mathcal{L} = \boldsymbol{\Sigma} \left(\frac{1}{1 - c^2} \partial_\mu \mathcal{L} - \frac{c}{1 - c^2} \partial_\alpha \mathcal{L} \right), \quad \partial_{m_\alpha} \mathcal{L} = \boldsymbol{\Sigma} \left(\frac{1}{1 - c^2} \partial_\alpha \mathcal{L} - \frac{c}{1 - c^2} \partial_\mu \mathcal{L} \right)$$

Therefore, the natural gradient w.r.t. $\boldsymbol{\lambda}^{[1]} = \begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\alpha} \end{bmatrix}$ is $\hat{\mathbf{g}}^{[1]} = \begin{bmatrix} \partial_{m_\mu} \mathcal{L} \\ \partial_{m_\alpha} \mathcal{L} \end{bmatrix}$ where the gradient $\partial_\mu \mathcal{L}$ and $\partial_\alpha \mathcal{L}$ can be computed as suggested by Lin et al. (2019a).

Likewise, the BC expectation parameter for block \mathbf{S} is

$$\mathbf{m}_{[2]} = \mathbb{E}_{q(w, z)} \left[-\frac{1}{2} \mathbf{z} \mathbf{z}^T + \mathbf{z} \left(\boldsymbol{\lambda}^{[1]} \right)^T \mathbf{Q}(w) \right] = -\frac{1}{2} \mathbf{S}^{-1} + \mathbb{E}_{q(w)} \left[\frac{1}{2} (\mathbf{Q}(w))^T \boldsymbol{\lambda}^{[1]} \left(\boldsymbol{\lambda}^{[1]} \right)^T \mathbf{Q}(w) \right]$$

Since $\boldsymbol{\lambda}^{[1]}$ is known, $\mathbb{E}_{q(w)} \left[\frac{1}{2} (\mathbf{Q}(w))^T \boldsymbol{\lambda}^{[1]} \left(\boldsymbol{\lambda}^{[1]} \right)^T \mathbf{Q}(w) \right]$ does not depend on \mathbf{S} . Therefore, the natural gradient w.r.t. \mathbf{S} is $\hat{\mathbf{g}}^{[2]} = \partial_{m_{[2]}} \mathcal{L} = -2\partial_{\mathbf{S}^{-1}} \mathcal{L} = -2\partial_{\boldsymbol{\Sigma}} \mathcal{L}$, where we compute the gradient $\partial_{\boldsymbol{\Sigma}} \mathcal{L}$ as suggested by Lin et al. (2019a).

L. More Results




Figure 5. The leftmost figure is MOG approximations for the banana distribution mentioned at Section 6.1, where the number indicates the number of components used in the approximations. The middle figure is a complete version of MOG approximations for the double banana distribution (the rightmost plot in Figure 2), where the number indicates the number of components used in the approximations. The rightmost figure is MOG approximations for the posterior $p(\mathbf{z}|y=1)$ of a BNN with a Gaussian prior $p(\mathbf{z}) = \mathcal{N}(\mathbf{z}|\mathbf{0}, \mathbf{I})$ and a NN likelihood $p(y|\mathbf{z}) = \mathcal{N}(y|3z_1^2(z_1^2 - 1) + z_2^2, 0.5^2)$, where the number indicates the number of components used in the approximations.




Figure 6. The leftmost plot is mean-field Gaussian approximations for the toy Bayesian logistic regression example considered at Section 6.1, where Vadam is proposed by Khan et al. (2018). The rightmost plot is a skew-Gaussian approximation with full covariance structure for the same example.




Figure 7. We compare our method to the unconstrained transformation method proposed by Salimbeni et al. (2018) in the Bayesian logistic model with a full Gaussian approximation, where we use the re-parametrization trick to avoid computing the Hessian ($\nabla_z^2 \bar{\ell}(\mathbf{z})$). Both methods are implemented and tested in the same environment. We tune the step size for each method by grid search. Our method can use a bigger step-size than theirs while still optimizing the objective function and maintaining numerical stability. Moreover, our method requires less memory than theirs. The leftmost plot shows the performance of both methods in terms of the number of iterations. The rightmost plot shows the performance of both methods from another perspective in terms of time, where both methods iterate 5,000 iterations. From this plot, we can clearly see that our method has a lower iteration cost than theirs, where the standard gradient computation time also is considered. If the standard gradient ($\nabla_z \bar{\ell}(\mathbf{z})$) is computed beforehand, our update is at least 6 ~ 10 times faster than theirs.




Figure 8. This is a complete version of the leftmost figure in Figure 2. The figure shows MOG approximation (with $K = 25$) to fit an MOG model with 10 components in a 20 dimensional problem.




Figure 9. This is the first 60 marginal distributions obtained from a MOG approximation with $K = 60$ for a 300-dimensional mixture of Student's T distributions with 20 components. We describe the problem at Section 6.1, where the approximation is obtained by our method at the 50,000-th iteration.




Figure 10. This is the second 60 marginal distributions obtained from a MOG approximation with $K = 60$ for a 300-dimensional mixture of Student's T distributions with 20 components. We describe the problem at Section 6.1, where the approximation is obtained by our method at the 50,000-th iteration.




Figure 11. This is the third 60 marginal distributions obtained from a MOG approximation with $K = 60$ for a 300-dimensional mixture of Student's T distributions with 20 components. We describe the problem at Section 6.1, where the approximation is obtained by our method at the 50,000-th iteration.




Figure 12. This is the fourth 60 marginal distributions obtained from a MOG approximation with $K = 60$ for a 300-dimensional mixture of Student's T distributions with 20 components. We describe the problem at Section 6.1, where the approximation is obtained by our method at the 50,000-th iteration.




Figure 13. This is the last 60 marginal distributions obtained from a MOG approximation with $K = 60$ for a 300-dimensional mixture of Student's T distributions with 20 components. We describe the problem at Section 6.1, where the approximation is obtained by our method at the 50,000-th iteration.